Parallel genetic adaptation amidst a background of changing effective population sizes in

divergent yellow perch (*Perca flavescens*) populations

Xiaoshen Yin[1,2,*], Claire E. Schraidt[3,4], Morgan M. Sparks[2,5], Peter T. Euclide[3,6], Tyler J. Hoyt[7],

Carl R. Ruetz III[7], Tomas O. Höök[3,6], Mark R. Christie[2,3,*]

[1] MOE Key Laboratory of Marine Genetics and Breeding, College of Marine Life Sciences,
Ocean University of China, Qingdao 266003, China

[2] Department of Biological Sciences, Purdue University; 915 W. State St., West Lafayette,
Indiana 47907-2054 USA

[3] Department of Forestry and Natural Resources, Purdue University; 715 W. State St., West
Lafayette, Indiana 47907-2054 USA

[4] Department of Ecology and Evolutionary Biology, Yale University; 165 Prospect Street, New
Haven, Connecticut 06511 USA

[5] US Forest Service, Rocky Mountain Research Station; 322 East Front Street, Boise, Idaho
83702 USA

[6] Illinois-Indiana Sea Grant, Purdue University, West Lafayette, Indiana 47907 USA

[7] Annis Water Resources Institute, Grand Valley State University; 740 W. Shoreline Dr.,
Muskegon, Michigan 49441 USA

* Corresponding authors:

Xiaoshen Yin; MOE Key Laboratory of Marine Genetics and Breeding, College of Marine Life
Sciences, Ocean University of China, Qingdao 266003; yinxiaoshen@ouc.edu.cn

Mark R. Christie; Department of Biological Sciences, Purdue University; 915 W. State St., West
Lafayette, Indiana 47907; christ99@purdue.edu, 765-494-2070

Running title: Adaptive divergence and *Ne*

Abstract

Aquatic ecosystems are highly dynamic environments vulnerable to natural and anthropogenic

disturbances. High-economic value fisheries are one of many ecosystem services affected by

these disturbances and it is critical to accurately characterize the genetic diversity and effective

population sizes of valuable fish stocks through time. We used genome-wide data to reconstruct

the demographic histories of economically important yellow perch (*Perca flavescens*)

populations. In two isolated and genetically divergent populations, we provide independent

evidence for simultaneous increases in effective population sizes over both historic and

contemporary time scales including negative genome-wide estimates of Tajima's D, 3.1 times

more SNPs than adjacent populations, and contemporary effective population sizes that have

increased 10- and 47-fold from their minimum, respectively. The excess of segregating sites and

negative Tajima's D values likely arose from mutations accompanying historic population

expansions with insufficient time for purifying selection, whereas linkage disequilibrium-based

estimates of Ne also suggest contemporary increases that may have been driven by reduced

fishing pressure or environmental remediation. We also identified parallel, genetic adaptation to

reduced visual clarity in the same two habitats. These results suggest that the synchrony of key

ecological and evolutionary processes can drive parallel demographic and evolutionary

trajectories across independent populations.

Keywords: demographic history, effective population size, fisheries, genetic adaptation, genetic

diversity, gene flow, parallel evolution

Introduction

Accurate estimates of effective population sizes (Ne) are critical for the successful management and conservation of exploited, threatened, and endangered species. The concept of Ne was first introduced in 1931, and often refers to the size of an idealized population with constant size and random mating that experiences an equivalent rate of genetic drift as the population under study (1). Consequently, estimates of effective population size can often be smaller than estimates of census population sizes (Ne/N ratio ranging from 10^{-5} to ~0.5) (2–13). From a conservation and management perspective, effective population size matters in several key aspects: 1.) Ne is a major determinant of nucleotide variation (*i.e.*, *4Neμ* under a neutral model), 2.) Ne measures the expected rate of genetic drift, a major, stochastic, evolutionary force acting on genetic diversity across the genome and through generations, and 3.) Ne provides a unified frame of reference for comparisons across species or populations (14,15). Thus, changes in effective population sizes through time can provide information on how populations are responding to novel stressors, such as climate change (16–19), changes in food web dynamics (20), habit alteration and destruction (17,20), and overexploitation (16–18,21,22).

Accurate estimates of effective population sizes are particularly needed in fisheries to guide sustainable harvest practices in economically important populations. Ne is a critical metric for the successful assessment and management of fisheries stocks as it not only provides a fisheries-independent approach, avoiding the biases inherent in some field- or catch-based methods (23), but also reflects the adaptative potential of different stocks under constantly changing environments (23,24). Estimates of effective population sizes may forecast overharvest and potential collapses in commercial fisheries, as signaled by either substantial reductions in effective population sizes or increasing ratios of effective to census population sizes (*i.e.*,

overharvest reduces both effective and census population sizes but reduces census population

size more quickly, so the ratio of effective to census population size goes up) (4,25–27).

Nevertheless, several challenges remain for estimating effective population sizes in fish and

fisheries. Estimates of *Ne* often include infinity when the signal for genetic variation arising from

genetic drift cannot be estimated with high precision and is instead overwhelmed by noise arising

from sampling error (28,29). Also, fish populations are often highly connected due to extensive

larval dispersal and juvenile or adult migration (30–33) and such migration and subsequent gene

flow can bias, or at the very least complicate estimates of *Ne* (14,34,35). Additional challenges

include overlapping generations (34–36), varied genomic architectures (34), rare alleles (34),

small sample sizes (34,35), sampling design (34), and sequencing or genotyping errors (34).

Despite these challenges, advances in high throughput sequencing and the influx of genomic data

have resulted in the development of many methods (37–41) that more accurately infer temporal

changes in effective population sizes for various data types, including polymorphic sites, short

sequence blocks, and whole-genome data (14).

Empirical assessments of *Ne* over contemporary time scales (*e.g.*, decades to centuries)

have largely demonstrated reductions in effective population sizes (16–22,42–45). Although

rarely documented (6,45,46), contemporary increases in effective population sizes exist and can

occur via one of three mechanisms. First, a reduction in the variance in reproductive success, for

example by equalizing mating success throughout a single generation, can result in larger

effective population sizes in the next generation (47,48), as demonstrated in captive breeding

programs where equalization of full-sib family sizes is often recommended to increase effective

population sizes (49–52). Second, introgression from neighboring, genetically divergent

populations may generate biases in estimating *Ne*, the direction of which is determined by the

magnitude and continuity of gene flow (34,35). Third, large, consistent increases in adult

abundance within a population, perhaps due to improved environmental conditions and

management policies, can also cause increases in effective population sizes (46). Over longer

time scales, these last two mechanisms may also be accompanied by several other genetic

indicators including negative values of Tajima's D, shifts in site frequency spectra, and larger

numbers of polymorphic loci with low minor allele frequency (14).

In this study, we explored changes in the effective population sizes and patterns of

adaptive divergence in yellow perch (*Perca flavescens*) populations found throughout the main

basin of Lake Michigan and two connected water bodies, Green Bay and Muskegon Lake (Fig.

1). Lake Michigan, with a drainage basin of about 118,000 km^2, is the fourth largest freshwater

lake in the world (53,54). Green Bay, an inlet of northwestern Lake Michigan, drains

approximately 40,468 km^2 while Muskegon Lake, a drowned river mouth (DRM) lake of 16.8

km^2, receives water from the Muskegon River and drains directly into Lake Michigan. DRM

lakes formed as the mouths of rivers were cut off from flowing into the Great Lakes following

the retreat of glaciers. The melting river water was trapped behind land formations, carving out

basins like Muskegon Lake, and eventually reconnecting after the buildup of water pressure

created channels allowing them to once again flow into the Great Lakes (55,56). The main basin

of Lake Michigan, Green Bay, and Muskegon Lake are all home to yellow perch, but are vastly

different in both abiotic (*e.g.*, water clarity) and biotic (*e.g.*, community composition)

characteristics and have distinct environmental histories. Yellow perch migrate between

nearshore Lake Michigan and Muskegon Lake seasonally, but they do not appear to interbreed

with resident populations (57,58). Previous work in this system using a RAD-Seq data set

revealed that yellow perch in the main basin of Lake Michigan are genetically divergent from

Green Bay (59,60), however the demographic and evolutionary histories of yellow perch from

these three highly differentiated habitats have not yet been investigated (57–60).

Yellow perch remain one of the most ecologically and economically important species

throughout the Great Lakes. They are an abundant nearshore fish species, serve as important

predators of small fishes and invertebrates, and are themselves important prey for larger fishes

(see "Yellow perch life history" in Supplementary Materials for relevant life history information)

(61). Historically, yellow perch supported commercially important fisheries throughout the Great

Lakes region; in Lake Michigan alone, peak annual commercial harvest would now represent

close to $16 million (US$) in dock-side value and much more at retail. However, Lake Michigan

yellow perch populations began declining during the late 1980s and early 1990s, which led to

closures of most yellow perch commercial fisheries. Here, we sequenced the complete genomes

and reconstructed the demographic histories for 210 yellow perch collected from seven sample

sites encompassing the main basin of Lake Michigan, Green Bay, and Muskegon Lake (Fig. 1a).

Using genome-wide data, we ask the following questions: 1.) what are the differences in genetic

differentiation and genetic diversity among sample sites?, 2.) what are the differences in Ne

among sample sites using multiple approaches that can estimate both historical and more recent

effective population sizes?, and 3.) what genes show signals of responding to selection in the

three environmentally distinct regions (*i.e.*, main basin, Green Bay, and Muskegon Lake)? We

identify three highly discrete populations, provide evidence for historic and contemporary

increases in Ne, and find signatures of parallel, genetic adaptation.

Methods

Data collection

We sampled 210 adult yellow perch from seven locations (Green Bay, Muskegon Lake, Grand

Haven, Michigan City, Milwaukee, Naubinway, Sutton's Bay; Fig. 1a, Supplementary Table 1)

in 2016, 2018 or 2019 using fyke netting, nighttime boat electrofishing, bottom trawls, gill

netting and creel surveys, and conducted whole-genome resequencing to a depth of $12.50\times$ after

aligning to the yellow perch genome (62) (ranging from $5.93\times$ to $18.58\times$; see "Whole-genome

sequencing" in Supplementary Materials). We called SNPs using Genome Analysis Toolkit

4.1.9.0 (GATK 4.1.9.0) (63) (see "Variant calling and filtering" in Supplementary Materials).

We filtered SNPs by minor allele frequency (MAF) in two ways to create two data sets of SNPs

suitable for different analyses (Supplementary Table 2): a **study-wide** data set and a **within-**

group data set (see "Variant calling and filtering" in Supplementary Materials) (64). Study-wide

MAF filtering is useful for delineating populations, while within-group filtering is essential for

identifying rare variants within each population. Thus, for population structure analyses, we used

the study-wide data set, whereas we used the within-group data sets for all subsequent analyses

(Supplementary Table 2).

Population structure analysis

To infer the phylogenetic relationship among the seven sample sites, we constructed

phylogenetic trees using 5,000 randomly selected SNPs. We first converted SNPs in VCF format

into PHYLIP format and then constructed a phylogenetic tree using RAxML version 8.2.12 (65).

When running RAxML, we selected the GTRGAMMA model, set the random number seed for

parsimony inferences to 12345 and the number of alternative runs on distinct starting trees to

1000, turned on rapid bootstrapping, ran rapid bootstrap analysis, and searched for the best-

scoring ML tree in each run. We viewed the phylogenetic tree using MEGA11 (66). To further

investigate the population structure, we first conducted principal coordinate analysis on all

individuals with 50,000 randomly selected SNPs in *hierfstat* in R 4.2.1 (67). Next, we conducted

principal coordinate analysis with pairwise F_{ST} using GenAlEx 6.51b2 (68). Weir and

Cockerham's F_{ST} was calculated for each pair of sample sites using VCFtools (0.1.16) (69).

Lastly, we estimated the ancestry for Green Bay, Muskegon Lake, and the five main basin

sample sites, separately and jointly, with 10,000 randomly selected SNPs using Admixture 1.3

(70) by setting *K* from 1 to 10.

Minor allele frequency, Tajima's D, and genetic diversity

Using the within-group data sets, we calculated minor allele frequencies, Tajima's D, and genetic

diversity: 1) we calculated minor allele frequencies for seven sample sites and viewed its

distribution by plotting density curves with unit bins of 0.015 with ggplot2 (71) and ggridges in

R 4.2.1 (67), 2) we calculated Tajima's D by setting the bin size to 5 MB with the within-group

data set filtered by MAF $\geq$ 0.02 (Supplementary Table 2) using VCFtools (0.1.16), and 3) we

estimated genetic diversity by calculating observed heterozygosity for every SNP (not including

invariant sites) and averaged observed heterozygosity across overlapping windows of 5 MB

using a step size of 2.5 MB across each chromosome. We also calculated nucleotide diversity (π)

for each population with biallelic sites that are covered by at least three reads in each sample

(including both polymorphic and invariant sites) using PIXY (72).

Effective population size and demographic history

Using the within-group data sets (Supplementary Table 2), we estimated effective population

size with 30,000 randomly selected SNPs (replicated 3 times per population) for each population

using the linkage disequilibrium method, as implemented in NeEstimator V2 (28), by selecting

the random mating model, setting the critical value to 0.05, and picking across chromosomes as

the locus pairing mode to restrict the analysis to pairs of loci on distinct chromosomes. To

further validate the *Ne* estimates, we estimated effective population size with 3,000 randomly

selected SNPs from a previously obtained RAD-Seq data set (59), again replicated 3 times per

sample site for each of 12 sample sites (five from Green Bay, two from Grand Haven, two from

Michigan City, one from Milwaukee, one from Naubinway, one from Sutton's Bay), using the

same options described above. The RAD-Seq data set was only used in this particular analysis

(Supplementary Table 2) and was employed to validate our findings with additional sample sites

and independent extraction, sequencing, and genotyping pipelines. Returning to our whole-

genome data sets, we first reconstructed demographic histories by running GONE (39) with all

SNPs in the within-group data sets 100 times for each sample site. Specifically, we reconstructed

the demographic history for Green Bay and Muskegon Lake by randomly selecting 30,000 SNPs

from the within-group data set for each sample site (replicated 1000 times per sample site;

Supplementary Table 2) and running GONE (39) for each of 1,000 randomly selected sets of

30,000 SNPs for each sample site using the default parameters. To examine changes in *Ne* over

longer time scales, we also ran Stairway Plot 2 (38), for each sample site, with all biallelic SNPs

at which genotype information is available for all samples in each of the seven sample sites,

where generation time was estimated as four years +/- one year (see "Yellow perch life history"

in Supplementary Materials).

Candidate genes and gene ontology (GO) hierarchy networks underlying rapid adaptation

To identify regions of the genome with overlapping windows of high F_{ST}, we first calculated

pairwise Weir and Cockerham's F_{ST} with separate VCF files for ten pairs of populations using

VCFtools (0.1.16) (69) and identified outlier windows from a minimum of eight out of the ten

pairwise population comparisons sharing at least one high F_{ST} SNP. We mapped SNPs on outlier

windows to the yellow perch genome (62) to identify candidate outlier genes potentially

explaining parallel genetic adaptation in Green Bay and Muskegon Lake in comparison to main

basin populations. We annotated candidate outlier genes using UniProtKB ID via UniProt

(https://www.uniprot.org/), with which we retrieved gene ontology (GO) terms for these outlier

genes from Gene Ontology and GO Annotations (https://www.ebi.ac.uk/QuickGO/). Lastly, we

constructed GO hierarchy networks of GO terms associated with at least two outlier genes in the

domain of "biological process" for each pairwise comparison (*i.e.*, Green Bay or Muskegon Lake

vs. main basin populations) using *metacoder* (73).

Water clarity, water quality, and phenotypic measurements

Because many of the outlier genes identified from the previous analysis were related to vision,

we investigated differences in water clarity, water quality, and yellow perch eye size between the

main basin, Green Bay, and Muskegon Lake sample sites. Differences in eye size among those

sample sites was not an *a priori* hypothesis but was rather an *a posteriori* hypothesis guided by

our candidate genes (see Discussion). Water clarity was estimated using Secchi depth readings

collected between June 1 and August 30 from Green Bay (320 readings), Muskegon Lake (72

readings), and the southern basin of Lake Michigan (180 measurements) between 2010 and 2022

(see "Water clarity, water quality, and phenotypic measurements" in Supplementary Materials).

We conducted an Analysis of Variance (ANOVA) to test for differences in Secchi depths among

Green Bay, Muskegon Lake, and the southern basin of Lake Michigan.

To determine whether eye diameter differed among Green Bay, Muskegon Lake, and

Lake Michigan, we collected yellow perch from the three locations, measured their eye diameter,

and analyzed yellow perch with total lengths of 150-250 mm to ensure comparisons of yellow

perch with similar sizes among the three locations (*n=31* for Menominee in Green Bay, *n=39* for

Muskegon Lake, and *n=27* for Grand Haven in Lake Michigan; see "Water clarity, water quality,

and phenotypic measurements" in Supplementary Materials). With this subset of fish, we used

Analysis of Covariance (ANCOVA) to test whether eye diameter differed among the three

locations while accounting for total length as a covariate (including interaction between location

and covariate). We compared least-squares means with Tukey's method using the "emmeans"

function in the *emmeans* package in R (version 1.8.9).

Results

Population structure

We mapped an average of 99.6% of 15.8 billion, 150 bp paired-end reads to the yellow perch

genome (62) and, on average, over 90% and 83% of reads had a mapping quality equal to or

greater than 20 or 60, respectively (Supplementary Table 3). The study-wide data set, which was

used for population genetic and stock identification analyses, resulted in 714,666 SNPs per

sample site while the within-group data set, which was used for demographic analyses, resulted

in an average of 1,160,504 SNPs with less than 0.9% of SNPs out of Hardy-Weinberg

equilibrium in each sample site (Supplementary Table 2, Supplementary Table 4). After inferring

phylogenetic relationships among populations, we found that Green Bay, Muskegon Lake, and

the main basin sample sites (*i.e.*, Grand Haven, Michigan City, Milwaukee, Naubinway, and

Sutton's Bay) form three distinct lineages (Fig. 1b, Supplementary Fig. 1). By conducting

principal coordinate analysis with all individuals, we found further support that Green Bay,

Muskegon Lake, and the main basin sample sites are clearly separated from each other (Fig. 1c,

Supplementary Fig. 2a,b,c), except for one Muskegon Lake sample clustering with main basin

populations and one Milwaukee main basin sample falling comparatively further away from the

main basin populations (Fig. 1c, Supplementary Fig. 2d,e,f; see "Yellow perch life history" and

"Genetic divergence among yellow perch sample sites" in Supplementary Materials). This clear

delineation of three populations was further supported by principal coordinate analysis on the

mean pairwise estimates of F_{ST} where we detected high genetic differentiation between Green

Bay and Muskegon Lake (F_{ST} = 0.0520), both of which are genetically divergent from the five

main basin sample sites (F_{ST} = 0.0550 to 0.0667, 0.0460 to 0.0592, respectively; Supplementary

Table 5, Supplementary Fig. 3a). By contrast, the five main basin sample sites are comparatively

similar to each other, with mean pairwise F_{ST} ranging from -0.0002 to 0.0041 (Supplementary

Table 5, Supplementary Fig. 3b). The optimal number of clusters/populations identified with

admixture analysis (70) for all populations was 3: Green Bay, Muskegon Lake, and the main

basin sample sites (Fig. 1d, Supplementary Fig. 4).

Numbers of SNPs, minor allele frequency, and Tajima's D

When using the within-group data set, we detected 2.8-3.4 times more polymorphic SNPs in

Green Bay and Muskegon Lake sample sites, respectively, in comparison to the main basin

sample sites (2,186,768 and 2,275,935 *vs.* 670,578 to 776,088 SNPs in main basin populations;

Fig. 2a). The median minor allele frequency of SNPs was 0.086 and 0.083 in Green Bay and

Muskegon Lake, respectively, which was 0.46 times lower than that in main basin populations,

which ranged from 0.183 to 0.185 (Fig. 2b). These differences in the number of SNPs and

median minor allele frequencies were not driven by differences in sample sizes (Supplementary

Table 1), sequencing quality (see "Whole-genome sequencing" in Supplementary Materials),

sequencing depth (Supplementary Table 6), read depth of aligned reads (Supplementary Table

7), or missing data (see "Variant calling and filtering" in Supplementary Materials). Using the

within-group data set, we observed consistently positive Tajima's D values in the main basin

populations, varying from 0.332 to 0.428 on average, which is suggestive of slight population

declines in the main basin (Fig. 2c). By contrast, Green Bay and Muskegon Lake populations had

a genome-wide average Tajima's D of -0.632 and -0.603, respectively (Fig. 2c), which is

suggestive of a recent (over evolutionary time scales) demographic expansion.

Effective population size, genetic diversity, and demographic history

In comparison to the main basin sample sites, Green Bay and Muskegon Lake have significantly

lower effective population sizes; the point estimates using whole-genome sequencing data for

Green Bay and Muskegon Lake were 23 and 90, respectively, by NeEstimator V2 and 46 and

159, respectively, by GONE (Fig. 3a; Supplementary Table 8). Effective population size

estimates by NeEstimator V2 using our RAD-Seq data set with larger numbers of samples (59),

but smaller numbers of loci, suggest that southern and middle Green Bay (SGB, MEN) have

smaller effective population sizes than northern Green Bay (BDN, LBD), with high levels of

concurrence between both data sets, suggesting that main basin sample sites have considerably

larger effective population sizes (Fig. 3b). Genetic diversity of main basin sample sites, with a

mean observed heterozygosity ranging from 0.312 to 0.323, was considerably higher than that of

Green Bay and Muskegon Lake populations, with a mean observed heterozygosity of 0.208 and

0.206, respectively (Fig. 3c, Supplementary Fig. 5). It should be noted that we calculated genetic

diversity (*i.e.*, observed heterozygosity) using only polymorphic sites found in each within-group

data set, meaning that the sets of SNPs for calculation differ among populations. Therefore,

despite Green Bay and Muskegon Lake having a greater number of SNPs, the lower observed

heterozygosity is predicted from theory given the much lower average minor allele frequencies.

By contrast, estimates of nucleotide diversity (π), which is calculated using both polymorphic

and invariant sites, were nearly three times higher in Green Bay and Muskegon Lake in

comparison to the Lake Michigan main basin samples (Supplementary Fig. 6). Using GONE, we

reconstructed contemporary demographic histories and found that Green Bay and Muskegon

Lake experienced rapid growth of Ne over the last 10 to 30 generations (Fig. 3d,e,

Supplementary Fig. 7). Stairway Plot 2 (38), based on site frequency spectra, suggests that

increases in Ne in Green Bay and Muskegon Lake also occurred around 2000 years ago and even

longer for main basin populations (Supplementary Fig. 8, Supplementary Fig. 9, Supplementary

Fig. 10).

Candidate genes underlying genetic adaptation

Given the high levels of genetic divergence between Green Bay, Muskegon Lake, and the main

basin sites, we used a genome-wide scan to determine if any regions of the genome are

suggestive of genetic adaptation to the different environments (74,75). We retained a total of 13

outlier windows from nine chromosomes that were shared in at least 80% of comparisons

between Green Bay and the main basin collection sites as well as Muskegon Lake and the main

basin populations (see Methods; Supplementary Table 9). The 13 outlier windows contained an

average of 1,427 SNPS (range: 363-3,203) and an average of 225 high F_{ST} SNPs ($F_{ST} \geq 0.6$;

range 82-360 SNPs per window) (Fig. 4a). SNPs within these 13 outlier windows mapped to 36

genes, among which three were located within a single outlier window on chromosome 4

(chromosome 4: 34762415-34852528). The three genes, *opn1sw*, *opn1lw*, and *opn1lw2*, encode

visual pigments, light-absorbing molecules mediating vision. Out of the three visual pigment-

encoding genes, *opn1lw2* has two outlier SNPs with $F_{ST} \geq 0.5$ in comparisons between Green

Bay and the main basin sample sites, and the exact same two outlier SNPs, one with $F_{ST} \geq 0.76$

and the other with $F_{ST} \geq 0.46$, were found in comparisons between Muskegon Lake and the main

basin sample sites (Fig. 4a). The observation that three visual pigment-encoding genes were

found on the same outlier window in all ten pairwise comparisons suggests that this region has

responded to recent positive and parallel selection. Across all outlier windows, seven major

categories of biological processes were detected between Green Bay and the main basin sample

sites and Muskegon Lake and the main basin sample sites: multicellular organismal process,

response to stimulus, biological regulation, cellular process, developmental process, localization,

and metabolic process (Fig. 4b). Multiple visual perception-related biological processes,

including detection of visible light, absorption of visible light, visual perception, cellular

response to light stimulus, and phototransduction, provide further support for parallel adaptive

differences in vision in Green Bay and Muskegon Lake yellow perch populations (Fig. 4b).

Using these candidate loci, we hypothesized that water clarity and eye size would differ between

the two sites (Green Bay and Muskegon Lake) and the main basin. We found significant

differences in water clarity between the main basin of Lake Michigan (low turbidity, high visual

clarity; mean Secchi depth $\pm$ s.d.: 6.10 $\pm$ 3.23 m) and both Green Bay (high turbidity, low visual

clarity; 1.59 $\pm$ 1.01 m) and Muskegon Lake (high turbidity, low visual clarity; 2.48 $\pm$ 1.23 m;

ANOVA, $P < 0.001$; Fig. 4c). We also found that yellow perch eye diameter differed among

locations (ANCOVA, $F_{2,91} = 12.03$; $P < 0.001$) and significantly increased with total length

(ANCOVA, $F_{1,91} = 45.67$; $P < 0.001$), but the interaction was not significant (ANCOVA, $F_{2,91} =$

0.11; $P = 0.779$; Supplementary Fig. 11). Yellow perch eye diameter was significantly larger for

Muskegon Lake yellow perch in comparison to fish collected from the main basin (Grand Haven;

$P < 0.001$) and Green Bay (near Menominee; $P < 0.001$). Eye diameter for yellow perch from

Green Bay was also significantly larger than fish from the main basin (Grand Haven; $P < 0.001$;

Fig. 4d).

Discussion

Using whole-genome sequence data from 210 yellow perch, we observed a clear delineation of

three populations and multiple genomic signatures associated with historic and contemporary

increases in Ne in Green Bay and Muskegon Lake where: (1) effective population sizes rapidly

increased over recent generations, (2) Tajima's D estimates were negative across the entire

genome, and (3) a factor of 2.8-3.4 more polymorphic SNPs were found in Green Bay and

Muskegon Lake than the main basin collection sites, where the majority of these additional SNPs

occur at low frequency. The additional SNPs likely come from an excess of mutations that

accumulated during population expansions that occurred over historic time scales (*i.e.*, thousands

of years). This last observation is consistent with a large population expansion over historic time

scales (*e.g.*, over thousands of years), as indicated by Stairway Plot 2 (Supplementary Fig. 9),

where there has not been sufficient time for purifying selection to remove deleterious alleles

(76). Given the substantial water quality and habitat improvements in Green Bay and Muskegon

Lake driven by environmental remediation initiatives implemented in the early 1970s

(Supplementary Fig. 12, Supplementary Fig. 13; see "Supplementary Discussion" in

Supplementary Materials), we should not have been surprised to see an increase in Ne.

Nevertheless, empirical documentation of contemporary increases in Ne remains rare and this

result was unexpected.

To what time period do our demographic inferences apply? Linkage disequilibrium based

methods of Ne, such as those depicted in Fig. 3, provide recent and contemporary estimates of

effective population size (77). We used the program GONE to estimate recent changes to Ne

over time and the results suggest that there have been contemporary increases in *Ne* in both

Green Bay and Muskegon Lake. However, the exact date for when these increases in *Ne* began

remains imprecise. A recent study with pink salmon (*Oncorhynchus gorbuscha*) and a known

date of an extreme founder event illustrated that GONE may have missed the timing of the event

by as much as six generations (45). Nevertheless, the known abundance patterns in salmon and

other species are well characterized with this approach (45,78) and our results clearly illustrate

that effective population sizes have been increasing in both Muskegon Lake and Green Bay in

the last 10 to 30 generations, with larger increases occurring in Muskegon Lake. Our inferences

based on Tajima's D, the number of segregating loci, and site frequency spectra suggest much

larger increases in *Ne* that occurred over historic time scales (centuries, not decades) (Fig. 2;

Supplementary Fig. 9). This result contrasts with those from main basin sites, which all show

increases in *Ne* starting more than 3000 years ago and no signs of a historic bottleneck

(Supplementary Fig. 9). We speculate that both the Green Bay and the Muskegon Lake yellow

perch populations were founded much more recently than the main basin populations or that both

went through a large, unknown, bottleneck event approximately 2,000 years ago.

We also found evidence of parallel evolution in response to reduced water clarity in both

Green Bay and Muskegon Lake yellow perch. In comparison to the main basin of Lake

Michigan, both Green Bay and Muskegon Lake exhibit high turbidity and low visual clarity, as

indicated by shallower Secchi depths. Such differences in water clarity and ambient light

conditions could have driven the significant allele frequency shifts in the coding sequences of

photopigment encoding genes and triggered a set of biological processes contributing to light

detection and absorption, visual perception, and phototransduction, a result that implies parallel

genetic adaptation in Green Bay and Muskegon Lake yellow perch in response to local

environmental conditions. Differences in vision and eye size, while accounting for allometry,

among those sample sites was not an *a priori* hypothesis, although it could have been had we

thought about the differences in water clarity among the sample sites. Nevertheless, using whole-

genome sequencing to guide phenotypic and evolutionary predictions does suggest a promising

avenue for the future of genomic tools (79). Future studies should investigate linkages between

candidate genes and their patterns of expression in addition to the structural and functional

analyses of the proteins they encode. Additional phenotyping efforts and comparisons among

multiple drowned river mouth lakes would also be useful. The observation that both genetic and

phenotypic changes to reduced water clarity occurred in genetically divergent populations

suggests that a match between vision-related traits and environmental characteristics is an

important component of fitness in yellow perch populations (80); environmental changes that

cause a mismatch may result in declines in abundance.

Several caveats are worth discussing. First, linkage disequilibrium-based estimates of

effective population sizes may not accurately estimate effective population sizes prior to

bottleneck events (45,81). As such, the effective population sizes of yellow perch in Green Bay

and Muskegon Lake prior to around 30 generations ago were likely much larger than present day

estimates but cannot be estimated using those approaches. Also, contemporary effective

population size may be underestimated by linkage disequilibrium-based methods due to

overlapping generations of yellow perch and the mixed-age adult samples under study (82).

Demographic estimates by GONE may also be unreliable for the most recent several generations.

However, despite the wide range of estimates for recent generations, the general trend still

indicates an increase in effective population size that appears to have persisted in Green Bay and

Muskegon Lake for the past 10 to 30 generations. Perhaps most importantly, some of our

demographic results could also be explained by gene flow from divergent populations (83,84);

however, we do not think this to be the case for three reasons. First, PCA, phylogenetic, and

Admixture analyses support a clear separation of Green Bay, Muskegon Lake, and Lake

Michigan main basin populations and suggest the absence of admixed individuals in both Green

Bay and Muskegon Lake. Even if all the main basin loci were introduced into Muskegon Lake

samples via gene flow (which is unlikely), this mechanism could only account for 24% of the

2,275,935 SNPs found in Muskegon Lake (*i.e.*, 535,860 SNPs shared by Muskegon Lake and

Grand Haven divided by 2,275,935 Muskegon Lake SNPs; Fig. 2a). Second, over shorter time

scales, estimates of contemporary admixture within Green Bay and Muskegon Lake samples

suggest a single, randomly mating population (as do the small number of loci out of Hardy-

Weinberg equilibrium) (Supplementary Fig. 4; Supplementary Table 4). Third, over longer time

scales, the high rates of introgression required to explain these results would also be likely to

swamp any signal of parallel genetic adaptation (Fig. 4) and would also leave more time for

purifying selection to remove deleterious loci. Note that other factors that can increase *Ne*

through time (*e.g.*, decreased variance in reproductive success) are unlikely to be accompanied

by the other genomic patterns we report (*i.e.*, negative Tajima's D, larger numbers of SNPs).

Thus, we think the most parsimonious explanation for our results may be large, simultaneous

increases in population sizes in Green Bay and Muskegon Lake yellow perch. Interestingly,

fisheries independent data suggest that yellow perch have been declining recently in the main

basin (85) and the positive, albeit small, genome-wide estimates of Tajima's D suggest that these

populations should be closely monitored in the coming decades. While we are aware that

correlation does not equal causation, we speculate that the recent increases in *Ne* in Green Bay

and Muskegon Lake could be a result of substantial water quality and habitat improvements

429 driven by a set of environmental remediation initiatives implemented in the early 1970s, as

430 evidenced by the strong relationship between decreases in nutrient loads and concomitant

431 increases in *Ne* (86,87) (Supplementary Fig. 12, Supplementary Fig. 13; see "Supplementary

432 Discussion" in Supplementary Materials).

433  In conclusion, we found that two populations of an economically important fish species

434 have had large, simultaneous increases in effective population sizes both historically and over

435 contemporary time scales. The two populations are separated by over 350 km and are genetically

436 divergent from one another, such that they represent two distinct evolutionary lineages (Fig. 1b).

437 We also found evidence of parallel adaptation to reduced visual clarity in each of the habitats,

438 which may also have been further magnified by centuries-long anthropogenic disturbances and

439 pollution, though we suspect that this genetic adaptation occurred over a longer time scale. The

440 contemporary estimates of *Ne* are orders of magnitude lower than historic estimates, suggesting

441 that recent anthropogenic disturbances may have reduced genetic diversity in Green Bay and

442 Muskegon Lake. While the recent increases in *Ne* suggest that the largest threats to these

443 populations may have passed, the overall point estimates remain low. We suggest that

444 conservation and management efforts targeting yellow perch in these regions should strive to

445 maintain genetic diversity and obtain accurate estimates of population abundances through time.

446 Taken together, these results suggest that the synchrony of key ecological and evolutionary

447 processes can drive parallel demographic and evolutionary trajectories across independent

448 populations. How frequently such synchrony occurs remains an open question in the fields of

449 ecology and evolution.

450 **References**

451 1. Wright S. Evolution in Mendelian Populations. Genetics. 1931 Mar;16(2):97–159.

2.   Hedgecock D, Pudovkin A. Sweepstakes Reproductive Success in Highly Fecund Marine
Fish and Shellfish: A Review and Commentary. Bulletin of Marine Science. 2011 Oct
1;87:971–1002.

3.   Hedgecock D. Does variance in reproductive success limit effective population sizes of
marine organisms? Genetics and Evolution of Aquatic Organisms. 1994 Jan 1;

4.   Hauser L, Adcock GJ, Smith PJ, Ramiréz JHB, Carvalho GR. Loss of microsatellite diversity
and low effective population size in an overexploited population of New Zealand snapper
(*Pagrus auratus*). Proceedings of the National Academy of Sciences of the United States of
America. 2002 Sep 3;99(18):11742–7.

5.   Hoarau G, Boon E, Jongma DN, Ferber S, Palsson J, Van der Veer HW, et al. Low effective
population size and evidence for inbreeding in an overexploited flatfish, plaice (*Pleuronectes*
platessa L.). Proc Biol Sci. 2005 Mar 7;272(1562):497–503.

6.   Miller LM, Kapuscinski AR. Historical analysis of genetic variation reveals low effective
population size in a northern pike (*Esox Lucius*) population. Genetics. 1997
Nov;147(3):1249–58.

7.   Turner TF, Wares JP, Gold JR. Genetic effective size is three orders of magnitude smaller
than adult census size in an abundant, Estuarine-dependent marine fish (*Sciaenops*
ocellatus). Genetics. 2002 Nov;162(3):1329–39.

8.   Hutchinson WF, van Oosterhout C, Rogers SI, Carvalho GR. Temporal analysis of archived
samples indicates marked genetic changes in declining North Sea cod (*Gadus morhua*). Proc
Biol Sci. 2003 Oct 22;270(1529):2125–32.

9.   Waples RS. Tiny estimates of the Ne/N ratio in marine fishes: Are they real? Journal of Fish
Biology. 2016;89(6):2479–504.

10. Waples RS, Grewe PM, Bravington MW, Hillary R, Feutry P. Robust estimates of a high
Ne/N ratio in a top marine predator, southern bluefin tuna. Science Advances. 2018 Jul
18;4(7):eaar7759.

11. Palstra FP, Fraser DJ. Effective/census population size ratio estimation: a compendium and
appraisal. Ecology and Evolution. 2012;2(9):2357–65.

12. Ruzzante DE, McCracken GR, Parmelee S, Hill K, Corrigan A, MacMillan J, et al. Effective
number of breeders, effective population size and their relationship with census size in an
iteroparous species, Salvelinus fontinalis. Proc Biol Sci. 2016 Jan 27;283(1823):20152601.

13. Clarke SH, Lawrence ER, Matte JM, Gallagher BK, Salisbury SJ, Michaelides SN, et al.
Global assessment of effective population sizes: Consistent taxonomic differences in meeting
the 50/500 rule. Mol Ecol. 2024 Jun;33(11):e17353.

14. Hahn M. Molecular Population Genetics. Oxford, New York: Oxford University Press; 2018.
352 p.

15. Waples RS. What Is *Ne*, Anyway? Murphy W, editor. Journal of Heredity. 2022 Jul
23;113(4):371–9.

16. Capblancq T, Butnor JR, Deyoung S, Thibault E, Munson H, Nelson DM, et al. Whole-
exome sequencing reveals a long-term decline in effective population size of red spruce
(*Picea rubens*). Evol Appl. 2020 May 22;13(9):2190–205.

17. Sharma R, Arora N, Goossens B, Nater A, Morf N, Salmona J, et al. Effective population
size dynamics and the demographic collapse of Bornean orang-utans. PLOS ONE. 2012 Nov
15;7(11):e49429.

18. Okello JBA, Wittemyer G, Rasmussen HB, Arctander P, Nyakaana S, Douglas-Hamilton I,
et al. Effective population size dynamics reveal impacts of historic climatic events and recent
anthropogenic pressure in African elephants. Mol Ecol. 2008 Sep;17(17):3788–99.

19. Ólafsdóttir GÁ, Westfall KM, Edvardsson R, Pálsson S. Historical DNA reveals the
demographic history of Atlantic cod (*Gadus morhua*) in medieval and early modern Iceland.
Proceedings of the Royal Society B: Biological Sciences. 2014 Feb 22;281(1777):20132976.

20. Lonsinger RC, Adams JR, Waits LP. Evaluating effective population size and genetic
diversity of a declining kit fox population using contemporary and historical specimens. Ecol
Evol. 2018 Nov 8;8(23):12011–21.

21. Abadía-Cardoso A, Freimer NB, Deiner K, Garza JC. Molecular population genetics of the
northern elephant seal *Mirounga angustirostris*. J Hered. 2017 Sep;108(6):618–27.

22. Pita A, Pérez M, Velasco F, Presa P. Trends of the genetic effective population size in the
Southern stock of the European hake. Fisheries Research. 2017 Jul 1;191:108–19.

23. Ovenden JR, Leigh GM, Blower DC, Jones AT, Moore A, Bustamante C, et al. Can
estimates of genetic effective population size contribute to fisheries stock assessments?
Journal of Fish Biology. 2016;89(6):2505–18.

24. Hare MP, Nunney L, Schwartz MK, Ruzzante DE, Burford M, Waples RS, et al.
Understanding and estimating effective population size for practical application in marine
species management. Conserv Biol. 2011 Jun;25(3):438–49.

25. Shrimpton JM, Heath DD. Census vs. effective population size in chinook salmon: large- and
small-scale environmental perturbation effects. Mol Ecol. 2003 Oct;12(10):2571–83.

26. Horreo J, Machado-Schiaffino G, Griffiths A, Bright D, Stevens J, Garcia-Vazquez E.
Atlantic salmon at risk: Apparent rapid declines in effective population size in southern
European populations. Transactions of the American Fisheries Society. 2011 May
1;140:605–10.

27. Kuparinen A, Hutchings JA, Waples RS. Harvest-induced evolution and effective population
size. Evolutionary Applications. 2016;9(5):658–72.

28. Do C, Waples RS, Peel D, Macbeth GM, Tillett BJ, Ovenden JR. NeEstimator v2: re-
implementation of software for the estimation of contemporary effective population size (*Ne*)
from genetic data. Molecular Ecology Resources. 2014;14(1):209–14.

29. Nomura T. Estimation of effective number of breeders from molecular coancestry of single
cohort sample. Evol Appl. 2008 Aug;1(3):462–74.

30. Cowen R, Gawarkiewicz G, Pineda J, Thorrold S, Werner F. Population connectivity in
marine systems: An overview. Oceanography. 2007 Sep 1;20.

31. Cowen RK, Sponaugle S. Larval dispersal and marine population connectivity. Annual
Review of Marine Science. 2009;1(1):443–66.

32. Waples R. Separating the wheat from the chaff: patterns of genetic differentiation in high
gene flow species. Journal of Heredity. 1998 Sep 1;89(5):438–50.

33. Christie MR, Tissot BN, Albins MA, Beets JP, Jia Y, Ortiz DM, et al. Larval connectivity in
an effective network of marine protected areas. PLOS ONE. 2010 Dec 21;5(12):e15715.

34. Waples RS. Practical application of the linkage disequilibrium method for estimating
contemporary effective population size: A review. Molecular Ecology Resources.
2024;24(1):e13879.

35. Palstra FP, Ruzzante DE. Genetic estimates of contemporary effective population size: What
can they tell us about the importance of genetic stochasticity for wild population persistence?
Mol Ecol. 2008 Aug;17(15):3428–47.

36. Jorde PE, Ryman N. Temporal allele frequency change and estimation of effective size in
populations with overlapping generations. Genetics. 1995 Feb;139(2):1077–90.

37. Liu X, Fu YX. Exploring population size changes using SNP frequency spectra. Nat Genet.
2015 May;47(5):555–9.

38. Liu X, Fu YX. Stairway Plot 2: demographic history inference with folded SNP frequency
spectra. Genome Biology. 2020 Nov 17;21(1):280.

39. Santiago E, Novo I, Pardiñas AF, Saura M, Wang J, Caballero A. Recent demographic
history inferred by high-resolution analysis of linkage disequilibrium. Molecular Biology
and Evolution. 2020 Dec 16;37(12):3642–53.

40. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the joint
demographic history of multiple populations from multidimensional SNP frequency data.
PLoS Genet. 2009 Oct;5(10):e1000695.

41. Li H, Durbin R. Inference of human population history from individual whole-genome
sequences. Nature. 2011 Jul;475(7357):493–6.

42. Laurent V, Planes S. Effective population size estimation on *Sardina pilchardus* in the Bay
of Biscay using a temporal genetic approach. Biological Journal of the Linnean Society.
2007 Apr 1;90(4):591–602.

43. Westbury MV, Hartmann S, Barlow A, Wiesel I, Leo V, Welch R, et al. Extended and
continuous decline in effective population size results in low genomic diversity in the
world's rarest hyena species, the brown hyena. Shapiro B, editor. Molecular Biology and
Evolution. 2018 May 1;35(5):1225–37.

44. Diaz M, Wethey D, Bulak J, Ely B. Effect of harvest and effective population size on genetic
diversity in a striped bass population. Transactions of the American Fisheries Society. 2000
Nov 1;129(6):1367–72.

45. Sparks MM, Schraidt CE, Yin X, Seeb LW, Christie MR. Rapid genetic adaptation to a
novel ecosystem despite a large founder event. Mol Ecol. 2023 Sep 5;

46. Hoey JA, Able KW, Pinsky ML. Genetic decline and recovery of a demographically rebuilt
fishery species. Molecular Ecology. 2022;31(22):5684–98.

47. Wang J, Santiago E, Caballero A. Prediction and estimation of effective population size.
Heredity. 2016 Oct;117(4):193–206.

48. Hedrick P. Large variance in reproductive success and the Ne/N ratio. Evolution.
2005;59(7):1596–9.

49. Allendorf FW. Delay of adaptation to captive breeding by equalizing family size.
Conservation Biology. 1993;7(2):416–9.

50. Borlase S, Loebel D, Frankham R, Nurthen R, Briscoe D, Daggard GE. Modeling problems
in conservation genetics using captive *Drosophila* populations: consequences of equalization
of family sizes. Conservation Biology. 2002 Feb 23;7:122–31.

51. Waples RS, Hindar K, Karlsson S, Hard JJ. Evaluating the Ryman–Laikre effect for marine
stock enhancement and aquaculture. Curr Zool. 2016 Dec;62(6):617–27.

52. Ryman N, Laikre L. Effects of supportive breeding on the genetically effective population
size. Conservation Biology. 1991;5(3):325–9.

53. Great Lakes Commission [Internet]. [cited 2024 Sep 22]. Lake Michigan. Available from:
https://www.glc.org/lakes/lake-michigan/

54. NOAA National Ecosystem Status [Internet]. [cited 2024 Sep 22]. Lake Michigan. Available
from: https://ecowatch.noaa.gov/index.php/regions/great-lakes/lake-michigan

55. Herdendorf CE. Great Lakes estuaries. Estuaries. 1990 Dec 1;13(4):493–503.

56. Mader MM, Ruetz CR, Woznicki SA, Steinman AD. Land cover and water quality of
drowned river mouths: Evidence of an environmental gradient along the eastern Lake
Michigan shoreline. Journal of Great Lakes Research. 2023 Dec 1;49(6):102237.

57. Senegal TJ, Ruetz CR, Chorak GM, Janetski DJ, Clapp DF, Bowen GJ, et al. Differential
habitat use patterns of yellow perch *Perca flavescens* in eastern Lake Michigan and
connected drowned river mouth lakes. Journal of Great Lakes Research. 2020 Oct
1;46(5):1412–22.

58. Chorak GM, Ruetz III CR, Thum RA, Partridge CG, Janetski DJ, Höök TO, et al. Yellow
perch genetic structure and habitat use among connected habitats in eastern Lake Michigan.
Ecology and Evolution. 2019;9(16):8922–32.

59. Schraidt CE, Ackiss AS, Larson WA, Rowe MD, Höök TO, Christie MR. Dispersive
currents explain patterns of population connectivity in an ecologically and economically
important fish. Evolutionary Applications. 2023;16(7):1284–301.

60. Miller LM. Microsatellite DNA loci reveal genetic structure of yellow perch in Lake
Michigan. Transactions of the American Fisheries Society. 2003;132(3):503–13.

61. Evans MS. Recent Major Declines in Zooplankton Populations in the Inshore Region of
Lake Michigan: Probable Causes and Implications. Can J Fish Aquat Sci. 1986
Jan;43(1):154–9.

62. Feron R, Zahm M, Cabau C, Klopp C, Roques C, Bouchez O, et al. Characterization of a Y-
specific duplication/insertion of the anti-Mullerian hormone type II receptor gene based on a
chromosome-scale genome assembly of yellow perch, *Perca flavescens*. Molecular Ecology
Resources. 2020;20(2):531–43.

63. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res [Internet]. 2010 Sep 1 [cited 2020 Jan 17];20(9):1297–303.
Available from: http://genome.cshlp.org/content/20/9/1297

64. Hemstrom W, Grummer JA, Luikart G, Christie MR. Next-generation data filtering in the
genomics era. Nat Rev Genet. 2024 Jun 14;1–18.

65. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 2014 May 1;30(9):1312–3.

66. Tamura K, Stecher G, Kumar S. MEGA11: Molecular evolutionary genetics analysis version
11. Molecular Biology and Evolution. 2021 Jul 1;38(7):3022–7.

67. R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/. 2020;

68. Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software
for teaching and research—an update. Bioinformatics. 2012 Oct 1;28(19):2537–9.

69. Weir BS, Cockerham CC. Estimating F-statistics for the analysis of population structure.
Evolution. 1984;38(6):1358–70.

70. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated
individuals. Genome Res. 2009 Sep;19(9):1655–64.

71. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 1st ed. 2009. Corr. 3rd printing
2010 edition. New York: Springer; 2010. 213 p.

72. Korunes KL, Samuk K. pixy: Unbiased estimation of nucleotide diversity and divergence in
the presence of missing data. Molecular Ecology Resources. 2021;21(4):1359–68.

73. Foster ZSL, Sharpton TJ, Grünwald NJ. Metacoder: An R package for visualization and
manipulation of community taxonomic diversity data. PLOS Computational Biology. 2017
Feb 21;13(2):e1005404.

74. Willoughby JR, Harder AM, Tennessen JA, Scribner KT, Christie MR. Rapid genetic
adaptation to a novel environment despite a genome-wide reduction in genetic diversity. Mol
Ecol. 2018;27(20):4041–51.

75. Bernatchez L, Ferchaud AL, Berger CS, Venney CJ, Xuereb A. Genomics for monitoring
and understanding species responses to global climate change. Nat Rev Genet. 2024
Mar;25(3):165–83.

76. Marth GT, Czabarka E, Murvai J, Sherry ST. The allele frequency spectrum in genome-wide
human variation data reveals signals of differential demographic history in three large world
populations. Genetics. 2004 Jan;166(1):351–72.

77. Waples RS, Do C. Linkage disequilibrium estimates of contemporary Ne using highly
variable genetic markers: a largely untapped resource for applied conservation and evolution.
Evol Appl. 2010 May;3(3):244–62.

78. Kardos M, Zhang Y, Parsons KM, A Y, Kang H, Xu X, et al. Inbreeding depression explains
killer whale population dynamics. Nat Ecol Evol. 2023 May;7(5):675–86.

79. Garner BA, Hand BK, Amish SJ, Bernatchez L, Foster JT, Miller KM, et al. Genomics in
Conservation: Case Studies and Bridging the Gap between Data and Application. Trends
Ecol Evol. 2016 Feb;31(2):81–3.

80. Andersson M, Scharnweber K, Eklöv P. Environmental and ecological drivers of eye size
variation in a freshwater predator: A trade-off between foraging and predation risk.
Functional Ecology. 2024 Sep 6;

81. Reid BN, Pinsky ML. Simulation-based evaluation of methods, data types, and temporal
sampling schemes for detecting recent population declines. Integr Comp Biol. 2022 Dec
30;62(6):1849–63.

82. Waples RS, Antao T, Luikart G. Effects of overlapping generations on linkage
disequilibrium estimates of effective population size. Genetics. 2014 Jun;197(2):769–80.

83. Novo I, Ordás P, Moraga N, Santiago E, Quesada H, Caballero A. Impact of population
structure in the estimation of recent historical effective population size by the software
GONE. Genet Sel Evol. 2023 Dec 4;55(1):86.

84. Paz-Vinas I, Quéméré E, Chikhi L, Loot G, Blanchet S. The demographic history of
populations experiencing asymmetric gene flow: combining simulated and empirical data.
Mol Ecol. 2013 Jun;22(12):3279–91.

85. Clapp D, Makauskas D. Status of yellow perch in Lake Michigan - 2019-2020. Report to the
Lake Michigan committee. 2021;

86. Harris HJ, Wenger RB, Sager PE, Val Klump J. The Green Bay saga: Environmental change,
scientific investigation, and watershed management. Journal of Great Lakes Research. 2018
Oct 1;44(5):829–36.

87. Liu B, McLean CE, Long DT, Steinman AD, Stevenson RJ. Eutrophication and recovery of
a lake inferred from sedimentary diatoms originating from different habitats. Science of The
Total Environment. 2018 Jul 1;628–629:1352–61.

Data availability

All raw data generated for this project are stored in the NCBI Short Read Archive (SRA)

PRJNA1037191 Great Lakes yellow perch whole genome sequencing. Additionally, unfiltered

and filtered VCF files (as described in Supplementary Table 2) are also available

(https://doi.org/10.6084/m9.figshare.24517471.v1). The scripts used in this project are hosted in

a public repository (https://github.com/ChristieLab/yellow_perch_demography_adaptation).

Acknowledgements

We thank the Michigan Department of Natural Resources, the Indiana Department of Natural

Resources, the Wisconsin Department of Natural Resources, the Illinois Natural History Survey,

and the Grand Traverse Band of Ottawa and Chippewa Indians for assistance with sample

collection. In particular, we thank Charles Roswell, Troy Zorn, Tammie Paoli, Dave Clapp, Pat

O'Neil, Dave Fielder, David Schindelholz, Erik Olsen, Jayson Beugly, Josey Cline, Taylor

Senegal, Greg Chorak, and Scott Koenigbauer. Field collections of yellow perch were authorized

by issuance of state-specific collector's permits and approved by Purdue University's Animal

Care and Use Committee protocol 1112000400. We thank Kurt Thompson for assistance with

the map of Muskegon Lake. Secchi depth data were collected and provided by Sarah Bartlett and

the NEW Water monitoring program in Green Bay and Sarah Stein and Les Warren of Purdue

University. This work was funded by the Great Lakes Fishery Commission (Project ID:

2018_CHR_44072) with additional support from the Great Lakes Fishery Trust (#1456 and

#1952) to sample Muskegon Lake.

Author contributions

MRC, XY, and TOH designed the project, CES and CRR collected samples with agency

assistance, XY performed molecular work, XY, MMS, PTE, TJH, CRR, and MRC analyzed the

data, and XY and MRC wrote the manuscript with input from all authors. All authors read and

approved the final manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary materials are available at XXXXXX.

Figure Legends

Fig. 1. Population structure of Lake Michigan yellow perch populations. Yellow perch collected from seven Lake Michigan sites (a), including the comparatively isolated Muskegon Lake, which is connected to the main basin by a narrow channel (see inset). A phylogenetic tree illustrates three distinct groups: Green Bay, Muskegon Lake, and the five main basin sample sites (b). Twenty-five main basin individuals were randomly sampled for visual clarity; see Supplementary Fig. 1 for the full phylogenetic tree with all 210 individuals. Both principal coordinate analysis of 210 individuals (c) and admixture analyses (d) further support the delineation of samples into three discrete populations. Notice that one individual sampled in Muskegon Lake likely swam in from the main basin (Supplementary Fig. 2; see "Genetic divergence among yellow perch sample sites" in Supplementary Materials).

Fig. 2. Numbers of single nucleotide polymorphisms (SNPs), minor allele frequency, and Tajima's D. Green Bay and Muskegon Lake populations have 2.8-3.4 times more SNPs than the five main basin sample sites (a), despite all sample sites having equal sample sizes, sequencing quality, read depth, and missing data. Both Green Bay and Muskegon Lake also have a lower median minor allele frequency compared to the five main basin populations (b) here illustrated with a density plot where the median minor allele frequency is depicted with a solid vertical line. The negative Tajima's D for both Green Bay and Muskegon Lake populations suggests recent population expansion (c). Collectively, these results suggest that the three populations (Green Bay, Muskegon Lake, and the main basin sample sites) have different demographic histories.

Fig. 3. Demographic history of Lake Michigan yellow perch populations. Green Bay and Muskegon Lake sample sites have substantially lower effective population sizes, which average 23 and 90 estimated by NeEstimator V2, respectively, in comparison to main basin populations (a, estimates based on whole-genome sequencing data obtained in this study; b, estimates based on RAD-Seq data obtained from Schraidt *et al.*; note that y-axis is on a base 10 logarithmic scale; SGB: South Green Bay, MUS: Muskegon Lake, GRH: Grand Haven, MIC: Michigan City, MIL: Milwaukee, NUB: Naubinway, SUT: Sutton's Bay, MEN: Menominee, BDN: Big Bay de Noc, LBD: Little Bay de Noc). Green Bay and Muskegon Lake have lower genetic diversity (c) than main basin populations, measured here as observed heterozygosity averaged across overlapping windows of 5 MB using a step size of 2.5 MB across each chromosome. Notice that the main basin heterozygosity estimates are highly correlated, confirming higher gene flow among main basin sample sites. The demographic history, as reconstructed by GONE, demonstrates a rapid increase in effective population size in Green Bay and Muskegon Lake since 10 to 30 generations ago (d,e).

Fig. 4. Candidate genes, biological processes, and phenotypes involved in parallel genetic adaptation. A total of 13 outlier windows were detected in comparisons between both Green Bay and the main basin sample sites and Muskegon Lake and the main basin sample sites (a). The outlier window on chromosome 4 contains visual pigment encoding genes, such as *opn1lw2*. The coding regions, as indicated by horizontal gray bars, of these outlier genes contain SNPs that are highly differentiated (a). Seven major categories of biological processes are potentially involved in adaptive differentiation in Green Bay and Muskegon Lake in comparison to main

basin sample sites, among which visual perception-related processes (highlighted with red
asterisks) further support parallel adaptive differences in vision in Green Bay and Muskegon
Lake (b). The genetic differentiation at *opn1lw2* is likely associated with water clarity, as
evidenced by significant differences in Secchi depths (c) and different eye sizes (d) between
Green Bay or Muskegon Lake and main basin populations (*i.e.*, Lake Michigan, Grand Haven).
The Secchi depth in Green Bay and Muskegon Lake is significantly smaller than that in the
southern basin of Lake Michigan ($P < 0.001$; c) and least-squares means of yellow perch eye
diameter in Green Bay and Muskegon Lake is significantly larger than that in Grand Haven ($P <$
0.001; d). We accounted for differences in total length when comparing eye diameter of yellow
perch collected from Green Bay, Muskegon Lake, and Grand Haven (Supplementary Fig. 11; see
Methods).

Figure 1

Figure 2

Figure 3

Figure 4

(a)

(b)

(c)

(d)