Accelerating antimalarial drug discovery with a new high-throughput screen for fast-killing

compounds

Takaya Sakura[1,2,†], Ryuta Ishii[3,4,†], Eri Yoshida[1], Kiyoshi Kita[2,5,6], Teruhisa Kato[4,7], Daniel Ken

Inaoka[1,2,5,6,*]

[1]Department of Molecular Infection Dynamics, Institute of Tropical Medicine (NEKKEN),

Nagasaki University, Nagasaki 852-8523, Japan. [2]School of Tropical Medicine and Global Health,

Nagasaki University, Sakamoto, Nagasaki 852-8523, Japan. [3]Department of Cellular Architecture

Studies, Institute of Tropical Medicine (NEKKEN), Nagasaki University, Nagasaki 852-8523,

Japan. [4]Laboratory for Drug Discovery and Disease Research, Shionogi & Co., Ltd., Osaka, 561-

0825, Japan. [5]Department of Infection Biochemistry, Institute of Tropical Medicine (NEKKEN),

Nagasaki University, Nagasaki 852-8523, Japan. [6]Department of Biomedical Chemistry, Graduate

School of Medicine, The University of Tokyo, Tokyo 113-0033, Japan. [7]Exploratory Research for

Drug Discovery, Institute of Tropical Medicine (NEKKEN), Nagasaki University, Nagasaki 852-

8523, Japan.

[†] These authors contributed equally to this work.

*Correspondence: Daniel Ken Inaoka, danielken@nagasaki-u.ac.jp. Department of Infection

Biochemistry, Institute of Tropical Medicine (NEKKEN), Nagasaki University, 1-12-4, Sakamoto,

Nagasaki 852-8523, Japan.

Abstract

The urgent need for rapidly acting compounds in the development of antimalarial drugs underscores the significance of such compounds in overcoming resistance issues and improving patient adherence to antimalarial treatments. The present study introduces a high-throughput screening (HTS) approach using 1536-well plates, employing *Plasmodium falciparum* lactate dehydrogenase (PfLDH) combined with nitroreductase (NTR) and fluorescent probes to evaluate inhibition of the growth of the asexual blood stage of malaria parasites. This method was adapted to efficiently measure the parasite reduction ratio (PRR) in a 384-well plate format, streamlining the traditionally time-consuming screening process. By successfully screening numerous compounds, this approach identified fast-killing hits early in the screening process, addressing challenges associated with artemisinin-based combination therapies. The high-throughput PRR method is expected to be of value in continuously monitoring fast-killing properties during structure-activity relationship studies, expediting the identification and development of novel, rapidly acting antimalarial drugs within phenotypic drug discovery campaigns.

Keywords: malaria, drug discovery, high-throughput screening, lactate dehydrogenase, nitroreductase, fast-killing,

Introduction

Malaria is a parasitic disease caused by *Plasmodium* spp. infection, and typically is seen in tropical and subtropical areas. In 2021, over 420,000 deaths, predominantly among children under 5 years old, were reported worldwide, along with more than 200 million cases[1]. Artemisinin-based combination therapies (ACTs) are common treatments for malaria in endemic areas. Unfortunately, artemisinin-resistant parasites have spread across Southeast Asia and Africa, highlighting the need for new antimalarial drugs to combat this life-threatening parasitic disease[2–4]. In the last decade, the combination of phenotypic screening, and *in vitro* evolution and whole-genome analysis (IVIEWGA) has played a significant role in antimalarial drug development. This approach has contributed to the identification of several novel drug targets for malarial parasites, facilitating target-based HTS campaigns [5,6]. Despite the strong efforts in target-based drug discovery being conducted by the Medicines for Malaria Venture (MMV) and collaborating researchers, the development of novel antimalarial medicines often has been hampered by resistance of the malarial parasites to the drug candidates. For example, recrudescent parasites emerged in Phase II clinical trials of both of DSM265 and cipargamin that targets dihydroorotate dehydrogenase (DHODH) and the *Plasmodium falciparum* Na^+ pump PfATP4, respectively[7–9]. Although target-based drug discovery is an effective method for generating lead compounds, the targets identified by IVIEWGA carry the risk of potential

resistance, a major concern in the later stages of drug development. Therefore, there has been

renewed interest in phenotypic screening, with several of the resulting candidates now listed in

the current drug development pipeline of MMV.

Single-exposure radical cures (SERCs) are highly appealing, since this strategy

prevents parasite resistance and simplifies patients' treatment. However, SERCs

characteristically require fast-killing, as seen in next-generation lead compounds such as

MMV688533, INE963, ZY19489, and GSK484, the subjects of ongoing clinical trials[10-13]. The

standard method employed to determine the speed of killing of parasites, the parasite reduction

ratio (PRR), first was described in 2012; since then PRR has been utilized widely to assess the

killing properties of candidate compounds[14,15]. The conventional PRR assay has a relatively low

throughput due to the complexity of the protocol, requiring at least one month to yield results.

This aspect means that one cannot determine whether a given hit compound is a fast killer until

the candidate selection step in the drug development pipeline. Despite the crucial importance of

the PRR properties of next-generation lead compounds, no scalable assay method for

determining the PRR values of multiple compounds has been reported to date.

Phenotypic screening of the asexual blood stage (ABS) of malaria parasites often

employs the *P. falciparum* lactate dehydrogenase (PfLDH) assay to quantify parasite growth[16].

Specifically, PfLDH (in the lysate of a parasite culture) is coupled to the *Clostridium kluyveri*

diaphorase; the bacterial enzyme catalyzes a redox reaction that converts nitroblue tetrazolium

(NBT) to nitroblue formazan (NBF) via 3-acetyl pyridine adenine dinucleotide (APAD+), a

nicotinamide adenine dinucleotide (NAD+) analogue that exhibits cofactor specificity for PfLDH.

In the past, GlaxoSmithKline (GSK) employed the PfLDH assay to screen approximately 2 million

compounds in a 384-well plate format. However, of late, the SYBR green I-based fluorescence

assay, which detects parasite DNA, has become more commonly used for phenotypic HTS of the

ABS of malaria parasites. This shift occurred because the fluorescence assay was the only

method available for use in the 1536-well format[17,18]. Nevertheless, the simple principle of a

PfLDH assay that involves a redox enzyme and a detection probe coupled with the LDH reaction,

allows for easy tuning of the assay protocol depending on the specific purpose of the experiment.

In the present report, we describe the development of a modified PfLDH assay using

the nitroreducatase (NTR) encoded by the *Escherichia coli nfsB* gene. NTR, a flavoprotein, uses

NAD(P)H as an electron donor to reduce a variety of nitroaromatics[19]. To enhance the

conventional PfLDH assay's performance, we incorporated into the assay system "turn-on"

fluorescence probes that include a nitrobenzyl moiety[20,21] that is reduced by NTR. The use of a

near-infrared (NIR) probe with an extended wavelength range (excitation/emission

wavelengths ($\lambda_{ex/em}$) = 615/690 nm) compared to haemoglobin absorption ($\sim$600 nm)

effectively decreased interference from haemoglobin absorption in the lysate, resulting in an

increased signal window. This innovative approach using the NIR probe enabled us to conduct

phenotypic screening of the ABS of malaria in a 1536-well format, and improved the assay

robustness compared to that of the conventional SYBR green I assay. Moreover, the use of the

optimized PfLDH-NTR assay system streamlined the time-consuming protocol of conventional

PRR. A feasibility study with representative antimalarial drugs showed results consistent with

those obtained using the conventional PRR method. Our high-throughput PRR (HT-PRR) assay

therefore is expected to serve as a powerful tool for identifying, in the early stages of drug

discovery, novel antimalarial compounds with fast-killing profiles.

Results

Coupling of NTR reaction with PfLDH

Histidine-tagged NTR expressed in *E. coli* was purified successfully using conventional nickel-

nitrilotriacetic acid (Ni-NTA) agarose purification, yielding a specific activity in the range of 3–5

μmol/min/mg (Fig. 1a). The recombinant NTR converted nitrobenzyl-umbelliferone 1 (NCOU1)

to umbelliferone, such that the fluorescent signal of the product ($\lambda_{ex/em}$ = 315/455 nm) exhibited

a time-dependent increase during room temperature incubation (Fig. 1b, c). For our assay, PfLDH

was coupled with the purified NTR, rather than with the bacterial diaphorase used in the

conventional PfLDH assay, and parasite growth was quantified by monitoring the fluorescence

signal of umbelliferone (Fig. 2a). As anticipated, the NTR quantitatively reduced NCOU1 by using

the reducing equivalents from the APADH produced by the PfLDH enzyme present in the parasite

lysate. A titration of cultured *P. falciparum* using the PfLDH-NTR assay demonstrated that the

fluorescent signal increased in a parasitemia-dependent manner, with the highest signal

intensity seen under the condition of 1% haematocrit (Fig. 2b and Supplementary Fig. 1).

Next, we compared the screening performance, in a 384-well plate format, of the

PfLDH-NTR assay compared to the conventional PfLDH-diaphorase assay. Specifically, we used

the two assays to assess the growth inhibition of ABS parasites by a test set of 3,837 compounds

from Shionogi & Co., Ltd., using a compound concentration of 2.5 μM. A correlation ($r^2 = 0.387$)

was observed between the results obtained by the two methods, as depicted in Fig. 2c. While the

average signal-to-background (S/B) ratio for the NTR method (5.02) was lower than that of the

diaphorase method (8.31), the two assays demonstrated similar Z'-factor values (0.71 and 0.76,

respectively) (Table 1). Notably, the diaphorase method produced a higher number of hits (261

hits) than did the NTR method (123 hits). The PfLDH-diaphorase method may have selected for

a larger number of pseudo-positive hits, given that this assay's percent inhibition data did not

exhibit a Gaussian distribution and did not correlate with data obtained by other methods

(Supplementary Fig. 2, 3, 4). These results indicated that our fluorescence-based PfLDH assay,

which incorporates NTR and NCOU1, is more robust than the conventional PfLDH assay using

diaphorase.

NIR probe with PfLDH-NTR assay improves screening robustness

The PfLDH-NTR-NCOU1 assay was not suitable for a 1536-well format due to the low S/B ratio.

Consequently, our next objective was to improve this assay method using another probe with a

different wavelength of fluorescence emission. Given that the parasite lysate contains a

substantial amount of haemoglobin from red blood cells, we conjectured that the absorption of

haemoglobin (~600 nm) interferes with the fluorescent signal of umbelliferone[20]. Therefore, we

introduced another NIR probe, one that exhibits longer excitation and emission wavelengths

after reduction by NTR (Fig. 3a). We observed that, following reduction of the NIR probe by NTR

in the assay solution, the signal intensity of the reduced species (NIR-P[red]), increased in a time-

dependent manner, permitting quantification of the malaria parasites (Fig. 3b, c). Given the read-

time need for measurement of output in the 1536-well format, we quenched the PfLDH reaction

by adding a high concentration of pyruvate (0.9 M), resulting in the complete inhibition of

NCOU1 reduction (Fig 3c). For reactions run in the 1536-well format, the parasite growth

inhibition detected by the PfLDH-NTR-NIR assay showed a strong correlation with that obtained

by the SYBR green I-based assay (Fig. 3d). Notably, the PfLDH-NTR-NIR assay achieved a higher

S/B ratio and Z'-factor (11.2 and 0.72, respectively) than those measured by the SYBR green I

assay (5.89 and 0.61), indicating that PfLDH-NTR-NIR assay is more robust than the existing

method when employed in a 1536-well format.

High-throughput PRR (HT-PRR) assay

In the ABS, LDH oxidizes NADH and provides NAD^+ to maintain the flux of anaerobic glycolysis

required for the synthesis of ATP, a function that is essential for parasite survival[22,23]. We

conceived the idea of using PfLDH activity as an indicator of parasite viability and applied the

PfLDH-NTR-NCOU1 method in a 384-well format to quantify the PRRs of test compounds. Mixed-

stage (non-synchronized) parasites at 1.5% parasitemia were incubated at 37 °C for 0, 24, 48,

72, and 96 hours with test compounds at concentrations of 10 μM. Live parasite signals were

quantified by the PfLDH-NTR-NCOU1 method and the PRR values were calculated for each

compound. The values then were normalized (as a percentage) to those of dihydroartemisinin

(DHA; defined as 100%), a representative fast-killing compound, and ELQ300 (defined as 0%),

an inhibitor of the electron transport chain (ETC) that is classified as a slow-killing compound

(Supplementary Fig. 5). At 24 hours, fast-killing compounds like ACT-451840, artemether, and

cipargamin exhibited PRR values exceeding 70%; chloroquine, MMV048, and ganaplacide,

categorized as moderate-speed-killing compounds, yielded PRR values in the approximate range

of 40‒70%. Other known slow-killing ETC inhibitors, DSM265 and atovaquone, yielded lower

PRR values (16.9 and 8.6%, respectively). These results are consistent with the data obtained by

the conventional PRR method. The sole exception was DDD107498, a compound known to be a

slow-killing inhibitor by conventional PRR[24], show high PRR value. Given that this compound is

an inhibitor of the eukaryotic translation elongation factor 2 (eEF2)[24]; we hypothesize that the

interruption of protein synthesis by DDD107498 may directly impact PfLDH synthesis, leading

to lowered enzymatic activity, which our assay detects as an elevated PRR.

Subsequently, we sought to optimize the level of parasitemia in the assay by employing

our method to assess the effects, on parasites at various densities, of 24- and 48-hour exposure

to the known inhibitors atovaquone, DHA, and cipargamin (Supplementary Fig. 6). We observed

that when the starting parasitemia exceeded 2%, the signal for vehicle (dimethylsulfoxide;

DMSO)-treated parasites achieved a virtual plateau after 48 hours of incubation. Therefore, we

fixed the starting parasitemia at 1.5% for further experiments.

Having completed the optimization of HT-PRR, we conducted PRR assays in an 8-point

dose-response experiment. The dose-response PRR experiment was implemented because

during the test run, we observed that the shape of PRR dose curves differed among the

compounds (Supplementary Fig. 7). The experiment assessed PRR at intervals of 0, 24, 48, and

72 hours; and determined the 50% effective concentration to reduce the PRR values (PRR_{50}) of

the representative test compounds, in addition to the 72-hour cultures for standard growth

inhibition (EC_{50}) assays (Fig. 4a). A comparison of the inhibition curves revealed that Hill slopes

and plateau levels of the PRR curves varied among compounds and time points. To quantify the

shape of the dose-response curve and utilize these data as an indicator of PRR, we calculated for

each compound the ratio of area under the curve (AUC) between the growth inhibition curve

and the PRR curve, yielding a parameter that we refer to as the "AUC%" (Table 2). We observed

that the AUC% values of the fast-killing compounds exceeded 80%. In contrast, the slow-killing

compound exhibited lower AUC% values (of approximately 50%); furthermore, the % inhibition

by slow-killing compounds did not reach 100%, even at the highest tested concentration (1 μM).

These data suggested that robust AUC% values, calculated using multiple data points, permitted

effective classification of compounds according to their killing-rate properties.

Discussion

In the present work, we established a novel PfLDH-based assay method for detecting

antimalarial candidates targeting the ABS by using NTR and "turn-on" fluorescent probes with a

nitrobenzyl moiety. The new method provides an assay with robustness superior to that of

existing screening methods, and we successfully conducted HTS with a small compound library

in a 1536-well format. Our method also was able to assess the parasite survival rate, as assessed

by PfLDH activity. The PfLDH-NTR assay permits quantification of the PRR for many compounds

in a 384-well format, significantly decreasing experimental time. The PRR classification of

known antimalarial drugs by our HT-PRR method was consistent with the data obtained by

conventional assays. Moreover, HT-PRR permitted the determination of the dose-dependency of

a compound's PRR, yielding a new parameter (the AUC%) indicating the speed of killing. Our

novel PfLDH-NTR-based assay is expected to be valuable for determining the PRR values of many

hit compounds in the early stages of drug discovery, facilitating the development of next-

generation fast-killing antimalarial drugs.

In the field of cancer research, NTR is widely used for the *in vivo* detection of tumour

cells and the activation of prodrugs, reflecting the fact that NTR expression is specifically

increased in tumour cells growing under hypoxic conditions[25–27]. Additionally, NTR plays a

crucial role in the treatment of Chagas disease, a condition caused by *Trypanosoma cruzi*

infection; benznidazole and nifurtimox, front-line drugs used against Chagas, are activated by

NTR activity. Resistance to these drugs is determined primarily by loss of function mutations in

the *T. cruzi* NTR-encoding gene (including homozygous nonsense mutations and the loss of one

copy) that lead to decreased levels of the trypanosomal enzyme[28,29]. Given NTR's biological

significance and diverse applications in biomedical research, we envisioned using NTR to

establish innovative assays for the discovery of antimalarial drugs. Moreover, the tissue

transparency of the $\lambda_{ex/em}$ of NIR, which fall within the bio-optical window (650–1100 nm),

makes these wavelengths useful in biomedical research, especially for *in vivo* imaging, due to

lower haemoglobin absorption[30]. This property of the NIR probe fluorophore mitigates

interference from the abundant haemoglobin in red blood cells, greatly enhancing the signal

strength of the PfLDH assay. Through the introduction of a coupling enzyme and probes, we

improved the performance of the conventional PfLDH assay for detecting growth inhibition of

the malaria parasite at the ABS. In the future, it will be interesting to introduce modifications to

existing assay systems, thereby enhancing robustness and reducing cost. This initiative has the

potential to pave the way for the creation of advanced HTS systems applicable to other infectious

diseases, including *Trypanosoma* spp. and *Mycobacteria* spp.; indeed, revisions to facilitate

screens for these systems are currently in progress.

For HTS of the ABS of the malaria parasite in a 1536-well format, the commonly used

assay employs SYBR green I, a fluorescent dye with excitation and emission wavelengths of 485

nm and 528 nm, respectively. Our PfLDH-NTR assay with NIR[ox] would be (to our knowledge) the

first system using NIR wavelengths to screen for antimalarial compounds. A noteworthy finding

is that a small number of hit compounds of the SYBR green I-based assay was not common with

that of the PfLDH-NTR with NIR[ox] assay (Supplementary Fig. 4). The optical properties of small

organic compounds in chemical libraries may impact the readout signal, resulting in false-

positives and -negatives[31,32]. Indeed, a small number of compounds showed less than 40%

inhibition as assessed by the SYBR green I-based assay, possibly reflecting the optical properties

of these compounds (Supplementary Fig. 2). The PfLDH-NTR with NIR^{ox} assay may serve as a

viable alternative screening method to prevent the loss of promising compounds as pseudo-

negatives in future phenotypic HTS of the malaria parasite at the ABS.

To enhance the throughput in assessing PRR, we developed the PfLDH-NTR assay

system, enabling the HTS evaluation of antimalarial compounds based on PRR values. Notably,

we observed that the dose dependent PRRs of fast-killing compounds typically demonstrate

curves that resembled the growth inhibition curve of the same chemicals. However, disparities

between these two types of curves are more pronounced for moderate- and slow-killing

compounds. To quantify these differences, and to classify multiple test compounds according to

the PRR value, we defined the AUC%. This parameter provides a succinct description of the

parasite killing rate of test compounds and is expected to accelerate structure-activity

relationship (SAR) studies, particularly for fast-killing compounds. At the same time, we note

that a limitation of HT-PRR is that PRR values for compounds targeting protein synthesis (e.g.,

DDD107498) may be overestimated due to the dependence of the assay on PfLDH activity.

However, this limitation can be addressed by cross-referencing results with conventional PRR or

classical Giemsa staining at later stages of drug discovery.

Conclusion

We developed, in-house, NTR-based methods for the identification of compounds with

activity against the ABS of malaria parasites, and successfully implemented one such assay for

HTS in a 1536-well format. The use of the PfLDH-NTR method for PRR determination represents

a novel assay of the parasite-killing rate of drug candidates. These innovative new HTS methods

are expected to permit the identification of fast-killing compounds with novel scaffolds and

properties, paving the way for the discovery of next-generation antimalarial drugs that may find

use in combatting drug resistance in malaria-endemic areas.

Methods

Preparation of recombinant enzymes

Prior to assay development, we prepared two coupling enzymes, *Clostridium kluyveri* diaphorase

and *Escherichia coli* NAD(P)H nitroreductase NfsB (NTR), as previously described[19,33]. Briefly,

codon-optimized genes were synthesized to encode proteins with N-terminal His10-SUMO tags

and C-terminal His10 tags. These genes with C-terminal His10 were cloned separately into the

pET101-D-TOPO expression vector (Thermo Fisher Scientific), and the resulting plasmids were

transformed into *E. coli* BL21™ Star (DE3) cells according to the manufacturer's protocol. The

resulting *E. coli* strains were pre-cultured (overnight at 37 °C with shaking at 200 rpm) in Luria–

Bertani medium supplemented with 100 μg/mL carbenicillin. The resulting cultures were used

as inocula for 600-mL volumes of Terrific-Broth medium supplemented with 100 μg/mL

carbenicillin and 0.4% glycerol, and the resulting cultures were incubated at 37 °C with shaking

at 200 rpm. When the optical density at 600 nm (OD_{600}) reached 0.4 to 0.6, protein expression

was induced by adding 50 μM isopropyl β-D-1-thiogalactopyranoside (IPTG; Sigma) and 1

mg/mL riboflavin (Sigma). After incubation for 16 h at 20 °C, the cells were harvested by

centrifugation at 7,000 × g for 10 min at 4 °C and re-suspended at a density of 0.4 g cell pellet/mL

in cold lysis buffer (50 mM potassium phosphate buffer, pH 8.0, 300 mM NaCl, 0.5 mM

ethylenediaminetetraacetic acid (EDTA) and 0.25 mM phenylmethylsulfonyl fluoride (PMSF)).

The suspended cells were lysed using a French press (Ohtake) at 180 MPa, and the supernatant

was collected by centrifugation at 40,000 × g for 30 min at 4 °C. The His-tagged proteins were

purified using Ni-NTA (Qiagen) following the manufacturer's protocol and eluted using lysis

buffer containing 200 mM imidazole. The eluate was concentrated using an Amicon Ultra

Centrifugal Filter 10 kDa Molecular Weight Cut-off (Merck) according to the manufacturer's

protocol, and the resulting purified enzyme was mixed with an equal volume of cold glycerol and

stored frozen at -30 °C. The protein concentrations in the final enzyme stocks were determined

by the Bradford assay using a Bio-Rad kit, and the specific activities of the enzymes were

quantified by measuring the reduction of 2,6-dichlorophenolindophenol (DCIP; Sigma) using a

UV760 spectrophotometer (Jasco). To confirm the purity of the enzymes, proteins were

separated by sodium dodecyl sulfate-polyacrylamide gel electrophoreses (SDS-PAGE) and

stained using GelCode™ Blue Safe Protein Stain (Thermo Fisher Scientific). The specific activities

of diaphorase and NTR were typically in the range of 100–150 and 3–5 μmol/min/mg,

respectively.

Quantification of the reduction of fluorescence probes by NTR

To test whether the purified NTR was capable of reducing fluorescence probes, we initially

measured NTR activity prior to using the enzyme for parasite quantification. Using a cuvette, 20

μM NCOU1 (Enamine) was combined with a mixture containing 5 μg/mL NTR, 50 μM NADH, and

0.25 % Triton X-100 in 30 mM Tris-HCl (pH 8.0), and the fluorescence of the umbelliferone

($\lambda_{ex}/_{em}$ = 315/455 nm) generated by the reaction was measured. In addition, we synthesized the

NIR[ox] probe according to the previously described route[21]. For the NIR reaction, 50 μM NIR-P[ox]

was combined with a mixture containing 0.5 μg/mL NTR and 50 μM NADH in 30 mM Tris-HCl

(pH 8.0), and the fluorescence of NIR-P[red] ($\lambda_{ex}/_{em}$ = 615/670 nm) was measured. The

fluorescence kinetics of both reduced probes were monitored using a FP-6300

spectrofluorometer (Jasco).

Parasite cultures

Plasmodium falciparum 3D7 parasites were maintained in RPMI1640 medium supplemented

with 25 mg/L gentamicin, 50 mg/L hypoxanthine, 23.8 mM sodium bicarbonate, and 0.5% (w/v)

Albumax II (Gibco). The culture was maintained at 2% haematocrit using O[+] human erythrocytes

obtained from The Japanese Red Cross Society. As per established protocols[34,35], the incubation

temperature was set at 37 °C and the environment consisted of mixed gas (5% O_2, 5% CO_2, 90%

N_2). Parasitemia was determined using either Giemsa staining or an XN-30 automated

haematology analyser (Sysmex)[36]. For plate assays, ring-stage synchronized parasites were

prepared using sorbitol treatment[37]. In the growth inhibition assay, the compound solvent DMSO

and a mixture of 1 μM artemisinin/atovaquone served as negative and positive controls,

respectively.

Diaphorase-NBT and NTR-NCOU1 assay in 384-well plates

Test compounds were pre-dispensed at a volume of 100 nL/well into 384-well plates. For the

diaphorase-NBT assay, ring-stage parasites at 0.3% parasitemia and 2% haematocrit were

dispensed into 384-well clear plates (Corning); for the NTR-NCOU1 assay, ring-stage parasites

at 0.3% parasitemia and 1% haematocrit were dispensed into 384-well black plates (Greiner).

Parasites then were dispensed at 25 μL/well into the compound-containing plates using a

Multidrop Combi with a small metal cassette (Thermo Fisher Scientific). The parasite cultures in

the assay plates then were incubated in a moist chamber filled with mixed gas for 72 hours and

subsequently frozen overnight (or longer) at -30 °C. The resulting frozen plates were thawed at

room temperature for at least 2 hours before the assay was performed. For the diaphorase-NBT

assay, the reaction mix was prepared in modified PfLDH buffer (100 mM Tris-HCl, pH 8.0, 150

mM lithium L-lactate, and 0.25% Triton X-100) supplemented with 75 μM $APAD^+$, 0.2 mg/mL

NBT, and 1 U/mL diaphorase (one unit was defined as the amount of enzyme required to reduce

1 μmol of APADH per minute). For this assay, the reaction mixture was dispensed into the assay

plates at 70 μL/well, and the plates then were incubated at room temperature with agitation at

650 rpm for 20 minutes. On the other hand, the reaction mixture for the NTR-NCOU1 assay

consisted of modified PfLDH buffer supplemented with 100 μM APAD$^+$, 100 μM NCOU1, 0.05

U/mL NTR, and 0.015% (v/v) KM-70 defoaming agent (Shin-Etsu). For this assay, the reaction

mix was dispensed into the assay plates at 35 μL/well, and the plates then were incubated under

the same condition as those used for the diaphorase-NBT assay. The absorption of nitro blue

formazan (650 nm; for the diaphorase-NBT assay) or the fluorescence of umbelliferone ($\lambda_{ex/em}$ =

360/465 nm; for the NTR-NCOU1 assay) was measured using a SpectraMax® Paradigm

spectrophotometer (Molecular Devices, Inc.).

NTR-NIR^{ox} assay in 1536-well plates

For the HTS in 1536-well format, ring-stage parasites at 0.3% parasitemia and 1%

haematocrit were dispensed at 4 μL/well into 1536-well black plates (Greiner) containing 20

nL/well of test compounds. Parasite-dispensed assay plates were maintained under the same

conditions as described above for the 384-well plate assays. The NIR^{ox} probe was initially

suspended at 1 mM in 90% acetonitrile. The reaction mixture for the NTR-NIR^{ox} assay was

prepared by supplementing modified PfLDH buffer with 500 μM APAD$^+$, 200 μM NIR^{ox} probe,

and 0.05 U/mL NTR. For this assay, the reaction mixture was dispensed into the assay plates at

3 μL/well, and the contents of each well were mixed at 1,200 rpm. The plates then were

incubated at room temperature for 3 minutes. The enzymatic reactions were quenched by

dispensing the "stop" mixture (0.9 M sodium pyruvate containing 0.015% KM-70) at 3 μL/well,

and the assay plates then were centrifuged at 90 × g for 1 min at room temperature. The

centrifuged plates were incubated for 1 hour at room temperature in a moist chamber, at which

point the NIR^{red} signal ($\lambda_{ex/em}$ = 615/690 nm) was measured using a PHERAstar Plus microplate

reader (BMG LABTECH).

SYBR green assay in 1536-well plates

For the SYBR green assay, ring-stage parasites at 0.3% parasitemia and 3% haematocrit were

dispensed at 4 μL/well into 1536-well black plates (Greiner) containing 20 nL/well of test

compounds. Parasite-dispensed assay plates were maintained under the same conditions as

described above for the 384-well plate assays. The reaction mixture for the SYBR green I assay

consisted of lysis buffer (20 mM Tris-HCl, pH 8.0, 5 mM EDTA, and 0.1% Triton X-100)

supplemented with 0.02% SYBR green I and 0.015% KM-70. This reaction mixture was

dispensed into the assay plates at 4 μL/well, and the contents of each well were mixed at 1,200

rpm. The plates then were incubated at room temperature for 1 hour. The fluorescence of SYBR

green I ($\lambda_{ex}/_{em}$ = 485/528 nm) was measured using a SpectraMax® Paradigm

spectrophotometer.

Data analysis

Hits of a test set of compounds were chosen by activity criteria [>3 standard deviation (SD)

cutoff of DMSO controls] that were calculated using Spotfire software (version 11.4.3; TIBCO). Z'

was calculated as $1 - (3 \times SD_{100\%} + 3 \times SD_{0\%})/(Mean_{100\%} - Mean_{0\%})$. The correlations

of %Inhibition and histograms were visualized with R (version 4.0.2). The inhibition curves and

EC_{50} values were determined using GraphPad Prism (version 8.4.3. GraphPad Software, Inc., San

Diego, California, USA) and Spotfire. Data visualization for the other figures was performed using

GraphPad Prism. The area under the curve (AUC) for both the HT-PRR assay and growth

inhibition assay in the 384-well plates was calculated using Spotfire software, and the AUC% was

calculated as $100 \times [(AUC$ of HT-PRR assay) / (AUC of NTR-NCOU1 assay)]. Venny (version 2.1)

was used to make a Venn diagram.

HT-PRR assay in 384-well plates

Test compounds were pre-dispensed at 100 nL/well in 384-well plates, and replicate plates were

prepared. Non-synchronized parasites at 1.5% parasitemia and 1% haematocrit were dispensed

into five 384-well black compound-containing plates at 25 μL/well using a Multidrop Combi

with a small metal cassette. One of the plates was frozen at -30 °C right immediately following

dispensing of parasites (control plate; 0 hour), while the other plates were incubated in a moist

chamber filled with mixed gas for 24, 48, 72, or 96 hours; at the indicated time points, growth

was terminated by freezing overnight (or longer) at -30 °C (PRR plates). Additional plates for the

generation of the growth inhibition curve were prepared and subjected to the NTR-NCOU1 assay

using the procedure described in the previous section, with a 72-hour incubation period. The

fluorescence of umbelliferone ($\lambda_{ex/em}$ = 360/465 nm) was measured using a PHERAstar

microplate reader, and the relative LDH activity was calculated by dividing the fluorescence of

the PRR plates by that of the respective control plate. The PRR inhibition percentage was

calculated by normalizing the primary values to the LDH activity obtained in the negative and

positive control reactions (compound solvent DMSO and 1 μM dihydroartemisinin, respectively).

In addition, the concentration of each compound to reduce the PRR value to 50% (PRR_{50}) was

calculated in each time point.

Acknowledgements

The authors thank the Japanese Red Cross Society for providing human red blood cells (RBCs;

Registry No. R030038) and Dr. Kenji Takaya (Laboratory for Medicinal Chemistry Research,

Shionogi & Co., Ltd.) for synthesizing the NIR probe.

Abbreviations Used

PfLDH, *Plasmodium falciparum* lactate dehydrogenase; NTR, nitroreductase; PRR, parasite

reduction ratio; ACTs, artemisinin-based combination therapies; IVIEWGA, *in vitro* evolution and

whole-genome analysis; MMV, Medicines for Malaria Venture; DHODH, dihydroorotate

dehydrogenase; SERCs, single-exposure radical cures; ABS, asexual blood stage; NBT, nitroblue

tetrazolium; NBF, nitroblue formazan; APAD$^+$, 3-acetyl pyridine adenine dinucleotide; NIR, near-

infrared; NCOU1, nitrobenzyl-umbelliferone 1; ETC, electron transport chain; DHA,

dihydroartemisinin; eEF2, elongation factor 2; PMSF, phenylmethylsulfonyl fluoride; DCIP, 2,6-

dichlorophenolindophenol

Fundings

This work was supported, in part, by the following funding sources: grants for Infectious Disease

Control from the Science and Technology Research Partnership for Sustainable Development

(SATREPS; No. JP10000284 to K.K., and No. JP14425718 to D.K.I. and T.S.); grants from the

Agency for Medical Research and Development (AMED); Grants-in-aid for Scientific Research (A)

(No. 20H00620 to D.K.I.), (B) (23H02711 to K.K. and D.K.I.), and (C) (No. JP22512047 to T.S.); a

grant from The Leading Initiative for Excellent Young Researchers (LEADER; No. JP16811362 to

D.K.I.) from the Japanese Ministry of Education, Science, Culture, Sports and Technology (MEXT);

and by Grants-in-aid for Research on Emerging and Re-emerging Infectious Diseases from the

Japanese Ministry of Health and Welfare (No. 21fk0108138 and No. 23fk0108680 to D.K.I.). This

study was conducted (in part) at the Joint Usage/Research Center on Tropical Disease, Institute

of Tropical Medicine, Nagasaki University, Japan. This work also was supported by Shionogi &

Co., Ltd., Osaka, Japan.

Author contributions

433   D.K.I., T.K., and K.K. directed the work. T.S., R.I., and E.Y. performed the experiments. T.S., R.I., and

434   E.Y. analysed the data. T.S., R.I., E.Y., and D.K.I. wrote the paper. All authors directly participated

435   in this work, including preparation of the manuscript and approval of the final version. All

436   authors have read and agreed to the published version of the manuscript.

Conflict of interest

RI and TK are employees of Shionogi & Co., Ltd..

Nagasaki University and Shionogi & Co., Ltd., launched the Shionogi Global Infectious Division

(SHINE) at the Institute of Tropical Medicine NEKKEN, Nagasaki University. Under the

collaboration agreement, Shionogi & Co., Ltd. provided the NIR-P reagent and compounds pre-

dispensed into the 384- and 1536-well plates for validation of HTS, and contributed to

discussions with the authors. Shionogi & Co., Ltd did not influence the experimental design, data

collection, data analysis, or interpretation.

References

1. World Health Organization. *World malaria report 2022*. (World Health Organization, 2022).

2. Ariey, F. *et al.* A molecular marker of artemisinin-resistant Plasmodium falciparum malaria. *Nature* **505**, 50–55 (2014).

3. Mbengue, A. *et al.* A molecular mechanism of artemisinin resistance in Plasmodium falciparum malaria. *Nature* **520**, 683–687 (2015).

4. Balikagala, B. *et al.* Evidence of Artemisinin-Resistant Malaria in Africa. *N. Engl. J. Med.* **385**, 1163–1171 (2021).

5. Cowell, A. N. *et al.* Mapping the malaria parasite druggable genome by using in vitro evolution and chemogenomics. *Science* **359**, 191–199 (2018).

6. Siqueira-Neto, J. L. *et al.* Antimalarial drug discovery: progress and approaches. *Nat. Rev. Drug Discov.* **22**, 807–826 (2023).

7. Llanos-Cuentas, A. *et al.* Antimalarial activity of single-dose DSM265, a novel plasmodium dihydroorotate dehydrogenase inhibitor, in patients with uncomplicated Plasmodium falciparum or Plasmodium vivax malaria infection: a proof-of-concept, open-label, phase 2a study. *Lancet Infect. Dis.* **18**, 874–883 (2018).

8. Schmitt, E. K. *et al.* Efficacy of Cipargamin (KAE609) in a Randomized, Phase II Dose-Escalation Study in Adults in Sub-Saharan Africa With Uncomplicated Plasmodium

falciparum Malaria. *Clin. Infect. Dis.* **74**, 1831–1839 (2022).

9.   Yang, T. *et al.* MalDA, Accelerating Malaria Drug Discovery. *Trends Parasitol.* **37**, 493–507

(2021).

10. Murithi, J. M. *et al.* The antimalarial MMV688533 provides potential for single-dose cures

with a high barrier to Plasmodium falciparum parasite resistance. *Sci. Transl. Med.* **13**,

(2021).

11. Taft, B. R. *et al.* Discovery and Preclinical Pharmacology of INE963, a Potent and Fast-Acting

Blood-Stage Antimalarial with a High Barrier to Resistance and Potential for Single-Dose

Cures in Uncomplicated Malaria. *J. Med. Chem.* **65**, 3798–3813 (2022).

12. Barber, B. E. *et al.* Safety, pharmacokinetics, and antimalarial activity of the novel

triaminopyrimidine ZY-19489: a first-in-human, randomised, placebo-controlled, double-

blind, single ascending dose study, pilot food-effect study, and volunteer infection study.

Lancet Infect. Dis. **22**, 879–890 (2022).

13. Gomes, P. & Guido, R. V. C. Editorial: Antimalarial chemotherapy in the XXIst century, volume

II. *Front. Pharmacol.* **14**, 1229764 (2023).

14. Sanz, L. M. *et al.* P. falciparum in vitro killing rates allow to discriminate between different

antimalarial mode-of-action. *PLoS One* **7**, e30949 (2012).

15. Walz, A. *et al.* The Parasite Reduction Ratio (PRR) Assay Version 2: Standardized

Assessment of Plasmodium falciparum Viability after Antimalarial Treatment In Vitro.

Pharmaceuticals **16**, (2023).

16. Makler, M. T. & Hinrichs, D. J. Measurement of the lactate dehydrogenase activity of

Plasmodium falciparum as an assessment of parasitemia. *Am. J. Trop. Med. Hyg.* **48**, 205–

210 (1993).

17. Gamo, F.-J. *et al.* Thousands of chemical starting points for antimalarial lead identification.

Nature **465**, 305–310 (2010).

18. Smilkstein, M., Sriwilaijaroen, N., Kelly, J. X., Wilairat, P. & Riscoe, M. Simple and inexpensive

fluorescence-based technique for high-throughput antimalarial drug screening. *Antimicrob.*

Agents Chemother. **48**, 1803–1806 (2004).

19. Zenno, S., Koike, H., Tanokura, M. & Saigo, K. Gene cloning, purification, and characterization

of NfsB, a minor oxygen-insensitive nitroreductase from Escherichia coli, similar in

biochemical properties to FRase I, the major flavin reductase in Vibrio fischeri. *J. Biochem.*

120, 736–744 (1996).

20. Sedgwick, A. C. *et al.* A simple umbelliferone based fluorescent probe for the detection of

nitroreductase. *Frontiers of Chemical Science and Engineering* **12**, 311–314 (2018).

21. Shi, Y., Zhang, S. & Zhang, X. A novel near-infrared fluorescent probe for selectively sensing

nitroreductase (NTR) in an aqueous medium. *Analyst* **138**, 1952–1955 (2013).

22. Zhang, M. *et al.* Uncovering the essential genes of the human malaria parasite Plasmodium

falciparum by saturation mutagenesis. *Science* **360**, (2018).

23. Shoemark, D. K., Cliff, M. J., Sessions, R. B. & Clarke, A. R. Enzymatic properties of the lactate

dehydrogenase enzyme from Plasmodium falciparum. *FEBS J.* **274**, 2738–2748 (2007).

24. Baragaña, B. *et al.* A novel multiple-stage antimalarial agent that inhibits protein synthesis.

Nature **522**, 315–320 (2015).

25. Li, Y. *et al.* Ultrasensitive near-infrared fluorescence-enhanced probe for in vivo

nitroreductase imaging. *J. Am. Chem. Soc.* **137**, 6407–6416 (2015).

26. Chevalier, A., Zhang, Y., Khdour, O. M., Kaye, J. B. & Hecht, S. M. Mitochondrial Nitroreductase

Activity Enables Selective Imaging and Therapeutic Targeting. *J. Am. Chem. Soc.* **138**, 12009–

12012 (2016).

27. Copp, J. N. *et al.* Engineering a Multifunctional Nitroreductase for Improved Activation of

Prodrugs and PET Probes for Cancer Gene Therapy. *Cell Chem Biol* **24**, 391–403 (2017).

28. Hall, B. S., Bot, C. & Wilkinson, S. R. Nifurtimox activation by trypanosomal type I

nitroreductases generates cytotoxic nitrile metabolites. *J. Biol. Chem.* **286**, 13088–13095

(2011).

29. Wilkinson, S. R., Taylor, M. C., Horn, D., Kelly, J. M. & Cheeseman, I. A mechanism for cross-

resistance to nifurtimox and benznidazole in trypanosomes. *Proc. Natl. Acad. Sci. U. S. A.* **105**,

5022–5027 (2008).

30. Hammerich, M. *et al.* Heterodiazocines: Synthesis and Photochromic Properties, Trans to

Cis Switching within the Bio-optical Window. *J. Am. Chem. Soc.* **138**, 13111–13114 (2016).

31. Joung, J. F., Han, M., Jeong, M. & Park, S. Experimental database of optical properties of

organic compounds. *Sci Data* **7**, 295 (2020).

32. Simeonov, A. *et al.* Fluorescence spectroscopic profiling of compound libraries. *J. Med. Chem.*

51, 2363–2371 (2008).

33. Chakraborty, S., Sakka, M., Kimura, T. & Sakka, K. Cloning and expression of a Clostridium

kluyveri gene responsible for diaphorase activity. *Biosci. Biotechnol. Biochem.* **72**, 735–741

(2008).

34. Komatsuya, K. *et al.* Siccanin Is a Dual-Target Inhibitor of Plasmodium falciparum

Mitochondrial Complex II and Complex III. *Pharmaceuticals* **15**, (2022).

35. Trager, W. & Jensen, J. B. Human malaria parasites in continuous culture. *Science* **193**, 673–

675 (1976).

36. Tougan, T. *et al.* An automated haematology analyzer XN-30 distinguishes developmental

stages of falciparum malaria parasite cultured in vitro. *Malar. J.* **17**, 1–8 (2018).

37.   Lambros, C. & Vanderberg, J. P. Synchronization of Plasmodium falciparum

erythrocytic stages in culture. *J. Parasitol.* **65**, 418–420 (1979).

Tables

Table 1: Parameters for each assay method

Method	Plate format	S/B	CV (%)		Z'	Number of hits	Hit rate (%)
			Negative	Positive			
Diaphorase-NBT	384	8.31	8.1	2.6	0.71	261	6.8
NTR-NCOU1	384	5.02	5.6	3.9	0.76	123	3.2
SYBR green	1536	5.89	9.6	6.2	0.61	66	1.7
NTR-NIR	1536	11.2	8.0	6.2	0.72	70	1.8

538 NBT: nitroblue tetrazolium, NTR: nitroreductase, NCOU1: nitrobenzyl-umbelliferone 1, NIR:

539 near-infrared, S/B: signal/background ratio, CV: coefficient of variation

541 **Table 2: Activity and AUC% of test compounds**

Compound	PRR_{50} (nM)			EC_{50} (nM)	AUC%		
	24 h	48 h	72 h		24 h	48 h	72 h
ACT-451840	2.77	2.72	4.07	1.95	89.2	95.6	90.6
Artemether	16.5	16.4	35.2	6.34	87.1	88.2	77.3
DHA	3.36	4.13	11.8	2.05	94.2	93.9	81.8
Cipargamin	1.61	1.17	1.15	1.29	88.6	98.2	100.1
DDD-107498	2.70	3.12	3.67	0.84	76.1	84.0	85.0
Chloroquine	16.9	11.3	10.4	4.85	78.6	86.6	89.1
Ganaplacide	117.5	30.0	15.9	7.05	55.1	71.9	80.0
Lumefantorine	24.7	15.3	15.0	7.28	67.9	67.4	66.7
Mefloquine	52.9	54.5	66.7	15.9	76.1	72.3	71.7
MMV048	276.9	209.6	23.5	23.9	62.6	72.1	78.3
Atovaquone	124.9	15.0	20.8	0.57	49.5	51.4	46.9
DSM265	1186	211.5	332.4	5.84	50.6	51.7	50.2
ELQ-300	9990	139.3	298.5	24.8	40.3	56.9	48.5
Pyrimethamine	9990	9990	9990	15.3	10.1	21.5	0.00
Cytochalasin D	9990	15.9	24.1	20.5	24.7	65.3	86.8

542 PRR_{50}: Concentration of the compounds required to reduce the PRR values to 50%; EC_{50}:

543 effective concentration to reduce the parasite growth to 50%; AUC: area under the curve; DHA:

544 dihydroartemisinin

Figures

Figure 1. Purification of recombinant nitroreductase (NTR) and quantification of

enzymatic activity. a Recombinant C-terminally His$_{10}$-tagged NTR was expressed in *E. coli* BL21

Star (DE3) and purified using Ni-NTA agarose. In the lanes denoted as Marker, Lysate,

Flowthrough, Wash1 and Wash2, a total of 10 μg protein were loaded. For Elution and

Concentrated lanes, 2 μg were loaded. Estimated protein size is 26.7 kDa. **b** Scheme of the

enzymatic reaction catalyzed by NTR reducing NCOU1 to umbelliferone ($\lambda_{ex/em}$ = 315/455 nm).

c The fluorescent spectra of umbelliferone. Time-dependent increases in the fluorescence

spectra were recorded for 60 minutes at room temperature in NTR assay buffer (30 mM Tris-

HCl, pH 8.0, 0.25 % Triton X-100) containing 5 μg/mL NTR, 20 μM NCOU1, and 50 μM NADH

($\lambda_{ex/em}$ = 315/350–600 nm over 0.5-nm intervals, recorded every 5 minutes). For better

visualization, only the spectra from time points at 0, 10, 20, 30, 45, and 60 minutes are shown

(grey: 0 min, orange: 10 min, yellow: 20 min, light blue: 30 min, blue: 45 min, purple: 60 min).

Fluorescence intensity (FI) over time ($\lambda_{ex/em}$ = 315/455 nm) is plotted in the inset. NCOU1:

nitrobenzyl-umbelliferone 1, NAD^+: oxidized nicotinamide adenine dinucleotide, NADH:

reduced nicotinamide adenine dinucleotide, a.u.: arbitrary unit.

Figure 2. Novel assay using nitroreductase (NTR) coupled with *Plasmodium falciparum*

lactate dehydrogenase (PfLDH).

a Scheme of the PfLDH-NTR coupled reaction used for quantification of parasite growth using

NCOU1 as the NTR probe. **b** Parasitemia titration measured by the NTR-NCOU1 assay. The

umbelliferone signal increased in a parasitemia-dependent manner. Each parasite culture (1%

haematocrit (Hct)) was assayed with PfLDH buffer (100 mM Tris-HCl (pH 8.0), 150 mM lithium

L-lactate, and 0.25 % Triton X-100) containing 100 µM APAD$^+$, 100 µM NCOU1, and 0.05–0.1

U/mL NTR at room temperature. **c** Correlation of %Inhibition between conventional

diaphorase-NBT and NTR-NCOU1 assays of PfLDH activity. The red line indicates unity (y = x).

NCOU1: nitrobenzyl-umbelliferone 1, $APAD^+$: oxidized 3-acetylpyridine adenine dinucleotide,

APADH: reduced 3-acetylpyridine adenine dinucleotide, RFU: relative fluorescence units. RBC:

red blood cell.

Figure 3. Nitroreductase (NTR) assay using near-infrared probe (NIR-P). a Scheme of

enzymatic reaction of NTR and NIR-P[ox] producing the fluorescent NIR-P[red] ($\lambda_{ex/em}$ = 615/670

nm**). b** Time-dependent fluorescent spectra of NIR-P[red] produced by NTR. The enzymatic

reaction catalyzed by NTR was monitored using 50 μM NIR-P^{ox} and 50 μM NADH in 30 mM Tris-

HCl, pH 8.0, buffer containing 0.5 μg/mL NTR. Fluorescence signals ($\lambda_{ex/em}$ = 605/620–740 nm

over 1-nm intervals) were measured over a 60-minute period at room temperature (orange: 0

min, yellow: 2 min, light green: 3 min, blue: 5 min, purple: 10 min, red: 60 min). Fluorescence

intensity (FI) over time is plotted in the inset. Note that the reduction of NIR-P by NTR is much

faster than the reaction observed using the NCOU1 substrate (Fig. 1c). **c** Parasitemia titration

measured using the NTR-NIR assay. Each parasite culture (1% haematocrit (Hct)) was assayed

in PfLDH buffer (100 mM Tris-HCl, pH8.0, 300 mM lithium L-lactate, and 0.25% Triton X-100)

containing 500 μM APAD$^+$, 200 μM NIR probe, and 0.05 U/mL NTR at room temperature. The

quenching ("stop") solution consisted of 0.9 M pyruvate. Data are presented as the mean ±

standard deviation (SD) (n=4). **d** Correlation of %Inhibition between conventional SYBR green

and NTR-NIR assays of PfLDH activity. The red line in the graph represents the line of equality

(y = x). a.u.: arbitrary unit, NAD$^+$: oxidized nicotinamide adenine dinucleotide, NADH: reduced

nicotinamide adenine dinucleotide, RFU: relative fluorescence units.

Figure 4. Schematic representation of high-throughput parasite reduction ratio (HT-PRR)

and assessment of well-characterized antimalarial agents. a To evaluate the parasite

reduction ratio (PRR) by established HT-PRR methods, non-synchronized 3D7 parasites

(haematocrit (Hct) 1%, parasitemia 1.5%) were plate-cultured for 0, 24, 48, and 72 hours in the

presence of compound. For the growth inhibition assay, 3D7 parasites were synchronized by

exposure to 5% sorbitol; ring stage parasites then were plate-cultured for 72 hours in the

presence of compound. Growth was terminated at the indicated time point by freezing (at -30 °

C) the respective assay plates; the number of live parasites then was quantified via an LDH assay

using retororeductase (NTR) and nitrobenzyl-umbelliferone 1 (NCOU1). For the growth

inhibition assay, the positive control consisted of a mixture of 1 μM atovaquone and artemisinin;

the negative control consisted of 0.4% dimethylsulfoxide (DMSO). For the HT-PRR assay, the

positive and negative controls consisted of 1 μM dihydroartemisinin (DHA) and 0.4% DMSO,

respectively. **b** Dose-response curves of Inhibition (%) and PRR (%) by test compounds were

generated for each assay condition (n =2).