A comprehensive collection of pain and addiction targets and compounds library for high throughput screening and AI-driven drug discovery

Xin Hu, Paul Shinn, Zina Itkin, Lin Ye, Ya-Qin Zhang, Min Shen, Stephanie Ford-Scheimer, Matthew D. Hall*

National Center for Advancing Translational Sciences (NCATS), National Institutes of Health,

9800 Medical Center Drive, Rockville, MD, 20850, United States

*To whom correspondence should be addressed:

Matthew D. Hall, Ph.D., hallma@mail.nih.gov

Abstract

As part of the NIH Helping to End Addiction Long-term (HEAL) Initiative, the National Center for Advancing Translational Sciences (NCATS) is dedicated to the development of new pharmacological tools and investigational drugs for managing and treating pain, as well as the prevention and treatment of opioid misuse and addiction. In line with these objectives, we created a comprehensive, annotated small molecule library including drugs, probes, and tool compounds that act on published pain and addiction-relevant targets. Nearly 3,000 small molecules associated with approximately 200 known and hypothesized HEAL targets have been assembled, curated, and annotated in one collection. Physical samples of the library compounds have been acquired and plated in 1536-well format, enabling rapid and efficient high throughput screen (HTS) against a wide range of assays. The creation of the HEAL Targets and Compounds Library, coupled with an integrated computational platform for AI-driven machine learning (ML), structural modeling, and virtual screening (VS), provides a valuable source for strategic drug repurposing, innovative profiling, and hypothesis testing of novel targets related to pain and addiction. The library is available to investigators for screening pain and addiction-relevant phenotypes.

Introduction

The opioid crisis highlights the urgent need for novel non-addictive pain medications, as well as improved treatments for opioid addiction and overdose [1, 2]. Advances in our understanding of the molecular and neural circuit physiology of pain, addiction, and overdose have unveiled many putative approaches for therapeutic intervention, such as pharmacological agents directed at novel targets to test new therapeutic hypotheses, the development of new drugs aimed at these targets, and the establishment of new testing systems with the potential to offer more accurate predictions of human clinical response compared to traditionally used models [3, 4]. Significant challenges remain for the creation of new therapeutics [5, 6], largely due to the absence of drug-like pharmacological probes for novel targets, hindering the exploration of new therapeutic hypotheses. Additionally, there is a lack of preclinical models capable of reliably predicting human responses, and the market attractiveness for addiction and overdose treatments is currently limited [7].

The Helping to End Addiction Long-term (HEAL) Initiative, launched in April 2018 as a collaborative effort across the National Institutes of Health (NIH), is dedicated to advancing national priorities in combating the opioid crisis through scientific endeavors [8-10]. As part of the HEAL Initiative, the National Center for Advancing Translational Sciences (NCATS) is committed to accelerating the process of discovery and demonstration of new treatments for opioid misuse, addiction, and pain. Our collaborative program actively engages in the development of novel chemistry, screening, and testing methodologies to discover new

pharmacological probes and investigational drugs tailored for addressing the challenges of pain and addiction.

Pharmacogenomics can be a valuable research approach for developing and evaluating cell-based models of pathologies such as those associated with the HEAL Initiative. This approach requires the availability of a library of small molecules with annotated mechanisms of action whose activity in biological assays can be related back to target biology. Ideally, such annotated libraries should contain a variety of molecules with redundant mechanisms of action to enhance confidence in their activity, and the target space covered by molecules present in the library is typically sourced from the existing literature. Beyond providing insight into biological mechanisms, screening with annotated libraries also offers the opportunity to identify potential drug repurposing candidates for the treatment of specific disorders. A range of annotated libraries have been reported, each designed with a particular focus. Some libraries are aiming at approved therapeutics (for example, the NCATS Pharmaceutical Collection, NPC [11]) or coverage of clinical space generally (such as the Broad Drug Re-purposing Hub [12]). Annotated libraries have also been designed around specific disease biology target space, as seen in the case of the NCATS MIPE library, which contains molecules with defined mechanisms of action associated with oncology indications [13].

As of now, there is no annotated library specifically designed to address the pain and addiction biological space. To fill this gap, we have created a comprehensive annotated HEAL library by curating known and potential targets related to pain and addiction, as well as assembling known drugs, chemical probes, and tool compounds associated with each of these

targets. The creation of the HEAL Targets and Compounds Library incorporates valuable insights gained from high-throughput screening and chemical library design at NCATS. This best-in-class repurposing library enables rapid testing of drug candidate compounds with established safety profiles in the context of pain and addiction, as well as the identification of targets and pharmacologies associated with mediating pain and addiction phenotypes. The collection can be made available in pre-spotted plates for investigators with laboratory assays able to test against novel biology in 384-well plates. Moreover, the HTS data generated at NCATS with this library will be openly shared with the scientific community, providing a valuable resource for researchers.

Here, we provide detailed information on the creation and composition of the HEAL library. A case study on several HEAL targets using quantitative HTS (qHTS) and *in silico* screening for probe and drug discovery as part of the collaborative program at NCATS is summarized. This case study illustrates the practical application of the HEAL targets and compounds as an integrated platform for AI-driven machine learning, *in silico* structural modeling, and large-scale database virtual screening for therapeutics development against pain and addiction.

Creating the HEAL Library target list

For the purposes of the HEAL Library design, potential HEAL targets could be broadly defined to include any targets that are involved in opioid and pain signaling pathways and implicated as a regulator of pain and inflammation. A survey of validated protein targets for addiction and pain in DrugBank and ChEMBL showed that only about 30-40 small molecule drug targets are associated with pain and addiction. These classical analgesic targets that provide

clinically validated efficacy are mostly nociceptors and neurological targets, including opioid receptors (e.g., mu opioid receptor MOR), dopamine receptors (e.g. D3R), serotonin receptors (e.g. 5-HT7R), adrenoceptor (A2A), cannabinoid receptors, and ion channels such as NMDA, GABA receptor and AChR. To tackle the problems and side effects associated with these traditional drugs and targets, novel approaches have been developed, such as the opioid receptors that block pain but don't affect breathing or reward signaling, or alternative opioid receptors that are not located in the reward center of the brain [5]. On the other hand, a large number of non-opioid pain targets have been identified for the development of novel analgesics. These novel targets, including P2X3 and P2X7 purinoceptors, NaV1.7 and NaV1.8 sodium-channels, orphan receptors GPR18/GPR35/GPR55, enzyme Triointsub receptor kinase A (TrkA), sepiapterin reductase (SPR), and soluble epoxide hydrolase (sEH) have been reviewed recently [14, 15].

In addition to assembling these classical pain and addiction-related targets in the HEAL Library's list, we also included a broader collection of novel and hypothesized targets that have been studied for pain and addiction treatment in the literature. For example, CRMP2 is an essential trafficking protein for NaV1.7, a valid pain target that is challenging to drug efficiently and specifically. Small molecules to modulate CRMP2 provide an alternative and novel strategy for blocking NaV1.7 function and reducing pain by preventing its trafficking to the neuronal plasma membrane [16]. The galanin 1 receptor (GAL1R) forms heteromers with MOR in the ventral tegmental area that modulates dopamine release. GAL1R ligands block MOR-mediated signaling by allosteric modulations within the MOR-GAL1R heterodimer. Targeting MOR-GAL1R heterodimers offers an appealing strategy for novel drug design to treat OUD while sparing the analgesic effect of opioids [17]. Some novel targets that are currently under investigation for high throughput screening for probe and drug development at NCATS are listed

in **Table 1**. A list of all targets, including novel and hypothesized targets used for the creation of the HEAL Library, is provided (**Supplementary S1**), and the details of library composition are discussed in the following section.

Identification of compounds included in the HEAL Library

As a first step towards creating the HEAL Library, we included 430 unique drugs, including experimental or clinical candidates with in-scope indications in pain that were found in DrugBank and NCATS Inxight Drugs [18], and a few drugs with indications for treating addiction, dependency, and overdose. The targets of these FDA-approved drugs are mostly classical pain targets, such as the opioid and dopamine receptors, cannabinoid receptor, and ion channels. Some of the pain and addiction-relevant drugs that do not have a clear protein target were also included in the HEAL Library. Secondly, new drug candidates in clinical trials associated with all HEAL targets were collected from public resources, including Chemble and a patent search. **Table 2** shows some examples of HEAL drug candidates in clinical trials. Compounds without disclosed structures (for example, drugs in Phase 1 clinical trials are not required to be entered to clinicaltrials.gov and often have not released the structure of the clinical candidate) were retrieved and annotated in the target library list for future updates (e.g. HEAL Library 2.0) once these molecules are commercially available.

Finally, we searched for small molecule tool compounds acting against targets in the curated target list from various public resources, like PubChem (pubchem.ncbi.nlm.nih.gov), BindingDB (www.bindingdb.org/), and IUPHAR/BPS (www.guidetopharmacology.org). All pharmacologic mechanisms of action (inhibitors, antagonist, agonist, and modulators) were included in this search, which resulted in an extensive assembly of compounds (over 4,000). To

prioritize the list, all compounds were manually annotated and curated. Structural clustering and molecule properties were evaluated so that compounds of structural representatives and the ones with CNS properties or implicated in pain/addiction pathway as well as neurodegenerative or inflammation diseases were selected. In addition, the availability of identified small molecules from NCATS in-house libraries and commercial vendors was also considered. This process resulted in approximately 2,400 compounds being identified: 850 were from the approved and clinical drugs list; 490 from the CNS drug-likeness list; and 1,033 from tool and probe molecules acting against targets of interest. Many of them were identified initially for various disease areas, such as oncology and infectious diseases, reinforcing the value of creating small molecule probes for every protein in the human genome [19].

Annotation of the HEAL Library

The workflow for creating the HEAL Library is illustrated in **Figure 1**. Like other NCATS annotated screening libraries [11, 20, 21], all compounds and targets were thoroughly curated, annotated, and deposited into NCATS probeDB, which is manageable through a HTS client tool with functionalities of searching, retrieving, updating, and structural analysis. Currently, nearly 3,000 drugs and probe compounds associated with approximately 240 known and hypothesized HEAL-relevant targets have been collected and plated in the HEAL Library (version 1.0). **Figure 2** shows the components of HEAL targets and compounds whose physical samples were plated for HTS (2,800 compounds). Forty-nine percent (49%, 1,375) of the plated library compounds were from the GPCR targets, and 25% (700) were from the ion channels. The least number of compounds was found with transporters (1%) and catalytic receptors (2%), while compounds from the kinases and other enzymes totaled 18% (512). This is generally in

agreement with the HEAL target distribution in the library, also reflecting the current status of

pain and addiction targets under research studies and drug development and the fact that most of

the genome remains 'undrugged'.

Figure 1. Collection of HEAL targets and compounds for quantitative high throughput screening

(qHTS) and AI-driven drug discovery.

Figure 2. Components of HEAL targets and compounds.

We analyzed the physicochemical properties and structural features of the HEAL Library compounds. **Figure 3A** shows the overall distributions of molecular weight (MW), H-bond acceptor (HBA) and H-bond donor (HBD) counts, and TPSA among these subsets of the HEAL compounds (clinical drugs, pain and additive drugs, CNS compounds, probes, and active compounds). Interestingly, the calculated CNS-MPO scores and BBB permeability (**Figure 3B and 3C**) of HEAL compounds are generally "CNS-druglike" and comparable to the subset of pain-related drug compounds. Since the CNS property is a vital consideration for most pain and addictive drug development, the "CNS-likeness" of the HEAL Library also makes it a valuable dataset for a broad HTS screening as well as modeling against CNS-related diseases.

We further compared the structural diversity and chemical space of the HEAL Library with other NCATS annotated libraries (**Figure 3D**). While the HEAL compounds mostly resembled the NPC approved drug collection, as aligned with Lipinki's "rule of 5s" [22], they also demonstrated an expansion into and overlap with the chemical diversity screening library

(Genesis MiniMe collection), as well as an annotated library with a broad collection of pharmacologically active compounds and natural products (NPACT). The diversity of structural chemotypes and chemical space of HEAL compounds, therefore, provided a valuable reservoir as starting points for drug repurposing and rational drug design for novel pain and addition targets.

Figure 3. Physicochemical, CNS and BBB properties of HEAL library compounds.

Compound plating for quantitative high throughput screening (qHTS)

Of 2,980 compounds collected, physical samples of 2,800 compounds were acquired. Except for a few controlled substances, all compounds were purchased from major chemical

suppliers. The same compound acquisition process was used for creating and updating the NPC

library [11]. Once compounds were procured, quality control (QC) was performed by liquid

chromatography-mass spectrometry (LC-MS) at NCATS. Compounds exhibiting less than 95%

purity were purified further to achieve a minimum value of purity. Stock solutions were prepared

at 10mL in DMSO after QC. All compounds were then plated in qHTS 1536-well plate format,

both in acoustic source plates and pin tool plates for qHTS with 1:5 dilution and 7

concentrations, thus enabling a rapid and large-scale screening against a wide range of assays on

NCATS qHTS robot system.

Profiling of HEAL Compounds for cell viability

We conducted a profiling of the HEAL Library using a cell killing assay against the

HeLa and HEK 293 cell lines to evaluate the cytotoxicity of the library's compounds. HeLa is

one of the commonly used human immortalized cell line, and the HEK 293 is a neuronal lineage

line. Both are extensively employed in scientific research for pain and addiction studies. The

HEAL Library compounds were tested with a 7-point dose-response CTG assay in these two cell

lines respectively. As shown in **Figure 4,** 20% and 24% of the library compounds were cytotoxic

to HeLa and HEK 293 cells, respectively. Of these actives, 126 compounds overlapped and

shared similar cytotoxic activity in both cell lines. Notably, the percentage of cytotoxic

compounds within the HEAL Library for these two cell lines was generally lower than other

annotated libraries we previously evaluated [23]. Further analysis showed that the majority of the

cytotoxic compounds were associated with the family of kinase targets, primarily comprising

probes and tool compounds. In contrast, cytotoxic compounds were less prevalent in the GPCR

and other families. This result is anticipated and consistent with the promiscuity score of these drug compounds associated with these targets (**Figure 4D**).

Figure 4. Profiling of HEAL library compounds A) cytotoxic compounds in Hella and HEK293 cells, B) overlap of cytotoxic compounds in HEK 293 and Hela cells, C) actives in each target class, D) promiscuity scores.

Application of HEAL library in qHTS/VS pilot screening

We have utilized the HEAL Library screening against several HEAL targets as part of active collaborations between NCATS scientists and extramural scientists, including GPCRs, ion channels, enzyme cyclase, and a metalloprotease (**Table 1**). Except for glutamate carboxypeptidase II (GCPII), which is a valid target for non-opioid drugs to treat pain (many potent but non-druggable small molecule inhibitors have been reported [24]), very few modulators are known for these HEAL target classes. Therefore, screening the HEAL Library

served as a pilot screen to identify active hits or tool compounds for further assay development and large-scale HTS. We conducted the screens with various biochemical or cell-based assays, including counter assays developed for each target, all of which were miniaturized for qHTS at NCATS. The hits were determined based on the dose-response curve class as well as efficacy [25]. As shown in **Table 3**, the hit rates across these 8 screens ranged from 0.15% - 6%. The hit rate associated with the two cyclase targets (NPR1 at 4.7% and AC1 at 6.0%) and PAC1R (5.7%) were relatively higher. Analysis showed that many of these "primary-assay-hits" may be non-selective or artifacts, suggesting that a secondary assay on their sister targets, for example, NPR2 and AC8, are needed to run and triage the hit list. The GCPII hits were mostly prior art compounds collected in the HEAL Library, indicating the robustness of the MS-LC assay used in the screen [26]. The hits from GALR1 and GPR151 screens showed some activity on both targets and appeared to be more appealing. GPR151 is also known as GALR4, sharing homology with the galanin receptor family [27]. The small molecule agonists of GALR1 and GPR151 identified from the HEAL Library provided novel candidates for further study of these two nociceptors for pain drug development.

In parallel, we also performed virtual screens (VS) for these HEAL targets using an AI-based platform integrated with HEAL compound structures. **Figure 5** shows the heatmap of VS hits in the HEAL Library in comparison with the hits from the experimental qHTS. The VS hit enrichment, defined as the qHTS hits pulled out from the top-ranked VS hitlist, varied from 1.0 (PZ2) to 7.94 (GCPII) (**Table 3**). This was in general agreement with the docking-based VS performance [28]. The notably higher enrichment with GCPII can be attributed to its well-defined active site of crystal structures bound with various inhibitors, whereas for other targets (GPR151, AC1, NPR1), homology models were used in the docking screen. The ion channel

PZ2 is a challenging target for assay development and screen, as well as structure-based VS. The 3D structures of PZ2 and PZ1 have been solved recently, but the small molecule binding site remains elusive [29, 30]. The predicted binding models of these potential hits may provide insight to probe the mechanism of action of ligand binding and activation. Comparison of the qHTS and VS hits in each target family also revealed some interesting findings. The robust and potent qHTS-hits were distributed more randomly across each HEAL compound class, while the VS hits showed target enrichment. For example, the top VS-hits with these four GPCR targets were typically found in the GPCR class of the HEAL Library, and the top VS-hits for the two cyclase NPR1 and AC1 were mostly identified in the enzyme class. These results reflected the similarity of the binding pocket within the target family, which could be used for further qHTS-hit prioritization.

Figure 5. Heatmap of qHTS and VS on 8 HEAL targets.

Outlook and Conclusions

While the HEAL Library plates are used routinely for HEAL project screening at NCATS, a pre-spotted copy of the HEAL Library in 384-well plate format (single concentration, 10mM) can be made available to the research community specifically for meritorious HEAL-related assays through NCATS Division of Preclinical Innovation (DPI) collaborations (https://ncats.nih.gov/research/research-activities/heal/expertise/library). Investigators interested in accessing the HEAL Library must submit a two-page proposal to NCATS DPI for evaluation.

If selected, an overview of the applicant's primary screening results will be shared and discussed with NCATS to determine whether spotted, cherry-picked plates of hits will be provided for counter screen assays. Plated copies of the HEAL Target and Compound Library as well as library cherry picks will only be provided as resources are available. All HEAL Library screening data are expected to be uploaded to PubChem within two years.

AI-driven machine learning (ML) and virtual screening (VS) are playing an ever-increasingly important role in for the field of rational drug discovery and development [31]. With the accumulation of qHTS and VS data on the HEAL Library, we have implemented and evaluated several noteworthy modules within a HEAL-focused AI-platform (**Figure 6**). For example, the 3D structures of all HEAL targets (Structure-DB) are collected from Protein Data Bank (PDB) or modeled by AlphaFold [32]. The binding pockets of representative structures for each target and family are clustered and extracted (Pocket-DB). Systematic docking of all HEAL Library compounds to these binding pockets is performed, and the predicted binding models of each target-drug complex are deposited (Binding-DB). Additionally, ligand-based machine learning models for each target and family are generated based on their qHTS bioassay data (ML-DB). The integration of these HEAL-focused modules, along with a qHTS open data portal [33], provides access to the abundant assay screening data generated from HEAL Library, and facilitates efficient virtual screening of ultra-large chemical databases, such as Enamine's REAL library containing billions of compounds.

Figure 6. An AI-driven platform of HEAL compounds and targets for structure-based drug discovery.

In summary, the creation of the HEAL Library combined with an AI-driven platform provided a valuable resource to establish assay screening and collaborations, strategically advancing drug development programs for pain and addiction. A notable feature of the HEAL Library is the assay-ready pre-spotted plates which have been made available to extramural sites for HEAL-related assays through NCATS. Currently, the HEAL library has been screened against various HEAL targets and other relevant neurodegenerative projects. Similar to other NCATS annotated libraries like NPC and NPACT, the HEAL Library serves as a versatile tool for assay and target validation, exploration of mechanisms of action (MOA), as well as pilot screening of novel candidates. This not only enhances our understanding of the intricate processes involved but also paves the way for further development of probes and drugs aimed at treating pain and addiction.

Acknowledgment

References

1. Stoicea, N., et al., *Current perspectives on the opioid crisis in the US healthcare system: A comprehensive literature review.* Medicine (Baltimore), 2019. **98**(20): p. e15425.
2. National Institute of Drug, A. and H. National Institutes of, *Pain, opioids, and addiction: an urgent problem for doctors and patients.* J Pain Palliat Care Pharmacother, 2007. **21**(4): p. 45-9.
3. Jovey, R.D., *Opioids, pain and addiction - practical strategies.* Br J Pain, 2012. **6**(1): p. 36-42.
4. Chaplan, S.R., I.W. Eckert, and N.I. Carruthers, *Drug Discovery and Development for Pain*, in *Translational Pain Research: From Mouse to Man*, L. Kruger and A.R. Light, Editors. 2010: Boca Raton (FL).
5. Obeng, S., et al., *Novel Approaches, Drug Candidates, and Targets in Pain Drug Discovery.* J Med Chem, 2021. **64**(10): p. 6523-6548.
6. Aronowitz, S.V., P. Compton, and H.D. Schmidt, *Innovative Approaches to Educating Future Clinicians about Opioids, Pain, Addiction and Health Policy.* Pain Manag Nurs, 2021. **22**(1): p. 11-14.
7. Henningfield, J.E., et al., *The nexus of opioids, pain, and addiction: Challenges and solutions.* Prev Med, 2019. **128**: p. 105852.
8. Collins, F.S., W.J. Koroshetz, and N.D. Volkow, *Helping to End Addiction Over the Long-term: The Research Plan for the NIH HEAL Initiative.* JAMA, 2018. **320**(2): p. 129-130.
9. Baker, R.G., W.J. Koroshetz, and N.D. Volkow, *The Helping to End Addiction Long-term (HEAL) Initiative of the National Institutes of Health.* JAMA, 2021. **326**(11): p. 1005-1006.
10. Coussens, N.P., et al., *The Opioid Crisis and the Future of Addiction and Pain Therapeutics.* J Pharmacol Exp Ther, 2019. **371**(2): p. 396-408.
11. Huang, R., et al., *The NCGC pharmaceutical collection: a comprehensive resource of clinically approved drugs enabling repurposing and chemical genomics.* Sci Transl Med, 2011. **3**(80): p. 80ps16.
12. Corsello, S.M., et al., *The Drug Repurposing Hub: a next-generation drug library and information resource.* Nat Med, 2017. **23**(4): p. 405-408.

13. Fernandez-Rodriguez, J., et al., *A High-Throughput Screening Platform Identifies Novel Combination Treatments for Malignant Peripheral Nerve Sheath Tumors.* Mol Cancer Ther, 2022. **21**(7): p. 1246-1258.

14. Woolf, C.J., *Capturing Novel Non-opioid Pain Targets.* Biol Psychiatry, 2020. **87**(1): p. 74-81.

15. Kumari, P., et al., *Targeting G protein coupled receptors for alleviating neuropathic pain.* Prog Mol Biol Transl Sci, 2022. **193**(1): p. 99-117.

16. Braden, K., et al., *Small molecule targeting NaV1.7 via inhibition of the CRMP2-Ubc9 interaction reduces and prevents pain chronification in a mouse model of oxaliplatin-induced neuropathic pain.* Neurobiol Pain, 2022. **11**: p. 100082.

17. Serafini, R.A. and V. Zachariou, *Opioid-galanin receptor heteromers differentiate the dopaminergic effects of morphine and methadone.* J Clin Invest, 2019. **129**(7): p. 2653-2654.

18. Siramshetty, V.B., et al., *NCATS Inxight Drugs: a comprehensive and curated portal for translational research.* Nucleic Acids Res, 2022. **50**(D1): p. D1307-D1316.

19. Müller, S., et al., *Target 2035-update on the quest for a probe for every protein.* Rsc Medicinal Chemistry, 2022. **13**(1): p. 13-21.

20. Huang, R., et al., *The NCATS Pharmaceutical Collection: a 10-year update.* Drug Discov Today, 2019. **24**(12): p. 2341-2349.

21. Huang, R., et al., *The NCATS BioPlanet - An Integrated Platform for Exploring the Universe of Cellular Signaling Pathways for Toxicology, Systems Biology, and Chemical Genomics.* Front Pharmacol, 2019. **10**: p. 445.

22. Lipinski, C.A., et al., *Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings.* Adv Drug Deliv Rev, 2001. **46**(1-3): p. 3-26.

23. Lee, O.W., et al., *Cytotoxic Profiling of Annotated and Diverse Chemical Libraries Using Quantitative High-Throughput Screening.* SLAS Discov, 2020. **25**(1): p. 9-20.

24. Pavlicek, J., J. Ptacek, and C. Barinka, *Glutamate carboxypeptidase II: an overview of structural studies and their importance for structure-based drug design and deciphering the reaction mechanism of the enzyme.* Curr Med Chem, 2012. **19**(9): p. 1300-9.

25. Inglese, J., et al., *Quantitative high-throughput screening: a titration-based approach that efficiently identifies biological activities in large chemical libraries.* Proc Natl Acad Sci U S A, 2006. **103**(31): p. 11473-8.

26. Hoxie, N.Q., Y.; Kales, S.C.; Schneider, R.; Hu, X.; Dalal, A.; Ford-Scheimer, S. L.; Wiseman,R.; Tsukamoto, T.; Wei, H.; Slusher, B. S.; Janiszewski, J. S.; Hall, M. D., *Development of a High-Throughput Dual Stream LC/MS/MS method to screen for inhibitors of Glutamate Carboxypeptidase II (GCPII).* Rapid Communications in Mass Spectrometry, 2023 (in press).

27. Liu, Z., et al., *Evolution of galanin receptor genes: insights from the deuterostome genomes.* J Biomol Struct Dyn, 2010. **28**(1): p. 97-106.

28. Cheng, T., et al., *Comparative assessment of scoring functions on a diverse test set.* J Chem Inf Model, 2009. **49**(4): p. 1079-93.

29. Wang, L., et al., *Structure and mechanogating of the mammalian tactile channel PIEZO2.* Nature, 2019. **573**(7773): p. 225-229.

30.   Lacroix, J.J., W.M. Botello-Smith, and Y. Luo, *Probing the gating mechanism of the mechanosensitive channel Piezo1 with the small molecule Yoda1.* Nat Commun, 2018. **9**(1): p. 2029.

31.   Vemula, D., et al., *CADD, AI and ML in drug discovery: A comprehensive review.* Eur J Pharm Sci, 2023. **181**: p. 106324.

32.   Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* Nature, 2021. **596**(7873): p. 583-589.

33.   Brimacombe, K.R., et al., *An OpenData portal to share COVID-19 drug repurposing data in real time.* bioRxiv, 2020.

Table 1. HEAL targets for pain and addiction.

HEAL Targets	Background	MOA	Status of Research
GALR1	Galanin receptor 1 (GALR1) forms heteromers with mu-opioid receptors (MOR) in the ventral tegmental area (VTA) that modulate dopamine release. GALR1 ligands block MOR-mediated signaling by allosteric modulations within the MOR-GALR1 heteromers.	Targeting MOR-GALR1 heteromers to treat OUD while sparing the analgesic effect of opioids	Structure of GAL1R bound with Gal1 peptide solved. No specific small molecules against GALR1 reported. HTS screening at NCATS
GPR151	Gpr151 is an orphan GPCR almost exclusively in the medial habenula (mHb), which is co-localized with μ opioid receptors. GPR151 modulates opioid receptor signaling in mHb.	Modulating opioid signaling in mHb by targeting GPR151 signaling as a novel treatment for opioid dependence in humans	No small molecules reported. HTS at NCATS
RXFP3	Relaxin-3 is a neuropeptide that is highly expressed in the nucleus incertus, a brain region that mostly consists of GABAergic projection neurons which innervate a broad range of forebrain regions involved in behavioral activation and the response to stress.	Small molecule agonists are a potential treatment for pain, anxiety, depression, and addiction.	A series of small molecule agonists reported. Probe compounds for family targets RXFP1 and RXFP2 have been identified.
PAC1R	The pituitary adenylate cyclase-activating polypeptide type I receptor (PAC1R) is widely distributed in the central neural system and peripheral organs. Abnormal activation of the receptor mediates trigeminovascular activation and sensitization is highly related to migraine.	PAC1R a potential therapeutic target for discovery of therapeutic drugs for neuronal disorders. Antagonist, overcome the CNS side effects	Structure bound with ligand PACUP peptide solved. A series of small molecule antagonist reported.
PIEZO2	PIEZO2 is a stretch-gated cell membrane ion channel expressed in sensory neurons, nociceptors, and mechanosensitive epithelial cells. Loss-of-function produces loss of proprioception, gentle touch, and mechanical allodynia.	Targeting PIEZO2 provides novel ways to treat tactile pain and chronic itch while preserving normal pain and temperature sensation. PIEZO2 agonists could ameliorate chronic itch and serve as powerful molecular tools	Target of ion channel structure available; no agonist/antagonist reported. A series of small molecule agonists identified for PIEZO1.
CRMP2	NaV1.7 and related ion channel targets have been challenging to drug efficiently and specifically. CRMP2 is an essential trafficking protein for NaV1.7, and SUMOylation of CRMP2 is required for membrane targeting.	Small molecule drug-like inhibitors provide an alternative strategy for blocking NaV1.7 function and reducing pain by preventing its trafficking to the neuronal plasma membrane	Structure available, no inhibitor reported.

NPR1	There are many diseases where both pain and itch coexist, producing major debilitating consequences. Opioids can elicit strong pruriceptive responses. Alternatives to opioids for treatment of pain would reduce the pruritic burden caused by their use.	Although itch is often considered a side-effect of a disease, it is difficult to treat effectively, reduces quality of life for patients, and is a major unmet health concern. Identify hNpr1 antagonist with higher potencies and greater selectivity that will be good candidates for preclinical testing	Small molecule antagonist reported but none in development
AC1	AC1 is highly expressed in neuronal tissues associated with pain processing and neuronal plasticity. Knockout animals reveal almost a complete loss in pain responsiveness in both inflammatory and neuropathic pain models.	Develop and characterize AC1-selective small molecule inhibitors as a novel, non-opioid treatment for chronic pain. Breakthrough approach of translational science to advance new treatments for the opioid crisis	Small molecule inhibitors reported but none in development
GCPII	Excess glutamatergic transmission is associated with acute and chronic pain caused by inflammation or brain injury. Drugs designed to target glutamatergic transmission have shown efficacy in animal models.	GCPII inhibitors that penetrate the brain are a major unrealized promise for the treatment of pain associated with glutamate excitotoxicity in that they will indirectly reduce glutamate neurotransmission by reducing glutamate levels without interfering with glutamatergic transmission	Structures and small molecule inhibitors extensively studied but none in further drug development

Table 2. Representatives of HEAL library targets and compounds.

HEAL ID	Compound	Condition	Deve. Status	Primary Target	Mechanism Type	Structure
NCGC00015476	MRS-1523	Pain	Probe	ADOR-A	Antagonist	
NCGC00689784	L-759633	Pain	Probe	CB2	Agonist	
NCGC00378701	STX209	Pain	Phase III	GABAB	Agonist	
NCGC00686991	MK-7622	Pain	Clinical	mAChR	PAM	
NCGC00685432	SB-612111	Pain	Probe	NOPR	Antagonist	

NCGC00024796	GW-9508	Pain	Probe	GPR120	Agonist
NCGC00370852	DSP-2230	Pain	Phase II	Nav1.7	Blocker
NCGC00371083	AZD-9056	Pain	Phase II	P2X	Antagonist
NCGC00379127	AMG-9810	Pain	Probe	TRPV	Antagonist
NCGC00025240	ARN-272	Pain	Probe	FAAH	Inhibitor
NCGC00344511	MFZ 10-7	Addiction	Probe	mGluR	NAM

NCGC00371069	CP-154526	Addiction	Investigation	CRF1	Antagonist	
NCGC00183870	LY-2456302	Addiction	Phase II	MOR	Antagonist	
NCGC00025250	Nepicastat	Addiction	Phase II	Dopamine beta-hydroxylase	Inhibitor	

Table 3. HEAL library screening against 8 targets using qHTS and VS. Enrichment is defined as the HTS hits pulled out from the top-ranked 5% of VS hitlist.

Target	HTS		VS	
	hits	hit rate (%)	hits	enrichment
GALR1	4	0.14	1	4.69
GPR151	18	0.64	5	5.21
RXFP3	77	2.73	21	5.12
PAC1R	159	5.65	25	2.95
PZ2	19	0.67	1	1.00
GCPII	26	0.92	11	7.94
NPR1	132	4.69	17	2.42
AC1	169	6.00	15	1.67