# Microbiota from young mice counteracts susceptibility to age-related gout through modulating butyric acid levels in aged mice

Ning Song [a,1], Hang Gao [b,1], Jianhao Li [a], Yi Liu [b], Mingze Wang [a], Zhiming Ma [c], Naisheng Zhang [a, *], Wenlong Zhang [a, *]

[a] Department of Clinical Veterinary Medicine, College of Veterinary Medicine, Jilin University, Changchun, 130062, China

[b] Department of Bone and Joint Surgery, No1 Hospital of Jilin University, Changchun, 130021, China

[c] Department of Gastrointestinal Nutrition and Hernia Surgery, The Second Hospital of Jilin University Changchun 130041, China

[1] These authors contributed equally to this work.

* Corresponding author. E-mail addresses: zhangns@jlu.edu.cn (N. Zhang); zwenlong123@126.com (W. Zhang)

**Abstract**

Gout is a prevalent form of inflammatory arthritis that occurs due to high levels of uric acid in the blood leading to the formation of urate crystals in and around the joints, particularly affecting the elderly. Recent research has provided evidence of distinct differences in the gut microbiota of patients with gout and hyperuricemia when compared to healthy individuals. However, the link between gut microbiota and age-related gout remained underexplored. Our study found that gut microbiota plays a crucial role in determining susceptibility to age-related gout. Specifically, we observed that age-related gut microbiota regulated the activation of the NLRP3 inflammasome pathway and modulated uric acid metabolism. More scrutiny highlighted the positive impact of "younger" microbiota on the gut microbiota structure of old or aged mice, enhancing butanoate metabolism and butyric acid content. Experimentation with butyrate supplementation indicated that butyric acid exerts a dual effect, inhibiting inflammation in acute gout and reducing serum uric acid levels. These insights emphasize the potential of gut microbiome

rejuvenation in mitigating senile gout, unraveling the intricate dynamics between microbiota, aging, and gout. It potentially serves as a therapeutic target for senile gout-related conditions.

**Key words** Senile gout; Gut microbiota; Hyperuricemia; Age; Butanoate metabolism

**Introduction**

Gout, the most common inflammatory arthritis in elderly individuals, results from the deposition of monosodium urate crystals in articular and nonarticular structures[1], particularly among individuals aged 75-84 years, and the occurrence rate in this population can reach 4%[2]. The development of gout is primarily attributed to the significant risk posed by a high serum urate concentration[2]. Pathological hyperuricemia is defined by a serum urate concentration exceeding 408 μmol/L [6.8 mg/dL], which forms monosodium urate crystals in vitro at physiological pH and temperature[3]. The reason why the elderly population commonly experiences gout is complicated by the fact that the population is also ageing, and the treatment of this disease is often intricate due to the presence of comorbidities and medications prescribed for concurrent conditions[2]. Although the basic principles for the prevention and treatment for gout remain the same across different age groups, elderly individuals often exhibit lower tolerance for medication dosages, types, side effects, and surgical procedures due to their physiological factors. Moreover, despite the increasing severity of gout among elderly individuals, research on this issue remains scarce.

The gut microbiota plays essential roles in regulating energy and metabolism, as revealed by recent studies[4, 5], which have shown that individuals with hyperuricemia and gout exhibit dysbiosis of the gut microbiota[6, 7]. Moreover, because the gut microbiota can directly participate in the metabolism of purines and uric acid[8], it may play a crucial role in gout and hyperuricemia development. Although numerous studies have examined

the relationship between the microbiome and early life stages, the impact of the microbiome's on ageing and frailty in later life needs to be explored. Furthermore, studies have shown that the structure of the gut microbiota undergoes a gradual "ageing" process with advancing age, and this process is characterized by a decrease in the microbial diversity(9, 10). In addition, transplantation of a young microbiota can improve central nervous system inflammation and retinal inflammation in aged mice(11), and counteract age-related behavioral deficits(12). We hypothesize that the high prevalence of gout in the elderly population may be closely related to its "ageing" gut microbiota. The association between the "ageing" gut microbiota and gout in elderly individuals has not been reported. Hence, we conducted a study to elucidate the influence of the ageing gut microbiota on the occurrence and progression of gout in elderly individuals.

In this study, we first tested the sensitivity to monosodium urate (MSU) in different age groups and conducted a microbiota clearance (Abx) on mice of different age groups to assess their sensitivity to MSU again. At the same time, we also tested their serum uric acid levels. We found that sensitivity to MSU increased with age, but changed after clearing the gut microbiota in terms of sensitivity to MSU and serum uric acid levels. Then we performed cross-age fecal microbiota transplantation (FMT) and subsequently stimulated the mice with MSU with the aim of how mice belonging to different age groups exhibit sensitivity to MSU after undergoing FMT. Because hyperuricemia is a necessary physiological factor for gout, we also evaluated the expression levels of uric acid-producing enzymes and uric acid transport proteins in the mice. Surprisingly, transplantation of the gut microbiota from aged mice into young mice, significantly increased their sensitivity to MSU. Conversely, the transplantation of the gut microbiota from young mice into aged mice, significantly decreased their sensitivity to MSU. These findings suggest that the gut microbiota of older individuals plays a promoting role in the occurrence and progression of gout. Moreover, an analysis of the serum uric acid levels of

the mice after cross-age FMT yielded similar results. To investigate the underlying mechanisms of how gut microbiota influences gout and hyperuricemia, we performed 16S rDNA sequencing and untargeted metabolomics analysis of fecal samples. We then, observed a significant increase in the abundance of *Bifidobacterium* and *Akkermansia* in the gut microbiota of young mice and old or aged mice after transplantation of the gut microbiota from young mice. To further investigate the potential biological pathways affected by the gut microbiota, we performed functional metagenomic analysis using Tax4FUN software and found that butanoate metabolism was more robust in young mice than in aged mice. Furthermore, we also observed that transplantation of the gut microbiota from young mice to aged mice enhanced the butanoate metabolism of the recipient mice. Due to the limitations of untargeted metabolomics, we did not observe any differences in the levels of butyric acid among the different groups. However, the pathway data obtained by fecal untargeted metabolomics also yielded similar results. Based on the above-described results, we hypothesize that butyrate may play a significant role in these processes. Excitingly, the results from a short-chain fatty acid (SCFA) analysis support our hypothesis. Furthermore, a supplementation experiment using butyrate revealed, that the results aligned well with the cross-age FMT findings, which suggests that transplantation of the gut microbiota from young mice into aged mice can effectively prevent gout and hyperuricemia, and that butyrate is likely the critical factor playing a crucial role. Our research findings demonstrate the potential of young gut microbiota in preventing gout in elderly individuals and provide new insights and perspectives for the prevention and treatment of gout in elderly individuals.

**Results**

**Gout susceptibility increases with age, related to gut microbiota**

In this study, we used the male C57BL/6 mice belonging to three age groups: young (~3 months), old (~18 months) and aged (~24-months) mice (Fig. 1a). To investigate the impact of age on gout, we used a mouse model in

which subcutaneous injections of MSU crystals were administered to the dorsal aspect of the hind paws (13-15). The result showed that the old group exhibited a significant increase in footpad swelling compared with the young group, and the levels of IL-1β was significantly elevated in the Old and Aged groups (Fig. 1b, c). A prominent correlation was found between the occurrence of gout and the concentration of serum uric acid, which exhibits an upwards trend with advancing age(2). Thus, we also measured the serum uric acid level, we found uric acid levels significantly increase with age (Fig. 1d). There are reports indicating a close relationship between gut microbiota and gout(6). Therefore, we simultaneously tested the gut microbiota of mice at different ages. The results showed that as age increases, the ASVs (Amplicon Sequence Variants) showed a declining trend (Fig. 1e). Moreover, the principal coordinates analysis (PCoA) results revealed distinctive differences in the phylogenetic community structures between these groups (Fig. 1f). Then we carried out antibiotics (ABX) cocktail on mice of different age groups, found changes in the footpad swelling, IL-1β and discovered no differences in the level of serum uric acid (Fig. 1g-i). These data highlight indicates that aging-associated changes in the gut microbiota exacerbate gout attacks.

**Aged-to-young FMT worsens acute gout, whereas young-to-aged FMT reduced this disease**

Based on the above results, we conducted fecal microbiota exchanges between male C57BL/6 mice also belonging to three age groups: young (~3 months), old (~18 months), and aged (~24-months) mice. The experimental design and timeline are presented in Figure 2a. To investigate the impact of cross-age FMT on gout, we used a gout mouse model as aforementioned. The results of supplementary figure 1a showed that young mice transplanted with fecal microbiota from mice in the old or aged group (Young+Old or Young+Aged) exhibited a significant increase in footpad swelling compared with the control group (Young+PBS). However, no significant difference in footpad swelling was found between old or aged mice transplanted with fecal

microbiota from mice in the young groups (Old+Young and Aged+Young) and the control groups (Old+PBS and Aged+PBS) (supplementary figure 1a). Subsequently, we conducted a comparative analysis of haematoxylin & eosin (H&E)-stained tissue sections (scar, 1000µm), derived from mice with cross-age FMT and MSU-induced acute gout. We found that the Young+Old and Young+Aged group showed more inflammatory cell infiltration in the subcutaneous soft tissues than the Young+PBS group, whereas the Old+Young and Aged+Young groups showed less inflammatory cell infiltration (Fig. 2b). The inflammatory factors in the foot tissue of the mice with cross-age FMT were then examined. The levels of IL-1β, IL-6, and TNFα were significantly elevated in the Young+Aged group and significantly lower in the Aged+Young group compared with the Aged+PBS group (Fig. 2c-e). Meanwhile, to investigate the impact of cross-age FMT on MSU-induced inflammation, we utilized an animal model of C57BL/6 mice administered an intraperitoneal injection of 2.5 mg of MSU (16). Six hours after MSU intraperitoneal injection, we washed the mouse peritoneal cavity with sterile PBS and collected the peritoneal fluid and peritoneal cells separately. Simultaneously, we also examined the levels of inflammatory cytokines in mouse serum. Consistent with the acute gout model, the Young+Old or Young+Aged group exhibited more pronounced activation of IL-1β, IL-6, and TNF-α in peritoneal fluid (Fig. 2f-h) and serum (Supplementary figure 1b-c). Although no differences in the IL-6 levels of serum and peritoneal fluid was found between the Old+PBS and Old+Young groups or between the Aged+Young and Aged+PBS groups (Supplementary figure 1c and Fig. 2g), a significant reduction in IL-1β and TNF-α was observed (Supplementary figure 1b, d and Fig. 2f, h). These findings suggest that the increased susceptibility to gout in the elderly may closely related to "aging" gut microbiota.

**"Younger" gut microbiota suppresses NLRP3 inflammasome pathway, "aging" gut microbiota promotes**

The pathogenesis of acute gout has been primarily linked to the activation

of proinflammatory pathways, notably NLRP3, and this activation instigates a
surge in the production of the inflammatory cytokines, which further
underscores their pivotal roles in the progression of the disease's (17). We
primarily examined proteins of foot tissue and peritoneal cells associated with
the NLRP3 inflammasome pathway. Interestingly, we discovered that the
Young+Aged group demonstrated more pronounced activation of NLRP3,
Pro-Caspase-1, Caspase-1, and IL-1β compared with the control group (Fig.
3a-b). However, the Old+Young and Aged+Young groups showed lower
protein levels of Caspase-1 and IL-1β compared with the control group (Fig.
3c-f), which suggests that the gut microbiota of elderly mice may make them
more sensitive to MSU stimulation, whereas the gut microbiota of young mice
can effectively resist the MSU stimulation, and inhibit Caspase-1 cleavage,
and IL-1β secretion. Meanwhile, the NLRP3 inflammasome pathway also was
investigated in peritoneal cells, and similar results were obtained in the gout
model. The fecal microbiota from mice in the old or aged group exacerbated
the activation of Pro- Caspase -1, Caspase-1, Pro-IL-1β, and IL-1β in
peritoneal cells, increasing the production of IL-1β (supplementary figure 2a-b).
In contrast, the fecal microbiota from mice in the young group inhibited the
cleavage of Caspase-1 and the secretion of IL-1β (supplementary figure 2c-f).

**Beneficial effects of FMT from young to aged mice on uric acid**
**metabolism**

Serum samples were collected from the the different groups, and their
serum uric acid levels were initially assessed. Surprisingly, we observed an
elevation in the average serum uric acid levels in the Young+Old or
Young+Aged group, whereas the Old+Young or Aged+Young group exhibited
decreases in the serum uric acid levels compared with the same-age control
group (Fig. 4a). Our study, showed that compared with those of the same-age
control group, the serum levels of AST and ALT were elevated in the
Young+Old and Young+Aged groups, although the differences were not
significant, whereas the Aged+Young groups exhibited modest decreases in

these levels (supplementary figure 3a-b). Furthermore, similar trends were found for indicators (Crea and BUN) of renal function (supplementary figure 3c-d). Hyperuricemia is attributed to increased uric acid synthesis and decreased uric acid excretion in the body. We first measured the activity of enzymes involved in uric acid synthesis, namely adenosine deaminase (ADA), guanine deaminase (GDA), and xanthine dehydrogenase (XOD), in the liver and that of (XOD) in the kidney. The activity of ADA, GDA, and XOD in the liver, and the activity of XOD in the kidney of the Young+Old and Young+Aged groups did not significantly differ from those of the same-age control groups (Fig. 4b-e). The fecal microbiota from mice in the young group induced a notable decrease in the activity of enzymes related to uric acid synthesis, and the most prominent reductions were found for ADA and XOD activity in the liver and XOD activity in the kidney (Fig. 4b, d and e). We then examined the mRNA levels of relevant proteins involved in uric acid transport. We initially measured the renal injury marker KIM-1, and observed that the fecal microbiota from mice in the young group contributed to attenuating the mRNA expression levels of this marker compared with the same-age control group (Fig. 4f). The urate transporter URAT1, which is responsible for uric acid reabsorption, exhibited a similar trend (Fig. 4g). However, cross-age FMT did not significantly impact the mRNA expression of another uric acid reabsorption protein, GLUT9 (Fig. 4h). The Young+Aged group showed lower mRNA expression levels of the uric acid excretion proteins OAT1 and OAT3 compared with the Young+PBS group (Fig. 4i-j). Although the mRNA expression levels of OAT1 and OAT3 did not show significant differences, a significant increasing trend was found for the other uric acid excretion protein, ABCG2, in aged mice transplanted with fecal microbiota from mice in the young group exhibited a significant increasing trend (Fig. 4k). Because ABCG2 is also expressed in the intestine, we examined its mRNA expression levels in the colon and found that the fecal microbiota from mice in the aged group could inhibit its expression (supplementary figure 3e). Similar trends were

found for the mRNA expression levels of mice in ZO-1 and JAMA in the colon of the Young+Aged group (supplementary figure 3g-h). No significant alterations in the colonic Occludin mRNA expression levels were observed among all the groups (supplementary figure 3f). However, the mRNA expression levels of JAMA in the Old+Young and Aged+Young groups were significantly different from those in the same-age control groups (supplementary figure 3h). These results suggests that the fecal microbiota from mice in the old or aged group leads to insufficient uric acid excretion, whereas the fecal microbiota from mice in the young group promotes uric acid elimination, inhibits reabsorption, and may contribute to the integrity of the intestinal barrier structure and the maintenance of normal physiological function.

**Modifications in the gut microbiota composition following cross-age FMT**

To characterize the age-related alterations in the gut microbiota and assess changes following transplantation, we conducted 16S rDNA amplicon sequencing of all the mice at the endpoint. We focused on investigating the gut microbiota of mice after cross-age FMT. We detected the relative abundance of the top 10 phylum levels in each group (Fig. 5a) and found no significant differences in the abundance of *Bacteroidetes* and *Firmicutes* in the transplanted groups compared with that of their respective control groups (supplementary figure 4a-b). No significant differences were found in the the ratio of *Firmicutes* to *Bacteroidetes* (supplementary figure 4c). The species richness, evenness, and rarity are fundamental components of biodiversity and are commonly quantified using indices such as Chao1, observed otus, Shannon, and Simpson indices. Although no significant differences in the Simpson index were found among these groups (supplementary figure 4e), and the indices of the Old+Young and Aged+Young groups were not significantly different from those of the Old+PBS and Aged+PBS groups, respectively, the Chao 1 (Fig. 5b), observed_otus (Fig. 5c), and Shannon

(supplementary figure 4d) indices of the Young+Aged group were significantly

lower than those of the Young+PBS group. These findings are in line with other

studies(18), indicating declines in the richness and diversity of the gut

microbiota during aging. Moreover, the principal coordinates analysis (PCoA)

results revealed distinctive differences in the phylogenetic community

structures between these groups. We showed that the Young+Old and

Young+Aged groups tended to be closer to the Old+PBS and Aged+PBS

groups, and the Old+Young and Aged+Young groups tended to be closer to

the Young+PBS group (Fig. 5d). We then compared the abundance of the top

15 genus level among all the groups (supplementary figure 4f). A Metastats

analysis, found that the abundance of *Bifidobacterium* significantly differed

between the Young+PBS group and the Young+Old and Young+Aged groups,

and that the abundance of *Lachnoclostridium* showed an increasing trend in

the Old+Young and Aged+Young groups (Fig. 5e). The ternary plot showed

that *Akkermansia* appears to be the dominant species in the Young+PBS,

Old+Young and Aged+Young group (Fig. 5f). According to previous

reports(19-22) on *Bifidobacterium* and *Akkermansia*, we hypothesize that

these genera or their metabolites may play a key role in resistance to gout and

hyperuricemia. Based on recent research findings(23) and a KEGG analysis of

bacterial community functions, we discovered that butanoate metabolism was

more robust in the Young+PBS group than in the Old+PBS and Aged+PBS

groups (Table 2a-b). However, no similar phenomenon was observed in the

Young+Old and Young+Aged groups compared with the Young+PBS group

(Table 2c-d). More interestingly, the Old+Young and Aged+Young groups

showed stronger butanoate metabolism than the Old+PBS and Aged+PBS

groups, respectively (Table 2e-f). Considering the results from the analysis

modifications in the gut microbiota composition after cross-age FMT, both

*Bifidobacterium* and *Akkermansia* metabolites include SCFAs. These findings

indicate that butyric acid derived from the young microbiome may be the

critical element responsible for controlling gout. Numerous studies have

reported the beneficial effects of butyric acid, but further research on the

relationship between butyric acid and gout is needed.

**Changes in fecal microbiota metabolism and pathways after cross-age**

**FMT**

We also performed an untargeted metabolomics analysis of samples

collected after cross-age FMT to investigate the research question. Chemical

classification of the metabolites identified in this study was performed, and a

pie chart was generated to reflect the distribution and numbers of metabolites

in each category. The pie chart for Class I metabolite classification and KEGG

pathway annotation is shown in supplementary figure 5a. Among the 42

samples analysed in this study, a total of 1009 metabolites were identified in

the positive ion mode, and 522 metabolites were identified in the negative ion

mode among the 42 samples analyzed in this project. The volcano plot

provides an overview of the distribution of differentially expressed metabolites

(supplementary figure 5b). We also used the KEGG database for metabolic

analysis and network research of the identified biological entities. The

enrichment results are based on KEGG pathway units, and hypergeometric

testing was performed to identify pathways enriched in differentially abundant

metabolites compared with all specified metabolite backgrounds. Through

pathway enrichment, we discerned and elucidated the principal biochemical

metabolic pathways and signal transduction pathways in which the differentialy

abundant metabolites actively participate. The metabolites showing differential

abundance between the Young+PBS and Old+PBS groups were enriched in

the secondary bile acid biosynthesis pathway (supplementary figure 6a).

Interestingly, the metabolites showing differential abundance between the

Young+PBS and Aged+PBS groups were also enriched in the butanoate

metabolism pathway (supplementary figure 6b). The same phenomenon was

found from the comparisons of the Young+PBS and Young+Old groups,

Young+PBS and Young+Aged groups, Old+PBS and Old+Young groups, and

Aged+Young and Aged+PBS groups: the metabolites identified as showing

differential abundance between the groups metabolites were enriched in the

butanoate metabolism pathway (Fig. 6a-d). Furthermore, we found that the

metabolites showing differential abundance between the Old+PBS and

Old+Young groups and between the Aged+Young and Aged+PBS groups were

enriched in the secondary bile acid biosynthesis pathway (Fig. 6c-d). Due to

the limitations of untargeted metabolomics, we did not observe any differences

in butyric acid. However, the results were consistent with the predicted

functional capabilities of microbial communities and corresponded to

butanoate metabolism. Based on the abovementioned results, butyrate may

play an important role.

**Butyrate supplementation inhibits gout and MSU-induced peritonitis**

Based on the predictions of the functional capabilities of microbial

communities and untargeted metabolomics results, we hypothesized that

butyrate might play a key role. Because butyric acid is a short-chain fatty acid,

we measured the levels of SCFAs (nk/g) in the Young+PBS, Aged+PBS, and

Aged+Young groups. Subsequently, we observed that the butyric acid content

in the faeces of Young+PBS group was significantly higher than that in the

Aged+PBS group. Simultaneously, the butyric acid content in the faeces of

mice in the Aged+Young was also significantly higher than that in the

Aged+PBS (Fig. 7a). 2-Methylvalerate and 3-Methylvalerate were not detected,

and 4-Methylvaleric was detected in only a subset of samples, thus, these

results are not presented. The difference analysis of the detected short chain

fatty acids is shown in supplementary figure 7a-h. Therefore, mice were

supplemented with butyrate for 14 days and then subjected to intraperitoneal

injections of MSU into the dorsal aspect of the hind paws to observe its

anti-inflammatory effects in acute gout. The experimental results showed that

butyrate exerted significant anti-inflammatory effects in the acute gout model

with inflammation induced by intraperitoneal injection of MSU. Mice with gout

model, supplemented with butyrate showed significant reductions in the foot

thickness ratio (Fig. 7b) and the levels of inflammatory factors (IL-1β, IL-6, and

TNF-α) in their foot tissue (Fig. 7c-e). Pathological sections revealed that the
foot tissue of mice supplemented with butyrate exhibited less inflammatory cell
infiltration than that of the control group (Fig. 7f). Moreover, after
supplementation with butyrate, mice that received an intraperitoneal injection
of MSU showed significant decreases in the levels of inflammatory factors
(IL-1β, IL-6, and TNF-α) in both serum (supplementary figure 7i-k) and
peritoneal fluid (Fig. 7g-i) compared with those in the control group.
Subsequently, we also analyzed the expression of proteins associated with the
NLRP3 inflammasome pathway. Although no significant change in the
expression of NLRP3 after supplementation with butyrate was observed in
both the gout and peritonitis models, supplementation with butyrate effectively
suppressed the production of Caspase-1 and IL-1β (Fig. 7j and supplementary
figure 7l). The anti-inflammatory effects of butyrate have been widely reported,
but limited studies have investigated its specific on gout. Here we demonstrate
that butyrate effectively inhibits inflammation induced by MSU stimulation.

**Serum uric acid-lowering effect of butyrate in old or aged mice**

This has been proved in figure 1d, that the serum uric acid levels in mice
of different age groups exhibited increase in their serum uric acid levels with
increase age. There is scarce research on the impact of butyrate on serum uric
acid levels, and the effects of butyrate on hepatic uric acid production or renal
uric acid excretion have not been investigated. Additionally, the impact of
butyrate on the serum uric acid levels in aged mice remains unexplored. Aged
mice were supplemented with butyrate for 14 days, and we found that butyrate
supplementation significantly reduced the serum uric acid levels in
18-month-old and 24-month-old mice (Fig. 8a). In this regard, we also
assessed the activity of enzymes implicated in uric acid synthesis, namely,
ADA, GDA, and XOD in the liver, as well as that of XOD in the kidney. In
addition to identifying that butyrate can reduce the activity of ADA in
24-month-old mice (Aged) (Fig. 8b), and supplementation with butyrate did not
have significant effects on the activities of enzymes involved in uric acid

production in both 18-month-old and 24-month-old mice (Fig. 8c-e). Butyrate
may not necessarily lower the serum uric acid levels by inhibiting uric acid
production. Because butyrate might exert its influence on serum uric acid
levels by affecting uric acid transport, we subsequently examined the
expression of relevant transporter mRNA in mouse kidneys and the expression
of the colonic ABCG2 transporter that facilitates uric acid excretion. The
expression of the renal injury marker KIM-1 was significantly decreased in
mice supplemented with butyrate (Fig. 8f). In contrast to the results obtained
cross-age FMT, no significant difference in the expression of URAT1 was
observed (Fig. 8g). However, the expression of GLUT9 was significantly
decreased in 24-month-old mice after butyrate supplementation, whereas no
significant difference was observed in the 18-month-old mice (Fig. 8h). We
also found that the supplementation of old or aged mice with butyrate resulted
in an increased mRNA expression level of uric acid transporters involved in
uric acid excretion and significantly increased the expression of OAT1 and
OAT3 (Fig. 8i-j). Although no significant difference in the expression of ABCG2
was detected in old mice, a significant increased in its expression was
observed in aged mice after supplementation with butyrate (Fig. 8k).
Meanwhile, the mRNA expression of ABCG2 in the colons was significantly
increased in old or aged mice after supplementation with butyrate (Fig. 8l).
Moreover, supplementation with butyrate significantly increased the mRNA
expression of ZO-1, Occludin, and JAMA in the colons of mice (supplementary
figure 8a-c). Based on the abovementioned results, we found that
supplementation with butyrate promotes uric acid excretion and improves the
intestinal tight junction integrity.

**Discussion**

The incidence of gout and hyperuricemia continually to increases as
individuals. However, research on gout among the elderly population is
relatively limited. Previous studies have confirmed a close relationship
between gout and the gut microbiota(24, 25), but the specific connection

between the gut microbiota of older individuals and gout remains unexplored. FMT from young to old mice also reportedly alleviates age-related stroke(26)and other benefits(11, 27). And there are also studies indicating Older patients demonstrate heightened inflammatory reactions during gout attacks(28). Based on this finding, we hypothesize that the gut microbiota of different age groups may exert varying effects on gout. Therefore, we conducted cross-age FMT to investigate the interaction between the gut microbiota in different age groups and gout.

This study demonstrated that FMT from young to aged mice effectively alleviated the inflammatory response caused by MSU and improved uric acid metabolism in elderly mice, reducing the symptoms of gout. In contrast, FMT from old or aged to young mice exacerbated gout, indicating the importance of the gut microbiota composition in gout development. Activation of the NLRP3 inflammasome has been implicated in the pathogenesis of gout, leading to the production of inflammatory cytokines such as IL-1β, IL-6, and TNF-α. The beneficial effects of FMT from young to aged mice could be attributed to inhibition of the NLRP3 inflammasome pathway. A "younger" gut microbiota could suppress the activation of NLRP3, Caspase-1 and IL-1β, thereby reduce the inflammatory response in gout. Furthermore, the modulation of uric acid metabolism in old or aged mice played a role in the effects of young gut microbiota transplantation. Additionally, the improvement in the intestinal tight junction integrity observed after transplantation of the gut microbiota from young to old or aged mice may contribute to enhanced elimination of uric acid. Recent studies have also shown a direct relationship between hyperuricemia and gut microbiota. Certain anaerobic microbial communities are capable of effectively degrading purines and uric acid and thereby regulate the abundance of purines in the body to maintain the body's uric acid balance(29-31). Our study also showed that the gut microbiota plays a role in maintaining the balance of serum uric acid in the body. The "ageing" gut microbiota is more likely to elevate the levels of serum uric acid in the body by

suppressing the expression of uric acid excretion transporter (OAT1, OAT3).
The "younger" gut microbiota helps to restrain the increase of serum uric acid
levels in old or aged mice by suppressing the expression of uric acid
reabsorption transporters (URAT1) and the activity of uric acid-producing
enzymes (mainly XOD) and also promotes the expression of uric acid
excretion transporters (ABCG2) in aged mice. We also found that old or aged
mice exhibited stronger purine metabolism (Table 2a, b). These results may
explain the reason for the high levels of uric acid detected in elderly
individuals.

Moreover, through 16S rDNA sequencing analysis, we found that the
abundance of *Bifidobacterium* and *Akkermansia* was higher in young mice.
Additionally, FMT from young to old or aged mice increased the abundance of
the *Akkermansia* genus. There are existing reports regarding the therapeutic
effects of *Bifidobacterium*(32) and *Akkermansia*(33) on gout and
hyperuricemia. Therefore, we performed further investigation, and using
microbiome-predicted functional and untargeted metabolomics data, we
observed a significant enhancement in butanoate metabolism in young mice.
Surprisingly, using untargeted metabolomics data, comparisons of the results
after FMT from young to old or aged mice with those of old or aged mice also
identified the secondary bile acid biosynthesis pathways. Studies have
indicated that transplantation of the gut microbiota from healthy mice to
progeroid mice can enhance the enrichment of secondary bile acid
biosynthesis pathways, and the restoration of secondary bile acids may
potentially contribute to extension of the healthspan and lifespan in progeroid
mice(34). The current investigation into the impact of secondary bile acids on
gout and hyperuricemia is still in its incipient phase, with the intrinsic
mechanisms remaining largely unelucidated, thereby rendering this domain a
pivotal subject of profound scientific inquiry. Moreover, old or aged mice
transplanted with gut microbiota from young mice also exhibited an increase in
butanoate metabolism. This finding led us to hypothesize that butyric acid

might play a role in this phenomenon. Subsequently, we selected the Young+PBS, Aged+PBS, and Aged+Young groups for analysis of the short-chain fatty acid levels and discovered that the fecal butyric acid content in young mice was higher than that in aged mice, and that FMT from young mice led to an increase in fecal butyric acid content in aged mice. Consistent with previous reports, the content of SCFAs in the faeces of elderly mice was lower than that in the faeces of young mice(35). This result shifted our focus to butyric acid. Butyric acid, a SCFA derived from the gut microbiota, has been found to play a crucial role in host health(36). Emerging evidence indicates that among SCFAs, butyrate plays a pivotal role as a regulator in mediating metabolic control of the microbiota(37). The anti-inflammatory effects of butyric acid have been extensively reported,(38, 39) but only a few studies have investigated its impact on gout. Similarly, research on the anti-hyperuricemia effects of butyric acid is relatively scarce. We then conducted experiments using gout and peritonitis models, and found that mice supplemented with butyrate exhibited relief in symptoms of gout and inflammation. Butyrate could also reduce the serum uric acid levels in old or aged mice by promoting the expression of uric acid excretion transporters.

The findings of this study demonstrate the positive effects of gut microbiota transplantation from young to aged mice in mitigating gout symptoms. Inhibition of the NLRP3 inflammasome pathway was found to modulate uric acid metabolism, and increase butyric acid levels. The gut microbiota of young mice effectively alleviated the inflammatory response and improved uric acid metabolism in aged mice. We enhanced Butanoate metabolism in the gut microbiota of young mice, while the opposite was observed in aged mice. Furthermore, we observed higher levels of butyrate in the feces of young mice. Moreover, butyrate has excellent anti-inflammatory and uric acid-lowering effects. These findings highlight the potential of targeting the gut microbiota as a therapeutic approach for the prevention and treatment of senile gout-related conditions.

**Materials and Methods**

**Preparation of MSU crystals**

A solution consisting of 10 mM uric acid and 154 mM NaCl (both from Sigma) was adjusted to pH 7.2 and agitated for 7 days to produce MSU crystals. For sterilization the needle-shaped crystals were washed with ethanol, dried under sterile conditions and heated at 180°C for 2 h. The crystals were stored at 20 mg/mL in sterile PBS for the experimental model.

**Animals**

All the mice in the different age groups were purchased from Longde Biotechnology Co., LTD (Changchun, Jilin, China). The mice were housed in a controlled environment with a temperature of 24±2°C and a 12-hour light/12-hour dark cycle, in a pathogen-free facility. These mice were provided ad libitum access to food and water. All animal experiments were conducted in compliance with the regulations set forth by the Administration of Affairs Concerning Experimental Animals in China. The Institutional Animal Care and Use Committee at Jilin University approved the entire protocol, including the mouse autopsy and sample collection (20170318).

**Animal and FMT**

Young (3 months), old (18 months) and aged (24 months) male mice were randomly reassigned to experimental cages one week prior to the experiment. The mice in the FMT groups were subjected to antibiotic interventions and depletion of the host gut microbiota by via oral gavage of a 3-day broad-spectrum antibiotic cocktail regimen (vancomycin, 100 mg/kg; metronidazole, 200 mg/kg; ampicillin 200 mg/kg, neomycin 200 mg/kg) to maximize potential engraftment of the donor microbiota. Those in the PBS groups were orally gavaged with isopycnic PBS.

Before antibiotic interventions, fecal slurries from the donor mice were prepared by pooling fecal pellet material from 12-15 mice per cage (chosen randomly) for each age group. At the time of pellet collection, at least three pellets from each mouse were collected into sterile centrifuge tubes, and the

mice were then returned to their cage. The fecal pellets from each age group
were mixed, and sterile PBS (five times the weight of the fecal pellets) was
added. The homogenate was centrifuged at 500 × g and 4°C for 10 min and
fecal slurry was collected. After addition of 10% glycerol (v/v) and the slurry
was stored at −80°C until use. After antibiotic washout, recipient mice were
reassigned to different groups and received heterochronic donor microbiota
twice 3 days apart by oral gavage of the fecal slurry preparation. Each mouse
received 200 µL the fecal slurry preparation during each gavage.

Control groups for 3-month-old, 18-month-old and 24-month-old receiving
heterochronic transfer were gavaged with PBS only, revealed antibiotic
treatment/PBS gavage only and were administered control microbiota
(obtained from a donor pool of the same age group).

Before MSU injection, fecal pellets were collected for sequencing and
untargeted metabolomics analysis on day 22, and were then stored at −80°C.
Blood and tissues were collected at the end of the experimental period. All
FMT interventions were performed at the same time of day across all the
cages to control for the circadian rhythm variabilities in feeding and the
microbiota/metabolite composition.

**Experimental model details**

**Gout model**

To establish this model, we administered 0.4 mg of preformed MSU
crystals, resuspended in 20 µL of sterile PBS via subcutaneous injection into
the left hind paws of the animals. We measured the hind paw thickness using
calipers (the minimum accuracy of which was 0.01 mm) before and 8 hours
after MSU injection. Eight hours after injection, mouse dorsal foot tissues were
collected for histopathological observation and protein extraction.

**Peritonitis model**

Peritonitis was induced in the mice by intraperitoneal injection of MSU
crystals at a dose of 2.5 mg in PBS. Six hours later, the peritoneal cells were
collected by lavage (3 mL of PBS), and the cell count was determined using a

hemocytometer to ensure consistent cell numbers. The lavage fluid was used to detect cytokines, and proteins were cells extracted from the cells.

**Butyrate supplementation experiment**

In this experiment, 3-month-old mice received a supplemental dose of butyrate (administered via gavage at a concentration of 200 mg/kg) for 14 days. Subsequently, these mice were subjected to monosodium urate (MSU) stimulation. The investigation was conducted utilizing both gout and peritonitis models to assess the effects of butyrate supplementation. Simultaneously, mice aged 18 months and 24 months were given the same dosage of supplemental butyrate as the 3-month-old mice for 14 days. Subsequently, their liver, kidneys, intestines, and blood samples were collected for analysis.

**Histological analysis**

The left hind paws were collected and immediately fixed with 4% paraformaldehyde for more than 48 h. The fixed tissues were prepared after paraffin embedding. At least three slices from each tissue were prepared. The prepared sections were then dewaxed with xylene and hydrated via different gradients of alcohol. Furthermore, all hydrated sections were stained with hematoxylin &eosin (H&E), detected under an optical microscope (Olympus, Tokyo, Japan) and analyzed with Caseviewer2.0 software.

**Western blotting**

The total proteins from the peritoneal cells and foot tissue samples were extracted using a protein extraction kit (Thermo Fisher Scientific, USA). The targeted proteins were separated by 10% or 15% SDS☐PAGE based on the molecular weight and then bonded to 0.22 μm PVDF membranes. After blocking with 5% skim milk for three hours at RT, specific antibodies, including NLRP3, Caspace-1, IL-1β and β-actin, were used to detected target proteins at an appropriate final concentration according to the manufacturer's instructions. The PVDF membranes were then incubated with goat anti-rabbit IgG (1:20,000), wash with TBST, and determined using the ECL plus western blotting detection system (Tanon, China). The grey values of the western

601 blotting bands were procured using Image-Pro Plus 6.0 software.

**RNA extraction and qPCR**

603 Tissue samples were collected and total RNA was extracted using Trizol (Takara, CA, JPN) as previously described, cDNA was reverse transcribed using a TransScript Uni All-in-One (TransGen Biotech, Beijing, China) and reacted with specific primers using a FastStart Universal SYBR Green Master Mix (ROX) (Roche, Switzerland, Basel) and Bio-Rad CFX96 (Bio-Rad Laboratories, CA, USA). The specific primers used in this study are shown in Table 1. The $2^{-\Delta\Delta Ct}$ method was used to calculate the relative expression levels of genes using the control group as the calibrator.

**Cytokine determination**

612 For detection of the IL-β, IL-6 and TNF-α concentrations in mouse foot tissue, foot tissue was harvested from different treatment groups, and 10% tissue homogenates were prepared using RIPA buffer. The prepared samples were then centrifuged at 3,000 × g and 4°C for 10 min, and supernatants were collected. All prepared foot tissue supernatants were determined using IL-β, IL-6 and TNF-α ELISA kits at proper dilutions according to the manufacturer's instructions (BioLegend, CA, USA). For the detection of cytokines in serum and peritoneal fluid, samples were separated by centrifugation (500 × g, 4°C, 10 min) and detected using IL-β, IL-6 and TNF-α ELISA kits according to the manufacturer's instructions (BioLegend, CA, USA).

**Serum biochemistry analysis**

623 The levels of serum uric acid (SUA), serum alanine aminotransferase (ALT), serum aspartate aminotransferase (AST), serum creatinine (Cr) and blood urea nitrogen (BUN) were measured using enzymatic kits from Nanjing Jiancheng Bioengineering Institute, Nanjing, China. The levels of serum lipopolysaccharide (LPS) and serum muramyl dipeptide (MDP) were measured using enzymatic kits from FEIYA BIOTECHNOLOGY.

**Assay of uric acid synthesis enzymes**

To detect the activity of Adenosine deaminase (ADA) and Guanine Deaminase (GDA) in the mouse liver, the liver was collected from mice in the various treatment groups and homogenized in sterile physiological saline solution (10-fold weight ratio of tissue). Following preparation, the samples were subjected to centrifugation at 3,000 × g and 4°C for 10 minutes, and the supernatants were collected. The activity of ADA and GDA in the liver supernatants was determined using ELISA kits (FEIYA BIOTECHNOLOGY, China) according to the manufacturer's instructions with appropriate dilutions. To measure the activity of xanthine oxidase (XOD) in the liver and kidney of mice, tissues were collected from mice in the different treatment groups and homogenized in the buffer provided in the kit (10-fold weight ratio of tissue). The samples were then centrifuged at 8,000 × g and 4°C for 10 minutes, and the supernatants were collected. The activity of XOD in the liver and kidney supernatants was assessed using an ELISA kit (Beijing Solarbio Science & Technology Co., Ltd, China) according to the manufacturer's instructions.

**Fecal microbiota composition analysis**

For specific details, please refer to Supplementary material 1.

**Untargeted metabolomic analysis**

For specific details, please refer to Supplementary Material 2.

**Analysis of short-chain fatty acids in faeces**

For specific details, please refer to Supplementary Material 3.

**Statistical analysis**

All values were expressed as the means ± SEM. Raw data were subjected to one-way ANOVA to evaluate statistical significance between at least three groups, and pairwise comparison was conducted using Student's t-test. The results were considered statistically significant at $p < 0.05$.

**Data and code availability**

The 16S rDNA sequencing data and untargeted metabolomic data presented in this study can be found in online repositories. The findings of this study have been deposited into the CNGB Sequence Archive (CNSA) of China

National GenBank Database (CNGBdb) under the accession number CNP0004751. The names of the repository/repositories and accession number(s) can be found below: https://db.cngb.org/cnsa/project/CNP0004751_46c178da/reviewlink/. The lead contact will provide original western blot images and microscopy data from this paper upon request.

No unique code was generated in this study.

For any further information needed to reanalyze the data presented in this paper, it can be obtained from the lead contact upon request.

**Declaration of interests**

The authors declare no competing interests.

**Acknowledgments**

This work was supported by the National Natural Science Foundation of China (Nos. 32202889 and 31972749) and the Natural Science Foundation of Jilin Province (No. 20220101304JC).

**Author contributions**

The study was designed by Naisheng Zhang, Wenlong Zhang, and Hang Gao. Ning Song conducted all mouse experiments and performed the statistical analysis. Mice sample collection was carried out by Jianhao Li, Mingze Wang and Yi Liu. Ning Song was responsible for software applications, investigation, and writing the original draft. Jianhao Li and Mingze Wang helped with validation and analysis. Ning Song and Yi Liu were responsible for resources and data curation. Wenlong Zhang and Zhiming Ma wrote and revised the final draft of the manuscript, and all authors participated in its revision and gave their approval.

**Reference**

1.    N. Dalbeth, T. R. Merriman and L. K. Stamp: Gout. *Lancet*, 388(10055), 2039-2052 (2016) doi:10.1016/S0140-6736(16)00346-9

2.    L. K. Stamp, S. J. D. Jordan and Aging: The Challenges of Gout Management in the Elderly, 28(8), 591-603 (2011)

3.    Ashika, Chhana, Gerald, Lee, Nicola and D. J. B. m. disorders: Factors influencing the crystallization of monosodium urate: a systematic literature review (2015)

4.    A. K. J. E.-B. N. Duttaroy, C. E. o. B. F. i. H. Health and Disease: Gut microbiota on human health and disease, 269-281 (2021)

5.    A. Adak, M. R. J. C. Khan and M. L. Sciences: An insight into gut microbiota and its functionalities, 76(11) (2019)

6.    Z. Guo, J. Zhang, Z. Wang, K. Y. Ang, S. Huang, Q. Hou, X. Su, J. Qiao, Y. Zheng and L. J. S. R. Wang: Intestinal Microbiota Distinguish Gout Patients from Healthy Humans, 6, 20602 (2016)

7.    S. Tiejuan, S. Li, L. Haichang, X. Zhijun, H. Zhixing and W. J. F. i. M. Chengping: Combined Signature of the Fecal Microbiome and Metabolome in Patients with Gout, 8(578527), 268 (2017)

8.    N. Yamada, C. Iwamoto, H. Kano, N. Yamaoka, T. Fukuuchi, K. Kaneko, Y. J. N. N. Asami and N. Acids: Evaluation of purine utilization by Lactobacillus gasseri strains with potential to decrease the absorption of food-derived purines in the human intestine, 35(10-12), 670-676 (2016)

9.    T. W. Buford: The Gut Microbiome and Aging. *J Gerontol A Biol Sci Med Sci*, 75(7), 1229-1231 (2020) doi:10.1093/gerona/glaa103

10.  S. V. Lynch and O. Pedersen: The Human Intestinal Microbiome in Health and Disease (2013)

11.  Aimée.Parker, S. Romano, R. Ansorge, A. Aboelnour, G. L. Gall, G. M. Savva, M. G. Pontifex, A. Telatin, D. Baker and E. J. M. Jones: Fecal microbiota transfer between young and aged mice reverses hallmarks of the aging gut, eye, and brain, 10(1), 68 (2022)

12.  M. Boehme, K. Guzzetta, T. F. Bastiaanssen and J. Cryan: Microbiota from young mice counteracts selective age-associated behavioral deficits (2021)

13. C. Schauer, C. Janko, L. E. Munoz, Y. Zhao, D. Kienhfer, B. Frey, M. Lell, B. Manger, J.
Rech and E. Naschberger: Aggregated neutrophil extracellular traps limit inflammation by
degrading cytokines and chemokines (2013)
14. Y. Gabsik, Y. S. Hyeon, L. H. Eun, K. H. Chang, C. Y. Yeon, L. H. Suk and L. J. J. R.
Young: Suppression of NLRP3 inflammasome by oral treatment with sulforaphane alleviates
acute gouty inflammation(4), 4
15. A. Mariotte, A. D. Cauwer, C. Po, C. Abou-Faycal and P. J. T. Georgel: A mouse model of
MSU-induced acute inflammation in vivo suggests imiquimod-dependent targeting of Il-1β as
relevant therapy for gout patients, 10(5), 2158-2171 (2020)
16. E. L. Goldberg, J. L. Asher, R. D. Molony, A. C. Shaw, C. J. Zeiss, C. Wang, L. A.
Morozova-Roche, R. I. Herzog, A. Iwasaki and V. D. J. C. R. Dixit: β-Hydroxybutyrate
Deactivates Neutrophil NLRP3 Inflammasome to Relieve Gout Flares, 18(9), 2077-2087
(2017)
17. F. Martinon, V. Pétrilli, A. Mayor, A. Tardivel and J. J. N. Tschopp: Gout-associated uric
acid crystals activate the NALP3 inflammasome, 440(7081), 237-41 (2006)
18. T. S. Ghosh, F. Shanahan and P. W. O'Toole: The gut microbiome as a modulator of
healthy ageing. *Nature Reviews Gastroenterology & Hepatology*, 19(9), 565-584 (2022)
doi:10.1038/s41575-022-00605-x
19. L. K. Sarao and M. Arora: Probiotics, prebiotics, and microencapsulation: A review.
*Critical Reviews in Food Science and Nutrition*, 57(2), 344-371 (2017)
doi:10.1080/10408398.2014.887055
20. A. Nowak, A. Paliwoda and J. J. C. R. C. C. R. i. F. T. Basiak: Anti-proliferative,
pro-apoptotic and anti-oxidative activity of Lactobacillus and Bifidobacterium strains: A review
of mechanisms and therapeutic perspectives
21. M. J. I. J. S. E. M. Derrien: Akkermansia muciniphila gen. nov., sp. nov., a human
intestinal mucin-degrading bacterium, 54 (2004)
22. T. Zhang, Q. Li, L. Cheng, H. Buch and F. J. M. B. Zhang: Akkermansia muciniphila is a
promising probiotic, 12(6) (2019)
23. J. Lee, J. d'Aigle, L. Atadja, V. Quaicoe, P. Honarpisheh, B. P. Ganesh, A. Hassan, J. Graf,
J. Petrosino, N. Putluri, L. Zhu, D. J. Durgan, R. M. Bryan, Jr., L. D. McCullough and V. R.
Venna: Gut Microbiota-Derived Short-Chain Fatty Acids Promote Poststroke Recovery in Aged
Mice. *Circ Res*, 127(4), 453-465 (2020) doi:10.1161/circresaha.119.316448
24. Y. Chu, S. Sun, Y. Huang, Q. Gao, X. Xie, P. Wang, J. Li, L. Liang, X. He, Y. Jiang, M.
Wang, J. Yang, X. Chen, C. Zhou, Y. Zhao, F. Ding, Y. Zhang, X. Wu, X. Bai, J. Wu, X. Wei, X.
Chen, Z. Yue, X. Fang, Q. Huang, Z. Wang and R. Huang: Metagenomic analysis revealed the
potential role of gut microbiome in gout. *NPJ Biofilms Microbiomes*, 7(1), 66 (2021)
doi:10.1038/s41522-021-00235-2
25. E. O. Méndez-Salazar and G. A. Martínez-Nava: Uric acid extrarenal excretion: the gut
microbiome as an evident yet understated factor in gout development. *Rheumatology*
*International*, 42(3), 403-412 (2022) doi:10.1007/s00296-021-05007-x
26. J. A. Lee, JohnAtadja, LouiseQuaicoe, VictoriaHonarpisheh, PedramGanesh, Bhanu
P.Hassan, AhmadGraf, JoergPetrosino, JosephPutluri, NagireddyZhu, LiangDurgan, David
J.Bryan, Robert M., Jr.McCullough, Louise D.Venna, Venugopal Reddy %J Circulation
research: a journal of the American Heart Association: Gut Microbiota-Derived Short-Chain

Fatty Acids Promote Poststroke Recovery in Aged Mice, 127(4) (2020)

27. K. H. Kim, Y. Chung, J.-W. Huh, D. J. Park, Y. Cho, Y. Oh, H. Jeong, J. Yoon, J.-H. Kang, H.-S. Shin, H.-C. Kim, S.-K. Kwon, K. Y. Seo, S. H. Oh, J. K. Seong, S.-J. Ha, K. T. Nam and J. F. Kim: Gut microbiota of the young ameliorates physical fitness of the aged in mice. *Microbiome*, 10(1), 238 (2022) doi:10.1186/s40168-022-01386-w

28. J. A. Y. ), J. H. L. ), E. Y. L. ), E. B. L. ), Y. W. S. ) and J. K. P. ): Poster Session : PS 0672 ; Rheumatology ; Higher Infl ammatory Response in Elderly Patients Dring Gout Attack

29. Y. Liu, J. B. Jarman, Y. S. Low, H. E. Augustijn, S. Huang, H. Chen, M. E. DeFeo, K. Sekiba, B.-H. Hou, X. Meng, A. M. Weakley, A. V. Cabrera, Z. Zhou, G. van Wezel, M. H. Medema, C. Ganesan, A. C. Pao, S. Gombar and D. Dodd: A widely distributed gene cluster compensates for uricase loss in hominids. *Cell*, 186(16), 3400-3413.e20 (2023) doi:https://doi.org/10.1016/j.cell.2023.06.010

30. K. Kasahara, R. L. Kerby, Q. Zhang, M. Pradhan, M. Mehrabian, A. J. Lusis, G. Bergström, F. Bäckhed and F. E. Rey: Gut bacterial metabolism contributes to host global purine homeostasis. *Cell Host & Microbe*, 31(6), 1038-1053.e10 (2023) doi:https://doi.org/10.1016/j.chom.2023.05.011

31. L. Pan, P. Han, S. Ma, R. Peng, C. Wang, W. Kong, L. Cong, J. Fu, Z. Zhang, H. Yu, Y. Wang and J. Jiang: Abnormal metabolism of gut microbiota reveals the possible molecular mechanism of nephropathy induced by hyperuricemia. *Acta Pharm Sin B*, 10(2), 249-261 (2020) doi:10.1016/j.apsb.2019.10.007

32. A. T. Vieira, I. Galvão, F. A. Amaral, M. M. Teixeira, J. R. Nicoli and F. S. Martins: Oral treatment with Bifidobacterium longum 51A reduced inflammation in a murine experimental model of gout. *Benef Microbes*, 6(6), 799-806 (2015) doi:10.3920/bm2015.0015

33. L. Zhang, J. Liu, T. Jin, N. Qin, X. Ren and X. Xia: Live and pasteurized Akkermansia muciniphila attenuate hyperuricemia in mice through modulating uric acid metabolism, inflammation, and gut microbiota. *Food Funct*, 13(23), 12412-12425 (2022) doi:10.1039/d2fo02702j

34. C. Bárcena, R. Valdés-Mas, P. Mayoral, C. Garabaya, S. Durand, F. Rodríguez, M. T. Fernández-García, N. Salazar, A. M. Nogacka, N. Garatachea, N. Bossut, F. Aprahamian, A. Lucia, G. Kroemer, J. M. P. Freije, P. M. Quirós and C. López-Otín: Healthspan and lifespan extension by fecal microbiota transplantation into progeroid mice. *Nat Med*, 25(8), 1234-1242 (2019) doi:10.1038/s41591-019-0504-5

35. J. Lee, J. D'Aigle, L. Atadja, V. Quaicoe and V. R. J. C. R. Venna: Gut Microbiota-Derived Short-Chain Fatty Acids Promote Post-Stroke Recovery in Aged Mice, 127(4) (2020)

36. M. A. Vinolo, H. G. Rodrigues, R. T. Nachbar and R. Curi: Regulation of inflammation by short chain fatty acids. *Nutrients*, 3(10), 858-76 (2011) doi:10.3390/nu3100858

37. L. Zhang, C. Liu, Q. Jiang and Y. Yin: Butyrate in Energy Metabolism: There Is Still More to Learn. *Trends Endocrinol Metab*, 32(3), 159-169 (2021) doi:10.1016/j.tem.2020.12.003

38. G. Li, J. Lin, C. Zhang, H. Gao, H. Lu, X. Gao, R. Zhu, Z. Li, M. Li and Z. Liu: Microbiota metabolite butyrate constrains neutrophil functions and ameliorates mucosal inflammation in inflammatory bowel disease. *Gut Microbes*, 13(1), 1968257 (2021) doi:10.1080/19490976.2021.1968257

39. G. Chen, X. Ran, B. Li, Y. Li, D. He, B. Huang, S. Fu, J. Liu and W. Wang: Sodium Butyrate Inhibits Inflammation and Maintains Epithelium Barrier Integrity in a TNBS-induced

817     Inflammatory   Bowel   Disease   Mice   Model.   *EBioMedicine*,   30,   317-325   (2018)

818     doi:10.1016/j.ebiom.2018.03.030

828     **Figure 1**

837 **Figure 2**

846 **Figure 3**

855 **Figure 4**

864 **Figure 5**

**Figure 6**

**Figure 7**

**Figure 8**

**Supplementary figure 1**

913     **Supplementary figure 2**

**Supplementary figure 3**

933 **Supplementary figure 4**

944 **Supplementary figure 5**

947 **Supplementary figure 6**

a

b

963 **Supplementary figure 7**

**Supplementary figure 8**

993 **Figure Legends**

**Fig 1 Gout susceptibility increases with age, related to gut microbiota**

(a) Mice of different ages.

(b-d) The foot thickness ratios (b), foot tissue's IL-1β concentrations (c) and serum concentrations of uric acid of three different age ranges groups (d) were tested after MSU administration (n=6).

(e-f) The three different age ranges groups' ASVs (Amplicon Sequence Variants) (e) and PcoA analysis (using Bray-Curtis dissimilarity and Jaccard distance) (f).

(g-i) The foot thickness ratios (g), foot tissue's IL-1β concentrations (h) and serum concentrations of uric acid of three different age ranges groups (i) (treated with antibiotics (ABX) cocktail) were tested after MSU administration (n=6).

Values are presented as the mean ± SEM. Differences were assessed by One-Way ANOVA and denoted as follows: $*p < 0.05$, $**p < 0.01$, and $***p < 0.001$, "ns" indicates no significant difference between groups.

**Fig 2 Aged-to-young FMT worsens acute gout, whereas young-to-aged FMT reduced this disease.**

(a) Overall experimental design and timeline for experiments.

(b) Representative H&E-stained images of left foot tissues. Scale bars 1000 μm and 3x magnification.

(c-e) Foot tissue inflammatory parameters, including IL-1β (c), IL-6 (d) and TNF-α (e) concentrations, from the indicated mice are shown (n=6).

(f-h) The peritoneal fluid concentrations of IL-1β (f), IL-6 (g) and TNF-α (h) inflammatory parameters were measured in the indicated mice(n=6).

Values are presented as the mean ± SEM. Differences were assessed by t-test or One-Way ANOVA and denoted as follows: $*p < 0.05$, $**p < 0.01$, and $***p < 0.001$, "ns" indicates no significant difference between groups.

**Fig 3 "Younger" gut microbiota suppresses NLRP3 inflammasome pathway, "aging" gut microbiota promotes**

(a-b) Representative western blot images and band density (Young+PBS,

Young+Old and Young+Aged) of foot tissue NLRP3 pathways proteins (n=3).

(c-d) Representative western blot images and band density (Old+PBS and

Old+Young) of foot tissue NLRP3 pathways proteins (n=3).

(e-f) Representative western blot images and band density (Aged+PBS and

Aged+Young) of foot tissue NLRP3 pathways proteins (n=3).

Values are presented as the mean ± SEM. Differences were assessed by t-test

or One-Way ANOVA and denoted as follows: $*p < 0.05$, $**p < 0.01$, and $***p <$

$0.001$, "ns" indicates no significant difference between groups.

**Fig 4 Beneficial effects of FMT from young to aged mice on uric acid**

**metabolism.**

(a) All groups' serum concentrations of uric acid (n=6).

(b-d) The activity of uric acid-producing enzymes of liver in the cross-age fecal

microbiota transplantation group and its control group (n=6), including ADA (b),

GDA (c) and XOD (d).

(e) The activity of XOD of kidney in the cross-age fecal microbiota

transplantation group and its control group (n=6).

(f) Relative kidney injury molecule-1 (KIM-1) expression in the indicated

groups by qPCR (n = 6).

(g and h) Relative renal genes for uric acid reabsorption expression in the

indicated groups by qPCR (n = 6), including URAT1 (g) and GLUT9 (h).

(i-k) Relative renal genes for uric acid excretion expression in the indicated

groups by qPCR (n = 6), including OAT1 (i), OAT3 (j) and ABCG2 (k).

Values are presented as the mean ± SEM. Differences were assessed by t-test

or One-Way ANOVA and denoted as follows: $*p < 0.05$, $**p < 0.01$, and $***p <$

$0.001$, "ns" indicates no significant difference between groups.

**Fig 5 Modifications in the gut microbiota composition following**

**cross-age FMT.**

(a) Bacterial composition at the phylum levels (top 10) of the indicated groups

(n=6).

(b-c) Alpha diversity indices including Chao1 (b) and observed_otus (c) index

in the indicated groups (n=6).

(d) β-diversity difference among the seven groups analyzed by the PcoA using
Bray-Curtis dissimilarity and Jaccard distance.

(e) Volcano Plot of inter-group significance analysis using metastat (t-test, $p <$
0.05).

(f) The ternary plot of three different groups among the seven groups at genus
levels (top 10).

**Fig 6 Changes in fecal microbiota metabolism and pathways after**
**cross-age FMT.**

(a) Comparison of differential KEGG enrichment bubble plots between
Young+PBS and Young+Old.

(b) Comparison of differential KEGG enrichment bubble plots between
Young+PBS and Young+Aged.

(c) Comparison of differential KEGG enrichment bubble plots between
Old+PBS and Old+Young.

(d) Comparison of differential KEGG enrichment bubble plots between
Aged+PBS and Aged+Young.

The enrichment analysis was performed at the KEGG Pathway level using a
hypergeometric test, as shown in the figure below. The pathways that were
significantly enriched in the differential metabolites compared to the
background of all identified metabolites. Pathway enrichment analysis enables
us to determine the major biochemical metabolic pathways and signaling
transduction pathways that are implicated by the differential metabolites.

**Fig 7 Butyrate supplementation inhibits gout and MSU-induced**
**peritonitis.**

(a) The concentration of primary short-chain fatty acids (SCFAs) in fecal
samples from both young (3 months), aged (24 months) mice and
Aged+Young (FMT from young to aged). Graphs were generated to illustrate
the changes in individual SCFAs, with a sample size of 6 in the Young+PBS,
Aged+PBS and Aged+Young group(n=6).

(b-e) The foot thickness ratios (b) were tested after MSU administration, foot tissue inflammatory parameters, including IL-1β (c), IL-6 (d) and TNF-α (e) concentrations, from the indicated mice are shown(n=6).

(f) Representative H&E-stained images of left foot tissues. Scale bars 1000 μm and 3x magnification.

(g-i) The peritoneal fluid concentrations of IL-1β (g), IL-6 (h) and TNF-α (i) inflammatory parameters were measured in the indicated mice (n=6).

(j) Representative western blot images of foot tissue NLRP3 pathways proteins and band density (n=3).

Values are presented as the mean ± SEM. Differences were assessed by t-test or One-Way ANOVA and denoted as follows: *p < 0.05, **p < 0.01, and ***p < 0.001, "ns" indicates no significant difference between groups.

**Fig 8 Serum uric acid-lowering effect of butyrate in old or aged mice.**

(a) The serum uric acid concentrations in Old+PBS and Old+Butyrate, Aged+PBS and Aged+Butyrate (n=6).

(b-d) The activity of uric acid-producing enzymes of liver in the old or aged control group and the old or aged group supplemented with butyrate (n=6), including ADA (b), GDA (c) and XOD (d).

(e) The activity of XOD of kidney in the Old or Aged control group and the old or aged group supplemented with butyrate (n=6).

(f) Relative kidney injury molecule-1 (KIM-1) expression in the indicated groups by qPCR (n = 6).

(g and h) Relative renal genes for uric acid reabsorption expression in the indicated groups by qPCR (n = 6), including URAT1 (g) and GLUT9 (h).

(i-k) Relative renal genes for uric acid excretion expression in the indicated groups by qPCR (n = 6), including OAT1 (i), OAT3 (j) and ABCG2 (k).

(l) Relative colonic genes for uric acid expression in the indicated groups by qPCR, including ABCG2 (n = 6).

Values are presented as the mean ± SEM. Differences were assessed by t-test or One-Way ANOVA and denoted as follows: *p < 0.05, **p < 0.01, and ***p <

0.001, "ns" indicates no significant difference between groups.

**Supplementary Fig. 1**

(a) All groups' foot thickness ratios were tested after MSU administration (n=6).

(b-d) The serum concentrations of IL-1β (b), IL-6 (c) and TNF-α (d)

inflammatory parameters were measured in the indicated mice(n=6).

Values are presented as the mean ± SEM. Differences were assessed by t-test

or One-Way ANOVA and denoted as follows: *p < 0.05, **p < 0.01, and ***p <

0.001, "ns" indicates no significant difference between groups.

**Supplementary Fig. 2**

(a-b) Representative western blot images and band density (Young+PBS,

Young+Old and Young+Aged) of peritoneal cells NLRP3 pathways proteins

(n=3).

(c-d) Representative western blot images and band density (Old+PBS and

Old+Young) of peritoneal cells NLRP3 pathways proteins (n=3).

(e-f) Representative western blot images and band density (Aged+PBS and

Aged+Young) of peritoneal cells NLRP3 pathways proteins (n=3).

Values are presented as the mean ± SEM. Differences were assessed by t-test

or One-Way ANOVA and denoted as follows: *p < 0.05, **p < 0.01, and ***p <

0.001, "ns" indicates no significant difference between groups.

**Supplementary Fig. 3**

(a and b) Serum AST (a) and ALT (b) concentrations in the cross-age fecal

microbiota transplantation group and its control group (n=6).

(c and d) Serum Crea (c) and BUN (d) concentrations in the cross-age fecal

microbiota transplantation group and its control group (n=6).

(e-h) Relative colonic gene expression in the indicated groups by qPCR (n = 6),

including ABCG2 (e), Occludin (f), ZO-1 (g) and JAMA (h).

Values are presented as the mean ± SEM. Differences were assessed by t-test

or One-Way ANOVA and denoted as follows: *p < 0.05, **p < 0.01, and ***p <

0.001, "ns" indicates no significant difference between groups.

**Supplementary Fig. 4**

(a) Bacteroidetes relative abundance of the indicated groups (n=6).

(b) Firmicutes relative abundance of the indicated groups (n=6).

(c) The Firm/Bac ratio in the different groups (n=6).

(d-e) Alpha diversity indices including Shannon (d) and Simpson (e) index in

the indicated groups (n=6).

(f) Bacterial composition at the genus levels (top 15) of the indicated

groups(n=6).

Values are presented as the mean ± SEM. Differences were assessed by t-test

or One-Way ANOVA and denoted as follows: $*p < 0.05$, $**p < 0.01$, and $***p <$

$0.001$, "ns" indicates no significant difference between groups.

**Supplementary Fig. 5**

(a) Metabolite classification of various groups.

(b) Volcano plots of differential metabolites in each group.

The horizontal axis of volcano plot represents the $\log_2$ (Fold Change) in

metabolite abundance between different groups, while the vertical axis of

volcano plot represents the significance level ($-\log_{10}$ (P-value)) of the

differences. Each point on the volcano plot represents a metabolite, with

significantly upregulated metabolites depicted as red points and significantly

downregulated metabolites depicted as green points. The size of the circles

corresponds to the VIP (Variable Importance in Projection) value.

**Supplementary Fig. 6**

(a) Comparison of differential metabolite volcano plots and KEGG enrichment

bubble plots between Young+PBS and Old+PBS.

(b) Comparison of differential metabolite volcano plots and KEGG enrichment

bubble plots between Young+PBS and Aged+PBS.

To visualize the differential metabolites resulting from pairwise comparisons,

we generated volcano plots that provide a clear representation of the

upregulation and downregulation of metabolites, especially those with

significant fold change differences. The enrichment analysis was performed at

the KEGG Pathway level using a hypergeometric test, as shown in the figure

below. The pathways that were significantly enriched in the differential metabolites compared to the background of all identified metabolites. Pathway enrichment analysis enables us to determine the major biochemical metabolic pathways and signaling transduction pathways that are implicated by the differential metabolites.

**Supplementary Fig. 7**

(a-h) The difference analysis of the detected short chain fatty acids, including acetic acid (a), propionic acid (b), isobutyric acid (c), butyric acid (d), 2-merhylbutyrate (e), isovaleric acid (f), valeric acid (g), hexanoic acid (h).

(i-k) The serum concentrations of IL-1β (i), IL-6 (j) and TNF-α (k) inflammatory parameters were measured in the indicated mice (n=6).

(l) Representative western blot images and band density of peritoneal cells NLRP3 pathways proteins (n=3)

Values are presented as the mean ± SEM. Differences were assessed by t-test or One-Way ANOVA and denoted as follows: *p < 0.05, **p < 0.01, and ***p < 0.001, "ns" indicates no significant difference between groups.

**Supplementary Fig. 8**

(a-c) Relative colonic gene expression in the indicated groups by qPCR (n = 6), including Occludin (a), ZO-1 (b) and JAMA (c).

Values are presented as the mean ± SEM. Differences were assessed by t-test or One-Way ANOVA and denoted as follows: *p < 0.05, **p < 0.01, and ***p < 0.001, "ns" indicates no significant difference between groups.

1204

1205 **Table1**

**Oligonucleotides**

| Gene | Primer | Sequence(5' to 3') |
|---|---|---|
| KIM-1 | sense | TGGTTGCCTTCCGTGTCTCT |
| KIM-1 | antisense | TCAGCTCGGGAATGCACAA |
| URAT1 | sense | GGCTCATCACAAAGACCCCA |
| URAT1 | antisense | CAGCATGTTCTGGGTGGTGA |
| OAT1 | sense | GCAGCCTATGCACCCAACTA |
| OAT1 | antisense | ACAGCATAGGCGATACCAGC |
| OAT3 | sense | GGGCTCGAGTGGGAAGTATG |
| OAT3 | antisense | CTGGTAAGGGCCGATTGAGG |
| GLUT9 | sense | GAAGGGTGTTCCTGGCTACC |
| GLUT9 | antisense | TGAACACGGACCCAAACCAG |
| ABCG2 | sense | TTGGACTCAAGCACAGCGAATG |
| ABCG2 | antisense | TCCGCAGGGTTGTTGTAGGG |
| ADA | sense | CATGCGGTTGTTCGCTTCAA |
| ADA | antisense | CTGCGTTGATGTTCAGTCGC |
| XDH | sense | ATGACGAGGACAACGGTAGATG |
| XDH | antisense | GAAGGCGGTCATACTTGGAGA |
| GDA | sense | CGACAGCGGCAAAATAGTGTT |
| GDA | antisense | GGGGCATGGATGTGGGTATC |
| ZO-1 | sense | TTTCAGAGTGGGGAAACCTCC |
| ZO-1 | antisense | CACTCTTCCTTAGCTGCTGAAC |
| JAMA | sense | TTGGCGTCTGGTTTGCCTAT |
| JAMA | antisense | CACTTCGAGTACTGGGCTGG |
| OCCLUDIN | sense | ATGCTCTCTCAGCCAGCGTA |
| OCCLUDIN | antisense | CCACACAGGCAAATATGGCG |
| GAPDH | sense | AGGTCGGTGTGAACGGATTTG |

| GAPDH | antisense | GGGGTCGTTGATGGCAACA |
| --- | --- | --- |

**Table 2 Changes in fecal microbiota metabolism and pathways after cross-age FMT.**

| Table 2a Young+PBS VS Old+PBS | | | | | | | | |
| --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Taxa | avg(Young+PBS) | sd(Young+PBS) | avg(Old+PBS) | sd(Old+PBS) | p.value | q.values | interval lower | interval upper |
| Genetic Information Processing;Translation;Transfer_RNA_biogenesis | 0.025388 | 0.000319 | 0.026282 | 0.000752 | 0.032527 | 0.095267 | -0.00169 | -1E-04 |
| Metabolism;Amino acid metabolism;Amino_acid_related_enzymes | 0.018844 | 0.000148 | 0.019137 | 0.000258 | 0.042118 | 0.102535 | -0.00057 | -1.3E-05 |
| Genetic Information Processing;Translation;Ribosome | 0.015723 | 0.000269 | 0.016458 | 0.000594 | 0.028278 | 0.090633 | -0.00137 | -0.0001 |
| Genetic Information Processing;Translation;Aminoacyl-tRNA_biosynthesis | 0.015499 | 0.000262 | 0.0164 | 0.000581 | 0.0106 | 0.060087 | -0.00152 | -0.00029 |
| Cellular Processes;Transport and catabolism;Exosome | 0.015496 | 0.000151 | 0.014721 | 0.000475 | 0.008904 | 0.057414 | 0.000277 | 0.001273 |
| Genetic Information Processing;Replication and repair;DNA_replication_proteins | 0.014807 | 0.000271 | 0.015186 | 0.000287 | 0.040656 | 0.101038 | -0.00074 | -2E-05 |
| Genetic Information Processing;Replication and repair;Chromosome_and_associated_proteins | 0.0131 | 0.000238 | 0.013673 | 0.000459 | 0.028126 | 0.090633 | -0.00107 | -8E-05 |
| Metabolism;Amino acid metabolism;Alanine,_aspartate_and_glutamate_metabolism | 0.011744 | 0.000114 | 0.011268 | 0.000326 | 0.014186 | 0.066361 | 0.000134 | 0.000819 |
| Metabolism;Amino acid metabolism;Cysteine_and_methionine_metabolism | 0.009602 | 7.4E-05 | 0.009046 | 0.000372 | 0.013822 | 0.065951 | 0.000166 | 0.000946 |
| Genetic Information Processing;Replication and repair;Nucleotide_excision_repair | 0.008149 | 0.000239 | 0.008575 | 0.000252 | 0.013177 | 0.065517 | -0.00074 | -0.00011 |
| Genetic Information Processing;Replication and repair;DNA_replication | 0.007972 | 0.000183 | 0.008423 | 0.000317 | 0.016667 | 0.072413 | -0.0008 | -0.00011 |
| Genetic Information | 0.0078 | 5.8E-0 | 0.008 | 0.00 | 0.0 | 0.1 | -0.00 | -1.4E |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Processing;Translation;Messenger_RNA_Biogenesis | 49 | 5 | 027 | 0156 | 378 77 | 000 3 | 034 | -05 |
| Metabolism;Amino acid metabolism;Glycine,_serine_and_threonine_metabolism | 0.007893 | 0.000123 | 0.007344 | 0.000264 | 0.002 38 | 0.042 25 | 0.000268 | 0.00083 |
| Metabolism;Carbohydrate metabolism;Pentose_phosphate_pathway | 0.007552 | 0.000179 | 0.007998 | 0.000372 | 0.032 208 | 0.095 267 | -0.00084 | -5E-05 |
| <mark>Metabolism;Carbohydrate metabolism;Butanoate_metabolism</mark> | <mark>0.007496</mark> | <mark>0.000432</mark> | <mark>0.006856</mark> | <mark>0.000234</mark> | <mark>0.013454</mark> | <mark>0.065517</mark> | <mark>0.000174</mark> | <mark>0.001105</mark> |
| Metabolism;Carbohydrate metabolism;Glyoxylate_and_dicarboxylate_metabolism | 0.007261 | 0.000272 | 0.006593 | 0.00058 | 0.037471 | 0.10003 | 5.11E-05 | 0.001284 |
| Metabolism;Amino acid metabolism;Lysine_biosynthesis | 0.006277 | 0.000198 | 0.006577 | 0.000222 | 0.033542 | 0.095267 | -0.00057 | -2.9E-05 |
| Metabolism;Amino acid metabolism;Arginine_biosynthesis | 0.005337 | 0.000206 | 0.004818 | 0.000269 | 0.004248 | 0.042225 | 0.000208 | 0.00083 |
| Metabolism;Lipid metabolism;Lipid_biosynthesis_proteins | 0.004748 | 0.000253 | 0.004195 | 0.000254 | 0.003621 | 0.042225 | 0.000227 | 0.00088 |
| Metabolism;Metabolism of terpenoids and polyketides;Terpenoid_backbone_biosynthesis | 0.004268 | 0.000118 | 0.004511 | 0.000132 | 0.007384 | 0.057414 | -0.0004 | -8.1E-05 |
| Metabolism;Lipid metabolism;Glycerophospholipid_metabolism | 0.004116 | 0.00012 | 0.00449 | 0.00018 | 0.002372 | 0.042225 | -0.00058 | -0.00017 |
| Metabolism;Energy metabolism;Nitrogen_metabolism | 0.004843 | 0.000271 | 0.004057 | 0.000366 | 0.002089 | 0.042225 | 0.000368 | 0.001206 |
| Metabolism;Metabolism of cofactors and vitamins;One_carbon_pool_by_folate | 0.004295 | 4.61E-05 | 0.003931 | 0.000257 | 0.017267 | 0.073561 | 9.46E-05 | 0.000634 |
| Genetic Information Processing;Replication and repair;Base_excision_repair | 0.003915 | 6.37E-05 | 0.004159 | 0.000206 | 0.033021 | 0.095267 | -0.00046 | -2.7E-05 |
| Metabolism;Metabolism of cofactors and vitamins;Pantothenate_and_CoA_biosynthesis | 0.004248 | 0.000113 | 0.003948 | 0.000163 | 0.00495 | 0.045425 | 0.000117 | 0.000483 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosyltransferases | 0.003709 | 0.000154 | 0.003904 | 0.000119 | 0.035217 | 0.098192 | -0.00037 | -1.7E-05 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Metabolism;Amino acid metabolism;Valine,_leucine_and_isoleucine_biosynthesis | 0.003878 | 0.000305 | 0.00317 | 0.0000297 | 0.0002248 | 0.0042225 | 0.0000321 | 0.001096 |
| Genetic Information Processing;Transcription;RNA_polymerase | 0.003305 | 7E-05 | 0.003533 | 0.0000182 | 0.0026747 | 0.0090633 | -0.00042 | -3.6E-05 |
| Metabolism;Carbohydrate metabolism;Pentose_and_glucuronate_interconversions | 0.003491 | 7.67E-05 | 0.002991 | 0.0000471 | 0.0048036 | 0.108091 | 6.23E-06 | 0.000993 |
| Metabolism;Glycan biosynthesis and metabolism;Other_glycan_degradation | 0.002999 | 0.000399 | 0.002122 | 0.0000652 | 0.0022059 | 0.0080739 | 0.0000162 | 0.001592 |
| Metabolism;Energy metabolism;Photosynthesis_proteins | 0.00274 | 0.000139 | 0.00297 | 8.72E-05 | 0.0008279 | 0.0057414 | -0.00038 | -7.7E-05 |
| Metabolism;Energy metabolism;Photosynthesis | 0.002731 | 0.000139 | 0.002958 | 8.61E-05 | 0.0008761 | 0.0057414 | -0.00038 | -7.4E-05 |
| Metabolism;Amino acid metabolism;Valine,_leucine_and_isoleucine_degradation | 0.00267 | 0.000206 | 0.002337 | 0.00011 | 0.0008698 | 0.0057414 | 0.0000112 | 0.000556 |
| Cellular Processes;Cell motility;Cytoskeleton_proteins | 0.002762 | 6.51E-05 | 0.002926 | 6.67E-05 | 0.0001549 | 0.0042225 | -0.00025 | -7.9E-05 |
| Cellular Processes;Cellular community - prokaryotes;Biofilm_formation-Escherichia_coli | 0.002944 | 0.00017 | 0.002446 | 0.0000315 | 0.0009901 | 0.0060087 | 0.0000158 | 0.000837 |
| Metabolism;Metabolism of other amino acids;Glutathione_metabolism | 0.002308 | 0.000124 | 0.002655 | 0.0000202 | 0.0006697 | 0.0057064 | -0.00057 | -0.00013 |
| Metabolism;Amino acid metabolism;Histidine_metabolism | 0.002726 | 0.000184 | 0.002115 | 0.0000282 | 0.0001788 | 0.0042225 | 0.0000299 | 0.000924 |
| Metabolism;Xenobiotics biodegradation and metabolism;Drug_metabolism-other_enzymes | 0.002595 | 4.74E-05 | 0.002414 | 0.0000157 | 0.0035973 | 0.0098192 | 1.66E-05 | 0.000346 |
| Metabolism;Metabolism of other amino acids;Cyanoamino_acid_metabolism | 0.002675 | 7.72E-05 | 0.002356 | 0.0000246 | 0.0023199 | 0.0082607 | 6.12E-05 | 0.000577 |
| Unclassified;Cellular processes and signaling;Cell_growth | 0.002913 | 0.000282 | 0.002442 | 0.0000367 | 0.0033195 | 0.0095267 | 4.65E-05 | 0.000897 |
| Cellular Processes;Cell growth and death;Necroptosis | 0.002668 | 0.000129 | 0.002208 | 0.0000181 | 0.0006 | 0.00422 | 0.0000255 | 0.000665 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | | 6 | 25 | | |
| Genetic Information Processing;Folding, sorting and degradation;Sulfur_relay_system | 0.002033 | 6.24E-05 | 0.002222 | 0.000115 | 0.0008311 | 0.0057414 | -0.00031 | -6.4E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Monobactam_biosynthesis | 0.002121 | 3.47E-05 | 0.001984 | 8.96E-05 | 0.0011592 | 0.0061459 | 4.23E-05 | 0.000231 |
| Metabolism;Carbohydrate metabolism;C5-Branched_dibasic_acid_metabolism | 0.002032 | 0.000102 | 0.001717 | 0.000241 | 0.0022336 | 0.0080739 | 6.07E-05 | 0.00057 |
| Cellular Processes;Cellular community - prokaryotes;Biofilm_formation-Vibrio_cholerae | 0.001887 | 0.0001 | 0.001714 | 0.00012 | 0.0022317 | 0.0080739 | 3.04E-05 | 0.000315 |
| Metabolism;Amino acid metabolism;Phenylalanine_metabolism | 0.001845 | 0.000192 | 0.00157 | 9.34E-05 | 0.0015359 | 0.0070468 | 7.01E-05 | 0.000479 |
| Human Diseases;Endocrine and metabolic diseases;Insulin_resistance | 0.001786 | 0.000103 | 0.001495 | 0.000116 | 0.0001045 | 0.0042225 | 0.000149 | 0.000433 |
| Metabolism;Metabolism of terpenoids and polyketides;Prenyltransferases | 0.001408 | 5.91E-05 | 0.001516 | 8.12E-05 | 0.0026094 | 0.0090224 | -0.0002 | -1.6E-05 |
| Metabolism;Metabolism of other amino acids;D-Glutamine_and_D-glutamate_metabolism | 0.001416 | 4.8E-05 | 0.001524 | 5.98E-05 | 0.0006665 | 0.0057064 | -0.00018 | -3.8E-05 |
| Environmental Information Processing;Signal transduction;HIF-1_signaling_pathway | 0.001241 | 3.17E-05 | 0.001343 | 9.34E-05 | 0.0043249 | 0.103127 | -0.0002 | -4.3E-06 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosphingolipid_biosynthesis-globo_and_isoglobo_series | 0.001423 | 0.000143 | 0.001106 | 0.000281 | 0.0041764 | 0.102535 | 1.54E-05 | 0.000618 |
| Metabolism;Biosynthesis of other secondary metabolites;Phenylpropanoid_biosynthesis | 0.001572 | 6.81E-05 | 0.001159 | 0.000289 | 0.0016311 | 0.0072413 | 0.00011 | 0.000715 |
| Organismal Systems;Nervous system;GABAergic_synapse | 0.001346 | 3.61E-05 | 0.001224 | 7.85E-05 | 0.0010521 | 0.0060087 | 3.86E-05 | 0.000205 |
| Metabolism;Metabolism of cofactors and vitamins;Vitamin_B6_metabolism | 0.00127 | 5.5E-05 | 0.001084 | 0.000119 | 0.0010047 | 0.0060087 | 6E-05 | 0.000312 |
| Organismal Systems;Nervous system;Glutamatergic_synapse | 0.001287 | 3.41E-05 | 0.001179 | 7.72E-05 | 0.0016694 | 0.0072413 | 2.65E-05 | 0.00019 |
| Metabolism;Enzyme | 0.0013 | 7.07E- | 0.001 | 0.00 | 0.0 | 0.0 | 5.64 | 0.000 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| families;Protein_phosphatase_and_associated_proteins | 01 | 05 | 114 | 012 | 10832 | 60087 | E-05 | 319 |
| Unclassified;Metabolism;Carbohydrate_metabolism | 0.001315 | 0.000147 | 0.001019 | 0.0000118 | 0.00351 | 0.042225 | 0.000123 | 0.000468 |
| Organismal Systems;Endocrine system;Insulin_signaling_pathway | 0.001273 | 8.54E-05 | 0.00102 | 0.0000129 | 0.0003262 | 0.042225 | 0.00011 | 0.000397 |
| Organismal Systems;Endocrine system;PPAR_signaling_pathway | 0.001209 | 6.31E-05 | 0.000931 | 0.0000165 | 0.0007484 | 0.057414 | 0.000104 | 0.000452 |
| Metabolism;Metabolism of other amino acids;beta-Alanine_metabolism | 0.001119 | 0.000151 | 0.000946 | 6.77E-05 | 0.0038154 | 0.10003 | 1.25E-05 | 0.000333 |
| Unclassified;Viral protein family;Viral_proteins | 0.00087 | 4.86E-05 | 0.000995 | 5.84E-05 | 0.0002547 | 0.042225 | -0.00019 | -5.6E-05 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_degradation | 0.000925 | 0.000151 | 0.000591 | 0.0000218 | 0.0013149 | 0.065517 | 8.89E-05 | 0.000579 |
| Metabolism;Biosynthesis of other secondary metabolites;Novobiocin_biosynthesis | 0.001016 | 5.43E-05 | 0.00082 | 0.0000101 | 0.0003339 | 0.042225 | 8.73E-05 | 0.000305 |
| Metabolism;Biosynthesis of other secondary metabolites;Tropane,_piperidine_and_pyridine_alkaloid_biosynthesis | 0.001043 | 6.63E-05 | 0.000805 | 7.65E-05 | 0.000202 | 0.042225 | 0.000145 | 0.00033 |
| Cellular Processes;Transport and catabolism;Lysosome | 0.00084 | 0.0002 | 0.000516 | 0.0000241 | 0.0030551 | 0.093446 | 3.74E-05 | 0.00061 |
| Metabolism;Carbohydrate metabolism;Ascorbate_and_aldarate_metabolism | 0.000764 | 6.21E-05 | 0.000622 | 8.97E-05 | 0.0011082 | 0.060087 | 4.14E-05 | 0.000243 |
| Metabolism;Biosynthesis of other secondary metabolites;Acarbose_and_validamycin_biosynthesis | 0.000751 | 2.85E-05 | 0.000653 | 9.16E-05 | 0.0045896 | 0.10735 | 2.48E-06 | 0.000194 |
| Metabolism;Lipid metabolism;Synthesis_and_degradation_of_ketone_bodies | 0.000626 | 3.57E-05 | 0.000706 | 7.54E-05 | 0.0049832 | 0.108091 | -0.00016 | -7.7E-08 |
| Environmental Information Processing;Signal transduction;AMPK_signaling_pathway | 0.000694 | 4.69E-05 | 0.000623 | 3.94E-05 | 0.0018262 | 0.076435 | 1.49E-05 | 0.000127 |
| Unclassified;Metabolism;Glycan_biosynthesis_and_metabolism | 0.000642 | 4.08E-05 | 0.000512 | 6.35E-05 | 0.0002595 | 0.042225 | 5.91E-05 | 0.0002 |

| Pathway | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Environmental Information Processing;Signal transduction;Phosphatidylinositol_signaling_system | 0.000471 | 3.16E-05 | 0.000509 | 1.46E-05 | 0.030116 | 0.093311 | -7.2E-05 | -4.9E-06 |
| Cellular Processes;Cell growth and death;Ferroptosis | 0.000572 | 3.2E-05 | 0.00033 | 0.0000121 | 0.0038 12 | 0.042225 | 0.0000114 | 0.000368 |
| Genetic Information Processing;Folding, sorting and degradation;Proteasome | 0.000566 | 5.42E-05 | 0.000444 | 0.0000112 | 0.047251 | 0.108059 | 1.93E-06 | 0.000241 |
| Organismal Systems;Endocrine system;Adipocytokine_signaling_pathway | 0.000557 | 4.04E-05 | 0.000303 | 0.00013 | 0.0038 | 0.042225 | 0.0000118 | 0.00039 |
| Metabolism;Biosynthesis of other secondary metabolites;Isoquinoline_alkaloid_biosynthesis | 0.000477 | 1.95E-05 | 0.000386 | 6.85E-05 | 0.020656 | 0.078274 | 1.99E-05 | 0.000163 |
| Metabolism;Lipid metabolism;Secondary_bile_acid_biosynthesis | 0.000335 | 2.13E-05 | 0.000417 | 6.43E-05 | 0.025372 | 0.089018 | -0.000015 | -1.4E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Glucosinolate_biosynthesis | 0.000341 | 3.28E-05 | 0.000265 | 2.64E-05 | 0.014 54 | 0.042225 | 3.73E-05 | 0.000114 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosylphosphatidylinositol_(GPI)-anchored_proteins | 0.000296 | 4.79E-05 | 0.000188 | 5.55E-05 | 0.048 04 | 0.045425 | 4.17E-05 | 0.000175 |
| Metabolism;Glycan biosynthesis and metabolism;N-Glycan_biosynthesis | 0.000331 | 9.81E-06 | 0.000253 | 4.94E-05 | 0.010931 | 0.060087 | 2.64E-05 | 0.00013 |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_siderophore_group_nonribosomal_peptides | 0.000249 | 6.71E-05 | 0.000164 | 3.97E-05 | 0.028492 | 0.090633 | 1.14E-05 | 0.000158 |
| Human Diseases;Cancers;Renal_cell_carcinoma | 0.000173 | 2.1E-05 | 0.000284 | 5.25E-05 | 0.002352 | 0.042225 | -0.000017 | -5.5E-05 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_binding_proteins | 0.000171 | 1.89E-05 | 0.000106 | 2.46E-05 | 0.005 44 | 0.042225 | 3.65E-05 | 9.35E-05 |
| Organismal Systems;Digestive system;Protein_digestion_and_absorption | 0.000241 | 5.39E-05 | 0.000103 | 0.000101 | 0.0194 33 | 0.078274 | 2.92E-05 | 0.000247 |
| Metabolism;Lipid metabolism;Steroid_hormone_biosynthesis | 0.000174 | 6.14E-05 | 8.44E-05 | 7.21E-05 | 0.043251 | 0.103127 | 3.33E-06 | 0.000176 |
| Environmental Information Processing;Signaling molecules and interaction;CD_Molecules | 0.000238 | 5.41E-05 | 9.98E-05 | 0.000103 | 0.0204 | 0.0782 | 2.82E-05 | 0.000249 |

| Description | | | | | | | | |
|---|---|---|---|---|---|---|---|
| | | | | | 03 | 74 | | |
| Organismal Systems;Digestive system;Carbohydrate_digestion_and_absorption | 0.000114 | 3.23E-05 | 7.38E-05 | 1.73E-05 | 0.028466 | 0.090633 | 5.52E-06 | 7.5E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Flavone_and_flavonol_biosynthesis | 0.000118 | 8.7E-06 | 9.64E-05 | 1.63E-05 | 0.020433 | 0.078274 | 4.45E-06 | 3.95E-05 |
| Human Diseases;Cancers;Choline_metabolism_in_cancer | 9.26E-05 | 7.18E-06 | 0.000109 | 1.08E-05 | 0.013456 | 0.065517 | -2.8E-05 | -4.3E-06 |
| Environmental Information Processing;Signal transduction;Phospholipase_D_signaling_pathway | 7.63E-05 | 6.87E-06 | 9.25E-05 | 1.34E-05 | 0.031262 | 0.094411 | -3.1E-05 | -1.9E-06 |
| Genetic Information Processing;Transcription;Basal_transcription_factors | 7.48E-05 | 9.1E-06 | 6.1E-05 | 1.11E-05 | 0.040494 | 0.101038 | 7.35E-07 | 2.69E-05 |
| Organismal Systems;Endocrine system;Renin-angiotensin_system | 6.03E-05 | 1.93E-05 | 2.88E-05 | 2.59E-05 | 0.039527 | 0.101038 | 1.87E-06 | 6.12E-05 |
| Human Diseases;Infectious diseases;African_trypanosomiasis | 3.1E-05 | 1.36E-05 | 7.93E-06 | 3.38E-06 | 0.00787 | 0.057414 | 8.83E-06 | 3.73E-05 |
| Human Diseases;Infectious diseases;Chagas_disease_(American_trypanosomiasis) | 2.83E-05 | 1.31E-05 | 5.73E-06 | 3.61E-06 | 0.007169 | 0.057414 | 8.87E-06 | 3.63E-05 |
| Human Diseases;Neurodegenerative diseases;Prion_diseases | 2.71E-05 | 3.06E-06 | 2.21E-05 | 4.03E-06 | 0.036218 | 0.098192 | 4.01E-07 | 9.7E-06 |
| Organismal Systems;Digestive system;Mineral_absorption | 1.16E-05 | 1.93E-06 | 8.81E-06 | 1.93E-06 | 0.029988 | 0.09331 | 3.34E-07 | 5.3E-06 |
| Metabolism;Metabolism of terpenoids and polyketides;Type_I_polyketide_structures | 9.57E-06 | 5.92E-06 | 3.27E-06 | 1.48E-06 | 0.047558 | 0.108059 | 9.56E-08 | 1.25E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Stilbenoid,_diarylheptanoid_and_gingerol_biosynthesis | 3.46E-06 | 5.49E-07 | 1.76E-06 | 9.04E-07 | 0.0040474 | 0.042225 | 7.08E-07 | 2.69E-06 |
| Metabolism;Biosynthesis of other secondary metabolites;Flavonoid_biosynthesis | 3.46E-06 | 5.49E-07 | 1.76E-06 | 9.04E-07 | 0.0040474 | 0.042225 | 7.08E-07 | 2.69E-06 |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_12-,_14-_and_16-membered_macrolides | 3.33E-06 | 2.17E-06 | 1.02E-06 | 4.86E-07 | 0.047456 | 0.108059 | 3.67E-08 | 4.57E-06 |
| Metabolism;Metabolism of terpenoids and | 2.58E- | 1.55E- | 8.51E | 4.94 | 0.0 | 0.1 | 1.03 | 3.35 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| polyketides;Biosynthesis_of_enediyne_antibiotics | 06 | 06 | -07 | E-07 | 40537 | 01038 | E-07 | E-06 |
| Human Diseases;Immune diseases;Systemic_lupus_erythematosus | 5.61E-07 | 2.55E-07 | 2.46E-07 | 9.81E-08 | 0.028032 | 0.090633 | 4.64E-08 | 5.83E-07 |
| Human Diseases;Infectious diseases;Vibrio_cholerae_infection | 7.42E-07 | 4.06E-07 | 2.66E-07 | 2.15E-07 | 0.036006 | 0.098192 | 4.04E-08 | 9.12E-07 |
| Human Diseases;Infectious diseases;Pathogenic_Escherichia_coli_infection | 6E-07 | 3.16E-07 | 1.71E-07 | 8.14E-08 | 0.019688 | 0.078274 | 9.83E-08 | 7.61E-07 |
| Organismal Systems;Endocrine system;Ovarian_Steroidogenesis | 3.18E-07 | 2.43E-07 | 6.29E-08 | 5.25E-08 | 0.049664 | 0.108091 | 5.32E-10 | 5.09E-07 |
| Environmental Information Processing;Signal transduction;Calcium_signaling_pathway | 7.35E-08 | 4.47E-08 | 1.4E-08 | 1.79E-08 | 0.02067 | 0.078274 | 1.24E-08 | 1.07E-07 |
| Environmental Information Processing;Signaling molecules and interaction;G_protein-coupled_receptors | 2.34E-08 | 1.07E-08 | 2.05E-09 | 3.26E-09 | 0.003543 | 0.042225 | 1.02E-08 | 3.26E-08 |
| Environmental Information Processing;Signaling molecules and interaction;Neuroactive_ligand-receptor_interaction | 5.88E-09 | 4.51E-09 | 1.14E-09 | 1.32E-09 | 0.049187 | 0.108091 | 2.35E-11 | 9.47E-09 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_biosynthesis-chondroitin_sulfate/dermatan_sulfate | 2.39E-09 | 1.68E-09 | 5.21E-10 | 8.36E-10 | 0.043658 | 0.103127 | 6.93E-11 | 3.66E-09 |
| Metabolism;Biosynthesis of other secondary metabolites;Clavulanic_acid_biosynthesis | 7.9E-10 | 6.83E-10 | 1.4E-11 | 1.12E-11 | 0.038774 | 0.100551 | 5.91E-11 | 1.49E-09 |

Table 2b Young+PBS VS Aged+PBS

| Taxa | avg(Young+PBS) | sd(Young+PBS) | avg(Aged+PBS) | sd(Aged+PBS) | p.value | q.values | interval lower | interval upper |
|---|---|---|---|---|---|---|---|---|
| Metabolism;Nucleotide metabolism;Purine_metabolism | 0.025653 | 0.000514 | 0.026306 | 0.00031 | 0.027754 | 0.018099 | -0.00122 | -9.1E-05 |
| Genetic Information Processing;Translation;Transfer_RNA_biogenesis | 0.025388 | 0.000319 | 0.025764 | 0.000192 | 0.037679 | 0.020034 | -0.00072 | -2.7E-05 |
| Metabolism;Nucleotide metabolism;Pyrimidine_metabolism | 0.020031 | 0.000437 | 0.020588 | 0.000135 | 0.024899 | 0.016912 | -0.00101 | -9.9E-05 |
| Genetic Information Processing;Translation;Ribosome | 0.015723 | 0.000269 | 0.016031 | 0.000179 | 0.045392 | 0.02079 | -0.00061 | -7.9E-06 |
| Genetic Information Processing;Translation;Aminoacyl-tRNA_biosynthesis | 0.015499 | 0.000262 | 0.015904 | 0.000183 | 0.012754 | 0.011118 | -0.0007 | -0.00011 |
| Cellular Processes;Transport and catabolism;Exosome | 0.015496 | 0.000151 | 0.01467 | 0.000299 | 0.000423 | 0.001644 | 0.000506 | 0.001146 |
| Genetic Information Processing;Replication and repair;DNA_replication_proteins | 0.014807 | 0.000271 | 0.015109 | 0.000133 | 0.042432 | 0.020702 | -0.00059 | -1.4E-05 |
| Genetic Information Processing;Replication and repair;Chromosome_and_associated_proteins | 0.0131 | 0.000238 | 0.013723 | 0.000145 | 0.000519 | 0.017088 | -0.00088 | -0.00036 |
| Metabolism;Amino acid metabolism;Alanine,_aspartate_and_glutamate_metabolism | 0.011744 | 0.000114 | 0.011119 | 0.000336 | 0.004746 | 0.006652 | 0.000273 | 0.000978 |
| Genetic Information Processing;Replication and repair;Mismatch_repair | 0.010868 | 0.000292 | 0.011191 | 0.000174 | 0.047672 | 0.020993 | -0.00064 | -4.2E-06 |
| Genetic Information Processing;Replication and repair;Homologous_recombination | 0.010494 | 0.000212 | 0.010894 | 0.000219 | 0.009155 | 0.009537 | -0.00068 | -0.00012 |
| Metabolism;Glycan biosynthesis and metabolism;Peptidoglycan_biosynthesis_and_degradation_proteins | 0.010172 | 0.000138 | 0.010573 | 0.000247 | 0.008688 | 0.009515 | -0.00067 | -0.00013 |
| Metabolism;Amino acid metabolism;Cysteine_and_methionine_metabolism | 0.009602 | 7.4E-05 | 0.009323 | 5.47E-05 | 3.49E-05 | 0.00528 | 0.000195 | 0.000364 |
| Metabolism;Energy | 0.0091 | 0.000 | 0.008 | 0.000 | 0.0 | 0.0 | 1.72 | 0.001 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| metabolism;Carbon_fixation_pathways_in_prokaryotes | 8 | 366 | 631 | 453 | 44292 | 20702 | E-05 | 082 |
| Metabolism;Glycan biosynthesis and metabolism;Peptidoglycan_biosynthesis | 0.008308 | 0.000207 | 0.008721 | 0.000328 | 0.029895 | 0.018642 | -0.00077 | -5.1E-05 |
| Genetic Information Processing;Replication and repair;DNA_replication | 0.007972 | 0.000183 | 0.00835 | 0.000115 | 0.02382 | 0.04186 | -0.00058 | -0.00018 |
| Genetic Information Processing;Translation;Messenger_RNA_Biogenesis | 0.007849 | 5.8E-05 | 0.007653 | 0.000128 | 0.01143 | 0.10236 | 6.06E-05 | 0.000332 |
| Metabolism;Amino acid metabolism;Glycine,_serine_and_threonine_metabolism | 0.007893 | 0.000123 | 0.0077 | 0.000166 | 0.047234 | 0.020993 | 2.91E-06 | 0.000383 |
| Metabolism;Carbohydrate metabolism;Pentose_phosphate_pathway | 0.007552 | 0.000179 | 0.008037 | 0.000253 | 0.004043 | 0.006168 | -0.00077 | -0.0002 |
| Genetic Information Processing;Folding, sorting and degradation;RNA_degradation | 0.007475 | 8.88E-05 | 0.007297 | 0.000143 | 0.030419 | 0.018733 | 2.15E-05 | 0.000336 |
| Metabolism;Carbohydrate metabolism;Butanoate_metabolism | 0.007496 | 0.000432 | 0.006828 | 0.000131 | 0.011301 | 0.01027 | 0.000216 | 0.00112 |
| Genetic Information Processing;Transcription;Transcription_machinery | 0.007439 | 9.11E-05 | 0.006994 | 0.00041 | 0.044514 | 0.020702 | 1.55E-05 | 0.000874 |
| Metabolism;Carbohydrate metabolism;Propanoate_metabolism | 0.006937 | 0.000132 | 0.007178 | 9.95E-05 | 0.005804 | 0.007512 | -0.00039 | -8.8E-05 |
| Metabolism;Carbohydrate metabolism;Glyoxylate_and_dicarboxylate_metabolism | 0.007261 | 0.000272 | 0.006487 | 0.000322 | 0.001233 | 0.003019 | 0.000389 | 0.001159 |
| Metabolism;Carbohydrate metabolism;Citrate_cycle_(TCA_cycle) | 0.006205 | 0.000286 | 0.005779 | 0.000294 | 0.029455 | 0.018636 | 5.2E-05 | 0.000798 |
| Unclassified;Metabolism;Energy_metabolism | 0.006315 | 0.000289 | 0.005571 | 0.00032 | 0.017779 | 0.034555 | 0.000352 | 0.001137 |
| Unclassified;Poorly characterized;General_function_prediction_only | 0.005257 | 0.000243 | 0.006104 | 0.00068 | 0.027117 | 0.017958 | -0.00156 | -0.00013 |
| Metabolism;Energy metabolism;Carbon_fixation_in_photosynthetic_organisms | 0.005094 | 4.39E-05 | 0.004982 | 4.4E-05 | 0.012780 | 0.030322 | 5.58E-05 | 0.000169 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Metabolism;Amino acid metabolism;Arginine_biosynthesis | 0.005337 | 0.000206 | 0.004855 | 0.000135 | 0.0011101 | 0.003019 | 0.000253 | 0.000711 |
| Metabolism;Amino acid metabolism;Phenylalanine,_tyrosine_and_tryptophan_biosynthesis | 0.004979 | 0.00031 | 0.004465 | 0.000266 | 0.0011991 | 0.01067 | 0.000141 | 0.000886 |
| Genetic Information Processing;Folding, sorting and degradation;Protein_export | 0.004501 | 8.55E-05 | 0.00462 | 8.29E-05 | 0.0034534 | 0.0020034 | -0.000023 | -1.1E-05 |
| Metabolism;Lipid metabolism;Lipid_biosynthesis_proteins | 0.004748 | 0.000253 | 0.004339 | 8.86E-05 | 0.0091 27 | 0.0095 37 | 0.000143 | 0.000675 |
| Metabolism;Metabolism of terpenoids and polyketides;Terpenoid_backbone_biosynthesis | 0.004268 | 0.000118 | 0.004426 | 4.29E-05 | 0.0205 07 | 0.0157 82 | -0.00028 | -3.4E-05 |
| Metabolism;Metabolism of cofactors and vitamins;Thiamine_metabolism | 0.004472 | 6.82E-05 | 0.004227 | 0.000161 | 0.0116 82 | 0.0105 05 | 7.47E-05 | 0.000415 |
| Metabolism;Lipid metabolism;Glycerophospholipid_metabolism | 0.004116 | 0.00012 | 0.004402 | 0.00011 | 0.0015 78 | 0.0033 06 | -0.00043 | -0.00014 |
| Metabolism;Energy metabolism;Nitrogen_metabolism | 0.004843 | 0.000271 | 0.004069 | 0.000223 | 0.0003 36 | 0.0015 53 | 0.000454 | 0.001095 |
| Metabolism;Metabolism of cofactors and vitamins;One_carbon_pool_by_folate | 0.004295 | 4.61E-05 | 0.003978 | 7.5E-05 | 1.69E-05 | 0.0005 28 | 0.000235 | 0.0004 |
| Genetic Information Processing;Replication and repair;Base_excision_repair | 0.003915 | 6.37E-05 | 0.004133 | 8.09E-05 | 0.0004 85 | 0.0016 58 | -0.00031 | -0.00012 |
| Metabolism;Metabolism of other amino acids;Selenocompound_metabolism | 0.004118 | 5.86E-05 | 0.004009 | 5.9E-05 | 0.0091 12 | 0.0095 37 | 3.38E-05 | 0.000185 |
| Metabolism;Metabolism of cofactors and vitamins;Pantothenate_and_CoA_biosynthesis | 0.004248 | 0.000113 | 0.003982 | 4.2E-05 | 0.0013 55 | 0.0030 46 | 0.000147 | 0.000384 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosyltransferases | 0.003709 | 0.000154 | 0.003989 | 0.000158 | 0.0111 38 | 0.0102 36 | -0.00048 | -7.9E-05 |
| Metabolism;Amino acid metabolism;Valine,_leucine_and_isoleucine_biosynthesis | 0.003878 | 0.000305 | 0.003302 | 0.000124 | 0.0041 86 | 0.0062 51 | 0.000254 | 0.000898 |
| Genetic Information Processing;Transcription;RNA_polymerase | 0.003305 | 7E-05 | 0.003427 | 4.68E-05 | 0.0065 | 0.0079 | -0.0002 | -4.4E-05 |

| Pathway | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | | 32 | | | |
| Metabolism;Carbohydrate metabolism;Pentose_and_glucuronate_interconversions | 0.003491 | 7.67E-05 | 0.003163 | 0.000104 | 0.00014 | 0.00098 | 0.000209 | 0.000446 |
| Metabolism;Lipid metabolism;Sphingolipid_metabolism | 0.003031 | 0.000248 | 0.00258 | 0.000251 | 0.01058 | 0.01043 | 0.000131 | 0.000773 |
| Metabolism;Metabolism of cofactors and vitamins;Folate_biosynthesis | 0.002563 | 5.22E-05 | 0.002807 | 0.000132 | 0.004674 | 0.006652 | -0.00038 | -0.00011 |
| Metabolism;Biosynthesis of other secondary metabolites;Streptomycin_biosynthesis | 0.002693 | 4.16E-05 | 0.002542 | 4.81E-05 | 0.000182 | 0.001049 | 9.31E-05 | 0.000209 |
| Cellular Processes;Cellular community - prokaryotes;Biofilm_formation-Escherichia_coli | 0.002944 | 0.00017 | 0.002448 | 6.37E-05 | 0.00042 | 0.001644 | 0.000317 | 0.000675 |
| Metabolism;Metabolism of other amino acids;Glutathione_metabolism | 0.002308 | 0.000124 | 0.002741 | 0.000188 | 0.001237 | 0.003019 | -0.00064 | -0.00022 |
| Metabolism;Energy metabolism;Sulfur_metabolism | 0.0026 | 9.62E-05 | 0.00237 | 6.26E-05 | 0.000966 | 0.002947 | 0.000123 | 0.000337 |
| Metabolism;Amino acid metabolism;Histidine_metabolism | 0.002726 | 0.000184 | 0.002265 | 6.44E-05 | 0.001014 | 0.002987 | 0.000269 | 0.000654 |
| Metabolism;Xenobiotics biodegradation and metabolism;Drug_metabolism-other_enzymes | 0.002595 | 4.74E-05 | 0.002421 | 5.19E-05 | 0.000123 | 0.00098 | 0.00011 | 0.000238 |
| Metabolism;Metabolism of other amino acids;Cyanoamino_acid_metabolism | 0.002675 | 7.72E-05 | 0.002357 | 0.000201 | 0.009879 | 0.009706 | 0.000106 | 0.00053 |
| Unclassified;Cellular processes and signaling;Cell_growth | 0.002913 | 0.000282 | 0.002427 | 0.000246 | 0.009998 | 0.009706 | 0.000145 | 0.000827 |
| Cellular Processes;Cell growth and death;Necroptosis | 0.002668 | 0.000129 | 0.002126 | 0.000197 | 0.000376 | 0.001607 | 0.000323 | 0.000761 |
| Genetic Information Processing;Folding, sorting and degradation;Sulfur_relay_system | 0.002033 | 6.24E-05 | 0.002168 | 1.8E-05 | 0.002388 | 0.004186 | -0.0002 | -7E-05 |
| Organismal Systems;Aging;Longevity_regulating_pathway-worm | 0.001962 | 1.51E-05 | 0.002049 | 4.11E-05 | 0.002488 | 0.004186 | -0.00013 | -4.3E-05 |
| Metabolism;Biosynthesis of other secondary | 0.0021 | 3.47E- | 0.001 | 8.39E | 0.0 | 0.0 | 6.45 | 0.000 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| metabolites;Monobactam_biosynthesis | 21 | 05 | 968 | -05 | 048 92 | 067 41 | E-05 | 242 |
| Metabolism;Carbohydrate metabolism;C5-Branched_dibasic_acid_metabolism | 0.002032 | 0.000102 | 0.001784 | 8.8E-05 | 0.0011 46 | 0.0030 19 | 0.000126 | 0.000371 |
| Unclassified;Cellular processes and signaling;Membrane_and_intracellular_structural_molecules | 0.001747 | 7.54E-05 | 0.001925 | 0.000136 | 0.0233 41 | 0.0169 12 | -0.00032 | -3.1E-05 |
| Cellular Processes;Cellular community - prokaryotes;Biofilm_formation-Vibrio_cholerae | 0.001887 | 0.0001 | 0.001767 | 6.28E-05 | 0.0355 97 | 0.0200 34 | 1.03E-05 | 0.000231 |
| Human Diseases;Drug resistance;Antimicrobial_resistance_genes | 0.001562 | 0.00013 | 0.001938 | 0.000318 | 0.0333 63 | 0.0195 22 | -0.00071 | -4E-05 |
| Human Diseases;Drug resistance;Cationic_antimicrobial_peptide_(CAMP)_resistance | 0.001578 | 0.000133 | 0.001938 | 0.000162 | 0.0019 51 | 0.0037 04 | -0.00055 | -0.00017 |
| Metabolism;Amino acid metabolism;Phenylalanine_metabolism | 0.001845 | 0.000192 | 0.001538 | 7.37E-05 | 0.0093 | 0.0095 72 | 0.000105 | 0.000509 |
| Human Diseases;Endocrine and metabolic diseases;Insulin_resistance | 0.001786 | 0.000103 | 0.001469 | 4.21E-05 | 0.0002 86 | 0.0014 39 | 0.000208 | 0.000426 |
| Metabolism;Metabolism of terpenoids and polyketides;Prenyltransferases | 0.001408 | 5.91E-05 | 0.001515 | 2.26E-05 | 0.0050 42 | 0.0068 37 | -0.00017 | -4.6E-05 |
| Metabolism;Metabolism of other amino acids;D-Glutamine_and_D-glutamate_metabolism | 0.001416 | 4.8E-05 | 0.001472 | 3.06E-05 | 0.0413 77 | 0.0207 02 | -0.00011 | -2.8E-06 |
| Environmental Information Processing;Signal transduction;HIF-1_signaling_pathway | 0.001241 | 3.17E-05 | 0.001378 | 8.59E-05 | 0.0095 58 | 0.0096 06 | -0.00023 | -4.7E-05 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosphingolipid_biosynthesis-globo_and_isoglobo_series | 0.001423 | 0.000143 | 0.001238 | 7.89E-05 | 0.0246 77 | 0.0169 12 | 3.06E-05 | 0.000339 |
| Metabolism;Metabolism of other amino acids;Taurine_and_hypotaurine_metabolism | 0.001248 | 2.41E-05 | 0.001392 | 6.39E-05 | 0.0017 2 | 0.0034 55 | -0.00021 | -7.7E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Phenylpropanoid_biosynthesis | 0.001572 | 6.81E-05 | 0.001167 | 0.000172 | 0.0013 39 | 0.0030 46 | 0.000223 | 0.000587 |
| Metabolism;Metabolism of other amino acids;D-Alanine_metabolism | 0.001168 | 4.12E-05 | 0.001297 | 8.82E-05 | 0.0138 15 | 0.0115 7 | -0.00022 | -3.5E-05 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Cellular Processes;Transport and catabolism;Peroxisome | 0.001301 | 6.36E-05 | 0.001189 | 8.51E-05 | 0.029136 | 0.018575 | 1.41E-05 | 0.00021 |
| Organismal Systems;Nervous system;GABAergic_synapse | 0.001346 | 3.61E-05 | 0.001162 | 0.000101 | 0.00527 | 0.007034 | 7.72E-05 | 0.00029 |
| Human Diseases;Infectious diseases;Staphylococcus_aureus_infection | 0.000788 | 0.000143 | 0.001567 | 0.000581 | 0.020721 | 0.015805 | -0.00139 | -0.00017 |
| Metabolism;Metabolism of terpenoids and polyketides;Polyketide_sugar_unit_biosynthesis | 0.001273 | 3.92E-05 | 0.001102 | 4.39E-05 | 3.58E-05 | 0.000528 | 0.000117 | 0.000224 |
| Metabolism;Metabolism of cofactors and vitamins;Vitamin_B6_metabolism | 0.00127 | 5.5E-05 | 0.001147 | 5.2E-05 | 0.002645 | 0.004345 | 5.39E-05 | 0.000192 |
| Organismal Systems;Nervous system;Glutamatergic_synapse | 0.001287 | 3.41E-05 | 0.001097 | 0.000116 | 0.008547 | 0.009483 | 6.98E-05 | 0.000312 |
| Metabolism;Enzyme families;Protein_phosphatase_and_associated_proteins | 0.001301 | 7.07E-05 | 0.001026 | 0.000148 | 0.004244 | 0.006251 | 0.000118 | 0.000433 |
| Unclassified;Metabolism;Carbohydrate_metabolism | 0.001315 | 0.000147 | 0.001103 | 3.74E-05 | 0.015668 | 0.012748 | 5.76E-05 | 0.000365 |
| Unclassified;Metabolism;Amino_acid_metabolism | 0.001195 | 5.78E-05 | 0.001107 | 4E-05 | 0.013495 | 0.011414 | 2.31E-05 | 0.000153 |
| Organismal Systems;Endocrine system;Insulin_signaling_pathway | 0.001273 | 8.54E-05 | 0.001 | 6.37E-05 | 0.000126 | 0.00098 | 0.000176 | 0.000372 |
| Metabolism;Metabolism of cofactors and vitamins;Ubiquinone_and_other_terpenoid-quinone_biosynthesis | 0.000957 | 9.74E-05 | 0.001156 | 6.87E-05 | 0.002744 | 0.004423 | -0.00031 | -8.9E-05 |
| Human Diseases;Cardiovascular diseases;Fluid_shear_stress_and_atherosclerosis | 0.001166 | 2.7E-05 | 0.001026 | 9.88E-05 | 0.016498 | 0.013297 | 3.66E-05 | 0.000243 |
| Organismal Systems;Endocrine system;PPAR_signaling_pathway | 0.001209 | 6.31E-05 | 0.000967 | 3.82E-05 | 3.71E-05 | 0.000528 | 0.000172 | 0.00031 |
| Metabolism;Metabolism of other amino acids;beta-Alanine_metabolism | 0.001119 | 0.000151 | 0.000955 | 4.08E-05 | 0.004471 | 0.020702 | 5.42E-06 | 0.000322 |
| Environmental Information Processing;Signaling molecules and interaction;Bacterial_toxins | 0.000828 | 8.51E-05 | 0.001118 | 0.000234 | 0.00277 | 0.00180 | -0.00054 | -4.4E-05 |

| | | | | | | 02 | 99 | | |
|---|---|---|---|---|---|---|---|---|
| Unclassified;Viral protein family;Viral_proteins | 0.00087 | 4.86E-05 | 0.001018 | 4.82E-05 | 0.000345 | 0.001553 | -0.00021 | -8.6E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Novobiocin_biosynthesis | 0.001016 | 5.43E-05 | 0.000857 | 5.38E-05 | 0.000468 | 0.001658 | 8.94E-05 | 0.000229 |
| Metabolism;Carbohydrate metabolism;Inositol_phosphate_metabolism | 0.000854 | 5.25E-05 | 0.000953 | 8.36E-05 | 0.00038817 | 0.00020034 | -0.00019 | -6.4E-06 |
| Metabolism;Biosynthesis of other secondary metabolites;Tropane,_piperidine_and_pyridine_alkaloid_biosynthesis | 0.001043 | 6.63E-05 | 0.000836 | 5.62E-05 | 0.000184 | 0.001049 | 0.000127 | 0.000286 |
| Cellular Processes;Cellular community - prokaryotes;Biofilm_formation-Pseudomonas_aeruginosa | 0.000931 | 9.62E-05 | 0.000813 | 5E-05 | 0.00030701 | 0.00018733 | 1.43E-05 | 0.000221 |
| Metabolism;Biosynthesis of other secondary metabolites;Acarbose_and_validamycin_biosynthesis | 0.000751 | 2.85E-05 | 0.000637 | 2.76E-05 | 3.4E-05 | 0.000528 | 7.86E-05 | 0.000151 |
| Metabolism;Lipid metabolism;Synthesis_and_degradation_of_ketone_bodies | 0.000626 | 3.57E-05 | 0.000714 | 6.61E-05 | 0.00021124 | 0.0001583 | -0.00016 | -1.7E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Carbapenem_biosynthesis | 0.000704 | 1.62E-05 | 0.000657 | 3.2E-05 | 0.00014556 | 0.00012073 | 1.22E-05 | 8.07E-05 |
| Environmental Information Processing;Signal transduction;AMPK_signaling_pathway | 0.000694 | 4.69E-05 | 0.000558 | 5.71E-05 | 0.00001228 | 0.00003019 | 6.87E-05 | 0.000204 |
| Metabolism;Xenobiotics biodegradation and metabolism;Drug_metabolism-cytochrome_P450 | 0.000515 | 3.98E-05 | 0.000629 | 0.000103 | 0.00042406 | 0.00020702 | -0.00022 | -5.3E-06 |
| Genetic Information Processing;Folding, sorting and degradation;Protein_processing_in_endoplasmic_reticulum | 0.000645 | 2.96E-05 | 0.000512 | 0.000115 | 0.00036719 | 0.00020034 | 1.16E-05 | 0.000253 |
| Metabolism;Xenobiotics biodegradation and metabolism;Metabolism_of_xenobiotics_by_cytochrome_P450 | 0.000478 | 4.11E-05 | 0.00059 | 0.000101 | 0.00041636 | 0.00020702 | -0.00022 | -5.7E-06 |
| Environmental Information Processing;Signal transduction;Phosphatidylinositol_signaling_system | 0.000471 | 3.16E-05 | 0.000551 | 3.83E-05 | 0.00002862 | 0.00004527 | -0.00013 | -3.5E-05 |
| Metabolism;Metabolism of cofactors and vitamins;Retinol_metabolism | 0.000448 | 4.19E-05 | 0.000565 | 9.17E-05 | 0.00024724 | 0.00016912 | -0.00021 | -2E-05 |

| Pathway | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Unclassified;Metabolism;Nucleotide_metabolism | 0.000424 | 6.16E-05 | 0.000559 | 7.62E-05 | 0.0007571 | 0.00874 | -0.00022 | -4.5E-05 |
| Unclassified;Metabolism;Biosynthesis_and_biodegradation_of_secondary_metabolites | 0.000552 | 8.08E-05 | 0.000446 | 4.36E-05 | 0.0022652 | 0.0016682 | 1.95E-05 | 0.000194 |
| Cellular Processes;Cell growth and death;Ferroptosis | 0.000572 | 3.2E-05 | 0.000384 | 4.92E-05 | 3.43E-05 | 0.00528 | 0.000133 | 0.000242 |
| Environmental Information Processing;Signal transduction;PI3K-Akt_signaling_pathway | 0.000531 | 3.76E-05 | 0.000404 | 0.000105 | 0.029838 | 0.018642 | 1.7E-05 | 0.000237 |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_vancomycin_group_antibiotics | 0.000483 | 1.79E-05 | 0.000414 | 2.67E-05 | 0.000537 | 0.0017 | 3.98E-05 | 9.93E-05 |
| Genetic Information Processing;Folding, sorting and degradation;Proteasome | 0.000566 | 5.42E-05 | 0.000391 | 0.000103 | 0.0006859 | 0.0008253 | 6.44E-05 | 0.000286 |
| Organismal Systems;Endocrine system;Adipocytokine_signaling_pathway | 0.000557 | 4.04E-05 | 0.00034 | 6.34E-05 | 7.79E-05 | 0.0008 32 | 0.000147 | 0.000287 |
| Metabolism;Biosynthesis of other secondary metabolites;Isoquinoline_alkaloid_biosynthesis | 0.000477 | 1.95E-05 | 0.000398 | 4.93E-05 | 0.0008874 | 0.0009537 | 2.78E-05 | 0.000132 |
| Human Diseases;Cancers;Prostate_cancer | 0.000502 | 2.81E-05 | 0.000392 | 0.000103 | 0.0046481 | 0.002079 | 2.4E-06 | 0.000217 |
| Organismal Systems;Immune system;IL-17_signaling_pathway | 0.000497 | 2.85E-05 | 0.000387 | 0.000101 | 0.0044528 | 0.0020702 | 3.75E-06 | 0.000216 |
| Organismal Systems;Endocrine system;Estrogen_signaling_pathway | 0.000497 | 2.84E-05 | 0.000387 | 0.000101 | 0.0044832 | 0.0020702 | 3.53E-06 | 0.000216 |
| Organismal Systems;Endocrine system;Progesterone-mediated_oocyte_maturation | 0.000497 | 2.84E-05 | 0.000387 | 0.000101 | 0.0044832 | 0.0020702 | 3.53E-06 | 0.000216 |
| Organismal Systems;Immune system;Th17_cell_differentiation | 0.000497 | 2.84E-05 | 0.000387 | 0.000101 | 0.0044832 | 0.0020702 | 3.53E-06 | 0.000216 |
| Organismal Systems;Immune system;Antigen_processing_and_presentation | 0.000497 | 2.84E-05 | 0.000387 | 0.000101 | 0.0044832 | 0.0020702 | 3.53E-06 | 0.000216 |
| Metabolism;Metabolism of other amino acids;Phosphonate_and_phosphinate_metabolism | 0.000382 | 2.97E-05 | 0.000323 | 2.78E-05 | 0.0056 | 0.0074 | 2.13E-05 | 9.54E-05 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | | 49 | 24 | | |
| Metabolism;Biosynthesis of other secondary metabolites;Glucosinolate_biosynthesis | 0.000341 | 3.28E-05 | 0.000272 | 1.21E-05 | 0.0024227 | 0.0004186 | 3.48E-05 | 0.000104 |
| Unclassified;Genetic information processing;Translation_proteins | 0.000322 | 2.64E-05 | 0.000271 | 1.91E-05 | 0.0036577 | 0.0005568 | 2.15E-05 | 8.15E-05 |
| Genetic Information Processing;Translation;Ribosome_biogenesis_in_eukaryotes | 0.000305 | 7.95E-06 | 0.000294 | 8E-06 | 0.0318838 | 0.0191988 | 1.21E-06 | 2.17E-05 |
| Environmental Information Processing;Signal transduction;FoxO_signaling_pathway | 0.0002 | 3.42E-05 | 0.000294 | 5.37E-05 | 0.0061994 | 0.0076690 | -0.00015 | -3.5E-05 |
| Metabolism;Glycan biosynthesis and metabolism;N-Glycan_biosynthesis | 0.000331 | 9.81E-06 | 0.000253 | 2.52E-05 | 0.0002830 | 0.0014390 | 5.17E-05 | 0.000105 |
| Organismal Systems;Aging;Longevity_regulating_pathway-mammal | 0.00017 | 2.6E-05 | 0.000284 | 5.38E-05 | 0.0022050 | 0.0040940 | -0.00017 | -5.6E-05 |
| Human Diseases;Cancers;Renal_cell_carcinoma | 0.000173 | 2.1E-05 | 0.000251 | 5.73E-06 | 0.0001498 | 0.0009980 | -0.0001 | -5.6E-05 |
| Environmental Information Processing;Signal transduction;MAPK_signaling_pathway-plant | 0.000144 | 2.46E-05 | 0.000255 | 4.18E-05 | 0.0004680 | 0.0016580 | -0.00016 | -6.6E-05 |
| Human Diseases;Neurodegenerative diseases;Huntingtons_disease | 0.000151 | 1.32E-05 | 0.000223 | 5.85E-05 | 0.0284290 | 0.0182660 | -0.00013 | -1.1E-05 |
| Metabolism;Metabolism of terpenoids and polyketides;Nonribosomal_peptide_structures | 0.000151 | 7.28E-06 | 0.000191 | 2.38E-05 | 0.0077760 | 0.0088570 | -6.5E-05 | -1.5E-05 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_binding_proteins | 0.000171 | 1.89E-05 | 0.000129 | 2.28E-05 | 0.0059560 | 0.0075940 | 1.53E-05 | 6.95E-05 |
| Organismal Systems;Digestive system;Protein_digestion_and_absorption | 0.000241 | 5.39E-05 | 0.000134 | 5.61E-05 | 0.0074950 | 0.008740 | 3.53E-05 | 0.000177 |
| Environmental Information Processing;Signaling molecules and interaction;CD_Molecules | 0.000238 | 5.41E-05 | 0.000129 | 5.88E-05 | 0.0073260 | 0.0086930 | 3.68E-05 | 0.000182 |
| Human Diseases;Neurodegenerative diseases;Amyotrophic_lateral_sclerosis_(ALS) | 9.5E-05 | 1.98E-05 | 0.000171 | 3.73E-05 | 0.0024999 | 0.0041860 | -0.00012 | -3.6E-05 |
| Genetic Information Processing;Folding, sorting | 0.0001 | 3.86E- | 0.000 | 4.01E | 0.0 | 0.0 | -1.2E | -2.2E |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| and degradation;Ubiquitin_system | 53 | 06 | 16 | -06 | 094 25 | 095 85 | -05 | -06 |
| Environmental Information Processing;Signal transduction;MAPK_signaling_pathway-yeast | 6.68E-05 | 1.96E-05 | 0.000151 | 3.85E-05 | 0.0017 46 | 0.0034 55 | -0.00 013 | -4.3E-05 |
| Environmental Information Processing;Signal transduction;MAPK_signaling_pathwayfly | 0.000104 | 8.44E-06 | 0.000133 | 1.86E-05 | 0.0099 17 | 0.0097 06 | -4.9E-05 | -9.5E-06 |
| Metabolism;Biosynthesis of other secondary metabolites;Flavone_and_flavonol_biosynthesis | 0.000118 | 8.7E-06 | 0.000105 | 5.92E-06 | 0.0129 97 | 0.0112 16 | 3.58E-06 | 2.31E-05 |
| Human Diseases;Cancers;Choline_metabolism_in_cancer | 9.26E-05 | 7.18E-06 | 0.000117 | 5.01E-06 | 9.04E-05 | 0.008 58 | -3.2E-05 | -1.6E-05 |
| Genetic Information Processing;Replication and repair;Non-homologous_end-joining | 9.66E-05 | 1.61E-05 | 7.88E-05 | 7.5E-06 | 0.0438 16 | 0.0207 02 | 6.49E-07 | 3.49E-05 |
| Environmental Information Processing;Signal transduction;Phospholipase_D_signaling_pathway | 7.63E-05 | 6.87E-06 | 0.000102 | 6.17E-06 | 4.38E-05 | 0.0005 35 | -3.4E-05 | -1.8E-05 |
| Genetic Information Processing;Transcription;Basal_transcription_factors | 7.48E-05 | 9.1E-06 | 5.9E-05 | 4.71E-06 | 0.0061 24 | 0.0076 69 | 6.02E-06 | 2.55E-05 |
| Metabolism;Lipid metabolism;alpha-Linolenic_acid_metabolism | 3.57E-05 | 1.21E-05 | 5.48E-05 | 1.65E-05 | 0.0474 7 | 0.0209 93 | -3.8E-05 | -2.6E-07 |
| Organismal Systems;Endocrine system;Renin-angiotensin_system | 6.03E-05 | 1.93E-05 | 2.79E-05 | 1.29E-05 | 0.0079 66 | 0.0089 54 | 1.09E-05 | 5.39E-05 |
| Cellular Processes;Cell growth and death;Meiosis-yeast | 3.84E-05 | 1.22E-05 | 1.95E-05 | 1.4E-06 | 0.0123 94 | 0.0109 16 | 6.11E-06 | 3.16E-05 |
| Metabolism;Xenobiotics biodegradation and metabolism;Steroid_degradation | 4.6E-05 | 2.1E-05 | 2.09E-05 | 2.37E-06 | 0.0326 | 0.0193 4 | 3.06E-06 | 4.71E-05 |
| Human Diseases;Infectious diseases;African_trypanosomiasis | 3.1E-05 | 1.36E-05 | 1.27E-05 | 5.55E-06 | 0.0199 18 | 0.0154 69 | 3.94E-06 | 3.26E-05 |
| Human Diseases;Infectious diseases;Chagas_disease_(American_trypanosomiasis) | 2.83E-05 | 1.31E-05 | 9.67E-06 | 5.34E-06 | 0.0156 49 | 0.0127 48 | 4.83E-06 | 3.25E-05 |
| Human Diseases;Neurodegenerative diseases;Prion_diseases | 2.71E-05 | 3.06E-06 | 1.99E-05 | 2.06E-06 | 0.0010 95 | 0.0030 19 | 3.76E-06 | 1.06E-05 |

| Pathway | | | | | | | |
|---|---|---|---|---|---|---|---|
| Metabolism;Metabolism of terpenoids and polyketides;Type_I_polyketide_structures | 9.57E-06 | 5.92E-06 | 1.81E-06 | 4.14E-07 | 0.02367 | 0.016912 | 1.54E-06 | 1.4E-05 |
| Metabolism;Lipid metabolism;Steroid_biosynthesis | 4.22E-06 | 2.71E-06 | 1.11E-06 | 4.34E-07 | 0.037436 | 0.020034 | 2.62E-07 | 5.95E-06 |
| Genetic Information Processing;Replication and repair;Fanconi_anemia_pathway | 1.83E-06 | 3.84E-07 | 1.22E-06 | 2.53E-07 | 0.0103 24 | 0.0099 1 | 1.85E-07 | 1.04E-06 |
| Human Diseases;Cancers;Bladder_cancer | 6.04E-06 | 3.41E-06 | 8.4E-07 | 2.57E-07 | 0.013351 | 0.011405 | 1.63E-06 | 8.78E-06 |
| Metabolism;Biosynthesis of other secondary metabolites;Stilbenoid,_diarylheptanoid_and_gingerol_biosynthesis | 3.46E-06 | 5.49E-07 | 2.61E-06 | 4.99E-07 | 0.019051 | 0.015042 | 1.71E-07 | 1.52E-06 |
| Metabolism;Biosynthesis of other secondary metabolites;Flavonoid_biosynthesis | 3.46E-06 | 5.49E-07 | 2.61E-06 | 4.99E-07 | 0.019051 | 0.015042 | 1.71E-07 | 1.52E-06 |
| Organismal Systems;Digestive system;Bile_secretion | 2.19E-06 | 1.48E-06 | 5.04E-07 | 1.75E-07 | 0.038054 | 0.020034 | 1.36E-07 | 3.24E-06 |
| Organismal Systems;Endocrine system;Oxytocin_signaling_pathway | 1.42E-06 | 9.88E-07 | 2.69E-07 | 1.46E-07 | 0.035613 | 0.020034 | 1.13E-07 | 2.18E-06 |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_12-,_14-_and_16-membered_macrolides | 3.33E-06 | 2.17E-06 | 5.4E-07 | 1.34E-07 | 0.025208 | 0.016956 | 5.16E-07 | 5.06E-06 |
| Genetic Information Processing;Translation;mRNA_surveillance_pathway | 6.58E-07 | 4.11E-07 | 2.08E-07 | 5.96E-08 | 0.043396 | 0.020702 | 1.94E-08 | 8.8E-07 |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_enediyne_antibiotics | 2.58E-06 | 1.55E-06 | 4.91E-07 | 2.05E-07 | 0.021067 | 0.01583 | 4.64E-07 | 3.71E-06 |
| Environmental Information Processing;Signal transduction;cAMP_signaling_pathway | 3.15E-06 | 2.21E-06 | 3.89E-07 | 1.16E-07 | 0.027999 | 0.018121 | 4.44E-07 | 5.08E-06 |
| Metabolism;Xenobiotics biodegradation and metabolism;Bisphenol_degradation | 2.34E-06 | 1.44E-06 | 5.92E-07 | 3.24E-07 | 0.030626 | 0.018733 | 2.34E-07 | 3.25E-06 |
| Metabolism;Biosynthesis of other secondary metabolites;Isoflavonoid_biosynthesis | 4.08E-07 | 1.61E-07 | 1.97E-06 | 1.41E-06 | 0.041675 | 0.020702 | -3E-06 | -8.6E-08 |
| Organismal Systems;Nervous system;Retrograde_endocannabinoid_signaling | 1.11E-06 | 7.26E-07 | 2.69E-07 | 1.64E-07 | 0.0350 | 0.0200 | 8.47E-08 | 1.6E-06 |

| | | | | | 54 | 34 | | |
|---|---|---|---|---|---|---|---|---|
| Organismal Systems;Endocrine system;Regulation_of_lipolysis_in_adipocyte | 1.11E-06 | 7.26E-07 | 2.69E-07 | 1.64E-07 | 0.035054 | 0.020034 | 8.47E-08 | 1.6E-06 |
| Human Diseases;Cardiovascular diseases;Hypertrophic_cardiomyopathy_(HCM) | 2.38E-07 | 9.33E-08 | 3.92E-09 | 8.5E-09 | 0.001587 | 0.0033 06 | 1.36E-07 | 3.32E-07 |
| Organismal Systems;Endocrine system;Renin_secretion | 2.38E-07 | 9.33E-08 | 3.92E-09 | 8.5E-09 | 0.001587 | 0.0033 06 | 1.36E-07 | 3.32E-07 |
| Environmental Information Processing;Signal transduction;Sphingolipid_signaling_pathway | 7.27E-07 | 4.34E-07 | 1.8E-07 | 8.06E-08 | 0.026428 | 0.017638 | 9.32E-08 | 1E-06 |
| Human Diseases;Infectious diseases;Vibrio_cholerae_infection | 7.42E-07 | 4.06E-07 | 2.59E-07 | 4.56E-08 | 0.032847 | 0.019352 | 5.79E-08 | 9.08E-07 |
| Organismal Systems;Immune system;Fc_gamma_R-mediated_phagocytosis | 7.14E-07 | 4.29E-07 | 1.63E-07 | 8.71E-08 | 0.02463 | 0.016912 | 1.02E-07 | 1E-06 |
| Cellular Processes;Transport and catabolism;Endocytosis | 7.14E-07 | 4.29E-07 | 1.63E-07 | 8.71E-08 | 0.02463 | 0.016912 | 1.02E-07 | 1E-06 |
| Organismal Systems;Endocrine system;GnRH_signaling_pathway | 7.14E-07 | 4.29E-07 | 1.63E-07 | 8.71E-08 | 0.02463 | 0.016912 | 1.02E-07 | 1E-06 |
| Environmental Information Processing;Signal transduction;Ras_signaling_pathway | 7.14E-07 | 4.29E-07 | 1.63E-07 | 8.71E-08 | 0.02463 | 0.016912 | 1.02E-07 | 1E-06 |
| Human Diseases;Infectious diseases;Pathogenic_Escherichia_coli_infection | 6E-07 | 3.16E-07 | 2.38E-07 | 1.56E-07 | 0.038692 | 0.020034 | 2.47E-08 | 7E-07 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosylphosphatidylinositol(GPI)-anchor_biosynthesis | 8.38E-07 | 7.9E-07 | 8.43E-09 | 6.81E-09 | 0.049832 | 0.021831 | 8.95E-10 | 1.66E-06 |
| Organismal Systems;Endocrine system;Ovarian_Steroidogenesis | 3.18E-07 | 2.43E-07 | 2.55E-08 | 1.25E-08 | 0.031973 | 0.0191 8 | 3.73E-08 | 5.47E-07 |
| Organismal Systems;Digestive system;Pancreatic_secretion | 2.51E-07 | 1.78E-07 | 4.81E-08 | 3.38E-08 | 0.037784 | 0.020034 | 1.65E-08 | 3.89E-07 |
| Environmental Information Processing;Signal transduction;cGMP-PKG_signaling_pathway | 2.45E-07 | 1.74E-07 | 4.81E-08 | 3.38E-08 | 0.038398 | 0.020034 | 1.52E-08 | 3.79E-07 |
| Organismal Systems;Circulatory | 2.45E- | 1.74E- | 4.81E- | 3.38E | 0.0 | 0.0 | 1.52 | 3.79 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| system;Adrenergic_signaling_in_cardiomyocytes | 07 | 07 | 08 | -08 | 0.0 383 98 | 0.0 200 34 | E-08 | E-07 |
| Organismal Systems;Endocrine system;Insulin_secretion | 2.45E-07 | 1.74E-07 | 4.81E-08 | 3.38E-08 | 0.0 383 98 | 0.0 200 34 | 1.52 E-08 | 3.79 E-07 |
| Organismal Systems;Digestive system;Salivary_secretion | 2.45E-07 | 1.74E-07 | 4.81E-08 | 3.38E-08 | 0.0 383 98 | 0.0 200 34 | 1.52 E-08 | 3.79 E-07 |
| Organismal Systems;Excretory system;Endocrine_and_other_factor-regulated_calcium_reabsorption | 2.45E-07 | 1.74E-07 | 4.81E-08 | 3.38E-08 | 0.0 383 98 | 0.0 200 34 | 1.52 E-08 | 3.79 E-07 |
| Organismal Systems;Digestive system;Gastric_acid_secretion | 2.45E-07 | 1.74E-07 | 4.81E-08 | 3.38E-08 | 0.0 383 98 | 0.0 200 34 | 1.52 E-08 | 3.79 E-07 |
| Organismal Systems;Excretory system;Aldosterone-regulated_sodium_reabsorption | 2.45E-07 | 1.74E-07 | 4.81E-08 | 3.38E-08 | 0.0 383 98 | 0.0 200 34 | 1.52 E-08 | 3.79 E-07 |
| Organismal Systems;Nervous system;Cholinergic_synapse | 4.28E-08 | 1.78E-08 | 2.62E-07 | 1.94E-07 | 0.0 389 28 | 0.0 200 34 | -4.2E -07 | -1.6E -08 |
| Environmental Information Processing;Signal transduction;Calcium_signaling_pathway | 7.35E-08 | 4.47E-08 | 1.74E-09 | 1.24E-09 | 0.0 110 02 | 0.0 102 36 | 2.49 E-08 | 1.19 E-07 |
| Cellular Processes;Transport and catabolism;Phagosome | 2.94E-08 | 2.02E-08 | 6.79E-09 | 5.38E-09 | 0.0 400 09 | 0.0 204 67 | 1.45 E-09 | 4.37 E-08 |
| Environmental Information Processing;Signal transduction;TNF_signaling_pathway | 2.23E-07 | 2.06E-07 | 1.38E-09 | 9.89E-10 | 0.0 464 51 | 0.0 207 9 | 5.12 E-09 | 4.37 E-07 |
| Human Diseases;Infectious diseases;Leishmaniasis | 2.23E-07 | 2.06E-07 | 1.38E-09 | 9.89E-10 | 0.0 464 51 | 0.0 207 9 | 5.12 E-09 | 4.37 E-07 |
| Environmental Information Processing;Signal transduction;VEGF_signaling_pathway | 2.23E-07 | 2.06E-07 | 1.38E-09 | 9.89E-10 | 0.0 464 51 | 0.0 207 9 | 5.12 E-09 | 4.37 E-07 |
| Environmental Information Processing;Signal transduction;NF-kappa_B_signaling_pathway | 2.23E-07 | 2.06E-07 | 1.38E-09 | 9.89E-10 | 0.0 464 51 | 0.0 207 9 | 5.12 E-09 | 4.37 E-07 |
| Environmental Information Processing;Signaling molecules and interaction;G_protein-coupled_receptors | 2.34E-08 | 1.07E-08 | 3.9E-09 | 5.34E-09 | 0.0 047 5 | 0.0 066 52 | 8.08 E-09 | 3.1E- 08 |
| Metabolism;Metabolism of terpenoids and polyketides;Monoterpenoid_biosynthesis | 1.94E-08 | 1.38E-08 | 1.86E-10 | 1.57E-10 | 0.0 191 93 | 0.0 150 42 | 4.7E- 09 | 3.37 E-08 |

| Organismal Systems;Excretory system;Vasopressin-regulated_water_reabsorption | 4.89E-08 | 3.88E-08 | 5.81E-09 | 5.23E-09 | 0.0417 | 0.020702 | 2.36E-09 | 8.37E-08 |
|---|---|---|---|---|---|---|---|---|
| Organismal Systems;Nervous system;Synaptic_vesicle_cycle | 4.89E-08 | 3.88E-08 | 5.81E-09 | 5.23E-09 | 0.0417 | 0.020702 | 2.36E-09 | 8.37E-08 |
| Organismal Systems;Immune system;Hematopoietic_cell_lineage | 3.63E-08 | 2.98E-08 | 4.63E-10 | 3.75E-10 | 0.032105 | 0.01918 | 4.55E-09 | 6.71E-08 |
| Environmental Information Processing;Signaling molecules and interaction;Neuroactive_ligand-receptor_interaction | 5.88E-09 | 4.51E-09 | 5.53E-11 | 5.24E-11 | 0.024944 | 0.016912 | 1.09E-09 | 1.06E-08 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_biosynthesis-chondroitin_sulfate/dermatan_sulfate | 2.39E-09 | 1.68E-09 | 1.75E-10 | 3.9E-10 | 0.022401 | 0.016641 | 4.52E-10 | 3.97E-09 |
| Metabolism;Biosynthesis of other secondary metabolites;Clavulanic_acid_biosynthesis | 7.9E-10 | 6.83E-10 | 3.61E-11 | 5.72E-11 | 0.042469 | 0.020702 | 3.76E-11 | 1.47E-09 |

| Table 2c Young+PBS VS Young+Old | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Taxa | avg(Young+PBS) | sd(Young+PBS) | avg(Young+Old) | sd(Young+Old) | p.value | q.values | interval lower | interval upper |
| Human Diseases;Endocrine and metabolic diseases;Insulin_resistance | 0.001786 | 0.000103 | 0.001621 | 0.000144 | 0.048676 | 1 | 1.19E-06 | 0.000329 |
| Organismal Systems;Nervous system;GABAergic_synapse | 0.001346 | 3.61E-05 | 0.001278 | 5.28E-05 | 0.030139 | 1 | 8.12E-06 | 0.000127 |
| Organismal Systems;Nervous system;Glutamatergic_synapse | 0.001287 | 3.41E-05 | 0.001227 | 4.1E-05 | 0.011934 | 1 | 1.85E-05 | 0.000116 |
| Human Diseases;Immune | 0.00037 | 1.16E-0 | 0.00035 | 1.56E- | 0.04 | 1 | 1.3E-0 | 3.59E- |

| diseases;Primary_immunodeficiency | | 5 | 2 | 05 | 867 1 | | 7 | 05 |

1234

| Table 2d Young+PBS VS Young+Aged | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Taxa | avg(Young+PBS) | sd(Young+PBS) | avg(Young+Aged) | sd(Young+Aged) | p.value | q.values | interval lower | interval upper |
| Genetic Information Processing;Replication and repair;DNA_repair_and_recombination_proteins | 0.035677 | 0.000782 | 0.036768 | 0.000855 | 0.043973 | 0.134546 | -0.00215 | ######## |
| Genetic Information Processing;Translation;Transfer_RNA_biogenesis | 0.025388 | 0.000319 | 0.026138 | 0.000489 | 0.012424 | 0.090629 | -0.00129 | -0.00021 |
| Metabolism;Nucleotide metabolism;Pyrimidine_metabolism | 0.020031 | 0.000437 | 0.020711 | 0.000134 | 0.01107 | 0.090629 | -0.00114 | -0.00022 |
| Genetic Information Processing;Translation;Ribosome | 0.015723 | 0.000269 | 0.016378 | 0.000499 | 0.023175 | 0.119455 | -0.00119 | -0.00012 |
| Genetic Information Processing;Translation;Aminoacyl-tRNA_biosynthesis | 0.015499 | 0.000262 | 0.016244 | 0.000466 | 0.009445 | 0.087226 | -0.00125 | -0.00024 |
| Genetic Information Processing;Replication and repair;DNA_replication_proteins | 0.014807 | 0.000271 | 0.015335 | 0.00017 | 0.0033 49 | 0.069354 | -0.00083 | -0.00023 |
| Genetic Information Processing;Replication and repair;Chromosome_and_associated_proteins | 0.0131 | 0.000238 | 0.013777 | 0.000364 | 0.004466 | 0.070819 | -0.00108 | -0.00027 |
| Metabolism;Carbohydrate metabolism;Amino_sugar_and_nucleotide_sugar_metabolism | 0.012636 | 0.000357 | 0.01204 | 0.000343 | 0.014709 | 0.094612 | 0.000145 | 0.001046 |
| Metabolism;Amino acid metabolism;Alanine,_aspartate_and_glutamate_metabolism | 0.011744 | 0.000114 | 0.011257 | 0.000426 | 0.037153 | 0.134546 | 4.10E-05 | 0.000934 |
| Genetic Information Processing;Replication and repair;Mismatch_repair | 0.010868 | 0.000292 | 0.011363 | 0.00034 | 0.022536 | 0.119455 | -0.00009 | ######## |
| Genetic Information Processing;Replication and repair;Homologous_recombination | 0.010494 | 0.000212 | 0.010857 | 0.00027 | 0.027869 | 0.123562 | -0.00068 | ######## |
| Metabolism;Glycan biosynthesis and metabolism;Peptidoglycan_biosynthesis_and_degradation_proteins | 0.010172 | 0.000138 | 0.010445 | 0.000238 | 0.041075 | 0.134546 | -0.00053 | ######## |

| Pathway | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Metabolism;Amino acid metabolism;Cysteine_and_methionine_metabolism | 0.009602 | 7.40E-05 | 0.009104 | 0.000299 | 0.0008535 | 0.087226 | 0.000185 | 0.000811 |
| Genetic Information Processing;Replication and repair;Nucleotide_excision_repair | 0.008149 | 0.000239 | 0.008565 | 0.00011 | 0.006039 | 0.07585 | -0.00067 | -0.00016 |
| Genetic Information Processing;Replication and repair;DNA_replication | 0.007972 | 0.000183 | 0.008364 | 0.000195 | 0.004922 | 0.07272 | -0.00064 | -0.00015 |
| Metabolism;Carbohydrate metabolism;Glyoxylate_and_dicarboxylate_metabolism | 0.007261 | 0.000272 | 0.006725 | 0.000464 | 0.040322 | 0.134546 | 3.01E-05 | 0.001041 |
| Metabolism;Amino acid metabolism;Arginine_biosynthesis | 0.005337 | 0.000206 | 0.004779 | 0.000157 | 0.000449 | 0.069354 | 0.00032 | 0.000796 |
| Metabolism;Metabolism of terpenoids and polyketides;Terpenoid_backbone_biosynthesis | 0.004268 | 0.000118 | 0.004527 | 0.000127 | 0.004455 | 0.070819 | -0.00042 | -0.0001 |
| Metabolism;Energy metabolism;Nitrogen_metabolism | 0.004843 | 0.000271 | 0.004204 | 0.000351 | 0.005994 | 0.07585 | 0.000232 | 0.001047 |
| Genetic Information Processing;Replication and repair;Base_excision_repair | 0.003915 | 6.37E-05 | 0.0041 | 8.30E-05 | 0.001761 | 0.069354 | -0.00028 | ###### |
| Metabolism;Lipid metabolism;Fatty_acid_biosynthesis | 0.004229 | 6.55E-05 | 0.004002 | 0.000185 | 0.028734 | 0.124876 | 3.25E-05 | 0.000422 |
| Metabolism;Metabolism of cofactors and vitamins;Pantothenate_and_CoA_biosynthesis | 0.004248 | 0.000113 | 0.00396 | 0.000119 | 0.001548 | 0.069354 | 0.000139 | 0.000437 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosyltransferases | 0.003709 | 0.000154 | 0.003935 | 0.000121 | 0.018867 | 0.10722 | -0.00041 | ###### |
| Metabolism;Amino acid metabolism;Valine,_leucine_and_isoleucine_biosynthesis | 0.003878 | 0.000305 | 0.003116 | 0.000348 | 0.002468 | 0.069354 | 0.000034 | 0.001184 |
| Genetic Information Processing;Transcription;RNA_polymerase | 0.003305 | 7.00E-05 | 0.003506 | 0.000183 | 0.042956 | 0.134546 | -0.00039 | ###### |
| Metabolism;Carbohydrate metabolism;Pentose_and_glucuronate_interconversions | 0.003491 | 7.67E-05 | 0.003023 | 0.000435 | 0.045648 | 0.134546 | 1.30E-05 | 0.000923 |
| Metabolism;Glycan biosynthesis and metabolism;Other_glycan_degradation | 0.002999 | 0.000399 | 0.00215 | 0.000652 | 0.025253 | 0.121218 | 0.000134 | 0.001565 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | | 13 | 16 | | |
| Metabolism;Lipid metabolism;Sphingolipid_metabolism | 0.003031 | 0.000248 | 0.002382 | 0.000423 | 0.011633 | 0.090629 | 0.0000189 | 0.001111 |
| Cellular Processes;Cell motility;Cytoskeleton_proteins | 0.002762 | 6.51E-05 | 0.002963 | 0.00012 | 0.00768 | 0.086047 | -0.000033 | ###### |
| Cellular Processes;Cellular community - prokaryotes;Biofilm_formation-Escherichia_coli | 0.002944 | 0.00017 | 0.002471 | 0.00033 | 0.015333 | 0.094612 | 0.00012 | 0.000827 |
| Metabolism;Metabolism of other amino acids;Glutathione_metabolism | 0.002308 | 0.000124 | 0.002636 | 0.000165 | 0.0034342 | 0.069354 | -0.00052 | -0.00014 |
| Metabolism;Energy metabolism;Sulfur_metabolism | 0.0026 | 9.62E-05 | 0.002303 | 0.000253 | 0.033449 | 0.134546 | 3.18E-05 | 0.000563 |
| Metabolism;Amino acid metabolism;Histidine_metabolism | 0.002726 | 0.000184 | 0.002175 | 0.000457 | 0.030885 | 0.131643 | 6.87E-05 | 0.001033 |
| Metabolism;Metabolism of other amino acids;Cyanoamino_acid_metabolism | 0.002675 | 7.72E-05 | 0.002398 | 0.000263 | 0.048672 | 0.134546 | 2.25E-06 | 0.000552 |
| Cellular Processes;Cell growth and death;Necroptosis | 0.002668 | 0.000129 | 0.00226 | 0.000271 | 0.012258 | 0.090629 | 0.0000119 | 0.000696 |
| Metabolism;Biosynthesis of other secondary metabolites;Monobactam_biosynthesis | 0.002121 | 3.47E-05 | 0.002002 | 8.44E-05 | 0.016158 | 0.094612 | 3.02E-05 | 0.000208 |
| Metabolism;Carbohydrate metabolism;C5-Branched_dibasic_acid_metabolism | 0.002032 | 0.000102 | 0.001725 | 0.000275 | 0.040128 | 0.134546 | 1.89E-05 | 0.000597 |
| Human Diseases;Endocrine and metabolic diseases;Insulin_resistance | 0.001786 | 0.000103 | 0.001511 | 0.000136 | 0.0030391 | 0.069354 | 0.0000119 | 0.000432 |
| Metabolism;Metabolism of terpenoids and polyketides;Prenyltransferases | 0.001408 | 5.91E-05 | 0.001536 | 0.000108 | 0.035041 | 0.134546 | -0.00024 | ###### |
| Metabolism;Metabolism of other amino acids;D-Glutamine_and_D-glutamate_metabolism | 0.001416 | 4.80E-05 | 0.001513 | 1.80E-05 | 0.0031363 | 0.069354 | -0.00015 | ###### |
| Metabolism;Glycan biosynthesis and metabolism;Glycosphingolipid_biosynthesis-globo_and_isoglobo_series | 0.001423 | 0.000143 | 0.001089 | 0.000203 | 0.0093881 | 0.087226 | 0.000104 | 0.000562 |
| Metabolism;Biosynthesis of other secondary | 0.0015 | 6.81E | 0.0011 | 0.000 | 0.0 | 0.0 | 0.00 | 0.00 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| metabolites;Phenylpropanoid_biosynthesis | 72 | -05 | 68 | 266 | 126 65 | 906 29 | 0125 | 0683 |
| Metabolism;Metabolism of other amino acids;D-Alanine_metabolism | 0.0011 68 | 4.12E -05 | 0.0012 31 | 5.04E- 05 | 0.0 391 06 | 0.1 345 46 | -0.00 012 | #### ## |
| Metabolism;Enzyme families;Protein_phosphatase_and_associated_prot eins | 0.0013 01 | 7.07E -05 | 0.0011 32 | 0.000 133 | 0.0 261 47 | 0.1 218 16 | 2.62 E-05 | 0.00 0312 |
| Unclassified;Metabolism;Carbohydrate_metabolism | 0.0013 15 | 0.000 147 | 0.0010 74 | 0.000 171 | 0.0 263 82 | 0.1 218 16 | 3.48 E-05 | 0.00 0446 |
| Unclassified;Metabolism;Amino_acid_metabolism | 0.0011 95 | 5.78E -05 | 0.0009 91 | 0.000 177 | 0.0 360 34 | 0.1 345 46 | 1.84 E-05 | 0.00 039 |
| Organismal Systems;Endocrine system;Insulin_signaling_pathway | 0.0012 73 | 8.54E -05 | 0.0010 24 | 0.000 108 | 0.0 014 65 | 0.0 693 54 | 0.00 0123 | 0.00 0375 |
| Human Diseases;Cardiovascular diseases;Fluid_shear_stress_and_atherosclerosis | 0.0011 66 | 2.70E -05 | 0.0010 97 | 5.46E- 05 | 0.0 250 61 | 0.1 218 16 | 1.15 E-05 | 0.00 0128 |
| Unclassified;Viral protein family;Viral_proteins | 0.0008 7 | 4.86E -05 | 0.0009 77 | 7.30E- 05 | 0.0 161 66 | 0.0 946 12 | -0.00 019 | #### ## |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_degradation | 0.0009 25 | 0.000 151 | 0.0006 13 | 0.000 289 | 0.0 483 34 | 0.1 345 46 | 2.96 E-06 | 0.00 0623 |
| Metabolism;Biosynthesis of other secondary metabolites;Novobiocin_biosynthesis | 0.0010 16 | 5.43E -05 | 0.0008 44 | 0.0001 11 | 0.0 104 74 | 0.0 906 29 | 5.42 E-05 | 0.00 0291 |
| Metabolism;Biosynthesis of other secondary metabolites;Tropane,_piperidine_and_pyridine_alkal oid_biosynthesis | 0.0010 43 | 6.63E -05 | 0.0008 39 | 0.0001 1 | 0.0 044 73 | 0.0 708 19 | 8.29 E-05 | 0.00 0324 |
| Cellular Processes;Transport and catabolism;Lysosome | 0.0008 4 | 0.000 2 | 0.0004 81 | 0.000 212 | 0.0 130 85 | 0.0 906 29 | 9.33 E-05 | 0.00 0623 |
| Metabolism;Carbohydrate metabolism;Ascorbate_and_aldarate_metabolism | 0.0007 64 | 6.21E -05 | 0.0005 73 | 0.000 134 | 0.0 152 69 | 0.0 946 12 | 4.95 E-05 | 0.00 0334 |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_ansamycins | 0.0008 44 | 2.70E -05 | 0.0007 95 | 3.53E- 05 | 0.0 229 16 | 0.1 194 55 | 8.50 E-06 | 9.01 E-05 |
| Environmental Information Processing;Signal transduction;AMPK_signaling_pathway | 0.0006 94 | 4.69E -05 | 0.0006 12 | 4.03E- 05 | 0.0 093 21 | 0.0 872 26 | 2.51 E-05 | 0.00 0138 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Unclassified;Metabolism;Glycan_biosynthesis_and_metabolism | 0.000642 | 4.08E-05 | 0.000535 | 5.04E-05 | 0.002677 | 0.069354 | 4.70E-05 | 0.000166 |
| Metabolism;Biosynthesis of other secondary metabolites;Neomycin,_kanamycin_and_gentamicin_biosynthesis | 0.000529 | 2.51E-05 | 0.000498 | 1.48E-05 | 0.034166 | 0.134546 | 2.89E-06 | 5.76E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Glucosinolate_biosynthesis | 0.000341 | 3.28E-05 | 0.000253 | 3.57E-05 | 0.012123 | 0.069354 | 4.43E-05 | 0.000133 |
| Human Diseases;Cancers;Renal_cell_carcinoma | 0.000173 | 2.10E-05 | 0.000265 | 7.42E-05 | 0.027875 | 0.123562 | -0.00017 | ###### |
| Metabolism;Lipid metabolism;Steroid_hormone_biosynthesis | 0.000174 | 6.14E-05 | 6.77E-05 | 3.33E-05 | 0.00616 | 0.07585 | 4.02E-05 | 0.000173 |
| Organismal Systems;Digestive system;Carbohydrate_digestion_and_absorption | 0.000114 | 3.23E-05 | 6.80E-05 | 2.74E-05 | 0.024285 | 0.121816 | 7.38E-06 | 8.47E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Flavone_and_flavonol_biosynthesis | 0.000118 | 8.70E-06 | 9.90E-05 | 1.79E-05 | 0.047375 | 0.134546 | 2.94E-07 | 3.85E-05 |
| Metabolism;Enzyme families;Cytochrome_P450 | 7.32E-05 | 2.56E-05 | 4.52E-05 | 1.33E-05 | 0.047224 | 0.134546 | 4.43E-07 | 5.55E-05 |
| Metabolism;Xenobiotics biodegradation and metabolism;Steroid_degradation | 4.60E-05 | 2.10E-05 | 2.35E-05 | 6.47E-06 | 0.04649 | 0.134546 | 4.85E-07 | 4.46E-05 |
| Human Diseases;Infectious diseases;African_trypanosomiasis | 3.10E-05 | 1.36E-05 | 1.14E-05 | 8.24E-06 | 0.016221 | 0.094612 | 4.65E-06 | 3.44E-05 |
| Human Diseases;Infectious diseases;Chagas_disease_(American_trypanosomiasis) | 2.83E-05 | 1.31E-05 | 8.78E-06 | 7.11E-06 | 0.013033 | 0.090629 | 5.41E-06 | 3.37E-05 |
| Human Diseases;Neurodegenerative diseases;Prion_diseases | 2.71E-05 | 3.06E-06 | 2.13E-05 | 4.88E-06 | 0.037345 | 0.134546 | 4.31E-07 | 1.12E-05 |
| Genetic Information Processing;Translation;mRNA_surveillance_pathway | 6.58E-07 | 4.11E-07 | 1.73E-07 | 1.07E-07 | 0.033088 | 0.134546 | 5.52E-08 | 9.15E-07 |
| Human Diseases;Immune diseases;Systemic_lupus_erythematosus | 5.61E-07 | 2.55E-07 | 2.84E-07 | 1.41E-07 | 0.049627 | 0.134546 | 5.77E-10 | 5.52E-07 |
| Human Diseases;Infectious diseases;Vibrio_cholerae_infection | 7.42E-07 | 4.06E-07 | 1.88E-07 | 8.80E-08 | 0.0194 | 0.1079 | 1.29E-07 | 9.79E-07 |

| | avg(Old+PBS) | sd(Old+PBS) | avg(Old+Young) | sd(Old+Young) | p.value | q.values | interval lower | interval upper |
|---|---|---|---|---|---|---|---|---|
| | | | | | 79 | 33 | | |
| Organismal Systems;Endocrine system;Ovarian_Steroidogenesis | 3.18E-07 | 2.43E-07 | 5.52E-08 | 4.16E-08 | 0.045175 | 0.134546 | 8.14E-09 | 5.17E-07 |
| Environmental Information Processing;Signaling molecules and interaction;G_protein-coupled_receptors | 2.34E-08 | 1.07E-08 | 4.33E-09 | 9.00E-09 | 0.007765 | 0.086047 | 6.32E-09 | 3.19E-08 |
| Organismal Systems;Excretory system;Vasopressin-regulated_water_reabsorption | 4.89E-08 | 3.88E-08 | 6.63E-09 | 6.05E-09 | 0.044322 | 0.134546 | 1.55E-09 | 8.29E-08 |
| Organismal Systems;Nervous system;Synaptic_vesicle_cycle | 4.89E-08 | 3.88E-08 | 6.63E-09 | 6.05E-09 | 0.044322 | 0.134546 | 1.55E-09 | 8.29E-08 |
| Organismal Systems;Immune system;Hematopoietic_cell_lineage | 3.63E-08 | 2.98E-08 | 3.89E-09 | 6.43E-09 | 0.044217 | 0.134546 | 1.20E-09 | 6.36E-08 |
| Environmental Information Processing;Signaling molecules and interaction;Neuroactive_ligand-receptor_interaction | 5.88E-09 | 4.51E-09 | 8.81E-10 | 1.21E-09 | 0.041204 | 0.134546 | 2.81E-10 | 9.72E-09 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_biosynthesis-chondroitin_sulfate/dermatan_sulfate | 2.39E-09 | 1.68E-09 | 5.30E-10 | 5.60E-10 | 0.041994 | 0.134546 | 9.26E-11 | 3.62E-09 |
| Metabolism;Biosynthesis of other secondary metabolites;Clavulanic_acid_biosynthesis | 7.90E-10 | 6.83E-10 | 1.51E-11 | 1.23E-11 | 0.038954 | 0.134546 | 5.80E-11 | 1.49E-09 |

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

| Table 2e Old+PBS VS Old+Young | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Taxa | avg(Old+PBS) | sd(Old+PBS) | avg(Old+Young) | sd(Old+Young) | p.value | q.values | interval lower | interval upper |

| Pathway | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Genetic Information Processing;Replication and repair;DNA_replication_proteins | 0.015186 | 0.000287 | 0.0148 66 | 9.1E-05 | 0.0040875 | 0.279366 | 1.84E-05 | 0.00062 |
| Metabolism;Carbohydrate metabolism;Pyruvate_metabolism | 0.012992 | 0.000385 | 0.013656 | 0.000423 | 0.001754 | 0.254192 | -0.00119 | -0.00014 |
| Genetic Information Processing;Replication and repair;Nucleotide_excision_repair | 0.008575 | 0.000252 | 0.008078 | 0.000267 | 0.0078 15 | 0.243757 | 0.000163 | 0.000832 |
| Metabolism;Amino acid metabolism;Glycine,_serine_and_threonine_metabolism | 0.007344 | 0.000264 | 0.00785 | 0.000189 | 0.004085 | 0.217068 | -0.00081 | -0.00021 |
| Metabolism;Carbohydrate metabolism;Butanoate_metabolism | 0.006856 | 0.000234 | 0.00723 | 0.000214 | 0.0163 88 | 0.248807 | -0.00066 | -8.5E-05 |
| Metabolism;Amino acid metabolism;Lysine_biosynthesis | 0.006577 | 0.000222 | 0.006169 | 0.000258 | 0.015102 | 0.248807 | 9.81E-05 | 0.00072 |
| Cellular Processes;Cell motility;Bacterial_chemotaxis | 0.006541 | 0.000982 | 0.005025 | 0.001103 | 0.030986 | 0.279366 | 0.00017 | 0.002862 |
| Metabolism;Carbohydrate metabolism;Fructose_and_mannose_metabolism | 0.005034 | 0.000448 | 0.005551 | 0.000339 | 0.049861 | 0.29823 | -0.00103 | -3.9E-07 |
| Cellular Processes;Cell growth and death;Cell_cycle-Caulobacter | 0.005301 | 0.000205 | 0.005009 | 0.000152 | 0.020387 | 0.26536 | 5.65E-05 | 0.000527 |
| Metabolism;Lipid metabolism;Lipid_biosynthesis_proteins | 0.004195 | 0.000254 | 0.004486 | 8.15E-05 | 0.036704 | 0.279366 | -0.00056 | -2.5E-05 |
| Metabolism;Metabolism of terpenoids and polyketides;Terpenoid_backbone_biosynthesis | 0.004511 | 0.000132 | 0.004343 | 3.68E-05 | 0.024962 | 0.274426 | 2.99E-05 | 0.000306 |
| Metabolism;Lipid metabolism;Glycerophospholipid_metabolism | 0.00449 | 0.00018 | 0.004242 | 0.000119 | 0.020808 | 0.26536 | 4.8E-05 | 0.000449 |
| Human Diseases;Drug resistance;beta-Lactam_resistance | 0.00351 | 0.000255 | 0.004224 | 0.000625 | 0.037756 | 0.279366 | -0.00137 | -5.4E-05 |
| Metabolism;Amino acid metabolism;Valine,_leucine_and_isoleucine_biosynthesis | 0.00317 | 0.000297 | 0.00354 | 8.59E-05 | 0.026938 | 0.279366 | -0.00068 | -6E-05 |
| Metabolism;Energy metabolism;Photosynthesis_proteins | 0.00297 | 8.72E-05 | 0.002711 | 0.000103 | 0.0009 | 0.1534 | 0.000136 | 0.000382 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | | 01 | 05 | | |
| Metabolism;Energy metabolism;Photosynthesis | 0.002958 | 8.61E-05 | 0.002693 | 0.000108 | 0.000962 | 0.153405 | 0.000138 | 0.000391 |
| Metabolism;Amino acid metabolism;Valine,_leucine_and_isoleucine_degradation | 0.002337 | 0.00011 | 0.002675 | 0.000237 | 0.015654 | 0.248807 | -0.00059 | -8.6E-05 |
| Cellular Processes;Cell motility;Cytoskeleton_proteins | 0.002926 | 6.67E-05 | 0.002601 | 0.000202 | 0.009175 | 0.243757 | 0.000114 | 0.000537 |
| Metabolism;Amino acid metabolism;Histidine_metabolism | 0.002115 | 0.0000282 | 0.002464 | 5.24E-05 | 0.028246 | 0.279366 | -0.00064 | -5.4E-05 |
| Metabolism;Amino acid metabolism;Phenylalanine_metabolism | 0.00157 | 9.34E-05 | 0.001676 | 4.86E-05 | 0.04133 | 0.279366 | -0.00021 | -5.4E-06 |
| Metabolism;Metabolism of other amino acids;D-Glutamine_and_D-glutamate_metabolism | 0.001524 | 5.98E-05 | 0.001445 | 3.28E-05 | 0.022738 | 0.274426 | 1.43E-05 | 0.000144 |
| Metabolism;Metabolism of cofactors and vitamins;Vitamin_B6_metabolism | 0.001084 | 0.0000119 | 0.001216 | 4.67E-05 | 0.041152 | 0.279366 | -0.00026 | -7.2E-06 |
| Unclassified;Metabolism;Carbohydrate_metabolism | 0.001019 | 0.0000118 | 0.001209 | 6.85E-05 | 0.008989 | 0.243757 | -0.00032 | -6.2E-05 |
| Metabolism;Metabolism of other amino acids;beta-Alanine_metabolism | 0.000946 | 6.77E-05 | 0.001021 | 3.03E-05 | 0.042008 | 0.279366 | -0.00015 | -3.6E-06 |
| Environmental Information Processing;Signaling molecules and interaction;Bacterial_toxins | 0.000835 | 0.0000179 | 0.001174 | 0.000293 | 0.041303 | 0.279366 | -0.00066 | -1.7E-05 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_degradation | 0.000591 | 0.0000218 | 0.000871 | 4.73E-05 | 0.02479 | 0.274426 | -0.00051 | -5.1E-05 |
| Metabolism;Biosynthesis of other secondary metabolites;Tropane,_piperidine_and_pyridine_alkaloid_biosynthesis | 0.000805 | 7.65E-05 | 0.0009 | 6.07E-05 | 0.039433 | 0.279366 | -0.00018 | -5.7E-06 |
| Cellular Processes;Transport and catabolism;Lysosome | 0.000516 | 0.0000241 | 0.000789 | 5.37E-05 | 0.03826 | 0.279366 | -0.00053 | -2.1E-05 |
| Metabolism;Carbohydrate metabolism;Ascorbate_and_aldarate_metabolism | 0.000622 | 8.97E-05 | 0.000961 | 0.000238 | 0.015635 | 0.248807 | -0.00059 | -8.9E-05 |
| Metabolism;Metabolism of terpenoids and | 0.000 | 5.17 | 0.0008 | 5.14E | 0.0 | 0.2 | -0.00 | -8.3E |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| polyketides;Biosynthesis_of_ansamycins | 79 | E-05 | 64 | -05 | 310 53 | 793 66 | 014 | -06 |
| Unclassified;Metabolism;Glycan_biosynthesis_and_metabolism | 0.000 512 | 6.35 E-05 | 0.0006 4 | 4.16E -05 | 0.0 028 66 | 0.2 170 68 | -0.00 02 | -5.7E -05 |
| Metabolism;Lipid metabolism;Secondary_bile_acid_biosynthesis | 0.000 417 | 6.43 E-05 | 0.0003 17 | 3.16E -05 | 0.0 108 32 | 0.2 488 07 | 3.07 E-05 | 0.000 168 |
| Metabolism;Biosynthesis of other secondary metabolites;Glucosinolate_biosynthesis | 0.000 265 | 2.64 E-05 | 0.0003 01 | 1.19E -05 | 0.0 190 76 | 0.2 644 35 | -6.4E -05 | -7.9E -06 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosylphosphatidylinositol_(GPI)-anchored_proteins | 0.000 188 | 5.55 E-05 | 0.0002 72 | 2.39E -05 | 0.0 120 06 | 0.2 488 07 | -0.00 014 | -2.5E -05 |
| Human Diseases;Cancers;Proteoglycans_in_cancer | 0.000 297 | 3.5E- 06 | 0.0002 86 | 9.84E -06 | 0.0 493 27 | 0.2 982 3 | 4.16 E-08 | 2.07 E-05 |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_siderophore_group_nonribosomal_peptides | 0.000 164 | 3.97 E-05 | 0.0002 68 | 5.31E -05 | 0.0 038 14 | 0.2 170 68 | -0.00 016 | -4.3E -05 |
| Human Diseases;Cancers;Renal_cell_carcinoma | 0.000 284 | 5.25 E-05 | 0.0002 24 | 9.02E -06 | 0.0 374 56 | 0.2 793 66 | 5.05 E-06 | 0.000 115 |
| Metabolism;Glycan biosynthesis and metabolism;Glycosaminoglycan_binding_proteins | 0.000 106 | 2.46 E-05 | 0.0001 66 | 2.76E -05 | 0.0 028 84 | 0.2 170 68 | -9.3E -05 | -2.6E -05 |
| Organismal Systems;Digestive system;Carbohydrate_digestion_and_absorption | 7.38E -05 | 1.73 E-05 | 0.0001 45 | 4.15E -05 | 0.0 067 76 | 0.2 437 57 | -0.00 011 | -2.7E -05 |
| Human Diseases;Infectious diseases;African_trypanosomiasis | 7.93E -06 | 3.38 E-06 | 1.96E- 05 | 6.83E -06 | 0.0 065 24 | 0.2 437 57 | -1.9E -05 | -4.4E -06 |
| Human Diseases;Infectious diseases;Chagas_disease_(American_trypanosomiasis) | 5.73E -06 | 3.61 E-06 | 1.58E- 05 | 7.11E -06 | 0.0 163 72 | 0.2 488 07 | -1.8E -05 | -2.4E -06 |
| Organismal Systems;Digestive system;Mineral_absorption | 8.81E -06 | 1.93 E-06 | 1.57E- 05 | 5.61E -06 | 0.0 285 09 | 0.2 793 66 | -1.3E -05 | -1E-0 6 |
| Environmental Information Processing;Signaling molecules and interaction;Cell_adhesion_molecules_and_their_ligands | 3.64E -06 | 2.05 E-06 | 8.03E- 06 | 4.03E -06 | 0.0 466 57 | 0.2 916 74 | -8.7E -06 | -8.4E -08 |
| Organismal Systems;Endocrine system;Relaxin_signaling_pathway | 3.44E -06 | 1.9E- 06 | 7.31E- 06 | 3.46E -06 | 0.0 438 | 0.2 793 | -7.6E -06 | -1.4E -07 |

| Taxa | avg | sd | avg | sd | p.value | q.values | interval lower | interval upper |
|---|---|---|---|---|---|---|---|---|
| | | | | | 12 | 66 | | |
| Environmental Information Processing;Signaling molecules and interaction;ECM-receptor_interaction | 3.44E-06 | 1.9E-06 | 7.31E-06 | 3.46E-06 | 0.043812 | 0.279366 | -7.6E-06 | -1.4E-07 |
| Cellular Processes;Cell growth and death;Focal_adhesion | 3.44E-06 | 1.9E-06 | 7.31E-06 | 3.46E-06 | 0.043812 | 0.279366 | -7.6E-06 | -1.4E-07 |
| Human Diseases;Endocrine and metabolic diseases;AGE-RAGE_signaling_pathway_in_diabetic_complications | 3.44E-06 | 1.9E-06 | 7.31E-06 | 3.46E-06 | 0.043812 | 0.279366 | -7.6E-06 | -1.4E-07 |
| Metabolism;Biosynthesis of other secondary metabolites;Stilbenoid,_diarylheptanoid_and_gingerol_biosynthesis | 1.76E-06 | 9.04E-07 | 3.18E-06 | 7.56E-07 | 0.015052 | 0.248807 | -2.5E-06 | -3.4E-07 |
| Metabolism;Biosynthesis of other secondary metabolites;Flavonoid_biosynthesis | 1.76E-06 | 9.04E-07 | 3.18E-06 | 7.56E-07 | 0.015052 | 0.248807 | -2.5E-06 | -3.4E-07 |
| Human Diseases;Immune diseases;Systemic_lupus_erythematosus | 2.46E-07 | 9.81E-08 | 8.68E-07 | 3.66E-07 | 0.007701 | 0.243757 | -1E-06 | -2.4E-07 |
| Human Diseases;Infectious diseases;Pathogenic_Escherichia_coli_infection | 1.71E-07 | 8.14E-08 | 3.99E-07 | 1.87E-07 | 0.029627 | 0.279366 | -4.3E-07 | -3E-08 |
| Organismal Systems;Endocrine system;Ovarian_Steroidogenesis | 6.29E-08 | 5.25E-08 | 1.3E-07 | 4.22E-08 | 0.036164 | 0.279366 | -1.3E-07 | -5.3E-09 |
| Environmental Information Processing;Signaling molecules and interaction;G_protein-coupled_receptors | 2.05E-09 | 3.26E-09 | 3.22E-08 | 2.31E-08 | 0.023419 | 0.274426 | -5.4E-08 | -6E-09 |

| Table 2f Aged+PBS VS Aged+Young | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Taxa | avg(Aged+PBS) | sd(Aged+PBS) | avg(Aged+Young) | sd(Aged+Young) | p.value | q.values | interval lower | interval upper |
| Genetic Information Processing;Replication and | 0.015 | 0.000 | 0.0144 | 0.0002 | 0.0 | 0.0 | 0.00 | 0.000 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| repair;DNA_replication_proteins | 109 | 133 | 68 | 77 | 01287 | 83073 | 0346 | 937 |
| Metabolism;Carbohydrate metabolism;Pyruvate_metabolism | 0.013256 | 0.000309 | 0.013857 | 0.000478 | 0.0030863 | 0.211941 | -0.00113 | ###### |
| Genetic Information Processing;Replication and repair;Mismatch_repair | 0.011191 | 0.000174 | 0.010727 | 0.000429 | 0.0045615 | 0.211941 | 1.22E-05 | 0.000917 |
| Environmental Information Processing;Membrane transport;Secretion_system | 0.010266 | 0.000857 | 0.00902 | 0.000808 | 0.0026975 | 0.211941 | 0.000174 | 0.002318 |
| Unclassified;Genetic information processing;Replication,_recombination_and_repair_proteins | 0.009811 | 0.000476 | 0.00863 | 0.001048 | 0.0403 | 0.211941 | 6.92E-05 | 0.002293 |
| Metabolism;Energy metabolism;Carbon_fixation_pathways_in_prokaryotes | 0.008631 | 0.000453 | 0.009592 | 0.000683 | 0.0019043 | 0.185973 | -0.00172 | -0.0002 |
| Genetic Information Processing;Translation;Messenger_RNA_Biogenesis | 0.007653 | 0.000128 | 0.008007 | 0.000249 | 0.0015963 | 0.185973 | -0.00062 | ###### |
| Genetic Information Processing;Folding, sorting and degradation;RNA_degradation | 0.007297 | 0.000143 | 0.007737 | 0.000271 | 0.0008643 | 0.164618 | -0.00073 | -0.00015 |
| Metabolism;Energy metabolism;Methane_metabolism | 0.007085 | 0.000104 | 0.00744 | 0.000299 | 0.0032733 | 0.211941 | -0.00067 | ###### |
| Metabolism;Carbohydrate metabolism;Butanoate_metabolism | 0.006828 | 0.000131 | 0.007411 | 0.000501 | 0.0034917 | 0.211941 | -0.00111 | ###### |
| Metabolism;Carbohydrate metabolism;Citrate_cycle_(TCA_cycle) | 0.005779 | 0.000294 | 0.006533 | 0.000662 | 0.0038681 | 0.211941 | -0.00146 | ###### |
| Unclassified;Metabolism;Energy_metabolism | 0.005571 | 0.00032 | 0.006071 | 0.000169 | 0.0010204 | 0.164618 | -0.00084 | -0.00016 |
| Metabolism;Energy metabolism;Carbon_fixation_in_photosynthetic_organisms | 0.004982 | 4.40E-05 | 0.005252 | 0.000113 | 0.0001201 | 0.083073 | -0.00039 | -0.00015 |
| Metabolism;Amino acid metabolism;Arginine_biosynthesis | 0.004855 | 0.000135 | 0.005175 | 0.000167 | 0.0004703 | 0.13006 | -0.00052 | -0.00012 |
| Metabolism;Lipid metabolism;Glycerophospholipid_metabolism | 0.004402 | 0.00011 | 0.00409 | 0.000279 | 0.004047 | 0.211941 | 1.83E-05 | 0.000607 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Metabolism;Glycan biosynthesis and metabolism;Glycosyltransferases | 0.003989 | 0.000158 | 0.003766 | 0.000147 | 0.030047 | 0.211941 | 2.63E-05 | 0.000419 |
| Metabolism;Amino acid metabolism;Valine,_leucine_and_isoleucine_biosynthesis | 0.003302 | 0.000124 | 0.003687 | 0.000289 | 0.020583 | 0.185973 | -0.00069 | #### |
| Metabolism;Energy metabolism;Sulfur_metabolism | 0.00237 | 6.26E-05 | 0.002678 | 0.000107 | 0.000281 | 0.054485 | -0.00042 | -0.00019 |
| Metabolism;Amino acid metabolism;Histidine_metabolism | 0.002265 | 6.44E-05 | 0.002736 | 0.000321 | 0.014798 | 0.185973 | -0.00081 | -0.00014 |
| Metabolism;Carbohydrate metabolism;C5-Branched_dibasic_acid_metabolism | 0.001784 | 8.80E-05 | 0.001976 | 0.000135 | 0.018159 | 0.185973 | -0.00034 | #### |
| Metabolism;Amino acid metabolism;Phenylalanine_metabolism | 0.001538 | 7.37E-05 | 0.00177 | 0.000109 | 0.01977 | 0.095695 | -0.00035 | -0.00011 |
| Unclassified;Metabolism;Amino_acid_metabolism | 0.001107 | 4.00E-05 | 0.001305 | 0.000104 | 0.004041 | 0.13006 | -0.00031 | #### |
| Unclassified;Viral protein family;Viral_proteins | 0.001018 | 4.82E-05 | 0.000951 | 5.44E-05 | 0.046274 | 0.211941 | 1.36E-06 | 0.000134 |
| Metabolism;Biosynthesis of other secondary metabolites;Novobiocin_biosynthesis | 0.000857 | 5.38E-05 | 0.000986 | 9.59E-05 | 0.021135 | 0.185973 | -0.00023 | #### |
| Metabolism;Biosynthesis of other secondary metabolites;Tropane,_piperidine_and_pyridine_alkaloid_biosynthesis | 0.000836 | 5.62E-05 | 0.000952 | 9.70E-05 | 0.034092 | 0.211941 | -0.00022 | #### |
| Human Diseases;Endocrine and metabolic diseases;Type_I_diabetes_mellitus | 0.000693 | 8.78E-06 | 0.000723 | 2.70E-05 | 0.038661 | 0.211941 | #### | #### |
| Metabolism;Biosynthesis of other secondary metabolites;Carbapenem_biosynthesis | 0.000657 | 3.20E-05 | 0.000715 | 3.61E-05 | 0.015544 | 0.185973 | -0.0001 | #### |
| Human Diseases;Immune diseases;Primary_immunodeficiency | 0.000376 | 1.44E-05 | 0.000344 | 2.31E-05 | 0.018877 | 0.185973 | 6.82E-06 | 5.76E-05 |
| Metabolism;Lipid metabolism;Secondary_bile_acid_biosynthesis | 0.000359 | 2.94E-05 | 0.000324 | 2.35E-05 | 0.045604 | 0.211941 | 8.46E-07 | 6.97E-05 |
| Metabolism;Biosynthesis of other | 0.000 | 1.21E | 0.0003 | 3.73E- | 0.0 | 0.1 | #### | #### |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| secondary metabolites;Glucosinolate_biosynthesis | 272 | -05 | 31 | 05 | 095 43 | 646 18 | ## | ## |
| Metabolism;Xenobiotics biodegradation and metabolism;Atrazine_degradation | 0.000 234 | 5.77E -05 | 0.0001 63 | 5.24E- 05 | 0.0 481 73 | 0.2 119 41 | 7.01 E-07 | 0.000 143 |
| Human Diseases;Neurodegenerative diseases;Huntingtons_disease | 0.000 223 | 5.85E -05 | 0.0001 61 | 1.05E- 05 | 0.0 479 24 | 0.2 119 41 | 8.15 E-07 | 0.000 123 |
| Metabolism;Xenobiotics biodegradation and metabolism;Xylene_degradation | 0.000 19 | 1.18E -05 | 0.0001 64 | 1.55E- 05 | 0.0 091 88 | 0.1 646 18 | 8.12 E-06 | 4.38 E-05 |
| Genetic Information Processing;Folding, sorting and degradation;Ubiquitin_system | 0.000 16 | 4.01E -06 | 0.0001 46 | 8.65E- 06 | 0.0 089 99 | 0.1 646 18 | 4.70 E-06 | 2.31 E-05 |
| Metabolism;Xenobiotics biodegradation and metabolism;Toluene_degradation | 0.0001 1 | 1.13E -05 | 8.93E- 05 | 1.90E- 05 | 0.0 462 8 | 0.2 119 41 | 4.43 E-07 | 4.19 E-05 |
| Metabolism;Xenobiotics biodegradation and metabolism;Caprolactam_degradation | 6.02E- 05 | 1.83E -05 | 9.04E- 05 | 2.37E- 05 | 0.0 343 98 | 0.2 119 41 | #### ## | #### ## |
| Metabolism;Lipid metabolism;Ether_lipid_metabolism | 6.40E- 05 | 2.06E -05 | 4.06E- 05 | 7.48E- 06 | 0.0 385 07 | 0.2 119 41 | 1.69 E-06 | 4.50 E-05 |
| Metabolism;Lipid metabolism;alpha-Linolenic_acid_metabolism | 5.48E- 05 | 1.65E -05 | 3.19E- 05 | 7.59E- 06 | 0.0 176 43 | 0.1 859 73 | 5.35 E-06 | 4.04 E-05 |
| Human Diseases;Cancers;Small_cell_lung_cancer | 4.15E- 06 | 1.57E -06 | 6.28E- 06 | 9.43E- 07 | 0.0 207 95 | 0.1 859 73 | #### ## | #### ## |
| Genetic Information Processing;Replication and repair;Fanconi_anemia_pathway | 1.22E- 06 | 2.53E -07 | 9.36E- 07 | 1.65E- 07 | 0.0 473 93 | 0.2 119 41 | 4.11 E-09 | 5.66 E-07 |
| Human Diseases;Cancers;Bladder_cancer | 8.40E- 07 | 2.57E -07 | 1.27E- 06 | 3.32E- 07 | 0.0 327 44 | 0.2 119 41 | #### ## | #### ## |
| Metabolism;Metabolism of terpenoids and polyketides;Biosynthesis_of_type_II_polyketide_bac kbone | 2.91E- 07 | 4.49E -08 | 6.86E- 07 | 1.91E- 07 | 0.0 032 27 | 0.1 249 25 | #### ## | #### ## |
| Organismal Systems;Excretory system;Vasopressin-regulated_water_reabsorption | 5.81E- 09 | 5.23E -09 | 2.11E- 08 | 1.43E- 08 | 0.0 481 09 | 0.2 119 41 | #### ## | #### ## |
| Organismal Systems;Nervous system;Synaptic_vesicle_cycle | 5.81E- 09 | 5.23E -09 | 2.11E- 08 | 1.43E- 08 | 0.0 481 09 | 0.2 119 41 | #### ## | #### ## |

1260