# Microglial MyD88-dependent pathways are regulated in a sex specific manner in the context of HMGB1-induced anxiety

Ashleigh Rawls[1]*, Dang Nyugen[2], Julia Dziabis[2], Dilara Anbarci[3], Madeline Clark[4], Kafui Dzirasa[4], Staci Bilbo[2]

## Abstract

Chronic stress is a major risk factor for development and recurrence of anxiety disorders. Chronic stress has been shown to impact the immune system, causing microglial activation in the medial prefrontal cortex (mPFC), a brain region involved in the pathogenesis of anxiety. HMGB1 is both an established modulator of neuronal firing and a potent pro-inflammatory stimulus that is released from neuronal and non-neuronal cells following stress. HMGB1 in the context of stress acts as a danger associated molecular pattern (DAMP) instigating robust proinflammatory responses throughout the brain; so much so, that localized drug delivery of HMGB1 alters behavior in the absence of any other forms of stress i.e. social isolation, or behavioral stress models. Few studies have investigated the molecular mechanisms which underlie HMGB1 associated behavioral effects in a cell-specific manner. The aim of this study is to investigate cellular and molecular mechanisms underlying HMGB1 induced behavioral dysfunction with regards to cell-type specificity and potential sex differences. Here, we report that both male and female mice exhibited anxiety-like behavior following increased HMGB1 in the mPFC as well as changes in microglial morphology. However, only female mice showed microglial activation characterized by increased phagocytic capacity and decreased Tmem119 expression. Moreover, when measuring RNA from isolated microglia and non-microglial cells from the frontal cortex we found that female HMGB1 treated mice displayed a robust increase of RAGE and MyD88. Given these findings, to further investigate the underlying molecular mechanisms associated with HMGB1 induced anxiety-like behavior and microglia activation in mice, we used cKO (conditional knockout) mice with conditional deletion of MyD88 in microglia and repeated the paradigm. For males, we saw that the genetic manipulation did not prevent behavioral deficits in response to HMGB1 treatment. However, female cKO mice were protected from HMGB1- induced behavioral deficits. This study supports the hypothesis that the MyD88 signaling in microglia may be a crucial mediator of stress response in adult female mice.

## Introduction

Anxiety is the most common cause of mental health related disability. Anxiety disorders describe a class of disorders in which the affected person experiences consistent and often unfocused anxiety and worry (Bandelow and Michaelis., 2015). Though generally high in prevalence, anxiety is twice as common in women throughout the lifetime (Mclean et al., 2011). Few biological mechanisms have been elucidated that explain the increased prevalence of mood disorders in women; however, we do understand that stress is a pivotal factor in the development and recurrence of anxiety disorders regardless of sex or age (McEwen and Akil., 2020). Stress instigates many alterations throughout the brain-body axis (Fonken et al., 2018). Specifically, changes in the peripheral and central immune system have been observed in patients with anxiety and comorbid depression. Preclinical models also show immune responses to physical and psychological stressors (Ménard et al., 2017). Stress acts as a sterile immune challenge that elicits proinflammatory responses centrally and peripherally (Hodes et al., 2021).

Sterile immune responses increase the release of danger associated molecular patterns (DAMPs) which are recognized by pattern recognition receptors (PRRs). This response has been demonstrated to be an essential mechanism by which stress elicits behavioral changes (Weber et al., 2018). High mobility group box one (HMGB1) is one such DAMP released during stress that activates multiple PRRs including Toll-like receptors

(TLRs) and the receptor for advanced glycation end products (RAGE) (Kang, 2014). Previous results demonstrate that stress is associated with increased HMGB1 release in the brain and subsequent activation of TLR2, TLR4, and RAGE signaling (Pluma-Pluma et al., 2023), (Chen, 2022). Even with this knowledge, the mechanisms by which HMGB1 can impact physiology in response to stressors remains largely undefined.

Within the brain, nearly all cell types express HMGB1 or one of its putative receptors. HMGB1's role as an instigator of pro-inflammatory signaling is well established; however, little work has been done to investigate the cell type specificity of HMGB1 signaling in stress responses in the brain. Microglia, the resident innate immune cells of the brain, play key roles in proinflammatory signaling in response to any disruption of homeostasis, including stress. Microglia can respond broadly to stressors via multiple functional alterations. Previous studies show that microglia undergo changes in phagocytic activity, cytokine release, as well as signaling (Fleshner et al., 2017).

The hypothesis that inflammation is a critical mechanism which drives behavioral changes after stress exposure has substantial precedent in the literature (Won and Kim. 2020). Previous studies demonstrate that anti-inflammatory treatments prevent, and reverse behavioral deficits caused by stress in preclinical models (Basset et al., 2021). However, it is becoming clear that non-specific anti-inflammatory treatment is not sufficient to treat mood disorders (Hellmann-Regen et al., 2022). Therefore, more work is needed to unravel mechanisms of stress induced neuroinflammation to determine if therapeutics can be more precisely targeted.

Stress induces neuroinflammation in key stress responsive regions (Carvahlo-Netton et al., 2011) including the medial prefrontal cortex (mPFC) (Hinwood et al., 2012; Tynan et al., 2010). HMGB1 is increased in response to acute and chronic stress in the hippocampus and frontal cortex (Weber et al., 2015). PRRs are a key driver of inflammatory stress responses and microglia express PRRs at high levels relative to other CNS cells. Microglial PRRs are so key in stress response that genetic deletion of TLR2/4 yields a highly stress resilient phenotype (Nie et al., 2018). Therefore, we hypothesize that microglia have a key role in the mechanisms by which increased HMGB1 causes neuroinflammation in the mPFC and alters behavior.

Disulfide HMGB1 (dsHMGB1) is an established pharmacological stressor that alters depressive-like behavior when circulated throughout the whole brain via direct infusion (Lian et al., 2017). Here we utilize localized administration of HMGB1 to incorporate a region-specific understanding of behavior with immune signaling in the context of stress. We found that HMGB1 locally administered to the mPFC produces an anxiety-like phenotype in both male and female adult mice. We also see striking sex-specific differences in both microglial morphology as well as transcriptional activity following HMGB1 infusion. Finally, we found that in females specifically, conditional knockout of myeloid differentiation adaptor protein 88 (MyD88) in microglia prevents anxiety like behavior following HMGB1 infusion.

## Results

### HMGB1 infusion to mPFC increases anxiety-like behavior in males and females

HMGB1 has been previously shown to increase despair and anhedonia like behavior (Wang et al., 2018). However, no studies to date have utilized HMGB1 as a pharmacological stressor in female subjects calling in to question how relevant HMGB1- associated mechanisms are to stress response in females. Here, we sought to first define the impact of a sub-chronic exposure to a stress-related danger signal on behavior in male and female mice. To do so, mice were infused via cannula with dsHMGB1 once daily for 5 consecutive days. Behaviors related to anxiety, sociability, and depression/anhedonia were assayed to clearly define how increased HMGB1 in the mPFC alters behavior. Anxiety-like behavior was assessed by measuring center exploration time in the Open field test **(Suppl Fig1A),** open arm exploration and latency to explore open arms of the elevated plus maze **(Fig 1B-C)**, and the latency to feed in the novelty suppressed feeding assay **(Fig1E)**. No changes in social preference or behavior in the tail suspension test were found **(Suppl Fig1B-C).** The results show that dsHMGB1 infusion to the mPFC selectively alters anxiety and this phenotype is shared between males and females. Furthermore, we found that females specifically had a significant change in weight associated with dsHMGB1 treatment **(Fig 1D).** Additionally, we investigated if hormonal changes related to estrus cycle were correlated with behavior and found no significant correlation **(Suppl Fig 2A-C)**. Taken together, these results demonstrate that HMGB1 elicits a shared behavioral phenotype of increased anxiety like behavior in male and female mice.

### Microglial activation is specific to females following dsHMGB1 infusion to mPFC

The mPFC is often described as a key node for stress responsivity and emotional processing; these are key processes in the context of anxiety. Sex differences in stress associated microglial activation are well documented (Bollinger et al., 2016). We sought to determine if microglia activation would be a shared phenotype between males and females underlying the similar behavioral outcomes. In a smaller cohort of mice, we repeated the implantation and dosing scheme as outlined in the previous section. However, 16 hours following the final infusion of dsHMGB1, mice completed the elevated plus maze assay and were immediately sacrificed for tissue collection. Again, an anxiety phenotype in the EPM was observed in males and females **(Suppl. Figure3A and 3B).** We triple stained Iba1, Tmem119, and CD68 to distinguish between surveillant, homeostatic, and phagocytic microglia respectively. We quantified volume of Iba1 and Tmem119 **(Fig 2D-G)**. There was no significant difference in Tmem119 volume for males or females; however, Iba1 volume differed significantly in females ($p<0.05$) **(Fig 2G)**. We also calculated the ratio of Tmem19 normalized to Iba1 for each reconstructed cell. We interpret this measurement as a ratio of homeostasis, as Tmem119 is often downregulated in the context of inflammation, injury, or disease progression (Spiteri et al., 2021). Female mice treated with dsHMGB1 showed decrease Tmem119:Iba1 relative to controls ($p<0.05$) **(Fig 2I)**. In addition to quantifying volume of Tmem119 and IBA1, we also calculated phagocytic capacity by normalizing CD68 volume to Tmem119 and IBA1 merged volume. Females treated with dsHMGB1 had significantly increased phagocytic capacity ($p<0.05$) **(Fig 2J)**. Finally, we examined changes in morphology by calculating the number of Sholl intersections from various distances from the radius of each cell soma. In both males and females, we sound significant changes in the number of intersections as you move distally from the soma **(Fig 2F and Fig2K)**. In both males and females, dsHMGB1 decreases the number of intersections and we interpret this to reflect a decrease in complexity and an increase in reactivity.

**dsHMGB1 exerts sex specific changes in RAGE/MyD88 signaling**

Noting the sex-specific effects of dsHMGB1 on microglial measures, we hypothesized there could also be sex-specific changes in proinflammatory signaling pathways. HMGB1 is a putative ligand for TLR4 and RAGE pathways (Xu et al., 2019; Pluma et al., 2020). Canonical activation of either of these pathways requires subsequent activation of MyD88. We utilized a CD11b+ cell specific isolation protocol where CD11b marks myeloid cells, which are solely expressed by microglia in the brain. CD11b- and CD11b+ cell fractions were isolated from mice 16 hours after the final dose to maintain the same timeframe from behavioral and immunohistochemical experiments. All cells were isolated from the frontal cortex. In males, no significant changes were found in TLR4, RAGE, or MyD88. Female CD11b+ cells showed significant increase in RAGE **(Fig 3D)** and MyD88 **(Fig 3F)**. Surprisingly an increase in MyD88 was also observed in the CD11b- cells for dsHMGB1 treated females **(Fig 3F).**

**Microglial MyD88 is necessary for HMGB1 induced anxiety in females**

We initially hypothesized that microglial MyD88 was necessary for anxiety-like behavioral responses to HMGB1. To test this hypothesis, we bred a previously characterized mouse line that has conditional deletion of the MyD88 gene for CX3CR1 expressing cell types (Rivera et al., 2019), which includes all microglia in the CNS. The experimental workflow is the same as previous behavioral aims. Both male and female control mice responded with an anxiety-like phenotype following dsHMGB1 treatment as we have seen previously. However, female cKO mice do not demonstrate an anxiety-like phenotype in the EPM or Novelty Suppressed Feeding (NSF) assay (\*\*$p < .01$, 2- way ANOVA) **(Figure 4C and 4E)**.

## Discussion

HMGB1 preclinical models are emerging as a robust and reliable way to induce stress associated behavioral effects (Huang et al., 2023). HMGB1 models exert stress associated behavioral changes and are reversed by antidepressants as well as minocycline (Lian et al., 2017; Wang et al., 2019). More recent studies, including this one, demonstrate HMGB1's utility as a pharmacological stressor which can impact specific behavior based on site of drug delivery. Moreover, HMGB1 does so while enacting other physiological changes associated with stress potentially providing novel insights into the pathophysiology of mood disorders.

Though mood disorders including anxiety and depression are twice as common in women, the current body of work related to stress and mood disorders has been primarily done in male subjects (Kalin., 2020; Farhane- Medina et al., 2019). Pre-clinical stress models do not utilize female subjects to a comparable degree as male subjects nor to a degree that mimics the increased prevalence of these disorders in women (Lopez and Bagot, 2021). Various models of stress are utilized throughout the field, each exerting both specific and generalizing effects (Saxena et al., 2021). A now growing body of work is revisiting these canonical mechanisms of stress response in female subjects and uncovering as much convergence as there is distinction. In the present study our findings echo this trend; for example, here male and female adult mice infused with HMGB1 in the mPFC exhibit a shared anxiety-like phenotype as in the EPM and NSF assay as well as significant morphological changes; however, increases in microglial activation were not observed in both sexes.

Increased microglial reactivity is a necessary mechanism by which stress alters behavior (Lehmann et al.,2019). Very little is established in the literature concerning understanding stress associated HMGB1 mechanisms in a cell specific manner. In this present study, we hypothesized that microglial activation would prove to be a crucial mechanism by which HMGB1 induces behavioral changes. We examined previously identified hallmarks of stress associated microglial pathology. Only females demonstrated significant alterations in microglia volume, phagocytic capacity, and homeostatic marker expression. Similarly, females specifically displayed significant increases in transcription of RAGE and MyD88 in microglial cells isolated from the frontal cortex. The role of microglial RAGE in the context of stress was first identified in male rats (Franklin et al., 2018). Here we demonstrate that adult female mice display increased activation of both microglial RAGE and MyD88 following HMGB1 infusion. Finally, we demonstrate that microglial MyD88 is necessary for HMGB1 induced anxiety by comparing behavioral responses to HMGB1 between control and cKO mice. Taken together this work provides illuminating findings concerning the cell specificity of HMGB1 signaling in the brain.

We must acknowledge several limitations regarding our study. Firstly, our intention was to examine how HMGB1 impacts behavior when administered with site-specificity in the mPFC. To do so, we infused 75nL of 20% dsHMGB1 to the mPFC once daily for five consecutive days. This dose was validated by its ability to significantly alter anxiety-like behaviors in males. We then utilized this same dosing scheme in the female experimental group. Previous studies utilize dsHMGB1 at a significantly higher dose and over a longer period report a similar behavioral effect (Du et al., 2022). It is possible that some findings may be dose dependent. Specifically, the transcriptional responses in male microglia trend in opposite directions than females. This was not surprising as it has been previously demonstrated that sub chronic stress transiently downregulates RAGE and TLR4 (Franklin et al., 2018). However, it highlights the importance of potentially utilizing this model in both sexes across a broader dosing range to better delineate the tunability of HMGB1 responses.

Secondly and possibly most importantly it would be exciting to draw broader conclusions about the role of microglial MyD88 in stress response. However, more work is needed to understand the role of microglial MyD88 in stress response in such a way that generalizes across stress models.

There are several interesting future directions based on the findings of this study. Now that we have

elucidated the importance of microglial MyD88 in female HMGB1 response, understanding the time course of resolution as well as investigating how this mechanism relates to stress-induced priming would be an interesting investigational aim. In line with this, we observed an almost bivalent effect of HMGB1 at this dose in both males and females. As previously stated, we utilized a significantly lower dose of HMGB1 that we believe allowed us to potentially capture a pseudo resilient v. susceptible phenotype. It would be interesting to follow up this observation with studies that examine how HMGB1 response correlates with response to stressors and subthreshold stressors.

This study adds to a growing body of work that demonstrates that similar behavioral response following stress may be the result of distinct sex-specific mechanisms. We also report the first use of HMGB1 as a model of mood disorders in female adult mice. Moreover, our findings provide further evidence for the hypothesis that the ability of microglia to confer to vulnerability to disease shifts from being male-biased in early life and development to female-biased as we age (Lynch., 2022). Models of localized HMGB1 drug delivery could be utilized to further probe the effects of neuroinflammation with increased specificity. Expanding our understanding of mechanisms upstream of HMGB1 release as well as downstream signaling could continue to provide novel insights into stress pathology with potential broader implications. In conclusion, the data found in this study supports utilization of a local HMGB1 model to provide insight into pathological stress with great potential for future work and possible utility for therapeutic development.

## Materials and Methods

### Mice

MyD88-floxed mice were purchased from Jackson Laboratories (Bar Harbor, ME; Stock # 00888). Cx3Cr1-CreBT (MW126GSat) mice were generated and provided by L. Kus (GENSAT BAC Transgenic Project, Rockefeller University, NY). All methods for breeding and genotyping transgenic mice were conducted according to Rivera et al., 2019. Cx3Cr1- CreBT mice were crossed with MyD88-floxed mice over 3 generations until all offspring were fully MyD88-floxed (F/F) and either Cre negative (Cre 0/0: no modification to microglial MyD88) or Cre positive (Cre tg/0: removal of microglial MyD88), and then back-crossed onto a fully Jackson background. Genotyping of transgenic animals was conducted using polymerase chain reaction (PCR) on tail-snip DNA. Primer sequences used for PCR can be found in Table 1. Wild-type (WT) C57Bl/6J mice (used as stimuli for behavioral experiments) were purchased from Jackson Laboratories (Bar Harbor, ME; Stock # 000664). All animals were group housed in standard mouse cages under standard laboratory conditions (12- hour light/dark cycle, 23°C, 60% humidity) with same-sex littermates.

### Surgery

Mice were anesthetized with a solution of ketamine/xylazine (100/10 mg/kg) and stereotaxically implanted with an intracerebral ventricular (ICV) guide cannula (26G, 2.25mm from the pedestal; Plastics One, VA) into the left medial prefrontal cortex using the following coordinates from bregma (33): 1.5mm anterior-posterior, −0.4mm medial-lateral, and
−2.25mm dorsal-ventral. After 10 days of recovery, the free moving mice were infused with 75nl of aCSF or recombinant HMGB1 (dsHMGB1 certified LPS free (0.2ug/ul); HMGBiotech) using a 26G internal cannula (Plastics One, VA) once daily for five consecutive days.

### Estrus Cycle Analysis

Smears were obtained by vaginal cytology collected at the end of behavioral tests and stained using the heamatoxylin- eosin method. Three cell types (nucleated epithelial cells, cornified epithelial cells and leukocytes) were counted to define the reproductive cycle (estrous), which is defined by the prevalence of each cell type: proestrus (nucleated), estrus (cornified), metestrus (all types in same proportion) and diestrus (leukocytes) (Westwood, 2008). Images were acquired with a light microscope Leica DM 4000 B (Leica, Wetzlar, Germany) with a ×10 objective lens (Plan 10×/0.25PH1).

### Weight Collections

Animals were weighed on the day of and the day preceding dosing. This weight was averaged to establish a baseline. Animals were weighed daily on dosing days and behavior days. The weight change reported for (Fig 1D) is the difference between baseline weight and weight 24 hours after dosing ends.

### Behavioral Tests

All behavioral tests were conducted approximately 12 hours apart from one another alternating between light and dark phase. Mice were habituated in the testing room for at least 30 min before the test. Examiners were blinded to the groups of mice. The testing room was sound-proof, and the temperature was maintained at 24–26 °C. After each trial, the mice were put back into their home cages and the experimental apparatus was cleaned with 75% alcohol and Rescue cleaning solution to eliminate the odor that may affect animal behavior. Behaviors were tested in order consistent with experimental outlines provided in Figure1A and Figure 4A.

**Open Field Test (OFT)**

The OFT was performed in an open rectangular box with a bottom edge length of 50 cm and a height of 60 cm. The box was placed directly on the floor. After placing the mouse in the intermediate region of the box, animal behavior was recorded by a camera for 10 minutes. The bottom of the box was divided into the central zone ($25 \times 25$ cm$^2$) and the rest of the outer zone by EthovisionXT. The time spent by the animal in each region as well as the moving distance in the box were calculated.

**Elevated Plus Maze (EPM)**

The apparatus used for the EPM test was a cross-shaped device consisting of an intermediate platform region ($5 \times 0.5$ cm$^2$) and two pairs of open arms and closed arms ($25 \times 5$ cm$^2$) connected to one another. The closed arms are surrounded by walls (16 cm), whereas the open arms have slight walls (0.5 cm). The entire maze was elevated to a height of 50 cm. The mouse was first placed in the intermediate region at the same position with its head toward closed arms, and its behavior in the device was then recorded by a camera for 5 min and the time spent by the animal in each arm was analyzed by EthovisionXT.

**Social Preference Assay**

Social preference was measured using a two-chamber assay in which animals explored a novel object or a novel mouse. The apparatus was a rectangular arena (61cm × 42.5cm × 22cm, L×W×H) constructed from clear plexiglass with a clear plexiglass wall dividing the arena into two equal chambers with an opening in the middle allowing free access between both chambers. Plastic, circular holding cages (8.3cm diameter and 12cm tall) were centered in each of the two chambers and were used to house either a novel object or sex- and age-matched C3H target mouse. The arena was evenly lit with indirect white light (~125 lux). Plastic toys and glass objects were used as novel objects with the object being between 3– 5cm in all directions.

**Novelty Suppressed Feeding**

Following the completion of EPM, mice were food-deprived 18 hours prior to the test, with free access to water. On the day of testing, mice were moved to the dimly lit testing room one hour before the test. Mice were placed into one corner of a clear plexiglass open field apparatus (17 in. × 17 in. × 12 in.) with an opaque white acrylic floor covered with a thin layer of home cage bedding material. The room was maintained with red light conditions. A food pellet was placed in the center of the open field and mice were placed in one corner. Latencies to approach and to begin eating were recorded with a limit of 8 minutes. As soon as the mouse was observed to eat, or the 8-minute time limit was reached, the mouse was removed from the open field. And placed in the home cage and observed until it began to eat in the home cage.

**Home Cage Feeding Assay**

Immediately following completion of the NSF trial. Mice were then placed in a cage with bedding material from their home cage to increase familiarity within the arena. Again, a mouse began the trial in the corner of the cage and the food pellet was placed in the center. Each trial was limited to 3 minutes and at the end of that time or after the mouse began its first feeding bout, the mouse was removed, and the trial completed. The latency to feed was recorded as the trial occurred and confirmed via video hand scoring.

**Light dark box (LDB) test**

The apparatus for LDB test was a rectangular box comprising two connected chambers: a light open chamber with white walls (~ 100 Lux, $15 \times 20 \times 25$ cm$^3$) and a dark covered chamber with black walls (~ 5 Lux, $30 \times 20 \times 25$ cm$^3$). The apparatus was placed on a table two feet off the ground. For every trial, the mouse was placed at the same position in the dark box. Animal behavior in the apparatus was then recorded for 8 min by a digital camera and the time the animal spent in each box was analyzed by EthovisionXT.

**Tail Suspension Test**

For TST, mice were suspended from the side of a behavioral arena with white background for video tracking. At the beginning of the test, the mouse was fixed to the side of the arena with tape 1.5cm wide and approximately 15cm long. Mice were taped at the distal end of the tail and suspended 15 cm from the base surface. The camera was then turned on to record the animal behavior for 6 min. Immobility was defined by mobility (> 2.8%) and velocity parameters (> 3 cm/s) set in EthovisionXT.

**Immunohistochemistry**

Mice were perfused intracardially with ice-cold saline followed by phosphate-buffered 4% paraformaldehyde (pH 7.4) after anesthetized with $CO_2$. The brain was removed and post-fixed overnight in the same fixative and dehydrated in 30% sucrose for 48 h at 4 °C. Coronal brain sections were cut at 30 μm by a cryostat (NX50, Thermo, USA). For immunohistochemical staining, the sections containing the mPFC, conventionally implicated in anxiety, were incubated in 0.1% Triton X-100 for 15 min and in 5% normal goat serum for 2 h firstly, and then were incubated with rabbit, chicken, or guinea pig anti-mouse primary antibodies against Iba1 (1:500, Abcam, UK) /TMEM119 (1:500, Millipore, USA) /Iba1 (1:500, Wako, Japan) or CD68 (1:500) overnight at 4 °C, then with anti-rabbit IgG-Alexa Fluor 488 or anti-rat IgG-Alexa Fluor 594 (1:1000, Invitrogen, USA) for 2 h at room temperature (RT). After repeated washing, the sections were then covered with glass coverslips and fluorescent images were captured by a fluorescence microscope (BX51, Olympus, Japan). The analysis of fluorescence intensity and cell counting were performed by Image J software (NIH, USA). Microglia reconstructions and individual cell analysis was performed utilizing IMARIS 10.0 software. Structural analysis included the total length and the 3D arrangement of microglial branching using the Sholl analysis (number of intersections with concentric circles positioned at radial intervals of 1 μm). For each animal, 6-8 microglia from mPFC were analyzed and microglia from the same animal were averaged. Observers were blinded to the experimental condition of each subject.

**RNA Extraction**

Samples were homogenized in Trizol® (Thermo-fisher scientific, NY) and then vortexed for 10 min at 2000 rpm. After 15 min resting at room temperature (RT), chloroform was added (1:5 with Trizol) and samples were vortexed for 2 min at 2000 rpm. Samples were next allowed to sit at RT for 3 min and then centrifuged (15 min at 11,800 rpm; 4°C). This resulted in a gradient from which the top, clear, aqueous phase was separated and placed in to a fresh tube. Isopropanol was then added to the new tube to precipitate RNA (1:1 with aqueous phase) and samples were again vortexed, allowed to set at RT for 10 min, and then centrifuged (15 min at 11,800 rpm; 4°C). Pellets obtained after this step were rinsed twice in ice-cold 75% Ethanol and then resuspended in 8 μl of nuclease-free water. RNA was frozen at −80°C until cDNA synthesis and qPCR. A NanoDrop Spectrophotometer (Thermo Scientific, Wilmington, DE) was used to determine RNA quantity and purity. RNA was considered pure enough for further use based on 260/280 (RNA:protein; range: 1.8–2.0) and 260/230 (RNA: Ethanol; range 1.6–2.0) ratios.

**qPCR**

cDNA was synthesized using the QuantiTect Reverse Transcription Kit (Quiagen, Hilden, Germany). Briefly, 200ng of RNA/12 μl of nuclease-free water was pre-treated with gDNase at 42°C for 2 min to remove genomic DNA contamination. Next, master mix containing both primer-mix and reverse transcriptase was added to each

sample and all samples were heated to 42°C for 30 min and then 95°C for 3 min (to inactivate the reaction) in the thermocycler. Quantitative real-time PCR (qPCR) was conducted using QuantiTect SYBR Green PCR Kit (Quiagen, Catalog #:204057, Hilden, Germany. To examine HMGB1-dependent signaling that is mediated by the MyD88-dependent pathways, we selected 4 genes within the signaling cascade of HMGB1 including putative receptors RAGE, TLR4, as well as MyD88 (Magna & Pisetsky, 2014). qPCR primers were designed in the lab and ordered from Integrated DNA technologies (Coralville, IA). All primer sequences can be found in Table 1. PCR product was monitored using a Mastercycler ep *realplex* (Eppendorf, Hauppauge, NY). Male and female samples were run on separate plates, and thus, cannot be directly compared. I8S was used as a house-keeping gene and relative gene expression was calculated using the $2-\Delta\Delta^{CT}$ method, relative to 18S and the lowest sample on the plate (Williamson et al., 2011; Livak & Schmittgen, 2001).

## Statistical Analysis

All data are expressed as the mean ± SEM. The required sample sizes were estimated based on our experience and Power analysis was used to justify the sample size. Statistical analysis was conducted by GraphPad Prism 10.2 (GraphPad Software, USA). The Shapiro–Wilk test was used to assess whether the data followed a normal distribution. If the data was normally distributed, one-tailed paired or unpaired Student's $t$ test was used for comparison between two groups; if not, Mann–Whitney test was used instead. For groups with a normal distribution, a two-way ANOVA was utilized to assess the significance of two factors and the interaction of those factors (ie: sex and treatment) either Bonferroni or Dunnett's post-hoc test was used. The significance level was set at $P < 0.05$.

## Acknowledgements
We acknowledge that the contents of this study do not represent the views of the National Institutes of Health, or the United States Government.

## Funding
Research reported in this publication was supported by NIH R01-ES033056 and NIH U01 AA029969 and the content in this publication does not reflect the view of the NIH.

# References

Amer SAAM, Fouad AM, El-Samahy M, et al. Mental Stress, Anxiety and Depressive Symptoms and Interleuken-6 Level among Healthcare Workers during the COVID-19 Pandemic. *J Prim Care Community Health*. 2021;12:21501327211027432. doi:10.1177/21501327211027432

Bandelow B, Michaelis S. Epidemiology of anxiety disorders in the 21st century. Dialogues Clin Neurosci. 2015;17(3):327-335. doi:10.31887/DCNS.2015.17.3/bbandelow

Bassett B, Subramaniyam S, Fan Y, et al. Minocycline alleviates depression-like symptoms by rescuing decrease in neurogenesis in dorsal hippocampus via blocking microglia activation/phagocytosis. Brain Behav Immun. 2021;91:519- 530. doi:10.1016/j.bbi.2020.11.009

Baugher BJ, Buckhaults K, Case J, Sullivan A, Huq SN, Sachs BD. Sub-chronic stress induces similar behavioral effects in male and female mice despite sex-specific molecular adaptations in the nucleus accumbens. Behav Brain Res. 2022;425:113811. doi:10.1016/j.bbr.2022.113811

Bauer ME, Teixeira AL. Inflammation in psychiatric disorders: what comes first?. *Ann N Y Acad Sci*. 2019;1437(1):57-67. doi:10.1111/nyas.13712

Bollinger JL, Bergeon Burns CM, Wellman CL. Differential effects of stress on microglial cell activation in male and female medial prefrontal cortex. *Brain Behav Immun*. 2016;52:88-97. doi:10.1016/j.bbi.2015.10.003

Bollinger JL, Wohleb ES. The formative role of microglia in stress-induced synaptic deficits and associated behavioral consequences. *Neurosci Lett*. 2019;711:134369. doi:10.1016/j.neulet.2019.134369

Brydges CR, Bhattacharyya S, Dehkordi SM, et al. Metabolomic and inflammatory signatures of symptom dimensions in major depression. *Brain Behav Immun*. 2022;102:42-52. doi:10.1016/j.bbi.2022.02.003

Butler MI, Long-Smith C, Moloney GM, et al. The immune-kynurenine pathway in social anxiety disorder. *Brain Behav Immun*. 2022;99:317-326. doi:10.1016/j.bbi.2021.10.020

Carvalho-Netto EF, Myers B, Jones K, Solomon MB, Herman JP. Sex differences in synaptic plasticity in stress- responsive brain regions following chronic variable stress. *Physiol Behav*. 2011;104(2):242-247. doi:10.1016/j.physbeh.2011.01.024

Cathomas F, Holt LM, Parise EM, et al. Beyond the neuron: Role of non-neuronal cells in stress disorders. *Neuron*. 2022;110(7):1116-1138. doi:10.1016/j.neuron.2022.01.033

Cathomas F, Lin HY, Chan KL, et al. Circulating myeloid-derived MMP8 in stress susceptibility and depression. *Nature*. 2024;626(8001):1108-1115. doi:10.1038/s41586-023-07015-2

Chan KL, Poller WC, Swirski FK, Russo SJ. Central regulation of stress-evoked peripheral immune responses. *Nat Rev Neurosci*. 2023;24(10):591-604. doi:10.1038/s41583-023-00729-2

Chen X, Cui QQ, Hu XH, et al. CD200 in dentate gyrus improves depressive-like behaviors of mice through enhancing hippocampal neurogenesis via alleviation of microglia hyperactivation. *J Neuroinflammation*. 2023;20(1):157. Published 2023 Jun 30. doi:10.1186/s12974-023-02836-4

Cooke JP. Inflammation and Its Role in Regeneration and Repair. *Circ Res*. 2019;124(8):1166-1168. doi:10.1161/CIRCRESAHA.118.314669

Deppermann S, Storchak H, Fallgatter AJ, Ehlis AC. Stress-induced neuroplasticity: (mal)adaptation to adverse life events in patients with PTSD--a critical overview. *Neuroscience*. 2014;283:166-177. doi:10.1016/j.neuroscience.2014.08.037

Dion-Albert L, Dudek KA, Russo SJ, Campbell M, Menard C. Neurovascular adaptations modulating cognition, mood, and stress responses. *Trends Neurosci*. 2023;46(4):276-292. doi:10.1016/j.tins.2023.01.005

Farhane-Medina NZ, Luque B, Tabernero C, Castillo-Mayén R. Factors associated with gender and sex differences in anxiety prevalence and comorbidity: A systematic review. *Sci Prog*. 2022;105(4):368504221135469. doi:10.1177/00368504221135469

Fleshner M, Frank M, Maier SF. Danger Signals and Inflammasomes: Stress-Evoked Sterile Inflammation in Mood Disorders. *Neuropsychopharmacology*. 2017;42(1):36-45. doi:10.1038/npp.2016.125

Fonken LK, Frank MG, Gaudet AD, et al. Neuroinflammatory priming to stress is differentially regulated in male and female rats. *Brain Behav Immun*. 2018;70:257-267. doi:10.1016/j.bbi.2018.03.005

Fonken LK, Frank MG, Gaudet AD, Maier SF. Stress and aging act through common mechanisms to elicit neuroinflammatory priming. *Brain Behav Immun*. 2018;73:133-148. doi:10.1016/j.bbi.2018.07.012

Frank MG, Fonken LK, Annis JL, Watkins LR, Maier SF. Stress disinhibits microglia via down-regulation of CD200R: A mechanism of neuroinflammatory priming. *Brain Behav Immun*. 2018;69:62-73. doi:10.1016/j.bbi.2017.11.001

Franklin TC, Wohleb ES, Zhang Y, Fogaça M, Hare B, Duman RS. Persistent Increase in Microglial RAGE Contributes to Chronic Stress-Induced Priming of Depressive-like Behavior. Biol Psychiatry. 2018;83(1):50-60. doi:10.1016/j.biopsych.2017.06.034

Franklin TC, Xu C, Duman RS. Depression and sterile inflammation: Essential role of danger associated molecular patterns. *Brain Behav Immun*. 2018;72:2-13. doi:10.1016/j.bbi.2017.10.025

Frederiksen HR, Haukedal H, Freude K. Cell Type Specific Expression of Toll-Like Receptors in Human Brains and Implications in Alzheimer's Disease. Biomed Res Int. 2019;2019:7420189. Published 2019 Jul 18. doi:10.1155/2019/7420189

González Ibáñez F, Halvorson T, Sharma K, et al. Ketogenic diet changes microglial morphology and the hippocampal lipidomic profile differently in stress susceptible versus resistant male mice upon repeated social defeat. *Brain Behav Immun*. 2023;114:383-406. doi:10.1016/j.bbi.2023.09.006

Hellmann-Regen J, Clemens V, Grözinger M, et al. Effect of Minocycline on Depressive Symptoms in Patients With Treatment-Resistant Depression: A Randomized Clinical Trial. JAMA Netw Open. 2022;5(9):e2230367. Published 2022 Sep 1. doi:10.1001/jamanetworkopen.2022.30367

Hinwood M, Morandini J, Day TA, Walker FR. Evidence that microglia mediate the neurobiological effects of chronic psychological stress on the medial prefrontal cortex. *Cereb Cortex*. 2012;22(6):1442-1454. doi:10.1093/cercor/bhr229

Hodes GE, Pfau ML, Leboeuf M, et al. Individual differences in the peripheral immune system promote resilience versus susceptibility to social stress. Proc Natl Acad Sci U S A. 2014;111(45):16136-16141. doi:10.1073/pnas.1415191111

Hodes GE, Kana V, Menard C, Merad M, Russo SJ. Neuroimmune mechanisms of depression. Nat Neurosci. 2015;18(10):1386-1393. doi:10.1038/nn.4113

Hopperton KE, Mohammad D, Trépanier MO, Giuliano V, Bazinet RP. Markers of microglia in post-mortem brain samples from patients with Alzheimer's disease: a systematic review. Mol Psychiatry. 2018;23(2):177-198. doi:10.1038/mp.2017.246

Lee EH, Park JY, Kwon HJ, Han PL. Repeated exposure with short-term behavioral stress resolves pre-existing stress- induced depressive-like behavior in mice. *Nat Commun*. 2021;12(1):6682. Published 2021 Nov 18. doi:10.1038/s41467- 021-26968-4

Lehmann, M.L., Weigel, T.K., Cooper, H.A. et al. Decoding microglia responses to psychosocial stress reveals blood- brain barrier breakdown that may drive stress susceptibility. Sci Rep 8, 11240 (2018). https://doi.org/10.1038/s41598-018- 28737-8

Lehmann ML, Weigel TK, Poffenberger CN, Herkenham M. The Behavioral Sequelae of Social Defeat Require Microglia and Are Driven by Oxidative Stress in Mice. *J Neurosci*. 2019;39(28):5594-5605. doi:10.1523/JNEUROSCI.0184- 19.2019

Lenz KM, Nelson LH. Microglia and Beyond: Innate Immune Cells As Regulators of Brain Development and Behavioral Function. *Front Immunol*. 2018;9:698. Published 2018 Apr 13. doi:10.3389/fimmu.2018.00698

Leuner B, Shors TJ. Stress, anxiety, and dendritic spines: what are the connections?. *Neuroscience*. 2013;251:108-119. doi:10.1016/j.neuroscience.2012.04.021

Lynch MA. Exploring Sex-Related Differences in Microglia May Be a Game-Changer in Precision Medicine. *Front Aging Neurosci*. 2022;14:868448. Published 2022 Mar 31. doi:10.3389/fnagi.2022.868448

Kalin NH. Novel Insights Into Pathological Anxiety and Anxiety-Related Disorders. *Am J Psychiatry*. 2020;177(3):187-189. doi:10.1176/appi.ajp.2020.20010057

Kim G, Lee Y, Jeong EY, et al. Acute stress responsive RGS proteins in the mouse brain. *Mol Cells*. 2010;30(2):161-165. doi:10.1007/s10059-010-0102-3

Kim SJ, Ryu MJ, Han J, et al. Non-cell autonomous modulation of tyrosine hydroxylase by HMGB1 released from astrocytes in an acute MPTP-induced Parkinsonian mouse model. *Lab Invest*. 2019;99(9):1389-1399. doi:10.1038/s41374- 019-0254-5

Kokkosis AG, Madeira MM, Hage Z, et al. Chronic psychosocial stress triggers microglial-/macrophage-induced inflammatory responses leading to neuronal dysfunction and depressive-related behavior. *Glia*. 2024;72(1):111-132. doi:10.1002/glia.24464

Labonté B, Engmann O, Purushothaman I, et al. Sex-specific transcriptional signatures in human depression [published correction appears in Nat Med. 2018 Apr 10;24(4):525]. *Nat Med*. 2017;23(9):1102-1111. doi:10.1038/nm.4386

Magna M, Pisetsky DS. The role of HMGB1 in the pathogenesis of inflammatory and autoimmune diseases. *Mol Med*. 2014;20(1):138-146. Published 2014 Mar 24. doi:10.2119/molmed.2013.00164

McEwen BS, Akil H. Revisiting the Stress Concept: Implications for Affective Disorders. *J Neurosci*. 2020;40(1):12-21. doi:10.1523/JNEUROSCI.0733-19.2019

McLean CP, Asnaani A, Litz BT, Hofmann SG. Gender differences in anxiety disorders: prevalence, course of illness, comorbidity and burden of illness. J Psychiatr Res. 2011;45(8):1027-1035. doi:10.1016/j.jpsychires.2011.03.006

Ménard C, Pfau ML, Hodes GE, Russo SJ. Immune and Neuroendocrine Mechanisms of Stress Vulnerability and Resilience. *Neuropsychopharmacology*. 2017;42(1):62-80. doi:10.1038/npp.2016.90

Mondelli V, Vernon AC, Turkheimer F, Dazzan P, Pariante CM. Brain microglia in psychiatric disorders. *Lancet Psychiatry*. 2017;4(7):563-572. doi:10.1016/S2215-0366(17)30101-3

Nimmerjahn A, Kirchhoff F, Helmchen F. Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. Science. 2005;308(5726):1314-1318. doi:10.1126/science.1110647

O'Donovan A, Hughes BM, Slavich GM, et al. Clinical anxiety, cortisol and interleukin-6: evidence for specificity in emotion-biology relationships. *Brain Behav Immun*. 2010;24(7):1074-1077. doi:10.1016/j.bbi.2010.03.003

Park SC, Kim YK. Anxiety Disorders in the DSM-5: Changes, Controversies, and Future Directions. *Adv Exp Med Biol*. 2020;1191:187-196. doi:10.1007/978-981-32-9705-0_12

Pluma-Pluma A, García G, Murbartián J. Chronic restraint stress and social transfer of stress produce tactile allodynia mediated by the HMGB1/TNFα/TNFR1 pathway in female and male rats. *Physiol Behav*. 2024;274:114418. doi:10.1016/j.physbeh.2023.114418

Rainville JR, Hodes GE. Inflaming sex differences in mood disorders. *Neuropsychopharmacology*. 2019;44(1):184-199. doi:10.1038/s41386-018-0124-7

Sessa L, Gatti E, Zeni F, et al. The receptor for advanced glycation end-products (RAGE) is only present in mammals, and belongs to a family of cell adhesion molecules (CAMs). *PLoS One*. 2014;9(1):e86903. Published 2014 Jan 27. doi:10.1371/journal.pone.0086903

Shimo Y, Cathomas F, Lin HY, et al. Social stress induces autoimmune responses against the brain. *Proc Natl Acad Sci U S A*. 2023;120(49):e2305778120. doi:10.1073/pnas.2305778120

Slavich GM, Sacher J. Stress, sex hormones, inflammation, and major depressive disorder: Extending Social Signal Transduction Theory of Depression to account for sex differences in mood disorders. *Psychopharmacology (Berl)*. 2019;236(10):3063-3079. doi:10.1007/s00213-019-05326-9

Sparvero LJ, Asafu-Adjei D, Kang R, et al. RAGE (Receptor for Advanced Glycation Endproducts), RAGE ligands, and their role in cancer and inflammation. *J Transl Med*. 2009;7:17. Published 2009 Mar 17. doi:10.1186/1479-5876-7-17

Tynan RJ, Naicker S, Hinwood M, et al. Chronic stress alters the density and morphology of microglia in a subset of stress-responsive brain regions. *Brain Behav Immun*. 2010;24(7):1058-1068. doi:10.1016/j.bbi.2010.02.001

Wang F, Yang X, Ren Z, Chen C, Liu C. Alternative splicing in mouse brains affected by psychological stress is enriched in the signaling, neural transmission and blood-brain barrier pathways. *Mol Psychiatry*. 2023;28(11):4707-4718. doi:10.1038/s41380-023-02103-1

Wang J, Russo SJ. Neurobiology of social stress and age-related neurodegeneration. *Neurosci Biobehav Rev*. 2024;156:105482. doi:10.1016/j.neubiorev.2023.105482

Weber MD, McKim DB, Niraula A, et al. The Influence of Microglial Elimination and Repopulation on Stress Sensitization Induced by Repeated Social Defeat. *Biol Psychiatry*. 2019;85(8):667-678. doi:10.1016/j.biopsych.2018.10.009

Westwood FR. The female rat reproductive cycle: a practical histological guide to staging. Toxicol Pathol. 2008;36(3):375-384. doi:10.1177/0192623308315665

Won E, Kim YK. Neuroinflammation-Associated Alterations of the Brain as Potential Neural Biomarkers in Anxiety Disorders. *Int J Mol Sci*. 2020;21(18):6546. Published 2020 Sep 7. doi:10.3390/ijms21186546

Woodburn SC, Bollinger JL, Wohleb ES. Synaptic and behavioral effects of chronic stress are linked to dynamic and sex- specific changes in microglia function and astrocyte dystrophy. *Neurobiol Stress*. 2021;14:100312. Published 2021 Mar 4. doi:10.1016/j.ynstr.2021.100312

Xu B, Zang SC, Li SZ, et al. HMGB1-mediated differential response on hippocampal neurotransmitter disorder and neuroinflammation in adolescent male and female mice following cold exposure. *Brain Behav Immun*. 2019;76:223-235. doi:10.1016/j.bbi.2018.11.313

Xue J, Suarez JS, Minaai M, et al. HMGB1 as a therapeutic target in disease. *J Cell Physiol*. 2021;236(5):3406-3419. doi:10.1002/jcp.30125
Yin W, Gallagher NR, Sawicki CM, McKim DB, Godbout JP, Sheridan JF. Repeated social defeat in female mice induces anxiety-like behavior associated with enhanced myelopoiesis and increased monocyte accumulation in the brain. *Brain Behav Immun*. 2019;78:131-142. doi:10.1016/j.bbi.2019.01.015

Zemskova M, Kurdyukov S, James J, McClain N, Rafikov R, Rafikova O. Sex-specific stress response and HMGB1 release in pulmonary endothelial cells. *PLoS One*. 2020;15(4):e0231267. Published 2020 Apr 9. doi:10.1371/journal.pone.0231267

Zhang S, Hu L, Jiang J, et al. HMGB1/RAGE axis mediates stress-induced RVLM neuroinflammation in mice via impairing mitophagy flux in microglia. *J Neuroinflammation*. 2020;17(1):15. Published 2020 Jan 10. doi:10.1186/s12974- 019-1673-3

Zhang Y, Xu H, Zhang F, et al. Deficiencies of microglia and TNFα in the mPFC-mediated cognitive inflexibility induced by social stress during adolescence. *Brain Behav Immun*. 2019;79:256-266. doi:10.1016/j.bbi.2019.02.010

Zhu J, Chen J, Zhang K. Clinical effect of flunarizine combined with duloxetine in the treatment of chronic migraine comorbidity of depression and anxiety disorder. *Brain Behav*. 2022;12(8):e2689. doi:10.1002/brb3.26

**Fig 1**. **HMGB1 infusion to mPFC causes specific anxiety-like behavior**
A) Schematic of Experimental Workflow  B) Open Arm Exploration C) Latency to Explore Open Arms D) Homecage Feeding Latency E) Female Change in Weight following Infusion Statistical analyses were performed using *2-way ANOVA to confirm primary effect of treatment and a Bonferroni post-hoc test is displayed on each graph* with the sexes grouped together. (*$p < 0.05$, **$p < 0.01$), n = 8-12 mice for each group Results are expressed as the mean ± SEM

***Fig. 2. HMGB1 infusion induces morphological changes in female microglia*** *A) Representative Images for IHC analysis (B-F) Microglial imaging analysis for male mice (B)Iba1 Volume Quantification (C) Tmem119 Volume Quantification (D) Tmem119:Iba1 Volume Ratio (E) Phagocytic Capacity Phagocytic capacity was calculated by normalizing total CD68 volume to the total volume of IBA1 and Tmem119 merged. (F) Male Microglia Have Significantly Altered Morphology following HMGB1 (G-K) Microglial imaging analysis for female mice. (G)Iba1 Volume Quantification (H) Tmem119 Volume Quantification (I) Tmem119:Iba1 Volume Ratio (J) Phagocytic Capacity Phagocytic capacity was calculated by normalizing total CD68 volume to the total volume of IBA1 and Tmem119 merged. (K) Female Microglia Have Significantly Altered Morphology following HMGB1 Statistical analyses were performed using* Mann-Whitney test *with the exception of Sholl Analysis. Statistical analysis for this measure was performed by comparing the Pearson r values of the aCSF condition and dsHMGB1 condition (\*p < 0.05, \*\*p < 0.01, n = 3-4 mice for each group, each mouse has an n=6-8 cells that were 3D reconstructed for analysis). Each dot represents an individual mouse. Results are expressed as the mean ± SEM.*

**Fig 3. Male and Female Display Differential Transcriptional Responses to HMGB1.** *16 hours following the final infusion of HMGB1, mice were sacrificed, and the frontal cortex was collected for cell isolation protocol as described in methods and materials section. CD11b+ cells are microglia and CD11b- cells are inclusive of neurons, astrocytes, and other glia (A) Male Expression of TLR4 (B) Female Expression of TLR4 (C) Male Expression of RAGE (D) Female Expression of RAGE (E) Male Expression of MyD88 (F) Female of Expression of MyD88. Statistical analyses were performed using Kruskal-Wallis Test (\*p < 0.05, \*\*p < 0.01, n = 4-5 mice for each group, each). Each dot represents an individual mouse. Results are expressed as the mean ± SEM*

*Fig 4. **Microglial MyD88 is necessary for HMGB1 response in females** (A) Schematic showing experimental outline (B)Male Time Spent in Open Arms of Elevated Plus Maze ,\*p < 0.05, \*\*p < 0.01 (C) Female Time Spent in Open Arms of Elevated Plus Maze Male (D) Male Latency to Feed in Novelty Suppressed Feeding Assay (E) Female Latency to Feed in Novelty Suppressed Feeding Assay ,\*p < 0.05, \*\*p < 0.01  N per group is 5-10, per treatment group. Statistical analyses were performed using 2-way ANOVA, Dunnett's post-hoc values displayed.*

***Supplemental Figure 1 HMGB1 does not impact OFT, Social Preference, or TST immobility** A) Center Exploration of the OFT is unchanged by HMGB1 B) Social Preference is not altered by HMGB1 C) TST Immobility is not significantly effected by HMGB1 Statistical analyses were performed using 2-way ANOVA to confirm primary effect of treatment and a Bonferroni post-hoc test is displayed on each graph (\*p < 0.05, \*\*p < 0.01, n = 8-12 mice for each group Results are expressed as the mean ± SEM.*

***Supplemental Figure 2 Observed Effects of HMGB1 in Females Do Not Significantly Correlate with Estrus Cycle***
*A) Weight Change Following HMGB1 does not correlate with Estrus cycle B) Open Arm Time in the EPM following HMGB1 does not correlate with Estrus cycle C) Home Cage Feeding Latency following HMGB1 does not correlate with Estrus cycle  Statistical analyses were performed using 2 way ANOVA test (\*p < 0.05, \*\*p < 0.01, n = 2-10 mice for each group. Results are expressed as the median.*

***Supplemental Figure 3. HMGB1 induced anxiety is robust in cohort utilized for microglia imaging experiments****(A) Open Arm Exploration for animals used for IHC imaging experiments (B) Latency to Explore Open Arms for animals used for IHC imaging experiments, n = 3-4 mice for each group. Each dot represents an individual mouse. Results are expressed as the mean ± SEM.*

**Supplemental Figure 4. No Changes in OFT, LDB, or TST were observed in response to dsHMGB1**
*A-C) Male Behavioral Data A) OFT Velocity is not changed by HMGB1 B) Light Dark Box (LDB) crossings are not impacted by HMGB1 C) TST Immobility remains constant following HMGB1 D-F) Female Behavioral Data D) OFT Velocity is not changed by HMGB1 E) Light Dark Box (LDB) crossings are not impacted by HMGB1 F) TST Immobility remains constant following HMGB1 \*p < 0.05, \*\*p < 0.01  N per group is 5-10, per treatment group. Statistical analyses were performed using 2-way ANOVA.*