The gatekeeper to gastric cancer; gastric microbiota invade the lamina propria in *Helicobacter pylori*-associated gastric carcinogenesis

Harriet J. Giddings[1], Ana Teodósio[2], Jack L. McMurray[2,3], Kelly Hunter[2], Zainab Abdawn[5], Jeffrey A. Cole[1], Claire D. Shannon-Lowe[4] and Amanda E. Rossiter-Pearson[1*].

Affiliations: [1]Institute of Microbiology and Infection, [2]Birmingham Tissue Analytics, [3] Rheumatology Research Group, [4]Institute of Immunology and Immunotherapy, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK.[5]Queen Elizabeth Hospital, University Hospital Birmingham NHS Foundation Trust, UK.

*Corresponding author: Amanda E. Rossiter-Pearson **Email:** rossiterae@gmail.com

Author Contributions: Conceptualisation: C.D.S-L. & A.E.R-P. Methodology: H.J.G, A.T., J.L.M, & A.E.R-P. Data analysis: H.J.G., A.T., J.L.M., K.H., Z.A. & A.E.R-P. Writing: H.J.G., J.A.C. & A.E.R-P. Funding acquisition: A.E.R-P.
Competing Interest Statement: The authors declare no competing interests.
Classification: Biological sciences; microbiology.
Keywords: *Helicobacter pylori,* gastric adenocarcinoma, microbiome.

Abstract

Stomach cancer is the fourth leading cause of cancer-related deaths worldwide. *Helicobacter pylori* is the main risk factor for gastric adenocarcinoma (GAC), yet the mechanism underpinning this association remains uncharacterised. Gastric intestinal metaplasia (GIM) represents the pre-cancerous stage and follows *H. pylori*-associated chronic gastritis (CG). Sequencing studies have revealed fewer *H. pylori* and more non-*H. pylori* bacteria in GAC. However, the spatial organisation of the gastric microbiota in health and disease is unknown. Here, we have combined RNA *in situ* hybridisation and immunohistochemistry to detect *H. pylori,* non-*H. pylori* bacteria and host cell markers (E-cadherin, Mucins 5AC and 2) from patients with CG (n=9), GIM (n=12), GAC and normal tissue adjacent to tumours (NATs) (n=3). Quantitative analysis of whole slide scans revealed significant correlations of *H. pylori* and other bacteria in CG and GIM samples. In contrast to sequencing studies, significantly fewer non-*H. pylori* bacteria were detected in *H. pylori*-negative patients. Importantly, whilst *H. pylori* exclusively colonised the gastric glands, non-*H. pylori* bacteria invaded the lamina propria in 3/4 CG and 5/6 GIM *H. pylori*-positive patients. Bacterial invasion was observed in 3/3 GAC samples and at higher levels than matched NATs. We propose that *H. pylori* 'holds the keys' to disrupt the gastric epithelial barrier, facilitating the opportunistic invasion of non-*H. pylori* bacteria to the lamina propria. Bacterial invasion could be a significant driver of inflammation in *H. pylori*-associated carcinogenesis. This proposed mechanism would both explain the synergistic roles of *H. pylori* and other bacteria and redirect attempts to prevent, diagnose and treat GAC.

Introduction

Gastric adenocarcinoma (GAC) is the fourth leading cause of cancer-related deaths, accounting for 7.7% of cancer mortalities worldwide (1). *Helicobacter pylori* infection is associated with approximately 70% of GAC cases and the global prevalence of *H. pylori* infection is 50% (1). Correa's cascade describes the histopathological changes to the gastric mucosa during progression to GAC (2). *H. pylori* first triggers chronic gastritis (CG), which can lead to gastric intestinal metaplasia (GIM), dysplasia and finally GAC. Although genotypes of *H. pylori* that encode the virulence factors CagA, VacA and HtrA (3) are more associated with severe disease outcomes, the mechanism by which a minority of *H. pylori* infections (1%) develop GAC remains unknown. In recent years, multiple 16S ribosomal RNA (16S rRNA) and metagenomic sequencing studies have profiled the human gastric microbiota during health and *H. pylori*-associated disease (4-7). The proposed model is that the healthy human stomach harbours a distinct microbial community structure and that upon *H. pylori* infection, this community structure shifts towards a dominance of *H. pylori* in CG. As carcinogenesis progresses, multiple other bacterial species displace *H. pylori* and dominate the GAC microbiota. However, sequencing studies are often prone to issues with contamination (8), do not offer spatial resolution and typically do not differentiate between transient and

persistent gastric bacteria (9). A longstanding question is whether the gastric microbiota play a causative or correlative role in GAC. Here, we provide high resolution spatial images of *H. pylori* and non-*H. pylori* bacteria in patients with *H. pylori*-positive or negative CG, GIM, GAC and normal tissue adjacent to tumour (NAT). Importantly, we provide direct evidence of non-*H. pylori* bacteria invading the lamina propria in all stages of Correa's cascade, suggesting that non-*H. pylori* bacteria might play a pivotal role in the early stages of *H. pylori*-associated carcinogenesis.

Results and Discussion
Localisation of *H. pylori* and the gastric microbiota in carcinogenesis Archived formalin-fixed paraffin-embedded gastric corpus tissue were retrieved from the Human Biomaterials Research Centre (n=21). The associated clinical pathology reports confirmed samples were *H. pylori*-positive CG (n=4) or GIM (n=6) and *H. pylori*-negative CG (n=5) or GIM (n=6). An independent pathologist scored all samples for inflammation (supporting dataset). Gastric corpus tissue was also obtained from surgical resections of GAC and NATs (n=3). To detect *H. pylori* or non-*H. pylori* bacteria, sections were stained using RNAScope *in situ* hybridisation (ISH) probes against *H. pylori*-specific or conserved sequences of the 16S rRNA gene, respectively. The probe 'Eubacteria' was used to detect non-*H. pylori* bacteria. Immunohistochemistry enabled detection of the host cell markers E-cadherin, Mucins 5AC (Muc5AC) and 2 (Muc2) using automated tissue staining for CG and GIM samples. Manual tissue staining was used for GAC and NAT. Goblet cells, present only in GIM, secrete Muc2, whereas Muc5AC is found in both healthy and diseased gastric mucus. Representative images and quantification of indicated markers are shown in Figure 1A-D and Figure 1E-H, respectively. Consistent with previous studies (10, 11), *H. pylori* exclusively colonised the gastric glands (Figure 1B). The presence of Eubacteria correlated with *H. pylori* infection in CG and GIM, but were significantly reduced in the absence of *H. pylori* infection (n=11) (Figure 1F). This contrasts with the current dogma that there is a unique microbiota in *H. pylori*-negative individuals (4). Comparable levels of Muc5AC were observed across the patient groups and, as expected, there was a significant increase in Muc2 in GIM patients (Figure 1 G&H). Eubacteria increases in GAC compared with matched NAT samples, whilst *H. pylori* increases in NAT compared to GAC (Figure 1E&F). Although this is not statistically significant, this is in agreement with sequencing studies that show a displacement of *H. pylori* by gastric microbiota in GAC tissue (5, 7).

***Helicobacter pylori* is associated with invasion of non-*H. pylori* bacteria to the lamina propria**
Bacterial invasion is associated with multiple diseases of the gastrointestinal tract, causing damage to the epithelial architecture by triggering inflammatory responses. Whilst *H. pylori* colonised the gastric glands, non-*H. pylori* bacteria invaded the lamina propria in 3/4 patients with *H. pylori*-associated CG and 5/6 *H. pylori*-positive patients with GIM (Figure 2A-C). In GAC, Eubacteria appear as dense regions of colonisation in one GAC patient (Figure 2F) or are more evenly distributed in the remaining two GAC tissue samples (Figure 2E). Eubacterial invasion increased progressively from CG to GIM and GAC compared to matched NATs (Figure 2H).

Next, bacterial invasion was confirmed at the cellular level using a modified Gram stain, which shows a mix of spindle shaped Gram-positive and Gram-negative bacteria in GAC (Figure 2F) and Gram-positive cocci in selected CG and GIM samples (data not shown). Invasive bacteria belonging to the Bacillota (Firmicute) phylum were detected in 3/5 CG *H. pylori*-positive patients. A representative image is shown in Figure 2G, where *H. pylori* is predominantly found to occupy glands, whereas Bacillotas and Eubacteria can be seen invading the lamina propria (Figure 2G). Further work is required to identify the precise identity of these invasive bacteria and the associated immune response. Invasion of *H. pylori* into the lamina propria is very rare across all patient samples, yet an instance of this can be seen in Figure 2C.

Recently, Sharafutdinov and colleagues reported that *H. pylori* strains encoding the trimeric form of HtrA, which proteolytically cleaves the cell junction proteins occludin, claudin-8 and E-cadherin, are associated with a higher GAC risk (3).
We propose that *H. pylori* acts as the 'gatekeeper' via HtrA-mediated cleavage of cell junction proteins of the gastric epithelial cell barrier (3), leading to infiltration of Eubacteria to the lamina propria (Figure 3). The prevalence of invasive bacteria in larger patient cohorts must be determined. Clinically, it should be

trialled whether antibiotic eradication of *H. pylori* clears invasive bacteria, particularly in patients with GIM where eradication does not significantly reduce GAC risk. The role of invasive bacteria should be explored for translation into the prevention, diagnosis and treatment of GAC.

Materials and Methods Archived human gastric biopsies were sectioned and stained as described in the extended methods. Images were acquired using a Vectra Polaris whole slide scanner and analysed on open source QuPath software (12).

Data and Materials Study data are included in the article and/or SI material.

Acknowledgments We thank the Royal Society and the University of Birmingham Centre Development fund from Cancer Research UK for funding (A.E.R.). We thank patients who have donated samples to HBRC. We also thank HBRC and Dr. Gary Reynolds for sample processing and sectioning.

References
1. H. Sung, J. Ferlay*, et al.*, Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*. 71(3), 209-49 (2021).
2. P. Correa, Human gastric carcinogenesis: a multistep and multifactorial process--First American Cancer Society Award Lecture on Cancer Epidemiology and Prevention. *Cancer Res*. 52(24), 6735-40 (1992).
3. I. Sharafutdinov, N. Tegtmeyer*, et al.*, A single-nucleotide polymorphism in *Helicobacter pylori* promotes gastric cancer development. *Cell Host Microbe*. 31(8), 1345-58 e6 (2023).
4. J. M. Noto, R. M. Peek, Jr., The gastric microbiome, its interaction with *Helicobacter pylori*, and its potential role in the progression to stomach cancer. *PLoS Pathog*. 13(10), e1006573 (2017).
5. R. M. Ferreira, J. Pereira-Marques*, et al.*, Gastric microbial community profiling reveals a dysbiotic cancer-associated microbiota. *Gut*. 67(2), 226-36 (2018).
6. S. Delgado, R. Cabrera-Rubio*, et al.*, Microbiological survey of the human gastric ecosystem using culturing and pyrosequencing methods. *Microb Ecol*. 65(3), 763-72 (2013).
7. Y. Li, Y. Ouyang*, et al.*, Research trends on the relationship between *Helicobacter pylori* and microbiota: A bibliometric analysis. *Helicobacter*. 28(6), e13021 (2023).
8. S. J. Salter, M. J. Cox*, et al.*, Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC Biol*. 12, 87 (2014).
9. M. R. Spiegelhauer, J. Kupcinskas*, et al.*, Transient and Persistent Gastric Microbiome: Adherence of Bacteria in Gastric Cancer and Dyspeptic Patient Biopsies after Washing. *J Clin Med*. 9(6), (2020).
10. C. Fung, S. Tan*, et al.*, High-resolution mapping reveals that microniches in the gastric glands control *Helicobacter pylori* colonization of the stomach. *PLoS Biol*. 17(5), e3000231 (2019).
11. D. Keilberg, Y. Zavros*, et al.*, Spatial and Temporal Shifts in Bacterial Biogeography and Gland Occupation during the Development of a Chronic Infection. *mBio*. 7(5), (2016).
12. P. Bankhead, M. B. Loughrey*, et al.*, QuPath: Open source software for digital pathology image analysis. *Sci Rep*. 7(1), 16878 (2017).

Figure 1. Quantification and spatial localisation of *H. pylori* and non-*H. pylori* bacteria in CG, GIM, NAT and GAC revealed by RNAScope *in situ* hybridisation (ISH) and immunohistochemistry (IHC). Whole slide scans of stained patient tissue sections were obtained using a Vectra whole slide scanner. Images were spectrally unmixed, viewed and quantified using QuPath. **A & C)** A representative whole slide scan of a GIM *H. pylori*-positive (A) or *H. pylori*-negative (C) section showing RNAScope ISH probes 'H. pylori' and 'Eubacteria' to detect *H. pylori* (green) and non-*H. pylori* bacteria (white), respectively. IHC staining against E-cadherin (red), Muc5AC (yellow) and Muc2 (turquoise) are also shown. **B & D)** A higher magnification of panel A and C, respectively. The coloured arrows show the indicated markers. **E-H)** Quantification of target markers as percentage coverage of tissue from whole slide scans for each patient group for *H. pylori* (E), Eubacteria (F) Muc5AC (G) and Muc2 (H). Lines between NAT and GAC data points show matched patient pairs. Horizontal bars indicate the median for each indicated patient group. A Mann-Whitney test was used to determine significance where * indicates *p*<0.05 and ** indicates *p*<0.01.

Figure 2. Non-*H. pylori* bacteria invade the gastric lamina propria in CG, GIM and GAC. Whole slide scans were obtained using a Vectra whole slide scanner. Images were spectrally unmixed and viewed using QuPath. **A-F)** Representative images showing Eubacterial (white) invasion in patients with CG (A-B), GIM (C), *H. pylori* negative CG (D) and GAC (E&F) (Inset in Panel F shows bacterial cells in a modified Gram stain). E-cadherin (red), *H. pylori* (green), Eubacteria (white) and DAPI (blue) are shown for visualisation purposes. Arrows indicate the target markers. Eubacterial invasion is highlighted in white ovals for CG and GIM. A rare occurrence of *H. pylori* invasion to the lamina propria is indicated by the green arrow in (C). **G)** Representative image showing Eubacterial (white) and Bacillota (yellow) invasion, and *H. pylori* (green) in a patient with CG. Bacillotas (Firmicutes) were detected using an RNAScope probe against target 16S rRNA sequences. **H)** Qualitative FISH scoring of Eubacterial invasion for each patient group; 0 = no invasion, 1 = sparse, 2 = moderate, 3 = high and 4 = colonisation. Horizontal bars indicate the mean for each indicated patient group. A Mann-Whitney test was used to determine significance where * indicates *p*<0.05. Lines between NAT and GAC data points show matched patient pairs.

Figure 3. Illustrative model proposing *H. pylori* as the 'gatekeeper' for bacterial invasion to the gastric lamina propria. Stages 1-5 are (1) Eubacteria and *H. pylori* in the gastric surface and gland, respectively. (2) *H. pylori* secretes HtrA, which cleaves tight junction protein E-cadherin. Stages 3-5 are as described.

Supplementary material

Extended methods

Sample preparation and pre-treatment protocol for automated tissue staining

Punch biopsy tissue samples were collected from consenting patients at the Queen Elizabeth Hospital, Birmingham fixed in formalin and prepared at the Human Biomaterial Resource Centre (HBRC), University of Birmingham by embedding in paraffin (Ethics #17-285). 4 µM thick tissue sections were then prepared for RNAScope, immunohistochemistry and H&E staining, using a Leica BOND RX Fully Automated Research Stainer. Prior to staining, sections were prepared in the BOND RX, following three short protocols: 1) Deparaffinisation, rehydration, hydrogen peroxide and distilled water wash 2) Heating to 100°C in target retrieval buffer ER2 (pH 9; AR9640) for 45 min and manual washing with distilled water followed by 100% ethanol and drying at 60°C. 3) Incubation in protease III solution for 30 min, followed by a final wash in distilled water.

Optimisation of RNAScope treatments and immunohistochemistry antibody concentrations for 5-plex automated staining

Automated RNAScope C1 and C2 probes against *H. pylori* (ACD-Bio, #542938) and Eubacteria (ACD-Bio; #464468), respectively, were first tested on healthy colon tissue in a fluorescent multiplex assay, using an RNAScope LS Multiplex Fluorescent Reagent Kit (ACD Bio; 322800), following the standard protocol recommended for this platform. The 50x C2 probe was diluted 1:50 in the C1 probe, according to manufacturer's instructions. Pre-treatment of tissue slides for automated tissue staining with lysozyme was not used as this was found to affect tissue integrity. However, automated tissue staining with lysozyme (micro bacteria detection protocol provided by ACD) and a 3-plex panel (E-Cadherin and RNAScope probes against Eubacteria and *H. pylori*) was used on all samples (n=21) to ensure bacterial signal was not underestimated in comparison to 5-plex stained images.

A positive and negative control probe section was used on every RNAScope run to validate and assess its quality and the sensitivity of the assay. The bacterial gene DapB (ACD Bio; 312038) was used as negative control to confirm the absence of background noise and a cocktail of housekeeping genes Polr2A C1, PPIB C2 and UBC C3 (ACD Bio; 320868) were used as positive control to validate the detection of the signal and the tissue integrity. *H. pylori* gene sequences can bind to both 16S rRNA probes against *H. pylori* and Eubacteria, whilst non-*H. pylori* bacteria stain with only the Eubacteria probe.

All antibodies used in IHC steps were optimised prior, using chromogenic DAB staining of healthy colon tissue in addition to a Leica Bond Polymer Refine Detection kit (Leica; DS9800). Antigen retrieval was tested using pH6 (Leica Bond TM Epitope Retrieval 1; AR9961) and pH9 (Leica Bond TM Epitope Retrieval

2; AR9640) buffers by heating to 100°C for 20 min. Three different dilutions were tested for each antibody, as recommended by manufacturer. Ideal staining pattern and intensity was assessed and approved by a pathologist, whereby slides were then used as reference throughout the validation process. All antibodies were then tested for compliance with the RNAScope pre-treatment, to ensure stability after protease III digestion. Once each antibody was assigned OPAL fluorophore single fluorescence assays were directly compared against DAB-stained colon tissue to optimise OPAL concentration. To assess epitope stability following heat steps and to define the order of the antibodies in the multiplex sequence, each antibody was tested individually in the different positions of the panel.

Automated 5-plex co-RNAScope *in situ* hybridisation/immunohistochemistry protocol

The automated RNAScope Multiplex Fluorescent LS assay (ACD Bio, 322800) was conducted according to manufacturer's instructions; all incubation periods were performed in a 40°C humidified environment. C1 and C2 probes against *H. pylori* and Eubacteria, respectively, were added to sections for 2 h. Next, a series of amplification steps were conducted by sequentially adding AMP1, AMP2 and AMP3 for 30 min, 30 min and 15 min respectively. Sections were then washed and HRP C1 was added for 15 min, followed by the corresponding, pre-diluted OPAL TSA 520 fluorophore assigned to the C1 probe. Sections were washed again and HRP blocker was added for 15 min. Next, the above steps are repeated whereby sections were washed and HRP C2 was added for 15 min, followed by the corresponding, pre-diluted OPAL TSA 620 fluorophore assigned to the C2 probe. Sections were washed again and HRP blocker was added for 15 min. Sections were further washed and the IHC protocol was immediately followed in the Leica BOND RX. Sections were washed, prior to heating at 100°C in ER2 (pH 9; AR9640) antigen retrieval solution for 10 min, followed by a further wash and blocking in peroxide for 10 min at room temperature. Next, pre-diluted primary antibody against E-Cadherin was added for 30 min at room temperature. Sections were washed, prior to the addition of OPAL polymer system antibody for 10 min. Sections were washed, and pre-diluted OPAL TSA 690 fluorophore was added to fluorescently label E-cadherin. Between each staining cycle, a heat induced stripping step with pH6 solution was added. The staining steps were repeated sequentially for MUC5AC and MUC2, using OPAL TSA 570 and 480 fluorophores, respectively.

Table 1. Automated 5-plex RNAScope probe and IHC antibody information

Marker	Marker type	Supplier	Clone	Cat. No.	Marker dilution	Fluorophore	Fluorophore cat. No.	Fluorophore dilution
H-pylori	RNAScope C1	ACD	N/A	N/A	RTU	520	FP1487001KT	1:150
Eubacteria	RNAScope C2	ACD	N/A	N/A	1:50	620	FP1495001KT	1:150

E-cadherin	Antibody	Cell Signalling Technology	24E10	3195S	1:250	690	FP1497001KT	1:200
MUC2	Antibody	Novus Biologicals	Polyclonal	NBP1-31231	1:1000	480	FP1500001KT	1:150
MUC5AC	Antibody	Invitrogen	2-25LE	MA119346	1:1000	570	FP1488001KT	1:150

Table 2. Exposure times for automated 5-plex RNAScope ISH/IHC assay

Filter	Exposure times (ms)
DAPI	1.13
480	2.47
520	36.06
570	5.61
620	29.58
690	7.46

Manual 3-plex sample preparation and pre-treatment protocol

Gastrectomy tissue samples were collected from consenting patients at the Queen Elizabeth Hospital, Birmingham, fixed in formalin and prepared at the Human Biomaterial Resource Centre (HBRC), University of Birmingham by embedding in paraffin (IRAS ethics #276525 and 20/NW/0001). 4 µM thick tissue sections were then prepared for staining at the Institute of Cancer and Genomic Sciences, University of Birmingham. Prior to staining, sections were oven-dried at 60°C for 1 h, followed by deparaffinisation in three 10-minute changes in xylene. Sections were then immediately rinsed twice in 100% ethanol and air-dried for 10 min at 60°C. Hydrogen peroxide was added to sections for 10 min at room temperature, before washing twice in distilled water. RNAScope target retrieval buffer was heated to ~98°C in Simport Easy Dip staining jars (Simport; M900-12W) in a microwave oven at 100% power. Once ~98°C was reached, sections were placed inside the jars and heated for a further 20 min at 20% power, with pauses every 2-3 min to check temperature and avoid boiling of sections. Sections were washed twice in distilled water, followed by dipping in 100% ethanol and drying at 60°C. Protease III solution was then added to sections for 20 min, followed by washing in distilled water.

Optimisation of RNAScope treatments and IHC antibody concentrations for manual 3-plex staining

Manual C1 (ACD Bio; 542931), C2 (ACD Bio; 464461-C2) and C3 (ACD Bio; 1112731-C3) probes were assigned to *H. pylori,* Eubacteria and Bacillota, respectively and prepared according to manufacturer's

instructions (the 50x C2 and/or C3 probe was diluted 1:50 in the C1 probe). The Bacillota probe was custom-designed by ACD Bio, according to strict criteria, including no cross-reactivity to the *H. pylori* 16S rRNA gene. Slides stained with the Bacillota probe were pre-treated with Lysozyme (2 mg/mL). Probes were tested alone and in combination with the micro bacteria detection protocol provided by ACD, including addition of lysozyme, using *in vitro* infection models. All antibodies used in IHC steps were optimised prior by the automated staining protocol and were further tested manually *in vitro* by staining healthy human gastric organoid models. Antigen retrieval pH and method was optimised in a water bath or microwave oven at ~98°C for 20-45 min to heat sections. pH6 (Perkin Elmer) and pH9 (Perkin Elmer) buffers were also compared during each heating method. To assess epitope stability following heat steps and to define the order of the antibodies in the manual IHC staining sequence, antibody staining was tested individually in the different positions of the panel.

Manual co-RNAScope ISH/IHC protocol

The RNAScope multiplex fluorescent v2 assay (ACD Bio; 323110) was conducted according to manufacturer's instructions; all incubation periods were at 40°C in a humidified incubator. Manual C1 and C2 probes against *H. pylori* and Eubacteria, respectively, were added to sections for 2 h. For Bacillota staining, the manual C3 probe against Bacillota was also included. Next, a series of amplification steps was carried out by sequentially adding AMP1, AMP2 and AMP3 for 30 min, 30 min and 15 min respectively. Sections were then washed and HRP C1 was then added for 15 min, followed by the corresponding, pre-diluted TSA vivid 520 fluorophore assigned to the C1 probe. Sections were washed again and HRP blocker was added for 15 min. Next, the above steps were repeated whereby sections were washed and HRP C2 was added for 15 min, followed by the corresponding, pre-diluted TSA vivid 570 fluorophore assigned to the C2 probe. For Bacillota staining, the above steps were repeated, whereby sections were washed and HRP C3 was added for 15 min, followed by the corresponding, pre-diluted OPAL 690 fluorophore assigned to the C3 probe. Sections were then washed again and HRP blocker was added for 15 min. Sections were then further washed in and blocked for 45 min in blocking buffer containing 0.3% Triton-X, 3% BSA, 0.3% Goat Serum and 1X PBS. Sections were then washed, and E-cadherin primary antibody was added to sections over night at 4°C. After 16-20 h, sections were washed and then incubated with secondary antibody Alexa Fluor 594, diluted 1:1000 in blocking buffer, for 1 h at room temperature. Exposure times varied between samples and were determined at the time of scanning.

Table 3. Manual RNAScope probe and IHC antibody information

Marker	Marker type	Supplier	Clone	Cat. No.	Marker dilution	Fluorophore	Fluorophore cat. no.	Fluorophore dilution
H-pylori	RNAScope C1	ACD	N/A	542931	RTU	TSA vivid 520	7526	1:1500

Eubacteria	RNAScope C2	ACD	N/A	464461-C2	1:50	TSA vivid 570	7523	1:1500
Bacillota	RNAScope C3	ACD	N/A	1112731-C3	1:50	OPAL TSA 690	FP1497001KT	1:250
E-cadherin	Antibody	Cell Signalling Technology	24E10	3195S	1:400	Alexa Fluor 594	A20185	1:1000

Manual 3-plex IHC protocol

Manual IHC staining was conducted using the Opal 7-colour manual IHC Kit (Perkin Elmer; NEL811001T), according to manufacturer's instructions. Following step 5, consecutive sections were washed prior to heating at ~98°C in Simport staining jars for 20 min in a microwave oven in pH 6 AR6 buffer (Perkin Elmer, AR600250ML). Sections were then washed twice in TBST and blocked with peroxide (Perkin Elmer ARD1001EA) for 10 min at room temperature. Next, primary antibody against E-Cadherin was added for 30 min at room temperature. Sections were then washed, prior to the addition of OPAL polymer HRP Ms+Rb secondary antibody for 10 min at room temperature. Sections were washed again, and pre-diluted OPAL 690 polymer was added to fluorescently label E-cadherin for 30 min at 40°C. Sections were further washed and the above was repeated twice more, in order to microwave heat-strip the previous antibody epitope, allowing for subsequent application of primary antibodies against MUC5AC and MUC2, respectively, which were then assigned OPAL fluorophores 620 and 480. Exposure times varied between samples and were determined at the time of scanning.

Table 4. Manual 3-plex IHC antibody information

Marker	Marker type	Supplier	Clone	Cat. no.	Marker dilution	Fluorophore	Fluorophore cat. no.	Fluorophore dilution
E-cadherin	Antibody	Cell Signalling Technology	24E10	3195S	1:400	OPAL TSA 690	FP1497001KT	1:100
MUC2	Antibody	Novus Biologicals	Polyclonal	NBP1-31231	1:1000	OPAL TSA 480	FP1500001KT	1:100
MUC5AC	Antibody	Invitrogen	2-25LE	MA119346	1:500	OPAL TSA 620	FP1495001KT	1:100

Counterstaining and section visualisation

All sections were counterstained with spectral DAPI (Akoya Biosciences) for 10 min, washed in distilled water and mounted with ProLong Diamond Antifade Mountant and 1.5 mm thickness glass coverslips. All mounted sections were stored in the dark at 4°C until viewing. During optimization steps, sections were visualised using a Zeiss LSM 900 confocal microscope or Mantra 2 Quantitative Pathology Digital Workstation (Akoya Biosciences) and all final images were obtained on a Vectra Polaris Multispectral

Whole Slide Scanner (Akoya Biosciences) and saved as a .qptiff file. Images were viewed at 40x magnification using Phenochart Whole Slide Viewer with PhenoImager HT (Akoya Biosciences) or QuPath software.

Spectral unmixing

 .qptiff image files were spectrally unmixed by importing and stamping in Phenochart Whole Slide Viewer (Akoya Biosciences), unmixed and exported in InForm software (Akoya Biosciences) and finally restitched as a BIGTIFF file using Visiopharm software (Visiopharm, Hørsholm, Denmark).

Qualitative image analysis

All images were viewed using Phenochart Whole Slide Viewer (Akoya Biosciences) or open source QuPath software (Version 0.4.3). The presence of each marker of interest (*H. pylori*, Eubacteria, E-Cadherin, MUC5AC and MUC2) were scored independently (H.J.G. and A.E.R-P.) whereby 1 = no/low staining intensity, 2 = medium staining intensity, 3 = high staining intensity. Invasion of Eubacteria was also qualitatively scored, whereby 0 = no invasion, 1 = sparse invasion, 2 = moderate invasion (patches of bacteria across sample), 3 = high invasion (multiple clear regions of bacterial invasion across sample) and 4 = dense colonisation of bacteria (large regions of sample colonised).

Quantitative image analysis

Manual image analysis was conducted using the free and open source Qupath software (Version 0.4.3). For each whole-slide scan, tissue regions were first annotated and defined as a region of interest (ROI). Tissue detection was then performed based on the average values of all channels using three separately created pixel thresholders. Each pixel thresholder was used to calculate tissue area (μM^2), which was exported, and mean tissue area value was obtained. Three separate pixel thresholders were created for each individual OPAL channel, corresponding to - *H. pylori*, Eubacteria, Bacillota, MUC5AC or MUC2. The thresholders were saved, and average area annotation measurements (μM^2) were obtained for each channel. To account for *H. pylori* double-staining and detection in both the *H. pylori* and Eubacteria channels, absolute *H. pylori* area was calculated by calculating total Eubacteria channel area minus combined *H. pylori* and Eubacteria channel areas. For each whole slide scan, average percentage area coverage of each marker of interest was calculated by dividing individual channel area over total tissue area x 100. Data were exported from Excel to GraphPad Prism 9 (Version 9.5.1). For statistical analysis, a Mann-Whitney test was performed, whereby * indicates $p<0.05$, ** indicates $p<0.01$ and ns indicates non-significant. Data are presented as median ± SEM, for each group.

H&E staining and assessment of inflammation

Additional sections were obtained and stained with H&E for further inflammation analysis and scoring by an independent pathologist.

Modified Gram Stain

Prior to staining, all sections underwent deparaffinisation and rehydration, with a series of incubations at room temperature in xylene followed by two 2-minute washes in 100% ethanol. A modified Gram stain was followed for staining of bacteria in tissue sections (1).

References

1) Taylor RD. Modification of the Brown and Brenn gram stain for the differential staining of Gram-positive and Gram-negative bacteria in tissue sections. *Am J Clin Pathol.* 1966;46(4):472-474. doi:10.1093/ajcp/46.4.472

Supplementary Table 1. Patient demographics, clinical inflammatory scores and raw values of target markers as % coverage of whole slide scan

Patient	Patient group	Gender	Diagnosis	PPI use	H&E Inflammation grade	Invasion Score (0-4)	*H. pylori* coverage (%)	Eubacteria coverage (%)	MUC5AC coverage (%)	MUC2 coverage (%)
1	CG H. pylori positive	Male	Mild active *H. pylori* associated gastritis	None	Moderate	3	0.236	0.218	2.793	0.377
2	CG H. pylori positive	Male	Active chronic gastritis	None	Mild	1	0.139	0.234	25.531	0.107
3	CG H. pylori positive	Male	Mild active *H. pylori* associated gastritis	None	Moderate	3	0.339	0.303	25.238	0.123
4	CG H. pylori positive	Male	Mild active chronic gastritis	Lansoprazole	Moderate	2	0.125	0.256	27.231	0.578
5	CG H. pylori negative	Male	Mild reactive gastropathy	None	Mild	1	0.003	0.003	17.926	0.834
6	CG H. pylori negative	Male	Mild chronic gastritis	None	Mild	0	0.006	0.005	10.44	0.0176
7	CG H. pylori negative	Male	Mild non-specific chronic inflammation	None	Mild	1	0.008	0.001	10.51	0.018
8	CG H. pylori negative	Male	Mild chronic inactive gastritis	Lansoprazole	Moderate	1	0.009	0.004	18.167	0.992
9	CG H. pylori negative	Male	Corrobative of inflamed Barretts mucosa	Lansoprazole	Moderate	0	0.004	0.007	7.662	0.534
10	GIM H. pylori positive	Male	Severe active chronic infection with gastritis/atrophy	None	Severe	2	0.112	0.128	7.856	10.028
11	GIM H. pylori positive	Male	Chronic active gastritis and intestinal metaplasia	None	Severe	3	0.005	0.217	57.06	4.321
12	GIM H. pylori positive	Male	Mild active *H. pylori* associated gastritis and intestinal metaplasi	None	Patchy severe	3	0.401	0.472	3.266	19.861
13	GIM H. pylori positive	Male	*H. pylori* associated gastritis and intestinal metaplasia	None	Moderate	1	0.85	0.003	28.602	6.103
14	GIM H. pylori positive	Male	Intestinal metaplasia with squamous and columnar mucosa	Lansoprazole	Moderate	2	0.332	0.214	29.725	0.419
15	GIM H. pylori positive	Male	*H. pylori* associated gastritis and intestinal metaplasia	None	Mild	3	0.321	0.685	29.032	6.36
16	GIM H. pylori negative	Male	Chronic gastritis and intestinal metaplasia	None	Moderate	0	0.003	0.075	39.776	11.663
17	GIM H. pylori negative	Male	N/C	None	Mild	0	0.004	0.082	8.84	6.16
18	GIM H. pylori negative	Female	Mild active gastritis and focal intestinal metaplasia	None	*Missing	1	0.008	0.01	9.479	24.095
19	GIM H. pylori negative	Female	Active chronic gastritis and intestinal metaplasia	None	*Missing	1	0.036	0.07	31.402	2.499
21	GIM H. pylori negative	Male	Focal active chronic gastritis and intestinal metaplasia/atrophy	None	Moderate	1	0.003	0.073	18.264	1.698
22	GIM H. pylori negative	Male	Mild chronic gastritis and intestinal metaplasia	Rabeprazole	Mild	2	0.072	0.138	26.334	8.198
23	GAC	Male	Gastric adenocarcinoma	N/A	N/A	4	0.22	4.163	12.525	9.595
23	NAT	Male	Healthy tissue adjacent to tumour	N/A	N/A	3	0.084	1.853	8.49	4.084
24	GAC	Male	Gastric adenocarcinoma	N/A	N/A	3	0.015	1.192	5.839	13.749
24	NAT	Male	Healthy tissue adjacent to tumour	N/A	N/A	0	0.028	0.175	26.362	14.853
25	GAC	Male	Gastric adenocarcinoma	N/A	N/A	3	0.021	1.122	11.587	4.441
25	NAT	Male	Healthy tissue adjacent to tumour	N/A	NA	1	0.114	1.58	13.18	2.553