How distributed subcortical integration of reward and threat may inform subsequent

approach-avoidance decisions

Abbreviated title: Reward-threat integration in approach-avoidance

Anneloes M. Hulsman[a,b], Felix H. Klaassen[a,b], Lycia D. de Voogd[a,b,c],

Karin Roelofs[a,b] & Floris Klumpers[a,b]

[a]Radboud University, Donders Centre for Cognitive Neuroimaging, Nijmegen, The Netherlands

[b]Radboud University, Behavioural Science Institute, Nijmegen, The Netherlands

[c]Leiden University, Institute of Psychology and Leiden Institute for Brain and Cognition, Leiden, The

Netherlands

Correspondence: anneloes.hulsman@ru.nl (AMH), floris.klumpers@ru.nl (FK)

Number of pages: 32

Number of figures: 8

Number of tables: 0

Number of words:

-   Abstract: 250

-   Introduction: 650

-   Discussion: 1235

This work was funded by the Netherlands Organization for Scientific Research (NWO)

Research Talent Grant #406-18-540 awarded to AMH and a European Research Council

(ERC) Consolidator Grant #ERC_CoG-2017_772337 awarded to KR.

ABSTRACT

Healthy and successful living involves carefully navigating rewarding and threatening

situations by balancing approach and avoidance behaviours. Excessive avoidance to evade

potential threats often leads to forfeiting potential rewards. However, little is known about how

reward and threat information is integrated neurally to inform approach or avoidance decisions.

In this preregistered study, participants ($N_{behaviour}$=31, 17F; N_{MRI}=28, 15F) made approach-

avoidance decisions under varying reward (monetary gains) and threat (electrical stimulations)

prospects during functional MRI scanning. In contrast to theorized *parallel subcortical*

processing of reward and threat information before cortical integration, Bayesian Multivariate

Multilevel analyses revealed subcortical reward and threat *integration* prior to indicating

approach-avoidance decisions. This integration occurred in the ventral striatum, thalamus, and

bed nucleus of the stria terminalis (BNST). When reward was low, risk-diminishing avoidance

decisions dominated, which was linked to more positive tracking of threat magnitude prior to

indicating avoidance than approach decisions across these regions. In contrast, the amygdala

exhibited dual sensitivity to reward and threat. While anticipating outcomes of risky approach

decisions, we observed positive tracking of threat magnitude within the salience network

(dorsal anterior cingulate cortex, thalamus, periaqueductal gray, BNST). Conversely, after risk-

diminishing avoidance, characterized by reduced reward prospects, we observed more

negative tracking of reward magnitude in the ventromedial prefrontal cortex and ventral

striatum. These findings shed light on the temporal dynamics of approach-avoidance decision-

making. Importantly, they demonstrate the role of subcortical integration of reward and threat

information in balancing approach and avoidance, challenging theories positing predominantly

separate subcortical processing prior to cortical integration.

INTRODUCTION

Healthy and successful living involves carefully navigating rewarding and threatening situations by adopting an appropriate balance between approach and avoidance behaviours. Disruption of this balance can result in severe consequences, as observed in anxiety-related disorders and addiction (American Psychiatric Association, 2013). In approach-avoidance decision-making, the weighing of potential reward and threat prospects is pivotal (Corr, 2013). However, little is known about how this weighing is neurally represented to inform approach or avoidance decisions. In this study, we leveraged Bayesian Multivariate Multilevel (BMM) analyses on trial-by-trial BOLD fMRI data to investigate how distributed processing and integration of reward and threat information in the human brain contribute to approach-avoidance decision-making.

Human neuroimaging studies and recent theoretical frameworks suggest that reward and threat information is processed in distinct *parallel* subcortical streams, followed by reward-threat *integration* in cortical regions (Yacubian et al., 2006; Aupperle and Paulus, 2010; Basten et al., 2010; Park et al., 2011; Schlund et al., 2016; Zorowitz et al., 2019; Azab and Hayden, 2020; Livermore et al., 2021). For example, the ventral striatum (vStriatum) plays a crucial role in processing positive value and rewards (Talmi et al., 2009; Aupperle and Paulus, 2010; Basten et al., 2010; Bartra et al., 2013), potentially driving approach behaviour (Alvarez & Ruarte, 2001; Beaufour et al., 2001; Pedersen et al., 2021). Conversely, the amygdala and bed nucleus of the stria terminalis (BNST) are widely recognized for their pivotal roles in threat processing (Shackman and Fox, 2016; Abivardi et al., 2020; Hur et al., 2020; Hulsman et al., 2021c) and facilitation of avoidance behaviour (Ch'ng et al., 2018; Duque-Wilckens et al., 2018; Giardino et al., 2018; Miller et al., 2019; Hu et al., 2020). Value signals, indicating reward and threat potential, subsequently need to be integrated to inform decision-making, as previously demonstrated in cortical regions including the ventromedial prefrontal cortex (vmPFC) and dorsal anterior cingulate cortex (dACC) (Basten et al., 2010; Park et al., 2011; Schlund et al., 2016; Zorowitz et al., 2019; Azab and Hayden, 2020).

However, preclinical studies have demonstrated that reward and threat processing may

be *intertwined* at the neuronal level in subcortical regions, with processing of opposite valence

occurring in neighbouring cells, e.g., within amygdala subregions (Gentry et al., 2016; Burgos-

Robles et al., 2017; Hamel et al., 2017; Beyeler et al., 2018). Moreover, recent rodent studies

have shown that subcortical regions, including the central nucleus of the amygdala and nucleus

accumbens can trigger both approach and avoidance motivation depending on the context and

input received from other structures (Warlow et al., 2020; Zhou et al., 2022) and that regions

of the thalamus respond particularly to situations requiring integration, such as approach-

avoidance conflict (Engelke et al., 2021). This aligns with recent human work showing that

regions of the salience network (thalamus, periaqueductal gray, dACC, and anterior insula)

are sensitive to both rewards and threats (Aupperle et al., 2015; Schlund et al., 2016;

Moughrabi et al., 2022). Strikingly, however, none of the aforementioned human studies

investigated how the interplay between reward and threat is represented across these brain

regions and how this balance shifts upon decisions to approach or avoid.

To investigate this, participants performed a Fearful Avoidance Task (Hulsman et al.,

2021a) in the MRI scanner. In this task, participants made approach-avoidance decisions

under varying prospects of reward (monetary gain) and threat (electrical shock). Based on

previous human literature, we hypothesized that prior to approach, activations would occur in

brain regions most strongly associated with appetitive processing (vStriatum, vmPFC).

Conversely, prior to avoidance, we expected activations in brain regions traditionally linked to

defensive responding (amygdala, BNST). Additionally, we expected that the vmPFC and dACC

would play a crucial role in reward-threat integration. Finally, following risky approach

decisions, leading to an increased chance of receiving shocks or rewards, we expected

increased activity in regions associated with appetitive (vStriatum, vmPFC), defensive

(amygdala, BNST), and salience processing (thalamus, periaqueductal gray, dACC, anterior

insula) compared to risk-diminishing avoidance.

MATERIALS AND METHODS

Research questions, hypotheses, and planned analyses were preregistered at the Open Science Framework: https://osf.io/7sm9k (Hulsman et al., 2021b) prior to data analyses. All data and code will be available at the Donders Institute for Brain Cognition and Behaviour repository upon publication: https://doi.org/10.34973/gxr8-9f24. Supplementary information is available at the Open Science Framework: https://osf.io/hbjp5.

Participants

The recruited sample consisted of 31 participants (17 females, M_{age} = 23.45, SD_{age} = 3.33). Participants had an average trait anxiety score of M_{STAI-T} = 38.03 (SD = 8.39, range = 27-55; one participant did not complete the State-Trait Anxiety Inventory (STAI; Spielberger et al., 1970). Three participants were a priori excluded from the MRI analyses due to a lack of avoidance behaviour (<10%, see also Hulsman et al., 2024). As a result, planned contrasts of approach vs. avoid trials could not be carried out in these participants. Exclusion of these participants led to a final sample size of N=28 (15 females, M_{age} = 23.25, SD_{age} = 3.27). Participants self-reported meeting the following inclusion criteria: between 18 and 35 years of age, normal/corrected-to-normal vision, sufficient comprehension of English or Dutch, no psychiatric disorders, no neurological or cardiological disease/treatment, no use of psychotropic medication, no current drug or alcohol abuse, and no epilepsy. Participants received EUR 10 for participation and an additional bonus between EUR 0-5 depending on their performance in the Fearful Avoidance Task (described below). All participants were MRI eligible and provided written informed consent. This study was carried out in compliance with the declaration of Helsinki and approved by the local ethics committee (CMO Arnhem-Nijmegen).

Procedure

Participants were informed that they had the opportunity to earn a monetary bonus based on their behaviour in the Fearful Avoidance Task (FAT, Hulsman et al., 2021a). They were asked

to write down 10 random numbers, which would later be linked to specific trial numbers using a mathematical formula that was disclosed to the participants only after the experiment. If they received a monetary reward during these trials, they would receive the corresponding sum as a bonus, capped at a maximum of EUR 5.

Next, participants underwent titration procedures to determine their individual reward and threat levels for the Fearful Avoidance Task. Initially, to establish shock intensity at a point of maximum discomfort without causing pain, participants underwent a standardized *shock work-up procedure* consisting of five shock administrations (Klumpers et al., 2010). Each shock was rated on a scale ranging from 1 (not painful/annoying) to 5 (very painful/annoying). Intensities were adjusted gradually to attain a subjective level that as maximally uncomfortable without being painful (aiming for a 4 on the 5-point scale). Electric shocks were delivered via two Ag/AgCl electrodes attached to the distal phalanges of the middle and ring finger of the right hand using a MAXTENS 2000 (Bio-Protech) device. Shock duration was 250ms at 150Hz, and intensity varied in 10 intensity steps between 0-80mA. We opted to use electric shocks in this paradigm because primary reinforcers may evoke stronger psychophysiological responses than for example affective pictures (Lissek et al., 2007; Delgado et al., 2011).

Next, we conducted a *reward-threat titration procedure* to determine the monetary reward required for individuals to accept the risk of receiving an electric shock of the predetermined shock work-up intensity. Amounts between EUR 0.20 and EUR 10 were presented in semi-random order. For each amount, participants were asked whether they would be willing to risk receiving electrical stimulation. No actual shock reinforcement was administered throughout this entire procedure. Participant's indifference point ($M = 1.17$, $SD = 1.84$) was subsequently used to calculate the reward values for the ensuing FAT (described below).

Finally, the Fearful Avoidance Task (FAT, Hulsman et al., 2021a) was employed to evaluate active approach-avoidance behaviour under competing reward (monetary gains) and threat (electric shocks) prospects. In this study, participants performed an adapted version of the FAT (see next section), optimized for use in the MRI scanner.

Fearful Avoidance Task (FAT)

Participants received on-screen instructions for the task in the MRI scanner. Each trial consisted of a cascade of three phases (see Figure 1A). In the *decision phase* participants were confronted with a combination of reward and threat. The reward level was a monetary amount displayed in numbers below the avatar and was relative to the individual's indifference point (IP) either low (IP - 40-50%), medium (IP ± 5%), or high (IP + 40-50%). The threat levels were displayed with different avatars (Figure 1B). In total, there were three threat levels associated with different durations of electrical shock (low: 5ms, medium: 35ms, and high: 250ms). Reward and threat were combined in a 3 x 3 full factorial manner, resulting in nine combinations of reward and threat. After trial onset, participants had 2.5 seconds to decide to approach or avoid by pulling the joystick towards themselves or pushing the joystick away from themselves, respectively. As confirmation of a timely response (<2.5s) a white line appeared below the reward for the remainder of the decision phase and the subsequent outcome anticipation phase.

Next, a 5-7s *outcome anticipation phase* started. In this phase, the avatar and monetary amount remained on the screen while participants anticipated the outcome of their decision.

Finally, participants received one of three possible outcomes: positive, negative, or neutral. In a positive outcome, the avatar offered a stack of banknotes. In a negative outcome, the avatar drew a gun and shot. Additionally, participants received an electric shock with a duration that corresponded to the threat level symbolized by the avatar. During a neutral outcome, the avatar kept its hands behind its back, resulting in omission of both reward and threat. The probability of receiving a particular outcome depended on the decision that participants made (Figure 1C). Approach led to a 40/40% chance of receiving a positive or negative outcome and a 20% chance on receiving a neutral outcome, whereas avoidance led to an 80% chance of receiving a neutral outcome and a 10/10% chance of receiving a positive or negative outcome. These differential outcome contingencies of approach and avoidance were modelled after real-life approach-avoidance conflicts. In such conflicts, approach decisions typically entail a higher risk of encountering perceived threats and rewards, while

avoidance decisions mitigate these threats but also forfeit potential rewards (Krypotos et al.,

2018). Participants were instructed that late responses (>2.5s) always led to the negative

outcome, i.e., getting shot and receiving an electrical shock of a duration that was symbolized

by the avatar. After participants received the outcome, a fixation cross was displayed during

the inter-trial interval (6-8s).

Since this study did not aim to investigate de novo contingency learning, participants

received explicit instructions about the association between the avatar and its corresponding

threat level, as well as the outcome probabilities of approach and avoidance decisions.

Providing explicit instructions has been shown to significantly enhance learning of

contingencies (Mertens et al., 2018), leading to stronger task effects, including those observed

in neuroimaging measures (Mechias et al., 2010). Thus, contingencies were all clearly and

repeatedly explained to the participants before starting the task, and comprehension was

verified during a short practice session of 9 trials in the MRI scanner. The use of avatars as

conditioned stimuli was intended to make the task more engaging. Previous research has

shown that such stimuli can enhance motivation (Sailer et al., 2017) and sustain attention

(Krath et al., 2021). The association between avatars and threat levels was counterbalanced

across participants. In total, participants completed 90 trials (10 trials for each combination of

reward and threat).

Figure 1. Overview of the Fearful Avoidance Task (FAT). (A) Example of the FAT trial structure. In

each trial, participants were first confronted with a combination of reward (displayed in text) and threat

208 (signalled by the avatar). Participants had 2.5s to decide to approach or avoid by pulling the joystick

209 towards themselves or pushing it away, respectively. Next, they anticipated the outcome of their

210 decision. Finally, the outcome of their decision was displayed. After receiving the outcome, a fixation

211 cross was presented during the inter-trial interval (ITI). **(B)** Overview of the reward levels: reward levels

212 were low (IP – 40-50%), medium (IP ± 5%), or high (IP + 40-50%), relative to the individual's indifference

213 point (IP). In this example, with an IP of €1.00, the low reward level ranged from €0.50 to €0.60, the

214 medium reward level ranged from €0.95 to €1.05, and the high reward level ranged from €1.40 to €1.50.

215 **(C)** Overview of the threat levels: three distinct avatars signalled shock duration. Contingencies between

216 avatars and threat levels were counterbalanced across participants. **(D)** Overview of the outcome

217 probabilities for approach and avoidance decisions. Approach decisions involved a high risk of receiving

218 electric shocks or earning monetary rewards, while avoidance decisions diminished the risk of electric

219 shocks but also reduced the potential for monetary rewards.

221 **MRI acquisition**

222 Data was acquired on a 3T MAGNETOM PrismaFit MR scanner (Siemens AG, Healthcare

223 Sector, Erlangen, Germany) using a 32-channel head coil. A T1-weighted scan was acquired

224 in the sagittal orientation using a 3D MPRAGE sequence (TR=2000ms, TE=3.03ms, 192

225 sagittal slices; 1.0mm isotropic voxels; FOV=256mm). T2-weighted volumes were acquired

226 using a multi-band multi-echo (MB3ME3) sequence, a fast sequence designed for whole brain

227 coverage with reduced artefacts and signal dropout in medial prefrontal and subcortical regions

228 (Cohen et al., 2018; Fazal et al., 2023) (TR = 1500ms, TE_{1-3} = 13.4/34.8/56.2ms, flip angle =

229 75°, 51 sagittal slices; 2.5mm isotropic voxels).

231 **Behavioural analyses**

232 First, we verified whether the data adhered to the preregistered inclusion criteria, which

233 stipulated a maximum of 50% missing responses and absence of atypical response patterns

234 as defined by Hulsman, Klaassen, et al., (2021). Atypical response patterns refer to behaviour

235 indicating that the participant may not have comprehended the task correctly or did not perform

236 the task seriously (e.g., when participants avoided substantially more on low threat trials than

high threat trials, see supplement for a detailed description of atypical response patterns). All participants met these inclusion criteria.

To investigate the effect of reward, threat, and their interaction on the decision (approach/avoid) that participants made, we ran a Bayesian multilevel model. For full model specifications, see equation (1) below. Other model specifications were as described in the next section.

$$Decision \; _{approach \; vs \; avoid} \sim Reward * Threat + (1 + Reward * Threat \,|\, Subject) \qquad (1)$$

Bayesian multilevel analyses (general)

As preregistered (Hulsman et al., 2021b), throughout behavioural and fMRI analyses, we employed linear Bayesian multilevel models. These linear Bayesian multilevel models were executed using R (Version 4.2.1; R Core team, 2022) within RStudio (Version 2022.12.0; RStudio Inc., 2009-2022) using the brms package (2.18.0, Bürkner, 2013 and Carpenter et al., 2017). All Bayesian multilevel models adhered to the following configurations: continuous predictors were standardized, and categorical predictors were coded using sum-to-zero contrasts. All models included a maximal random effects structure, consisting of a random intercept for each participant along with random slopes for all predictors and their interactions. Models with a binomial dependent variable (i.e., decision: approach/avoid) were modelled using a Bernoulli distribution, whereas models with continuous dependent variables (i.e., fMRI beta values) were modelled using a Gaussian distribution. Models were fitted using 4 chains with 4000 iterations each (2000 warm-up). A coefficient was deemed statistically significant when the associated ≥95% posterior credible interval was non-overlapping with zero. As recommended for analyses with an effective sample size <10.000 (Kruschke, 2014) this was supplemented with 90% posterior credible intervals when the 95% intervals were overlapping with 0, given that these intervals may yield more stable results (Makowski et al., 2019). These latter results are reported as trends when the 90% interval was non-overlapping with zero. When effects were non-overlapping with zero at the 90 or 95% interval, we additionally tested

99% and 99.9% intervals to further explore the likelihood of these effects (akin to reporting

$p<.01$ and $p<.001$, respectively, rather than only $p<.05$).

fMRI analyses

The fMRI data were processed using SPM12 (Wellcome Trust Centre for Neuroimaging,

London, UK) following the standard preprocessing procedures. Functional scans were

combined into a single image using the PAID-weighting method (Poser et al., 2006), co-

registered to the anatomical scan, and normalized to the Montreal Neurological Institute (MNI)

152 T1-template using nonlinear transformations based on SPM's tissue probability map. The

normalized images (2mm isotropic) were then smoothed with an isotropic 3D Gaussian kernel

with 6mm full width at half maximum (FWHM). For all models, high-pass filtering at 1 /128 Hz

and a first-order autoregressive model were used as standard.

Our preregistered regions of interest (ROIs, see Figure 2) included the anterior insula

(Shirer et al., 2012), amygdala (Rolls et al., 2020), bed nucleus of the stria terminalis (Avery et

al., 2014), dorsal anterior cingulate cortex (Shirer et al., 2012), periaqueductal gray (Lojowska

et al., 2015), thalamus (Rolls et al., 2020), ventral medial prefrontal cortex (Rolls et al., 2020),

and ventral striatum (Piray et al., 2017).

Figure 2. Regions of interest (ROIs). aInsula = anterior insula, amygdala, BNST = bed nucleus of the stria terminalis, dACC = dorsal anterior cingulate cortex, PAG = periaqueductal gray, thalamus, vmPFC = ventral medial prefrontal cortex, vStriatum = ventral striatum, CMA = central medial amygdala, BLA = basolateral amygdala, SFA = superficial amygdala, AStr = amygdalostriatal region. We hypothesized that the aInsula, dACC, PAG, and thalamus show sensitivity to both reward and threat. Additionally, we hypothesized that the amygdala and BNST are predominantly sensitive to threat, while the vmPFC and vStriatum are predominantly sensitive to reward. Furthermore, we expected that the dACC and vmPFC are involved in the integration of reward and threat.

To gain global insight into the neural activation patterns probed by the task, independent of the level of reward, the level of threat, and the decision, we first report the results of the whole-brain voxel-wise analysis contrasting neural activity during the *decision phase* against the baseline (i.e., the intertrial interval). All fMRI figures were created in MRIcroGL, with the use of the MNI 152 template image. As preregistered, whole-brain results

were familywise (FWE) corrected for multiple comparisons according to random field theory at

the cluster level (*p*<.05) after an initial cluster-defining threshold of *p*<.001 uncorrected.

Bayesian Multivariate Multilevel (BMM) fMRI analysis: effects of reward, threat, and

decision

To investigate the effects of reward, threat, decision, and their interactions on neural activity

within our regions of interest, we leveraged Bayesian Multivariate Multilevel (BMM) analyses.

In our task, we expected that approach and avoidance decisions would depend on specific

task conditions, and indeed, in some individuals, conditions and behaviour were strongly

correlated. For instance, in the low reward/high threat condition, several participants

consistently decided to avoid. Given these circumstances, conventional MRI analyses are less

suited to accurately disentangle the effect of reward/threat from the effect of the decision to

approach or avoid). Particularly for these intricate designs, BMM analyses surpass

conventional MRI analyses by effectively managing data dependency and unbalanced data,

leading to increased accuracy of results (Chen et al., 2019a, 2019b). Therefore, BMM

analyses emerged as the most optimal choice to investigate the effects of reward, threat,

decision, and their interactions. Given that conducting BMM analyses on whole-brain voxel-

wise level is not computationally feasible, we performed the BMM analyses on our

preregistered regions of interest. In line with our preregistration, we also conducted two

additional whole-brain voxel-wise analyses: one parametric modulation analysis focussing on

stimulus effects (i.e., the effect of reward/threat) and one general linear model (GLM) analysis

focussing on response effects (i.e., the effect of the decision to approach or avoid). However,

these additional analyses partially overlap with the BMM analyses, which, as explained above,

are better suited for capturing higher-order interactions between task conditions and the

observed behaviour. Therefore, we report the results of the BMM analyses. For completeness,

voxel-wise model specifications and results are provided at: https://osf.io/hbjp5.

Subject-level analysis

After preprocessing, we employed subject-level models using SPM12 (Wellcome Trust Centre

for Neuroimaging, London, UK). For each trial, we fitted one regressor for the *decision phase*

(i.e., from trial onset until the response) and one regressor for the *outcome anticipation phase*

(i.e., from response until the outcome), leading to 90 regressors for the *decision phase*, 90

regressors for the *outcome anticipation phase*. Additionally, we added regressors for each type

of outcome (positive, negative, neutral) and motion regressors (3 translational, 3 rotational).

This resulted in 3 additional regressors for the outcomes, 6 motion regressors, and 1 constant,

leading to a total of 190 regressors. There was no censoring of data applied. The collinearity

among all regressors of the *decision phase* and *outcome anticipation phase* was within an

acceptable range ($M = 0.0135$, $Mdn = 0.0056$, $SD = 0.0368$, $min = 0$, $max = 0.4025$, see

Mumford et al., 2015), with the highest collinearity observed between the regressors of the

decision phase and *outcome anticipation phase* within the same trial. For each trial, we

contrasted the regressor of each phase against the regressors of the corresponding phase in

all other trials. Finally, trial-by-trial mean betas for each phase were extracted for each ROI

using the MarsBaR toolbox (Brett et al., 2002). To ensure that the findings from our ROI

analyses mainly stem from within-region voxels and not adjacent white matter or cerebrospinal

fluid, we repeated our BMM analyses on the unsmoothed data from our ROI masks. These

control analyses confirmed all key findings. Consequently, we report the results of the

preregistered analyses on the smoothed data, as this approach accounts for anatomical

variability and thereby yields more statistical power (Mikl et al., 2008).

Group-level analysis

For each phase, we employed a Bayesian Multivariate Multilevel (BMM) analysis where we

used the trial-by-trial betas of our ROIs as dependent variables and reward, threat, decision,

and their interactions as predictors. Given the correlational nature of our analysis, the choice

whether variables are entered as dependent variables or predictors does not impact the

results. However, since neural activity across different ROIs is correlated, using a multivariate

model that accounts for this variance provides the best model fit and most accurate inferences.

For full model specifications, see equation (2) below. Other model specifications were as

described in the section 'Bayesian multilevel analyses (general)' above.

$$B_{aInsula}, B_{AMY}, B_{BNST}, B_{dACC}, B_{PAG}, B_{Thalamus}, B_{vStriatum}, B_{VMPFC} \sim \qquad (2)$$

$$Reward * Threat * Decision + (1 + Reward * Threat * Decision \,|\, Subject)$$

354   Given that the amygdala consists of functionally distinct anatomical subregions (McDonald,

355   1998; Price, 2003; Bzdok et al., 2013), we explored whether significant findings in the

356   amygdala could be attributed to distinct contributions from its subnuclei: amygdalostriatal

357   (AStr), basolateral (BLA), centromedial (CMA), and superficial (SFA) as defined in the SPM

358   Anatomy Toolbox (Amunts et al., 2005; Eickhoff et al., 2007), see equation (3) below.

$$B_{AStr,}, B_{BLA,}, B_{CMA}, B_{SFA} \sim \qquad (3)$$

$$Reward * Threat * Decision + (1 + Reward * Threat * Decision \,|\, Subject)$$

In the event of any significant interactions between predictors in our BMMs, planned

comparisons were conducted using the emmeans package (Lenth et al., 2018) to clarify the

nature of these interactions. Specifically, we investigated the interactions between reward and

decision, as well as threat and decision, by examining whether the slope of reward was

significant for each type of decision (approach/avoid) and whether the slope of threat was

significant for each type of decision (approach/avoid). For interactions between reward, threat,

and decision, we assessed whether the slope of threat was significantly different between

approach and avoidance decisions in low and high reward conditions separately. If a significant

difference was found within a specific reward level, we further examined whether the slope of

threat was significant for each type of decision (approach/avoid) within that reward level. Betas

and posterior credible intervals of all main effects, interaction effects, and post-hoc

comparisons are provided at: https://osf.io/hbjp5.

RESULTS

Avoidance behaviour

As expected, participants avoided significantly more with decreasing reward levels (B = -2.01, 99.9% CI [-3.49, -0.74]) and increasing threat levels (B = 1.79, 99.9% CI [0.86, 2.75], see Figure 3). In low threat conditions, the effect of reward appears to be less pronounced, as participants typically approach, while in high reward conditions, participants tend to approach regardless of threat. However, this was not reflected in a significant interaction between reward and threat on avoidance (B = -.01, 90% CI [-0.19, 0.15]).

Figure 3. Overview task effects on avoidance behaviour. The bars represent the mean proportion avoidance as function of reward and threat level, overlaid with individual data points to illustrate variance between participants. Results showed the hypothesized opposing influences of reward and threat on avoidance.

fMRI: decision phase

Whole-brain voxel-wise analysis: decision phase vs baseline

Initial whole-brain voxel-wise analyses confirmed global activation across regions of interest hypothesized to be associated with weighing of reward and threat (aInsula, amygdala, BNST, dACC, PAG, thalamus, and vStriatum) when making approach-avoidance decisions (vs baseline; see Figure 4). Subsequently, to capture the weighing of rewards and threats for approach and avoidance decisions for each region of interest, we employed Bayesian Multivariate Multilevel (BMM) analyses.

Figure 4. Whole-brain fMRI results demonstrating widespread activations during approach-

avoidance decision-making (contrast: decision phase vs baseline). Images are shown at a cluster

threshold of $p<.05$ corrected for multiple comparisons (FWE) after an initial threshold of $p<.001$

uncorrected. All marked regions of interests reach significance after correction for multiple comparisons

($p_{FWE}<.05$). aInsula = anterior insula, AMY = amygdala, BNST = bed nucleus of the stria terminalis,

dACC = dorsal anterior cingulate cortex, PAG = periaqueductal gray, vStriatum = ventral striatum.

Bayesian Multivariate Multilevel fMRI analysis: reward, threat, and decision effects

First, we investigated the effects of reward, threat, decision, and their interactions on BOLD

activity within our ROIs during the *decision phase*, i.e., prior to indicating approach-avoidance

decisions (see Figure 5). We found that BOLD activity in the bed nucleus of the stria terminalis

(BNST), periaqueductal gray (PAG), and ventral striatum increased as a function of reward,

whereas BOLD activity in the dorsal anterior cingulate (dACC) increased as a function of threat.

We did not find significant differences in overall BOLD activity between approach and

avoidance decisions, nor significant interactions between reward and threat for any ROIs

independent of decision (all 90% credible intervals overlapping with 0).

Figure 5. Posterior estimates and credible intervals for task-related activations in the regions of

interest during the *decision phase*. This figure illustrates the posterior estimates (dots) with respect

to the null (vertical dashed line). Bold lines represent 90% posterior credible intervals, while thin lines

represent 95% posterior credible intervals. Significant effects, defined as ≥90% posterior credible

intervals non-overlapping with zero, are highlighted in color. aInsula = anterior insula, BNST = bed

nucleus of the stria terminalis, dACC = dorsal anterior cingulate cortex, PAG = periaqueductal gray,

vmPFC = ventral medial prefrontal cortex, vStriatum = ventral striatum.

More importantly, we discovered that in several regions, the effect of reward and threat

on BOLD activity differed between subsequent approach and avoidance decisions.

Specifically, both the amygdala and vmPFC exhibited stronger BOLD activity with increasing

reward preceding risky approach, but not risk-diminishing avoidance decisions (reward x

decision; see Figure 6A), in line with the idea that activity in these regions may drive approach

responses with increasing reward. Exploratory analyses indicated that this effect was present

for all amygdala subnuclei (B_{Astr} = 0.05, 95% CI [0.00, 0.10]; B_{BLA} = 0.05, 95% CI [0.00, 0.10];

B_{CMA} = 0.05, 90% CI [0.00, 0.07]; B_{SFA} = 0.05, 90% CI [0.01, 0.09]). Moreover, stronger BOLD

activity was observed in the amygdala in response to increasing threat for risk-diminishing

avoidance relative to risky approach decisions (threat x decision; see Figure 6B), thus

suggestive of a dual role for the amygdala in processing both rewards and threats. This effect

was particularly driven by the amygdalostriatal subregion of the amygdala (B_{Astr} = -0.05, 95%

CI [-0.09, -0.01], 90% credible intervals for all other subregions overlapping with 0).

Interestingly, in addition to these two-way interactions, BOLD activity in the BNST, thalamus

and ventral striatum exhibited a pattern suggesting that the integration of reward and threat

differed for approach and avoidance decisions (reward x threat x decision; see Figure 6C).

Specifically, when participants decided to avoid in low reward conditions, BOLD activity in the

thalamus and ventral striatum was stronger with increasing threat. In contrast, prior to

indicating risky approach decisions in low reward conditions, BOLD activity in the BNST and

ventral striatum was decreased with increasing threat. Conversely, no differences in threat

reactivity between approach and avoidance decisions were found when reward was high and

the impact of threat on decisions much smaller (see Figure 3).

In conclusion, our findings reveal that BOLD activity as a function of reward and threat

preceding approach-avoidance decisions varied depending on the subsequent decision.

Interestingly and contrary to theoretical predictions, several subcortical regions exhibited a

pattern suggesting an integrative role.

Figure 6. Effects of reward and threat on neural activity during approach-avoidance decision-making. The figure presents marginal effects plots for interactions between reward, threat, and decision in the *decision phase*. These plots depict separately for approach (solid lines) and avoidance (dashed lines): **(A)** the effect of *reward*, **(B)** the effect of *threat*, and **(C)** the *interaction between reward and threat* across all regions of interest where these interactions were observed. For each region of interest, only the highest-order interaction is plotted as it conveys most information. The asterisks indicate the

449 significance levels of the slopes of **(A)** reward and **(B, C)** threat for approach and avoidance separately.

450 #≥90%, *≥95%, **≥99%, or ***≥99.9% posterior credible intervals non-overlapping with zero. vmPFC =

451 ventomedial prefrontal cortex, BNST = bed nucleus of the stria terminalis, vStriatum = ventral striatum.

fMRI: outcome anticipation phase

Bayesian Multivariate Multilevel fMRI analysis: reward, threat, and decision effects

Recent research suggests involvement of specific brain regions may depend on the specific

stage of the approach-avoidance decision-making process. Therefore, we subsequently

probed the contribution of our ROIs at the stage following approach-avoidance decisions, when

participants were passively anticipating the outcome. As indicated above, risky approach

decisions were associated with a relatively high chance of receiving either electric stimulation

or monetary reward (40% for each; high risk/high gain), whereas risk-diminishing avoidance

decisions were associated with a relatively low chance of receiving these outcomes (10% for

each; low risk/low gain). During this *outcome anticipation phase*, BOLD activity in the

amygdala, BNST, PAG, and thalamus was more pronounced following risky approach

decisions compared to risk-diminishing avoidance decisions (see Figure 7). Moreover, we

observed stronger BOLD activity in the aInsula and PAG with increasing threat. We did not

find any significant main effects of reward nor interactions between reward and threat in any

ROI (all 90% credible intervals overlapping with 0).

Figure 7. Posterior estimates and credible intervals for task-related activations in the regions of

interest during the *outcome anticipation phase*. This figure illustrates the posterior estimates (dots)

with respect to the null (vertical dashed line). Bold lines represent 90% posterior credible intervals, while

thin lines represent 95% posterior credible intervals. Significant effects, defined as ≥90% posterior

credible intervals non-overlapping with zero, are highlighted in color. aInsula = anterior insula, BNST =

bed nucleus of the stria terminalis, dACC = dorsal anterior cingulate cortex, PAG = periaqueductal gray,

vmPFC = ventral medial prefrontal cortex, vStriatum = ventral striatum.

Interestingly, in several regions, the influence of reward and threat on the BOLD

response during the outcome anticipation phase depended on the preceding decision.

Following risk-diminishing avoidance decisions, when the chance of receiving shocks or

rewards was low, BOLD activity within the vStriatum and vmPFC significantly decreased with

increasing reward. Conversely, no such effects were observed following risky approach

decisions, when the chance of receiving shocks or rewards was relatively high (reward x

decision; see Figure 8A). Furthermore, we found that following risk-diminishing avoidance

decisions, BOLD activity in the thalamus decreased with increasing threat. In contrast,

following risky approach decisions, when the risk of receiving electrical stimulation was high,

BOLD activity in the aInsula, BNST, dACC, PAG, and thalamus increased with increasing

threat (threat x decision; see Figure 8B). This finding further aligns with the presumed role of

these regions in anticipating impending threats. Notably, there was integration of reward and

threat in the aInsula, exhibiting variations between approach and avoidance decisions (reward

x threat x decision; see Figure 8C). This interaction was driven by the fact that while

anticipating the outcome of avoidance decisions (low risk), BOLD activity in the aInsula was

marginally decreased as a function of threat in low reward conditions. In contrast, while

anticipating the outcome of approach decisions (high risk), BOLD activity in the aInsula

increased as a function of threat in low reward conditions. Thus, the pattern of BOLD activity

in the aInsula was similar to patterns observed in other threat anticipation regions. However, it

was the only region that exclusively exhibited this pattern when reward was low.

In conclusion, following avoidance decisions, when the chance of receiving rewards

was relatively low, regions within the reward network negatively tracked reward magnitude.

Conversely, after approach decisions, when the risk of receiving electric shock was relatively high, regions in the salience network positively tracked threat magnitude, with the aInsula only doing so when reward was low.

Figure 8. Effects of reward and threat on neural activity during anticipation of the outcome of approach-avoidance decisions. The figure presents marginal effects plots for interactions between reward, threat, and decision in the *outcome anticipation phase*. These plots depict separately for approach (solid lines) and avoidance (dashed lines): **(A)** the effect of *reward*, **(B)** the effect of *threat*, and **(C)** the *interaction between reward and threat* across all regions of interest where these interactions were observed. For each region of interest, only the highest-order interaction is plotted as it conveys most information. The asterisks indicate the significance levels of the slopes of **(A)** reward and **(B, C)** threat for approach and avoidance separately. #≥90%, *≥95%, **≥99%, or ***≥99.9% posterior credible intervals non-overlapping with zero. vStriatum = ventral striatum, vmPFC = ventromedial prefrontal

cortex, BNST = bed nucleus of the stria terminalis, dACC = dorsal anterior cingulate cortex, PAG =

periaqueductal gray, aInsula = anterior insula.

DISCUSSION

In this study, we investigated how the decision to engage in risky approach or risk-diminishing

avoidance may arise from distributed neural processing of reward and threat information. We

aimed to provide a better understanding of the neural dynamics governing approach-

avoidance decisions by 1) investigating whether the decision to approach or avoid is preceded

by parallel and/or integrated neural processing of reward and threat across brain regions

previously implicated in approach-avoidance decision-making and 2) evaluated how such

processing and integration evolve over time following approach-avoidance decisions during

outcome anticipation. Our preregistered Bayesian Multivariate Multilevel (BMM) analyses on

fMRI data revealed two key findings. First, in contrast to the notion of *parallel processing* of

reward and threat in subcortical regions traditionally associated predominantly with either

reward (vStriatum, vmPFC) or threat (amygdala, BNST) processing, we found evidence for

dual and *integrated processing*. Specifically, there was *dual processing* of reward and threat

information in the amygdala and evidence for *integration* in other subcortical regions, including

the vStriatum, thalamus, and BNST. Critically, the weighing of reward and threat in these

regions varied as a function of the subsequent decision to approach or avoid. Second, after

indicating the decision to approach or avoid, we observed that neural activity in regions

traditionally associated with reward (vmPFC, vStriatum) and threat (BNST) processing, as well

as the salience network (dACC, thalamus, PAG), was influenced the magnitudes of reward

and threat, consistent with previous theories. Together, these findings illuminate neural

dynamics of approach-avoidance decision-making and suggest distributed subcortical

integration of reward and threat as a potential driver of subsequent approach-avoidance

decisions.

In accordance with prior research, we found that when making decisions involving risky

approach or risk-dimishing avoidance, individuals carefully balance potential rewards and

threats (Talmi et al., 2009; Basten et al., 2010; Park et al., 2011; Aupperle et al., 2015; Schlund et al., 2016; Hulsman et al., 2021a, 2024; Klaassen et al., 2021; Moughrabi et al., 2022). To delve deeper into the distributed neural patterns underlying this decision-making process, we employed Bayesian Multivariate Multilevel (BMM) analyses. The observed neural patterns deviated from conventional perspectives that primarily suggest isolated reward and threat processing in subcortical regions, followed by the integration of reward and threat in cortical regions such as the ACC or vmPFC (Yacubian et al., 2006; Talmi et al., 2009; Basten et al., 2010; Park et al., 2011; Azab and Hayden, 2020; Livermore et al., 2021). Instead, our findings suggest that the decision to approach or avoid is supported by partially overlapping subcortical regions sensitive to both reward and threat information. Specifically, the amygdalostriatal subnucleus of the amygdala, BNST, thalamus, and vStriatum showed a pattern of increasing neural activity as function of threat prior to indicating risk-diminishing avoidance as compared to risky approach decisions. In contrast, the vmPFC uniquely displayed a reward-by-decision effect without being moderated by threat, consistent with its hypothesized role in reward valuation (Talmi et al., 2009; Aupperle and Paulus, 2010; Basten et al., 2010; Bartra et al., 2013) and subsequent approach decisions (Pedersen et al., 2021). The convergence of brain regions engaged in processing both reward and threat corresponds with recent findings demonstrating that the BNST, vStriatum, and PAG respond to both reward and threat information (Murty et al., 2023) and that conflict between approach and avoidance tendencies is represented at the level of the vStriatum (Ironside et al., 2020). Additionally, converging findings have shown that the amygdala responds to both aversive and appetitive stimuli (Baxter and Murray, 2002; O'Doherty, 2004; Schultz, 2006; Pessoa, 2010).

Our findings, which highlight subcortical integration of reward and threat, provide a new perspective that contrasts with prior human neuroimaging research. However, they do resonate with long-standing reports in rodents (Gentry et al., 2016; Burgos-Robles et al., 2017; Hamel et al., 2017; Beyeler et al., 2018), collectively suggesting that the traditional notion of these regions responding primarily to either reward or threat is untenable. Nevertheless, it remains possible that depending on the context certain regions predominantly encode reward

or threat information, as observed here, with vmPFC and PAG demonstrating sensitivity to

reward and the dACC to threat. Tentatively, such signals could serve as input for regions

integrating both reward and threat inputs, such as the BNST, vStriatum, and thalamus. This

perspective aligns with theories proposing that these regions, due to their connectivity profile,

may serve as intermediaries between lower downstream subcortical and upstream cortical

regions (Avery et al., 2014; Goode and Maren, 2017; De Groote and de Kerchove d'Exaerde,

2021; Hammack et al., 2021; Sieveritz and Raghavan, 2021), akin to striatal-thalamo-cortical

loops in motor control (Pessoa, 2023).

Furthermore, the observed increase in threat-related activity in the vStriatum,

thalamus, and BNST preceding avoidance relative to approach under low reward, aligns with

recent findings that increased threat representations in these regions during decision-making

were linked to avoidance (Moughrabi et al., 2022). Behaviourally, the presence of high reward

attenuated the impact of threat on subsequent decisions. This attenuation was also reflected

in neural responses within the vStriatum, thalamus, and BNST. These findings further align

with previous research outside the field of value-based decision making that demonstrate

competition between reward and threat processing in various brain regions, including the

BNST (Choi et al., 2014).

Finally, we demonstrated shifts in these neural patterns of reward/threat processing

following approach-avoidance decisions. Following risky approach decisions, we observed

positive tracking of threat in the salience and threat network, encompassing the BNST,

thalamus, dACC, and PAG, all of which have consistently been associated with threat

anticipation (Mechias et al., 2010; Fullana et al., 2016; Klumpers et al., 2017; Andrzejewski et

al., 2019; Patrick et al., 2019). Interestingly, in the aInsula we only found increased activity as

a function of threat following approach decisions when rewards were low. This aligns with

previous research suggesting that the presence of reward can suppress the effect of threat in

the aInsula (Cristofori et al., 2015). Conversely, following risk-diminishing avoidance decisions

we found negative tracking of reward in the reward network, including the vStriatum and

vmPFC. These findings appear to reflect decreased reward expectancies after avoidance

decisions (Heekeren et al., 2007; Jia et al., 2016; Pujara et al., 2016; Rehbein et al., 2023).

Despite the implementation of a fast multiband-multiecho fMRI sequence (MB3ME3),

further delineation the temporal dynamics of approach-avoidance decisions would require

methods with superior temporal resolution, such as MEG or EEG (Khemka et al., 2017;

McFadyen et al., 2023). Another limitation of our study is a relatively small sample size (N=28

included in MRI analyses). Our analyses were preregistered, and we employed BMM analyses

to investigate neural responses to minimize the risk of false positives (Kajimura et al., 2023).

Still, replication of our findings is desirable, particularly in clinical populations that show

approach-avoidance deficits to gain further knowledge of how reward-threat balances across

distinct regions may be disturbed (Ironside et al., 2020; Smith et al., 2021b, 2021a, 2023;

McDermott et al., 2022).

In conclusion, our findings unveiled distributed cortico-subcortical processing and

subcortical integration of reward and threat prior to the decision to approach or avoid. These

findings suggest a departure from traditional, yet widely professed, views that segregate brain

regions as either predominantly reward-sensitive or threat-sensitive, and assign the integration

of reward and threat primarily to cortical regions. Following approach-avoidance decisions,

however, brain regions traditionally associated with reward and threat processing reflected

reward and threat tracking contingent on the decision that was made. These findings provide

new theoretical insight into the unfolding neural dynamics of approach-avoidance decision-

making.

References

References

Abivardi A, Khemka S, Bach DR (2020) Hippocampal Representation of Threat Features and
Behavior in a Human Approach–Avoidance Conflict Anxiety Task. J Neurosci
40:6748–6758.

Alvarez EO, Ruarte MB (2001) Role of glutamate receptors in the nucleus accumbens on
behavioural responses to novel conflictive and non-conflictive environments in the rat.
Behav Brain Res 123:143–153.

American Psychiatric Association (2013) Diagnostic and statistical manual of mental
disorders, 5th ed. Washington, DC. Available at:
https://doi.org/10.1176/appi.books.9780890425596.

Amunts K, Kedo O, Kindler M, Pieperhoff P, Mohlberg H, Shah NJ, Habel U, Schneider F,
Zilles K (2005) Cytoarchitectonic mapping of the human amygdala, hippocampal
region and entorhinal cortex: Intersubject variability and probability maps. Anat
Embryol (Berl) 210:343–352.

Andrzejewski JA, Greenberg T, Carlson JM (2019) Neural correlates of aversive anticipation:
An activation likelihood estimate meta-analysis across multiple sensory modalities.
Cogn Affect Behav Neurosci 19:1379–1390.

Aupperle RL, Melrose AJ, Francisco A, Paulus MP, Stein MB (2015) Neural substrates of
approach-avoidance conflict decision-making. Hum Brain Mapp 36:449–462.

Aupperle RL, Paulus MP (2010) Neural systems underlying approach and avoidance in
anxiety disorders. Dialogues Clin Neurosci 12:517–531.

Avery SN, Clauss JA, Winder DG, Woodward N, Heckers S, Blackford JU (2014) BNST
neurocircuitry in humans. NeuroImage 91:311–323.

Azab H, Hayden B (2020) Partial integration of the components of value in anterior cingulate
cortex. Behav Neurosci 134:296–308.

Bartra O, McGuire JT, Kable JW (2013) The valuation system: A coordinate-based meta-
analysis of BOLD fMRI experiments examining neural correlates of subjective value.
NeuroImage 76:412–427.

Basten U, Biele G, Heekeren HR, Fiebach CJ (2010) How the brain integrates costs and
benefits during decision making. Proc Natl Acad Sci 107:21767–21772.

Baxter MG, Murray EA (2002) The amygdala and reward. Nat Rev Neurosci 3:563–573.

Beaufour CC, Le Bihan C, Hamon M, Thiébot MH (2001) Extracellular serotonin is enhanced
in the striatum, but not in the dorsal hippocampus or prefrontal cortex, in rats
subjected to an operant conflict procedure. Behav Neurosci 115:125–137.

Beyeler A, Chang C-J, Silvestre M, Lévêque C, Namburi P, Wildes CP, Tye KM (2018)
Organization of Valence-Encoding and Projection-Defined Neurons in the Basolateral
Amygdala. Cell Rep 22:905–918.

Brett M, Anton J-L, Valabregue R, Poline J-B (2002) Region of interest analysis using an
SPM toolbox [abstract].

Burgos-Robles A, Kimchi EY, Izadmehr EM, Porzenheim MJ, Ramos-Guasp WA, Nieh EH,
Felix-Ortiz AC, Namburi P, Leppla CA, Presbrey KN, Anandalingam KK, Pagan-
Rivera PA, Anahtar M, Beyeler A, Tye KM (2017) Amygdala inputs to prefrontal
cortex guide behavior amid conflicting cues of reward and punishment. Nat Neurosci
20:824–835.

Bürkner P-C (2013) brms : An R Package for Bayesian Multilevel Models using Stan. J Stat
Softw 80:1–28.

Bzdok D, Laird AR, Zilles K, Fox PT, Eickhoff SB (2013) An investigation of the structural,
connectional, and functional subspecialization in the human amygdala. Hum Brain
Mapp 34:3247–3266.

Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker MA, Li P,
Riddell A (2017) Stan : A Probabilistic Programming Language. J Stat Softw 76.

Chen G, Paul PB, Zhihao AT, Lijun L, Glen DR, Kinnison J, Cox RW, Pessoa L (2019a) An
integrative Bayesian approach to matrix-based analysis in neuroimaging. Hum Brain
Mapp:4072–4090.

Chen G, Xiao Y, Taylor PA, Rajendra JK, Riggins T, Geng F, Redcay E, Cox RW (2019b)
Handling Multiplicity in Neuroimaging Through Bayesian Lenses with Multilevel
Modeling. Neuroinformatics 17:515–545.

Ch'ng S, Fu J, Brown RM, McDougall SJ, Lawrence AJ (2018) The intersection of stress and
reward: BNST modulation of aversive and appetitive states. Prog
Neuropsychopharmacol Biol Psychiatry 87:108–125.

Choi JM, Padmala S, Spechler P, Pessoa L (2014) Pervasive competition between threat
and reward in the brain. Soc Cogn Affect Neurosci 9:737–750.

Cohen AD, Nencka AS, Wang Y (2018) Multiband multi-echo simultaneous ASL/ BOLD for
task-induced functional MRI. PLoS ONE 13:1–21.

Corr PJ (2013) Approach and Avoidance Behaviour: Multiple Systems and their Interactions.
Emot Rev 5:285–290.

Cristofori I, Harquel S, Isnard J, Mauguière F, Sirigu A (2015) Monetary reward suppresses
anterior insula activity during social pain. Soc Cogn Affect Neurosci 10:1668–1676.

De Groote A, de Kerchove d'Exaerde A (2021) Thalamo-Nucleus Accumbens Projections in
Motivated Behaviors and Addiction. Front Syst Neurosci 15:711350.

Delgado MR, Jou RL, Phelps EA (2011) Neural systems underlying aversive conditioning in
humans with primary and secondary reinforcers. Front Neurosci 5:1–10.

Duque-Wilkens N, Steinman MQ, Busnelli M, Chini B, Yokoyama S, Pham M, Laredo SA,
Hao R, Perkeybile AM, Minie VA, Tan PB, Bales KL, Trainor BC (2018) Oxytocin
Receptors in the Anteromedial Bed Nucleus of the Stria Terminalis Promote Stress-
Induced Social Avoidance in Female California Mice. Biol Psychiatry 83:203–213.

Eickhoff SB, Paus T, Caspers S, Grosbras M-H, Evans AC, Zilles K, Amunts K (2007)
Assignment of functional activations to probabilistic cytoarchitectonic areas revisited.
NeuroImage 36:511–521.

Engelke DS, Zhang XO, O'Malley JJ, Fernandez-Leon JA, Li S, Kirouac GJ, Beierlein M, Do-
Monte FH (2021) A hypothalamic-thalamostriatal circuit that controls approach-
avoidance conflict in rats. Nat Commun 12:2517.

Fazal Z, Gomez DEP, Llera A, Marques JPRF, Beck T, Poser BA, Norris DG (2023) A
comparison of multiband and multiband multiecho gradient-echo EPI for task fMRI at
3 T. Hum Brain Mapp 44:82–93.

Fullana MA, Harrison BJ, Vervliet B, Cardoner N, Radua J (2016) Neural signatures of
human fear conditioning: an updated and extended meta-analysis of fMRI studies.
Mol Psychiatry 21:500–508.

Gentry RN, Lee B, Roesch MR (2016) Phasic dopamine release in the rat nucleus
accumbens predicts approach and avoidance performance. Nat Commun 7:13154.

Giardino WJ, Eban-Rothschild A, Christoffel DJ, Li S-B, Malenka RC, de Lecea L (2018)
Parallel circuits from the bed nuclei of stria terminalis to the lateral hypothalamus
drive opposing emotional states. Nat Neurosci 21:1084–1095.

Goode TD, Maren S (2017) Role of the bed nucleus of the stria terminalis in aversive
learning and memory. Learn Mem 24:480–491.

Hamel L, Thangarasa T, Samadi O, Ito R (2017) Caudal Nucleus Accumbens Core Is Critical
in the Regulation of Cue-Elicited Approach-Avoidance Decisions. eNeuro 4.

Hammack SE, Braas KM, May V (2021) Chemoarchitecture of the BNST: neurophenotypic
diversity and function. Handb Clin Neurol 179:385–402.

Heekeren HR, Wartenburger I, Marschner A, Mell T, Villringer A, Reischies FM (2007) Role
of ventral striatum in reward-based decision making. NeuroReport 18:951.

Hu P, Liu J, Maita I, Kwok C, Gu E, Gergues MM, Kelada F, Phan M, Zhou J-N, Swaab DF,
Pang ZP, Lucassen PJ, Roepke TA, Samuels BA (2020) Chronic Stress Induces
Maladaptive Behaviors by Activating Corticotropin-Releasing Hormone Signaling in
the Mouse Oval Bed Nucleus of the Stria Terminalis. J Neurosci 40:2519–2537.

Hulsman AM, Kaldewaij R, Hashemi MM, Zhang W, Koch SBJ, Figner B, Roelofs K,
Klumpers F (2021a) Individual differences in costly fearful avoidance and the relation
to psychophysiology. Behav Res Ther 137:103788.

Hulsman AM, Klaassen FH, Roelofs K, Klumpers F (2021b) Neural Mechanisms Underlying
Fearful Avoidance Behaviour. Available at: osf.io/7sm9k.

Hulsman AM, Terburg D, Roelofs K, Klumpers F (2021c) Chapter 28 Roles of the bed
nucleus of the stria terminalis and amygdala in fear reactions (Swaab DF, Kreier F,
Lucassen PJ, Salehi A, Buijs R, eds).

Hulsman AM, van de Pavert I, Roelofs K, Klumpers F (2024) Tackling costly fearful
avoidance using Pavlovian counterconditioning. Behav Ther 55:361–375.

Hur J, Smith JF, DeYoung KA, Anderson AS, Kuang J, Kim HC, Tillman RM, Kuhn M, Fox
AS, Shackman AJ (2020) Anxiety and the neurobiology of temporally uncertain threat
anticipation. J Neurosci 40:7949–7964.

Ironside M, Amemori K, McGrath CL, Pedersen ML, Kang MS, Amemori S, Frank MJ,
Graybiel AM, Pizzagalli DA (2020) Approach-Avoidance Conflict in Major Depressive

Disorder: Congruent Neural Findings in Humans and Nonhuman Primates. Biol
Psychiatry 87:399–408.

Jia T et al. (2016) Neural basis of reward anticipation and its genetic determinants. Proc Natl
Acad Sci 113:3879–3884.

Kajimura S, Hoshino T, Murayama K (2023) Stimulus-specific random effects inflate false-
positive classification accuracy in multivariate-voxel-pattern-analysis: A solution with
generalized mixed-effects modelling. NeuroImage 269:119901.

Khemka S, Barnes G, Dolan RJ, Bach DR (2017) Dissecting the Function of Hippocampal
Oscillations in a Human Anxiety Model. J Neurosci 37:6869–6876.

Klaassen FH, Held L, Figner B, O'Reilly JX, Klumpers F, de Voogd LD, Roelofs K (2021)
Defensive freezing and its relation to approach–avoidance decision-making under
threat. Sci Rep 11:1–12.

Klumpers F, Kroes MCW, Baas J, Fernández G (2017) How human amygdala and bed
nucleus of the stria terminalis may drive distinct defensive responses. J Neurosci
37:3830–16.

Klumpers F, Raemaekers MAHL, Ruigrok ANV, Hermans EJ, Kenemans JL, Baas JMP
(2010) Prefrontal mechanisms of fear reduction after threat offset. Biol Psychiatry
68:1031–1038.

Krath J, Schürmann L, von Korflesch HFO (2021) Revealing the theoretical basis of
gamification: A systematic review and analysis of theory in research on gamification,
serious games and game-based learning. Comput Hum Behav 125:106963.

Kruschke JK (2014) Doing Bayesian data analysis: A tutorial with R, JAGS, and Stan.
Academic Press.

Krypotos A-M, Vervliet B, Engelhard IM (2018) The validity of human avoidance paradigms.
Behav Res Ther 111:99–105.

Lenth R, Singmann H, Love J, Buerkner P, Herve M (2018) Emmeans. R Package Version
115-15 34:216–221.

Lissek S, Orme K, Mcdowell DJ, Johnson LL, Luckenbaugh DA, Baas JM, Cornwell BR,
Grillon C (2007) Emotion regulation and potentiated startle across affective picture
and threat-of-shock paradigms. Biol Psychol 76:124–133.

Livermore JJA, Klaassen FH, Bramson B, Hulsman AM, Meijer SW, Held L, Klumpers F, de
Voogd LD, Roelofs K (2021) Approach-Avoidance Decisions Under Threat: The Role
of Autonomic Psychophysiological States. Front Neurosci 15.

Lojowska M, Gladwin TE, Hermans EJ, Roelofs K (2015) Freezing promotes perception of
coarse visual features. J Exp Psychol Gen 144:1080–1088.

Makowski D, Ben-Shachar M, Lüdecke D (2019) bayestestR: Describing Effects and their
Uncertainty, Existence and Significance within the Bayesian Framework. J Open
Source Softw 4:1541.

McDermott TJ, Berg H, Touthang J, Akeman E, Cannon MJ, Santiago J, Cosgrove KT,
Clausen AN, Kirlic N, Smith R, Craske MG, Abelson JL, Paulus MP, Aupperle RL
(2022) Striatal reactivity during emotion and reward relates to approach–avoidance

conflict behaviour and is altered in adults with anxiety or depression. J Psychiatry
Neurosci 47:E311–E322.

McDonald AJ (1998) Cortical pathways to the mammalian amygdala. Prog Neurobiol
55:257–332.

McFadyen J, Liu Y, Dolan RJ (2023) Differential replay of reward and punishment paths
predicts approach and avoidance. Nat Neurosci 26:627–637.

Mechias M-L, Etkin A, Kalisch R (2010) A meta-analysis of instructed fear studies:
Implications for conscious appraisal of threat. NeuroImage 49:1760–1768.

Mertens G, Boddez Y, Sevenster D, Engelhard IM, De Houwer J (2018) A review on the
effects of verbal instructions in human fear conditioning: Empirical findings,
theoretical considerations, and future directions. Biol Psychol 137:49–64.

Mikl M, Mareček R, Hluštík P, Pavlicová M, Drastich A, Chlebus P, Brázdil M, Krupa P (2008)
Effects of spatial smoothing on fMRI group inferences. Magn Reson Imaging 26:490–
503.

Miller SM, Marcotulli D, Shen A, Zweifel LS (2019) Divergent medial amygdala projections
regulate approach-avoidance conflict behavior. Nat Neurosci 22.

Moughrabi N, Botsford C, Gruichich TS, Azar A, Heilicher M, Hiser J, Crombie KM,
Dunsmoor JE, Stowe Z, Cisler JM (2022) Large-scale neural network computations
and multivariate representations during approach-avoidance conflict decision-making.
NeuroImage 264:119709.

Mumford JA, Poline JB, Poldrack RA (2015) Orthogonalization of regressors in fMRI models.
PLoS ONE 10:1–11.

Murty DVPS, Song S, Surampudi SG, Pessoa L (2023) Threat and Reward Imminence
Processing in the Human Brain. J Neurosci 43:2973–2987.

O'Doherty JP (2004) Reward representations and reward-related learning in the human
brain: insights from neuroimaging. Curr Opin Neurobiol 14:769–776.

Park SQ, Kahnt T, Rieskamp J, Heekeren HR (2011) Neurobiology of Value Integration:
When Value Impacts Valuation. J Neurosci 31:9307–9314.

Patrick F, Kempton MJ, Marwood L, Williams SCR, Young AH, Perkins AM (2019) Brain
activation during human defensive behaviour: A systematic review and preliminary
meta-analysis. Neurosci Biobehav Rev 98:71–84.

Pedersen ML, Ironside M, Amemori KI, McGrath CL, Kang MS, Graybiel AM, Pizzagalli DA,
Frank MJ (2021) Computational phenotyping of brain-behavior dynamics underlying
approach-avoidance conflict in major depressive disorder. PLoS Comput Biol 17:1–
21.

Pessoa L (2010) Emotion and cognition and the amygdala: From "what is it?" to "what's to be
done?" Neuropsychologia 48:3416–3429.

Pessoa L (2023) How many brain regions are needed to elucidate the neural bases of fear
and anxiety? Neurosci Biobehav Rev 146:105039.

Piray P, den Ouden HEM, van der Schaaf ME, Toni I, Cools R (2017) Dopaminergic
Modulation of the Functional Ventrodorsal Architecture of the Human Striatum. Cereb
Cortex 27:485–495.

Poser BA, Versluis MJ, Hoogduin JM, Norris DG (2006) BOLD contrast sensitivity
enhancement and artifact reduction with multiecho EPI: Parallel-acquired
inhomogeneity-desensitized fMRI. Magn Reson Med 55:1227–1235.

Price JL (2003) Comparative aspects of amygdala connectivity. Ann N Y Acad Sci 985:50–
58.

Pujara MS, Philippi CL, Motzkin JC, Baskaya MK, Koenigs M (2016) Ventromedial Prefrontal
Cortex Damage Is Associated with Decreased Ventral Striatum Volume and
Response to Reward. J Neurosci 36:5047–5054.

Rehbein MA, Kroker T, Winker C, Ziehfreund L, Reschke A, Bölte J, Wyczesany M,
Roesmann K, Wessing I, Junghöfer M (2023) Non-invasive stimulation reveals
ventromedial prefrontal cortex function in reward prediction and reward processing.
Front Neurosci 17:1219029.

Rolls ET, Huang C-C, Lin C-P, Feng J, Joliot M (2020) Automated anatomical labelling atlas
3. NeuroImage 206:116189.

Sailer M, Hense JU, Mayr SK, Mandl H (2017) How gamification motivates: An experimental
study of the effects of specific game design elements on psychological need
satisfaction. Comput Hum Behav 69:371–380.

Schlund MW, Brewer AT, Magee SK, Richman DM, Solomon S, Ludlum MD, Dymond S
(2016) The tipping point: Value differences and parallel dorsal-ventral frontal circuits
gating human approach-avoidance behavior. NeuroImage 136:94–105.

Schultz W (2006) Behavioral Theories and the Neurophysiology of Reward. Annu Rev
Psychol 57:87–115.

Shackman AJ, Fox AS (2016) Contributions of the Central Extended Amygdala to Fear and
Anxiety. J Neurosci 36:8050–8063.

Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD (2012) Decoding Subject-Driven
Cognitive States with Whole-Brain Connectivity Patterns. Cereb Cortex 22:158–165.

Sieveritz B, Raghavan RT (2021) The Central Thalamus: Gatekeeper or Processing Hub? J
Neurosci 41:4954–4956.

Smith R, Kirlic N, Stewart JL, Touthang J, Kuplicki R, Khalsa SS, Feinstein J, Paulus MP,
Aupperle RL (2021a) Greater decision uncertainty characterizes a transdiagnostic
patient sample during approach-avoidance conflict: a computational modelling
approach. J Psychiatry Neurosci 46:E74–E87.

Smith R, Kirlic N, Stewart JL, Touthang J, Kuplicki R, McDermott TJ, Taylor S, Khalsa SS,
Paulus MP, Aupperle RL (2021b) Long-term stability of computational parameters
during approach-avoidance conflict in a transdiagnostic psychiatric patient sample.
Sci Rep 11:1–13.

Smith R, Lavalley CA, Taylor S, Stewart JL, Khalsa SS, Berg H, Ironside M, Paulus MP,
Aupperle R (2023) Elevated decision uncertainty and reduced avoidance drives in

depression, anxiety and substance use disorders during approach–avoidance
conflict: a replication study. J Psychiatry Neurosci 48:E217–E231.

Spielberger CD, Gorusch RL, Lushene RE (1970) Manual for the State-Trait Anxiety
Inventory. Palo Alto, CA: Consulting Psychologists Press.

Talmi D, Dayan P, Kiebel SJ, Frith CD, Dolan RJ (2009) How Humans Integrate the
Prospects of Pain and Reward during Choice. J Neurosci 29:14617–14626.

Warlow SM, Naffziger EE, Berridge KC (2020) The central amygdala recruits
mesocorticolimbic circuitry for pursuit of reward or pain. Nat Commun 11:2716.

Yacubian J, Gläscher J, Schroeder K, Sommer T, Braus DF, Büchel C (2006) Dissociable
systems for gain- and loss-related value predictions and errors of prediction in the
human brain. J Neurosci 26:9530–9537.

Zhou K, Xu H, Lu S, Jiang S, Hou G, Deng X, He M, Zhu Y (2022) Reward and aversion
processing by input-defined parallel nucleus accumbens circuits in mice. Nat
Commun 13:6244.

Zorowitz S, Rockhill AP, Ellard KK, Link KE, Herrington T, Pizzagalli DA, Widge AS,
Deckersbach T, Dougherty DD (2019) The Neural Basis of Approach-Avoidance
Conflict: A Model Based Analysis. eneuro 6:1–12.