The genomic consequences and persistence of sociality in spiders

Jilong Ma[1,2*], Jesper Bechsgaard[2], Anne Aagaard[2], Palle Villesen[1]. Trine Bilde[2*$], Mikkel Heide Schierup[2*$]

1. Bioinformatics Research Centre, Aarhus University, DK-8000 Aarhus C, Denmark
2. Department of Biology, Aarhus University, DK-8000 Aarhus C, Denmark

* Corresponding authors: aujilongm@birc.au.dk, trine.bilde@bio.au.dk, mheide@birc.au.dk
$ shared last authorship

1 Abstract

In cooperatively breeding social animals, a few individuals account for all reproduction. In some taxa,

sociality is accompanied by a transition from outcrossing to inbreeding, in concert, these traits act to

reduce effective population size, potentially rendering transitions to sociality 'evolutionarily

dead-ends'. We addressed this hypothesis in a comparative genomic study in spiders, where sociality

has evolved independently at least 23 times, but social species are recent and evolutionarily

short-lived. We present genomic evidence for the evolutionary dead-end hypothesis in three

independent transitions to sociality in the spider genus *Stegodyphus*. We sequenced, assembled and

annotated high-quality, chromosome-level reference genomes from three pairs of closely related social

and subsocial *Stegodyphus* species. Genome sizes range from 2.65 Gb to 3.32 Gb with high synteny,

and we identify 10,065 single-copy orthologous genes. We timed the divergence between the social

and subsocial species pairs to be from 1.3 to 1.8 million years. Social evolution in spiders involves a

shift from outcrossing to inbreeding and from equal to female-biased sex ratio, causing severe

reductions in effective population size and decreased efficacy of selection. Based on analyses of

purifying selection, we determined whether transitions to sociality co-occurred with divergence. We

show that transitions to sociality only had full effect on purifying selection at 119 kya (95CI: 71 kya

-169 kya), 260 kya (95CI: 231 ky - 289 kya) and 279 kya (95CI: 230 kya - 332 kya) respectively, and

follow remarkably similar convergent trajectories of progressive loss of diversity and shifts to an

increasingly female-biased sex ratio. This almost deterministic genomic response to sociality may

explain why social spider species do not persist. What causes species extinction is not clear, but could

be either selfish meiotic drive that eliminates the production of males, or an inability to retain genome

integrity in the face of extremely reduced efficacy of selection.

Introduction

The evolution of cooperative breeding, where some individuals forego their own reproduction to become helpers that assist other individuals in raising the offspring, is widespread in the animal kingdom (Rubenstein and Abbot 2017; Alexander 1974). In the most extreme cases, as in eusocial insects, strict division of labour results in highly complex societies with one or a few reproducing queens and countless non-reproducing helpers that are further organised into specialised sterile worker castes that each have specific tasks to perform(Crespi and Yanega 1995). In most other cooperatively breeding taxa, individuals can switch between roles of reproducer and helper depending on age, hierarchy and ecological conditions (Alexander 1974; Koenig et al. 1992; Ross and Keller 1995; Faulkes et al. 1997; Lubin and Bilde 2007). Cooperative breeding has evolved independently in divergent taxa including insects, arachnids, birds, mammals and fish. Common to all social cooperative species is that the evolution of cooperative breeding occurs in family groups and that helpers gain inclusive fitness benefits by assisting related individuals in reproducing, highlighting the importance of relatedness and kin selection in the evolution of sociality(Hamilton 1964).

As only a subset of individuals accounts for all reproduction in the social group, cooperative breeding invariably implies a reduction in effective population size (Settepani et al. 2017; Romiguier et al. 2014; Nomura and Takahashi 2012). Most cooperatively breeding species maintain a predominantly outcrossing mating system governed by pre-mating or sex-biased dispersal that takes place prior to mating. However, in a subset of species, the transition to sociality is accompanied by a switch from outcrossing to strict inbreeding through the elimination of pre-mating dispersal and reproduction among highly related group members. These taxa show a 'social syndrome' characterised by convergent independent transitions to cooperative breeding, regular inbreeding, reproductive skew (i.e. a subset of adult females reproduce), and a primary female biased sex ratio(Bilde and Lubin 2011; Settepani et al. 2017; Vanthournout et al. 2018). This combination of traits has evolved convergently multiple times in spiders (Lubin and Bilde 2007), thrips (Chapman et al. 2000), aphids

(Johnson et al. 2000), beetles (Keller et al. 2011) and possibly also mole rats(Reeve et al. 1990). In concert, these traits further act to reduce effective population sizes, and this is predicted to have negative consequences for population genetic diversity and the efficacy of selection through elevated genetic drift (Charlesworth and Wright 2001; Charlesworth 2003; Bechsgaard et al. 2019).

Spiders are an evolutionarily old group (>300 million years) (Garwood et al. 2016), and signs of social behavior were reported from fossil evidence in 100 million-year-old amber (Poinar and Buckley 2012). Although sociality in spiders is rare, the combination of cooperative breeding and inbreeding mating system has evolved independently at least 23 times (Lubin and Bilde 2007) in 6 spider families, constituting ~23 of more than 50,000 extant spider species (World Spider Catalog). Phylogenetic analyses show that social spider lineages are evolutionarily young as they are found at the terminal branches. Furthermore, there are no described cases of speciation following transitions to sociality (Lubin and Bilde 2007; Agnarsson et al. 2006), in contrast to the social insects, where relatively few transitions to sociality are followed by rich diversifications (Danforth 2002; Moreau et al. 2006). These phylogenetic patterns substantiate the hypothesis that the joint transitions to sociality and inbreeding may represent an "evolutionary dead-end"(Takebayashi and Morrell 2001; Settepani et al. 2017), where social transitions are driven by ecological benefits, while long term costs of repeated inbreeding cause the demise of social lineages.

Sociality in spiders involves living in communal nests where they cooperate in web building, prey capture, and offspring care (Avilés 2021). Although there is a degree of behavioral differentiation (Settepani et al. 2013), they do not show eusociality like in ants and bees (Crespi and Yanega 1995), as there are no casts or strict division of labor (Lubin and Bilde 2007). Phylogenetic relationships indicate that sociality in spiders evolves from subsocial ancestors (Kullmann 1972; Avilés 2010; Agnarsson et al. 2006). In subsocial species, females provide extensive maternal care for their offspring over a prolonged period after hatching (Lubin and Bilde 2007; Grinsted et al. 2014), and the juveniles of subsocial spider species spend the first weeks in the natal nest before they disperse to establish a solitary life (Bilde et al. 2005). While both social and subsocial species show elaborate

maternal care of the offspring, subsocial species maintain premating dispersal and consequently an outcrossing mating system (Bilde et al. 2005). In contrast, sociality is associated with the complete elimination of premating dispersal, which results in obligatory inbreeding as siblings reproduce within the group (Lubin and Bilde 2007). This is accompanied by the evolution of female biassed sex ratios (Vanthournout et al. 2018; Bechsgaard et al. 2019), cooperative breeding where female helpers provide extensive allomaternal care (Junghanns et al. 2017, 2019) and reproductive skew (Lubin and Bilde 2007). The convergent evolution of traits that constitute the 'social syndrome' and the phylogenetic relationships between social and subsocial species makes this system ideal for understanding causes and consequences of transitions to sociality and inbreeding.

In the spider genus *Stegodyphus*, which contains more than 25 species, sociality has evolved three times independently (Johannesen et al. 2007; Settepani et al. 2016), and each social species has a closely related subsocial sister-species (see Fig 1A) suggesting that sociality evolved recently (Bechsgaard et al. 2019; Settepani et al. 2016). The social *Stegodyphus* species express the "social syndrome" characteristics, which are predicted to reduce their effective population sizes dramatically (Bechsgaard et al. 2019) which increases the strength of drift and reduces the efficacy of natural selection (Kimura et al. 1963). This substantiates the question of whether consequences of low Ne impacts on the evolutionary persistence of social lineages. Reports of substantially reduced neutral genetic diversity (Settepani et al. 2017) and elevated rates of nonsynonymous to synonymous substitution rates (Mattila et al. 2012; Settepani et al. 2016; Bechsgaard et al. 2019, 2022) in social *Stegodyphus* species are consistent with reduced effective population sizes. Repeated inbreeding over evolutionary time is likely to have resulted in purging but also elevated rates of fixation of segregating recessive deleterious alleles, this is supported by the absence of inbreeding depression and evidence for outbreeding depression (lower hatching success when individuals from isolated populations were mated) in one of the social species *S. dumicola* (Berger-Tal et al. 2014).

Here we performed whole genome assembly and comprehensive comparative analyses of the three social *Stegodyphus* species and their subsocial sister-species to assess the timing of evolutionary

transitions and genomic consequences of social evolution. Based on analyses of Ne trajectories and

purifying selection, we determined the divergence time of lineages and the age of individual

transitions to sociality, to assess whether speciation coincides with the evolution of sociality. This

information further serves to investigate the progression of genomic consequences caused by the

transition to sociality via comparisons of social spider lineages of different ages. These data sets

enable us to assess the hypothesis that transitions to sociality are evolutionary dead-ends in spiders.

Results

***De novo* assembly and genome annotation**

Six species of *Stegodyphus* (3 social, 3 subsocial) were sequenced to ~30X coverage using Pacbio HiFi long reads. Combining this with Hi-C contact maps based on 30X Illumina sequencing, we generated chromosome-level, *de novo* assemblies. (Table 1, Supplementary Figure S1). The sex chromosomes of each species were identified by the read coverage of male individual alignments. We find two sex chromosomes in all species except for the outgroup species *S. lineatus* which has three X chromosomes (Figure S8). For each species, the inferred number of linkage groups matched the number of chromosomes found by cytological visualisation (Avilés et al. 1999; Král et al. 2011). We evaluated possible assembly errors by comparing the independent assemblies by dot plots using dotPlotly (Supplementary Figure S4-S6). For the most closely related species pairs (1-3% divergent at the genome level), assemblies were essentially syntenic, suggesting that the rate of assembly error is very low and also that the synteny breaks for more divergent species pairs indicate real chromosomal rearrangements and/or inversions. Annotation using BRAKER2 (Brůna et al. 2021) independently for each assembly revealed similar numbers of protein-coding genes ranging from 32,093 to 35,724, except for *S. sarasinorum* where 47,066 protein-coding genes were annotated. This recent gene number expansion in *S. sarisinorum* is not an artefact of annotation and is the focus of a separate study. BUSCO (Simão et al. 2015) analyses show high completeness of the genomes ranging from 91.1% to 93.3%. The GC content ranges from 33.26% to 34.18%, and the proportion of the genomes constituting repeats ranges from 57.17% to 65.20%. We found 10,065 single-copy orthologous genes present in all six species using our *de novo* assemblies, synteny information and annotations (Table 1, Table S2). Mercury (Rhie et al. 2020) estimation on our genomes reveal k-mer completeness ranging from 97.65% to 99.41% and a consensus quality QV ranging from 62.54 to 73.86, which is higher than the Vertebrate Genomes Project (VGP) benchmarking recommendations of QV > 40 and k-mer completeness > 95% (Rhie et al. 2021). When incorporating the *de novo* assembled transcriptome

from *S. africanus* and short reads DNA sequences from *S. pacificus*, our final dataset ends up with 2,303 and 348 reliable single copy orthologs for autosomes and X chromosomes, respectively, across all 8 species.

Species	Chrom_Num	Sex_chrom	Scaffolded Length	GC level	RepeatMasked	Gene_Num	BUSCO (S.C)
S.lineatus	23	X1X2X3O	2658234335	33.26%	58.72%	32093	94.10%
S.mimosarum	16	X1X2O	2910110412	33.92%	57.17%	35480	92.40%
S.tentoriicola	14	X1X2O	2984342755	33.42%	64.59%	35725	91.30%
S.dumicola	14	X1X2O	3146466829	34.18%	64.53%	34096	91.00%
S.bicolor	16	X1X2O	2984548140	33.78%	61.98%	35096	92.30%
S.sarasinorum	16	X1X2O	3321123892	34.06%	65.20%	47066	93.10%

Table 1 Assembly and annotation statistics. BUSCO completeness is only reported for the single complete fraction checked with protein sequences from annotated genes, complete BUSCO results reported in Table S2.

High gene synteny and few chromosomal rearrangements in the *Stegodyphus* genus

To investigate genome rearrangements across *Stegodyphus* from the accurate *de novo* assemblies, we used GENESPACE (Lovell et al. 2022) to retrieve the pair-wise synteny blocks based on the gene order of orthologs found in each *de novo* annotation (Figure 1 outlines the results). In general, synteny blocks suggest few chromosomal translocations and further support that the assemblies are accurate.

We identify a number of large-scale structural rearrangements including splits and joins of chromosomes unique to certain lineages of the phylogeny. For example, the outgroup species *S. lineatus* has 23 chromosomes including three X chromosomes, while the other species contain 14 to 16 chromosomes and only two X chromosomes, suggesting that multiple chromosome splits or joins have happened in the *S. lineatus* lineage, or that chromosome joins have occurred after the split with

S. lineatus in the common ancestor of the other species (Table 1). We find that the third X chromosome in *S. lineatus* appears to originate from a "cut and paste" non-homologous recombination process, which joins a part of the ancestral X chromosome X1 and a part of the ancestral X chromosome X2 (Fig 1. D). Thus, the three X chromosomes in the *S. lineatus* lineage have almost the same gene complement as the two X chromosomes in the other species. In addition, we find three chromosome join events assigned to the ancestral branch of *S. dumicola* and *S. tentoriicol* (chromosome 1, 2, and 3 are homologous to 6 separate chromosomes in other species), and one chromosome join event (chromosome 2 are homologous to 2 separate chromosomes in other species) and a non-homologous recombination event (chromosome 3 contain extra components that are homologous to another chromosome in other species) assigned to the ancestral branch of *S. bicolor* and *S. sarasinorum.*

Figure 1. (A) Phylogeny of the spider genus *Stegodpyhus* with social species in red. The divergence time is estimated from d_S of autosomal single-copy orthologous genes and a yearly mutation rate of 5e-09 per site per year. (B) Sampling locations of the species included in our study. The gene synteny plots among the six *de novo* chromosome-level assemblies are shown using GENESPACE, separately for autosomes (C) and X chromosomes (D). The length of chromosomes in the gene synteny plot represents the number of genes. The id of chromosomes in (C) and (D) are labelled as the number we assigned in the assembled genome of each species. Stars(*) mark chromosomes that are complementary-reversed in the synteny analysis.

Estimating the beginning and end of the transitions to sociality

We first estimated divergence times at each node of the *Stegodyphus* phylogeny using the resampled branch-wise d_S from autosomal branches and assuming a mutation rate of 5e-09 per site per generation (Wang and Obbard 2023). Among the three pairs of social and subsocial sister species, the estimated divergence times are 1.78±0.22 million years between *S. mimosarum* and *S. africanus*, 1.31±0.18 million years between *S. dumicola* and *S. tentoriicola*, and 1.38±0.14 million years between *S. sarasinorum* and *S. pacificus*.

Transitions to sociality in spiders are associated with a shift to inbreeding and reduced effective population size (Ne) causing natural selection to be less efficient. We assume that a substantial less efficient selection and Ne reduction in the social spider species are more likely caused by evolution of sociality than other factors. We then used our genome data to estimate the time when these transitions to sociality occurred from the expected signals in purifying selection and effective population size (Ne). We did this in two separate ways, first from the estimated changes in purifying selection estimated from d_N/d_S ratios, and second, from the inferred changes in effective population size through time estimated from coalescent analyses.

Intensity of purifying selection: We estimated the transition time to social behaviour from d_N/d_S ratios on the terminal branches of each social/subsocial species pair based on the following assumptions: a) d_N/d_S ratios of the social species was the same as for the subsocial species until social transitions on the terminal branches leading to the social species, b) after transition to sociality, the d_N/d_S ratio increased due to less efficient purifying selection, c) this 'social d_N/d_S', can be approximated by the pi_N/pi_S ratio estimated between divergent and isolated extant populations of each social species (see Methods). This d_N/d_S analysis suggests that the effects of a transition to sociality on purifying selection are very recent (Figure 2 A-D). For example, the d_N/d_S over the entire *S. mimosarum* branch is 0.1259±0.0044, and for its subsocial sister-species *S. africanus* d_N/d_S is 0.0927±0.0007. The d_N/d_S ratio since sociality evolved is estimated at 0.3043±0.0018 comparing two *S. mimosarum* individuals from Madagascar and mainland Africa. From the divergence time estimates for this species pair, this generates an estimated time since transition to sociality of *S. mimosarum* of 279 kya (95% CI: 230

kya - 332 kya). Using the same calculation, the social d_N/d_S ratio of *S.dumicola* is estimated to be

0.3366±0.0037 by comparing individuals from Otavi North Namibia and Karasburg South Namibia.

Combining the d_N/d_S ratio being 0.1776±0.0039 and 0.1615±0.0054 for the *S. dumicola* and *S.*

tentoriicola branches respectively, the estimation of the transition to socialility is 119 kya (95% CI:

72 kya - 170 kya) in *S. dumicola*.

Figure 2. The intensity of purifying selection was assessed using d_N/d_S from the coding sequences of

2302 single-copy orthologous genes. (A) The locations of the Sri Lanka *S.sarasinorum* and Himalayas

S.sarasinorum samples, along with their pairwise d_N/d_S estimations for the social period. (B) The

locations of the Namibia South *S.dumicola* and Namibia North *S.dumicola* samples, with

corresponding pairwise d_N/d_S estimations for the social period. (C) The locations of the South Africa

S.mimosarum and Madagascar *S.mimosarum* samples, and their pairwise d_N/d_S estimations for the

social period. (D) Branch-wise d_N/d_S estimations using PAML, with the 95% confidence interval from

resampling. Red, blue, and grey denote social species terminal branches, subsocial species terminal

branches, and internal branches, respectively. The d_N/d_S estimation, indicated by points with round

and triangle denote autosomes and X chromosomes respectively, is aligned to the phylogeny that corresponds to the respective branches. The phylogeny in (D) provides an indication of the topology without branch length information.

For the social *S. sarasinorum,* we performed two separate estimations by comparing *S. sarasinorum* to the subsocial species *S. pacificus,* which diverged 1.38±0.14 million years ago, and the subsocial species *S. bicolor*, which diverged 2.67±0.19 million years ago. In *S. sarasinorum*, we estimated the social d_N/d_S ratio to be 0.2864±0.0023 based on two individuals from Himalaya and Sri Lanka. The d_N/d_S ratio on the terminal branch of *S. sarasinorm* is 0.1349±0.0009. Compared to the 0.0996±0.0016 of the *S. pacificus* branch and the 0.1211±0.0007 of the *S. bicolor* branch, the final estimation for the transition to sociality of *S. sarasinorum* is 260 kya (95% CI: 231 kya - 289 kya) and 223 kya (95% CI: 199 kya - 247 kya), respectively.

We have attributed the observed higher d_N/d_S ratio in the terminal branches of social species to a decrease in purifying selection. However, increased adaptive selection acting on social species could have the same effect. RELAX analyses (Wertheim et al. 2015) suggest that the elevated d_N/d_S ratios in social species is almost entirely due to relaxed selection in autosomes (Figure S3), as has previously been suggested (Tong et al. 2022). We also show elevated d_N/d_S ratios on X chromosomes compared to autosomes across all branches in the phylogeny, except for the terminal branches of *S. pacificus*. Such elevated d_N/d_S ratios can be attributed to a decrease in purifying selection acting on X chromosomes, which is consistent with the expected outcome of reduced Ne in X chromosomes compared to autosomes (Ne_X/Ne_Auto is expected to be 0.75 in diploid species with equal sex ratio). Alternatively, but not mutually exclusive, an adaptive explanation for elevated d_N/d_S ratios on the X chromosomes is the so-called faster-X effect (Charlesworth et al. 1987). Recessive variants on the X chromosomes are directly exposed to selection in males that are haploid for X chromosomes, and under certain conditions this will lead to more adaptive substitutions on X chromosomes compared to autosomes.

Additionally, we note that our implication of using pi_N/pi_S between divergent populations of the same species could potentially lead to a systematically higher "social d_N/d_S" due to the unfixed polymorphisms rather than reflect the actual selection strength when the selection strength acts differently on synonymous polymorphism sites and non-synonymous polymorphism sites within species. We benchmarked our strategy by comparing the site frequency spectrum of missense variants (N-sites) and synonymous variants (S-sites) in *S. dumicola* built from 9 individuals. Site frequency spectrums of the N-sites and S-sites reveal no visually significant difference in shape, indicating genetic drift being the major force acting on both N-sites and S-sites (Figure S7). Mattila et al 2012 also quantified the d_N/d_S and pi_N/pi_S in three *Stegodyphus* species, including one social species (*S. mimosarum)* and two subsocial species (*S. tentoriicola* and *S. lineatus*) (Mattila et al. 2012). Mattila et al. reveal that the increase of pi_N/pi_S compared to d_N/d_S is much higher in the social spider species (3-fold) than subsocial species (1-1.5 fold)(Mattila et al. 2012). It is also worth noting that the pi_N/pi_S in Mattila et al 2012 is based on transcriptome data from a single individual, where the unfixed polymorphisms will increase the tendency of inherently higher pi_N/pi_S than d_N/d_S at a larger effect than using isolated populations. Thus, we conclude that pi_N/pi_S to a good approximation can be used as a proxy for the d_N/d_S ratio after sociality ("social d_N/d_S").

Coalescent analysis: Next, we used PSMC (Li and Durbin 2011) analysis to reconstruct the effective population sizes and the sex ratio in the populations over time by analysing autosomes and X chromosomes separately in each species. For the social species, which are all highly inbred, a diploid genome was created by merging two haploid genomes sampled from different divergent and isolated populations, i.e. an artificial diploid. This allows coalescent events further back in time to be inferred. The transition time to sociality is then estimated by graphically determining the time period when the effective population size of the social species became severely reduced (Figure 3). Compared to subsocial species, the population trajectories of all the social species share a gradual reduction of Ne from more than 300,000 to less than 50,000, but starting and ending at different time points within the past 900 ky. In the social species *S. dumicola*, the Ne is reduced from 300,000 around 420 kya to around 50,000 at 300 kya, and even lower today. In the social species *S. mimosarum*, the Ne is

1 reduced from 1.4 million to 57,500 in the time interval between 900kya and 350kya. The reduction of

2 Ne in the social species *S. sarasinorum* starts 600 kya at 800,000 to less than 30,000 around 300 kya.

7 Figure 3. The historical effective population size inferred from the Pairwise Sequentially Markovian

8 Coalescent (PSMC) model, for different *Stegodyphus* species with chromosome-level assembly. In

9 each panel, the left figure depicts PSMC results, while the right figure displays the effective

10 population size ratio between X chromosomes and autosomes. Each species' trajectory is shown with

thicker lines for autosomes and thinner lines for X chromosomes. The type of species are distinguished by colour: red for social species and blue for subsocial species. Red shadows mark the estimated social transition period based on the decreasing effective population size in social species. Vertical lines indicate estimated time points by comparing the social species with its closest sister subsocial counterpart. Red lines correspond to the estimated social transition time, derived from relaxed selection in social species, while blue lines represent the estimated divergence time between a social species and its closest subsocial sister species. Solid lines provide point estimates, and dashed lines delineate the 95% confidence intervals. The PSMC results are truncated in the near recent past for better comparison across species. Bootstrapped PSMC results without time truncation are provided in Figure S9. (A) Displays data for the social species *S.sarasinorum* and the subsocial species *S.bicolor*. Estimates for social transition and divergence time are based on a comparison between *S.sarasinorum* and its subsocial sister species, *S.pacificus*. (B) Presents findings from a comparison between the social species *S.dumicola* and its subsocial sister species *S.tentoriicola*. (C) Shows data for the social species *S.mimosarum*, with social transition and divergence time estimates based on a comparison with its subsocial sister species, *S. africanus*. Note that we do not have an effective population size estimation for *S. africanus* due to missing whole genome data. (D) Presents historical effective population size for the outgroup subsocial species *S.lineatus*.

Since the PSMC analyses of the social species are made from pseudodiploid genomes combined from separate genetically isolated populations, they can additionally provide an estimate of the time of separation of these populations, since coalescence can not occur when they are separate and the Ne estimate will thus be very large (Figure S9). We observe such an uptick in Ne in recent times in the three social species, and estimate that the two *S. mimosarum* populations from Madagascar and South Africa diverged around 200 kya, the two *S. dumicola* populations in North and South Namibia diverged 20 kya, and the two *S. sarasinorum* populations from Himalaya and Sri Lanka diverged approximately 110 ky ago.

The pseudo-diploid of social species collapsed two heavily inbred diploid individuals. We estimate the run out of heterozygosity (ROH) regions takes 59.83%-86.27% of their genomes (Table S4), but the individuals are still not 100% homozygous. Hence, they are not identical as a haploid in practice. Thus, we can not rule out the fact that pseudo-diploid actually capture more than two haploids for part of the genomes outside the ROH region. However, we expect this will only increase a local coalescence rate that is randomly distributed in the genome, resulting in a lower estimation of Ne overall, without biassing the Ne reduction information retrieved from the Ne trajectories.

Evolution of sex ratio bias

The sex determination system of spiders is X0, and as there are four autosomes for each three X chromosomes in the population with 1-to-1 sex ratio, the effective population size of X chromosomes is expected to be 75% of the autosomes. In subsocial species, PSMC analyses of X chromosomes and autosomes showed that although the estimated Ne for X chromosomes is indeed lower, the ratio of X chromosomes to autosome effective population size is around 50-60% instead of the expected 75%. This can be due to a lower mutation rate and/or stronger selection on the X chromosomes, as was previously suggested based on diversity estimates on X chromosomes and autosomes in *S. africanus* *(Bechsgaard et al. 2019)*. In contrast to the subsocial species, the social species show strongly female biassed primary sex ratios (Lubin and Bilde 2007) and this has been shown to be genetically determined (Vanthournout et al. 2018). When a species evolves from an unbiased to a female biassed sex ratio, the effective population size of X chromosomes should approach that of autosomes. In concordance with this, the PSMC analyses reveal that X chromosome and autosome Ne are almost identical in the most recent time intervals of the PSMC, as expected when sociality has fully evolved. The evolution of a female biassed sex ratio can therefore also be timed using PSMC analyses under the assumption that it occurred after the transition to sociality, by comparing the Ne of X chromosomes and autosomes back in time. We found elevated Ne(X)/Ne(autosomes) ratios in all social species associated with reductions in Ne, which links the evolution of sex-ratio bias with social transitions. This suggests that

the evolution of a female biassed sex ratio in the three social species occurred early in the

evolutionary transition to sociality.

The evolution of female bias is predicted under strong continuous inbreeding and lack of pre-mating

dispersal, characteristics of social spider species and other social arthropods (Johnson et al. 2000;

Chapman et al. 2000; Keller et al. 2011), to reduce competition among brothers for fertilisation of

their sisters (Bulmer and Taylor 1980; West 2009). Furthermore, allocation to daughters as the helping

sex in social species is expected by the local resource enhancement model (Trivers and Willard 1973).

Biassed sex ratios are often caused by sex chromosome meiotic drive (Lindholm et al. 2016), and the

driving loci are selfish genetic elements that would usually trigger adaptations to restore even sex

ratio (Lindholm et al. 2016). However, selection against female bias is not expected here, because of

the adaptive benefits of female biassed sex allocation (Vanthournout et al. 2018) generating

predictions of rapid fixation of the sex-linked meiotic driver. If stronger female bias keeps being

adaptive, selection will lead to an increasingly female biassed sex ratio following transition to

sociality. The results from the PSMC analyses showing that NeX/NeA continues to increase after

transition to sociality suggest that stronger female bias continues to be adaptive. Furthermore, our

estimation for the transition to sociality matches with observations of increasing female bias with age

of sociality in the three species. The oldest social species *S. mimosarum* has the strongest sex ratio

bias with 9% males, the younger social species *S. dumicola* and *S. sarasinorum* have a male

percentage of 12% and 8.5-33%, respectively (Lubin and Bilde 2007). The correlation between sex

ratio bias and age of the social period in the three transitions to sociality therefore indicate that the

continued loss of males is genetically determined.

Figure 4. Estimated species divergence (Td), and the time of transition to sociality/cooperative breeding (indicated in red) in three independent social transitions within the *Stegodyphus* spider genus. Hypothetical speciation time (Ts) between each species pair is marked as the dashed lines. The scale of the plots is symbolic and not proportional. The widths of the terminal branches represent variation in effective population size, with red symbolising social periods and blue indicating subsocial periods. Light blue branches represent species without chromosome-level assemblies for PSMC. For each lineage where a social transition occurs, point estimates and 95% confidence intervals for social transition times are presented on the left, while social transition periods derived from population size history are displayed on the right. In PSMC estimations, the start of social transition periods corresponds to the initial decrease in effective population size, while the end coincides with Ne dropping below 50,000.

Discussion

Divergence times and transition to sociality

To elucidate genomic consequences of repeated, independent transitions to sociality, we estimated species divergence times and the time of speciation using different approaches (Figure 4). An obvious question is whether the speciation itself was caused by one of two diverging populations evolving sociality. The large time interval between estimates of species divergence (Td = 1.3-1.8 my, Figure 4) and the beginning of social transitions (400-900 ky) as derived from strong reductions in Ne at first suggests that species separation took place first with subsequent evolution of sociality and cooperative breeding. However, coalescent theory predicts that it takes 2Ne generations on average for ancestral polymorphisms to find most common ancestry. Since the effective population sizes of subsocial species are very large (300,000-600,000), divergence times (Td in Figure 4) are expected to be much more ancient than the species separation times (Ts in Figure 4) due to long coalescence times in the ancestral species. With a generation time of one year, average coalescence time in the ancestral species are between 600,000 and 1.2 million years, and our best estimate of the speciation times (Ts in Figure 4) are therefore almost 1 million years closer to the present than the divergence times. This corresponds to the time when we estimate the beginning of the social transitions to (from PSMC results), thus suggesting that social behaviour emerging in a population of a species may facilitate separation of this population into a new species.

There is, however, also some evidence from the social *S. dumicola* that sociality can arise independently after the speciation process. *Stegodyphus dumicola* and *S. sarasinorum* diverged from their closest subsocial sister species at a similar time scale of 1.30 Mya and 1.37 Mya respectively. The ancestral Ne for *S. dumicola* (300,000) is smaller than the *S.sarasinorum* (600,000). Thus, the speciation time of *S. dumicola* (Ts2 in Figure 4) is expected to be closer to the species divergence time than the speciation time of *S. sarsinroum* (Ts1 in Figure 4) based on coalescence theory (Figure S11). Meanwhile, both analysis from PSMC and d_N/d_S show that the age of sociality is younger in *S. dumicola* than *S. sarasinorum*. The combination of the earlier speciation time and the younger age of

sociality in *S. dumicola* compared to *S. sarasinorum* suggests that sociality can arise as a separate event from the initiation of speciation. Nevertheless, evidence for separation of the two events in *S. dumicola* could be confounded by other factors, including (a) a missing subsocial sister species (unidentified or extinct) to *S. dumicola* that is not included in the analysis, which can bias the analyses to overestimate the divergence time for *S. dumicola* lineage, (b) observed reduction of Ne in recent past of *S. tentoriicola* from PSMC (Figure S9) and elevated d_N/d_S in *S. tentoriicola* compared to other subsocial species (Figure 2), which can bias the analyses to underestimate the age of sociality in *S. dumicola*. Nevertheless, the combination of data presented above substantiates that sociality and characteristics of the 'social syndrome' evolved much more recently than the divergence estimate.

High extinction risk of social spiders

The genomic consequences of transitions to sociality in spiders are consistent with expectations of strong reductions in Ne when the evolution of sociality is accompanied by a transition from outcrossing to inbreeding and the development of a female biased sex ratio (Vanthournout et al. 2018). The relatively young age of social lineages indicate that sociality evolves rapidly and frequently in the spider phylogeny and comes with a subsequent strong signal of decreased purifying selection caused by strict inbreeding and much reduced effective population size. Overall, there are at least 23 independent convergent origins of sociality (Lubin and Bilde 2007; Agnarsson et al. 2006). The recurring and rapid evolution of cooperative breeding, coupled with the absence of lineages with a long history of sociality, is strong indirect evidence that sociality in spiders represents an evolutionary impasse, unlikely to persist beyond a timespan of 1 millions years once established. Thus, it is plausible that numerous social spider species have emerged and gone extinct throughout evolutionary history, but we can only observe the presence of extant and recently developed social spider species.

The risk of extinction in social spiders is hypothesised to be much higher than in eusocial insects owing to the 'social syndrome' which causes reductions in Ne. Phylogenetic studies show that diversification of species is only associated with a few old transitions to eusociality in the order *Hymenoptera*, i.e. ants, bees and wasps(Wcislo and Fewell 2017; Hunt and Toth 2017; Heinze et al.

2017), which is different from the repeated and recent convergent evolution of sociality in the spider phylogeny (Agnarsson et al. 2006; Settepani et al. 2016). We propose that a key factor contributing to the difference in longevity of social lineages among major taxa is the elimination of pre-mating dispersal which results in an obligatory inbreeding mating system. In eusocial insects, new colonies are established by mated queens, and outcrossing is maintained by pre-mating dispersal of reproductive individuals (Wcislo and Fewell 2017; Heinze et al. 2017). Thus, eusocial insects may show reduced Ne and relaxed selection compared to their nonsocial counterparts (Kapheim et al. 2015), but the reduction in Ne reduction is mitigated by the maintenance of outcrossing. In contrast, we here show that transitions to sociality in independently evolved spider lineages are associated with dramatic declines in Ne.

Chromosomal systems, mating systems and the evolutionary longevity of social lineages

Some of the most successful social lineages have evolved in the *Hymenoptera*, where the entire clade is haplodiploid with haploid males and diploid females (Grimaldi and Engel 2005). These lineages are evolutionarily old, for example in ants where eusociality is dated at least to the Cretaceous (Table 2) and they have since diversified dramatically both in species and abundance to become one of the most ecologically dominating taxonomic groups in the world (Wilson and Hölldobler 2005). The evolutionary success of haplodiploid social lineages, which is also found in *Thysanoptera* (thrips) and sporadically in some spider mites, *Hemiptera*, and *Coleoptera*, has long pointed to an importance of haplodiploidy in the evolution of sociality (Hamilton 1964; Trivers and Hare 1976; Joshi and Wiens 2023). However, phylogenetic patterns are debated and perhaps inconclusive, as eusociality has evolved repeatedly within the haplodiploid *Hymenoptera* and *Thysanoptera*, whereas other successful independently evolved social lineages are diploid, such as in termites and aphids.

A key question that arises from our study is whether haplodiploidy represents a more robust system for governing the evolutionary longevity of social lineages. Termites and *Hymenoptera* represent the

oldest social lineages and represent both chromosomal systems (Table 2). These taxa also maintain predominantly outcrossing mating systems, which acts to maintain higher effective population sizes that are less exposed to genetic drift. In taxa where social evolution is accompanied by a transition to inbreeding, the diploid social spiders show the most short-lived and recent transitions (Settepani et al. 2017), whereas thrips and beetles with a haplodiploid system may have relatively older social lineages (Table 2). In combination, these patterns suggest that whether or not sociality is associated with a transition from outcrossing to inbreeding is perhaps the most important factor for determining their evolutionary longevity. The taxa that evolve inbreeding mating systems are simultaneously characterised by female biased sex ratio which further acts to reduce effective population size (Bechsgaard et al. 2019). Our study provides genomic support for the hypothesis that the independent evolution of sociality accompanied by an inbreeding mating system and female-biased sex ratios results in drastic reductions in Ne and relaxed efficacy of selection, which likely contribute to drive lineage extinctions.

Ploidy	Mating system	Age of social lineages	Social Complexity*(Fewell and Abbot 2018)	Taxonomy range	Taxa examples
Diploid	Strictly inbreeding (sib-mating)	Young (Sociality only has full impact on selection within the past 300ky from this study)	Cooperative communal groups	Broad, often multiphyletic for the young social species lineages	Spiders(Agnarsson et al. 2006; Settepani et al. 2016)
	Outcrossing	Various? (oldest can be 130 million year in termites(Krishna et al. 2013))	Advanced eusocial, Primitive eusocial, Cooperative communal groups		Termites(Legendre et al. 2008; Korb and Thorne 2017), Snapping shrimps(Chak and Rubenstein 2019), Mole rats(Kverková et al. 2018; Faulkes and Bennett 2021), Aphids(Stern 1994)

Haplodiploid	Inbreeding (sib-mating), occasionally outcrossing	Less Old? (10 million years in thrips, in term of species lineage(Abbot and Chapman 2017))	Primitive eusocial, Cooperative communal groups	Limited (*order Hymenoptera* and *Thysanoptera, tribe Xyleborini*), often monophyletic for the old social species lineage.	Thrips(Chapman et al. 2000), Beetles(Johnson et al. 2018)
	Outcrossing	Old (The phylogeny of ants dates back to 140 - 168 million years ago(Moreau et al. 2006))	Advanced eusocial, Primitive eusocial		Ants, bees, wasps

Table 2. Summary of the characteristics of sociality in diploid and haplodiploid organisms with

different mating systems. *The social complexity is a broad qualitative summary with the terminology

used for describing social diversity in insects in Fewell and Abbot 2018.

Genomics determinants of evolutionary dead ends in social spiders

We note that even though the three social transitions we report were initiated at different times, they

have followed remarkably similar trajectories in population size reduction, increased d_N/d_S, and biased

sex ratios that complements the strong similarities in behaviour and life history associated with

transitions to sociality. This almost deterministic genomic response to sociality may also hold the key

to understanding why social spider species do not seem to persist for very long. Several possibilities

may explain the demise. These include a) loss of adaptive potential due to loss of genetic diversity, b)

fixation of deleterious variation, c) loss of males, and d) loss of genome integrity. While a) and b) may

definitely make sociality a doomed evolutionary strategy, it is questionable whether these processes can

happen within a few hundred thousand years since the fixation of slightly deleterious variants occurs

over a very large number of generations (Lynch et al. 1995). Loss of males could carry on until

population collapse, and it is intriguing that we see fewer males in the social transition estimated to be

the oldest (Figure 4). Finally, loss of genome integrity by, for example transposable elements (TEs),

increased mutation rate, poorer genomic repair of damage, and double strand breaks, are all global

mechanisms that could impact the longevity of the social species. TEs are expected to accumulate in

species with reduced Ne. Although the question of whether the TE fraction increases with reduced Ne

remains inconsistent in several tested organisms (Glémin et al. 2019), evidence for increased amounts

of TEs has been seen in eusocial snapping shrimps with reduced Ne (Chak et al. 2021). This idea warrants future research in social spiders.

Methods

Sample collection across the *Stegodyphu*s genus

The spider genus *Stegodyphus* is distributed widely across Eurasia and Africa and we collected seven species including three social and four subsocial species in this study (Supplementary Table 1). Samples were collected from multiple regions for social species (Fig. 1 A). We collected *S. dumicola* samples from Otavi and Betta in Namibia (2017), *S. mimosarum* samples from Antanario in Madagascar and Weenen in South Africa (2011), and *S. sarasinorum* samples from the Himalaya region and Sri Lanka (2014). For subsocial species, we collected *S.lineatus* from the Negev Desert in Israel (2020), *S. bicolor* from Betta in Namibia (2021), *S. tentoriicola* from Tierberg in South Africa (2015). *Stegodyphus africanus* RNA data was previously published in Bechsgaard et al (2019).

Reference genome: DNA extraction, sequencing and assembly

DNA was extracted from single individuals using the MagAttract HWM DNA kit from Qiagen (Hilden, Germany). Sequencing was done by PacBio HiFi technology to obtain from 69 to 99 GB data.We assembled haplotype resolved contigs using Hifiasm(Cheng et al. 2021) with Pacbio HiFi reads and Hi-C reads. Hi-C reads were aligned to the contigs from the longer haplotype of each species using Juicer pipeline for scaffolding chromosome-level assemblies using 3D-DNA pipeline (Dudchenko et al. 2017). The chromosome-level assemblies were further manually checked and curated using Juicebox Assembly Tools(Durand et al. 2016) based on consistency of Hi-C contact pattern.

Resequencing: DNA extraction and sequencing

DNA was extracted from single individuals using the Qiagen Blood and Tissue kit, Qiagen (Hilden, Germany), and the DNA was sequenced using DNBSEQ-G400.

Genome Annotation

We annotated six species with chromosome-level assemblies using BRAKER2 with RNA-seq and protein homology evidence combined. For each species, we used RepeatModeler2(Flynn et al. 2020) to generate a repeat library and combine it with the repeat library for Arthopoda from Repbase(Bao et al. 2015). The chromosome-level assemblies are masked using Repeatmasker(Tarailo-Graovac and Chen 2009) and their corresponding repeat libraries. We used STAR (Dobin et al. 2013) to align RNA-seq reads to the repeat-masked genome as transcriptome evidence for gene prediction in BRAKER2. For protein homology evidence, we used protein sequence from a previous *S.dumicola* annotation (Liu et al. 2019) and arthropoda protein from OrthoDB v10 (Kriventseva et al. 2019). In the end, the completeness of the genome annotations is evaluated with BUSCO together with a set of 1013 *Arthropoda* genes from OrthoDB v10.

Ortholog groups finding and multiple genome alignments

We performed GENESPACE to check the gene synteny among the six hifi chromosome-level assemblies. During the GENESPACE analysis, OrthoFinder (Emms and Kelly 2019) was applied to identify orthologous groups across the six species. The 10065 single-copy orthologous groups out of 48143 orthologous groups identified from the six species were further used in evolutionary analysis.

We do not have the genome assembled from *S. africanus*. To include *S. africanus* in the analyses, we downloaded the transcriptome data of *S. africanus* from Bechsgaard et.al 2019 (Bechsgaard et al. 2019) and used Trinity (Haas et al. 2013) to assemble transcripts. We used DIAMOND(Buchfink et al. 2021) blastx mode to align the transcripts to the database of all the translated amino acid sequences from the 10065 single-copy orthologous groups of all six species. 5590 out of 10065 single-copy orthologous groups find one or more hits from the trinity transcriptome assemblies. We further check the aligned percentage of the hitted orthologous genes and filtered for a percentage of 75% to consider a transcript from *S. africanus* being a valid match to a certain single-copy ortholog groups identified, which ended up with 2649 single-copy orthologous groups.

We also do not have a genome assembled from *S. pacificus*. To include *S. pacificus* genes into the 2649 single-copy orthologous groups identified, we first used bwa-mem2 (Vasimuddin et al. 2019) to aligned short-read DNA sequence from a *S. pacificus* individual to the *S.sarasinorum* reference genome, which is the closest sister species of it. Then we used SAMtools (Danecek et al. 2021) consensus to call the consensus sequence as the genome of *S. pacificus* based on the bam file. The sequences of 2649 ortholog genes in *S. pacifcus* are then retrieved using the genome annotation file of *S. sarasinorum*. This process can be challenging when there are indels found in *S. pacificus* compared to *S. sarasinorum* reference when we construct the consensus sequence of *S. pacificus*. The sequence of *S. pacificus* and *S. sarasinorum* will not be in alignment as they have different total lengths with different genome coordinate systems. In practice, we fill gaps for deletions and remove insertions in *S. pacificus* based on the reference of *S. sarasinorum* when we construct the consensus sequences. This practice is achieved by setting parameters for SAMtools being "consensus --show-del yes --show-ins no". By ignoring the indels in such a way, we maintain the same coordinate systems between the *S. pacificus* and *S. sarasinorum*. Orthologous sequences in S. pacificus can be then retrieved directly using the genome annotation file of *S. sarasinorum* to build multiple sequence alignments across all 8 species for d_N/d_S estimations.

Branch-wise d_N/d_S estimation

We used CodeML in PAML(Yang 2007) to estimate d_N and d_S in branch-wise mode based on the 2649 single-copy orthologous groups. The 2649 orthologous groups were split up into an autosome set containing 2302 genes and a X chromosome set containing 347 genes. We randomly sample 500 genes from the 2302 autosomal without replacement in each resampling run, and we run the resampling independently 500 times to have an estimation of d_N/d_S of each branch in the phylogeny with confidence intervals. For X chromosomes, we randomly sample 100 genes independently 100 times instead.

We do codon aware alignment in gene-wise for d_N/d_S estimation. We used alignSequence from

MACSE (Ranwez et al. 2018) to align coding sequences of each gene from all the species excluding

S.pacificus, which was attached to the alignment according to the *S.sarasinorum* in the end. We also

examined and filtered each gene alignment for local mis-alignment. The size of each alignment block

without gap should have a length for at least 300 nucleotide base pairs as well as a fraction of

polymorphism sites under 15% (Figure S2). The passing alignment blocks of the randomly chosen

genes were concatenated using GOalign (Lemoine and Gascuel 2021) for d_N/d_S estimation.

d_N/d_S (empirical pi_N/pi_S) estimation for social species

We used CodeML in PAML to estimate a pairwise d_N/d_S for each social species with two separate

populations. We aligned one individual from different populations to the species reference separately

using BWA-MEM2. After calling the SNP using platypus (Rimmer et al. 2014), we retrieve the

alternative nucleotide to substitute the species reference as the reference genome of each

sub-population. Sequences from the 2649 single-copy orthologous genes were retrieved from the

sub-population references for pairwise d_N/d_S estimation. We used the same 500 resampled sets of

genes as used in the branch-wise d_N/d_S estimation before to estimate the pi_N/pi_S with 95% confidence

interval between separate social species populations for autosomes.

d_N/d_S and estimation of speciation divergence and social transition

For the divergence time between lineages, we used the dS branch length, a constant mutation rate of

5e-09 per site per generation, and a generation time of one year to scale the d_S length into years. We

use the average d_S branch length between a split as an estimation of divergence time except for the

split between *S.sarasinorum* and *S.pacificus*. The exception is due to the underestimation of d_S length

of *S. pacificus* lineage from ignoring of indels, thus we use the d_S length of *S.sarasinorum* branch

alone for divergence time estimation between *S.sarasinorum* and *S.pacificus*.

Assuming the transition to sociality is an instant process. The d_N/d_S estimated for the social species

terminal branch is an average of the subsocial period d_N/d_S and social period d_N/d_S weighted by the

length of each period on that branch. The same idea has been applied to estimate the age of selfing in *Arabidopsis thaliana* compared to selfing incompatible *Arabidopsis lyrata (Bechsgaard et al. 2006)*.

In our case, the subsocial period can be substituted approximately with the d_N/d_S for the branch of the subsocial sister species. The social period d_N/d_S can be estimated by pairwise comparing two individuals from two isolated and divergent populations of the social species.

For each resampling run of autosomal genes (500 genes out of 2303 genes), we obtained four estimations for each pair of social species and its subsocial counterpart using PAML, include

    (a) d_N/d_S of the subsocial lineage

    (b) d_N/d_S of the social lineage

    (c) pi_N/pi_S between two isolated populations of social species

    (d) dS branch length between social and subsocial species

The distributions of the d_N/d_S and pi_N/pi_S from the resampling process for each species pair are provided in Figure S10. Confidence intervals for mean estimations of d_N/d_S and pi_N/pi_S are retrieved using the 2.5% -97.5% quantile from a normal distribution built according to the mean and standard error. The point estimation for species divergence time is estimated from the mean of value (d) from the 500 rounds of resampling. Thus the 95% confidence interval of species divergence time is retrieved using the 2.5% -97.5% quantile from a normal distribution built according to the mean and standard deviation of dS. For the time of transition to sociality, we sampled 10000 sets of observations, each of which consists of a set of values including one observation of (a), (b), (c) and (d) sampled from their distribution built by previous resampling process of 500 random autosomal genes. We then solved the social transition time for each sampled observation set. The point estimation for transition time is obtained from the mean of 10000 sampled calculations. The 95% confidence interval for transition time is retrieved as 2.5%-97.5% quantile from the empirical distribution of 10000 rounds. This will account for all uncertainty of the genome-wide selection strength and species divergence time estimated using the resampling process.

Under the assumption of our calculation for the social period fraction, the social transition time is estimated as the time point when the selection becomes as relaxed as what we observe today between the isolated populations of social species. This level of relaxation in selection intensity would be viewed as the last signal to arise after the completion of social transitions. The social transition times estimated with our d_N/d_S calculation reflecting sociality have fully evolved based on its impact on purifying selection, despite the start of social transition may be earlier than the estimations.

$$t_2 = T_1 \times \frac{(\omega_{sub-social} - \omega_{sub})}{(\omega_{social} - \omega_{sub})}$$

Figure M1. Illustration for calculating the social period length for current social species as an instant process. t_2 denotes the length of the social period. T_1 denotes the divergence time between social species and its sister subsocial species. ω_{sub} denotes the d_N/d_S estimated from the subsocial species branch in branch-wise PAML estimations. $\omega_{sub-social}$ denotes the d_N/d_S estimated from the social species branch in branch-wise PAML estimations. ω_{social} denotes the d_N/d_S (In practice, pi_N/pi_S) estimated from the pairwise PAML estimations of two separate social species populations.

PSMC estimation of social transition time

We used PSMC to reconstruct the historic effective population size for all the six species with a chromosome-level genome. The PSMC method relies on a single diploid individual, where the social species runs out of heterozygosity due to inbreeding. We combine the DNA-seq of two individuals from two separate and diverged populations to retain the loss of heterozygosity.

We benchmarked the fraction of running out of heterozygosity region in the predicted youngest social species *S. dumicola*. We used the GATK pipeline to genotype 9 individuals (5 females and 4 males) from *S. dumicola* and only kept bi-allelic SNP variants among the individuals with a genotype quality filter of 30. We did a sliding window heterozygosity with 100kb window size and 10kb step size on each individual. Windows of an individual without any variants are further classified as run out of homozygosity regions and summarised into ROH fraction by individual (Table S4).

We ran the PSMC method on autosomes and X chromosomes separately. The generation time is fixed to 1 year per generation. The mutation rate used for autosomes is 5e-09 per site per generation and the rate for X chromosomes is scaled to 3.8e-09 with a factor of 0.76 based on the piX/piA being 0.57 instead of 0.75 in subsocial species *S. africanus (Bechsgaard et al. 2019)*. To align the estimated historical Ne estimated between autosomes and X chromosomes, we retrieve all the time points with an estimation of Ne either for autosomal or X chromosomes, the missing value is filled with the linear interpolate according to the closest values on both sides. Then we calculated the historical NeX/NeA as an indicator for evolution of sex-ratio bias. In the end, the curve of NeX/NeA was smoothed using a spline fit in R. The period of accomplished social transition by retrieving the time interval where the decrease of Ne in social species slows down.

Testing for selection relaxation

Based on the alignment of coding sequences for all filtered autosomal genes, we use HyPhy RELAX to detect the relaxation of intensification of selection on the social species branches. We ran the RELAX test with four different 'test - reference' pair settings for the 2303 autosomal genes and 247 X

chromosome genes separately. We have one pair that uses all social species terminal branches as the 'test' and all subsocial species terminal branches as the 'reference' to evaluate the overall effect on selection from the social transition. We have three pairs including each terminal branch of a social species as the 'test' and its sister subsocial species terminal branches as well as common ancestor branches as the 'reference' to test for relaxed selection in each social transition (Supplementary Figure S3).

Data access

All of the raw sequencing data was uploaded to NCBI Bioproject PRJNA994315.

The assembled genomes are available in NCBI with the under the following genome accession numbers:

Stegodyphus dumicola: CP160598-CP160611

Stegodyphus tentoriicola: CP160584-CP160597

Stegodyphus bicolor: CP160568-CP160583

Stegodyphus sarasinorum: CP160513-CP160528

Stegodyphus mimosarum: CP160529-CP160544

Stegodyphus lineatus: CP160545-CP160567

Codes for analysis pipeline is available through GitHub:

https://github.com/Jilong-Jerome/sociality-in-spiders-dead-end

Competing interest statement

The authors declare no competing interests.

Acknowledgement

We thank the support from Independent Research Fund Denmark (Grant number: 0135-00201B). We thank Yael Lubin, Tharina Bird, Virginia Settepani, Lena Grinsted and Clemence Rose for help in collecting samples included in this study.

Author contribution:

Jilong Ma: Analysis design, data analysis, manuscript writing.
Jesper Bechsgaard: Analysis design, manuscript writing.
Anne Aagaard: Analysis design, manuscript refining.
Palle Villesen: Analysis design, manuscript refining.
Trine Bilde: Experiment design, manuscript writing
Mikkel Heide Schierup: Analysis design, manuscript writing, supervision.

References

Abbot P, Chapman T. 2017. Sociality in Aphids and Thrips. In *Comparative Social Evolution* (eds. D.R. Rubenstein and P. Abbot), pp. 154–187, Cambridge University Press, Cambridge.

Agnarsson I, Avilés L, Coddington JA, Maddison WP. 2006. Sociality in theridiid spiders: repeated origins of an evolutionary dead end. *Evolution* **60**: 2342–2351.

Alexander RD. 1974. The Evolution of Social Behavior. *Annu Rev Ecol Syst* **5**: 325–383.

Avilés L. 2010. Causes and consequences of cooperation and permanent-sociality in spiders. In *The Evolution of Social Behavior in Insects and Arachnids* (eds. J.C. Choe and B.J. Crespi), pp. 476–498, Cambridge University Press, Cambridge.

Avilés L. 2021. Social Spiders. In *Encyclopedia of Social Insects* (ed. C.K. Starr), pp. 858–868, Springer International Publishing, Cham.

Avilés L, Varas C, Dyreson E. 1999. Does the African social spider Stegodyphus dumicola control the sex of individual offspring? *Behav Ecol Sociobiol* **46**: 237–243.

Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mob DNA* **6**: 11.

Bechsgaard J, Jorgensen TH, Jønsson AK, Schou M, Bilde T. 2022. Impaired immune function accompanies social evolution in spiders. *Biol Lett* **18**: 20220331.

Bechsgaard JS, Castric V, Charlesworth D, Vekemans X, Schierup MH. 2006. The transition to self-compatibility in Arabidopsis thaliana and evolution within S-haplotypes over 10 Myr. *Mol Biol Evol* **23**: 1741–1750.

Bechsgaard J, Schou MF, Vanthournout B, Hendrickx F, Knudsen B, Settepani V, Schierup MH, Bilde T. 2019. Evidence for Faster X Chromosome Evolution in Spiders. *Mol Biol Evol* **36**: 1281–1293.

Berger-Tal R, Tuni C, Lubin Y, Smith D, Bilde T. 2014. Fitness consequences of outcrossing in a social spider with an inbreeding mating system. *Evolution* **68**: 343–351.

Bilde T, Lubin Y. 2011. Group living in spiders: cooperative breeding and coloniality. In *Spider Behaviour* (ed. M.E. Herberstein), pp. 275–306, Cambridge University Press, Cambridge.

Bilde T, Lubin Y, Smith D, Schneider JM, Maklakov AA. 2005. The transition to social inbred mating systems in spiders: role of inbreeding tolerance in a subsocial predecessor. *Evolution* **59**: 160–174.

Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. 2021. BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. *NAR Genom Bioinform* **3**: lqaa108.

Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* **18**: 366–368.

Bulmer MG, Taylor PD. 1980. Sex ratio under the haystack model. *J Theor Biol* **86**: 83–89.

Chak STC, Harris SE, Hultgren KM, Jeffery NW, Rubenstein DR. 2021. Eusociality in snapping shrimps is associated with larger genomes and an accumulation of transposable elements. *Proceedings of the National Academy of Sciences* **118**: e2025051118.

Chak ST, Rubenstein DR. 2019. Social transitions in sponge-dwelling snapping shrimp. *Curr Opin Insect Sci* **34**: 33–39.

Chapman TW, Crespi BJ, Kranz BD, Schwarz MP. 2000. High relatedness and inbreeding at the origin of eusociality in gall-inducing thrips. *Proceedings of the National Academy of Sciences* **97**: 1648–1650.

Charlesworth B, Coyne JA, Barton NH. 1987. The Relative Rates of Evolution of Sex Chromosomes and Autosomes. *Am Nat* **130**: 113–146.

Charlesworth D. 2003. Effects of inbreeding on the genetic diversity of populations. *Philos Trans R Soc Lond B Biol Sci* **358**: 1051–1070.

Charlesworth D, Wright SI. 2001. Breeding systems and genome evolution. *Curr Opin Genet Dev* **11**: 685–690.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nat Methods* **18**: 170–175.

Crespi BJ, Yanega D. 1995. The definition of eusociality. *Behav Ecol* **6**: 109–115.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools. *Gigascience* **10**. http://dx.doi.org/10.1093/gigascience/giab008.

Danforth BN. 2002. Evolution of sociality in a primitively eusocial lineage of bees. *Proc Natl Acad Sci U S A* **99**: 286–290.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol I, Lander ES, Aiden AP, et al. 2017. De novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds. *Science* **356**: 92–95.

Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL. 2016. Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. *Cell Syst* **3**: 99–101.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol* **20**: 238.

Faulkes CG, Bennett NC. 2021. Social evolution in African mole-rats - A comparative overview. *Adv Exp Med Biol* **1319**: 1–33.

Faulkes CG, Bennett NC, Bruford MW, O'brien HP, Aguilar GH, Jarvis JUM. 1997. Ecological constraints drive social evolution in the African mole–rats. *Proceedings of the Royal Society of London Series B: Biological Sciences* **264**: 1619–1627.

Fewell J, Abbot P. 2018. *Sociality*. Oxford University Press.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020. RepeatModeler2 for automated genomic discovery of transposable element families. *Proc Natl Acad Sci U S A* **117**: 9451–9457.

Garwood RJ, Dunlop JA, Selden PA, Spencer ART, Atwood RC, Vo NT, Drakopoulos M. 2016. Almost a spider: a 305-million-year-old fossil arachnid and spider origins. *Proc Biol Sci* **283**: 20160125.

Glémin S, François CM, Galtier N. 2019. Genome Evolution in Outcrossing vs. Selfing vs. Asexual Species. *Methods Mol Biol* **1910**: 331–369.

Grimaldi D, Engel MS. 2005. *Cambridge evolution series: Evolution of the insects*. Cambridge University Press, Cambridge, England.

Grinsted L, Breuker CJ, Bilde T. 2014. Cooperative breeding favors maternal investment in size over number of eggs in spiders. *Evolution* **68**: 1961–1973.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, et al. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc* **8**: 1494–1512.

Hamilton WD. 1964. The genetical evolution of social behaviour. II. *J Theor Biol* **7**: 17–52.

Heinze J, Kellner K, Seal J. 2017. Sociality in Ants. In *Comparative Social Evolution* (eds. D.R. Rubenstein and P. Abbot), pp. 21–49, Cambridge University Press, Cambridge.

Hunt JH, Toth AL. 2017. Sociality in Wasps. In *Comparative Social Evolution* (eds. D.R. Rubenstein and P. Abbot), pp. 84–123, Cambridge University Press, Cambridge.

Johannesen J, Lubin Y, Smith DR, Bilde T, Schneider JM. 2007. The age and evolution of sociality in Stegodyphus spiders: a molecular phylogenetic perspective. *Proc Biol Sci* **274**: 231–237.

Johnson AJ, McKenna DD, Jordal BH, Cognato AI, Smith SM, Lemmon AR, Lemmon EM, Hulcr J. 2018. Phylogenomics clarifies repeated evolutionary origins of inbreeding and fungus farming in bark beetles (Curculionidae, Scolytinae). *Mol Phylogenet Evol* **127**: 229–238.

Johnson PC, Llewellyn KS, Amos W. 2000. Microsatellite loci for studying clonal mixing, population structure and inbreeding in a social aphid, Pemphigus spyrothecae (Hemiptera: pemphigidae). *Mol Ecol* **9**: 1445–1446.

Joshi CH, Wiens JJ. 2023. Does haplodiploidy help drive the evolution of insect eusociality? *Front Ecol Evol* **11**: 1118748.

Junghanns A, Holm C, Schou MF, Overgaard J, Malte H, Uhl G, Bilde T. 2019. Physiological Adaptations to Extreme Maternal and Allomaternal Care in Spiders. *Frontiers in Ecology and Evolution* **7**. https://www.frontiersin.org/articles/10.3389/fevo.2019.00305.

Junghanns A, Holm C, Schou MF, Sørensen AB, Uhl G, Bilde T. 2017. Extreme allomaternal care and unequal task participation by unmated females in a cooperatively breeding spider. *Anim Behav* **132**: 101–107.

Kapheim KM, Pan H, Li C, Salzberg SL, Puiu D, Magoc T, Robertson HM, Hudson ME, Venkat A, Fischman BJ, et al. 2015. Social evolution. Genomic signatures of evolutionary transitions from solitary to group living. *Science* **348**: 1139–1143.

Keller L, Peer K, Bernasconi C, Taborsky M, Shuker DM. 2011. Inbreeding and selection on sex ratio in the bark beetle Xylosandrus germanus. *BMC Evol Biol* **11**: 359.

Kimura M, Maruyama T, Crow JF. 1963. THE MUTATION LOAD IN SMALL POPULATIONS. *Genetics* **48**: 1303–1312.

Koenig WD, Pitelka FA, Carmen WJ, Mumme RL, Stanback MT. 1992. The evolution of delayed dispersal in cooperative breeders. *Q Rev Biol* **67**: 111–150.

Korb J, Thorne B. 2017. Sociality in Termites. In *Comparative Social Evolution* (eds. D.R. Rubenstein and P. Abbot), pp. 124–153, Cambridge University Press, Cambridge.

Král J, Kořínková T, Forman M, Krkavcová L. 2011. Insights into the meiotic behavior and evolution of multiple sex chromosome systems in spiders. *Cytogenet Genome Res* **133**: 43–66.

Krishna K, Grimaldi DA, Krishna V, Engel MS. 2013. Treatise on the Isoptera of the world: Volume 1 introduction. *Bull Am Mus Nat Hist* **377**: 1–200.

Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, Zdobnov EM. 2019. OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for evolutionary and functional annotations of orthologs. *Nucleic Acids Res* **47**: D807–D811.

Kullmann EJ. 1972. Evolution of social behavior in spiders (Araneae; Eresidae and Theridiidae). *Am Zool* **12**: 419–426.

Kverková K, Bělíková T, Olkowicz S, Pavelková Z, O'Riain MJ, Šumbera R, Burda H, Bennett NC, Němec P. 2018. Sociality does not drive the evolution of large brains in eusocial African mole-rats. *Sci Rep* **8**: 9203.

Legendre F, Whiting MF, Bordereau C, Cancello EM, Evans TA, Grandcolas P. 2008. The phylogeny of termites (Dictyoptera: Isoptera) based on mitochondrial and nuclear markers: Implications for the evolution of the worker and pseudergate castes, and foraging behaviors. *Mol Phylogenet Evol* **48**: 615–627.

Lemoine F, Gascuel O. 2021. Gotree/Goalign: toolkit and Go API to facilitate the development of phylogenetic workflows. *NAR Genom Bioinform* **3**: lqab075.

Li H, Durbin R. 2011. Inference of human population history from individual whole-genome sequences. *Nature* **475**: 493–496.

Lindholm AK, Dyer KA, Firman RC, Fishman L, Forstmeier W, Holman L, Johannesson H, Knief U, Kokko H, Larracuente AM, et al. 2016. The Ecology and Evolutionary Dynamics of Meiotic Drive. *Trends Ecol Evol* **31**: 315–326.

Liu S, Aageaard A, Bechsgaard J, Bilde T. 2019. DNA Methylation Patterns in the Social Spider, Stegodyphus dumicola. *Genes* **10**. http://dx.doi.org/10.3390/genes10020137.

Lovell JT, Sreedasyam A, Schranz ME, Wilson M, Carlson JW, Harkess A, Emms D, Goodstein DM, Schmutz J. 2022. GENESPACE tracks regions of interest and gene copy number variation across multiple genomes. *Elife* **11**. http://dx.doi.org/10.7554/eLife.78526.

Lubin Y, Bilde T. 2007. The Evolution of Sociality in Spiders. In *Advances in the Study of Behavior*, Vol. 37 of, pp. 83–145, Academic Press.

Lynch M, Conery J, Burger R. 1995. Mutation Accumulation and the Extinction of Small Populations. *Am Nat* **146**: 489–518.

Mattila TM, Bechsgaard JS, Hansen TT, Schierup MH, Bilde T. 2012. Orthologous genes identified by transcriptome sequencing in the spider genus Stegodyphus. *BMC Genomics* **13**: 70.

Moreau CS, Bell CD, Vila R, Archibald SB, Pierce NE. 2006. Phylogeny of the ants: diversification in the age of angiosperms. *Science* **312**: 101–104.

Nomura T, Takahashi J. 2012. Effective population size in eusocial Hymenoptera with worker-produced males. *Heredity (Edinb)* **109**: 261–268.

Poinar G, Buckley R. 2012. Predatory behaviour of the social orb-weaver spider, Geratonephila burmanica n. gen., n. sp. (Araneae: Nephilidae) with its wasp prey, Cascoscelio incassus n. gen., n. sp. (Hymenoptera: Platygastridae) in Early Cretaceous Burmese amber. *Hist Biol* **24**: 519–525.

Ranwez V, Douzery EJP, Cambon C, Chantret N, Delsuc F. 2018. MACSE v2: Toolkit for the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. *Mol Biol Evol* **35**: 2582–2584.

Reeve HK, Westneat DF, Noon WA, Sherman PW, Aquadro CF. 1990. DNA "fingerprinting" reveals high levels of inbreeding in colonies of the eusocial naked mole-rat. *Proc Natl Acad Sci U S A* **87**: 2496–2500.

Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, Uliano-Silva M, Chow W, Fungtammasan A, Kim J, et al. 2021. Towards complete and error-free genome assemblies of all vertebrate species. *Nature* **592**: 737–746.

Rhie A, Walenz BP, Koren S, Phillippy AM. 2020. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biol* **21**: 245.

Rimmer A, Phan H, Mathieson I, Iqbal Z, Twigg SRF, WGS500 Consortium, Wilkie AOM, McVean G, Lunter G. 2014. Integrating mapping-, assembly- and haplotype-based approaches for calling variants in clinical sequencing applications. *Nat Genet* **46**: 912–918.

Romiguier J, Lourenco J, Gayral P, Faivre N, Weinert LA, Ravel S, Ballenghien M, Cahais V, Bernard A, Loire E, et al. 2014. Population genomics of eusocial insects: the costs of a vertebrate-like effective population size. *J Evol Biol* **27**: 593–603.

Ross KG, Keller L. 1995. ECOLOGY AND EVOLUTION OF SOCIAL ORGANIZATION: Insights from Fire Ants and Other Highly Eusocial Insects. *Annu Rev Ecol Syst* **26**: 631–656.

Rubenstein DR, Abbot P. 2017. *Comparative social evolution*. Cambridge University Press.

Settepani V, Bechsgaard J, Bilde T. 2016. Phylogenetic analysis suggests that sociality is associated with reduced effectiveness of selection. *Ecol Evol* **6**: 469–477.

Settepani V, Grinsted L, Granfeldt J, Jensen JL, Bilde T. 2013. Task specialization in two social spiders, Stegodyphus sarasinorum (Eresidae) and Anelosimus eximius (Theridiidae). *J Evol Biol* **26**: 51–62.

Settepani V, Schou MF, Greve M, Grinsted L, Bechsgaard J, Bilde T. 2017. Evolution of sociality in spiders leads to depleted genomic diversity at both population and species

levels. *Mol Ecol* **26**: 4197–4210.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**: 3210–3212.

Stern DL. 1994. A phylogenetic analysis of soldier evolution in the aphid family Hormaphididae. *Proc Biol Sci* **256**: 203–209.

Takebayashi N, Morrell PL. 2001. Is self-fertilization an evolutionary dead end? Revisiting an old hypothesis with genetic theories and a macroevolutionary approach. *Am J Bot* **88**: 1143–1150.

Tarailo-Graovac M, Chen N. 2009. Using RepeatMasker to identify repetitive elements in genomic sequences. *Curr Protoc Bioinformatics* **Chapter 4**: 4.10.1–4.10.14.

Tong C, Avilés L, Rayor LS, Mikheyev AS, Linksvayer TA. 2022. Genomic signatures of recent convergent transitions to social life in spiders. *Nat Commun* **13**: 6967.

Trivers RL, Hare H. 1976. Haploidploidy and the evolution of the social insect: The unusual traits of the social insects are uniquely explained by Hamilton's kinship theory. *Science* **191**: 249–263.

Trivers RL, Willard DE. 1973. Natural selection of parental ability to vary the sex ratio of offspring. *Science* **179**: 90–92.

Vanthournout B, Busck MM, Bechsgaard J, Hendrickx F, Schramm A, Bilde T. 2018. Male spiders control offspring sex ratio through greater production of female-determining sperm. *Proc Biol Sci* **285**. http://dx.doi.org/10.1098/rspb.2017.2887.

Vasimuddin M, Misra S, Li H, Aluru S. 2019. Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems. In *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)*, pp. 314–324.

Wang Y, Obbard DJ. 2023. Experimental estimates of germline mutation rate in eukaryotes: a phylogenetic meta-analysis. *bioRxiv* 2023.01.24.525323. https://www.biorxiv.org/content/biorxiv/early/2023/05/10/2023.01.24.525323 (Accessed July 17, 2023).

Wcislo W, Fewell JH. 2017. Sociality in Bees. In *Comparative Social Evolution* (eds. D.R. Rubenstein and P. Abbot), pp. 50–83, Cambridge University Press, Cambridge.

Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. 2015. RELAX: detecting relaxed selection in a phylogenetic framework. *Mol Biol Evol* **32**: 820–832.

West S. 2009. *Sex Allocation*. Princeton University Press.

Wilson EO, Hölldobler B. 2005. The rise of the ants: a phylogenetic and ecological explanation. *Proc Natl Acad Sci U S A* **102**: 7411–7414.

World Spider Catalog WS. World spider catalog, version 21.0. *Natural History Museum Bern*.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**: 1586–1591.