Deviation from typical brain activity during naturalistic stimulation is related to personality traits

Lucia Jajcay[1,2,3], **David Tomeček**[1,2,3], **Renáta Androvičová**[1], **Iveta Fajnerová**[1], **Filip Děchtěrenko**[4], **Jan Rydlo**[1,5], **Jaroslav Tintěra**[1,5], **Jiří Lukavský**[1,4], **Jiří Horáček**[1,6], and **Jaroslav Hlinka**[1,2,✉]

[1]National Institute of Mental Health, Czech Republic
[2]Department of Complex Systems, Institute of Computer Science of the Czech Academy of Sciences, Czech Republic
[3]Department of Cybernetics, Faculty of Electrical Engineering, Czech Technical University in Prague, Czech Republic
[4]Department of Cognitive Psychology, Institute of Psychology of the Czech Academy of Sciences, Czech Republic
[5]Institute for Clinical and Experimental Medicine, Czech Republic
[6]Department of Psychiatry and Medical Psychology, Third Faculty of Medicine, Charles University, Czech Republic

The relationship between personality and brain activity has been an increasingly popular topic of neuroscientific research. However, the limitations of both personality measures and neuroimaging, as well as methodological issues, continue to pose challenges to its understanding. The naturalistic viewing condition has been shown to enhance individual differences and might, therefore, be of benefit to the endeavor. Here, we thus examine this relationship using naturalistic fMRI of 82 healthy subjects. We implemented a simple dimensionality reduction method to characterize brain activity by its 'typicality', assessed a range of personality traits using widely-used personality inventories, and tested the relationship between the two. We found that there is, indeed, a relationship between personality and the typicality of brain activity, which appears to be most consistently manifested by lower typicality in subjects with higher Neuroticism/Harm Avoidance. Our results highlight the usefulness of naturalistic viewing data for exploring the relationship between individual differences in personality and brain activity.

personality traits | NEO-FFI | TCI | naturalistic stimuli | fMRI

Correspondence: *hlinka@cs.cas.cz*

Introduction

Personality refers to a set of cognitive, emotional, and behavioral patterns characteristic of an individual (1). These patterns are shaped by a combination of biological and environmental factors and tend to be relatively consistent and stable over time (2). While there is no general agreement on a definition of personality, most definitions emphasize the role of motivation and interaction with the environment in shaping an individual's unique behavioral and emotional responses to their surroundings, and revolve around personality traits.

Measuring personality is a complex endeavor that requires reliable and valid assessment tools. Personality questionnaires, or inventories, such as the NEO Five-Factor Inventory (NEO-FFI) (3), which assesses the so-called Big Five personality traits (4), present one way of doing so (2). They are standardized measures designed to assess the personality traits of an individual by asking them to rate themselves on a series of statements. While such inventories can provide valuable insights into an individual's personality characteristics, they

are subject to a range of limitations. Similarly to the concept of personality itself, the dimensions of personality assessed by these measures are not universally agreed upon. Moreover, different inventories may emphasize different aspects of the same personality trait, leading to differences in how traits are defined and measured across studies. Personality inventories also inherently involve the potential for a number of biases, such as self-reporting, response-style, or cultural biases. However, their advantages – mainly ease of use and interpretation – appear to outweigh the limitations, and personality inventories remain the dominant method of personality assessment (2).

While traditional approaches to understanding personality have predominantly focused on behavioral and psychological perspectives, in recent years, there has been an increased interest in linking individual differences in personality to anatomical and functional brain markers. Due to its high spatial resolution, availability, and non-invasive nature, such 'personality neuroscience' studies often rely on functional magnetic resonance imaging (fMRI) (5, 6). However, fMRI analysis also involves several challenges stemming from the nature of the signal. Notably, fMRI is an indirect method. Rather than measuring neuronal activity itself, it measures the metabolic demands of active neurons. Although the two are coupled, the relationship between them is neither straightforward nor perfectly understood (7, 8). Moreover, the slowness of the hemodynamic response, together with the generally rather sparse temporal acquisition of consecutive scans, makes the temporal resolution of fMRI comparably low.

Consequently, despite general agreement that personality stems from brain function, the limitations of both personality measures and fMRI make it challenging to detect relationships between the two.

Nevertheless, a growing amount of research has successfully detected such relationships. Notably, several studies have used resting-state fMRI to relate brain functional connectivity (FC) to personality traits. FC captures statistical dependencies of the activity of spatially separated brain regions (9). It has been suggested that the FC matrix may be a useful representation of brain activity dynamics and provide behaviorally or clinically relevant markers (10, 11). FC has also

been shown to act as a unique and reliable 'fingerprint' capable of distinguishing an individual by capturing their intrinsic, characteristic connectivity profile (12). Aghajani et al. 2014 found Neuroticism and Extraversion to be associated with different amygdala resting-state functional connectivity patterns. A study of the 'neurotic brain' by Servaas et al. 2015 concluded that the brains of highly neurotic individuals have less than optimal functional network organization and show signs of functional disconnectivity. Sampaio et al. 2014 revealed correlations between each of the Big Five personality dimensions and the default mode network (DMN). Similarly, Simon et al. 2020 found Openness to be positively associated with connectivity in the DMN. In addition, they found negative associations between Neuroticism and both the ventral and dorsal attention networks and Agreeableness and the dorsal attention network. They also showed that the association between Conscientiousness and the frontoparietal network becomes stronger with age. Moreover, the relationships between personality traits and resting-state functional connectivity also appear to be gender-specific, as demonstrated by Nostro et al. 2018. Wang et al. 2018 used an exploratory approach based on independent component analysis to identify a parietal network and found the inter-subject similarity of this network to be associated with Openness. Using connectome-based predictive modeling, Hsu et al. 2018 identified networks consisting of functional connections correlated with Neuroticism and Extraversion scores, then successfully predicted these scores for a different sample of individuals.

A more recent study by Cai et al. 2020 applied a similar approach to a large Human Connectome Project (HCP) dataset and could reliably predict four of the Big Five personality traits (all but Extraversion). Also capitalizing on the large HCP database, Liu et al. 2019 demonstrated that subjects with similar personality profiles have similar whole-brain connectivity patterns. Lastly, using individual FC matrices of 884 young, healthy adults, Dubois et al. 2018 reliably predicted Openness to experience. In addition, they also performed principal component analysis (PCA) on the personality scores to derive two orthogonal higher-order personality dimensions. They could reliably predict 5% of the variance in the score of the second component (β), which loaded mostly on Openness to experience. It, therefore, appears that personality can indeed be successfully investigated using fMRI. However, as argued by Tomeček et al. 2020, due to methodological issues, the reflection of personality in the brain's intrinsic functional architecture remains elusive, and findings of such studies should be interpreted with great caution.

Note that all of the above-mentioned studies relied on resting-state functional connectivity in conjunction with the five-factor model of personality (4). The naturalistic viewing (movie-watching) condition presents an increasingly popular alternative. Since it provides rich stimulation and, simultaneously, sufficient space for spontaneous responses, it might elicit highly individual responses and inter-regional interactions. Indeed, while naturalistic stimuli make individual activation profiles more similar to one another, observable in-

dividual differences have been shown to persist and even become easier to identify (22–27).

Here, we therefore analyzed fMRI data obtained while subjects were watching movies. The more controlled nature of the naturalistic viewing condition allowed us to go beyond the FC approach commonly applied to resting-state data and focus on comparing the time courses of brain activations across subjects. We implemented a robust dimensionality reduction method to characterize brain activity during the condition by its 'typicality' and tested its relationship to personality traits. The latter were assessed using two well-established personality inventories. In addition to the widely-used NEO-FFI, we also used a second inventory based on a psychobiological model.

We expect that the naturalistic-viewing condition combined with the comparably large dataset and comprehensive assessment of personality traits will allow us to examine the elusive relationship between personality and brain activity.

Materials and Methods

Participants. Eighty-two healthy volunteers participated in the study (77 right-handed, 48 male, mean age ± SD: 30.96 ± 8.54 years). Participants were informed about the experimental procedures and provided written informed consent. The study design was approved by the local Ethics Committee of the Institute for Clinical and Experimental Medicine. Each participant underwent MRI scanning that included functional magnetic resonance imaging acquisitions during an audiovisual stimulation by segments of naturalistic stimuli (3 video segments, total length 29'45") and 10 min in the resting-state condition with eyes closed, as well as the acquisition of a T1-weighted and T2-weighted anatomical scan, and a diffusion-weighted imaging scan.

Personality inventories. The participants filled out Czech versions of two widely-used personality inventories – the NEO-FFI (3) and the Temperament and Character Inventory (TCI) (28) – providing a total of 12 scores representing different personality traits.

The NEO-FFI consists of 60 items and measures five personality traits: Neuroticism, Extraversion, Openness to experience, Agreeableness, and Conscientiousness – the so-called Big Five. The five-factor model of personality is based on the lexical hypothesis, which assumes that fundamental individual differences are represented in the natural language by trait descriptive adjectives and was proposed as a universal and comprehensive framework for describing personality (4). The TCI (240 items) is a seven-factor model of personality based on a psychobiological theory. It assesses four temperament dimensions assumed to be related to biology, heritable, and stable through the life-span: Novelty Seeking, Harm Avoidance, Reward Dependence, and Persistence; and three character dimensions affected by learning and culture, which are less heritable and mature with age: Self-Directedness, Cooperativeness, and Self-Transcendence (29, 30). Upon visual inspection, the distributions of most of the personality scores in our subject sample appear to be close to

normal (see Fig. 1), yet a formal statistical test rejected normality for two of them (see Analysis).

Fig. 1. Distributions of the 12 personality traits in our subject sample ($N = 82$), with fitting normal distributions. *Top*: NEO-FFI scores; *bottom*: TCI scores.

Note that the Big Five are not fully orthogonal (31). Similarly, certain TCI dimensions tend to correlate (29). Moreover, since the two inventories are measures of two competing theories of personality and might be measuring similar personality constructs with different scales, similarities and correlations naturally exist across the two. Most notably, Harm Avoidance (defined by Cloninger 1987 as 'a heritable tendency to respond intensely to signals of aversive stimuli, thereby learning to inhibit behavior to avoid punishment, novelty, and frustrative nonreward') appears to be highly positively correlated with Neuroticism (the tendency to experience negative emotions) and negatively correlated with Extraversion (a measure of how sociable, outgoing and energetic a person is) (30, 33).

As shown in Fig. 2, significant correlations across the 12 personality traits also exist in our sample. In particular, we found strong positive Pearson correlations between Neuroticism and Harm Avoidance ($r = 0.751$, $p < 0.001$) and Agreeableness and Cooperativeness ($r = 0.721$, $p < 0.001$) and strong negative correlations between Extraversion and Harm Avoidance ($r = -0.590$, $p < 0.001$) and Neuroticism and Self-Directedness ($r = -0.585$, $p < 0.001$). The relationships between the traits in our final sample are, therefore, in line with the findings of an analogous study mentioned above (30).

Fig. 2. Relationships between the 12 personality traits in our subject sample ($N = 82$). Pearson correlations between traits assessed by NEO-FFI (N, E, O, A, C) and TCI (NS, HA, RD, Pe, SD, Co, ST). * $p < 0.01$, ** $p < 0.001$ (uncorrected).

Stimuli. We used 3 video segments, in the following order:

1. an uninterrupted excerpt from 'The Good, the Bad and the Ugly' (16:48-26:48), a feature film by Sergio Leone (1966) which has been used in previous studies of the naturalistic viewing condition (34, 35) (further denoted as 'western', duration 10 min);

2. an excerpt from a local TV reality show 'Hledá se táta a máma' (season 3, episode 2), which showed a strong activation profile in our previous, unpublished study ('reality show', 8 min and 5 s);

3. a collection of six short clips intended to elicit a range of emotions – excerpts from mostly Hollywood movies, from the FilmStim database (36) ('emotional clips', 11 min and 40 s).

Our selection of stimuli thus consists of materials high in social content and likely to elicit strong emotions and, similarly to other naturalistic viewing studies, includes well-known Hollywood movies (23, 27).

MRI data acquisition. Scanning was performed using a Siemens TrioTim 3T MRI machine located at the Institute for Clinical and Experimental Medicine in Prague, Czech Republic. A high-resolution T1-weighted image (MP-RAGE

3D, TR = 2300 ms, TE = 4.63 ms, TI = 900 ms, flip angle = 10°, voxel size = 1×1×1 mm, FOV 256×256, PAT-factor = 2, BW = 130 Hz/pix, 224 sagittal slices) covering the entire brain was acquired for anatomical reference. Functional images were then obtained using T2*-weighted echo-planar imaging (EPI) with blood oxygenation level-dependent (BOLD) contrast. GE-EPIs (TR = 2500 ms, TE = 30 ms, flip angle = 90°, voxel size = 3×3×3 mm, FOV 192×192, BW = 1594 Hz/pix, 44 axial slices) covering the entire cerebrum were acquired continuously in ascending order. T2-weighted and diffusion-weighted images were also acquired but not used in the current study.

The video segments were concatenated into a single MP4 file without breaks between the segments. They were back-projected onto a screen using an Epson EB-G5100 projector with an ELPLM04 (Middle Throw 1) lens. The subjects viewed the 47×35 cm large screen via a mirror mounted on the head coil at a 22° visual angle from a distance of 1210 mm. The audio was delivered using MRI-compatible headphones (MR CONFON OPTIME 1), with its intensity amplified, allowing the subjects to hear it over the scanner noise. For further technical details, such as the resolution and frame rate of each video, see Hlinka et al. 2022.

Preprocessing. Preprocessing was performed using the SPM12 software package (Wellcome Centre for Human Neuroimaging, UCL) implemented in MATLAB (The Mathworks, Inc.). The preprocessing pipeline consisted of motion correction, slice timing correction, spatial normalization into standard stereotaxic space (Montreal Neurological Institute, MNI), and spatial smoothing with an 8 mm FWHM kernel.

Subsequently, two types of time series to be used in further analyses were extracted from the data. Firstly, using the GIFT toolbox, Infomax-based independent component analysis (ICA) was performed, decomposing the data into 39 independent components. The number of components was estimated using the minimum description length (MDL) criterion. The signal was band-pass filtered using the Butterworth filter (0.009-0.1 Hz).

Secondly – for an alternative, region-based processing pipeline – mean BOLD time series from the 90 regions of interest (ROIs) of the Automated Anatomical Labeling (AAL) atlas were extracted. The anatomical positions of the ROIs are described in Tzourio-Mazoyer et al. 2002. The signal was band-pass filtered using the same Butterworth filter (0.009–0.1 Hz).

Analysis.

ICA analyses. We first calculated the 'typical time series' (average across all subjects) for each of the 39 independent components. For each subject, we then correlated each of their time series with the respective typical time series. The mean of these Pearson's correlation coefficients (39 'similarity indices') presents the subject's individual 'typicality index' – a single number characterizing the typicality of their brain activity.

Using the interquartile range method, selected for its robustness, we identified three subjects with low typicality indices as outliers. To prevent extreme values from skewing the results, we removed these subjects and then repeated the analysis to obtain the final typicality indices.

Since demographic factors such as age and sex have been reported to be related to the relationship between personality traits and brain activity, we examined the relationship between the typicality indices and these two potential confounds before proceeding with the analyses (16, 17). While the typicality indices are normally distributed, the null hypothesis of a normal distribution was rejected for subject age and sex (using the Anderson-Darling test implemented in MATLAB; $p < 0.05$). We thus assessed the relationship between the typicality indices and these two variables by computing Spearman's rank correlations.

As discussed above, similarities and overlaps across the traits assessed by the two personality inventories have been shown to exist in the general population as well as in our sample. To examine the relationship between the typicality indices and personality traits, we thus reduced the dimensionality of the personality data by performing PCA. In our primary analysis, we calculated Spearman's rank correlation between the typicality indices and a subset of the principal components. The decision to use *Spearman's rank* correlation was made because, using the Anderson-Darling test implemented in MATLAB, the null hypothesis of a normal distribution was rejected for Agreeableness (NEO-FFI) and Harm Avoidance (TCI). We performed the analysis for all stimuli combined (using the entire time series corresponding to all three videos) as well as for individual stimuli (using the corresponding segment of the entire time series) to assess robustness with respect to stimulus selection.

We then performed a number of exploratory follow-up analyses. Due to their exploratory nature, the results of these analyses are presented uncorrected for multiple testing. We present them nonetheless, as they provide useful context for interpreting the main findings.

First, after observing a significant relationship between the high-level (PCA-based) personality traits and the typicality of brain activity, we explored the result in further detail by calculating Spearman's rank correlation between subjects' typicality indices and their scores on each of the twelve, albeit overlapping, personality traits. Again, we performed this analysis for all stimuli combined as well as for individual stimuli.

Next, we delved into even more detail and examined the relationship between each ICA component and the personality traits – meaning we computed Spearman's rank correlation between the latter and the 39 correlation coefficients of each subject's time series with the respective typical time series (their 'similarity indices' rather than their summarized 'typicality index').

The doubly multivariate nature of the data (with multiplicity in terms of both personality traits and ICA components) invites the application of a more advanced technique. In addition to simply computing pairwise correlations, we thus

also performed a canonical correlation analysis (39, 40). The latter is an alternative method of measuring the association among the two sets of variables – the 12 personality scores on the one hand and the 39 similarity indices on the other. It finds linear combinations of variables from the two sets (canonical variables) with maximum correlations.

AAL analyses. We repeated the analyses using the mean BOLD time series from the 90 ROIs derived from the AAL atlas rather than the ICA components. The canonical correlation analysis could not be performed on the AAL data due to rank deficiency, resulting from having more features (90 time series) than observations (81 subjects). Note that only one of the three outliers removed in the ICA analysis was removed in the AAL analysis.

Results

ICA analyses. Since neither of the correlations between subject typicality indices and subject age or sex is statistically significant (age: $\rho = -0.139$, $p = 0.220$; sex: $\rho = -0.120$, $p = 0.293$), the following results do not take age or sex into account.

The principal component analysis performed on the personality data resulted in 12 principal components. Based on Fig. 3, the first two principal components capture most of the relevant between-subject variability. We thus computed Spearman's rank correlation between the typicality indices and the *first two* principal components. Note that Fig. 4 shows that the first principal component is influenced negatively by Neuroticism and Harm Avoidance and positively primarily by Cooperativeness and Agreeableness – the two pairs of personality traits shown above to be the most strongly correlated among the 12 traits.

Fig. 3. Percentage of total variance explained by the principal components (ICA analysis). The bars show the percentage of variance explained by each component. The line plot shows the cumulative explained variance percentages. The plot only shows the first nine components that together explain 95% of the total variance. A bend in the line occurs after the first two components, which together explain more than half (55%) of the total variance. Only the first two components were therefore used in the subsequent analysis.

Fig. 4. PCA biplot showing the influence of each personality trait on the principal components (ICA analysis). The 12 variables (personality traits) are depicted as vectors with their direction and length indicating how each trait contributes to the first two principal components (depicted on the x-axis and the y-axis, respectively) – principal component *coefficients* or loadings. In addition, the 79 observations (participants) are represented as points with their coordinates indicating each participant's *scores* for the two principal components. The points are scaled with respect to the maximum score value and maximum coefficient length.

We found the typicality of brain activity during the experimental condition to be positively correlated with the first principal component, which represents the maximum variance direction of the data along the personality axis ($\rho = 0.324$, $p = 0.004$). The negative correlation with the second principal component is not statistically significant ($\rho = -0.171$, $p = 0.131$). Similarly, when considering individual video segments, only the correlation with the first principal component is statistically significant (1st PC: western: $\rho = 0.230$, $p = 0.042$; reality show: $\rho = 0.363$, $p = 0.001$; emotional clips: $\rho = 0.254$, $p = 0.024$. 2nd PC: western: $\rho = -0.154$, $p = 0.176$; reality show: $\rho = -0.147$, $p = 0.195$; emotional clips: $\rho = -0.158$, $p = 0.163$). For comparison and full transparency, the Appendix also includes the results of this analysis without removing outliers (Table A1) and with age and sex included as covariates (Table A3).

Upon closer inspection of individual personality traits, we found that the typicality of brain activity during the experimental condition appears to be significantly negatively correlated with Neuroticism ($\rho = -0.290$, $p = 0.010$) and Harm Avoidance ($\rho = -0.307$, $p = 0.006$) and positively correlated with Extraversion ($\rho = 0.265$, $p = 0.018$) and Cooperativeness ($\rho = 0.281$, $p = 0.012$). These relationships are depicted in Fig. 5. No significant relationship with the other personality traits was observed. Looking at the individual video segments, we found that the relationship between the typicality of brain activity and the personality traits is in line with the results for all three videos combined. In particular, the brain activity corresponding to the reality show is related to all of the personality traits associated with the three video segments combined. In the case of the western and the emotional clips, the relationships are descriptively similar, albeit not statistically significant for each personality trait. For de-

tailed results, see Table 1.

Figure 6 shows how each of the 39 ICA components relates to each of the 12 personality traits. Note that the pattern is primarily driven by the personality traits rather than by the similarity indices corresponding to the independent components of brain activity. In numerical terms, the five personality traits assessed by NEO-FFI are significantly ($p < 0.05$) correlated with similarity indices corresponding to 13, 17, 0, 1, and 5 of the 39 independent components, respectively, while the seven traits assessed by TCI are significantly correlated with 2, 15, 0, 7, 8, 12, and 0 similarity indices. The values are, naturally, affected by statistical power being weaker in the case of individual similarity indices relative to the summarized typicality index. Nevertheless, the direction of the correlations remains consistent with the previous analysis: all significant correlations with Neuroticism and Harm Avoidance are negative, while those with Extraversion and Cooperativeness are positive.

Pairs of canonical variables with the strongest correlations resulting from the canonical correlation analysis are shown in Fig. 7 (three pairs out of a total of 12, with $r = 0.911$, $r = 0.903$, and $r = 0.840$, respectively). Notably, none of the canonical correlations are statistically significant ($F_{(468,364)} = 1.110$, $p = 0.146$; $F_{(418,342)} = 0.994$, $p = 0.526$, and $F_{(370,319)} = 0.865$, $p = 0.911$, respectively).

AAL analyses. Note that, due to the removal of different subsets of outliers, the final subject sample for the AAL analyses ($N = 81$) is slightly larger than that used in the ICA analyses ($N = 79$). Nevertheless, neither of the correlations between the typicality indices and subject age and sex is statistically significant (age: $\rho = -0.181$, $p = 0.106$; sex: $\rho = -0.018$, $p = 0.871$). The following analyses, thus, again, do not take these two variables into account.

The first two principal components, again, explain 55% of the total variance and were thus selected for the primary analysis. Here, the first principal component is influenced negatively by Neuroticism and Harm Avoidance and positively primarily by Cooperativeness and Self-Directedness, i.e., with this data, the latter surpasses Agreeableness.

Using the 90 regions of the AAL atlas instead of the 39 ICA components, the experimental condition is, again, positively correlated with the first principal component ($\rho = 0.276$, $p = 0.013$) and the negative correlation with the second principal component, again, is not statistically significant ($\rho = -0.092$, $p = 0.413$). Similarly, when considering individual video segments, only the correlation with the first principal component is statistically significant – and only for the reality show (1st PC: western: $\rho = 0.193$, $p = 0.084$; reality show: $\rho = 0.297$, $p = 0.007$; emotional clips: $\rho = 0.154$, $p = 0.170$. 2nd PC: western: $\rho = -0.069$, $p = 0.537$; reality show: $\rho = -0.046$, $p = 0.685$; emotional clips: $\rho = -0.081$, $p = 0.474$). Again, the Appendix also includes the results of this analysis without removing outliers (Table A2) and with age and sex included as covariates (Table A4).

An examination of individual personality traits showed significant negative correlations with Neuroticism ($\rho = -0.226$, $p = 0.042$) and Harm Avoidance ($\rho = -0.250$, $p = 0.024$) and a positive correlation with Self-Directedness ($\rho = 0.249$, $p = 0.025$), as depicted in Fig. 8. No significant relationship with the other personality traits was observed. The results for individual video segments are in line with the combined results for all three videos combined. For details, see Table 2. The five personality traits assessed by NEO-FFI are significantly ($p < 0.05$) correlated with similarity indices corresponding to 17, 16, 0, 0, and 0 of the 90 ROIs, respectively. The seven traits assessed by TCI are significantly correlated with 2, 28, 0, 2, 23, 15, and 0 similarity indices, respectively. In line with the results of the previous analysis, all of the significant correlations with Neuroticism and Harm Avoidance are negative and the significant correlations with Self-Directedness are all positive. Figure 9 shows the correlations of each personality trait with each ROI in detail.

The findings from the AAL analyses thus closely mirror those obtained using ICA.

Discussion

Using naturalistic fMRI thus helped us detect relationships between brain activity and personality. While the personality traits associated with the typicality of brain activity vary slightly across the analyses, the relationship between lower typicality and Neuroticism/Harm Avoidance – two highly correlated personality traits that likely measure similar aspects of personality – emerges consistently in all of them, including the primary analysis where the first principal component loads mainly on these traits. Moreover, the ICA analyses suggest a positive relationship with Extraversion – perhaps unsurprisingly, given that Extraversion and Neuroticism are strongly negatively correlated. However, as previously noted, findings regarding specific personality traits are based on exploratory analyses, uncorrected for multiple comparisons, and should therefore be interpreted cautiously.

The results of our primary analysis reveal a relationship between personality and the typicality of brain activity, detectable with ICA as well as AAL data.

Comparisons across the video segments suggest that the relationship with personality traits slightly varies across naturalistic stimuli. However, we did not explicitly test for differences. Moreover, since the videos may differ in a range of parameters, we do not speculate on the generalizability of the observations to entire genres. We note, nevertheless, that the results of these comparisons are in line with previous reports of social clips predicting behaviour better than other types of videos (27). Moreover, the advantage of the reality show could also be due to its comparably high split-half reliability (37).

We further narrowed the results down to individual personality traits and even related those to individual independent components/ROIs. Some of the relationships appear to be significant. However, these results were presented uncorrected. The recommendation to interpret findings of studies relating personality to brain activity with great caution also applies here and we refrain from drawing conclusions about the role of different ICs or ROIs in relation to individual personality traits.

Fig. 5. Relationship between the typicality of brain activity measured across the three video segments and selected personality traits (ICA analysis). The scatterplots show relationships between the typicality indices (horizontal axes) and the scores on certain personality dimensions of the two personality inventories (vertical axes) across subjects. Only personality traits for which the result is statistically significant ($p < 0.05$) are shown. *Left* (NEO-FFI): *top*: Neuroticism ($\rho = -0.290$, $p = 0.010$); *bottom*: Extraversion ($\rho = 0.265$, $p = 0.018$). *Right* (TCI): *top*: Harm Avoidance ($\rho = -0.307$, $p = 0.006$); *bottom*: Cooperativeness ($\rho = 0.281$, $p = 0.012$).

		all	**western**	**reality show**	**emotional clips**
	Neuroticism	**-0.290 (0.010)**	**-0.243 (0.031)**	**-0.290 (0.010)**	-0.201 (0.075)
	Extraversion	**0.265 (0.018)**	**0.233 (0.039)**	**0.360 (0.001)**	0.143 (0.208)
NEO-FFI	**Openness to experience**	0.003 (0.980)	-0.091 (0.426)	0.053 (0.641)	-0.033 (0.770)
	Agreeableness	0.169 (0.136)	0.107 (0.349)	0.201 (0.075)	0.141 (0.214)
	Conscientiousness	0.149 (0.190)	0.073 (0.525)	0.181 (0.110)	0.150 (0.186)
	Novelty Seeking	0.046 (0.685)	0.062 (0.589)	0.068 (0.552)	0.001 (0.996)
	Harm Avoidance	**-0.307 (0.006)**	**-0.277 (0.014)**	**-0.304 (0.006)**	-0.192 (0.089)
	Reward Dependence	0.022 (0.847)	0.006 (0.958)	0.077 (0.501)	0.003 (0.982)
TCI	**Persistence**	0.216 (0.056)	0.149 (0.190)	0.218 (0.054)	0.213 (0.059)
	Self-Directedness	0.215 (0.057)	0.100 (0.378)	**0.264 (0.019)**	0.169 (0.137)
	Cooperativeness	**0.281 (0.012)**	0.198 (0.080)	**0.315 (0.005)**	**0.233 (0.039)**
	Self-Transcendence	-0.049 (0.671)	-0.044 (0.702)	-0.030 (0.795)	-0.071 (0.532)

Table 1. Relationship between each personality trait and the typicality of brain activity for individual video segments (ICA analysis). Spearman's rho (*p*-value) for each video is shown. Results across all videos ('all') are also included for comparison. Statistically significant ($p < 0.05$) results are highlighted.

Indeed, the more stringent canonical correlation analysis did not detect a relationship between personality traits and similarities of brain activations to those of an average subject. Given the large number of degrees of freedom – stemming from the large number of independent variables on both sides (personality traits and typicality indices) and the relatively small sample size – the fact that canonical correlation analysis lacked the power to detect the relationship and failed to replicate the previous findings was, arguably, to be expected.

While the main findings remained largely similar when time series derived from regions of the AAL atlas were used instead of independent components, statistically significant correlations were generally fewer and weaker. Including all anatomic regions of the brain will inevitably introduce more noise than the output of independent component analysis. The weaker correlations are, therefore, unsurprising and highlight the appropriateness of ICA for our study.

As explained above, the typicality index of a subject was

Fig. 6. Spearman correlations between the 12 personality traits and the similarity indices corresponding to the 39 independent components (ICA analysis), $* \ p < 0.05$, $** \ p < 0.01$.

		all	western	reality show	emotional clips
NEO-FFI	**Neuroticism**	**-0.226 (0.042)**	-0.195 (0.082)	**-0.262 (0.018)**	-0.086 (0.447)
	Extraversion	0.183 (0.102)	0.141 (0.209)	**0.275 (0.013)**	0.012 (0.916)
	Openness to experience	0.045 (0.693)	-0.006 (0.958)	0.116 (0.304)	-0.023 (0.839)
	Agreeableness	0.103 (0.361)	0.065 (0.567)	0.083 (0.461)	0.081 (0.471)
	Conscientiousness	0.081 (0.472)	0.022 (0.842)	0.139 (0.217)	0.054 (0.635)
TCI	**Novelty Seeking**	0.086 (0.447)	0.176 (0.116)	0.068 (0.547)	-0.071 (0.531)
	Harm Avoidance	**-0.250 (0.024)**	**-0.288 (0.009)**	**-0.299 (0.007)**	-0.006 (0.956)
	Reward Dependence	0.000 (0.997)	-0.056 (0.620)	0.073 (0.517)	-0.009 (0.936)
	Persistence	0.132 (0.240)	0.051 (0.648)	0.184 (0.101)	0.047 (0.677)
	Self-Directedness	**0.249 (0.025)**	0.175 (0.119)	**0.278 (0.012)**	0.124 (0.271)
	Cooperativeness	0.216 (0.053)	0.123 (0.274)	0.179 (0.110)	0.166 (0.140)
	Self-Transcendence	-0.023 (0.837)	0.040 (0.721)	-0.022 (0.844)	-0.040 (0.724)

Table 2. Relationship between each personality trait and the typicality of brain activity for individual video segments (AAL analysis). Presentation as in Table 1.

computed by correlating their time series to the typical time series. The fact that a typical time series was computed as the mean across all subjects, including the subject in question, could arguably have had an impact on the results. While it clearly biases typicality upwards, the case for it also systematically affecting the relationship between personality and typicality is rather unclear. Still, out of an abundance of caution, we also performed a version of the analyses where a subject was omitted from the calculation of the typical time series when their typicality index was being quantified. As expected, the impact of the inclusion or removal of the subject on the result was negligible.

Since the unconstrained resting-state condition does not allow for meaningful comparisons of time series across subjects, previous studies relating personality to brain activity all analysed functional connectivity. For comparison, we also

Fig. 7. Relationships between the canonical scores of subject similarity indices and personality scores (ICA analysis). The pairs of canonical variables ($similarity_i, personality_i$) depicted in subplots along the diagonal are ordered by the strength of their correlations – strongest to weakest, with $r = 0.911$, $r = 0.903$, and $r = 0.840$, respectively. (Out of 12 pairs, only the three pairs with the strongest canonical correlations are shown. None of these are statistically significant.) All the other pairs (off-diagonal subplots) are independent.

repeated our primary analysis using functional connectivity – rather than computing the typicality of the time series, here, we computed the typicality of the FC matrices derived from them (both ICA and AAL). Interestingly, using this approach, we found subject age to be (negatively) correlated with subject typicality indices. This finding is in line with the previously reported age-related differences discussed above. The association between personality and FC typicality was therefore measured using Spearman's *partial* correlation, controlling for age. We did not detect any statistically significant results. This highlights the importance of temporal dynamics and, consequently, the advantage of the naturalistic viewing condition for studies of the relationship between personality and brain activity.

Conclusion

We have shown that a relationship between personality traits and the typicality of brain activity, indeed, exists, and can be detected using naturalistic stimuli. While a number of personality traits seem to be related to the typicality of brain activity, this relationship is most evident in subjects with higher Neuroticism and Harm Avoidance whose brain activity tended to show lower typicality.

Our findings highlight the importance of temporal dynamics for studies of the relationship between personality and brain

activity. Consequently, the fact that the naturalistic viewing condition allows for taking temporal dynamics into account presents an important advantage over rest.

Further research is necessary to enhance our understanding of the nature of this relationship. Datasets with a greater range of personality scores might further increase signal variance, thus increasing the likelihood of detecting associations. Validating the results using different, preferably larger, datasets could also allow for a more comprehensive examination of the relationship between personality traits and brain activation patterns, and more confidence in drawing conclusions regarding specific personality traits, as well as an improved understanding of the variability across different audiovisual stimuli. Deliberately choosing stimuli that differ with respect to a certain parameter, while being similar in other aspects, might provide more insights into the role of various parameters of the stimuli. This would, in turn, allow for better informed selection of suitable stimuli in future experiments.

AUTHOR CONTRIBUTIONS

Lucia Jajcay: Methodology, Software, Validation, Formal analysis, Investigation, Writing - original draft, Writing - review & editing, Visualization. **David Tomeček:** Software, Formal analysis, Investigation, Writing - review & editing. **Renáta Androvičová:** Investigation, Data curation, Writing - review & editing, Project administration. **Iveta Fajnerová:** Investigation, Data curation, Writing - review & editing, Project administration. **Filip Dechtěrenko:** Investigation, Data curation, Writing - review & editing. **Jan Rydlo:** Investigation, Writing - review & editing. **Jaroslav Tintěra:** Investigation, Resources, Writing - review & editing, Project administration, Funding acquisition. **Jiří Lukavský:** Investigation,

Fig. 8. Relationship between the typicality of brain activity measured across the three video segments and selected personality traits (AAL analysis). The scatterplots show relationships between the typicality indices (horizontal axes) and the scores on certain personality dimensions of the two personality inventories (vertical axes) across subjects. Only personality traits for which the result is statistically significant ($p < 0.05$) are shown. *Left* (NEO-FFI): Neuroticism ($\rho = -0.226$, $p = 0.042$). *Right* (TCI): *top*: Harm Avoidance ($\rho = -0.250$, $p = 0.024$); *bottom*: Self-Directedness ($\rho = 0.249$, $p = 0.025$).

Resources, Writing - review & editing, Supervision, Project administration, Funding acquisition. **Jiří Horáček:** Investigation, Resources, Writing - review & editing, Supervision, Project administration, Funding acquisition. **Jaroslav Hlinka:** Conceptualization, Methodology, Validation, Formal analysis, Investigation, Resources, Data curation, Writing - review & editing, Supervision, Project administration, Funding acquisition.

ACKNOWLEDGMENTS
This work was supported by the European Regional Development Fund (ERDF) – project Brain dynamics (project No. CZ.02.01.01/00/22_008/0004643); the Czech Science Foundation (project No. 21-32608S); and by the Grant Agency of the Czech Technical University in Prague (grant No. SGS23/119/OHK3/2T/13).

CONFLICT OF INTEREST
The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT
The fMRI and personality data are stored in an OSF repository at https://osf.io/795hb/ and will be made publicly available once the manuscript has been accepted for publication.

Bibliography

1. American Psychological Association. Personality. APA Dictionary of Psychology. Updated April 19, 2018. Accessed April 23, 2024. https://dictionary.apa.org/personality.

2. Philip J. Corr and Gerald (Eds.) Matthews. *The Cambridge Handbook of Personality Psychology*. Cambridge University Press, 2nd edition, 2020.

3. Paul T Costa Jr. *The NEO-PI-R Professional Manual: Revised NEO Five-Factor Inventory (NEO-FFI)*. Psychological Assessment Resources, 1992.

4. J M Digman. Personality structure: Emergence of the five-factor model. *Annual Review of Psychology*, 41(1):417–440, 1990. doi: https://doi.org/10.1146/annurev.ps.41.020190.002221.

5. Julien Dubois, Paola Galdi, Yanting Han, Lynn K Paul, and Ralph Adolphs. Resting-state functional brain connectivity best predicts the personality dimension of openness to experience. *Personality Neuroscience*, 1:e6, 2018. doi: https://doi.org/10.1017/pen.2018.8.

6. David Tomeček, Renata Androvičová, Iveta Fajnerová, Filip Děchtěrenko, Jan Rydlo, Jiří Horáček, Jiří Lukavský, Jaroslav Tintěra, and Jaroslav Hlinka. Personality reflection in the brain's intrinsic functional architecture remains elusive. *PLoS ONE*, 15(6):e0232570, 2020. doi: https://doi.org/10.1371/journal.pone.0232570.

7. Arne Ekstrom. How and when the fMRI BOLD signal relates to underlying neural activity: The danger in dissociation. *Brain Research Reviews*, 62(2):233–244, 2010. doi: https://doi.org/10.1016/j.brainresrev.2009.12.004.

8. John C Gore et al. Principles and practice of functional MRI of the human brain. *The Journal of Clinical Investigation*, 112(1):4–9, 2003. doi: https://doi.org/10.1172/JCI19010.

9. Karl J. Friston. Functional and effective connectivity in neuroimaging: A synthesis. *Human Brain Mapping*, 2(1–2):56–78, 1994. ISSN 10659471. doi: https://doi.org/10.1002/hbm.460020107.

10. Bharat B. Biswal, Maarten Mennes, Xi-Nian Zuo, Suril Gohel, Clare Kelly, Steve M. Smith, Christian F. Beckmann, Jonathan S. Adelstein, Randy L. Buckner, Stan Colcombe, Anne-Marie Dogonowski, Monique Ernst, Damien Fair, Michelle Hampson, Matthew J. Hoptman, James S. Hyde, Vesa J. Kiviniemi, Rolf Kotter, Shi-Jiang Li, Ching-Po Lin, Mark J. Lowe, Clare Mackay, David J. Madden, Kristoffer H. Madsen, Daniel S. Margulies, Helen S. Mayberg, Katie McMahon, Christopher S. Monk, Stewart H. Mostofsky, Bonnie J. Nagel, James J. Pekar, Scott J. Peltier, Steven E. Petersen, Valentin Riedl, Serge A. R. B. Rombouts, Bart Rypma, Bradley L. Schlaggar, Sein Schmidt, Rachael D. Seidler, Greg J. Siegle, Christian Sorg, Gao-Jun Teng, Juha Veijola, Arno Villringer, Martin Walter, Lihong Wang, Xu-Chu Weng, Susan Whitfield-Gabrieli, Peter Williamson, Christian Windischberger, Yu-Feng Zang, Hong-Ying Zhang, F. Xavier Castellanos, and Michael P. Milham. Toward discovery science of human brain function. *Proceedings of the National Academy of Sciences of the United States of America*, 107(10):4734–4739, 2010. ISSN 0027-8424. doi: https://doi.org/10.1073/pnas.0911855107.

11. Koene R. A. van Dijk, Trey Hedden, Archana Venkataraman, Karleyton C. Evans, Sara W. Lazar, and Randy L. Buckner. Intrinsic functional connectivity as a tool for human connectomics: Theory, properties, and optimization. *Journal of Neurophysiology*, 103(1):297–321, 2010. ISSN 0022-3077. doi: https://doi.org/10.1152/jn.00783.2009.

12. Emily S. Finn, Xilin Shen, Dustin Scheinost, Monica D. Rosenberg, Jessica Huang, Marvin M. Chun, Xenophon Papademetris, and R. Todd Constable. Functional connectome fingerprinting: Identifying individuals using patterns of brain connectivity. *Nature Neuroscience*, 18(11):1664–1671, 2015. doi: https://doi.org/10.1038/nn.4135.

13. Moji Aghajani, Ilya M Veer, Marie-José Van Tol, André Aleman, Mark A Van Buchem, Dick J Veltman, Serge ARB Rombouts, and Nic J van der Wee. Neuroticism and extraversion are associated with amygdala resting-state functional connectivity. *Cognitive, Affective, & Behavioral Neuroscience*, 14:836–848, 2014. doi: https://doi.org/10.3758/s13415-013-0224-0.

14. Michelle N Servaas, Linda Geerligs, Remco J Renken, Jan-Bernard C Marsman, Johan Ormel, Harriëtte Riese, and André Aleman. Connectomics and neuroticism: An altered functional network organization. *Neuropsychopharmacology*, 40(2):296–304, 2015. doi: https://doi.org/10.1038/npp.2014.169.

15. Adriana Sampaio, José Miguel Soares, Joana Coutinho, Nuno Sousa, and Óscar F Gonçalves. The Big Five default brain: Functional evidence. *Brain Structure and Func-

Fig. 9. Spearman correlations between the 12 personality traits and the similarity indices corresponding to the 90 regions of interest (AAL analysis), $* \ p < 0.05$, $** \ p < 0.01$.

tion, 219:1913–1922, 2014. doi: https://doi.org/10.1007/s00429-013-0610-y.

16. Sharon S Simon, Eleanna Varangis, and Yaakov Stern. Associations between personality and whole-brain functional connectivity at rest: Evidence across the adult lifespan. *Brain and Behavior*, 10(2):e01515, 2020. doi: https://doi.org/10.1002/brb3.1515.

17. Alessandra D Nostro, Veronika I Müller, Deepthi P Varikuti, Rachel N Pläschke, Felix Hoffstaedter, Robert Langner, Kaustubh R Patil, and Simon B Eickhoff. Predicting personality from network-based resting-state functional connectivity. *Brain Structure and Function*, 223: 2699–2719, 2018. doi: https://doi.org/10.1007/s00429-018-1651-z.

18. Junjie Wang, Yang Hu, Hong Li, Ling Ge, Jing Li, Long Cheng, Zhi Yang, Xinian Zuo, and Yong Xu. Connecting openness and the resting-state brain network: A discover-validate approach. *Frontiers in Neuroscience*, 12:762, 2018. doi: https://doi.org/10.3389/fnins.2018.00762.

19. Wei-Ting Hsu, Monica D Rosenberg, Dustin Scheinost, R Todd Constable, and Marvin M Chun. Resting-state functional connectivity predicts neuroticism and extraversion in novel individuals. *Social Cognitive and Affective Neuroscience*, 13(2):224–232, 2018. doi: https://doi.org/10.1093/scan/nsy002.

20. Huanhuan Cai, Jiajia Zhu, and Yongqiang Yu. Robust prediction of individual personality from brain functional connectome. *Social Cognitive and Affective Neuroscience*, 15(3):359–369, 2020. doi: https://doi.org/10.1093/scan/nsaa044.

21. Wei Liu, Nils Kohn, and Guillén Fernández. Intersubject similarity of personality is associated with intersubject similarity of brain connectivity patterns. *NeuroImage*, 186:56–69, 2019. doi: https://doi.org/10.1016/j.neuroimage.2018.10.062.

22. Linda Geerligs, Mikail Rubinov, Cam-CAN, and Richard N. Henson. State and trait components of functional connectivity: Individual differences vary with mental state. *Journal of Neuroscience*, 35(41):13949–13961, 2015. ISSN 0270-6474. doi: https://doi.org/10.1523/JNEUROSCI.1324-15.2015.

23. Tamara Vanderwal, Jeffrey Eilbott, Emily S. Finn, R. Cameron Craddock, Adam Turnbull, and F. Xavier Castellanos. Individual differences in functional connectivity during naturalistic viewing conditions. *NeuroImage*, 157:521–530, 2017. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2017.06.027.

24. Jiahui Wang, Yudan Ren, Xintao Hu, Vinh Thai Nguyen, Lei Guo, Junwei Han, and Christine Cong Guo. Test-retest reliability of functional connectivity networks during naturalistic fMRI paradigms. *Human Brain Mapping*, 38(4):2226–2241, 2017. ISSN 1097-0193. doi: https://doi.org/10.1002/hbm.23517.

25. Emily S. Finn, Dustin Scheinost, Daniel M. Finn, Xilin Shen, Xenophon Papademetris, and R. Todd Constable. Can brain state be manipulated to emphasize individual differences in functional connectivity? *NeuroImage*, 160:140–151, 2017. ISSN 1053-8119. doi: https:

//doi.org/10.1016/j.neuroimage.2017.03.064.

26. Ma Feilong, Samuel A. Nastase, J. Swaroop Guntupalli, and James V. Haxby. Reliable individual differences in fine-grained cortical functional architecture. *NeuroImage*, 183:375–386, 2018. doi: https://doi.org/10.1016/j.neuroimage.2018.08.029.

27. Emily S. Finn and Peter A. Bandettini. Movie-watching outperforms rest for functional connectivity-based prediction of behavior. *NeuroImage*, 235:117963, 2021. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2021.117963.

28. C Robert Cloninger, Thomas R Przybeck, Dragan M Svrakic, and Richard D Wetzel. *The Temperament and Character Inventory (TCI): A guide to its development and use*. Center for Psychobiology of Personality, 1994.

29. Anton Aluja and Angel Blanch. The five and seven factors personality models: Differences and similitude between the TCI-R, NEO-FFI-R and ZKPQ-50-CC. *The Spanish Journal of Psychology*, 14(2):659–666, 2011. doi: https://doi.org/10.5209/rev_sjop.2011.v14.n2.14.

30. Filip De Fruyt, Lieve Van de Wiele, and Cornelis Van Heeringen. Cloninger's psychobiological model of temperament and character and the five-factor model of personality. *Personality and Individual Differences*, 29(3):441–452, 2000. doi: https://doi.org/10.1016/S0191-8869(99)00204-4.

31. Gerard Saucier. Orthogonal markers for orthogonal factors: The case of the Big Five. *Journal of Research in Personality*, 36(1):1–31, 2002. doi: https://doi.org/10.1006/jrpe.2001.2335.

32. C Robert Cloninger. A systematic method for clinical description and classification of personality variants: A proposal. *Archives of General Psychiatry*, 44(6):573–588, 1987. doi: https://10.1001/archpsyc.1987.01800180093014.

33. Paul T Costa and Robert R McCrae. *The NEO Personality Inventory manual*. Psychological Assessment Resources, 1985.

34. Uri Hasson, Y Nir, I Levy, G Fuhrmann, and R Malach. Intersubject synchronization of cortical activity during natural vision. *Science*, 303(5664):1634–1640, 2004. ISSN 0036-8075. doi: https://doi.org/10.1126/science.1089506.

35. Dante Mantini, Uri Hasson, Viviana Betti, Mauro G. Perrucci, Gian Luca Romani, Maurizio Corbetta, Guy A. Orban, and Wim Vanduffel. Interspecies activity correlations reveal functional correspondence between monkey and human brain areas. *Nature Methods*, 9(3): 277–282, 2012. ISSN 1548-7091. doi: https://doi.org/10.1038/nmeth.1868.

36. Alexandre Schaefer, Frederic Nils, Xavier Sanchez, and Pierre Philippot. Assessing the effectiveness of a large database of emotion-eliciting films: A new tool for emotion researchers. *Cognition & Emotion*, 24(7):1153–1172, 2010. ISSN 0269-9931. doi: https://doi.org/10.1080/02699930903274322.

37. Jaroslav Hlinka, Filip Děchtěrenko, Jan Rydlo, Renáta Androvičová, Martin Vejmelka, Lucia

Jajcay, Jaroslav Tintěra, Jiří Lukavskỳ, and Jiří Horáček. The intra-session reliability of functional connectivity during naturalistic viewing conditions. *Psychophysiology*, 59(10): e14075, 2022. doi: https://doi.org/10.1111/psyp.14075.

38. Nathalie Tzourio-Mazoyer, Brigitte Landeau, Dimitri Papathanassiou, Fabrice Crivello, Octave Etard, Nicolas Delcroix, Bernard Mazoyer, and Marc Joliot. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*, 15(1):273–289, 2002. doi: https://doi.org/10.1006/nimg.2001.0978.

39. Wojtek J Krzanowski and Wojtek Krzanowski. *Principles of multivariate analysis: A user's perspective*. Oxford University Press, 1988.

40. George AF Seber. *Multivariate observations*. John Wiley & Sons, 1984.

Supporting information

Additional details can be found in the Appendix.