An imbalance between proliferation and differentiation underlies the development of microRNA-defective pineoblastoma

Claudette R. Fraire[1], Kavita Desai[2,3], Uma A. Obalapuram[4], Lindsay K. Mendyka[5], Veena Rajaram[6], Teja Sebastian[1], Yemin Wang[7], Kenan Onel[8], Jeon Lee[9], Kenneth S. Chen[1,10] *

[1] Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas, TX USA

[2] Division of Oncology, Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, PA USA

[3] University of Pennsylvania Perelman School of Medicine, Philadelphia, PA USA

[4] Emory University, Atlanta, GA USA

[5] Geisel School of Medicine, Dartmouth College, Hanover, NH USA

[6] Department of Pathology, University of Texas Southwestern Medical Center, Dallas, TX USA

[7] Department of Pathology and Laboratory Medicine, University of British Columbia and Department of Molecular Oncology, British Columbia Cancer Research Institute, Vancouver, BC, Canada

[8] Department of Cancer Prevention and Control, Roswell Park Comprehensive Cancer Center, Buffalo, New York

[9] Lyda Hill Department of Bioinformatics, UT Southwestern Medical Center, Dallas, TX USA

[10] Children's Medical Center Research Institute, University of Texas Southwestern Medical Center, Dallas, TX USA

*Correspondence: Kenneth S. Chen, kenneth.chen@utsouthwestern.edu, +1-214-648-8291

The authors have no competing interests to declare.

ABSTRACT

Mutations in the microRNA processing genes *DICER1* and *DROSHA* drive several cancers that resemble embryonic progenitors. To understand how microRNAs regulate tumorigenesis, we ablated *Drosha* or *Dicer1* in the developing pineal gland to emulate the pathogenesis of pineoblastoma, a brain tumor that resembles undifferentiated precursors of the pineal gland. Accordingly, these mice develop pineal tumors marked by loss of microRNAs, including the let-7/miR-98-5p family, and de-repression of microRNA target genes. Pineal tumors driven by loss of *Drosha* or *Dicer1* mimic tumors driven by *Rb1* loss, as they exhibit upregulation of S-phase genes and homeobox transcription factors that regulate pineal development. Blocking proliferation of these tumors facilitates expression of pinealocyte maturation markers, with a concomitant reduction in embryonic markers. Select embryonic markers remain elevated, however, as the microRNAs that normally repress these target genes remain absent. One such microRNA target gene is the oncofetal transcription factor *Plagl2*, which regulates expression of pro-growth genes, and inhibiting their signaling impairs tumor growth. Thus, we demonstrate that tumors driven by loss of microRNA processing may be therapeutically targeted by inhibiting downstream drivers of proliferation.

INTRODUCTION

Mutations in microRNA processing genes, such as *DROSHA*, *DICER1*, and *DGCR8*, drive ~2% of childhood cancers [1, 2]. These mutations arise in a family of cancers that resemble embryonic progenitors, including Wilms tumor, pleuropulmonary blastoma, rhabdomyosarcoma, and the brain tumor pineoblastoma [3-11]. It is thought that loss of microRNA production leads to de-repression of the genes that are normally repressed by these microRNAs, leading to tumor formation [12, 13]. However, microRNA loss leads to transcriptome-wide changes, with different effects in each cell type, and it is challenging to separate genes upregulated through loss of direct microRNA targeting from secondary effects. In addition, little is known about the specific microRNAs or microRNA target genes that regulate the balance between proliferation and differentiation in each cancer type. As a result, there is currently no way to therapeutically target these mutations.

Pineoblastoma is a rare brain tumor that resembles embryonic neurogenic progenitors of the pineal gland, and it is treated with chemotherapy, surgery, and radiation [14]. However, the morbidities associated with each of these treatment modalities limits their utility; cranial radiation therapy, for instance, is generally avoided in young children. Most other *DICER1/DROSHA*-related tumors, which exhibit point mutations in these genes that incompletely impair microRNA production, but pineoblastoma can uniquely exhibit complete loss of *DROSHA* or *DICER1*, leading to complete loss of microRNA production [11, 15]. Specifically, recent molecular studies have identified three mutational patterns in pineoblastoma: (1) bi-allelic loss of *DROSHA*, *DICER1*, or *DGCR8*; (2) bi-allelic loss of *RB1*; or (3) activation of *MYC* or *FOXR2* [16-19]. The RB1 tumor suppressor normally regulates E2F transcription factors to allow progression through the G1/S cell cycle checkpoint [20], and MYC orchestrates tumorigenesis by binding "E-box" motifs in promoters or enhancers of tumor-promoting genes [21]. However, the microRNAs and microRNA target genes that mediate tumorigenesis in tumors driven by loss of *DROSHA* or *DICER1* remain poorly understood.

Previous studies have used *Rb1* loss, cyclin D1 upregulation, and *Dicer1* loss to model the development of pineoblastoma in mice [22-27]. To dissect how microRNA loss drives formation of pineoblastoma, we generated mice with loss of microRNAs in the developing pineal gland. Here we show that loss of *Drosha* or *Dicer1* mimics loss of the *Rb1* tumor suppressor by driving cell cycle progression through de-repression of cyclin D2 (*Ccnd2*). Uncontrolled proliferation itself restrains tumor cells in an undifferentiated state, and restraining proliferation

leads them to upregulate differentiation markers. In addition to being directly targeted by microRNAs, we find that *Ccnd2* is also indirectly regulated by microRNAs through the transcription factor *Plagl2,* and blocking other Plagl2 target genes also slows tumor growth. Lastly, we show that human pineoblastoma tumors exhibit parallel expression patterns.

RESULTS

Pineal tumors driven by loss of Drosha, Dicer1, or Rb1 resemble embryonic pineal progenitors

Prior studies had demonstrated that mice develop pineoblastoma upon pineal-directed ablation of *Rb1* and *p53* (*IrbpCre*;*p53*$^{flox/flox}$;*Rb1*$^{flox/flox}$, hereafter abbreviated as IPRb1) [27]. These studies used a Cre recombinase line driven by the promoter of interphotoreceptor retinoid binding protein (*Irbp*, also known as *Rbp3*), which is expressed in retinal and pineal progenitors [28, 29]. To understand how microRNAs regulate pineal tumor formation, we developed mice with analogous loss of *Drosha* and *p53* (*IrbpCre*;*p53*$^{flox/flox}$;*Drosha*$^{flox/flox}$, abbreviated as IPDrosha) (**Fig. 1A,B**). These mice were healthy at birth, but at 5-10 months of age, they began to exhibit signs of brain tumor formation. Within days of developing an enlarged skull, they become moribund (**Suppl. Fig. S1A**). Similarly, pineal tumors also arise upon conditionally deleting *Dicer1* and *p53* (*IrbpCre*;*p53*$^{flox/flox}$;*Dicer1*$^{flox/flox}$, or IPDicer1) (**Fig. 1A,B**). By microscopic evaluation, IPDrosha, IPDicer1, and IPRb1 tumors contain large, pleomorphic, primitive cells with a high nucleus-to-cytoplasm ratio and rare inconspicuous nucleoli that resemble embryonic pineal progenitors. These cells are arranged as sheets, in a vaguely nodular pattern, with nuclear molding, frequent mitosis, and areas of necrosis. These cells stain positive for markers of proliferation and neuronal lineage (Ki-67 and synaptophysin, respectively) and negative for a glial lineage marker (glial fibrillary acidic protein, or GFAP) (**Fig. 1C,D**). Furthermore, just as *DROSHA/DICER1*-related pineoblastomas arise later in childhood than *RB1*-driven pineoblastoma, IPDrosha and IPDicer1 tumors appeared to arise at later ages than IPRb1 tumors (median age of onset 269 days, 239 days, and 182 days, respectively), though we observed tumors as early as ~5 months of age in all three groups. We did not observe any histological abnormalities in the retina, where *IrbpCre* is also active (not shown).

To understand how these tumors arise, we next profiled gene expression and histone modification patterns, focusing on acetylation of histone H3 lysine 27 (H3K27ac), which marks active promoters and enhancers, and trimethylation of histone H3 lysine 4 (H3K4me3), which marks active promoters. We found that IPDrosha and IPRb1 tumors retain the gene expression patterns of embryonic pineal progenitors, with enrichment for embryonic pineal markers and depletion for adult markers (**Fig. 1E-F**, **Suppl. Fig. S1B-E**). Specifically, the pineal homeobox transcription factors *Otx2, Crx, Pax6*, and *Rax* are expressed at high levels in IPDrosha, IPDicer1, and IPRb1 tumors, reflecting their shared pineal progenitor identity (**Fig. 1G**) [30]. Accordingly, the genomic regions surrounding *Otx2, Crx, Rax*, and *Pax6* are marked by broad

H3K27ac signal, sometimes termed "super-enhancers" [31] (**Suppl. Fig. S1F**). Broad H3K27ac signal at these homeobox transcription factors indicates the importance of these genes to maintain embryonic identity in these pineal tumors. Together, these results indicate that IPDrosha, IPDicer1, and IPRb1 tumors all arise from a similar pineal gland progenitor.

Drosha-driven tumors lack microRNAs and overexpress microRNA target genes

To understand how IPRb1 tumors form with intact microRNA processing and, conversely, how IPDrosha/IPDicer1 tumors form with intact Rb1 function, we next investigated differences between IPRb1 and IPDrosha/IPDicer1 tumors. We first performed small RNA sequencing in IPDrosha tumors, IPRb1 tumors, matched normal brain, and age-matched normal adult pineal glands to identify the microRNAs regulating normal pineal gland development. As expected, IPDrosha tumors are devoid of canonical microRNAs (**Fig. 2A,C**, **Suppl. Fig. S2A**, **Suppl. Data 1**). Some microRNAs can be produced without *Drosha*, through a non-canonical pathway, if they are processed by the splicing machinery (called "miRtrons") or if they are transcribed as a capped microRNA [32, 33]. These "Drosha-independent" microRNAs remain expressed in IPDrosha tumors at levels similar to matched normal brain, age-matched adult pineal glands, or IPRb1 tumors (**Suppl. Fig. 2B-D**). These findings are consistent with *Drosha* loss.

We identified the most abundant microRNAs in age-matched adult pineal glands and in IPRb1 tumors, which resemble embryonic pineal progenitors. Members of the let-7/miR-98-5p, miR-99/100-5p, and miR-125/351-5p families are among the most abundant microRNA families in both IPRb1 tumors and adult pineal glands (**Fig. 2A,C, Suppl. Fig. S2E-F**). These microRNAs are co-transcribed from the same genomic loci in an evolutionarily conserved manner [34]. They are not expressed in neural progenitors but are induced late in neural or retinal development to downregulate progenitor genes [35-37]. In IPRb1 tumors and adult pineal glands, we detected nearly all let-7/miR-98-5p family members (**Fig. 2B,D**). As expected, IPDicer1 tumors similarly expressed undetectable levels of microRNAs (**Fig. 2A,C,E**). The depletion of mature let-7/miR-98-5p microRNAs was seen in IPDrosha tumors despite broad H3K27ac and H3K4me3 at the host genes of let-7/miR-98-5p transcripts (**Suppl. Fig. S2G**).

To understand which microRNAs have the greatest effect on gene expression in IPDrosha tumors, we next examined the expression of genes predicted to be targeted by the 50

microRNAs expressed in IPRb1 tumors but not in IPDrosha tumors. Based on TargetScan calculations of sequence context and conservation, we found 781 genes predicted to be targeted by these 50 microRNAs [38]. These predicted microRNA targets are expressed at higher levels in IPDrosha tumors than IPRb1 tumors, consistent with a loss of microRNA repression (**Fig. 2F**, **Suppl. Data 2**). To understand which microRNAs are the key contributors, we examined enrichment for the genes targeted by each individual microRNA family. Predicted gene targets of the let-7/miR-98-5p and miR-125/351-5p families were the most significantly enriched sets of microRNA targets in IPDrosha compared to IPRb1 tumors or adult pineal glands, consistent with the loss of their expression in IPDrosha tumors (**Fig. 2G**, **Suppl. Fig. S2H-J**, **Suppl. Data 2-3**).

However, observing loss of a microRNA and upregulation of a predicted target transcript cannot fully distinguish between the direct and indirect effects of microRNA targeting. Some genes may be upregulated as a secondary effect of another de-repressed microRNA target gene. Thus, we next used microRNA enhanced cross-linking immunoprecipitation (miR-eCLIP) sequencing to confirm which genes are directly targeted by microRNAs in IPRb1 tumors [39]. This technique directly identifies microRNA-mRNA interactions through immunoprecipitation of Argonaute 2 (AGO2) and ligation of microRNA-mRNA "chimeras". We performed miR-eCLIP sequencing in two independent IPRb1 tumors, and we examined the non-chimeric peaks and chimeric peaks that were reproduced between the two samples (**Suppl. Fig. S3A**, **Suppl. Data 4**). As expected, reproduced reads clustered primarily in the 3' untranslated regions (3'UTRs) of coding genes, confirming enrichment for regions captured by microRNA targeting [12] (**Suppl. Fig. S3B**). In non-chimeric transcripts immunoprecipitated with AGO2, two of the three most significantly enriched motifs were let-7/miR-98-5p recognition sequences (**Suppl. Fig. S3C**). Furthermore, let-7/miR-98-5p family members were commonly identified in the microRNA side of chimeric reads (**Suppl. Fig. S3D**). Altogether, these results highlight the let-7/miR-98-5p family as key regulators of gene expression in IPRb1 tumors.

Next, we examined whether the genes identified as direct microRNA targets in miR-eCLIP sequencing of IPRb1 tumors are repressed compared to IPDrosha tumors. Genes represented in microRNA-mRNA chimeric reads were slightly upregulated in IPDrosha tumors, and they were more likely to be upregulated in IPDrosha tumors than downregulated (**Suppl. Fig. S3E,F**). Furthermore, while transcripts represented in any chimeric reads were upregulated in IPDrosha tumors, chimeras which corresponded to a true 7- or 8-mer perfect seed match were upregulated even further (**Suppl. Fig. S3E**). We thus considered genes with true seed

matches to be "confirmed top microRNA targets". These 1,454 confirmed top microRNA genes were even more upregulated than the microRNA targets predicted by TargetScan (**Fig. 2F,H**). Similarly, the confirmed top targets of the let-7/miR-98-5p family were more enriched than the set of predicted let-7/miR-98-5p targets (**Fig. 2G,I**).

Together, these results demonstrate that de-repression of direct microRNA targets explain many of the differences between IPDrosha and IPRb1 tumors. Nevertheless, genes represented in microRNA-mRNA chimeras only constituted 25.1% of all genes overexpressed in IPDrosha tumors (**Suppl. Fig. S3F**). In other words, while microRNAs directly drive many of the transcriptomic differences between IPDrosha and IPRb1 tumors, up to 75% of the genes overexpressed in IPDrosha tumors may be attributable to other effects.

Proliferation restrains differentiation in Drosha-driven tumors

Thus, we next examined other gene sets upregulated by loss of *Drosha*, whether directly or indirectly. Specifically, we identified "hallmark" gene sets differentially enriched in IPDrosha tumors, IPRb1 tumors, and normal adult pineal glands. We noted that four of the top five most enriched hallmark gene sets overlapped between IPDrosha and IPRb1 tumors, demonstrating their overlapping paths to tumor formation (**Fig. 3A, , Suppl. Data 5**). These similarities in overall transcriptome, despite independent molecular mechanisms, suggests that *Drosha* loss and *Rb1* loss converge to drive cancer formation through overlapping downstream effects.

As expected, since Rb1 normally regulates E2F transcription factors, the most enriched "hallmark" gene set in IPRb1 tumors is "E2F target genes" (**Fig. 3A**, **Suppl. Fig. S4B**). However, E2F target genes were also the most enriched gene set in IPDrosha tumors (**Fig. 3A**, **Suppl. Fig. S4A**). In cells with intact Rb1, D-type cyclins (cyclin D1, D2, or D3) partner with cyclin-dependent kinases 4 and 6 (CDK4/6) to phosphorylate Rb1 and release it from inhibiting E2Fs [20, 40]. It had previously been shown that *Ccnd1* overexpression was sufficient to replace Rb1 loss and drive pineal tumor formation [23, 41]. Thus, we examined the expression of D-type cyclins in IPDrosha and IPDicer1 tumors and found that cyclin D2 (*Ccnd2*) was the only one significantly overexpressed (**Fig. 3B**, **Suppl. Fig. S4C**). *Ccnd2* is a target of several microRNAs; for example, it is thought to drive proliferation of leukemias driven by loss of miR-16 [42, 43]. We confirmed that it is regulated by several microRNAs in our miR-eCLIP sequencing (it was also the most overexpressed predicted let-7/miR-98-5p target gene in IPDrosha tumors,

though we did not detect any chimeras demonstrating let-7/miR-98-5p microRNAs interacting *Ccnd2* transcript in miR-eCLIP, **Suppl. Fig. S4D**). Using immunohistochemistry, we also confirmed that IPDrosha and IPDicer1 tumors exhibit high levels of CDK4/6 activity, as nearly every cell stained positive for phosphorylated Rb1 (**Fig. 3C**). Thus, *Drosha* loss and *Rb1* loss converge on the same proliferation-driving pathway to drive cancer cells through the G1/S cell cycle checkpoint.

In the developing brain, proliferation maintains progenitors in an undifferentiated state, and differentiation is allowed to occur once proliferation slows down [44]. Accordingly, IPRb1 tumors remain undifferentiated solely because of unrestrained proliferation of these embryonic progenitors. Thus, to understand whether IPDrosha/IPDicer1 tumors with high levels of *Ccnd2* are dependent on CDK4/6 activity to remain undifferentiated, we treated them with the clinically approved CDK4/6 inhibitor palbociclib [45]. Specifically, we implanted tumor cells as subcutaneous allografts and administered either vehicle or palbociclib to recipient mice. Over two weeks, palbociclib suppressed proliferation and extended survival in IPDrosha and IPDicer1 tumors (**Fig. 3D** and **Suppl. Fig. S4E**). Palbociclib-treated tumors exhibited loss of Rb1 phosphorylation, consistent with a loss of CDK4/6 activity (**Fig. 3E**). Histologically, the palbociclib-treated IPDrosha and IPDicer1 tumors exhibit increased karyorrhexis and increased background neuropil, a sign of differentiation.

Next, we used RNA-seq to understand how palbociclib affected IPDrosha tumor cells. Consistent with inhibition of CDK4/6, E2F target genes were the most downregulated hallmark gene set in palbociclib-treated tumors (**Fig. 3F**). In contrast, interferon alpha and gamma response pathways were enriched in palbociclib-treated tumors, consistent with reports that CDK4/6 inhibitors can induce antitumor immunity [46-49] (**Suppl. Fig. S4F**). When we examined pineal differentiation markers, we found that palbociclib-treated tumors were enriched for adult pineal markers such as *Chrna3* and *Nefh* (**Fig. 3G,H, Suppl. Fig. S4G**). On the other hand, embryonic markers were mildly depleted as a whole (**Fig. 3G**, **Suppl. Fig. S4H**). Many embryonic genes remain highly expressed upon palbociclib treatment, such as the embryonic transcription factor *Plagl2* (**Fig. 3H**). In other words, blocking the effect of *Ccnd2* partially restores differentiation in IPDrosha tumors, but microRNAs remain absent, allowing some embryonic pineal markers to remain highly expressed.

The microRNA target gene Plagl2 regulates proliferation drivers Igf2 and Ccnd2

We next identified the proliferation-independent microRNA target genes that contribute to maintaining an embryonic expression program. Specifically, we next identified embryonic pineal markers that were overexpressed in IPDrosha tumors, confirmed "top" let-7/miR-98-5p target genes by miR-eCLIP, and independent of palbociclib. Three genes met these criteria: *Bach1*, *Tgfbr1*, and *Plagl2*. We focused on *Plagl2* (*PLAG1*-like gene 2) because of previous evidence that it is an oncogenic, developmental transcription factor, and because its most well-characterized target gene, *Igf2*, was the single most overexpressed gene in IPDrosha tumors [50-52] (**Fig. 4A-B**). Our miR-eCLIP sequencing demonstrated that the 3' UTR of *Plagl2* interacted with several microRNAs (**Fig. 4C**). In fact, let-7/miR-98-5p microRNAs interacted with *Plagl2* at three sites, including one highly conserved site with a 7-mer perfect match.

Histone marks at the *Plagl2* locus were similar in IPDrosha and IPRb1 tumors, consistent with a gene whose expression is primarily regulated by microRNAs post-transcriptionally (**Suppl. Fig. S5A**). At the *Ccnd2* locus, however, IPDrosha tumors had much higher levels of H3K27ac and H3K4me3, suggesting that *Ccnd2* upregulation is due to a combination of transcriptional and post-transcriptional regulation (**Fig. 4D**). Since the *Ccnd2* promoter also contains two clusters of predicted Plagl2 binding sites, we examined whether Plagl2 resides at the *Ccnd2* promoter in IPDrosha tumors (**Fig. 4D**). Using ChIP-qPCR, we verified Plagl2 binding at a region of the *Ccnd2* promoter with four Plagl2 binding motifs clustered in a 300bp range ("region 2" in **Fig. 4D,E**). Lastly, we noted that levels of *Plagl2*, *Igf2,* and *Ccnd2* are high in embryonic pineal glands and fall in parallel during pineal development, suggesting coordinated regulation (**Suppl. Fig. S5B**). Together, these results suggest that *Ccnd2* is upregulated by both increased transcription and decreased microRNA repression, and that the microRNA target gene *Plagl2* contributes to the maintenance of the embryonic expression program, in part by regulating *Igf2* and *Ccnd2*.

Igf2, whose overexpression drives many other childhood cancers, signals through the insulin signaling pathway primarily by binding the insulin-like growth factor 1 receptor (Igf1r), leading to activation of mTORC1 and other kinase pathways [53]. Accordingly, we observed that the "insulin signaling" and "mTORC1 signaling" gene sets were significantly enriched in IPDrosha tumors (**Fig. 5A** and **Fig. 3A**). Thus, to study the importance of Igf2 signaling, we treated tumor-bearing mice with ceritinib, a small-molecule Igf1r inhibitor [54]. In both IPDrosha and IPDicer1 tumors, we found that ceritinib significantly impaired tumor growth (**Fig. 5B**).

Vehicle- and ceritinib-treated IPDrosha and IPDicer1 tumors are histologically similar, though ceritinib-treated tumors appeared to have slightly more background neuropil (**Fig. 5C**). By immunostains, ceritinib-treated tumors exhibited decreased phosphorylation of S6 (a measure of Igf1r/mTORC1 activity) and Rb1 (**Fig. 5C**). Similarly, RNA sequencing revealed depletion of the insulin signaling genes in ceritinib-treated IPDrosha tumors (**Fig. 5D**).

E2F targets are enriched in human pineoblastoma tumors with low DROSHA

Lastly, we examined RNA-seq data from human pineoblastoma tumors. Among these tumors, we noted that *DROSHA* expression was essentially undetectable in two of nine tumors. We compared expression patterns in these two "*DROSHA*-low" tumors to the other seven "*DROSHA*-high" tumors. Consistent with impairment in microRNA processing, we found that *DROSHA*-low tumors are enriched for genes predicted to be targeted by let-7/miR-98-5p and other microRNAs (**Suppl. Fig. S6A,B**). Furthermore, consistent with our findings in IPDrosha tumors, the most enriched hallmark gene set in *DROSHA*-low tumors is "E2F targets" (**Suppl. Fig. 6C,D**, **Suppl. Data 6**). Lastly, although they did not reach statistical significance, the two *DROSHA*-low tumors expressed slightly higher levels of *CCND1, CCND2*, *PLAG1*, and *PLAGL2* (**Suppl. Fig. S6E**). (*IGF2* was not consistently detected in these tumors). In sum, these data in human tumors are consistent with findings from our mouse model that impaired microRNA processing dysregulates the G1/S checkpoint.

DISCUSSION

Drosha loss leads to de-repression of microRNA target genes, which has cascading effects on the expression of other genes. In this study, we dissect *Drosha*-dependent gene networks with a combination of microRNA sequencing, RNA-sequencing, miR-eCLIP sequencing and histone profiling. Our study shows that loss of *Drosha* drives pineal tumors by mimicking *RB1* loss. Together, these studies revealed *Plagl2* and *Ccnd2* as microRNA target genes driving IPDrosha tumors and small-molecule inhibitors of these effectors that can block tumor growth (**Suppl. Fig. S6F**). In other words, while loss of microRNAs cannot be therapeutically targeted currently, the microRNA target genes that drive tumorigenesis could be blocked with available drugs.

Our results shed light on the cells from which pineoblastoma arise. IPDrosha, IPDicer1, and IPRb1 tumors express high levels of *Pax6, Otx2, Crx*, and *Rax* – the homeobox transcription factors that cooperate to regulate pinealocyte development. In particular, *Pax6* and *Otx2* are expressed in rapidly dividing cells at the earliest stages of pineal development [30, 55]. *Otx2* regulates the expression of *Crx*, a specific marker for retina and the pineal gland that may even be used as a diagnostic marker for retinoblastoma and pineoblastoma [19, 56-58]. *Otx2* and *Crx* recognize the same binding motif and thus activate the same target genes, including *Irbp* and *Rax* [30, 59, 60]. In fact, *OTX2* and *CRX* are key oncogenic drivers in group 3 medulloblastoma and are likely to play similar roles in pineoblastoma [61, 62]. Together, these results suggest that pineoblastomas driven by loss of *Drosha*, *Dicer1*, or *Rb1* arise from an embryonic pineal progenitor marked by *Pax6*, *Otx2*, *Crx*, *Rax*, and *Irbp*.

Using a Cre driver initially developed for the mammary gland, Chung et al. previously showed that *Dicer1* ablation also drives development of pineal tumors; to our knowledge, our study is the first report of pineal tumors arising in mice through loss of *Drosha* [26, 63]. Our findings recapitulate many of their findings, including incomplete penetrance and delayed onset compared with *Rb1*-driven tumors, suggesting that tumors from the two Cre drivers represent the same cell of origin. Nevertheless, the incomplete penetrance and delayed onset of *Drosha/Dicer1*-driven tumors suggest that other factors might also contribute to or restrain tumor development. For instance, in both studies, loss of *Rb1*, *Dicer1,* or *Drosha* alone was not sufficient to drive tumor development. In other words, while these genetic events are critical drivers for tumor development, other tumor-suppressive mechanisms, such as p53 and p18^{Ink4c}, may compensate to restrain cell proliferation, as previously reported in a cyclin D1-driven

mouse model of pineal tumors [23]. The role of such mechanisms is unclear in human pineoblastoma, which rarely exhibits loss or mutation of p53. Future studies are thus required to characterize the genetic status of these genes in human pineoblastoma and understand the effect of p53 loss on the proliferative potential of pineal cells losing *Rb1*, *Dicer1,* or Drosha.

Previous studies showed that pineal tumors arise through dysregulation of the G1/S checkpoint, either by ablating *Rb1* or upregulating *Ccnd1* [23, 41]. By demonstrating that IPDrosha/IPDicer1 tumors drive proliferation by upregulating D-type cyclins, we build upon these studies and show similarities between loss of *Drosha/Dicer1* and loss of *Rb1*. CDK4/6 inhibitors are clinically approved for use in hormone receptor-positive breast cancer and under investigation for treatment of other undifferentiated pediatric and adult tumors that retain intact RB1-E2F responses [20, 40, 64, 65]. Specifically, CDK4/6 inhibitors have been found to induce differentiation in neuroblastoma [66], rhabdomyosarcoma [22], medulloblastoma [67], and prostate cancer [68]. To date, the only known genomic predictors of sensitivity to CDK4/6 inhibition directly drive overexpression of *CCND1*, *CCND2*, or *CCND3* [69]. These may arise through translocation, amplification, or variants that remove or alter microRNA binding sites in the 3' UTR. Our results suggest that removal of the microRNAs themselves could also contribute to overexpression of D-type cyclins and thus sensitivity to CDK4/6 inhibitors.

We have also previously reported overlapping results in Wilms tumor, a pediatric cancer of embryonic renal progenitors frequently driven by mutations in microRNA processing genes [3-7]. Unlike the complete loss of microRNA production seen in pineoblastoma, Wilms tumors most commonly exhibit point mutations in *DROSHA* that lead to incomplete impairment of microRNA processing. In two separate studies, we previously found that *DROSHA* regulates *CCND2* and *PLAG1/IGF2* in Wilms tumor; it remains to be seen whether *PLAG1* or *PLAGL2* interacts with *CCND2* in Wilms tumor [70, 71]. Regardless, together, our observations suggest that despite arising from different cell types and through different effects on microRNA processing, *DROSHA*-mutant Wilms tumor and pineoblastoma harness similar pathways to drive proliferation of embryonic progenitors.

Another family of tumors, including pleuropulmonary blastoma, Sertoli-Leydig cell tumor, and rhabdomyosarcoma, arises through point mutations in the RNase IIIB domain of *DICER1,* which cleaves the 5p arm of pre-microRNA hairpins [9, 10]. This results in loss of 5p-derived microRNAs but retention of 3p-derived microRNAs. Many of these *DICER1*-related tumors also resemble highly proliferative, undifferentiated progenitors. In fact, in a recently published mouse

model of *Dicer1* RNase IIIb mutation, the gynecologic tumors that arose exhibited loss of 5p microRNAs and upregulation of E2F and Myc target genes, similar to our findings [72]. Furthermore, many of the microRNAs we identified as key regulators of gene expression in pineoblastoma are 5p-derived and thus lost in tumors with *DICER1* missense variants, including members of the let-7/miR-98-5p, miR-99/100-5p, and miR-125/352-5p families. It remains to be seen whether D-type cyclins and *PLAG1* family members are also important in these cancer types.

Recently, a subset of embryonal brain tumors was found to be defined by amplification and overexpression of *PLAGL2*, and our study expands on its oncogenic importance [73]. *PLAGL2* was initially described as a homolog of the *PLAG1* transcription factor, whose translocation-driven overexpression defines pleomorphic adenoma of the salivary glands [74]. In the embryonic brain and in brain tumors, *Plagl2* regulates the balance between differentiation and proliferation, and its overexpression shifts neural progenitors towards proliferation [75, 76]. Outside the brain, *PLAGL2* also contributes to oncogenesis in lung cancer [77, 78], colorectal cancer [79-82], gastric cancer [83], breast cancer [84, 85], and acute myeloid leukemia [86]. In these cancers, *PLAGL2* has been shown to contribute to proliferation, epithelial-mesenchymal transition, and "stemness" by regulating target genes such as *Igf2* and regulators of Wnt signaling [73, 76, 79, 80, 82, 85]. To our knowledge, however, this is the first description of *PLAGL2* upregulation in *DROSHA*- or *DICER1*-related pediatric cancers.

In this study, we identify two small molecules that impaired tumor proliferation by blocking signaling pathways normally regulated by microRNAs. Although they are both clinically approved, it remains to be seen whether they will be effective in children with these cancers. In our models, neither drug was cytotoxic, and they may need to be combined with other therapies to harness clinical benefit. For instance, previous studies have found synergy between CDK4/6 inhibitors and immune checkpoint blockade, and we also found that palbociclib upregulated immune gene signatures [46-49] (**Suppl. Fig. S4E**). Alternatively, human tumors may harness other oncogenic signaling pathways or growth factors, such as RAS/MAP kinase; indeed, RAS pathway genes are the most frequently mutated genes in *DICER1*-related tumors after *TP53* [87-89]. Future studies will be needed to identify the mitogenic signaling pathways used in human pineoblastomas. Nevertheless, our results demonstrate the potential for the signals upregulated by microRNA loss to be therapeutic targets in cancers driven by impairment of *DROSHA* or *DICER1*.

METHODS

RNA-seq

Flash frozen mouse tissue was disrupted using an electric pestle and homogenized using a Qiashredder column (Qiagen, cat. no. 79654). Total RNA was extracted using the miRNeasy mini kit (Qiagen, cat. no. 217004). For IPDrosha and IPRb1 pineal tumors, single-end whole transcriptome sequencing was performed at the UT Southwestern McDermott Next Generation Sequencing (NGS) Core. For palbociclib and ceritinib treated tumor allografts, single-end strand-specific mRNA sequencing was performed at the UT Southwestern McDermott Next Generation Sequencing (NGS) Core. For IPDrosha tumors, matched normal brain, and age-matched normal adult pineal glands, paired-end whole transcriptome sequencing was performed at DNA Link Inc. For control samples from normal adult pineal glands, pineal glands were pooled from two age and genotype matched adults to ensure sufficient RNA sample for sequencing.

Reads from each sample were mapped to the reference genome using STAR (v2.5.3a) [90], and read counts were generated using featureCounts [91]. Differential gene expression calculations were then computed from raw counts using the DESeq2 package (v1.42.0) [92]. The Wald statistic output for protein-coding genes from DESeq2 was used as input for gene set enrichment analysis (GSEA) using the fgsea package (v1.28.0) [93]. Hallmark gene sets were derived from MSigDB v2023.1.Mm or v2023.1.Hs for mouse or human, respectively [94]. Predicted microRNA target genes were defined as genes with Aggregate $P_{CT} \geq 0.9$ and Total context++ score < -0.4 based on TargetScan (v8.0) [38].

For external data, RNA-seq FPKM tables from rat pineal glands across different stages of development were downloaded from GEO dataset GSE46127 [95]. Differential expression was calculated using t-tests after log_2 transformation. Marker genes at each timepoint were defined as the 300 most overexpressed protein-coding genes with mouse homologs, compared to other timepoints. RNA-seq counts tables from pineoblastoma patient samples were obtained from the St. Jude Cloud data repository (https://www.stjude.cloud) [96] and processed as above.

Quantitative PCR

Flash-frozen mouse tissue was disrupted using an electric pestle and homogenized using a Qiashredder column (Qiagen, cat. no. 79654). RNA was extracted using the miRNeasy

mini kit (Qiagen, cat. no. 217004), and cDNA was prepared using RT^2 HT First Strand kit (Qiagen, cat. no. 330411) or iScript Reverse Transcription Supermix for RT-qPCR (Bio-Rad, cat. no. 1708840). iTaq Universal SYBR Green Supermix (Bio-Rad, cat. no. 1725125) was used for qPCR in four replicates per condition. The oligos used for qPCR are referenced in **Supplementary Table 1**.

For miRNA TaqMan qPCR, total RNA was extracted using the miRNeasy mini kit (Qiagen, cat. no. 217004). cDNA synthesis was performed with the TaqMan MicroRNA Reverse Transcription kit (Applied Biosystems, cat. no. 4366596). The following assays were used for cDNA synthesis and qPCR detection (ThermoFisher cat. no. PN4427975): hsa-let-7a (Thermo Fisher, assay id 000377), hsa-let-7b (Thermo Fisher, assay id 002619), hsa-let-7i (Thermo Fisher, assay id 002221), hsa-miR-100 (Thermo Fisher, assay id 000437), mmu-miR-183 (Thermo Fisher, assay id 002269), and U6 snRNA (Thermo Fisher, assay id 001973). TaqMan Universal PCR Master Mix (Applied Biosystems, cat. no. 4304437) was used for qPCR, and each sample had three replicates per condition. Each miRNA assay was normalized to U6 and displayed as $2^{-\Delta\Delta Ct}$, and p values were calculated using two-tailed *t* tests.

Small RNA sequencing

For IPDrosha and IPRb1 tumors, total RNA was submitted for small RNA sequencing to the UT Southwestern McDermott NGS Core using the NEBNext Small RNA kit (New England Biolabs, cat. no. E7330), with size selection between 15bp and 30bp. Reads were aligned and counted using miRge3.0 [97]. Filtered reads aligning to microRNAs were normalized to reads aligning to tRNAs and snoRNAs as an internal control.

For small RNA sequencing and analysis from IPDrosha tumors, pineal gland, and brain samples, total RNA was submitted to DNA Link. QIAseq miRNA Library QC Spike-ins (Qiagen cat. no. 331535) were used as a spike-in control. Analysis was performed using mirDeep2 [98] for alignment, quantification, and annotation. Counts for each microRNA were normalized to spike-in control, and microRNAs of the same family were grouped per TargetScan [38]. Known mirtrons were annotated from mirtronDB [99]. Differential expression for individual microRNAs was performed using t-tests of log$_2$-transformed, normalized read counts.

CUT&RUN

Pineal tumors from IPDrosha/IPDicer1 mice were dissociated into single-cell suspension

using the Brain Tumor Dissociation Kit (P) (Miltenyi Biotec, cat. no. 130-095-942), and then "Cleavage Under Targets and Release Using Nuclease" (CUT&RUN) [100, 101] libraries were prepared from dissociated tumor cells using the ChIC/CUT&RUN Kit (EpiCypher, cat. no. 14-1048) according to manufacturer recommendations. Briefly, single-cell suspensions were immobilized onto beads, permeabilized, and incubated in Epicypher antibody buffer at 1:50 for H3K4me3, H3K27ac, or Rabbit IgG. (See **Supplementary Table 2** for more details.) Target-DNA complexes were cleaved using pAG-MNase, and libraries were prepared using NEBNext Ultra II DNA Library Kit from Illumina (New England Biolabs, cat. no. E7645S).

Library quality checks and sequencing were performed at the UT Southwestern McDermott NGS Core. Sequencing reads were analyzed using the nf-core/cutandrun bioinformatic pipeline v2.0 [102] on the UT Southwestern BioHPC Astrocyte platform. Briefly, after trimmed reads are aligned to the target genome and the spike-in using Bowtie 2 (v2.4.4) [103], peaks are called using the SEACR (v1.3) [104] and MACS2 (v2.2.7) [105] pipelines. Downstream visualization is performed using the Integrative Genomics Viewer (IGV, v2.17.1) [106], deepTools (v3.5.1) [107], or Gviz (v1.46.1) [108] toolkits.

miR-eCLIP sequencing

Two frozen IPRb1 tumor fragments underwent miR-eCLIP sequencing at EclipseBio, as previously described [109]. Briefly, Ago2 was immunoprecipitated from crosslinked RNA. After ligation of miRNA-mRNA chimeras, input and immunoprecipitated RNA were analyzed by sequencing on the Illumina NovaSeq 6000 platform. Chimeric and non-chimeric eCLIP sequences were processed using the pipeline described [109] and compared to input in each sample using the CLIPper algorithm (v2.0) [110]. Chimeric and non-chimeric peaks were compared between the two samples to identify reproducible peaks. Reproducible non-chimeric peaks were those enriched in each IP with a $\log_2$-fold-change ≥ 3 and p value ≤ 0.001. HOMER *de novo* motif analysis [111] was used to identify enriched motifs in these peaks. Reproducible microRNA-miRNA chimeras were those with ≥3 chimeric reads and a reproducibility index ≥ 0.5. Seed matches were then limited to perfect-match, canonical 7- or 8-mers (based on TargetScan definitions [38]) in the 3'UTRs of their target genes.

Immunohistochemistry

Formalin-fixed tissues were processed at the UT Southwestern Tissue Management Shared Resource or Mouse Histology core, and fixed sections were deparaffinized using Histo-

Clear II (VWR, cat. no. 101412-884) or xylene (Sigma, cat. no. 534056) with gradient ethanol washes. Heat antigen retrieval was performed using Trilogy (Sigma, cat. no. 920P) or 1M citrate buffer pH 6.0. Tissue slides were stained with Ki-67, phospho-Rb, synaptophysin, phospho-S6, and GFAP (see **Supplementary Table 2** for dilutions and catalog numbers). Signal was obtained using diaminobenzadine (DAB) (Millipore, cat. no. 281751), and hematoxylin (Sigma Aldrich, cat. no. GHS 132-1L) was used for counterstaining.

ChIP-qPCR

ChIP-qPCR was performed using the ChIP-IT High Sensitivity kit (Active Motif, cat. no. 53040). Flash-frozen tumors were fixed with 1% formaldehyde, and 6.75 µg chromatin was sonicated using a Diagenode Bioruptor and then immunoprecipitated with 2.5 µg of primary antibodies recognizing Plagl2 (Sigma Aldrich, cat. no. SAB3500815) or Rabbit IgG (Cell Signalling Technology, cat. no. 2729). Control input chromatin was also prepared with the same kit. Quantitative PCR was performed with iTaq Universal SYBR Green Supermix (Bio-Rad, cat. no. 1725125), and each sample had three replicates per condition. The oligos used for qPCR are shown in **Supplementary Table 1**. Results were calculated as percent input, and p values were calculated using two-tailed *t* tests.

Animal breeding and husbandry

Mice of the following strains were obtained from Jackson laboratories: Irbp-Cre (also known as Rbp3-Cre, cat. no. 003967, RRID IMSR_JAX:003967), $Rb1^{flox}$ (cat. no. 026563, RRID IMSR_JAX:026563), $Drosha^{flox}$ (cat. no. 008592, RRID IMSR_JAX:008592), $Dicer1^{flox}$ (cat. no. 006366, IMSR_JAX:006366) and $p53^{flox}$ (cat. no. 008462, RRID IMSR_JAX:008462). These were maintained on a C57BL6/J background. Nod-scid gamma (NSG) mice were also purchased from Jackson laboratories (cat. no. 005557, RRID IMSR_JAX:005557) and maintained under the UT Southwestern Immunocompromised Breeding Program. Animal experiments were performed under oversight from the University of Texas Southwestern Institutional Animal Care and Use Committee protocols 2019-102689 and 2019-102680. Mice were monitored visually twice weekly for a domed head and given moist chow upon tumor development. Tumors were harvested 3 to 5 days later. To compare tumor-free survival of mice with different genotypes, p values were calculated using the log-rank test.

Allografts and in vivo drug treatments

Pineal tumors from IPDrosha/IPDicer1 mice were dissociated using the Brain Tumor Dissociation Kit (P) (Miltenyi Biotec, cat. no. 130-095-942). Male and female NSG mice at 6-8 weeks of age were subcutaneously injected with 0.5-1.5 million tumor cells, suspended in 50% cold PBS and 50% matrigel (Corning, cat. no. CB-40234). Tumor volume was measured by calipers twice a week. When tumors reached 150 mm^3, mice were administered palbociclib (150 mg/kg, LC Laboratories, cat. no. P-7788) or an equivalent volume of vehicle (50 mM sodium L-lactate, Sigma cat. no. 71718) by oral gavage, five days a week for up to 3 weeks. Tumors were harvested when they reached 1,000 mm^3, 4 hours after the last dose of drug, and rinsed in PBS before flash-freezing or fixation. Tumor sizes were compared using two-tailed t-tests at each timepoint, and survival was compared using the log-rank test.

For ceritinib treatment, pineal tumor cells were injected into NSG mice as described above. Once tumors reached 150 mm^3, mice were administered ceritinib (50 mg/kg, Sellekchem, cat. no. S4967) or an equivalent volume of vehicle (0.5% methylcellulose and 0.5% Tween-80, Fisher Scientific cat. no. AC428430500 & Sigma cat. no. P8192) by oral gavage, five days a week for up to 3 weeks. Tumors were harvested when they reached 1,000 mm^3, 3 hours after the last dose of drug, and rinsed in PBS before flash-freezing or fixation.

Data availability

Raw and processed sequencing data are available at the GEO SuperSeries GSE255914.

ACKNOWLEDGMENTS

This work was supported by funding from the V Foundation for Cancer Research (V2022-010 to K.S.C.); National Cancer Institute (K08CA207849 to K.S.C. and Cancer Center Support Grant P30CA142543); and Cancer Prevention and Research Institute of Texas (RR180071 to K.S.C., RP210041 to C.F.). This research used computational resources provided by the BioHPC supercomputing facility in the Lyda Hill Department of Bioinformatics at UT Southwestern, which is supported by Cancer Prevention and Research Institute of Texas (RP150596), and tissue processing at the UT Southwestern Simmons Comprehensive Cancer Center Tissue Management Shared Resource, which is supported by the National Cancer Institute (P30CA142543). We appreciate the past, current, and future mentoring from James Amatruda, Joshua Mendell, Stephen Skapek, and many others.

AUTHOR CONTRIBUTIONS

C.R.F., K.D, L.K.M., U.O., T.S., K.O., and K.S.C. conceived and performed experiments. C.R.F., Y.W., and K.S.C. wrote the manuscript. V.R. provided expertise, feedback, and analysis. J.L. and K.S.C. performed data analysis and secured funding.

FIGURE LEGENDS

Figure 1. Pineal tumors driven by loss of *Drosha, Dicer1,* or *Rb1* resemble embryonic pineal progenitors.

(A) Kaplan-Meier plots showing tumor-free survival of IPDrosha, IPDicer1, and IPRb1 tumors. IrbpCre;Drosha$^{f/f}$;p53$^{f/f}$ (IPDrosha, dark green) mice have a tumor incidence of 36% at 1 year; IrbpCre;Dicer1$^{f/f}$;p53$^{f/f}$ (IPDicer1, light green) mice have a tumor incidence of 34% at 1 year; IrbpCre;Rb1$^{f/f}$;p53$^{f/f}$ (IPRb1, purple) mice have a tumor incidence of 90% at 1 year (***p<0.001 and ****p<0.0001; vs. control genotypes; by log-rank test). **(B)** Photomicrographs of brains from control (Drosha$^{f/f}$;p53$^{f/f}$), IPDrosha, IPDicer1, and IPRb1 mice show expansile pineal tumors expanding into the skull above and normal brain below (scale bar 1000 μm). **(C)** Photomicrographs of IPDrosha, IPDicer1, and IPRb1 tumors (scale bar 50 μm). Top row shows H&E, and bottom row shows immunostaining for Ki-67. **(D)** Photomicrographs of IPDrosha, IPDicer1, and IPRb1 tumors (scale bar 50 μm), with immunostaining for synaptophysin (top) and GFAP (bottom). **(E)** Gene set enrichment analysis (GSEA) of enrichment for pineal development markers comparing IPDrosha tumor with normal adult pineal gland. **(F)** GSEA of enrichment for pineal development markers comparing IPRb1 tumor with normal adult pineal gland. **(G)** qPCR for pineal homeobox genes (*Otx2, Crx, Pax6*, and *Rax*) in brain, IPDrosha, IPDicer1, and IPRb1 tumors. Values shown are mean ± SD (**p < 0.01 and **p < 0.001; vs. normal brain; two-sided t-test). See also Supplementary Figure S1.

Figure 2. *Drosha*-driven tumors lack microRNAs and overexpress microRNA target genes.

(A) Small RNA sequencing displaying the most abundant microRNA families in IPRb1 tumors and IPDrosha tumors. **(B)** Expression profile of individual let-7/miR-98-5p family microRNAs in IPRb1 and IPDrosha tumors. **(C)** Small RNA sequencing displaying the most abundant microRNA families in adult pineal glands and IPDrosha tumors. **(D)** Expression profile of let-7/miR-98-5p family microRNAs in IPDrosha tumors and adult pineal glands. **(E)** TaqMan miRNA expression of let-7a-5p, let-7d-5p, let-7i-5p, miR-100-5p, and miR-183-5p in normal adult brain, IPDrosha, IPDicer1, and IPRb1 tumors. **(F,G)** GSEA of enrichment for predicted targets of all microRNAs expressed in IPRb1 tumors **(F)** or predicted let-7/miR-98-5p targets **(G)** in IPDrosha

vs. IPRb1 tumors. **(H,I)** GSEA for enrichment of confirmed top targets of microRNAs **(H)** or let-7/miR-98-5p **(I)** in IPDrosha vs. IPRb1 tumors. See also Supplementary Figures S2 and S3.

Figure 3. Proliferation restrains differentiation in *Drosha*-driven tumors.

(A) Hallmark gene sets most enriched in IPDrosha tumors vs. age-matched adult pineal gland, IPRb1 tumors vs. age-matched adult pineal gland, and IPDrosha tumors vs. IPRb1 tumors. **(B)** qPCR for *Ccnd2* in normal adult brain, IPDrosha, IPDicer1, and IPRb1 tumors. Values shown are mean ± SD (*p< 0.05; vs. normal brain; two-sided t-test). **(C)** Photomicrographs of IPDrosha, IPDicer1, and IPRb1 tumors stained for phospho-Rb1 (Ser807/811) (scale bar 50 μm). **(D)** Tumor volumes for IPDrosha or IPDicer1 tumors in mice dosed with vehicle or palbociclib. Values shown are mean ± SD (*p<0.05 and *p<0.01; vs. vehicle-treated; two-sided t-test). **(E)** Photomicrographs of vehicle- or palbociclib-treated IPDrosha and IPDicer1 tumors (scale bar 50 μm). Top row shows H&E, and bottom row shows immunostaining for phospho-Rb1 (Ser807/811). **(F)** GSEA for E2F target genes in RNA sequencing of IPDrosha tumor allografts treated with palbociclib vs. vehicle. **(G)** GSEA of enrichment for pineal development genes in IPDrosha tumors treated with palbociclib vs. vehicle. **(H)** qPCR for adult pineal marker genes (*Chrna3* and *Nefh*) and an embryonic pineal marker gene (*Plagl2*) in IPDrosha and IPDicer1 tumors treated with palbociclib vs. vehicle. Values shown are mean ±SD (*p<0.05; vs. vehicle-treated; two-sided t-test). See also Supplementary Figure S4.

Figure 4. The microRNA target gene *Plagl2* regulates proliferation drivers *Igf2* and *Ccnd2*.

(A) qPCR for *Igf2* and *Plagl2* in normal adult brain, IPDrosha, IPDicer1, and IPRb1 tumors. Values shown are mean ± SD (*p< 0.05 and ***p<0.001; vs. normal adult brain; two-sided t-tests). **(B)** Heat map from total RNA sequencing of IPDrosha and IPRb1 tumors highlighting *Igf2*. **(C)** miR-eCLIP chimeric reads from IPRb1 tumors of the 3' UTR of *Plagl2,* with microRNA chimeric read peaks denoted below, including three peaks containing let-7/miR-98-5p family members. The conservation of the second let-7/miR-98-5p binding site (marked with asterisk) is featured on the right. **(D)** H3K27ac and H3K4me3 signal at the *Ccnd2* promoter in IPDrosha and IPRb1 tumors, with predicted Plagl2 binding sites and regions targeted by qPCR marked below. **(E)** ChIP-qPCR detecting Plagl2 occupancy in IPDrosha and IPRb1 tumors at the mouse *Ccnd2*

promoter, *Igf2* promoter, and a negative-control intergenic region. Values shown are mean ± SD (*p<0.05 and *p<0.01; vs. IgG; two-sided t-test). See also Supplementary Figure S5.

Figure 5. The Igf inhibitor ceritinib suppresses tumor growth.

(A) GSEA of enrichment for insulin signaling in IPDrosha vs. IPRb1 tumors (Wikipathway WP481). **(B)** Tumor volumes for IPDrosha and IPDicer1 tumors in mice dosed with vehicle or ceritinib. Values shown are mean ± SD (*p< 0.05; vs. vehicle treated; two-sided t-test). **(C)** Photomicrographs of vehicle- or ceritinib-treated IPDrosha and IPDicer1 tumors (scale 50 µm). Top row shows H&E, the middle row shows phospho-S6 (Ser235/236), and the bottom row shows phospho-Rb1 (Ser807/811). **(D)** GSEA of enrichment for insulin signaling gene set in IPDrosha tumors treated with ceritinib vs. vehicle.

REFERENCES

1.   Grobner, S.N., et al., *The landscape of genomic alterations across childhood cancers.* Nature, 2018. **555**(7696): p. 321-327.

2.   Ma, X., et al., *Pan-cancer genome and transcriptome analyses of 1,699 paediatric leukaemias and solid tumours.* Nature, 2018. **555**(7696): p. 371-376.

3.   Rakheja, D., et al., *Somatic mutations in DROSHA and DICER1 impair microRNA biogenesis through distinct mechanisms in Wilms tumours.* Nat Commun, 2014. **2**: p. 4802.

4.   Torrezan, G.T., et al., *Recurrent somatic mutation in DROSHA induces microRNA profile changes in Wilms tumour.* Nat Commun, 2014. **5**: p. 4039.

5.   Walz, A.L., et al., *Recurrent DGCR8, DROSHA, and SIX homeodomain mutations in favorable histology Wilms tumors.* Cancer Cell, 2015. **27**(2): p. 286-97.

6.   Wegert, J., et al., *Mutations in the SIX1/2 pathway and the DROSHA/DGCR8 miRNA microprocessor complex underlie high-risk blastemal type Wilms tumors.* Cancer Cell, 2015. **27**(2): p. 298-311.

7.   Gadd, S., et al., *A Children's Oncology Group and TARGET initiative exploring the genetic landscape of Wilms tumor.* Nat Genet, 2017. **49**(10): p. 1487-1494.

8.   Hill, D.A., et al., *DICER1 mutations in familial pleuropulmonary blastoma.* Science, 2009. **325**(5943): p. 965.

9.   Brenneman, M., et al., *Temporal order of RNase IIIb and loss-of-function mutations during development determines phenotype in pleuropulmonary blastoma / DICER1 syndrome: a unique variant of the two-hit tumor suppression model.* F1000Res, 2015. **4**: p. 214.

10.   de Kock, L., M.K. Wu, and W.D. Foulkes, *Ten years of DICER1 mutations: Provenance, distribution, and associated phenotypes.* Hum Mutat, 2019. **40**(11): p. 1939-1953.

11.   Snuderl, M., et al., *Recurrent homozygous deletion of DROSHA and microduplication of PDE4DIP in pineoblastoma.* Nat Commun, 2018. **9**(1): p. 2868.

12.   Bartel, D.P., *MicroRNAs: genomics, biogenesis, mechanism, and function.* Cell, 2004. **116**(2): p. 281-97.

13.   Flynt, A.S. and E.C. Lai, *Biological principles of microRNA-mediated regulation: shared themes amid diversity.* Nat Rev Genet, 2008. **9**(11): p. 831-42.

14.   Hansford, J.R., et al., *Pediatric pineoblastoma: A pooled outcome study of North American and Australian therapeutic data.* Neurooncol Adv, 2022. **4**(1): p. vdac056.

15.   de Kock, L., B. Rivera, and W.D. Foulkes, *Pineoblastoma is uniquely tolerant of mutually exclusive loss of DICER1, DROSHA or DGCR8.* Acta Neuropathol, 2020. **139**(6): p. 1115-1118.

16.   Li, B.K., et al., *Pineoblastoma segregates into molecular sub-groups with distinct clinico-pathologic features: a Rare Brain Tumor Consortium registry study.* Acta Neuropathol, 2020. **139**(2): p. 223-241.

17.   Liu, A.P.Y., et al., *Risk-adapted therapy and biological heterogeneity in pineoblastoma: integrated clinico-pathological analysis from the prospective, multi-center SJMB03 and SJYC07 trials.* Acta Neuropathol, 2020. **139**(2): p. 259-271.

18.   Pfaff, E., et al., *Molecular subgrouping of primary pineal parenchymal tumors reveals distinct subtypes correlated with clinical parameters and genetic alterations.* Acta Neuropathol, 2020. **139**(2): p. 243-257.

19.   Liu, A.P.Y., et al., *Clinical and molecular heterogeneity of pineal parenchymal tumors: a consensus study.* Acta Neuropathol, 2021. **141**(5): p. 771-785.

20.   Goel, S., J.S. Bergholz, and J.J. Zhao, *Targeting CDK4 and CDK6 in cancer.* Nat Rev Cancer, 2022. **22**(6): p. 356-372.

21.   Dang, C.V., *MYC on the path to cancer.* Cell, 2012. **149**(1): p. 22-35.

22.   Saab, R., et al., *Pharmacologic inhibition of cyclin-dependent kinase 4/6 activity arrests proliferation in myoblasts and rhabdomyosarcoma-derived cells.* Mol Cancer Ther, 2006. **5**(5): p. 1299-308.

23.   Saab, R., et al., *p18Ink4c and p53 Act as tumor suppressors in cyclin D1-driven primitive neuroectodermal tumor.* Cancer Res, 2009. **69**(2): p. 440-8.

24.   Harajly, M., et al., *p53 Restoration in Induction and Maintenance of Senescence: Differential Effects in Premalignant and Malignant Tumor Cells.* Mol Cell Biol, 2016. **36**(3): p. 438-51.

25.   Idriss, S., et al., *A temporal in vivo catalog of chromatin accessibility and expression profiles in pineoblastoma reveals a prevalent role for repressor elements.* Genome Res, 2023. **33**(2): p. 269-282.

26.   Chung, P.E.D., et al., *Modeling germline mutations in pineoblastoma uncovers lysosome disruption-based therapy.* Nat Commun, 2020. **11**(1): p. 1825.

27.   Vooijs, M., et al., *Tumor formation in mice with somatic inactivation of the retinoblastoma gene in interphotoreceptor retinol binding protein-expressing cells.* Oncogene, 2002. **21**(30): p. 4635-45.

28. Rodrigues, M.M., et al., *Interphotoreceptor retinoid-binding protein in retinal rod cells and pineal gland.* Invest Ophthalmol Vis Sci, 1986. **27**(5): p. 844-50.

29. Marszalek, J.R., et al., *Genetic evidence for selective transport of opsin and arrestin by kinesin-II in mammalian photoreceptors.* Cell, 2000. **102**(2): p. 175-87.

30. Rath, M.F., et al., *Homeobox genes in the rodent pineal gland: roles in development and phenotype maintenance.* Neurochem Res, 2013. **38**(6): p. 1100-12.

31. Whyte, W.A., et al., *Master transcription factors and mediator establish super-enhancers at key cell identity genes.* Cell, 2013. **153**(2): p. 307-19.

32. Wen, J., et al., *Analysis of Nearly One Thousand Mammalian Mirtrons Reveals Novel Features of Dicer Substrates.* PLoS Comput Biol, 2015. **11**(9): p. e1004441.

33. Kim, Y.K., B. Kim, and V.N. Kim, *Re-evaluation of the roles of DROSHA, Export in 5, and DICER in microRNA biogenesis.* Proc Natl Acad Sci U S A, 2016. **113**(13): p. E1881-9.

34. Roush, S. and F.J. Slack, *The let-7 family of microRNAs.* Trends Cell Biol, 2008. **18**(10): p. 505-16.

35. La Torre, A., S. Georgi, and T.A. Reh, *Conserved microRNA pathway regulates developmental timing of retinal neurogenesis.* Proc Natl Acad Sci U S A, 2013. **110**(26): p. E2362-70.

36. Chawla, G., et al., *A let-7-to-miR-125 MicroRNA Switch Regulates Neuronal Integrity and Lifespan in Drosophila.* PLoS Genet, 2016. **12**(8): p. e1006247.

37. Wu, Y.C., et al., *Let-7-complex microRNAs regulate the temporal identity of Drosophila mushroom body neurons via chinmo.* Dev Cell, 2012. **23**(1): p. 202-9.

38. Agarwal, V., et al., *Predicting effective microRNA target sites in mammalian mRNAs.* Elife, 2015. **4**: p. e05005.

39. Manakov, S.A., et al., *Scalable and deep profiling of mRNA targets for individual microRNAs with chimeric eCLIP.* bioRxiv, 2022: p. 2022.02.13.480296.

40. Fassl, A., Y. Geng, and P. Sicinski, *CDK4 and CDK6 kinases: From basic science to cancer therapy.* Science, 2022. **375**(6577): p. eabc1495.

41. Zalzali, H., et al., *Temporally distinct roles for tumor suppressor pathways in cell cycle arrest and cellular senescence in Cyclin D1-driven tumor.* Mol Cancer, 2012. **11**: p. 28.

42. Kawano, M., et al., *c-Myc Represses Tumor-Suppressive microRNAs, let-7a, miR-16 and miR-29b, and Induces Cyclin D2-Mediated Cell Proliferation in Ewing's Sarcoma Cell Line.* PLoS One, 2015. **10**(9): p. e0138560.

43.   Lovat, F., et al., *miR-15b/16-2 deletion promotes B-cell malignancies.* Proc Natl Acad Sci U S A, 2015. **112**(37): p. 11636-41.

44.   Liu, L., et al., *The cell cycle in stem cell proliferation, pluripotency and differentiation.* Nat Cell Biol, 2019. **21**(9): p. 1060-1067.

45.   Fry, D.W., et al., *Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991 and associated antitumor activity in human tumor xenografts.* Mol Cancer Ther, 2004. **3**(11): p. 1427-38.

46.   Goel, S., et al., *CDK4/6 inhibition triggers anti-tumour immunity.* Nature, 2017. **548**(7668): p. 471-475.

47.   Deng, J., et al., *CDK4/6 Inhibition Augments Antitumor Immunity by Enhancing T-cell Activation.* Cancer Discov, 2018. **8**(2): p. 216-233.

48.   Zhang, J., et al., *Cyclin D-CDK4 kinase destabilizes PD-L1 via cullin 3-SPOP to control cancer immune surveillance.* Nature, 2018. **553**(7686): p. 91-95.

49.   Schaer, D.A., et al., *The CDK4/6 Inhibitor Abemaciclib Induces a T Cell Inflamed Tumor Microenvironment and Enhances the Efficacy of PD-L1 Checkpoint Blockade.* Cell Rep, 2018. **22**(11): p. 2978-2994.

50.   Strubberg, A.M., et al., *The Zinc Finger Transcription Factor PLAGL2 Enhances Stem Cell Fate and Activates Expression of ASCL2 in Intestinal Epithelial Cells.* Stem Cell Reports, 2018. **11**(2): p. 410-424.

51.   Alam, S., et al., *Members of the Plag gene family are expressed in complementary and overlapping regions in the developing murine nervous system.* Dev Dyn, 2005. **234**(3): p. 772-82.

52.   Shenoy, A., M. Danial, and R.H. Blelloch, *Let-7 and miR-125 cooperate to prime progenitors for astrogliogenesis.* EMBO J, 2015. **34**(9): p. 1180-94.

53.   Werner, H. and D. LeRoith, *Hallmarks of cancer: The insulin-like growth factors perspective.* Front Oncol, 2022. **12**: p. 1055589.

54.   Kuenzi, B.M., et al., *Polypharmacology-based ceritinib repurposing using integrated functional proteomics.* Nat Chem Biol, 2017. **13**(12): p. 1222-1231.

55.   Ibanez Rodriguez, M.P., S.C. Noctor, and E.M. Munoz, *Cellular Basis of Pineal Gland Development: Emerging Role of Microglia as Phenotype Regulator.* PLoS One, 2016. **11**(11): p. e0167063.

56.   Santagata, S., et al., *CRX is a diagnostic marker of retinal and pineal lineage tumors.* PLoS One, 2009. **4**(11): p. e7932.

57.   Chen, S., et al., *Crx, a novel Otx-like paired-homeodomain protein, binds to and transactivates photoreceptor cell-specific genes.* Neuron, 1997. **19**(5): p. 1017-30.

58.   Nishida, A., et al., *Otx2 homeobox gene controls retinal photoreceptor cell fate and pineal gland development.* Nat Neurosci, 2003. **6**(12): p. 1255-63.

59.   Kimura, A., et al., *Both PCE-1/RX and OTX/CRX interactions are necessary for photoreceptor-specific gene expression.* J Biol Chem, 2000. **275**(2): p. 1152-60.

60.   Boatright, J.H., et al., *Endogenous CRX expression and IRBP promoter activity in retinoblastoma cells.* Brain Res, 2001. **916**(1-2): p. 136-42.

61.   Adamson, D.C., et al., *OTX2 is critical for the maintenance and progression of Shh-independent medulloblastomas.* Cancer Res, 2010. **70**(1): p. 181-91.

62.   Garancher, A., et al., *NRL and CRX Define Photoreceptor Identity and Reveal Subgroup-Specific Dependencies in Medulloblastoma.* Cancer Cell, 2018. **33**(3): p. 435-449 e6.

63.   Wagner, K.U., et al., *Cre-mediated gene deletion in the mammary gland.* Nucleic Acids Res, 1997. **25**(21): p. 4323-30.

64.   Pu, D., et al., *Efficacy of CDK4/6 inhibitors combined with endocrine therapy in HR+/HER2- breast cancer: an umbrella review.* J Cancer Res Clin Oncol, 2024. **150**(1): p. 16.

65.   Schubert, N.A., et al., *Target actionability review to evaluate CDK4/6 as a therapeutic target in paediatric solid and brain tumours.* Eur J Cancer, 2022. **170**: p. 196-208.

66.   Ferguson, K.M., et al., *Palbociclib releases the latent differentiation capacity of neuroblastoma cells.* Dev Cell, 2023. **58**(19): p. 1967-1982 e8.

67.   Vo, T., et al., *Spatial transcriptomic analysis of Sonic hedgehog medulloblastoma identifies that the loss of heterogeneity and promotion of differentiation underlies the response to CDK4/6 inhibition.* Genome Med, 2023. **15**(1): p. 29.

68.   Wu, C., et al., *PARP and CDK4/6 Inhibitor Combination Therapy Induces Apoptosis and Suppresses Neuroendocrine Differentiation in Prostate Cancer.* Mol Cancer Ther, 2021. **20**(9): p. 1680-1691.

69.   Gong, X., et al., *Genomic Aberrations that Activate D-type Cyclins Are Associated with Enhanced Sensitivity to the CDK4 and CDK6 Inhibitor Abemaciclib.* Cancer Cell, 2017. **32**(6): p. 761-776 e6.

70.   Xu, L., et al., *Wilms Tumor Mutational Subclasses Converge to Drive Ccnd2 Overexpression.* medRxiv, 2023.

71.   Chen, K.S., et al., *Mutations in microRNA processing genes in Wilms tumors derepress the IGF2 regulator PLAG1.* Genes Dev, 2018. **32**(15-16): p. 996-1007.

72.   Wang, Y., et al., *Biallelic Dicer1 Mutations in the Gynecologic Tract of Mice Drive Lineage-Specific Development of DICER1 Syndrome-Associated Cancer.* Cancer Res, 2023. **83**(21): p. 3517-3528.

73.   Keck, M.K., et al., *Amplification of the PLAG-family genes-PLAGL1 and PLAGL2-is a key feature of the novel tumor type CNS embryonal tumor with PLAGL amplification.* Acta Neuropathol, 2023. **145**(1): p. 49-69.

74.   Hensen, K., et al., *The tumorigenic diversity of the three PLAG family members is associated with different DNA binding capacities.* Cancer Research, 2002. **62**(5): p. 1510-1517.

75.   Adnani, L., et al., *Plag1 and Plagl2 have overlapping and distinct functions in telencephalic development.* Biol Open, 2018. **7**(11).

76.   Zheng, H., et al., *PLAGL2 regulates Wnt signaling to impede differentiation in neural stem cells and gliomas.* Cancer Cell, 2010. **17**(5): p. 497-509.

77.   Yang, Y.S., M.C. Yang, and J.C. Weissler, *Pleiomorphic adenoma gene-like 2 expression is associated with the development of lung adenocarcinoma and emphysema.* Lung Cancer, 2011. **74**(1): p. 12-24.

78.   Lin, Y., et al., *PLAGL2 promotes the stemness and is upregulated by transcription factor E2F1 in human lung cancer.* Environ Toxicol, 2023. **38**(4): p. 941-949.

79.   Fischer, A.D., et al., *The oncogenic function of PLAGL2 is mediated via ASCL2 and IGF2 and a Wnt-independent mechanism in colorectal cancer.* Am J Physiol Gastrointest Liver Physiol, 2023. **325**(2): p. G196-G211.

80.   Li, N., et al., *Overexpressed PLAGL2 transcriptionally activates Wnt6 and promotes cancer development in colorectal cancer.* Oncol Rep, 2019. **41**(2): p. 875-884.

81.   Li, D., et al., *PLAGL2 and POFUT1 are regulated by an evolutionarily conserved bidirectional promoter and are collaboratively involved in colorectal cancer by maintaining stemness.* EBioMedicine, 2019. **45**: p. 124-138.

82.   Liu, X., et al., *DNA-methylation-mediated silencing of miR-486-5p promotes colorectal cancer proliferation and migration through activation of PLAGL2/IGF2/beta-catenin signal pathways.* Cell Death Dis, 2018. **9**(10): p. 1037.

83.   Chen, W., et al., *PLAGL2 promotes Snail expression and gastric cancer progression via UCA1/miR-145-5p/YTHDF1 axis.* Carcinogenesis, 2023. **44**(4): p. 328-340.

84. Xu, B., et al., *MiR-449a suppresses cell migration and invasion by targeting PLAGL2 in breast cancer.* Pathol Res Pract, 2018. **214**(5): p. 790-795.

85. Li, Y., et al., *PLAGL2 increases adriamycin resistance and EMT in breast cancer cells by activating the Wnt pathway.* Genes Genomics, 2023. **45**(1): p. 49-57.

86. Landrette, S.F., et al., *Plag1 and Plagl2 are oncogenes that induce acute myeloid leukemia in cooperation with Cbfb-MYH11.* Blood, 2005. **105**(7): p. 2900-7.

87. Pugh, T.J., et al., *Exome sequencing of pleuropulmonary blastoma reveals frequent biallelic loss of TP53 and two hits in DICER1 resulting in retention of 5p-derived miRNA hairpin loop sequences.* Oncogene, 2014. **33**(45): p. 5295-302.

88. Bae, J.S., et al., *High Prevalence of DICER1 Mutations and Low Frequency of Gene Fusions in Pediatric Follicular-Patterned Tumors of the Thyroid.* Endocr Pathol, 2021. **32**(3): p. 336-346.

89. Kommoss, F.K.F., et al., *Genomic characterization of DICER1-associated neoplasms uncovers molecular classes.* Nat Commun, 2023. **14**(1): p. 1677.

90. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.

91. Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program for assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-30.

92. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

93. Korotkevich, G., et al., *An algorithm for fast preranked gene set enrichment analysis using cumulative statistic calculation.* bioRxiv, 2021: p. 060012.

94. Liberzon, A., et al., *The Molecular Signatures Database (MSigDB) hallmark gene set collection.* Cell Syst, 2015. **1**(6): p. 417-425.

95. Yamazaki, F., et al., *The Lhx9 homeobox gene controls pineal gland development and prevents postnatal hydrocephalus.* Brain Struct Funct, 2015. **220**(3): p. 1497-509.

96. McLeod, C., et al., *St. Jude Cloud: A Pediatric Cancer Genomic Data-Sharing Ecosystem.* Cancer Discov, 2021. **11**(5): p. 1082-1099.

97. Patil, A.H. and M.K. Halushka, *miRge3.0: a comprehensive microRNA and tRF sequencing analysis pipeline.* NAR Genom Bioinform, 2021. **3**(3): p. lqab068.

98. Friedlander, M.R., et al., *miRDeep2 accurately identifies known and hundreds of novel microRNA genes in seven animal clades.* Nucleic Acids Res, 2012. **40**(1): p. 37-52.

99.   Da Fonseca, B.H.R., D.S. Domingues, and A.R. Paschoal, *mirtronDB: a mirtron knowledge base.* Bioinformatics, 2019. **35**(19): p. 3873-3874.

100.   Skene, P.J. and S. Henikoff, *An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites.* Elife, 2017. **6**: p. e21856.

101.   Meers, M.P., et al., *Improved CUT&RUN chromatin profiling tools.* Elife, 2019. **8**: p. e46314.

102.   Cheshire, C., et al., *nf-core/cutandrun: nf-core/cutandrun v3.2.2 Iridium Ibis (3.2.2).* Zenodo.

103.   Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat Methods, 2012. **9**(4): p. 357-9.

104.   Meers, M.P., D. Tenenbaum, and S. Henikoff, *Peak calling by Sparse Enrichment Analysis for CUT&RUN chromatin profiling.* Epigenetics Chromatin, 2019. **12**(1): p. 42.

105.   Zhang, Y., et al., *Model-based analysis of ChIP-Seq (MACS).* Genome Biol, 2008. **9**(9): p. R137.

106.   Robinson, J.T., et al., *Integrative genomics viewer.* Nat Biotechnol, 2011. **29**(1): p. 24-6.

107.   Ramirez, F., et al., *deepTools2: a next generation web server for deep-sequencing data analysis.* Nucleic Acids Res, 2016. **44**(W1): p. W160-5.

108.   Hahne, F. and R. Ivanek, *Visualizing Genomic Data Using Gviz and Bioconductor*, in *Statistical Genomics: Methods and Protocols*, E. Mathé and S. Davis, Editors. 2016, Springer New York: New York, NY. p. 335-351.

109.   Manakov, S.A., et al., *Scalable and deep profiling of mRNA targets for individual microRNAs with chimeric eCLIP.* bioRxiv, 2022: p. 2022.02.13.480296.

110.   Lovci, M.T., et al., *Rbfox proteins regulate alternative mRNA splicing through evolutionarily conserved RNA bridges.* Nat Struct Mol Biol, 2013. **20**(12): p. 1434-42.

111.   Heinz, S., et al., *Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities.* Mol Cell, 2010. **38**(4): p. 576-89.

Fig 1. Pineal tumors driven by loss of *Drosha*, *Dicer1*, or *Rb1* resemble embryonic pineal progenitors.

Fig 2. *Drosha*-driven tumors lack microRNAs and overexpress microRNA target genes.

Fig 3. Proliferation restrains differentiation in *Drosha*-driven tumors.

Fig 4. The microRNA target gene *Plagl2* regulates proliferation drivers *Igf2* and *Ccnd2*.

Fig 5. The Igf inhibitor ceritinib suppresses tumor growth.

Fig S1. IPDrosha and IPRb1 pineal tumors retain embryonic pineal expression pathways.

Fig S2. Drosha tumors exhibit de-repression of let-7/miR-98-5p target genes.

Fig S3. Patterns of microRNA regulation in miR-eCLIP sequencing of IPRb1 tumors.

A

Non-chimeric reads

Reproduced
4,362 peaks

Sample 1
5,640 peaks

Sample 2
4,669 peaks

Chimeric reads

Reproduced
13,524 peaks

Sample 1
16,765 peaks

Sample 2
14,987 peaks

B

Non-chimeric reads

5' UTR | CDS | 3' UTR

Chimeric reads

5' UTR | CDS | 3' UTR

Average peaks

Relative position on gene

Relative position on gene

— Reproducible peaks
— IPRb1, sample 1
— IPRb1, sample 2

C

Top motifs in non-chimeric reads

Motif	Seed match	% of peaks with motif	$-\log_{10}(p)$
HWGTGCCTA	miR-124-3p	8.85%	748.9
NHHCTACCTCAD	let-7/miR-98-5p	8.57%	745.6
ATCTACCTCA	let-7/miR-98-5p	6.54%	715.9
CTAAGTGCCTA	miR-124-3p	8.47%	709.2

D

Top microRNAs in chimeric reads

Average RPM

miR-124-3p, miR-466i-5p, miR-16-5p, let-7c-5p, let-7b-5p, miR-15a-5p, let-7a-5p, miR-19b-3p, let-7f-5p, miR-92a-3p, miR-30b-5p, miR-17-5p, miR-182-5p, miR-15b-5p, miR-30d-5p, miR-30c-5p, miR-125b-5p, miR-20a-5p, let-7d-5p

E

Expression of microRNA target genes

cumulative distribution function

$\log_2$ fold-change (IPDrosha vs. IPRb1)

— other genes (n=8,122)
— imperfect microRNA targets (n=381)
— top microRNA targets (n=1,454)

F

Differential expression of microRNA targets

genes upregulated in IPDrosha (n=2,642)

genes downregulated in IPDrosha (n=1,591)

$p = 8.7 \times 10^{-21}$

fraction of genes

■ top microRNA targets
(chimeric peaks with 7- or 8-mer perfect matches)
■ imperfect microRNA targets
(chimeric peaks without 7- or 8-mer perfect matches)
□ other genes

Fig S4. IPDrosha tumors are highly proliferative.

Fig S5. Epigenetic regulation of *Ccnd2.*

Fig S6. *DROSHA* expression is negatively correlated with let-7/miR-98-5p targets and E2F targets in human tumors.