Sperm derived H2AK119ub1 is required for embryonic development in *Xenopus laevis*

Valentin Francois--Campion[1,^], Florian Berger[1,^], Mami Oikawa[2], Maissa Goumeidane[1], Nolwenn Mouniée[1], Vanessa Chenouard[1], Kseniya Petrova[3], Jose G Abreu[3,4], Cynthia Fourgeux[1], Jeremie Poschmann[1], Leonid Peshkin[3], Romain Gibeaux[5], and Jérôme Jullien[1,*]

AFFILIATIONS

[1] Nantes Université, CHU Nantes, INSERM, Center for Research in Transplantation and Translational Immunology, UMR 1064, F-44000 Nantes, France

[2] Laboratory of Regenerative Medicine, Tokyo University of Pharmacy and Life Science, Hachioji, Tokyo, 192-0392, Japan

[3] Systems Biology, Harvard Medical School, Boston, MA, 02115, USA.

[4] Instituto de Ciências Biomédicas, Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, 21941-590, Brazil

[5] Univ Rennes, CNRS, IGDR (Institut de Génétique et Développement de Rennes) - UMR 6290, F-35000 Rennes, France

^equal contribution

*Corresponding author email: jerome.jullien@inserm.fr

ABSTRACT [140words]

Deposition of H2AK119ub1 by the polycomb repressive complexe-1 plays a key role in the initiation of facultative heterochromatin formation in somatic cells. Here we evaluate the contribution of sperm derived H2AK119ub1 to embryo development. In *Xenopus laevis* we found that H2AK119ub1 is present during spermiogenesis and into early embryonic development, highlighting its credential for a role in the transmission of epigenetic information from the sperm to the embryo. *In vitro* treatment of sperm with USP21, a H2AK119ub1 deubiquitylase, just prior to injection to egg, results in developmental defects associated with gene upregulation. Sperm H2AK119ub1 editing disrupts egg factor mediated paternal chromatin remodelling processes. It leads to post-replication accumulation of H2AK119ub1 on repeat element of the genome instead of CpG islands. This shift in post-replication H2AK119ub1 distribution triggered by sperm epigenome editing entails a loss of H2AK119ub1 from genes misregulated in embryos derived from USP21 treated sperm. We conclude that sperm derived H2AK119ub1 instructs egg factor mediated epigenetic remodelling of paternal chromatin and is required for embryonic development.

INTRODUCTION

Several studies point towards an epigenetic contribution of sperm for embryonic development. First, in several species paternal exposure to various environmental insults results in changes to the offspring [1–5]. In addition to such environmentally induced epigenetic changes, sperm is also proposed to naturally harbour epigenetic cues that are required for development. Histones have been considered as possible vectors for such epigenetic information. Indeed, in several species including human, modified histones such as H3K4me3 and H3K27me3 are found in the sperm chromatin, in particular around developmental genes and in a pattern reminiscent of that found in embryonic stem cells[6–9]. In a previous work we provided evidence that in the frog, *Xenopus laevis,* some of these modified histones are homogenously distributed in a population of sperm cells, hence indicating that they have the required attributes for a role in the regulation of embryonic development[10]. Lastly, interference with histone variant or post-translational modifications during the formation of the gametes, or after fertilisation, lead to embryonic gene misregulation[9,11,12]. Although these later experiments support a functional role for sperm-derived modified histone in embryonic development, indirect effect of the interference on the process of spermiogenesis or early embryos development cannot be ruled out. A direct assay to test for the requirement of sperm modified histones in embryo development is still lacking. This is due in part to the difficulty to access the chromatin in the highly compacted sperm nucleus. In this work, we aim to fill this gap and devised a strategy to alter the sperm epigenome just prior to its introduction to the egg. We specifically considered H2AK119ub1 as a possible carrier of necessary epigenetic information from the sperm to the embryos. Indeed, recent works points towards a key role of polycomb repressive complexe-1 (PRC1), upstream of PRC2, in the establishment of facultative heterochromatin, both in cultured cells and in embryos[13]. The H2AK119 ubiquitylation activity of the PRC1 complex appears to be integral part of this mechanism[13,14]. In *Xenopus laevis* we found that H2AK119ub1 dynamic during spermiogenesis and early embryonic development is compatible with a role in the epigenetic programming of sperm for embryonic development. We applied an *in vitro* epigenetic editing assay that uses USP21 deubiquitylase activity [15–18] to erase H2AK119ub1 from sperm chromatin. Embryos derived from such epigenetically edited sperm are developmentally impaired. Transcriptome and epigenome analysis indicates that H2AK119ub1, in combination with H3K4me3, is enriched on a set of embryonic genes misregulated in embryos derived from USP21 treated sperm. Such USP21 sensitive genes preferentially retain H2AK119ub1 upon sperm chromatin exposure to maternal factors. Lastly, we provide evidence that egg factor mediated deposition of H2AK119ub1 on paternal chromatin is influenced by pre-existing sperm H2AK119ub1.We conclude that sperm derived H2AK119ub1 instructs egg factor mediated epigenetic remodelling of paternal chromatin and is required for embryonic development.

RESULTS

We first evaluated the overall levels of H2AK119ub1 in chromatin during spermiogenesis and early embryonic development. Indeed, the presence of such modified histone in sperm seems to differ between species, its presence documented in that of mouse[19] while it appears absent in that of zebrafish[20]. In frogs the last stage of spermatogenesis is associated with a remodelling of nucleosomes resulting in a partial loss of H2A/H2B while a full complement of H3/H4 is retained on sperm chromatin

(**Fig. 1A**)[10]. Western Blot (WB) analysis captures this loss of H2A and reveal an overall
decrease in H2AK119ub1 in sperm compared to spermatid (**Fig. 1B and Suppl Fig.S1-**
A, see H2A/H4 and H2AK119ub1/H4 ratio, respectively). Importantly however, the
H2AK119ub1 to H2A ratio is similar in sperm compared to spermatid, indicating that
in this context of spermiogenesis-associated H2A loss, a significant level of
H2AK119ub1 is maintained in sperm (H2AK119ub1/H2A, **Fig. 1B**). We next assessed
H2AK119ub1 occurrence after fertilisation: in early frog embryos around ZGA
(blastulae stage) and later during gastrulation and neurulation. Interestingly, contrary
to what is seen with H3K27me3, a PRC2 associated modification, H2AK119ub1 is
detected throughout early embryogenesis (**Fig. 1C**). Altogether, these observations
highlight H2AK119ub1 credentials for a potential role in the transmission of epigenetic
information from sperm to the embryos.

We then carried out H2AK119ub1 ChIP to map the distribution of this histone
post-translational modification (PTM) on the sperm genome. Biological triplicates
were well correlated and subsequently pooled for downstream analysis (**Suppl Fig. 2A-**
D). We first assessed the distribution of H2AK119ub1 around the transcription start of
all sperm genes. H2AK119ub1 accumulates around the TSS of a fraction of sperm
genes (**Fig. 2A**), similar to what is observed in somatic cells[21] and embryos[14,20].
Noticeably, we observed in the top 10% TSS harbouring H2AK119ub1 an enrichment
for genes associated with developmental function (**Fig. 2A, Suppl Table 1**), and
reminiscent of the association reported with H3K27me3 in many species[6–9]. We next
sought to more generally identify regions associated with H2AK119ub1 genome wide.
H2AK119ub1 is a pervasive histone modification typically present on ~10% of nuclear
H2A in somatic cell[22,23] and up to ~40% in germ cells[19]. In *Xenopus laevis* sperm, it
accumulates over region covering larger part of the genome than other modified
histones with broad distribution such as H3K27me3 (**Suppl Fig. 2E**). To identify region
of H2AK119ub1 genomic enrichment we therefore used RECOGNICER, a coarse-
graining approach designed to capture enrichment over broad scale[24] (**Suppl Fig. 3,**
Table S1). Although we observed peaks of accumulation around genes, such as
illustrated for *wnt1.L*, *pax6.L* or the *hoxa.S* cluster (**Fig. 2B**), a large proportion of
H2AK119ub1 also accumulates on the sperm genome in distal region away from genes
(**Fig. 2C**). Noticeably we observed that H2AK119ub1 is significantly associated with
sperm gene regulatory elements (promoters and enhancers, **Fig. 2D**). Epigenetic control
of gene expression often entails a combination of epigenetic cues such as those
observed in X inactivation[25] or in transcriptional reprogramming[26]. We therefore
investigated the relationship between genomic regions enriched in sperm for
H2AK119ub1 and for other histone post-translational modifications. We focused on
H3K27me3[10], associated with the activity of Polycomb Repressive Complex 2, and
H3K4me3[10], a histone modification associated with transcription (Peaks, *e.g.* around
pax6.L and *hoxa.S* loci, **Fig. 2B**). We then determined the genome-wide association of
H2AK119ub1 with methylated K4 and/or K27 on histone H3. While most
H2AK119ub1 peaks exist independently of these methylated histones, ~24% of
H2AK119ub1 peaks overlap with either H3K4me3 and/or H3K27me3 peaks (**Fig. 2E**).
We observed a very strong promoter bias for peaks of H2AK119ub1 co-occurring with
H3K4me3, an expected trend given that H3K4me3 is known to strongly associate with
gene promoter (**Fig. 2F**). By contrast, when occurring in the absence of H3K4me3,
H2AK119ub1 is located away from promoter and in intergenic regions. Analysis of
fragment length obtained following sequencing of ChIP and Input fractions allow the
discrimination of DNA fragment associated with nucleosomes (full complement of

histones protecting 150bp of DNA) from those associated with subnucleosomes
(nucleosomes lacking one or two H2A-H2B dimer and protecting 110 or 70bp of
DNA) in sperm chromatin (**Suppl Fig. 4**). Noticeably, in sperm chromatin on the whole,
H2AK119ub1 is mostly associated to nucleosomes while H3K27me3 and H3K4me3
tend to be associated with subnucleosomes. So, while we observe co-enrichment of
H2AK119ub1 with methylated histone H3 at a subset of genomic loci, those histone
modifications are possibly present at those sites on adjacent
nucleosomal/subnucleosomal nucleoprotein complexes rather than on the same particle.
Altogether, this survey of H2AK119ub1 distribution in sperm reveals its enrichment on
gene regulatory region further strengthening its potential for gene regulation across
generation.

To further evaluate how sperm H2AK119ub1 could have an impact on
embryonic development, we next sought to follow its fate after fertilisation. Following
delivery to the egg, protamines are removed from sperm chromatin and a canonical
nucleosomal structure is re-instated. In conjunction with these events the sperm
chromatin is replicated. Therefore, the extensive changes to sperm chromatin that
follow fertilization have the potential to profoundly alter the sperm derived modified
histone landscape. To determine the fate of sperm modified histones we turned to
interphase egg extracts that can mimic the sperm chromatin remodelling events that
follow fertilization, including protamine to histone transition and DNA replication[10].
We carried out H2AK119ub1 and H3K4me3 ChIP-seq on untreated sperm, or sperm
having gone through egg extract mediated replication (**Fig. 3A**). Combining this dataset
with existing spermatid and gastrulae ChIP-seq[9,10,27] we obtain a global view of
H2AK119ub1, H3K4me3, and H3K27me3 dynamic during spermiogenesis and early
embryogenesis (**Fig. 3B**). Remarkably, while H3K4me3 and H3K27me3 are mostly
stable, H2AK119ub1 exhibits drastic redistribution throughout this developmental
period. Indeed, a large set of genomic location acquires H2AK119ub1 following sperm
chromatin replication (cluster 4, **Fig. 3B, Suppl Table 2**), and includes a large
proportion of gene TSSs (**Suppl Fig. 5A**). To better delineate the principle of this
developmental dynamic, we then compared H2AK119ub1 and H3K4me3 peaks in
untreated (sperm) *versus* egg extract treated sperm (replicated sperm). We confirm a
marked disparity in replication dynamics between sperm H3K4me3, mostly stable
(77% of sperm peaks conserved after replication (19345/24982) and 52% stable peaks
overall (19345/36770)), and sperm H2AK119ub1, mostly changing (23.5% of sperm
peaks maintained after replication (15796/67233) and 15% stable peaks overall
(15796/108740)) (**Fig. 3C, Suppl Fig. 5B**). Most of the replication redistribution of
H2AK119ub1 happens in intergenic region, while H2AK119ub1 peaks stable through
this process are mostly located on promoter (**Suppl Fig. 5D**). In this dynamic landscape,
the subset of H2AK119ub1 genomic location associated with H3K4me3 in sperm
appears selectively conserved (93% of sperm bivalent peaks are maintained after
replication (5222/5597) (**Fig. 3C**). Broader H2AK119ub1 peaks with higher level of
H2A retention discriminate location where H2AK119ub1/ H3K4me3 is maintained
following exposure to egg factors from those where it is lost (**Suppl Fig. 5C**). Lastly,
we confirmed that the H2AK119ub1 configuration observed following *in vitro*
treatment of sperm with egg extract is also detected in blastulae stage embryo at the
time of ZGA (**Suppl Fig. 6A-C**). We conclude that sperm exposure to egg factors
profoundly remodels paternal H2AK119ub1 genomic distribution.

We next evaluated to which extent such egg factor-mediated H2AK119ub1
remodelling is dependent on incoming sperm chromatin component. To directly

evaluate sperm H2AK119ub1 functional relevance to this process, we set out to erase
this histone modification, after completion of spermatogenesis, and prior to exposure
to egg factors. To that end, we incubated mature sperm in an oocyte extract containing
USP21, a H2A deubiquitylase[15,28] (**Fig. 4A**). In such extract, the sperm nucleus is
partially decondensed, allowing efficient deubiquitylation of H2AK119ub1 while the
levels of trimethylated H3K4, H3K9 and H3K27 remain unchanged (**Fig. 4B, Suppl**
Fig. 7A, Suppl Fig. 1B). Importantly, MNAse digestion of oocyte extract USP21-
treated sperm yields nucleosomal and subnucleosomal size DNA fragments typical to
that of naïve *Xenopus* sperm[9] (**Fig. 4C**, **Suppl Fig. 7B, and 7C left**). This indicates that
the overall sperm chromatin structure is maintained after oocyte extract treatment and
does not convert to a somatic type of chromatin organisation as triggered by egg extract
treatment[29] (**Fig. 4C, Suppl Fig. 7**). We next asked to what extend USP21 mediated
erasure of H2AK119ub1 from sperm chromatin influences its egg factor mediated
redistribution. To that end, we carried out modified histone analysis in sperm treated
with control or USP21 extract oocyte with or without a subsequent incubation in egg
extract (**Suppl Fig. 8A**). Interestingly, WB analysis indicates that H2AK119ub1, erased
from sperm chromatin by USP21 treatment, is at least partially reinstated by subsequent
exposure to egg extract (**Fig. 4D**), indicating the existence of a maternal H2AK119
ubiquitylation activity independent of pre-existing sperm H2AK119ub1. Using ChIP-
seq, we next ask if sperm H2AK119ub1 erasure prior to exposure to egg components
affects the maternal factor mediated remodelling of H2AK119ub1 on paternal
chromatin. We first focused on sperm genomic sites where we previously detected
accumulation of H3K4me3, H3K27me3, or H2AK119ub1 (as in **Fig. 3B**). This analysis
confirmed that USP21 efficiently erases H2AK119ub1 from its genomic site in sperm
(**Suppl Fig. 8B, Suppl Table 3**). We observed an overall reduction in egg factor
mediated H2AK119ub1 deposition on chromatin derived from USP21+ sperm
compared to that from USP21-sperm (**Suppl Fig. 8B**). We then compared
H2AK119ub1 peaks in samples originating from replicated USP21-treated sperm to
that from replicated control-treated sperm (**Fig. 4E**). Surprisingly, we observed that
sperm USP21 treatment leads not only to loss of H2AK119ub1 peaks post-replication
(lost peaks) but also triggers the appearance of a large set of peaks that is not deposited
by maternal factors in control sperm samples (gained peaks) (**Fig. 4E**). Comparatively,
USP21 treatment impact on post-replication H3K4me3 is much milder (**Suppl Fig.**
8C). To better characterize this shift, we first evaluated if H2AK119 ubiquitylation sites
detected following egg extract exposure of sperm harbour genomic features associated
with variant PRC1 (vPRC1) targeting in somatic cell (reviewed in[30]). Those
H2AK119ub1 sites that are specific to egg extract treated control sperm (lost peaks,
Fig. 4E) are associated with cluster 4 in Fig.3B and are indeed enriched for TF motifs
associated with vPRC1 targeting (*i.e.* USF1/2 E2F, E-box, **Suppl Fig. 8D, Suppl Table**
2). We also detected a strong association of these sites with unmethylated CpG island
(**Fig. 4F-G**), a genomic feature also shown to attract vPRC1 activity. By contrast, sperm
treatment with USP21 prior to egg factor exposure redirects H2AK119ub1
accumulation towards repeat elements and away from such targets (**Fig. 4H-I, Suppl**
Fig. 9, Suppl Table 5). We conclude that maternal factors mediated redistribution of
sperm H2AK119ub1 is instructed by pre-existing H2AK119ub1 derived from paternal
chromatin. In the absence of sperm H2AK119ub1, maternal factor mediated H2A
ubiquitylation activity is shifted from CpG islands to repeat elements.

To then determine the functional relevance of sperm derived H2AK119ub1 to
embryo development, we took advantage of the availability of paternally derived

embryos in frog. As described previously, such embryos develop solely on chromatin
derived from sperm and provide a direct measure of the paternal contribution to early
development[9]. We generated paternally derived embryos from sperm pre-incubated in
control or USP21 containing oocyte extracts and monitored their development (**Fig.**
5A)[31]. From gastrulation onwards, we observed a decrease in the developmental
potential of USP21 treated- compared to control treated- sperm derived embryos (**Fig.**
5B). We conclude that besides its DNA content, sperm also contributes to the embryo
an USP21 sensitive signal that is necessary for development. The observed defects
occur around the time of zygotic gene activation suggesting that USP21 treatment
associated developmental failure might relate to defective regulation of embryonic gene
expression. To evaluate this possibility, we carried out transcriptome analysis of early
gastrulae generated from pool of untreated, control extract-treated or USP21 extract-
treated haploid embryos. We performed RNA-seq analysis using sperm, oocyte extract,
and egg originating from different preparations for each of the biological replicates.
Differential gene expression analysis indicates that the treatment of sperm with a
control oocyte extract has minor effect on the resulting embryo transcriptome when
compared to embryos generated from untreated sperm (**Fig. 5C**, left; **Suppl Fig. 10A-**
B). By contrast, USP21 extract treatment leads to embryonic misregulation of 269
genes (**Fig. 5C**, right), the majority of which are upregulated in embryos derived from
USP21 extract treated sperm compared to embryos derived from control extract treated
sperm (74%, 200/269, **Fig. 5D, Suppl Fig. 10C, Suppl Table 4**). Thereafter, this set of
genes found differentially expressed in embryos derived from sperm treated with
USP21 oocyte will be referred to as USP21 sensitive genes. GO enrichment analysis
highlight genes related to cell cycle processes in this gene set (**Suppl Table 4**). We next
asked to which extend the expression pattern of USP21 sensitive genes identified on
pooled embryos reflect the behaviour in single embryos. To that end, we carried out
transcriptome analysis in single paternally derived gastrulae. We find that the
expression pattern of USP21 sensitive gene observed in pooled embryos is replicated
in half of the single embryos derived from USP21-treated sperm (4/8 USP21 treated
sperm derived embryos, **Suppl Fig. 10D**). This heterogeneous transcriptome pattern
matches the observed developmental outcome, with a fraction of gastrulae derived from
USP21-treated sperm developing to the tadpole stage (**Fig. 5B**) and is reminiscent of
the transcriptome heterogeneity detected in cloned embryos[32]. We then interrogated the
developmental dynamic of USP21 sensitive genes. We observed a clear distinction
between up- and down regulated USP21 sensitive genes (**Suppl Fig. 10E**). During
development of embryos generated by *in vitro* fertilisation[33], upregulated genes exhibit
a strong maternal contribution followed by clearing prior to gastrulation (stage 10-12)
and re-expression at tadpole stage (stage 25-40) while downregulated genes show
increased expression around gastrulation. A recent study indicates that at ZGA,
transcription occurs on genes with (maternal-zygotic, "MZ genes") or without (zygotic,
"Z genes") a maternal contribution of transcripts[34,35]. Interestingly, we observed that
USP21 sensitive genes are enriched for MZ genes, indicating that these genes exhibit
zygotic transcription in the context of a maternally derived transcripts (**Suppl Fig. 10E-**
F). Taken together, these analyses support the hypothesis whereby USP21 mediated
erasure of the repressive histone modifications H2AK119ub1 from sperm leads to
precocious embryonic expression of a subset of genes that could in turn impact critical
embryonic process such as cell division.

To evaluate this possibility further, we investigated to which extent
H2AK119ub1 genomic distribution associate with embryonic gene sensitivity to sperm

USP21 treatment. We observed an enrichment of sperm H2AK119ub1 peaks on USP21
sensitive gene body (**Suppl Fig. 10G**). Additionally, using a genome partitioning
approach (hidden Markov model) we also observe that USP21 sensitive genes are
associated with genomic regions exhibiting H2AK119ub1 either in sperm or in sperm
following exposure to egg extract (**Suppl Fig. 10H**). However, many more genes
harbour this modification in sperm, without being affected by USP21 treatment. We
therefore next interrogated the association of USP21 sensitive genes with different
types of H2AK119ub1 peaks stratified according to association with other histone
PTMs, namely H3K4me3 and H3K27me3 (as in **Fig. 2E-F**). We detect an association
of USP21 sensitive gene features with either H2AK119ub1 only peaks
(H2AK119ub1+; H3K4me3-; H3K27me3-) or trivalent H2AK119ub1 peaks
(H2AK119ub1+; H3K4me3+; H3K27me3+) (**Fig. 6A**). Strikingly however, USP21
sensitive genes show a strong enrichment for bivalent H2AK119ub1/H3K4me3 and
depletion for bivalent H2AK119ub1/H3K27me3 chromatin configuration. This
analysis suggests that the effect on embryonic gene expression of sperm USP21
treatment arise from erasure of H2AK119ub1 in the context of a bivalent
H2AK119ub1/H3K4me3 chromatin status. We also observed that, post-replication,
H2AK119ub1 and H3K4me3 signals around USP21 sensitive gene TSSs are higher
than around the bulk of *X. laevis* genes TSSs (**Fig. 6B**). Importantly, the significant
enrichment of USP21 sensitive genes promoter and gene body for a bivalent
H2AK119ub1/H3K4me3 chromatin configuration reported in sperm holds true after
sperm replication (**Suppl Fig. 10I**). This association of USP21 sensitive genes with
H2AK119ub1 is also observed in embryos at ZGA, with such trend still present but less
pronounced later during embryogenesis (**Suppl Fig. 6D**). We then took advantage of a
published dataset describing the consequences of expressing the H2A deubiquitylase
BAP1 in early mouse embryos to investigate if such a relationship is conserved across
specie[19]. Using a hidden Markov model, we indeed found that gene identified as
BAP1 sensitive in mouse two cell embryos are enriched in genome domains
exhibiting H2AK119ub1 marking in sperm (**Suppl Fig. 10J-K**). Altogether these
observations support a role for sperm derived H2AK119ub1 in the regulation of the
expression of a subset of embryonic genes.

We next considered the possibility that while a local accumulation of
H2AK119ub1 could be maintained post-replication, sperm modified histones could
have undergone replicative dilution. In other word, during replication the local
accumulation of histone along the genome could be maintained albeit at a reduced level,
for example through redistribution of sperm derived modified histones between the
replicated strands of DNA. Mechanistically, this is important as in frog ~8 rounds of
replication separate fertilisation form the main wave of ZGA and could therefore leads
to complete loss of sperm derived modified histones. Therefore, to compare
H2AK119ub1 level pre- and post- replication we calibrated the signal *via* semi-
synthetic H2AK119ub1 nucleosome spiked-in used in the ChIP-seq (see M&M)
(**Fig.6C**). We observed a global decrease of H2AK119ub1 signal on TSSs upon egg
extract treatment, suggesting that sperm chromatin replication induces redistribution of
this modified histones at these locations (black line, **Fig. 6C**). When considering USP21
sensitive gene however, we observed a different trend. While an overall decrease of
H2AK119ub1 signal is also detected upon replication, H2AK119ub1 loss is clearly less
pronounced on the TSSs of this subset of gene than seen genome wide (red line, **Fig.**
6C), a feature shared with the broader set of gene that have maternal-zygotic
contribution (green line, **Suppl Fig. 10L**). These analyses indicate that USP21 sensitive

genes are indeed associated with a sperm H2AK119ub1 configuration that persists
through egg factor mediated remodelling of the paternal epigenome. Lastly, we asked
how sperm USP21 treatment induced disruption of paternal epigenome remodelling by
egg factors affects USP21 sensitive genes. We found that the shift of H2AK119ub1
away from CpG towards repeat elements induced by USP21 treatment (as in **Fig. 4E**)
significantly deplete post replication association of USP21 sensitive genes from this
histone modification (**Fig. 6D**). Analysis of gene TSSs also showed that the preferential
post replication retention of H2AK119ub1 on USP21 sensitive genes is lost in response
to H2AK119ub1 sperm editing (**Fig. 6E**).

Taken together our results reveal that the setup of paternally derived zygotic
chromatin is dictated by component of the sperm epigenome and that this process is
required for proper expression of a subset of embryonic genes.

DISCUSSION

In this work, we found that in *Xenopus laevis*, paternal H2AK119ub1 genomic
distribution is extensively remodelled upon exposure to egg factors, reminiscent of the
situation found in the mouse[19]. In this dynamic epigenetic landscape, a subset of
genomic location is nevertheless found to maintain H2AK119ub1 following sperm
replication in extract as well as following early embryonic division. This suggests that
the mechanism recently described in cultured cells and demonstrating recycling of
H2AK119ub1 between daughter cells[67] could apply to the transmission of H2A
associated epigenetic cue from the gametes to the dividing cell of the frog embryos.
Besides those stable sites of H2AK119ub1 accumulation, sperm incubation in egg
extract also triggers H2AK119ub1 deposition on new genomic sites. Whether this
remodelling requires replication is not known since sperm treatment with egg
extract triggers many processes besides replication. Once established those new
sites appear stable through replication as they could be detected in late cleavage stage
embryos. However, our study is limited by the lack of knowledge about H2AK119ub1
genomic distribution during xenopus laevis pre-ZGA developmental period.
Particularly, we are lacking in vivo data describing the H2AK119ub1 landscape
immediately after fertilisation and during the first round of embryonic divisions.
Therefore, mechanistically we don't know if the remodelled H2AK119ub1 distribution
found after in vitro sperm replication, and in vivo after 8 round of cell division, is
faithfully transmitted throughout cleavage stage or transiently adopt a different
configuration during that period.

Egg factor dependent accumulation of H2AK119ub1 occurs around
unmethylated CpG island and/or TF binding sites (*i.e.* USF1/2), genomic features that
are known target of PRC1 activity in other systems[30]. Besides transcription factors,
Kdm2b a component of vPRC1 is involved in more broadly targeting the H2AK119
ubiquitylation activity of this complex to unmethylated CpG island in ESC[30].
Kdm2b is a likely candidate for a role in the observed H2AK119ub1 remodelling
following exposure to egg components as it is maternally provided as protein in
frog
(https://www.xenbase.org/xenbase/gene/geneExpressionChart.do?method=dr
[awProtein](https://www.xenbase.org/xenbase/gene/geneExpressionChart.do?method=drawProtein)). In mouse embryonic stem cell, the establishment of polycomb repressive
domain occurs in a stepwise manner with a variant PRC1 complex initiating their

formation through ubiquitylation of H2AK119, followed by recruitment of the PRC2
complex and subsequent methylation of H3K27me3. Such sequential events have also
been described during Neural Progenitor Cell differentiation[36] and X chromosome
inactivation[37]. Recently, this sequence of events have been investigated in the mouse
and zebrafish embryos[19,20,38]. In these embryos H2AK119ub1 is also required for the
deposition of H3K27me3 and transcriptional repression around the time of zygotic gene
activation. Our observation that H2AK119ub1 is present in pre-ZGA embryo in the
absence of H3K27me3 argues for the existence of a similar mechanism in frog[39,40].

Interference with H2AK119ub1 in the mouse, either by knocking down PRC1
component[38] in oocyte, or erasing H2AK119ub1 from embryos[19,20] leads to mis-
regulation of genes at ZGA and is associated with embryonic lethality. While clearly
demonstrating a role for H2AK119ub1 in regulation of transcription around ZGA, these
approaches entail prolonged epigenetic interference during either gametogenesis or
early embryogenesis and, as a consequence, do not enable to determine the extent to
which embryonic H2AK119ub1 is instructed/inherited from sperm and/or oocyte
H2AK119ub1[9,11,12,19,20,38]. Here, by editing the sperm epigenome, after completion of
spermiogenesis and prior to exposure to egg factors, we circumvent these limitations.
We demonstrate that sperm contribute to the embryo a USP21 sensitive cue required
for paternal H2AK119ub1 remodelling by egg factor, embryonic gene expression, and
developmental competence (**Fig. 7**). We find that egg factor mediated accumulation on
paternal CpG island depends on H2AK119ub1 delivered by the sperm. Strikingly, we
observe that erasure of H2AK119ub1 from sperm leads to a global shift in genomic
accumulation from CpG islands to repeat elements. Given the propensity of vPRC1 to
be targeted to and to spread from existing H2AK119ub1 location[41] it raises the
possibility that pre-existing sperm derived "docking sites" could limit chromatin
sampling by the polycomb complex and prevent H2AK119ub1 deposition at genomic
location where targeting is usually limited such as the repeat elements. Embryos
derived from H2AK119ub1 edited sperm exhibit transcriptional misregulation, mostly
upregulation, compatible with the repressive function of this histone PTM on
transcription. A similar observation is observed in mouse when H2AK119ub1
deubiquitylase is expressed at fertilization[19], or in zebrafish when Rnf2 is chemically
inhibited[20]. Given the extent of egg factor mediated H2AK119ub1 accumulation on
gene promoters observed in frog, mouse or zebrafish, it is somewhat surprising that
only a limited number of genes are mis-regulated when its distribution is massively
altered. We identify an association of misregulated genes with H2AK119ub1 in frog
sperm and find that this association is preferentially maintained on these genes upon
sperm chromatin replication, further supporting a direct role for this histone post-
translational modification in the epigenetic programming of sperm for regulation of
embryonic transcription. We cannot rule out that a part of the genes is indirectly
affected, for example as downstream target of a "primary" H2AK119ub1 target.
Alternatively, we provide evidence that gene sensitivity to H2AK119ub1 erasure from
sperm occurs when genes are associated with H3K4me3, a histone PTM known to act
as a vector of intergenerational information[9–11,42,43]. It is likely that additional epigenetic
features besides H3K4me3 contribute to gene sensitivity to H2AK119ub1 erasure from
sperm. Additionally, we observe that most affected genes are normally expressed late
in embryogenesis (tadpole stage) a feature also seen in mouse with a large proportion
of gene sensitive to H2AK119ub1 depletion expressed post-implantation[8]. It is
therefore possible that the effect of H2AK119ub1 expression is only partially revealed

at early embryonic stage since transcription factors targeting these genes might not be
present.

Our work advance knowledge form previous studies by showing that part of
H2AK119ub1 found in embryos is paternally instructed. This necessary function of
sperm derived modified histones has important bearing on our understanding of
inter/trans generational transmission of epigenetic information. One could
indeedenvisage that such a naturally occurring mechanism of epigenetic information
transmission to the embryos could be hijacked to transmit environmentally induced
sperm epigenetic cues. Based on the very dynamic nature of histone PTMs around
fertilisation, it is likely that transmission to the embryos will be context- (combination
of marks) and location- dependent. Future studies will aim at evaluating to which extent
sperm histones PTMs differ between individuals and to which extent they are
maintained in embryos.

FIGURE LEGENDS

Figure 1: *H2AK119ub1 dynamics during spermiogenesis and early embryonic*
development

(**A**) Schematic representation of *Xenopus laevis* nucleosome remodelling during
spermiogenesis (**B**) Left, Western blot analysis of H2AK119ub1, H2A and H4 histones.
Right quantitation of signal ratio of H2A to H4, H2AK119ub1 to H4, and H2AK119ub1
to H2A, in spermatid and sperm. * $p<0.05$, ratio paired t test (n=4). Median is shown.
(**C**) Western blot analysis of H2AK119ub1, H3K27me3, H3K9me3, H3K4me3, H3 and
H4 in *Xenopus laevis* embryo pre- (blastulae) and post- (gastrulae, and neurula) ZGA.

Figure 2: *H2AK119ub1 associates with gene regulatory regions in sperm chromatin*

(**A**) Heatmap of H2AK119ub1 signal (log(ChIP/Input)) on all *Xenopus laevis* genes
TSS +/-5kB. GO enrichment is shown for the top 10% genes. (**B**) Genome browser
view of log(ChIP/Input) of sperm H2AK119ub1, H3K4me3, and H3K27me3 around
wt1.L, pax6.L, snx10.s and *hoxa.S* loci. Boxed area underneath the gene track indicates
regions of enrichment (peaks). (**C**) Distribution of sperm H2AK119ub1 peaks on
genomic features. (**D**) Fold enrichment of H2AK119Ub1 peaks on different genomic
features, *** $p<0.001$, permutation test. (**E**) Pie chart indicating the proportion of
H2AK119ub1 peaks either alone, or overlapping H3K4me3 and/or H3K27me3 peaks
in sperm. (**F**) Genomic distribution of sperm H2AK119ub1 peaks according to overlap
with H3K4me3 and/or H3K27me3 peaks.

Figure 3: *Sperm H2AK119ub1 is remodelled following exposure to egg factors*

(**A**) Experimental design for monitoring modified histones fate following sperm
exposure to egg factors. (**B**) Heatmap of H2AK119ub1, H3K4me3, or H3K27me3
signal (log(ChIP/Input)) in spermatid, sperm, replicated sperm, and gastrula. All

genomic locations enriched for one of these histone modifications (peaks) in at least
one of these developmental stages are represented. (**C**) Venn diagram indicating the
changes to H2AK119ub1 (left), H3K4me3 (middle) or overlapping
H3K4me3/H2AK119ub1(right) peaks during sperm chromatin replication in egg
extract. Percentages of sperm peaks that are maintained after replication are indicated
underneath.

Figure 4: *H2AK119ub1 remodelling by egg factors is altered by sperm pre-treatment*
with a H2A deubiquitylase

(**A**) Schematic representation of sperm treatment with oocyte extract containing USP21.
(**B**) Left, Western blot of H2AK119ub1, H3K4me3, H3K9me3, H3K27me3 and
histone H4 on untreated sperm, USP21 treated sperm and oocyte extract treated sperm.
Right, signal ratio of H2AK119ub1 to H4 in untreated sperm, USP21 treated sperm, or
oocyte extract treated sperm. *** $p<0.005$, **** $p<0.001$, ratio paired t test (n=6).
Median is shown. (**C**) DNA fragments generated by MNase digestion of *Xenopus*
laevis untreated sperm, oocyte control extract treated sperm, oocyte USP21 extract
treated sperm, egg extract treated sperm and somatic cell. (**D**) Western blot analysis of
H2AK119ub1 and histone H4 on sperm after treatment with control- or USP21- oocyte
extract without (left) or with (right) subsequent replication in egg extract. (**E**) Heatmap
of H2AK119ub1 and H3K4me3 signal (log(ChIP/Input)) in sperm after treatment with
control- or USP21- oocyte extract without or with subsequent replication in egg extract.
All genomic locations enriched for H2AK119ub1 (peaks) in replicated samples are
represented. (**F-I**) DNAme and H2AK119ub1 signal (log(ChIP/Input)) on *Xenopus*
laevis genome CpG island (F) and repeat element (H). Z-score of H2AK119ub1
enrichment on CpG island (G) and repeat element (I) in sperm, replicated sperm and
replicated USP21 sperm. *** $p<0.005$, permutation test. Numbers of H2AK119ub1
peak overlapping the genomic feature are indicated for each cell type.

Figure 5: *Sperm pre-treatment with a H2A deubiquitylase decreases developmental*
potential and leads to embryonic gene misregulation

(**A**) Experimental design for the generation of haploid embryos from sperm or USP21
treated sperm. *Xenopus* illustrations © Natalya Zahn (2022) from Xenbase
(www.xenbase.org RRID:SCR_003280). (**B**) Developmental progression to blastula,
gastrula or swimming tadpole stage of haploid embryos derived from a USP21 or
control oocyte extract treated sperm. Chi-squared test, * p-value<0.05. Median and SD
values are shown. (**C**) MA-Plot showing as y-axis the log foldchange (Log2FC) of gene
expression between control oocyte extract treated sperm *versus* untreated sperm (left)
and between USP21 oocyte extract treated sperm derived embryo and control oocyte
extract treated sperm (right). x-axis is the averaged gene expression across samples.

Figure 6: *Genes sensitive to sperm deubiquitylase treatment are H2AK119ub1 target*

(**A**) Heatmap showing expression of differentially expressed gene in gastrulae derived
from USP21 oocyte extract treated sperm *versus* that derived from control oocyte
extract treated sperm. (**B**) Z-score of different H2AK119ub1 peaks categories

enrichment on TSS, enhancer, and gene body of USP21 sensitive genes. * p-
value<0.05; ** p-value < 0.01, permutation test. Number of H2AK119ub1 peak
overlapping the USP21 sensitive gene feature are indicated for each category. **(C)**
H2AK119ub1 and H3K4me3 signal on TSS +/- 5kB for all genes (black), USP21
sensitive genes (red) before and after sperm replication. *** p-value<0.001, Wilcoxon-
Mann-Whitney test on TSS+/1kb interval (shaded orange area). **(D)** Calibrated
H2AK119ub1 ChIP-seq signal on TSS +/- 5kB for all genes (black), USP21 sensitive
genes (red) before and after sperm replication. *** p-value<0.001, Wilcoxon-Mann-
Whitney test. **(E)** Z-score of USP21 sensitive gene enrichment on H2AK119ub1 peaks
gained (USP21 specific), lost (Control specific) or maintained (common) in replicated
USP21 treated sperm *versus* replicated control sperm. ***p-value<0.001, permutation
test. Number of USP21 sensitive gene overlapping each category is shown. **(F)**
Calibrated H2AK119ub1 ChIP-seq signal on TSS +/- 5kB for all genes (black), USP21
sensitive genes (red) before and after replication using a control oocyte extract treated
sperm or USP21 treated sperm. *** p-value<0.001, Wilcoxon-Mann-Whitney test.

Figure 7: *Model of H2AK119ub1 epigenetic programming of sperm for embryonic*
development in Xenopus laevis

Supplemental Figure 1: *Quantification of Western blot data.*

(A) Ratio of fluorescence intensity detected by WB for H2AK119ub1/H2A (left)
H2AK119ub1/H4 (right) and H2A/H4 (middle). The same data as Fig.1G are plotted
with the addition of a solid line linking paired sample quantified in the same WB
experiment. **(B)** H2AK119ub1 fluorescence intensity detected by WB in untreated,
control oocyte extract treated sperm, USP21 extract treated sperm. The same data as
Fig.2E are plotted with the addition of a solid line linking paired sample quantified in
the same WB experiment.

Supplemental Figure 2: *Pooling strategy for ChIP-seq replicates*

(A) Genome browser view of the log(ChIP/Input) for H2AK119ub1 around the *hoxa.L*
cluster on 3 sperm biological replicates. (**B**) Spearman correlation of sperm and
replicated sperm H2AK119ub1, H3K27me3 and H3K4me3 ChIP-seq replicates. (**C**)
(**D**) Heatmap illustrating the distribution of H2AK119ub1 and H3K4me3 in sperm and
replicated sperm replicates and in the corresponding pooled sample obtained by
subsampling. Log(ChIP/Input) signal is shown for TSS (C) and CpG (D), respectively.
(**E**) Fingerprint plot of Input, H3K4me3, H3K27me3, and H2AK119ub1 sperm ChIP
samples.

Supplemental Figure 3: *Peak calling using RECOGNICER and SICER/MACS2.*

(A) Overlap of peaks called using RECOGNICER, SICER, and MACS2. Numbers in
red indicate RECOGNICER specific peaks *versus* those overlapping with those of
SICER and/or MACS2. Diagram scaling is based on RECOGNICER peaks. Blue and
green numbers correspond to the same comparison using SICER or MACS2 as the
reference set of peaks, respectively. (**B**) Heatmap of H2AK119ub1, H3K27me3,

H3K4me3 signal (log(ChIP/Input)) for RECOGNICER, SICER, and MACS2 set of peaks. (**C**) Comparison of Z-scores obtained from permutation test using set of peaks based on peaks set on peak calling using RECOGNICER (top row, corresponding to data presented in Fig. 2K, Fig. 2M, Fig. 3H and Suppl Fig. 10G) or SICER (bottom row). (**D**) Genomic distribution of sperm H2AK119ub1 peaks according to overlap with H3K4me3 and/or H3K27me3 peaks when calling peak using RECOGNICER (top, corresponding to data presented in Fig. 1I) or SICER (bottom) (**E**) Z-score indicating enrichment of different H2AK119ub1 peaks categories on TSS +/-5kb, enhancer, and gene body of USP21 sensitive genes. Top row is using RECOGNICER (already shown in Fig. 3E) while bottom row is using SICER for H2AK119ub1 peak calling.

Supplemental Figure 4: *H2AK119ub1 is mostly associated with nucleosomes while H3K4me3/H3K27me3 are mostly associated with sub-nucleosomes in Xenopus laevis sperm.* (**A**) Schematic representation of *Xenopus laevis* chromatin remodelling from nucleosomes to a mixture of nucleosomes and sub-nucleosomes during spermiogenesis. (**B**) Fragment size distribution obtained after paired-end sequencing of chromatin immunoprecipitated with antibodies against H2AK119ub1 (pink) in sperm and spermatid, and against H3K4me3 (green) and H3K27me3 (yellow) in sperm. Matching Input fraction are shown in grey.

Supplemental Figure 5: *Sperm H2AK119ub1 fate following exposure to egg factors*

(**A**) Heatmap of H2AK119ub1 (pink), H3K4me3 (green), and H3K27me3 (yellow) log(ChIP/Input) signal ratio on genes TSS from spermatid, sperm prior and after egg extract treatment, and gastrula stage 12 embryo. (**B**) Scaled Venn diagram of H2AK119ub1, H3K4me3 or overlapping H2AK119ub1/H3K4me3 peaks in spermatid, sperm, and sperm exposed to egg factors. Percentage of peaks that are maintained at each developmental transition are indicated underneath. (**C**) Top: boxplot showing calibrated H2AK119ub1 on H2AK119ub1/H3K4me3 peaks lost (375), maintained (5222) or gained (10442) following exposure to egg factors (top). Bottom: peak size (blue dots) as well as H2A (red dots) in each peak category. (**D**) Genomic distribution of sperm peaks before or after replication, as well as the subset of peaks lost, maintained, or gained during replication**.**

Supplemental Figure 6: *H2AK119ub1 distribution from spermiogenesis to late embryogenesis.*

(**A**) Pearson correlation of H2AK119ub1 ChIP in spermatid, sperm, sperm replicated in egg extract, blastulae at ZGA, early gastrula, and late neurula. (**B**) (**C**) Heatmap of H2AK119ub1 log(ChIP/Input) signal for TSSs (B) and CpG islands (C) in spermatid, sperm, sperm replicated in egg extract, blastulae at ZGA, early gastrula, and late neurula. (**D**) Metaplot analysis of H2AK119ub1 log(ChIP/Input) signal for TSSs of all genes (black) or USP21 sensitive genes (red) in spermatid, sperm, sperm replicated in egg extract, blastulae at ZGA, early gastrula, and late neurula. *** p-value<0.001, Wilcoxon-Mann-Whitney test on TSS+/1kb interval (shaded orange area).

Supplemental Figure 7: *Effect of oocyte extract treatment on sperm chromatin*

(**A**) Quantitation of H3K4me3, H3K9me3 and H3K27me3 to H4 WB signal ratio in untreated sperm, USP21 treated sperm or control oocyte extract treated sperm, (**B**) Capillary electrophoresis analysis of DNA fragments obtained following MNAse digestion of somatic cell (yellow), egg extract treated sperm (dark blue), sperm (blue),

control oocyte extract treated sperm (cyan), and USP21 extract treated sperm (red). (**C**)
Fragment size distribution obtained after paired-end sequencing of DNA fragments
obtained following MNase digestion of sperm (top row), control oocyte extract treated
sperm (middle row) or USP21 treated sperm (bottom row) without (left column) or with
(right column) subsequent incubation in egg extract. (**D**)(**E**) Heatmap of H2AK119ub1
and H3K4me3 log(ChIP/Input) signal for TSSs (D) and CpG islands (E) in sperm and
sperm treated with control oocyte extract.

Supplemental Figure 8: *Fate of sperm H2AK119ub1 and H3K4me3 following*
exposure to egg factors.

(**A**) Experimental design for monitoring modified histones fate from control oocyte
extract or USP21 treated sperm. (**B**) Heatmap of H2AK119ub1 (pink), H3K4me3
(green) and H3K27me3 (yellow) log(ChIP/Input) signal in spermatid, control oocyte
extract- and USP21- treated sperm prior and after egg extract treatment, and gastrula
stage 12 embryo. Heatmap includes the same genomic location as in **Fig. 2B**. (**C**)
Heatmap of H2AK119ub1 and H3K4me3 signal log(ChIP/Input) in sperm after
treatment with control oocyte extract or USP21 oocyte extract without or with
subsequent replication in egg extract. All genomic locations enriched for H3K4me3
(peaks) in replicated samples are shown. (**D**) Transcription factor motifs enriched in
peaks that are lost, maintained, or gained upon replication of control versus USP21
treated sperm (clusters from Figure 4E).

Supplemental Figure 9: *H2AK119ub1 is enriched on repeat elements of the genome*
following exposure of USP21 treated sperm to egg factors.

(**A**) Metaplot analysis of H2AK119ub1 log(ChIP/Input) signal on all CpG islands
(10630 elements) and all repeat elements of the genome (313338 elements). * p-
value<0.005, Wilcoxon-Mann-Whitney test is indicated with an asterisk in black
(replicated control sperm *versus* replicated USP21 sperm), blue (replicated control
versus control sperm) or red (replicated USP21 sperm *versus* control sperm). (**B**) same
analysis as in (**A**) but splitting repeat elements according to subtypes.

Supplemental Figure 10: *Genes misexpressed in embryos upon H2AK119ub1*
removal by deubiquitylases are associated with H2AK119ub1 in sperm.

(**A**) Principal Component analysis of replicate RNA-seq samples used in this study
(USP21, USP21 extract treated sperm derived gastrulae; Control, control oocyte extract
treated sperm derived gastrulae; Untreated, untreated sperm derived gastrulae). (**B**)
Hierarchical clustering of replicate samples used in this analysis. (**C**) Heatmap
depicting expression of USP21 sensitive genes in replicates from USP21 extract treated
sperm derived gastrulae, control extract treated sperm derived gastrulae, as well as
gastrulae derived from untreated sperm. (**D**) Heatmap of gene expression in single
haploid gastrula derived from control- or USP21-treated sperm. Expression is shown
for the USP21 sensitive gene set identified on embryos pool (**Fig. 3D**). (**E**) USP21
sensitive gene set expression during early *Xenopus laevis* embryo development (data
from[33]). Expression is shown for oocyte, egg, blastulae (stage 8-9), gastrulae (stage 10-
12), neurula (stage 15-20), as well as early to late tailbud (stage 25-40) are indicated.
The right-hand side panel shows the expression of USP21 sensitive gene in gastrulae
derived from control- and USP21-treated sperm. (**F**) Violin plot of gene expression in

egg (log of normalised gene count) for all genes (black), USP21 sensitive genes (red),
maternal genes (cyan), maternal and zygotic genes (green) and zygotic only gene
(orange). USP21 sensitive genes are enriched for genes with maternal and zygotic
expression in *Xenopus laevis*. The table below indicates p-Value resulting from a
permutation test to evaluate enrichment of USP21 sensitive gene for either maternal-
zygotic (MZ) or zygotic only (Z) genes. MZ and Z dataset taken from[34,35]. **(G)** Z-Score
of H2AK119ub1 peak enrichment on TSS, enhancer, and gene body of USP21 sensitive
genes. *p-value<0.05; permutation test. **(H)** Left, Heatmap representing H2AK119ub1
dynamics of each ChromHMM type in spermatid, sperm, replicated sperm, and
gastrulae embryos. Color intensity represents H2AK119ub1 signal for a given state and
cell type. Right, Z-score indicating enrichment for USP21 sensitive genes in genomic
regions associated with the different chromHMM states. *p-value<0.05; **p-value <
0.01, ***p-value<0.001, permutation test. **(I)** Z-score of different H2AK119ub1 peaks
sets enrichment on TSS, enhancer, and gene body of USP21 sensitive genes. Z-score
are shown for sets corresponding to H2AK119ub1 peaks that are lost, maintained or
gained following incubation of sperm in egg extract as well as for the subset of
maintained H2AK119ub1 overlapping H3K4me3 peaks. *p-value<0.05; **p-value <
0.01, ***p-value<0.001, permutation test. **(J)** Experimental scheme used in Chen *et al.*,
Nature genetics, 2021[19]. **(K)** Left, Heatmap representing H2AK119ub1 dynamics of
each ChromHMM type in mouse embryos and gametes based on data generated[19].
Color intensity represents H2AK119ub1 percentage for a given state and cell type.
Right, Z-score indicating enrichment for BAP1 sensitive genes identified in[19] in
genomic regions associated with the different chromHMM states. *p-value<0.05; **p-
value < 0.01, ***p-value<0.001, permutation test. **(L)** Calibrated H2AK119ub1 ChIP
signal on TSS +/- 5kb for all genes (black), USP21 sensitive genes (red), maternal genes
(cyan), maternal zygotic genes (green) and zygotic genes (orange) before and after
sperm replication. *** p-value<0.001, Wilcoxon-Mann-Whitney test.

<u>**Supplemental Table 1**</u>*: Histone PTMs ChIP-seq*

<u>**Supplemental Table 2**</u>: *Gene Ontology and TF motif analysis related to Fig.2*

<u>**Supplemental Table 3**</u>: *Gene Ontology related to Supplemental Fig.8*

<u>**Supplemental Table 4**</u>: *Control- and epigenetically edited- sperm derived embryos*
RNA-seq data

<u>**Supplemental Table 5**</u>: *Repeat element analysis*

<u>**Supplemental Table 6**</u>: *Extended Permutation test data*

MATERIALS AND METHODS

Contact for Reagent and Resource Sharing. Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Dr J. Jullien (jerome.jullien@inserm.fr).

Experimental Model and Subject Details. Mature *Xenopus laevis* females were obtained from Tefor Paris-saclays (151 Rte de la Rotonde, 91400 Saclay, France, https://tefor.net/tefor-paris-saclay-nos-equipes/tefor-paris-saclay-aqua/). Our work with Xenopus complied with all relevant ethical regulations for animal testing and research. Work with frog was covered under the licence #202212131604754/APAFIS 39794V2 and frog husbandry and all experiments were performed according to the relevant regulatory standards. Animals were kept in a recirculating system (ALS, https://www.aquariumlocationservices.fr/professionnels-elevages-aquatique-aquariophilie/) at a density of one adult/3l, with ~10% water change per day. Water was sequentially filtered with mechanical pad sump filter, nitrifying bacteria filter, and UV sterilized. Water quality parameters were as follow: temperature 17-22°C; PH 6-8. Photoperiod was set to 12h ON/12h OFF. Frogs are fed twice per week with NEO MARIN coul 4 - X4 - S20 (Le Gouessant, https://www.legouessant.com/). Unconsumed food was removed 30 min after the start of feeding.

Oocyte extract. Fresh ovaries were manually dissociated in 15-25 oocytes clumps. 3-5 mL of dissociated ovary fragments were transferred into 12.5 mL 1X Modified Barth's Saline (MBS, 88 mM NaCl, 1 mM KCl, 10 mM Hepes, 2.4 mM NaHCO3, 1.7 mM MgSO4, 1.3 mM Ca(NO3)2, 0.8 mM CaCl2) supplemented with 250 µL of 5 mg/mL Liberase (Roche, 05 401 127 001) and incubated for 2h30-3h at room temperature (RT) with gentle rocking. Defolliculated oocytes were washed three times with 1X MBS and healthy stage 5-6 oocytes manually sorted. Oocytes were then injected in the vegetal pole with 25 ng of USP21-NLS-HA mRNA, or control GFP mRNA and incubated for 3 days at 18°C in 1X MBS, renewing medium and discarding dead cells every day. ~500-1000 oocytes were collected in an Eppendorf tube on ice, the excess 1X MBS discarded and replaced with 1X MBS supplemented with Protease Inhibitor Cocktail (Sigma, P2714-1BTL). The oocytes were then stacked up by centrifugation at 100 g for 1 min at RT and excess medium carefully discarded. Oocytes were then thoroughly vortexed and centrifuged at 21 000g for 10 min at 4°C. The clear intermediate phase between yolk (top) and pigment (bottom) was retrieved. The collected fraction was further cleared by an additional centrifugation at 21 000 g for 10 min at 4°C. The oocyte extract was then aliquoted and stored at -80°C.

Oocyte extract treatment. Permeabilized sperm stored in SSB at -80°C were thawed, washed with two volumes of ice cold 1X PBS and centrifuged at 800 g for 8 min at 4°C. 2M sperm in 1X PBS were centrifuged at 21000 g for 1 min at 4°C. The supernatant was discarded, and the pelleted sperm resuspended in 10 µL of oocyte extract on ice. The sperm incubated in the oocyte extract for 20-40 min at 23°C on a heating block. Treatment was stopped by addition of 90 µL of ice cold 1X PBS and centrifugation at 21000 g for 1 min at 4°C. The supernatant discarded and the remaining pellets resuspended in SSB for -80°C long term storage.

Egg extract treatment. Interphasic extracts were generated as previously described[45] .For the quantification of DNA replication in individual nuclei in egg extract, 33 µM rhodamine dUTP (Roche, 11534378910) was added at the indicated

times (20, 40, 60, 80 mins after incubation) for 5 min as indicated, and the reaction was
stopped with 1X PBS and fixed in 4% paraformaldehyde. The reaction was spun
through a 30 % sucrose cushion in PBS. The sperm cells were rinsed twice in 1X PBS
for 10 min and mounted in Vectashield mounting medium with DAPI (Vector
laboratories, H-1200). Counting of rhodamine positive cells were performed on an
epifluorescence microscope.

Protein extraction and Western Blot analysis. Sperm samples stored in SSB were
washed in 1X PBS and 1M sperm pellets were resuspended in 30 µL 2X Blue Eye (600
mM NaCl, 1 % Triton X100, 0.5 % Na-Deoxycholate, 4 % Sodium Dodecyl Sulfate,
20 % Glycerol, 0.0004 % Bromophenol blue, 175 mM Tris-HCl, 10 % Betamercapto
ethanol), boiled at 96°C for 5 min and centrifuged at 21000g for 5 min at 4°C. The
supernatant was transferred to a new tube and directly use for analysis or stored at -
20°C. 4-20 % mini-PROTEAN® TGX (Biorad, #4561094) were loaded with 10 µL of
sample and western blot analysis were performed according to standard protocols. For
blotting nitrocellulose membranes, a semi-dry transfer was used (Biorad, Trans-Blot®
SD Semi-Dry Transfer Cell) 30-40 min at 25V. The following primary antibodies were
used: anti-H2A (1:2500, Cell Signalling, 3636), anti-H2AK119ub1 (1:2500, Cell
Signalling, 8240), anti-H3K4me3 (1:2500, Abcam, ab8580), anti H3K9me3 (1:2500,
Abcam, ab8898), anti-H3K27me3(1:2500, Cell Signalling, 9735S), anti-H4 (1:2500,
Abcam, ab31830) and anti-HA (1:2500, Sigma, H9658). Membranes were incubated
with primary antibodies 2 h at RT or overnight (ON) at 4 °C. Secondary antibody are
anti-rabbit (1:25000, ThermoFisher, A21076), anti-mouse IgG Alexa Fluor 680
(1:25000, ThermoFisher, A21058), Goat anti-rabbit Alexa 800 (1:25,000,
ThermoFisher, A32735), or IRDye 800CW Goat Anti-Mouse IgG (H+L) (LI-COR,
926-32210) were used. The membranes were imaged either on a ChemiDoc MP
Imaging System (Biorad) or an Odyssey scanner (LI-COR).

Sperm Chromatin Immunoprecipitation (ChIP). Chromatin fractionation and
chromatin immunoprecipitation (ChIP) were performed as described before[10,44] with
slight modifications. 80 µL of Magnetic beads (M280 Sheep anti-rabbit IgG, Invitrogen,
#11204D) were used per reaction and all wash steps were carried out with a magnet.
Beads were washed with 1 mL of TE pH 8.0, lysis buffer (LB)(1:1 mixture of Buffer 1
and MNase buffer, see below), and incubated in 1 mL of LB with 100 µL of 10 mg/mL
BSA at 4°C for 30 min ("pre-blocking"). Pre-blocked beads were washed with 1 mL of
LB twice and resuspend in 80 µL of LB. Half of beads were incubated with antibody
overnight and leftovers were stored in 4°C for "pre-clear chromatin" step. For ChIP,
the following antibodies were used: anti-H3K4me3 (Abcam, ab8580), anti-
H2AK119ub1 (Cell Signaling, 8240). One million digitonin permeabilised sperm
nuclei or egg extract treated sperm nuclei were resuspended in 50 µL of Buffer 1 (0.3
M Sucrose, 15 mM Tris pH 7.5, 60 mM KCl, 15 mM NaCl, 5 mM MgCl2, 0.1 mM
EGTA, 0.5 mM DTT) containing 0.125 ng H2AK119ub biotinyltated nucleosome
(Epicypher #16-0020), followed by addition of 50 µL of Buffer 1 with detergent (Buffer
1 including 0.5 % NP40 and 1 % NaDOC). Samples were incubated for 10 min on ice.
100 µL of MNase buffer (0.3 M Sucrose, 85 mM Tris, 3 mM MgCl2, 2 mM CaCl2, 2.5
U of microccocal nuclease: Roche 10107921001) was added in to each tube (0.5 million
of cells per tube). Tubes were incubated at 37°C for 30 min in pre-warmed water bath.
Reaction was stopped by adding 2 µL of 0.5 M EDTA pH 8.0 in the same order as
started. Tubes were vortexed and placed on ice for at least 5 min. Supernatant and pellet
were separated by centrifugation at 13000 rpm for 10 min at room temperature. Beads

(stored at 4C for pre-clear chromatin) were washed with 1 mL of LB twice. Supernatant
and EDTA-free Protease Inhibitor Cocktail were added into the beads and rotating on
the wheel at 4°C for 60 min. 10% of volume of pre-cleared chromatin was taken as
"Input" and stored at 4°C. Antibody conjugated beads were washed with 1 mL of LB
twice. Pre-cleared chromatin was added to washed beads and incubated at 4°C for 6 h.
After incubation, beads were washed with 1 mL of Washing buffer A (50 mM Tris-HCl
pH 7.5, 10 mM EDTA and 75 mM NaCl) on the wheel at 4C for 5 min and buffer was
discarded carefully. Subsequently, beads were washed wish 1 ml of Washing buffer B
(50 mM Tris-HCl pH 7.5, 10 mM EDTA and 125 mM NaCl) and buffer was discarded
carefully. Another 1 mL of Washing buffer B was added and transferred to 1.5 mL
Protein lo-bind tubes (Eppendorf, 0030108116) with beads. Buffer was discarded
carefully. 150 µL of elution buffer (1:9 mixture of 10 % SDS and TE) was added. Tubes
were placed at 25°C on the Eppendorf thermomixer and shaken at 800 rpm for 15 min.
Supernatant were taken and transferred to the new tube using magnetic stand. Another
150 µL of elution buffer was added and repeated this step (in total 300 µl). Input
samples were diluted to 300 µL total by adding TE. 2 µL of 2 mg/ml RNase A (heat-
inactivated or DNase-free) was added and incubated at 37°C for 30 min. 4 µL of 10
mg/ml Proteinase K was added and incubated at 55°C overnight. After incubation,
tubes were shortly spun. 300 µL of Phenol/chloroform were added and vortexed for 30
sec. Samples were centrifuged at 13000 rpm for 10 min at room temperature. 1 µL of
Glycogen, 1/10 volume of NaAcetate and 2 volumes of 100% EtOH were added and
vortexed. Tubes were stored on dry ice for 20-30 min. Once frozen, tubes were
centrifuged at 13000 rpm for 30 min at 4°C. Supernatant was discarded carefully and
500 µL of 70% EtOH was added. Tubes were centrifuged at 13000 rpm for 10 min at
4°C. Pellets were dried and suspended in 30 µL of ddH$_2$O.

Embryo Chromatin Immunoprecipitation (ChIP). Batches of 200 in vitro fertilized
embryos were fixed in 15 ml 1 % Paraformaldehyde in 1X MMR for 20min at room
temperature followed by 4 washes with 1X MMR. Embryos were split in two and
equilibrated at 4°C overnight in 500 µL HEG solution (50 mM HEPES-KOH pH 7.5,
1 mM EDTA, 20 % Glycerol), then excess buffer was discarded, and samples were
transferred to -80°C. For chromatin extraction embryos were resuspended in 700 µL of
sonication buffer (20 mM Tris-HCl pH 8.0, 70 mM KCl, 1 mM EDTA pH 8.0, 10%
Glycerol, 5 mM DTT, 0.125% NP40, 1X complete protease inhibitors) and
homogenized by pipetting up and down. 200 µL embryo extract were distributed in 1.5
mL tubes and sonicated in two sets of 15 min cycles (with 30 s on/off cycles). Sonicated
samples were centrifuged at 12000 rpm for 5 min at 4 °C a Supernatants were retrieved
and the centrifugation was repeated. Before ChIP, 180 µL (2x 10%) of the chromatin
extract was taken as an Input control and stored at -20°C. For each ChIP, 720 µL of
chromatin extract was mixed with 2 µL of anti-H2AK119ub1 antibodies (Cell
Signalling, 8240) and incubated overnight on a rotating wheel. The next day, 40 µL (20
µL per IP) of Magnetic beads (M280 Sheep anti-rabbit IgG, Invitrogen, #11204D) were
washed with 1X PBS using a magnet and incubated with 1mL of 1X PBS containing 1
mg/mL BSA for 2 h at 4°C on a rotating wheel. Blocking solution was discarded from
the beads using a magnet and beads were resuspended in 200 µL (100 µl per IP) of
incubation buffer (50 mM Tris-HCl pH 8, 100 mM NaCl, 2 mM EDTA, 1% NP40, 1
mM DTT, 1X Protease inhibitor). 100 µL of incubation buffer containing beads is
transferred to chromatin/antibody mix and incubated for 4 h at 4°C. Samples were
washed with Buffer 1 (50 mM Tris-HCl pH 8, 100 mM NaCl, 2 mM EDTA, 1 mM
DTT, 1 % NP40, 0.1 % sodium deoxycholate, 1X complete protease inhibitors) for 5

min at 4°C on rotating wheel. Using a magnet, supernatant was discarded and beads
subsequently washed with Buffer 2 (50 mM Tris-HCl pH 8, 500 mM NaCl, 2 mM
EDTA, 1 mM DTT, 1 % NP40, 0.1 % sodium deoxycholate, 1X complete protease
inhibitors), Buffer 3 (50 mM Tris-HCl pH 8, 100 mM NaCl, 250 mM LiCl, 2 mM
EDTA, 1 mM DTT, 1 % NP40, 0.1 % sodium deoxycholate, 1X complete protease
inhibitors), Buffer 1 after which immunoprecipitated chromatin was eluted using 400
μL (310 μL for Input fraction) of Elution buffer (0.1 M NaHCO3, 1 % SDS) overnight
at 65°C. The next day, 4 μL of 20 mg/mL RNAse A was added to the chromatin solution
and incubated for 2 h at 37°C. After incubation, tubes were shortly spun. 300 μL of
Phenol/chloroform were added and vortexed for 30 sec. Samples were centrifuged at
13,000 rpm for 10 min at room temperature. 1 ul of Glycogen, 1/10 volume of
NaAcetate and 2 volumes of 100% EtOH were added and vortexed. Tubes were stored
on dry ice for 20-30 min. Once frozen, tubes were centrifuged at 13,000 rpm for 30 min
at 4C. Supernatant was discarded carefully and 500 μL of 70% EtOH was added. Tubes
were centrifuged at 13,000 rpm for 10 min at 4C. Pellets were dried and suspended in
30 μl of ddH2O.

Xenopus **sperm collection** *Xenopus* **sperm collection.** *Xenopus* sperm collection was
performed as described before[10,44]. For each round of sperm purification, testes from
six adult *Xenopus laevis* males were isolated and manually cleaned from blood vessels
and fat bodies in 1X Marc's Modified Ringers (MMR, 100 mM NaCl, 2 mM KCl, 1
mM MgSO$_4$, 2 mM CaCl$_2$, 3 mM HEPES, pH 7.4) using forceps and paper tissues.
Subsequently, testes were torn into small pieces with forceps and homogenized with 2–
3 strokes of a Dounce homogenizer (one testis at a time). The cell suspension was then
filtered to remove tissue debris and cell clumps using a 60 μm Mesh (CellTrics, cat.
04-0042-2317) and spun down at 800 rcf, 4°C, for 20 min. Supernatant was discarded
and the cell pellet was resuspended in 12 mL of 1 × MMR. If any red blood cells were
visible at the bottom of the pellet (as a result of incomplete removal of blood vessels),
only the uncontaminated part of the pellet was recovered, taking extreme care not to
disturb the red blood cells. Subsequently, step gradients of iodixanol (Optiprep; Sigma,
D1556; 60% iodixanol in water) in 1X MMR final were manually prepared in pre-
chilled 14 mL ultra- clear centrifuge tubes (Beckman Coulter, #344060) in the
following order from the bottom to the top of the tube: 4 mL of 30 % iodixanol, 1 mL
of 20 % iodixanol, 5 mL of 12 % iodixanol (all in 1 × MMR), and 2 mL of cell
suspension in 1X MMR on top. Gradients were spun down in a pre-chilled SW40Ti
rotor at 7500 rpm (10,000g), 4°C, for 15 min, deceleration without brake (Beckman
Coulter Ultracentrifuge, Optima L-100XP). The pelleted fraction, containing mature
sperm, was collected. Collected fractions were diluted six times with 1X MMR and
collected by spinning first at 805 rcf, 4°C, for 20 min and then at 3220 rcf, 4°C, for 20
min to pellet remaining cells. Pelleted cells were subsequently permeabilized with
Digitonin (10 mg/mL as a final concentration, Sigma, D141) for 5 min at room
temperature as detailed before (Smith et al., 2006).

Library preparation and sequencing. ChIP samples were subjected to ChIP-seq
library preparation using QIAseq Ultralow Input Lib UDI-A Kit (Qiagen, 180497) and
Agencourt AMPure XP beads (Beckman coulter, A63881). Samples pre- and post-
library protocol were analyzed on a LabChip GX (PerkinElmer) using DNA 1KHS and
1K Assay LabChIP, respectively. Libraries were quantitated using NEBNext Library
Quant Kit (NEB, E7630,). For RNA sequencing, RNAs were purified using Qiagen
RNeasy Mini Kit (#74004) and used to generate cDNA sequencing libraries using an
Illumina TruSeq Kit (#RS-122-2001), according to the manufacturer's specifications.

Paired-end sequencing was carried out either on a NOVA-seq 6000 or HiSeq 1500
sequencer (Illumina).

RNA-Seq analysis. Adapters were trimmed and low-quality reads filtered using
cutadapt (v1.18)[46]. Sample quality was confirmed with fastqc (v0.11.8)[47]. fastq were
aligned to *Xenopus laevis* genome (v9.2) using STAR aligner (v2.7.3.a)[48]. Count table
was generated using HTSeq-count (v0.12.3)[49] and exon as genes reference. MultiQC
(v1.9)[50] summarized data pre-processing. Differential gene expression analysis of the
count table was performed on R (v3.6)[51] using DESeq2tools[52]. Details of the pipeline
used can be found at https://gitlab.univ-nantes.fr/E114424Z/BulkRNAseq. Gene
ontology analysis were carried out with gprofiler[53] using human orthologs of *Xenopus*
laevis genes recovered from Xenbase (https://www.xenbase.org/xenbase/static-
xenbase/ftpDatafiles.jsp)[54]. *Xenopus laevis* development RNA-seq data were from
published dataset GSE73430.

ChIP-Seq analysis. Adapters were trimmed and low-quality reads filtered using
cutadapt (v1.18)[46]. Sample quality was confirmed with fastqc (v0.11.9)[47]. Alignment
was performed with bwa mem[55] against *Xenopus laevis* (v9.2) or against a *Xenopus*
laevis (v9.2) - 601DNA hybrid genome generated with bwa index (v0.7.17-r1188).
Reads extraction and analysis was performed with samtools view and samtools flagstat
(v1.12)[56]. ChIP-Seq sample quality was assessed by Deeptools (v3.5.1)[57] using
plotFingerprint function and bamPEFragmentSize. Replicates quality was visualised
with plotCorrelation using spearman correlation after genome fragmentation into bins
of 10Kb. PCR duplicates were marked by Picard tools using MarkDuplicates command
(https://broadinstitute.github.io/picard) and then removed with samtools view -F 0x400.
Replicates were pooled by subsampling the same amount of reads per replicates using
sambamba (v0.8.1)[58] and samtools view (-s option). Peak calling was processed with
MACS2 (https://github.com/macs3-project/MACS)[59] (-q 0.01 --nomodel --extsize 147
--gsize=2.6e9) for H3K4me3, with MACS2 broad option for H3K27me3 (--broad --
broad-cutoff 0.1 -q 0.01 --nomodel --extsize 147 --gsize=2.6e9) and RECOGNICER[24]
(qvalue 0.001) /MACS2 (--broad --broad-cutoff 0.1 -q 0.01 --nomodel --extsize 147)
for H2AK119ub1. SICER peak calling was performed using the following parameters
-s XL -w 200 -rt 1 -f 150 -egf 0.74 -fdr 0.001 -g 600 -e 1000. Peaks overlap was defined
using bedtools intersect having at least 1 bp of overlap between peaks. H2AK119ub1
signal was calibrated using reads mapping to the 601DNA and corresponding to the
fully ubiquitylated H2AK119 nucleosome spiked-in in samples prior to ChIP (see ChIP
M&M section). To evaluate difference in modifed histone ChIP-seq signal around the
TSS of gene sets, Wilcoxon Mann-Whitney test was performed on the +/-1 kb window
around the TSSs. ChIPSeeker[60] and ChIPPeakAnno[61,62] were used to annotate peaks on
R[51]. The RegioneR[63] package was used to perform peaks permutation and assess
enrichment on USP21 sensitive gene features (*i.e.* TSS, enhancer, or gene body)
compared to a random set of genes (using 10000 iterations of the resampleRegions
function, , for Fig.3E, .3H, .S3C two right panels, .S3E, .S10G, .S10I) and to assess
peak enrichment on genomic features (using 1000 iterations of the randomizeRegions
function, for Fig 1G, 2K, 2M, S3C two left panels, S10H, S10K). Log10 (ChIP/Input)
was computed using Deeptools with BamCompare using 50bp binsize. Heatmap and
TSS plot were produced with computeMatrix and plotHeatmap or plotProfile. IGV was
used for peaks and log10(ChIP/Input) signal visualisation. Heatmap were produced

with computeMatrix using 50bp binsize +/-5kb of TSS or peak center. Heatmaps were
generated with plotHeatmap function using unsupervised clustering (--hclust option),
and metaplot using plotProfile function. IGV[64] was used to visualize output signal
tracks. Bedtools getfasta[65] was used to extract FASTA sequences corresponding to
peaks from a selected heatmap cluster. To identify transcription factors motif in the
selected peak set, Homer[66] was then used with the findMotifs.pl function and all other
peaks of the heatmap as a background set. CpG islands and repeats elements were
recovered from UCSC genome annotation database
(https://hgdownload.soe.ucsc.edu/goldenPath/xenLae2/database/). For genome
partitioning approach, chromHMM model was initialized using BinarizeBam on ChIP
and Input sample file, using binsize of 150bp (-b 150). The model was trained using a
random initiation (-init random), over binsize of 150bp (-b 150) and a maximum of
5000 iterations (-r 5000). SNPsplit was used for reads assignment to paternal and
maternal copies as described in[19].

Data Availability. The *Xenopus laevis* ChIP-seq and RNA-seq data generated for this
study will be referenced as PRJEB56442 at the European Genome-phenome Archive
(EGA). Additional dataset used are from GSE73430 (Xenopus embryo RNA-seq),
GSE125982 (sperm and replicated sperm H3K27me3 ChIP-seq), GSE126599
(Gastrulae H3K27me3, , and H3K4me3 ChIP-seq), and GSE75164 (sperm MBD-seq,
spermatid H3K27me3), and GSE153531 (H2AK119ub1Cut&Run mouse embryos and
gametes).

Code Availability. All scripts used for these analyses are available on Github
(https://github.com/FCValentin/H2AubPaper).

AUTHORS CONTRIBUTION

Conceptualization: JJ; Methodology: JJ, VF-C, FB, LP; Validation: JJ, VF-C, FB;
Formal analysis: VF-C, FB, MO, MG, VC; Investigation: JJ, VF-C, FB, NM, CF, JP,
RG, MO, BH, KP, JGA; Writing original draft: JJ; Writing review and editing: JJ, FB,
VFC. Visualisation: FB, VF-C; Supervision: JJ; Project administration: JJ; Funding
acquisition: JJ.

COMPETING INTERESTS

The authors declare no competing interests

ACKOWLEDGEMENTS

J.J group is funded by ANR-21-CE13-0031-01 grant from the Agence Nationale de la
Recherche and a Connect Talent grant from Ville de Nantes and Region pays de la
Loire. M.O is funded by a postdoctoral fellowship from the Japan Society for the
Promotion of Science (JSPS). R.G was supported for this work by a Career
Development Award grant (CDA00019/2019-C) form the Human Frontier Science

Program.. L.P group is supported by NIH award R24 OD031956. We would like to
acknowledge the support of Xenbase.
966
967

REFERENCE

1. Öst, A. *et al.* Paternal diet defines offspring chromatin state and
intergenerational obesity. *Cell* (2014) doi:10.1016/j.cell.2014.11.005.

2. Dias, B. G. & Ressler, K. J. Parental olfactory experience influences behavior
and neural structure in subsequent generations. *Nat. Neurosci.* (2014)
doi:10.1038/nn.3594.

3. Radford, E. J. *et al.* In utero undernourishment perturbs the adult sperm
methylome and is linked to metabolic disease transmission. *Science (80-.).*
345, 1–26 (2015).

4. Ben Maamar, M., Beck, D., Nilsson, E. E., Kubsad, D. & Skinner, M. K.
Epigenome-wide association study for glyphosate induced transgenerational
sperm DNA methylation and histone retention epigenetic biomarkers for
disease. *Epigenetics* **16**, (2021).

5. Lismer, A., Siklenka, K., Lafleur, C., Dumeaux, V. & Kimmins, S. Sperm
histone H3 lysine 4 trimethylation is altered in a genetic mouse model of
transgenerational epigenetic inheritance. *Nucleic Acids Res.* **48**, (2020).

6. U., B. *et al.* Repressive and active histone methylation mark distinct promoters
in human and mouse spermatozoa. *Nature Structural and Molecular Biology*
(2010).

7. Hammoud, S. S. *et al.* Distinctive chromatin in human sperm packages genes
for embryo development. *Nature* (2009) doi:10.1038/nature08162.

8. Wu, S. F., Zhang, H. & Cairns, B. R. Genes for embryo development are
packaged in blocks of multivalent chromatin in zebrafish sperm. *Genome Res.*
(2011) doi:10.1101/gr.113167.110.

9. Teperek, M. *et al.* Sperm is epigenetically programmed to regulate gene
transcription in embryos. *Genome Res.* **26**, (2016).

10. Oikawa, M. *et al.* Epigenetic homogeneity in histone methylation underlies
sperm programming for embryonic transcription. *Nat. Commun.* (2020)
doi:10.1038/s41467-020-17238-w.

11. Siklenka, K. *et al.* Disruption of histone methylation in developing sperm
impairs offspring health transgenerationally. *Science (80-.).* (2015)
doi:10.1126/science.aab2006.

12. Murphy, P. J., Wu, S. F., James, C. R., Wike, C. L. & Cairns, B. R. Placeholder
Nucleosomes Underlie Germline-to-Embryo DNA Methylation
Reprogramming. *Cell* (2018) doi:10.1016/j.cell.2018.01.022.

13. Blackledge, N. P. *et al.* Variant PRC1 complex-dependent H2A ubiquitylation
drives PRC2 recruitment and polycomb domain formation. *Cell* **157**, (2014).

14.   Tamburri, S. *et al.* Histone H2AK119 Mono-Ubiquitination Is Essential for
1006   Polycomb-Mediated Transcriptional Repression. *Mol. Cell* **77**, (2020).

1007   15.   Nakagawa, T. *et al.* Deubiquitylation of histone H2A activates transcriptional
1008   initiation via trans-histone cross-talk with H3K4 di- and trimethylation. *Genes*
1009   *Dev.* (2008) doi:10.1101/gad.1609708.

1010   16.   Joo, H. Y. *et al.* Regulation of cell cycle progression and gene expression by
1011   H2A deubiquitination. *Nature* (2007) doi:10.1038/nature06256.

1012   17.   Cao, J. & Yan, Q. Histone ubiquitination and deubiquitination in transcription,
1013   DNA damage response, and cancer. *Frontiers in Oncology* (2012)
1014   doi:10.3389/fonc.2012.00026.

1015   18.   Chen, D., Dai, C. & Jiang, Y. Histone H2A and H2B Deubiquitinase in
1016   Developmental Disease and Cancer. *Cancer Transl. Med.* (2015)
1017   doi:10.4103/2395-3977.168578.

1018   19.   Chen, Z., Djekidel, M. N. & Zhang, Y. Distinct dynamics and functions of
1019   H2AK119ub1 and H3K27me3 in mouse preimplantation embryos. *Nat. Genet.*
1020   **53**, (2021).

1021   20.   Hickey, G. J. M. *et al.* Establishment of developmental gene silencing by
1022   ordered polycomb complex recruitment in early zebrafish embryos. *Elife* **11**,
1023   (2022).

1024   21.   Conway, E. *et al.* BAP1 enhances Polycomb repression by counteracting
1025   widespread H2AK119ub1 deposition and chromatin condensation. *Mol. Cell*
1026   **81**, (2021).

1027   22.   Albright, S. C., Nelson, P. P. & Garrard, W. T. Histone molar ratios among
1028   different electrophoretic forms of mono- and dinucleosomes. *J. Biol. Chem.*
1029   **254**, (1979).

1030   23.   Matsui, S., Seon, B. K. & Sandberg, A. A. Disappearance of a structural
1031   chromatin protein A24 in mitosis: Implications for molecular basis of
1032   chromatin condensation. *Proc. Natl. Acad. Sci. U. S. A.* **76**, (1979).

1033   24.   Zang, C., Wang, Y. & Peng, W. RECOGNICER: A coarse-graining approach
1034   for identifying broad domains from ChIP-seq data. *Quant. Biol.* **8**, (2020).

1035   25.   Heard, E. & Disteche, C. M. Dosage compensation in mammals: Fine-tuning
1036   the expression of the X chromosome. *Genes and Development* vol. 20 (2006).

1037   26.   Jullien, J. *et al.* Gene Resistance to Transcriptional Reprogramming following
1038   Nuclear Transfer Is Directly Mediated by Multiple Chromatin-Repressive
1039   Pathways. *Mol. Cell* **65**, (2017).

1040   27.   Kuznetsov, J. N. *et al.* BAP1 regulates epigenetic switch from pluripotency to
1041   differentiation in developmental lineages giving rise to BAP1-mutant cancers.
1042   *Sci. Adv.* **5**, (2019).

1043   28.   Jin, J. *et al.* The deubiquitinase USP21 maintains the stemness of mouse
1044   embryonic stem cells via stabilization of Nanog. *Nat. Commun.* (2016)

1045   doi:10.1038/ncomms13594.

1046   29.   Gaggioli, V., Le Viet, B., Germe, T. & Hyrien, O. DNA topoisomerase IIα
1047   controls replication origin cluster licensing and firing time in Xenopus egg
1048   extracts. *Nucleic Acids Res.* **41**, (2013).

1049   30.   Blackledge, N. P. & Klose, R. J. The molecular principles of gene regulation
1050   by Polycomb repressive complexes. *Nature Reviews Molecular Cell Biology*
1051   vol. 22 (2021).

1052   31.   Zahn, N., Levin, M. & Adams, D. S. The zahn drawings: New illustrations of
1053   xenopus embryo and tadpole stages for studies of craniofacial development.
1054   *Dev.* **144**, (2017).

1055   32.   Hörmanseder, E. *et al.* H3K4 Methylation-Dependent Memory of Somatic Cell
1056   Identity Inhibits Reprogramming and Development of Nuclear Transfer
1057   Embryos. *Cell Stem Cell* **21**, (2017).

1058   33.   Session, A. M. *et al.* Genome evolution in the allotetraploid frog Xenopus
1059   laevis. *Nature* **538**, (2016).

1060   34.   Chen, H. & Good, M. C. Imaging nascent transcription in wholemount
1061   vertebrate embryos to characterize zygotic genome activation. in *Methods in*
1062   *Enzymology* vol. 638 (2020).

1063   35.   Chen, H., Einstein, L. C., Little, S. C. & Good, M. C. Spatiotemporal
1064   Patterning of Zygotic Genome Activation in a Model Vertebrate Embryo. *Dev.*
1065   *Cell* **49**, (2019).

1066   36.   Tsuboi, M. *et al.* Ubiquitination-Independent Repression of PRC1 Targets
1067   during Neuronal Fate Restriction in the Developing Mouse Neocortex. *Dev.*
1068   *Cell* **47**, (2018).

1069   37.   Żylicz, J. J. *et al.* The Implication of Early Chromatin Changes in X
1070   Chromosome Inactivation. *Cell* **176**, (2019).

1071   38.   Mei, H. *et al.* H2AK119ub1 guides maternal inheritance and zygotic deposition
1072   of H3K27me3 in mouse embryos. *Nat. Genet.* **53**, (2021).

1073   39.   Van Heeringen, S. J. *et al.* Principles of nucleation of h3k27 methylation
1074   during embryonic development. *Genome Res.* (2014)
1075   doi:10.1101/gr.159608.113.

1076   40.   Akkers, R. C. *et al.* A Hierarchy of H3K4me3 and H3K27me3 Acquisition in
1077   Spatial Gene Regulation in Xenopus Embryos. *Dev. Cell* (2009)
1078   doi:10.1016/j.devcel.2009.08.005.

1079   41.   Zhao, J. *et al.* RYBP/YAF2-PRC1 complexes and histone H1-dependent
1080   chromatin compaction mediate propagation of H2AK119ub1 during cell
1081   division. *Nat. Cell Biol.* **22**, (2020).

1082   42.   Lismer, A. *et al.* Histone H3 lysine 4 trimethylation in sperm is transmitted to
1083   the embryo and associated with diet-induced phenotypes in the offspring. *Dev.*
1084   *Cell* **56**, (2021).

43. Pepin, A. S., Lafleur, C., Lambrot, R., Dumeaux, V. & Kimmins, S. Sperm
histone H3 lysine 4 tri-methylation serves as a metabolic sensor of paternal
obesity and is associated with the inheritance of metabolic dysfunction. *Mol.*
Metab. **59**, (2022).

44. Teperek, M. *et al.* Sperm and spermatids contain different proteins and bind
distinct egg factors. *Int. J. Mol. Sci.* **15**, (2014).

45. Hannak, E. & Heald, R. Investigating mitotic spindle assembly and function in
vitro using Xenopus laevis egg extracts. *Nat. Protoc.* **1**, (2006).

46. Martin, M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet.journal* **17**, (2011).

47. Andrews, S. & others. FastQC: a quality control tool for high throughput
sequence data. 2010.
Https://Www.Bioinformatics.Babraham.Ac.Uk/Projects/Fastqc/ (2019).

48. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*
29, (2013).

49. Anders, S., Pyl, P. T. & Huber, W. HTSeq-A Python framework to work with
high-throughput sequencing data. *Bioinformatics* **31**, (2015).

50. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize
analysis results for multiple tools and samples in a single report. *Bioinformatics*
32, (2016).

51. Dessau, R. B. & Pipper, C. B. [''R"--project for statistical computing]. *Ugeskr.*
Laeger **170**, (2008).

52. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, (2014).

53. Raudvere, U. *et al.* G:Profiler: A web server for functional enrichment analysis
and conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, (2019).

54. Fisher, M. *et al.* Xenbase: Key features and resources of the Xenopus model
organism knowledgebase. *Genetics* **224**, (2023).

55. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-
Wheeler transform. *Bioinformatics* **25**, (2009).

56. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,
(2021).

57. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res.* **44**, (2016).

58. Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J. & Prins, P. Sambamba:
Fast processing of NGS alignment formats. *Bioinformatics* **31**, (2015).

59. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**,
(2008).

60. Wang, Q. *et al.* Exploring Epigenomic Datasets by ChIPseeker. *Curr. Protoc.*
2, (2022).

61. Zhu, L. J. *et al.* ChIPpeakAnno: A Bioconductor package to annotate ChIP-seq
and ChIP-chip data. *BMC Bioinformatics* **11**, (2010).

62. Zhu, L. J. Integrative analysis of ChIP-chip and ChIP-seq dataset. *Methods*
Mol. Biol. **1067**, (2013).

63. Gel, B. *et al.* RegioneR: An R/Bioconductor package for the association
analysis of genomic regions based on permutation tests. *Bioinformatics* **32**,
(2016).

64. Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics
Viewer (IGV): High-performance genomics data visualization and exploration.
Brief. Bioinform. **14**, (2013).

65. Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for
comparing genomic features. *Bioinformatics* **26**, (2010).

66. Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription
Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell
Identities. *Mol. Cell* **38**, (2010).

67. Flury,V. *et al.* Recycling of modified H2A-H2B provides short-term memory
of chromatin states. *Cell* **186**, (2023).

1142

1143

(A)

(B)

(C)

FIGURE 1

FIGURE 2

(A)

(B)

(C)

FIGURE 3

FIGURE 4

(A)

(B)

(C)

(D)

FIGURE 5

FIGURE 6

FIGURE 7