Sperm derived H2AK119ub1 is required for embryonic development in *Xenopus Laevis*

Valentin Francois--Campion[1,^], Florian Berger[1,^], Mami Oikawa[2], Maissa Goumeidane[1], Romain Gibeaux[3], and Jérôme Jullien[1,*]

AFFILIATIONS

[1] Nantes Université, CHU Nantes, INSERM, Center for Research in Transplantation and Translational Immunology, UMR 1064, F-44000 Nantes, France

[2] Laboratory of Regenerative Medicine, Tokyo University of Pharmacy and Life Science, Hachioji, Tokyo, 192-0392, Japan

[3] Univ Rennes, CNRS, IGDR (Institut de Génétique et Développement de Rennes) - UMR 6290, F-35000 Rennes, France

^equal contribution

*Corresponding author email: jerome.jullien@inserm.fr

ABSTRACT [140words]

Deposition of H2AK119ub1 by the polycomb repressive complexe-1 plays a key role in the initiation of facultative heterochromatin formation in somatic cells. Here we evaluate the contribution of sperm derived H2AK119ub1 to embryo development. In *Xenopus laevis* we found that H2AK119ub1 is retained during spermiogenesis and into early embryonic development, highlighting its credential for a role in the transmission of epigenetic information from the sperm to the embryo. *In vitro* treatment of sperm with USP21, a H2AK119ub1 deubiquitylase, just prior to injection to egg, results in developmental defects associated with gene upregulation. Sperm H2AK119ub1 editing disrupts egg factor mediated paternal chromatin remodelling processes. It leads to post-replication accumulation of H2AK119ub1 on repeat element of the genome instead of CpG islands. This shift in post-replication H2AK119ub1 distribution triggered by sperm epigenome editing entails a loss of H2AK119ub1 from genes misregulated in embryos derived from USP21 treated sperm. We conclude that sperm derived H2AK119ub1 instructs egg factor mediated epigenetic remodelling of paternal chromatin and is required for embryonic development.

INTRODUCTION

Several studies point towards an epigenetic contribution of sperm for embryonic development. First, in several species paternal exposure to various environmental insults results in changes to the offspring [1–5]. In addition to such environmentally

induced epigenetic changes, sperm is also proposed to naturally harbour epigenetic cues
that are required for development. Histones have been considered as possible vectors
for such epigenetic information. Indeed, in several species including human, modified
histones such as H3K4me3 and H3K27me3 are found in the sperm chromatin, in
particular around developmental genes and in a pattern reminiscent of that found in
embryonic stem cells[6–9]. In a previous work we provided evidence that in the frog,
Xenopus laevis some of these modified histones are homogenously distributed in a
population of sperm cells, hence indicating that they have the required attributes for a
role in the regulation of embryonic development[10]. Lastly, interference with histone
variant or post-translational modifications during the formation of the gametes, or after
fertilisation, lead to embryonic gene misregulation[9,11,12]. Although these later
experiments support a functional role for sperm-derived modified histone in embryonic
development, indirect effect of the interference on the process of spermiogenesis or
early embryos development cannot be ruled out. A direct assay to test for the
requirement of sperm modified histones in embryo development is still lacking. This is
due in part to the difficulty to access the chromatin in the highly compacted sperm
nucleus. In this work, we aim to fill this gap and devised a strategy to alter the sperm
epigenome just prior to its introduction to the egg. We specifically considered
H2AK119ub1 as a possible carrier of necessary epigenetic information from the sperm
to the embryos. Indeed, recent works points towards a key role of polycomb repressive
complexe-1 (PRC1), upstream of PRC2, in the establishment of facultative
heterochromatin, both in cultured cells and in embryos[13]. The H2AK119 ubiquitylation
activity of the PRC1 complex appears to be integral part of this mechanism[13,14]. In
Xenopus laevis we found that H2AK119ub1 dynamic during spermiogenesis and early
embryonic development is compatible with a role in the epigenetic programming of
sperm for embryonic development. We applied an *in vitro* epigenetic editing assay that
uses USP21 deubiquitylase activiy [15–18] to erase H2AK119ub1 from sperm chromatin.
Embryos derived from such epigenetically edited sperm are developmentally impaired.
Transcriptome and epigenome analysis indicates that H2AK119ub1, in combination
with H3K4me3, is enriched on a set of embryonic genes misregulated in embryos
derived from USP21 treated sperm. Such USP21 sensitive genes preferentially retain
H2AK119ub1 upon sperm chromatin exposure to maternal factors. Lastly, we provide
evidence that egg factor mediated deposition of H2AK119ub1 on paternal chromatin is
influenced by pre-existing sperm H2AK119ub1.We conclude that sperm derived
H2AK119ub1 instructs egg factor mediated epigenetic remodelling of paternal
chromatin and is required for embryonic development.

RESULTS
We first evaluated the overall levels of H2AK119ub1 in chromatin during
spermiogenesis and early embryonic development. Indeed, the presence of such
modified histone in sperm seems to differ between species, its presence documented in
that of mouse[19] while it appears absent in that of zebrafish[20]. In frogs the last stage of
spermatogenesis is associated with a remodelling of nucleosomes resulting in a partial
loss of H2A/H2B while a full complement of H3/H4 is retained on sperm chromatin
(**Fig. 1A**)[10]. Western Blot (WB) analysis captures this loss of H2A and reveal an overall
decrease in H2AK119ub1 in sperm compared to spermatid (**Fig. 1B**, see H2A/H4 and
H2AK119ub/H4 ratio, respectively). Importantly however, the H2AK119ub1 to H2A
ratio is similar in sperm compared to spermatid, indicating that in this context of
spermiogenesis-associated H2A loss, a significant level of H2AK119ub1 is maintained
in sperm (H2AK119ub1/H2A, **Fig. 1B**). We next assessed H2AK119ub1 occurrence

after fertilisation: in early frog embryos around ZGA (blastulae stage) and later during gastrulation and neurulation. Interestingly, contrary to what is seen with H3K27me3, a PRC2 associated modification, H2AK119ub1 is detected throughout early embryogenesis (**Fig. 1C**). Altogether, these observations highlight H2AK119ub1 credentials for a potential role in the transmission of epigenetic information from sperm to the embryos.

We then carried out H2AK119ub1 ChIP to map the distribution of this histone post-translational modification (PTM) on the sperm genome. Biological triplicates were well correlated and subsequently pooled for downstream analysis (**Suppl Fig.1A-B**). We first assessed the distribution of H2AK119ub1 around the transcription start of all sperm genes. H2AK119ub1 accumulates around the TSS of a fraction of sperm genes (**Fig. 1D**), similar to what is observed in somatic cells[21] and embryos[14,20]. Noticeably, we observed in the top 10% TSS harbouring H2AK119ub1 an enrichment for genes associated with developmental function (**Fig.1D, Table S1**), and reminiscent of the association reported with H3K27me3 in many species[6–9]. We next sought to more generally identify regions associated with H2AK119ub1 genome-wide. H2AK119ub1 is a pervasive histone modification typically present on ~10% of nuclear H2A in somatic cell[22,23] and up to ~40% in germ cells[19]. In *Xenopus* sperm, it accumulates over region covering larger part of the genome than other modified histones with broad distribution such as H3K27me3 (**Suppl Fig.1C**). To identify region of H2AK119ub1 genomic enrichment we therefore used RECOGNICER, a coarse-graining approach designed to capture enrichment over broad scale[24] (**Suppl Fig.1C-F, Table S1**). Although we observed peaks of accumulation around genes, such as illustrated for Wnt1.L, Pax6.L or the Hoxa.S cluster (**Fig. 1E**), a large proportion of H2AK119ub1 also accumulates on the sperm genome in distal region away from genes (**Fig. 1F**). Noticeably we observed that H2AK119ub1 is significantly associated with sperm gene regulatory elements (TSSs and enhancers (**Fig. 1G**)). Epigenetic control of gene expression often entails a combination of epigenetic cues such as those observed in X inactivation[25] or in transcriptional reprogramming[26]. We therefore investigated the relationship between genomic regions enriched in sperm for H2AK119ub1 and for other histone post-translational modifications. We focused on H3K27me3[10], associated with the activity of Polycomb Repressive Complex 1, and H3K4me3[10], a histone modification associated with transcription (Peaks, *e.g.* around Pax6.L and HoxA loci, **Fig. 1E**). We then determined the genome-wide association of H2AK119ub1 with methylated K4 and/or K27 on histone H3. While most H2AK119ub1 peaks exist independently of these methylated histones, ~24% of H2AK119ub1 peaks overlap with either H3K4me3 and/or H3K27me3 peaks (**Fig. 1H**). We observed a very strong promoter bias for peaks of H2AK119ub1 co-occurring with methylated H3K4me3, an expected trend given that H3K4me3 is known to strongly associate with gene promoter (**Fig. 1I**). By contrast, when occurring in the absence of H3K4me3, H2AK119ub1 is located away from promoter and in intergenic regions. Importantly, in sperm chromatin on the whole, H2AK119ub1 is associated to different type of nucleoprotein complexes than that associated with H3K27me3 and H3K4me3 (**Suppl Fig.1G**). So, while we observe co-enrichment of H2AK119ub1 with methylated histone H3 at a subset of genomic loci, those histone modifications are possibly present at those sites on adjacent nucleosomal/subnucleosomal nucleoprotein complexes rather than on the same particle. Altogether, this survey of H2AK119ub1 distribution in sperm reveals its enrichment on gene regulatory region further strengthening its potential for gene regulation across generation.

To further evaluate how sperm H2AK119ub1 could have an impact on
embryonic development, we next sought to follow its fate after fertilisation. Following
delivery to the egg, protamines are removed from sperm chromatin and a canonical
nucleosomal structure is re-instated. In conjunction with these events the sperm
chromatin is replicated. Therefore, the extensive changes to sperm chromatin that
follow fertilization have the potential to profoundly alter the sperm derived modified
histone landscape. To determine the fate of sperm modified histones we turned to
interphase egg extracts that can mimic the sperm chromatin remodelling events that
follow fertilization[10]. We carried out H2AK119ub1 and H3K4me3 ChIP-seq on
untreated sperm, or sperm having gone through egg extract mediated replication (**Fig.
2A**). Combining this dataset with existing spermatid and gastrulae ChIP-seq[9,10,27] we
obtain a global view of H2AK119ub1, H3K4me3, and H3K27me3 dynamic during
spermiogenesis and early embryogenesis (**Fig.2B**). Remarkably, while H3K4me3 and
H3K27me3 are mostly stable, H2AK119ub1 exhibits drastic redistribution throughout
this developmental period. Indeed, a large set of genomic location acquires
H2AK119ub1 following sperm chromatin replication (cluster 4, **Fig.2B**), and includes
a large proportion of gene TSSs (**Suppl Fig. 2A**). To better delineate the principle of
this developmental dynamic, we then compared H2AK119ub1 and H3K4me3 peaks in
untreated (sperm) *versus* egg extract treated sperm (replicated sperm). We confirm a
marked disparity in replication dynamics between sperm H3K4me3, mostly stable
(77% of sperm peaks conserved after replication (19345/24982) and 51% stable peaks
overall (19345/37747)), and sperm H2AK119ub1, mostly changing (23.5% of sperm
peaks maintained after replication (15796/67233) and 15% stable peaks overall
(15796/104956)) (**Fig. 2C, Suppl Fig. 2B**). Most of the replication-induced
redistribution of H2AK119ub1 happens in intergenic region, while H2AK119ub1
peaks stable through this process are mostly located on promoter (**Suppl Fig. 2D**). In
this dynamic landscape, the subset of H2AK119ub1 genomic location associated with
H3K4me3 in sperm appears selectively conserved (93% of sperm bivalent peaks are
maintained after replication (5222/5597) (**Fig. 2C**). Broader H2AK119ub1peaks with
higher level of H2A retention discriminate location where H2AK119ub1/H3K4me3 is
maintained through replication from those where it is lost (**Suppl Fig. 2C**). We
conclude that sperm exposure to egg factors profoundly remodels paternal
H2AK119ub1 genomic distribution.

We next evaluated to which extent such egg factor-mediated H2AK119ub1
remodelling is dependent on incoming sperm chromatin component. To directly
evaluate sperm H2AK119ub1 functional relevance to this process, we set out to erase
this histone modification, after completion of spermatogenesis, and prior to exposure
to egg factors. To that end, we incubated mature sperm in an oocyte extract containing
USP21, a H2A deubiquitylase[15,28] (**Fig.2 D**). In such extract, the sperm nucleus is
partially decondensed, allowing efficient deubiquitylation of H2AK119ub1 while the
levels of trimethylated H3K4, H3K9 and H3K27 remain unchanged (**Fig. 2E, Suppl
Fig. 2E**). Importantly, MNAse digestion of oocyte extract USP21-treated sperm yields
nucleosomal and subnucleosomal size DNA fragments typical to that of naïve *Xenopus*
sperm[9] (**Fig.2F**, lane 1&2; **Suppl Fig. 2F**). This indicates that the overall sperm
chromatin structure is maintained after oocyte extract treatment and does not convert
to a somatic type of chromatin organisation as triggered by egg extract treatment[29]
(**Fig.2F,** lane 3&4; **Suppl Fig. 2F**). We next asked to what extend USP21 mediated
erasure of H2AK119ub1 from sperm chromatin influences its egg factor mediated
redistribution. To that end, we carried out modified histone analysis in sperm treated

with control or USP21 extract oocyte with or without a subsequent incubation in egg
extract (**Suppl Fig. 2G**). Interestingly, WB analysis indicates that H2AK119ub1,
erased from sperm chromatin by USP21 treatment, is at least partially reinstated by
subsequent exposure to egg extract (**Fig. 2G**), indicating the existence of a maternal
H2AK119 ubiquitylation activity independent of pre-existing sperm H2AK119ub1.
Using ChIP-seq, we next ask if sperm H2AK119ub1 erasure prior to exposure to egg
factors affects the post-replication genomic location of H2AK119ub1 on paternal
chromatin. We first focused on sperm genomic sites where we previously detected
accumulation of H3K4me3, H3K27me3, or H2AK119ub1 (as in **Fig. 2B**). This analysis
confirmed that USP21 efficiently erases H2AK119ub1 from its genomic site in sperm
(**Suppl Fig. 2H**). We observed an overall reduction in egg factor mediated
H2AK119ub1 deposition on chromatin derived from USP21+ sperm compared to that
from USP21-sperm (**Suppl Fig. 2H**). We then compared H2AK119ub1 peaks in
samples originating from replicated USP21-treated sperm to that from replicated
control-treated sperm (**Fig. 2H**). Surprisingly, we observed that sperm USP21 treatment
leads not only to loss of H2AK119ub1 peaks post-replication (lost peaks) but also
triggers the appearance of a large set of peaks that is not deposited by maternal factors
in control sperm samples (gained peaks) (**Fig. 2H**). Comparatively, USP21 treatment
impact on postreplication H3K4me3 is much milder (**Suppl Fig. 2I**). To better
characterize this shift, we first evaluated if H2AK119 ubiquitylation sites detected
following egg extract exposure of sperm harbour genomic features associated with
vPRC1 targeting in somatic cell (reviewed in[30]). H2AK119ub1 sites that are specific to
egg extract treated control sperm (lost peaks, **Fig. 2H**) are indeed enriched for TF motif
associated with vPRC1 targeting (*i.e.* USF1/2 E2F, E-box, **Fig. 2I, Table S1**). We also
detected a strong association of these sites with unmethylated CpG island (**Fig. 2J-K**),
a genomic feature also shown to attract vPRC1 activity. By contrast, sperm treatment
with USP21 prior to egg factor exposure redirects H2AK119ub1 accumulation towards
repeat elements and away from such targets (**Fig. 2L-M**). We conclude that maternal
factors mediated redistribution of sperm H2AK119ub1 is instructed by pre-existing
H2AK119ub1 derived from paternal chromatin. In the absence of sperm H2AK119ub1,
maternal factor mediated H2A ubiquitylation activity is shifted from CpG islands to
repeat elements.

To then determine the functional relevance of sperm derived H2AK119ub1 to
embryo development, we took advantage of the availability of paternally derived
embryos in frog. As described previously, such embryos develop solely on chromatin
derived from sperm and provide a direct measure of the paternal contribution to early
development[9]. We generated paternally derived embryos from sperm pre-incubated in
control or USP21 containing oocyte extracts and monitored their development
(**Fig.3A**)[31]. From gastrulation onwards, we observed a decrease in the developmental
potential of USP21 treated- compared to control treated- sperm derived embryos
(**Fig.3B**). We conclude that besides its DNA content, sperm also contributes to the
embryo an USP21 sensitive signal that is necessary for development. The observed
defects occur around the time of zygotic gene activation suggesting that USP21
treatment associated developmental failure might relate to defective regulation of
embryonic gene expression. To evaluate this possibility, we carried out transcriptome
analysis of early gastrulae generated from pool of untreated, control extract-treated or
USP21 extract-treated haploid embryos. We performed RNA-seq analysis using sperm,
oocyte extract, and egg originating from different preparations for each of the biological
replicates. Differential gene expression analysis indicates that the treatment of sperm

with a control oocyte extract has minor effect on the resulting embryo transcriptome
when compared to embryos generated from untreated sperm (**Fig.3C**, left; **Suppl Fig.**
3A-B). By contrast, USP21 extract treatment leads to embryonic misregulation of 269
genes (**Fig.3C**, right), the majority of which are upregulated in embryos derived from
USP21 extract treated sperm compared to embryos derived from control extract treated
sperm (74%, 200/269, **Fig.3D, Suppl Fig. 3C, Table S2**). Thereafter, this set of genes
found differentially expressed in embryos derived from sperm treated with USP21
oocyte will be referred to as USP21 sensitive genes. GO enrichment analysis highlight
genes related to cell cycle processes in this gene set (**Table S2**). We next asked to which
extend the expression pattern of USP21 sensitive genes identified on pooled embryos
reflect the behaviour in single embryos. To that end, we carried out transcriptome
analysis in single paternally derived gastrulae. We find that the expression pattern of
USP21 sensitive gene observed in pooled embryos is replicated in half of the single
embryos derived from USP21-treated sperm (4/8 USP21 treated sperm derived
embryos, **Suppl Fig. 3D**). This heterogeneous transcriptome pattern matches the
observed developmental outcome, with a fraction of gastrulae derived from USP21-
treated sperm developing to the tadpole stage (**Fig.3B**) and is reminiscent of the
transcriptome heterogeneity detected in cloned embryos[32]. We then interrogated the
developmental dynamic of USP21 sensitive genes. We observed a clear distinction
between up- and down regulated USP21 sensitive genes (**Suppl Fig. 3E**). During
development of embryos generated by *in vitro* fertilisation[33], upregulated genes exhibit
a strong maternal contribution followed by clearing prior to gastrulation (stage 10-12)
and re-expression at tadpole stage (stage 25-40) while downregulated genes show
increased expression around gastrulation. A recent study indicates that at ZGA,
transcription occurs on genes with (maternal-zygotic, "MZ genes") or without (zygotic,
"Z genes") a maternal contribution of transcripts[34,35]. Interestingly, we observed that
USP21 sensitive genes are enriched for MZ genes, indicating that these genes exhibit
zygotic transcription in the context of a maternally derived transcripts (**Suppl Fig. 3E-**
F). Taken together, these analyses support the hypothesis whereby USP21 mediated
erasure of the repressive histone modifications H2AK119ub1 from sperm leads to
precocious embryonic expression of a subset of genes that could in turn impact critical
embryonic process such as cell division.

To evaluate this possibility further, we instigated to which extent H2AK119ub1
genomic distribution associate with embryonic gene sensitivity to sperm USP21
treatment. We observed an enrichment of sperm H2AK119ub1 peaks on USP21
sensitive gene body (**Suppl Fig. 3G**). However, many more genes harbour this
modification in sperm, without being affected by USP21 treatment. We therefore next
interrogated the association of USP21 sensitive genes with different types of
H2AK119ub1 peaks stratified according to association with other histone PTMs,
namely H3K4me3 and H3K27me3 (as in **Fig.1G**). We detect an association of USP21
sensitive gene features with either H2AK119ub1 only peaks (H2AK119ub1+;
H3K4me3-; H3K27me3-) or trivalent H2AK119ub1 peaks (H2AK119ub1+;
H3K4me3+; H3K27me3+) (**Fig. 3E**). Strikingly however, USP21 sensitive genes show
a strong enrichment for bivalent H2AK119ub1/H3K4me3 and depletion for bivalent
H2AK119ub1/H3K27me3 chromatin configuration. This analysis suggests that the
effect on embryonic gene expression of sperm USP21 treatment arise from erasure of
H2AK119ub1 in the context of a bivalent H2AK119ub1/H3K4me3 chromatin status.
We also observed that, post-replication, H2AK119ub1 and H3K4me3 signals around
USP21 sensitive gene TSSs are higher than around the bulk of *X. Laevis* genes TSSs

(**Fig. 3F**). Importantly, the significant enrichment of USP21 sensitive genes promoter
and gene body for a bivalent H2AK119ub1/H3K4me3 chromatin configuration
reported in sperm holds true after sperm replication (**Suppl Fig. 3H**). We next
considered the possibility that while a local accumulation of H2AK119ub1 could be
maintained post-replication, sperm modified histones could have undergone replicative
dilution. In other word, during replication the local accumulation of histone along the
genome could be maintained albeit at a reduced level, for example through
redistribution of sperm derived modified histones between the replicated strands of
DNA. Mechanistically, this is important as in frog ~8 rounds of replication separate
fertilisation form the main wave of ZGA and could therefore leads to complete loss of
sperm derived modified histones. Therefore, to compare H2AK119ub1 level pre- and
post- replication we calibrated the signal *via* semi-synthetic H2AK119ub1 nucleosome
spiked-in used in the ChIP-seq (see M&M) (**Fig. 3G**). We observed a global decrease
of H2AK119ub1 signal on TSSs upon egg extract treatment, suggesting that sperm
chromatin replication induces redistribution of this modified histones at these locations
(black line, **Fig. 3G**). When considering USP21 sensitive gene however, we observed
a different trend. While an overall decrease of H2AK119ub1 signal is also detected
upon replication, H2AK119ub1 loss is clearly less pronounced on the TSSs of this
subset of gene than seen genome wide (**Fig. 3G**), a feature shared with the broader set
of gene that have maternal-zygotic contribution (**Suppl Fig. 3I**). These analyses
indicate that USP21 sensitive genes are indeed associated with a sperm H2AK119ub1
configuration that persists through egg factor mediated remodelling of the paternal
epigenome. Lastly, we asked how sperm USP21 treatment induced disruption of
paternal epigenome remodelling by egg factors affects USP21 sensitive genes. We
found that the shift of H2AK119ub1 away from CpG towards repeat elements induced
by USP21 treatment (as in **Fig. 2H**) significantly depleted post replication association
of USP21 sensitive genes from this histone modification (**fig. 3H**). Analysis of gene
TSSs also showed that the preferential post replication retention of H2AK119ub1 on
USP21 sensitive genes is lost in response to H2AK119ub1 sperm editing (**Fig. 3I**).

Taken together our results reveal that the setup of paternally derived zygotic
chromatin is dictated by component of the sperm epigenome and that this process is
required for proper expression of a subset of embryonic genes.

DISCUSSION

In this work, we found that in *Xenopus Laevis*, paternal H2AK119ub1 genomic
distribution is extensively remodelled upon exposure to egg factors, reminiscent of the
situation found in the mouse[19]. This remodelling is characterized by an accumulation
of H2AK119ub1 around unmethylated CpG island and/or TF binding sites (*i.e.*
USF1/2), genomic features that are known target of PRC1 activity. In mouse embryonic
stem cell, the establishment of polycomb repressive domain occurs in a stepwise
manner with a variant PRC1 complex initiating their formation through ubiquitylation
of H2AK119, followed by recruitment of the PRC2 complex and subsequent
methylation of H3K27me3. Such sequential events have also been described during
Neural Progenitor Cell differentiation[36] and X chromosome inactivation[37]. Recently,
this sequence of events have been investigated in the mouse and zebrafish
embryos[19,20,38]. In these embryos H2AK119ub1 is also required for the deposition of
H3K27me3 and transcriptional repression around the time of zygotic gene activation.

Our observation that H2AK119ub1 is present in pre-ZGA embryo in the absence of
H3K27me3 argues for the existence of a similar mechanism in frog[39,40]. Interference
with H2AK119ub1 in the mouse, either by knocking down PRC1 component[38] in
oocyte, or erasing H2AK119ub1 from embryos[19,20] leads to mis-regulation of genes at
ZGA and is associated with embryonic lethality. While clearly demonstrating a role for
H2AK119ub1 in regulation of transcription around ZGA, these approaches entail
prolonged epigenetic interference during either gametogenesis or early embryogenesis
and, as a consequence, do not enable to determine the extent to which embryonic
H2AK119ub1 is instructed/inherited from sperm and/or oocyte
H2AK119ub1[9,11,12,19,20,38]. Here, by editing the sperm epigenome, after completion of
spermiogenesis and prior to exposure to egg factors, we circumvent these limitations.
We demonstrate that sperm contribute to the embryo a USP21 sensitive cue required
for paternal H2AK119ub1 remodelling by egg factor, embryonic gene expression, and
developmental competence (**Fig. 3J**). We find that egg factor mediated accumulation
on paternal CpG island depends on H2AK119ub1 delivered by the sperm. Strikingly,
we observe that erasure of H2AK119ub1 from sperm leads to a global shift in genomic
accumulation from CpG islands to repeat elements. Given the propensity of vPRC1 to
be targeted and to spread from existing H2AK119ub1 location[41] it raises the possibility
that pre-existing sperm derived ''docking sites'' could limit chromatin sampling by the
polycomb complex and prevent H2AK119ub1 deposition at abnormal locations.
Embryos derived from H2AK119ub1 edited sperm exhibit transcriptional
misregulation, mostly upregulation, compatible with the repressive function of this
histone PTM on transcription. A similar observation is observed in mouse when
H2AK119ub1 deubiquitylase is expressed at fertiization[19], or in zebrafish when Rnf2
is chemically inhibited[20]. Given the extent of egg factor mediated H2AK119ub1
accumulation on gene promoters observed in frog, mouse or zebrafish, it is somewhat
surprising that only a limited number of genes are mis-regulated when its distribution
is massively altered. We identify an association of misregulated genes with
H2AK119ub1 in frog sperm and find that this association is preferentially maintained
on these genes upon sperm chromatin replication, further supporting a direct role for
this histone post-translational modification in the epigenetic programming of sperm for
regulation of embryonic transcription. We cannot rule out that a part of the genes is
indirectly affected, for example as downstream target of a "primary" H2AK119ub1
target. Alternatively, we provide evidence that gene sensitivity to H2AK119ub1 erasure
from sperm occurs when genes are associated with H3K4me3, a histone PTM known
to act as a vector of intergenerational information[9–11,42,43]. It is likely that additional
epigenetic features besides H3K4me3 contribute to gene sensitivity to H2AK119ub1
erasure from sperm. Additionally, we observe that most affected genes are normally
expressed late in embryogenesis (tadpole stage) a feature also seen in mouse with a
large proportion of gene sensitive to H2AK119ub1 depletion expressed post-
implantation[8]. It is therefore possible that the effect of H2AK119ub1 expression is only
partially revealed at early embryonic stage since transcription factors targeting these
genes might not be present.

Our work advance knowledge form previous studies by showing that part of
H2AK119ub1 found in embryos is paternally instructed. This necessary function of
sperm derived modified histones has important bearing on our understanding of
inter/trans generational transmission of epigenetic information. One could indeed
envisage that such a naturally occurring mechanism of epigenetic information
transmission to the embryos could be hijacked to transmit environmentally induced

sperm epigenetic cues. Based on the very dynamic nature of histone PTMs around
fertilisation, it is likely that transmission to the embryos will be context- (combination
of marks) and location- dependent. Future studies will aim at evaluating to which sperm
histones PTMs differ between individuals and to which extent they are maintained in
embryos.

FIGURE LEGENDS

<u>**Figure 1:**</u> *H2AK119ub1 associates with gene regulatory regions in sperm chromatin*

(**A**) Schematic representation of *Xenopus laevis* nucleosome remodelling during
spermiogenesis (**B**) Left, Western blot analysis of H2AK119ub1, H2A and H4 histones.
Right quantitation of signal ratio of H2A to H4, H2AK119ub1 to H4, and H2AK119ub1
to H2A, in spermatid and sperm. * $p<0.05$, ratio paired t test (n=4). (**C**) Western blot
analysis of H2AK119ub1, H3K27me3, H3K9me3, H3K4me3, and H4 in *Xenopus*
laevis embryo pre- (blastulae) and post- (gastrulae, and neurula) ZGA. (**D**) Heatmap of
H2AK119ub1 signal (log (ChIP/Input)) on all *Xenopus Laevis* genes TSS +/-5kB. GO
enrichment is shown for the top 10% genes. (**E**) Genome browser view of log ratio of
ChIP to input signal of sperm H2AK119ub1, H3K4me3, and H3K27me3 around *wt1.L,*
pax6.L, snx10.s and *hoxa.S* loci. Boxed area underneath the gene track indicates regions
of enrichment (peaks). (**F**) Distribution of sperm H2AK119ub1 peaks on genomic
features. (**G**) Fold enrichment of H2AK119Ub1 peaks on different genomic features,
*** $p<0.001$, permutation test. (**H**) Pie chart indicating the proportion of H2AK119ub1
peaks either alone, or overlapping H3K4me3 and/or H3K27me3 peaks in sperm. (**I**)
Genomic distribution of sperm H2AK119ub1 peaks according to overlap with
H3K4me3 and/or H3K27me3 peaks.

<u>**Figure 2:**</u> *Sperm H2AK119ub1 is required for the remodelling of paternal chromatin*
by egg factors.

(**A**) Experimental design for monitoring modified histones fate following sperm
exposure to egg factors. (**B**) Heatmap of H2AK119ub1, H3K4me3, or H3K27me3
signal in spermatid, sperm, replicated sperm, and gastrula. All genomic locations
enriched for one of this histone modification (peaks) in at least one of these
developmental stages are represented. (**C**) Venn diagram indicating the changes to
H3K4me3, H2AK119ub1, or overlapping H3K4me3/H2AK119ub1 peaks during
sperm chromatin replication. Left, all H2AK119ub1 peaks; Middle all H3K4me3
peaks; Right, overlapping H2AK119ub1/H3K4me3 peaks. Percentages of sperm peaks
that are maintained after replication are indicated underneath. (**D**) Schematic
representation of sperm treatment with oocyte extract containing USP21. (**E**) Left,
Western blot of H2AK119ub1, H3K4me3, H3K9me3, H3K27me3 and histone H4 on
oocyte extract treated sperm, USP21 treated sperm, or untreated sperm. Right, signal
ratio of H2AK119ub1 to H4 in oocyte extract treated sperm, USP21 treated sperm, or
untreated sperm. ** $p<0.01$, *** $p<0.005$, ratio paired t test (n=4). (n=6) (**F**) DNA
fragments generated by MNase digestion of *Xenopus laevis* sperm, oocyte extract
treated sperm, egg extract treated sperm and somatic cell.
(**G**) Western blot analysis of H2AK119ub1 and histone H4 on sperm after treatment
with control- or USP21- oocyte extract without (left) or with (right) subsequent
replication in egg extract. (**H**) Heatmap of H2AK119ub1 and H3K4me3 signal in sperm
after treatment with control- or USP21- oocyte extract without or with subsequent

replication in egg extract. All genomic locations enriched for H2AK119ub1 (peaks) in
replicated samples are represented. (**I**) Transcription factor motifs enriched in peaks
that are lost, maintained, or gained upon replication of control versus USP21 treated
sperm. (**J-M**) DNAme and H2AK119ub1signal on *X.Laevis* genome CpG island (J)
and repeat element (L). Fold enrichment of H2AK119ub1 peaks on CpG island (K)
and repeat element (M).

Figure 3: *Sperm H2AK119ub1 is required for optimal embryonic development and*
gene expression.

(**A**) Experimental design for the generation of haploid embryos from sperm or USP21
treated sperm. *Xenopus* illustrations © Natalya Zahn (2022) from Xenbase
(www.xenbase.org RRID:SCR_003280). (**B**) Developmental progression to blastula,
gastrula or swimming tadpole stage of haploid embryos derived from control or a
USP21 oocyte extract treated sperm. Chi-squared test, * p-value<0.05. (**C**) MA-Plot
showing as y-axis the log foldchange (LogFC) of gene expression between control
oocyte extract treated sperm *versus* untreated sperm (left) and between USP21 oocyte
extract treated sperm derived embryo and untreated sperm (right). X-axis values are
averaged gene expression across samples. (**D**) Heatmap showing expression of
differentially expressed gene in gastrulae derived from USP21 oocyte extract treated
sperm *versus* that derived from control oocyte extract treated sperm. (**E**) Fold
enrichment of different H2AK119ub1 peaks categories on promoter, enhancer, and
gene body of USP21 sensitive genes. * p-value<0.05; ** p-value < 0.01, permutation
test. (**F**) H2AK119ub1 and H3K4me3 signal on TSS +/- 5kB for all genes (black),
USP21sensitive genes (red) before and after sperm replication. *** p-value<0.001,
Wilcoxon-Mann-Whitney test on TSS+/1kb interval (shaded orange area). (**G**)
Calibrated H2AK119ub1 ChIP-seq signal on TSS +/- 5kB for all genes (black), USP21
genes (red) before and after sperm replication. *** p-value<0.001, Wilcoxon-Mann-
Whitney test. (**H**) Fold enrichment of lost, maintained, or gained H2AK119ub1 peaks
on promoter, enhancer, and gene body of maintained, loss or conserved peaks in
replicated USP21 treated sperm *versus* control sperm. * p-value<0.05; ** p-value<0.01,
***p-value<0.001, permutation test. (**I**) Calibrated H2AK119ub1 ChIP-seq signal on
TSS +/- 5kB for all genes (black), USP21 genes (red) before and after replication using
a WT sperm or USP21 treated sperm. *** p-value<0.001, Wilcoxon-Mann-Whitney
test. (**J**) Model of H2AK119ub1 epigenetic programming of sperm for embryonic
development in *Xenopus Laevis*.

Supplemental Figure S1: *ChIP-seq analysis of H2AK119ub1 distribution of*
H2AK119ub1in sperm chromatin

(**A**) Genome browser view of the log ratio of H2AK119ub1 ChIP to input signal around
the *hoxa*.L cluster on 3 sperm biological replicates. (**B**) Spearman correlation of sperm
and replicated sperm H2AK119ub1, H3K27me3 and H3K4me3 ChIP-seq replicates.
(**C**) Fingerprint plot of input, H3K4me3, H3K27me3, and H2AK119ub1 ChIP samples.
(**D**) Overlap between RECOGNICER and MACS2 peaks for H2AK119ub1, H3K7me3,
and H3K4me3 peaks. (**E**) Heatmap displaying H2AK119ub1 log(ChIP/Input) signal
+/- 5kb from RECOGNICER peaks center (middle) *versus* an equivalent number of
genomic regions randomly selected (left). On the right RECOGNICER peaks are
spitted between those identified by RECOGNICER only versus those detected in both
RECOGNICER and MACS2. (**F**) Genome track showing logratio of ChIP to input
signal for H2AK119ub1 and H3K27me3 as well as the region where peaks are detected

by MACS2 and RECOGNICER (boxed area, red for H2AK119ub1, yellow for
H3K27me3) at different scale view. **(G)** ChIP-seq fragment size distribution for
H2AK119ub1 (red) in sperm and spermatid, and for H3K4me3 (green) and H3K27me3
(yellow) in sperm. Matching inputs are shown in grey.

<u>**Supplemental Figure S2:**</u> *Sperm H2AK119ub1 fate following exposure to egg factors*

(A) Heatmap of H2AK119ub1 (red), H3K4me3 (green), and H3K27me3 (yellow)
log(ChIP/Input) signal ratio on genes TSS from spermatid, sperm prior and after egg
extract treatment, and gastrula stage 12 embryo. **(B)** Scaled Venn diagram of
H2AK119ub1, H3K4me3 or overlapping H2AK119ub1/H3K4me3 peaks in spermatid,
sperm, and sperm exposed to egg factors. Percentage of peaks that are maintained at
each developmental transition are indicated underneath. **(C)** Top: boxplot showing
calibrated H2AK119ub1 on H2AK119ub1/H3K4me3 peaks lost (375), maintained
(5222) or gained (10442) following exposure to egg factors (top). Bottom: peak size
(blue dots) as well as peak H2A enrichment particle (red dots) in each peak category.
(D) Genomic distribution of sperm peaks before or after replication, as well as the
subset of peaks lost, maintained, or gained during replication. **(E)** Quantitation of
H3K4me3, H3K9me3 and H3K27me3 to H4 WB signal ratio in untreated sperm,
oocyte extract treated sperm or USP21 treated sperm. **(F)** Capillary electrophoresis
profiles of DNA size obtained following MNAse digestion of sperm (blue), oocyte
extract treated sperm (cyan), egg extract treated sperm (green) or somatic cell (purple).
(G) Experimental design for monitoring modified histones fate from control or USP21
treated sperm. **(H)** Heatmap of H2AK119ub1 (red), H3K4me3 (green) and H3K27me3
(green) log(ChIP/Input) signal ratio in spermatid, control- and USP21- treated sperm
prior and after egg extract treatment, and gastrula stage 12 embryo. Heatmap includes
the same genomic location as in figure 2B. **(I)** Heatmap of H2AK119ub1 and H3K4me3
signal in sperm after treatment with control- or USP21- oocyte extract without or with
subsequent replication in egg extract. All genomic locations enriched for H3K4me3
(peaks) in replicated samples are shown.

<u>**Supplemental Figure S3:**</u> *Gene expression in gastrulae derived from control or*
USP21 treated sperm.

(A) Principal Component analysis of replicate RNA-seq samples used in this study
(USP21, USP21 extract treated sperm derived gastrulae; Control, control extract treated
sperm derived gastrulae; Untreated, Untreated sperm derived gastrulae. **(B)**
Hierarchical clustering of replicate samples used in this analysis. **(C)** Heatmap
depicting expression of USP21 sensitive genes in replicates from USP21 extract treated
sperm derived gastrulae, control extract treated sperm derived gastrulae, as well as
gastrulae derived from untreated sperm. **(D)** Heat map of gene expression in single
haploid gastrula derived from control- or USP21- treated sperm. Expression is shown
for the USP21 sensitive gene set identified on embryos pool (fig 3D). **(E)** USP21
sensitive gene set expression during early Xenopus embryo development (data from).
Expression is shown for oocyte, egg, blastulae (stage 8-9), gastrulae (stage 10-12),
neurula (stage 15-20), as well as early to late tailbud (stage 25-40) are indicated. The
right-hand side panel shows the expression of USP21 sensitive gene in gastrulae
derived from control- and USP21-treated sperm. **(F)** Violin plot of gene expression in
egg (log of normalised gene count) for all genes (black), USP21 sensitive genes (red),
maternal genes (cyan), maternal and zygotic genes (green) and zygotic only gene
(orange). USP21 sensitive genes are enriched for genes with maternal and zygotic

expression in *Xenopus Laevis*. The table below indicates p-Value resulting from an
hypergeometric test to evaluate enrichment of USP21 sensitive gene for either
maternal-zygotic (MZ) or zygotic only (Z) genes. MZ and Z dataset taken from[34,35].
(G) Fold enrichment of H2AK119ub1 peaks on promoter, enhancer, and gene body of
USP21 sensitive genes. *p-value<0.05; permutation test. **(H)** Fold enrichment of
different H2AK119ub1 peaks categories on promoter, enhancer, and gene body of
USP21 sensitive genes. *p-value<0.05; **p-value < 0.01, ***p-value<0.001,
permutation test. **(I)** Calibrated H2AK119ub1 ChIP signal on TSS +/- 5kb for all genes
(black), USP21 sensitive genes (red), maternal genes (cyan), maternal zygotic genes
(green) and zygotic genes (orange) before and after sperm replication. *** p-
value<0.001, Wilcoxon-Mann-Whitney test.

Table S1*: Germ cell histone PTMs ChIP-seq data*

Table S2: *Control- and epigenetically edited- sperm derived embryos RNA-seq data*

MATERIALS AND METHODS

Contact for Reagent and Resource Sharing. Further information and requests for
resources and reagents should be directed to and will be fulfilled by the Lead Contact,
Dr J. Jullien (jerome.jullien@inserm.fr).

Experimental Model and Subject Details. Mature *Xenopus laevis* females were
obtained from Tefor Paris-saclays (151 Rte de la Rotonde, 91400 Saclay, France,
https://tefor.net/tefor-paris-saclay-nos-equipes/tefor-paris-saclay-aqua/). Our work
with Xenopus complied with all relevant ethical regulations for animal testing and
research. Work with frog was covered under the licence #202212131604754/APAFIS
39794V2 and frog husbandry and all experiments were performed according to the
relevant regulatory standards. Animals were kept in a recirculating system (ALS,
https://www.aquariumlocationservices.fr/professionnels-elevages-aquatique-
aquariophilie/) at a density of one adult/3l, with ~10% water change per day. Water
was sequentially filtered with mechanical pad sump filter, nitrifying bacteria filter, and
UV sterilized. Water quality parameters were as follow: temperature 17-22°C; PH 6-8.
Photoperiod was set to 12h ON/12h OFF. Frogs are fed twice per week with NEO
MARIN coul 4 - X4 - S20 (Le Gouessant, https://www.legouessant.com/).
Unconsumed food was removed 30 min after the start of feeding.

Oocyte extract. Fresh ovaries were manually dissociated in 15-25 oocytes clumps. 3-
5 mL of dissociated ovary fragments were transferred into 12.5mL 1X Modified Barth's
Saline (MBS, 88mM NaCl, 1 mM KCl, 10 mM Hepes, 2.4 mM NaHCO3, 1.7 mM
MgSO4, 1.3 mM Ca(NO3)2, 0.8mM CaCl2) supplemented with 250uL of 5mg/mL
Liberase (Roche, 05 401 127 001) and incubated for 2h30-3h at room temperature (RT)
with gentle rocking. Defolliculated oocytes were washed three times with 1X MBS and
healthy stage 5-6 oocytes manually sorted. Oocytes were then injected in the vegetal
pole with 25ng of USP21-NLS-HA mRNA, or control GFP mRNA and incubated for
3 days at 18°C in 1X MBS, renewing medium and discarding dead cells every day.
~500-1000 oocytes were collected in an Eppendorf tube on ice, the excess 1X MBS
discarded and replaced with 1X MBS supplemented with Protease Inhibitor Cocktail

(Sigma, P2714-1BTL). The oocytes were then stacked up by centrifugation at 100g for 1min at RT and excess medium carefully discarded. Oocytes were then thoroughly vortexed and centrifuged at 21 000g for 10min at 4°C. The clear intermediate phase between yolk (top) and pigment (bottom) was retrieved. The collected fraction was further cleared by an additional centrifugation at 21 000g for 10min at 4°C. The oocyte extract was then aliquoted and stored at -80°C.

Oocyte extract treatment. Permeabilized sperm stored in SSB at -80°C were thawed, washed with two volumes of ice cold 1XPBS and centrifuged at 800g for 8min at 4°C. 2M sperm in 1X PBS were centrifuged at 21 000g for 1min at 4°C. The supernatant was discarded, and the pelleted sperm resuspended in 10uL of oocyte extract on ice. The sperm incubated in the oocyte extract for 20-40min at 23°C on a heating block. Treatment was stopped by addition of 90uL of ice cold 1X PBS and centrifugation at 21 000g for 1min at 4°C. The supernatant discarded and the remaining pellets resuspended in SSB for -80°C long term storage.

Protein extraction and Western Blot analysis. Sperm samples stored in SSB were washed in 1X PBS and 1M sperm pellets were resuspended in 30uL 2X Blue Eye (600mM NaCl, 1% Triton X100, 0.5% Na-Deoxycholate, 4% Sodium Dodecyl Sulfate, 20% Glycerol, 0.0004% Bromophenol blue, 175mM Tris-HCl, 10% Betamercapto ethanol), boiled at 96°C for 5min and centrifuged at 21 000g for 5min at 4°C. The supernatant was transferred to a new tube and directly use for analysis or stored at -20°C. 4-20% mini-PROTEAN® TGX (Biorad, #4561094) were loaded with 10uL of sample and western blot analysis were performed according to standard protocols. For blotting nitrocellulose membranes, a semi-dry transfer was used (Biorad, Trans-Blot® SD Semi-Dry Transfer Cell) 30-40min at 25V. The following primary antibodies were used: anti-H2A (1:2500, Cell Signalling, 3636), anti-H2AK119ub1 (1:2500, Cell Signalling, 8240), anti-H3K4me3 (1:2500, Abcam, ab8580), anti H3K9me3 (1:2500, Abcam, ab8898), anti-H3K27me3(1:2500, Cell Signalling, 9735S), anti-H4 (1:2500, Abcam, ab31830) and anti-HA (1:2500, Sigma, H9658). Membranes were incubated with primary antibodies 2h at RT or overnight (ON) at 4 °C. Secondary antibody are anti-rabbit (1:25000, ThermoFisher, A21076) , anti-mouse IgG Alexa Fluor 680 (1:25000, ThermoFisher ,A21058), Goat anti-rabbit Alexa 800 (1:25,000, ThermoFisher, A32735), or IRDye 800CW Goat Anti-Mouse IgG (H+L) (LI-COR, 926-32210) were used. The membranes were imaged either on a ChemiDoc MP Imaging System (biorad) or an Odyssey scanner (LI-COR).

Sperm Chromatin Immunoprecipitation (ChIP). Chromatin fractionation and chromatin immunoprecipitation (ChIP) were performed as described before[10,44] with slight modifications. 80 µl of Magnetic beads (M280 Sheep anti-rabbit IgG, Invitrogen, #11204D) were used per reaction and all wash steps were carried out with a magnet. Beads were washed with 1 ml of TE pH 8.0, lysis buffer (LB)(1:1 mixture of Buffer 1 and MNase buffer, see below), and incubated in 1 ml of LB with 100 µl of 10 mg/ml BSA at 4C for 30 min ("pre-blocking"). Pre-blocked beads were washed with 1 ml of LB twice and resuspend in 80 µl of LB. Half of beads were incubated with antibody overnight and leftovers were stored in 4C for "pre-clear chromatin" step. For ChIP, the following antibodies were used: anti-H3K4me3 (Abcam, ab8580), anti-H2AK119ub1 (Cell Signaling, 8240). One million digitonin permeabilised sperm nuclei or egg extract treated sperm nuclei were resuspended in 50 µl of Buffer 1 (0.3 M Sucrose, 15 mM Tris pH 7.5, 60 mM KCl, 15 mM NaCl, 5 mM MgCl2, 0.1 mM EGTA, 0.5 mM DTT)

containing 0.125ng H2AK119ub biotinyltated nucleosome (Epicypher #16-0020),
followed by addition of 50 μl of Buffer 1 with detergent (Buffer 1 including 0.5% NP40
and 1% NaDOC). Samples were incubated for 10 min on ice. 100 μl of MNase buffer
(0.3 M Sucrose, 85 mM Tris, 3 mM MgCl$_2$, 2   mM CaCl$_2$, 2.5U of micrococcal
nuclease: Roche 10107921001) was added in to each tube (0.5 millions of cells per
tube). Tubes were incubated at 37C for 30 min in pre-warmed water bath. Reaction was
stopped by adding 2 μl of 0.5 M EDTA pH8.0 in the same order as started. Tubes were
vortexed and placed on ice for at least 5 min. Supernatant and pellet were separated by
centrifugation at 13,000 rpm for 10 min at room temperature. Beads (stored at 4C for
pre-clear chromatin) were washed with 1 ml of LB twice. Supernatant and EDTA-free
Protease Inhibitor Cocktail were added into the beads and rotating on the wheel at 4C
for 60 min. 10% of volume of pre-cleared chromatin was taken as "input" and stored at
4C. Antibody conjugated beads were washed with 1 ml of LB twice. Pre-cleared
chromatin was added to washed beads and incubated at 4C for 6 h. After incubation,
beads were washed with 1 ml of Washing buffer A (50 mM Tris-HCl pH 7.5, 10 mM
EDTA and 75 mM NaCl) on the wheel at 4C for 5 min and buffer was discarded
carefully. Subsequently, beads were washed wish 1 ml of Washing buffer B (50 mM
Tris-HCl pH 7.5, 10 mM EDTA and 125 mM NaCl) and buffer was discarded carefully.
Another 1 ml of Washing buffer B was added and transferred to 1.5 ml Protein lo-bind
tubes (Eppendorf, 0030108116) with beads. Buffer was discarded carefully. 150 μl of
elution buffer (1:9 mixture of 10% SDS and TE) was added. Tubes were placed at 25C
on the Eppendorf thermomixer and shaken at 800 rpm for 15 min. Supernatant were
taken and transferred to the new tube using magnetic stand. Another 150 μl of elution
buffer was added and repeated this step (in total 300   l). Input samples were diluted to
300 μl total by adding TE. 2 μl of 2 mg/ml RNase A (heat-inactivated or DNase-free)
was added and incubated at 37C for 30 min. 4 μl of 10 mg/ml Proteinase K was added
and incubated at 55C overnight. After incubation, tubes were shortly spun. 300 μl of
Phenol/chloroform were added and vortexed for 30 sec. Samples were centrifuged at
13,000 rpm for 10 min at room temperature. 1 ul of Glycogen, 1/10 volume of
NaAcetate and 2 volumes of 100% EtOH were added and vortexed. Tubes were stored
on dry ice for 20-30 min. Once frozen, tubes were centrifuged at 13,000 rpm for 30 min
at 4C. Supernatant was discarded carefully and 500 μl of 70% EtOH was added. Tubes
were centrifuged at 13,000 rpm for 10 min at 4C. Pellets were dried and suspended in
30 μl of ddH$_2$O.

Egg extract treatment. Interphasic extracts were generated as previously
described[45] .For the quantification of DNA replication in individual nuclei in egg
extract, 33 μM rhodamine dUTP (Roche, 11534378910) was added at the indicated
times (20, 40, 60, 80 mins after incubation) for 5 min as indicated, and the reaction was
stopped with 1xPBS and fixed in 4% paraformaldehyde. The reaction was spun through
a 30% sucrose cushion in PBS. The sperm cells were rinsed twice in 1x PBS for 10 min
and mounted in Vectashield mounting medium with DAPI (Vector laboratories, H-
1200). Counting of rhodamine positive cells were performed on an epifluorescence
microscope.

***Xenopus* sperm collection *Xenopus* sperm collection.** *Xenopus* sperm collection was
performed as described before[10,44]. For each round of sperm purification, testes from
six adult *Xenopus laevis* males were isolated and manually cleaned from blood vessels
and fat bodies in 1 × Marc's Modified Ringers (MMR, 100 mM NaCl, 2 mM KCl, 1
mM MgSO$_4$, 2 mM CaCl$_2$, 3 mM HEPES, pH 7.4) using forceps and paper tissues.
Subsequently, testes were torn into small pieces with forceps and homogenized with 2–

3 strokes of a Dounce homogenizer (one testis at a time). The cell suspension was then
filtered to remove tissue debris and cell clumps using a 60μm Mesh (CellTrics, cat. 04-
0042-2317) and spun down at 800 rcf, 4°C, for 20 min. Supernatant was discarded and
the cell pellet was resuspended in 12 mL of 1 × MMR. If any red blood cells were
visible at the bottom of the pellet (as a result of incomplete removal of blood vessels),
only the uncontaminated part of the pellet was recovered, taking extreme care not to
disturb the red blood cells. Subsequently, step gradients of iodixanol (Optiprep; Sigma,
D1556; 60% iodixanol in water) in 1×MMR final were manually prepared in pre-chilled
14 mL ultra- clear centrifuge tubes (Beckman Coulter, #344060) in the following order
from the bottom to the top of the tube: 4 mL of 30% iodixanol, 1 mL of 20% iodixanol,
5 mL of 12% iodixanol (all in 1 × MMR), and 2 mL of cell suspension in 1 × MMR on
top. Gradients were spun down in a pre-chilled SW40Ti rotor at 7500 rpm (10,000g),
4°C, for 15 min, deceleration without brake (Beckman Coulter Ultracentrifuge, Optima
L-100XP). The pelleted fraction, containing mature sperm, was collected. Collected
fractions were diluted six times with 1× MMR and collected by spinning first at 805
rcf, 4°C, for 20 min and then at 3220 rcf, 4°C, for 20 min to pellet remaining cells.
Pelleted cells were subsequently permeabilized with Digitonin (10 mg/mL as a final
concentration, Sigma, D141) for 5 min at room temperature as detailed before (Smith
et al., 2006).

Library preparation and sequencing. ChIP samples were subjected to ChIP-seq
library preparation using QIAseq Ultralow Input Lib UDI-A Kit (Qiagen, 180497)
and Agencourt AMPure XP beads (Beckman coulter, A63881). Samples pre- and
post-library protocol were analyzed on a LabChip GX (PerkinElmer) using DNA
1KHS and 1K Assay LabChIP, respectively. Libraries were quantitated using
NEBNext Library Quant Kit (NEB, E7630,). For RNA sequencing, RNAs were
purified using Qiagen RNeasy Mini Kit (#74004) and used to generate cDNA
sequencing libraries using an Illumina TruSeq Kit (#RS-122-2001), according to the
manufacturer's specifications. Paired-end sequencing was carried out either on a
NOVA-seq 6000 or HiSeq 1500 sequencer (Illumina).

RNA-Seq analysis. Adapters were trimmed and low-quality reads filtered using
cutadapt (v1.18)[46]. Sample quality was confirmed with fastqc (v0.11.8)[47]. fastq were
aligned to *Xenopus Laevis* genome (v9.2) using STAR aligner (v2.7.3.a)[48]. Count table
was generated using HTSeq-count (v0.12.3)[49] and exon as genes reference. MultiQC
(v1.9)[50] summarized data pre-processing. Differential gene expression analysis of the
count table was performed on R (v3.6)[51] using DESeq2tools[52]. Details of the pipeline
used can be found at https://gitlab.univ-nantes.fr/E114424Z/BulkRNAseq. Gene
ontology analysis were carried out with gprofiler[53] using human orthologs of *Xenopus
laevis* genes recovered from Xenbase (https://www.xenbase.org/xenbase/static-
xenbase/ftpDatafiles.jsp)[54]. *Xenopus laevis* development RNA-seq data were from
published dataset GSE73430.

ChIP-Seq analysis. Adapters were trimmed and low-quality reads filtered using
cutadapt (v1.18)[46]. Sample quality was confirmed with fastqc (v0.11.9)[47]. Alignment
was performed with bwa mem[55] against *Xenopus laevis* (v9.2) or against a *Xenopus
laevis* (v9.2) - 601DNA hybrid genome generated with bwa index (v0.7.17-r1188).
Reads extraction and analysis was performed with samtools view and samtools flagstat

(v1.12)[56]. ChIP-Seq sample quality was assessed by Deeptools (v3.5.1)[57] using
plotFingerprint function and bamPEFragmentSize. Replicates quality was visualised
with plotCorrelation using spearman correlation after genome fragmentation into bins
of 10Kb. PCR duplicates were marked by Picard tools using MarkDuplicates command
(https://broadinstitute.github.io/picard) and then removed with samtools view -F 0x400.
Replicates were pooled by subsampling the same amount of reads per replicates using
sambamba (v0.8.1)[58] and samtools view (-s option). Peak calling was processed with
MACS2 (https://github.com/macs3-project/MACS)[59] (-q 0.01 --nomodel --extsize 147
--gsize=2.6e9) for H3K4me3, with MACS2 broad option for H3K27me3 (--broad --
broad-cutoff 0.1 -q 0.01 --nomodel --extsize 147 --gsize=2.6e9) and RECOGNICER[24]
(qvalue 0.001) /MACS2 (--broad --broad-cutoff 0.1 -q 0.01 --nomodel --extsize 147)
for H2AK119ub1. Peaks overlap was defined using bedtools intersect having at least 1
bp of overlap between peaks. H2AK119ub1 signal was calibrated using reads mapping
to the 601DNA and corresponding to the fully ubiquitylated H2AK119 nucleosome
spiked-in in samples prior to ChIP (see ChIP M&M section). To evaluate difference in
modifed histone ChIP-seq signal around the TSS of gene sets, Wilcoxon Mann-
Whitney test was performed on the +/-1 kb window around the TSSs. ChIPSeeker[60] and
ChIPPeakAnno[61,62] were used to annotate peaks on R[51]. RegioneR[63] was used to
perform peaks permutation to assess enrichment on USP21 sensitive gene features (*i.e.*
TSS, enhancer, or gene body) compared to random set of genes (resampleRegions,
10000 iterations) and to assess peak enrichment on genomic features
(randomizeRegions, 1000 iterations) Log10 chip over input was computed using
Deeptools with BamCompare using 50bp binsize. Heatmap and TSS plot were
produced with computeMatrix and plotHeatmap or plotProfile. IGV was used for peaks
and log10(chip/input) signal visualisation. Heatmap were produced with
computeMatrix using 50bp binsize +/-5kb of TSS or peak center. Heatmaps were
generated with plotHeatmap function using unsupervised clustering (--hclust option),
and metaplot using plotProfile function. IGV[64] was used to visualize output signal
tracks. Bedtools getfasta[65] was used to extract FASTA sequences corresponding to
peaks from a selected heatmap cluster. To identify transcription factors motif in the
selected peak set, Homer[66] was then used with the findMotifs.pl function and all other
peaks of the heatmap as a background set. CpG islands and repeats elements were
recovered from UCSC genome annotation database
(https://hgdownload.soe.ucsc.edu/goldenPath/xenLae2/database/). All scripts used for
these analysis are available on Github (https://github.com/FCValentin/H2AubPaper).

Data Availability. The *Xenopus laevis* ChIP-seq and RNA-seq data generated for this
study will be referenced as PRJEB56442 at the European Genome-phenome Archive
(EGA). Additional dataset used are from GSE73430 (Xenopus embryo RNA-seq),
GSE125982 (sperm and replicated sperm H3K27me3 ChIP-seq), GSE126599
(Gastrulae H3K27me3, H2AK119ub1, and H3K4me3 ChIP-seq), and GSE75164
(sperm MBD-seq).

AUTHORS CONTRIBUTION

Conceptualization: JJ; Methodology: JJ, VF-C, FB; Validation: JJ, VF-C, FB; Formal
analysis: VF-C, FB, MO, MG; Investigation: JJ, VF-C, FB, RG, MO, BH; Writing

original draft: JJ; Writing review and editing: JJ, FB, VFC. Visualisation: FB, VF-C;
Supervision: JJ; Project administration: JJ; Funding acquisition: JJ.

COMPETING INTERESTS

759 The authors declare no competing interests

ACKOWLEDGEMENTS

J.J group is funded by ANR-21-CE13-0031-01 grant from the Agence Nationale de la
Recherche and a Connect Talent grant from Ville de Nantes and Region pays de la
Loire. M.O is funded by a postdoctoral fellowship from the Japan Society for the
Promotion of Science (JSPS). R.G was supported for this work by a Career
Development Award grant (CDA00019/2019-C) form the Human Frontier Science
Program. We would like to acknowledge the support of Xenbase.

768
769

REFERENCE

1. Öst, A. *et al.* Paternal diet defines offspring chromatin state and
intergenerational obesity. *Cell* (2014) doi:10.1016/j.cell.2014.11.005.

2. Dias, B. G. & Ressler, K. J. Parental olfactory experience influences behavior
and neural structure in subsequent generations. *Nat. Neurosci.* (2014)
doi:10.1038/nn.3594.

3. Radford, E. J. *et al.* In utero undernourishment perturbs the adult sperm
methylome and is linked to metabolic disease transmission. *Science (80-.).*
345, 1–26 (2015).

4. Ben Maamar, M., Beck, D., Nilsson, E. E., Kubsad, D. & Skinner, M. K.
Epigenome-wide association study for glyphosate induced transgenerational
sperm DNA methylation and histone retention epigenetic biomarkers for
disease. *Epigenetics* **16**, (2021).

5. Lismer, A., Siklenka, K., Lafleur, C., Dumeaux, V. & Kimmins, S. Sperm
histone H3 lysine 4 trimethylation is altered in a genetic mouse model of
transgenerational epigenetic inheritance. *Nucleic Acids Res.* **48**, (2020).

6. U., B. *et al.* Repressive and active histone methylation mark distinct promoters
in human and mouse spermatozoa. *Nature Structural and Molecular Biology*
(2010).

7. Hammoud, S. S. *et al.* Distinctive chromatin in human sperm packages genes
for embryo development. *Nature* (2009) doi:10.1038/nature08162.

8. Wu, S. F., Zhang, H. & Cairns, B. R. Genes for embryo development are
packaged in blocks of multivalent chromatin in zebrafish sperm. *Genome Res.*
(2011) doi:10.1101/gr.113167.110.

9.   Teperek, M. *et al.* Sperm is epigenetically programmed to regulate gene
transcription in embryos. *Genome Res.* **26**, (2016).

10.   Oikawa, M. *et al.* Epigenetic homogeneity in histone methylation underlies
sperm programming for embryonic transcription. *Nat. Commun.* (2020)
doi:10.1038/s41467-020-17238-w.

11.   Siklenka, K. *et al.* Disruption of histone methylation in developing sperm
impairs offspring health transgenerationally. *Science (80-.).* (2015)
doi:10.1126/science.aab2006.

12.   Murphy, P. J., Wu, S. F., James, C. R., Wike, C. L. & Cairns, B. R. Placeholder
Nucleosomes Underlie Germline-to-Embryo DNA Methylation
Reprogramming. *Cell* (2018) doi:10.1016/j.cell.2018.01.022.

13.   Blackledge, N. P. *et al.* Variant PRC1 complex-dependent H2A ubiquitylation
drives PRC2 recruitment and polycomb domain formation. *Cell* **157**, (2014).

14.   Tamburri, S. *et al.* Histone H2AK119 Mono-Ubiquitination Is Essential for
Polycomb-Mediated Transcriptional Repression. *Mol. Cell* **77**, (2020).

15.   Nakagawa, T. *et al.* Deubiquitylation of histone H2A activates transcriptional
initiation via trans-histone cross-talk with H3K4 di- and trimethylation. *Genes*
Dev. (2008) doi:10.1101/gad.1609708.

16.   Joo, H. Y. *et al.* Regulation of cell cycle progression and gene expression by
H2A deubiquitination. *Nature* (2007) doi:10.1038/nature06256.

17.   Cao, J. & Yan, Q. Histone ubiquitination and deubiquitination in transcription,
DNA damage response, and cancer. *Frontiers in Oncology* (2012)
doi:10.3389/fonc.2012.00026.

18.   Chen, D., Dai, C. & Jiang, Y. Histone H2A and H2B Deubiquitinase in
Developmental Disease and Cancer. *Cancer Transl. Med.* (2015)
doi:10.4103/2395-3977.168578.

19.   Chen, Z., Djekidel, M. N. & Zhang, Y. Distinct dynamics and functions of
H2AK119ub1 and H3K27me3 in mouse preimplantation embryos. *Nat. Genet.*
53, (2021).

20.   Hickey, G. J. M. *et al.* Establishment of developmental gene silencing by
ordered polycomb complex recruitment in early zebrafish embryos. *Elife* **11**,
(2022).

21.   Conway, E. *et al.* BAP1 enhances Polycomb repression by counteracting
widespread H2AK119ub1 deposition and chromatin condensation. *Mol. Cell*
81, (2021).

22.   Albright, S. C., Nelson, P. P. & Garrard, W. T. Histone molar ratios among
different electrophoretic forms of mono- and dinucleosomes. *J. Biol. Chem.*
254, (1979).

23.   Matsui, S., Seon, B. K. & Sandberg, A. A. Disappearance of a structural
chromatin protein A24 in mitosis: Implications for molecular basis of

chromatin condensation. *Proc. Natl. Acad. Sci. U. S. A.* **76**, (1979).

24.   Zang, C., Wang, Y. & Peng, W. RECOGNICER: A coarse-graining approach
for identifying broad domains from ChIP-seq data. *Quant. Biol.* **8**, (2020).

25.   Heard, E. & Disteche, C. M. Dosage compensation in mammals: Fine-tuning
the expression of the X chromosome. *Genes and Development* vol. 20 (2006).

26.   Jullien, J. *et al.* Gene Resistance to Transcriptional Reprogramming following
Nuclear Transfer Is Directly Mediated by Multiple Chromatin-Repressive
Pathways. *Mol. Cell* **65**, (2017).

27.   Kuznetsov, J. N. *et al.* BAP1 regulates epigenetic switch from pluripotency to
differentiation in developmental lineages giving rise to BAP1-mutant cancers.
Sci. Adv. **5**, (2019).

28.   Jin, J. *et al.* The deubiquitinase USP21 maintains the stemness of mouse
embryonic stem cells via stabilization of Nanog. *Nat. Commun.* (2016)
doi:10.1038/ncomms13594.

29.   Gaggioli, V., Le Viet, B., Germe, T. & Hyrien, O. DNA topoisomerase IIα
controls replication origin cluster licensing and firing time in Xenopus egg
extracts. *Nucleic Acids Res.* **41**, (2013).

30.   Blackledge, N. P. & Klose, R. J. The molecular principles of gene regulation
by Polycomb repressive complexes. *Nature Reviews Molecular Cell Biology*
vol. 22 (2021).

31.   Zahn, N., Levin, M. & Adams, D. S. The zahn drawings: New illustrations of
xenopus embryo and tadpole stages for studies of craniofacial development.
Dev. **144**, (2017).

32.   Hörmanseder, E. *et al.* H3K4 Methylation-Dependent Memory of Somatic Cell
Identity Inhibits Reprogramming and Development of Nuclear Transfer
Embryos. *Cell Stem Cell* **21**, (2017).

33.   Session, A. M. *et al.* Genome evolution in the allotetraploid frog Xenopus
laevis. *Nature* **538**, (2016).

34.   Chen, H. & Good, M. C. Imaging nascent transcription in wholemount
vertebrate embryos to characterize zygotic genome activation. in *Methods in*
Enzymology vol. 638 (2020).

35.   Chen, H., Einstein, L. C., Little, S. C. & Good, M. C. Spatiotemporal
Patterning of Zygotic Genome Activation in a Model Vertebrate Embryo. *Dev.*
Cell **49**, (2019).

36.   Tsuboi, M. *et al.* Ubiquitination-Independent Repression of PRC1 Targets
during Neuronal Fate Restriction in the Developing Mouse Neocortex. *Dev.*
Cell **47**, (2018).

37.   Żylicz, J. J. *et al.* The Implication of Early Chromatin Changes in X
Chromosome Inactivation. *Cell* **176**, (2019).

38. Mei, H. *et al.* H2AK119ub1 guides maternal inheritance and zygotic deposition
of H3K27me3 in mouse embryos. *Nat. Genet.* **53**, (2021).

39. Van Heeringen, S. J. *et al.* Principles of nucleation of h3k27 methylation
during embryonic development. *Genome Res.* (2014)
doi:10.1101/gr.159608.113.

40. Akkers, R. C. *et al.* A Hierarchy of H3K4me3 and H3K27me3 Acquisition in
Spatial Gene Regulation in Xenopus Embryos. *Dev. Cell* (2009)
doi:10.1016/j.devcel.2009.08.005.

41. Zhao, J. *et al.* RYBP/YAF2-PRC1 complexes and histone H1-dependent
chromatin compaction mediate propagation of H2AK119ub1 during cell
division. *Nat. Cell Biol.* **22**, (2020).

42. Lismer, A. *et al.* Histone H3 lysine 4 trimethylation in sperm is transmitted to
the embryo and associated with diet-induced phenotypes in the offspring. *Dev.*
Cell **56**, (2021).

43. Pepin, A. S., Lafleur, C., Lambrot, R., Dumeaux, V. & Kimmins, S. Sperm
histone H3 lysine 4 tri-methylation serves as a metabolic sensor of paternal
obesity and is associated with the inheritance of metabolic dysfunction. *Mol.*
Metab. **59**, (2022).

44. Teperek, M. *et al.* Sperm and spermatids contain different proteins and bind
distinct egg factors. *Int. J. Mol. Sci.* **15**, (2014).

45. Hannak, E. & Heald, R. Investigating mitotic spindle assembly and function in
vitro using Xenopus laevis egg extracts. *Nat. Protoc.* **1**, (2006).

46. Martin, M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet.journal* **17**, (2011).

47. Andrews, S. & others. FastQC: a quality control tool for high throughput
sequence data. 2010.
Https://Www.Bioinformatics.Babraham.Ac.Uk/Projects/Fastqc/ (2019).

48. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*
29, (2013).

49. Anders, S., Pyl, P. T. & Huber, W. HTSeq-A Python framework to work with
high-throughput sequencing data. *Bioinformatics* **31**, (2015).

50. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize
analysis results for multiple tools and samples in a single report. *Bioinformatics*
32, (2016).

51. Dessau, R. B. & Pipper, C. B. [''R"--project for statistical computing]. *Ugeskr.*
Laeger **170**, (2008).

52. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, (2014).

53. Raudvere, U. *et al.* G:Profiler: A web server for functional enrichment analysis

and conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, (2019).

54. Fisher, M. *et al.* Xenbase: Key features and resources of the Xenopus model
organism knowledgebase. *Genetics* **224**, (2023).

55. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-
Wheeler transform. *Bioinformatics* **25**, (2009).

56. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,
(2021).

57. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res.* **44**, (2016).

58. Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J. & Prins, P. Sambamba:
Fast processing of NGS alignment formats. *Bioinformatics* **31**, (2015).

59. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**,
(2008).

60. Wang, Q. *et al.* Exploring Epigenomic Datasets by ChIPseeker. *Curr. Protoc.*
2, (2022).

61. Zhu, L. J. *et al.* ChIPpeakAnno: A Bioconductor package to annotate ChIP-seq
and ChIP-chip data. *BMC Bioinformatics* **11**, (2010).

62. Zhu, L. J. Integrative analysis of ChIP-chip and ChIP-seq dataset. *Methods*
Mol. Biol. **1067**, (2013).

63. Gel, B. *et al.* RegioneR: An R/Bioconductor package for the association
analysis of genomic regions based on permutation tests. *Bioinformatics* **32**,
(2016).

64. Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics
Viewer (IGV): High-performance genomics data visualization and exploration.
Brief. Bioinform. **14**, (2013).

65. Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for
comparing genomic features. *Bioinformatics* **26**, (2010).

66. Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription
Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell
Identities. *Mol. Cell* **38**, (2010).

FIGURE 1

FIGURE 2

FIGURE 3