The robust, high-throughput, and temporally regulated roxCre and loxCre reporting

systems for genetic modifications *in vivo*

Mengyang Shi[1,6], Jie Li[1,6], Xiuxiu Liu[1], Kuo Liu[2], Lingjuan He[3], Wenjuan Pu[1], Wendong Weng[1],

Shaohua Zhang[1], Huan Zhao[1], Kathy O. Lui[4,*], Bin Zhou[1,2,5,7,*]

[1]CAS CEMCS-CUHK Joint Laboratories, New Cornerstone Science Laboratory, State Key

Laboratory of Cell Biology, CAS Center for Excellence in Molecular Cell Science, Shanghai

Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences, University of Chinese

Academy of Sciences, Shanghai, China

[2]Key Laboratory of Systems Health Science of Zhejiang Province, School of Life Science,

Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou,

China

[3]School of Life Sciences, Westlake University, Hangzhou, China

[4]Department of Chemical Pathology, Li Ka Shing Institute of Health Sciences, The Chinese

University of Hong Kong, Prince of Wales Hospital, Hong Kong, China

[5]School of Life Science and Technology, ShanghaiTech University, Shanghai, China

[6]These authors contributed equally

[7]Lead contact

[*]Correspondence: kathyolui@cuhk.edu.hk or zhoubin@sibs.ac.cn

Abstract

Cre-loxP technology, a cornerstone in fate mapping and *in vivo* gene function studies, faces

challenges in achieving precise and efficient conditional mutagenesis through inducible systems.

This study introduces two innovative genetic tools designed to overcome these limitations. The

first, roxCre, enables DreER-mediated Cre release, paving the way for intersectional genetic

manipulation that permits increased precision and efficiency. The second, loxCre, facilitates

conditional gene targeting by allowing any CreER line to induce Cre expression with

significantly enhanced efficiency. These tools incorporate a fluorescent reporter for genetic

lineage tracing, simultaneously revealing efficient gene knockout in cells marked by the reporter.

These strategies hold great potential for precise and efficient exploration of lineage-specific gene

functions, marking a significant advancement in genetic research methodologies.

Introduction

Genetic modification, such as gene knockout in a precise spatiotemporal manner forms the basis

for understanding cellular and molecular mechanisms in multiple biological processes[1, 2]. The

Cre-loxP recombination system is widely employed to knock out or over-express specific genes

in vivo[3]. Cre recombinase, derived from the bacteriophage P1 gene, targets short palindrome

DNA sequence loxP (34bp) and recombines two loxP-flanked sequences for deletion[4, 5]. To date,

most of the mouse genes have been successfully targeted by two loxP sites (floxed allele), which

could be recombined by cell-type specific Cre lines. Application of the Cre-loxP system in

mouse genetics revolutionizes gene functional analysis and significantly advances our

exploration of many developmental and pathophysiological processes at the molecular level[6, 7].

While constitutively active Cre is robust and efficient in recombination for gene deletion, Cre

lacks temporal control as the promoter driving Cre is active from the embryonic to the adult

stage. Therefore, a temporally active Cre is needed to avoid early mortality and realize the

ablation of an indispensable gene at a key developmental stage. To overcome the temporal

limitation of Cre, CreER, the fusion of Cre with a mutated estrogen receptor, is generated to

enable its activation only after tamoxifen (Tam) treatment, and the interaction of ER with Tam

induces nuclear translocation of CreER from the cytoplasm[8]. As a result, Tam treatment restricts

CreER activity at a specific time window and permits gene functional analysis at a higher

spatiotemporal resolution than Cre[9]. As the loxP-flanked sequences and their locations in

genome or chromatin influence its accessibility for recombination, the recombination efficiency

of CreER varies significantly among multiple floxed alleles, ranging from easy- to inert-to-

recombine alleles[10]. For example, CreER has high efficiency in targeting the easy-to-recombine

alleles, such as generic reporter *R26-tdT*[11], but exhibits low efficiency on some other reporters

such as *R26-Confetti*[12], or may not excise some floxed gene alleles that are inert or resistant for

targeting, leading to failure of gene deletion in some tdT$^+$ cells and causing false-positive tracing

fate (Figure 1A). In other words, tdT expression may not always necessarily denote gene deletion

in the labeled cells. The notion that the pattern of Cre-expression based on a reporter (e.g., *R26-*

tdT) represents the pattern of Cre-mediated recombination is less rigorous[13]. Therefore, it is not

precise to assume the successful deletion of codirectional-loxP-flanked sequences in the gene of

interest based on the deletion in another gene (or reporter) by the same CreER mouse line.

One way to increase the recombination efficiency of CreER is to treat mice with many times of

Tam, as the high dosage of Tam in theory could increase the chances of CreER-mediated

recombination. However, the increased dosage of Tam treatment is toxic and could lead to many

side effects on the phenotype, creating confounding effects on the study[14, 15]. To increase the

inducible recombination efficiency of floxed alleles, Lao et al. reported a mosaic mutant analysis

with spatial and temporal control of recombination (MASTR), in which initial FlpoER-mediated

recombination induces GFPCre expression, that subsequently recombines floxed alleles

effectively in GFP$^+$ cells[16]. Similarly, a Flp-induced mosaic analysis system with Cre or Tomato

(MASCOT) has been reported to express Cre and reporter simultaneously in a cell[17]. While

MASTR and MASCOT enable spatiotemporal labeling of mosaic mutant cells utilizing the

current resources of floxed mouse lines, it may not be suitable for effective gene knockout at a

tissue/population level due to low recombination efficiency initiated by Flp recombinase. In

order to improve the inducible recombination efficiency, Tian et al. have also reported a self-

cleaved inducible CreER (sCreER) that switches inducible CreER into a constitutively active Cre,

effectively recombining floxed allele for gene manipulation[10]. However, self-cleaved inducible

Cre mice have to be newly generated each time for targeting different promoters, which is time-

consuming and the system is not compatible with current resources largely based on CreER

mouse lines. A recent study has reported an iSuRe-Cre strategy to induce and report Cre-

dependent genetic modifications utilizing conventional CreER tools[18]. The reporter expression

reflects gene knockout in cells by iSuRe-Cre. However, two limitations hinder its widespread

applications. First, without any induction, the iSuRe-Cre transgene is leaky in some tissues, such

as heart and skeletal muscle. Second, the initial recombination for releasing the iSuRe-Cre allele

induced by CreER is far from efficient in some tissues compared with easy-to-recombine alleles

such as *R26-tdT*[18], limiting its usage for efficient gene knockout at a tissue level. Thus, a new

method is needed to ensure gene knockout in cells as effectively and efficiently as recombination

on easy-to-recombine alleles, thus allowing efficient gene deletion at a population level.

Conventional Cre-loxP system uses tissue-specific promoter to drive Cre, thus the precision of

genetic targeting solely depends on promoter activity. Now it is known that many promoters are

not as specific as previously reported[19–21]. The ectopic or unwanted Cre expression in other cell

types may result in unspecific cell labeling or gene deletion, leading to many confounding issues

or controversies in multiple fields of research[22–24]. In addition, some cell types cannot be clearly

distinguished by one marker from another, and thus could not be specifically and genetically

targeted by only one gene promoter-driven recombinase[25]. To circumvent this limit, dual

recombinases-mediated genetic targeting utilizes two promoters to independently drive Cre and

Dre[26]. Dre is another bacteriophage recombinase that targets rox sites and is orthogonal to the

Cre-loxP system[27]. While cells of interest could be labeled by specific reporters that are

responsive to dual recombinases, gene deletion requires a final readout on singular Cre

recombinase for targeting floxed gene alleles.

In this study, we developed two genetic strategies: roxCre and loxCre, in which the *rox-Stop-rox*

(RSR) and *loxP-Stop-loxP* cassettes are respectively inserted into the *Cre* coding region, such

that Cre transcription and translation are not properly carried out before the removal of the *Stop*

cassette. There is no obviously spontaneous leakiness by roxCre or loxCre. We found that

CreER-induced loxCre effectively deletes genes in reporter-labeled cells, ensuring efficient gene

knockout for the evaluation of lineage-specific gene function. In addition, DreER-induced

roxCre efficiently deletes genes in cells for both specific and efficient intersectional genetic

studies. We expect that these tools would overcome the issues of inefficient and nonspecific

genetic modifications, enhancing the ability to more precisely manipulate genes in a particular

cell lineage for a better understanding of gene functions in multiple biomedical fields.

Results

Design of roxCre for DreER-induced mCre expression

A sequential genetic approach has been recently reported to delete genes[28], which used Dre-

induced CreER expression by removing the *rox-Stop-rox* sequence ahead. As aforementioned,

the recombination carried out by the released CreER for excising floxed gene alleles may not be

as efficient as reporters (e.g., *R26-tdT*). A possible strategy to control Cre activity is to engineer a

rox-flanked *Stop* sequence ahead of its DNA (*R26-RSR-Cre*) that will be removed after Dre-rox

recombination (Figure 1B). However, such a strategy is not ideal, as the upstream transcriptional

Stop sequence might not completely prevent *Cre* transcription, leading to leakiness (Figure 1C).

We then inserted a longer *Stop* sequence or reversed the direction of the *Cre* DNA by generating

R26-RSR-Cre2 and *R26-R-reverseCre-R* lines, respectively. The whole-mount fluorescence

imaging results displayed heavy leakiness, which demonstrates that Cre was independently

activated for recombination without being crossed with a Dre mouse line (Figure 1D-G). To

enable efficient dual recombinases-mediated gene knockout, we need to design a strategy for

DreER-induced expression of a constitutively active Cre, and the system should not exhibit

leakiness.

We first generated roxCre, in which an RSR cassette was inserted into the coding region of *Cre*,

splitting *Cre* into N- and C-terminal segments (Figure 1H). After removal of RSR by Dre-rox

recombination, Cre is recombined containing one remaining rox site within its coding sequence,

hereafter termed modified Cre (mCre, Figure 1H). To ensure that mCre has the same

recombination efficiency as the conventional Cre, we inserted a rox sequence at 12 different sites

of *Cre* individually, most of which were located between helix domains (Figure 1I). As

insertions of additional amino acids of rox would change the original sequence of Cre,

potentially leading to reduced activity, we screened for their recombination efficiency by

transfecting cells with 12 versions of mCre (mCre1 to mCre12) and the responding GFP reporter

(Figure 1J). We found that mCre1, 2, 3, 4, 6, 7, 10, and 11 displayed a comparable efficiency as

the conventional Cre; whereas mCre5, 8, 9, and 12 exhibited impaired recombination efficiency

(Figure 1K, L). These data demonstrate that some of the mCre generated in our study are as

efficient as the conventional Cre *in vitro*.

Next, we sought to screen for the most robust version of mCre *in vivo*. We generated four roxCre

knock-in mice based on mCre1, 4, 7, and 10 as determined from the above *in vitro* screening

(Figure S1-S2). In these roxCre lines, RSR inserted into *Cre* was removed by Dre-rox

recombination resulting in the generation of mCre (Figure 1J). We then used hepatocyte- and

endothelial cell-specific promoters albumin (Alb) and VE-cadherin (Cdh5) to drive roxCre, and

generate *Alb-roxCre1-tdT*, *Cdh5-roxCre4-tdT*, *Alb-roxCre7-GFP*, and *Cdh5-roxCre10-GFP*

knock-in mice, respectively (Figure S1-S2). No spontaneous fluorescence signal was detected in

Alb-roxCre1-tdT, *Cdh5-roxCre4-tdT*, *Alb-roxCre7-GFP*, and *Cdh5-roxCre10-GFP* knock-in

mice (Figure S1-S2), demonstrating no reporter leakiness without Dre-rox recombination. Nor

did we detect leakiness as a result of any Cre activity when they were crossed with *R26-GFP*[29] or

R26-tdT[11] reporter (Figure S1-S2). These data demonstrate that roxCre mouse lines are

functionally efficient yet non-leaky.

DreER-induced mCre robustly recombines inert alleles

To more rigorously evaluate the recombination effectiveness among different versions of mCre,

we used an inducible DreER to temporally and specifically release mCre in hepatocytes or

endothelial cells. Generic reporter mouse lines *R26-GFP* and *R26-tdT* were used to indicate the

difference in recombination efficiency between roxCre1 and roxCre7 first. The fluorescence-

activated cell sorting (FACS) and immunostaining results showed no leakiness in *R26-*

DreER;Alb-roxCre1-tdT;R26-GFP and *R26-DreER;Alb-roxCre7-GFP;R26-tdT* (Figure S3C-F).

Nevertheless, both two versions of mCre efficiently labeled virtually all targeted cells (Figure

S3C-E, G) due to the easy-to-recombine characteristic of *R26-GFP* and *R26-tdT*.

To evaluate a strong mCre, we attempted to target one of the most inert alleles for recombination,

R26-Confetti[12], which is often used as reporters for clonal analysis due to its sparse labeling with

rare recombination events[10, 12]. According to the Cre-loxP recombination principle, Cre can

remove the sequence between two loxP sites that have the same transcription direction and turn

over the sequence in the middle of two loxP sites that have the opposite transcription direction.

As the recombination efficiency of CreER is limited, *R26-Confetti* can only be recombined into YFP, or nGFP (nuclear GFP), or mCFP (membrane CFP), or RFP (Figure S4A).

We crossed *R26-DreER;R26-Confetti* mice with *Alb-roxCre1-tdT* (group 1) and *Alb-roxCre7-GFP* (group 2) mice, and Tam-induced Dre-rox recombination yielded *Alb-mCre1-tdT* and *Alb-mCre7-GFP* mice, respectively (Figure 2A). In this system, the constitutively active mCre would completely remove the sequence between two loxP sites that have the same transcription direction and recombine *R26-Confetti* into two sets of reporter pairs: YFP-nGFP pair and mCFP-RFP pair, which could be detected in a single hepatocyte due to the poly-nuclear or polyploidy feature of hepatocytes (Figure 2A). In each pair, reporters can equally express, as mCre is strong enough to turn over the sequence in the middle of two loxP sites that have the opposite transcription direction (Figure S4B). Therefore, we used YFP/mCFP to detect mCre recombination efficiency and tdT/RFP or GFP to trace its endogenous reporter activity that was driven by Alb promoter in *Alb-mCre1-tdT* and *Alb-mCre7-GFP* mice, respectively (Figure 2A). Besides, *Alb-CreER[26];R26-Confetti* was used as a control under the same Tam treatment (group 3, Figure 2A).

In fact, DreER-rox recombination in hepatocytes is not 100%. Therefore, we quantified YFP and mCFP expression in hepatocytes with DreER-rox recombination to evaluate the strength of mCre activity. We found sparse YFP$^+$ or mCFP$^+$ hepatocytes ($0.39 \pm 0.04\%$) in *Alb-CreER;R26-Confetti* mice. However, $99.94 \pm 0.02\%$ of tdT$^+$ hepatocytes (DreER recombined) were YFP$^+$ or mCFP$^+$ in *R26-DreER;Alb-roxCre1-tdT;R26-Confetti* mice; and $76.83 \pm 3.20\%$ of GFP$^+$ hepatocytes (DreER recombined) were YFP$^+$ or mCFP$^+$ in *R26-DreER;Alb-roxCre7-GFP;R26-Confetti* mice (Figure 2B, C). Similarly, we found that $39.81 \pm 2.61\%$ of tdT$^+$ endothelial cells and $74.15 \pm 3.64\%$ of GFP$^+$ endothelial cells in small intestine were YFP$^+$ or mCFP$^+$ in *R26-DreER;Cdh5-roxCre4-tdT;R26-Confetti* and *R26-DreER;Cdh5-roxCre10-GFP;R26-Confetti* mice, respectively, compared with sparse labeling in *Cdh5-CreER;R26-Confetti* mice (Figure 2D-F). Similar results were also observed in other tissues and organs (Figure S5). Furthermore, *R26-DreER;Alb-roxCre1-tdT;R26-Confetti*, *R26-DreER;Alb-roxCre7-GFP;R26-Confetti*, *R26-DreER;Cdh5-roxCre4-tdT;R26-Confetti*, *and R26-DreER;Cdh5-roxCre10-GFP;R26-Confetti* were not leaky (Figure S4C). Together, these data demonstrate that inducible DreER controls

roxCre activation, and mCre1 is the most efficient recombinase assisting the recombination of
inert alleles *in vivo*. Hereafter, roxCre1 will be referred to as roxCre for brevity.

Dre-induced mCre realizing cell subpopulation-specific gene manipulation

To substantiate the precision and specificity of roxCre targeting cells, we employed
intersectional genetic targeting using dual recombinases. We generated the *Cyp2e1-DreER*
mouse line which labeled peri-central hepatocytes and also renal epithelial cells (Figure S6).
Crossing *Cyp2e1-DreER* with hepatocyte-specific *Alb-roxCre-tdT* mice achieved genetic
targeting of peri-central hepatocytes exclusively, circumventing issues associated with ectopic
targeting of renal cells by *Cyp2e1-DreER* line (Figure S6) or unwanted targeting of peri-portal
hepatocytes by *Alb-CreER* line[26]. Wnt-β-catenin plays an important role in liver development,
zonation, homeostasis, and diseases[30]. Therefore, we generated the *Alb-roxCre-tdT;Ctnnb1$^{flox/flox}$*
mouse line to examine if Dre-induced mCre could efficiently delete *Ctnnb1* gene that encodes β-
catenin (*Ctnnb1^{flox}*)[31].

We generated *Cyp2e1-DreER;Alb-roxCre-tdT;Ctnnb1$^{flox/flox}$* triple knock-in mice (mutant), in
which Tam-induced DreER-rox recombination yielded mCre that subsequently targeted floxed
Ctnnb1 allele (Figure 3A). We treated mutant mice and their littermate control *Cyp2e1-*
DreER;Alb-roxCre-tdT;Ctnnb1$^{flox/+}$ mice with Tam at 8 weeks of age, and sorted tdT$^+$
hepatocytes for analysis at 3 days after Tam treatment (Figure 3B). qRT-PCR analysis showed
significantly reduced *Ctnnb1* and *Glul* in tdT$^+$ hepatocytes of the mutant group, compared to that
of the control group (Figure 3C, D). We then collected liver samples at 3 days (D) and 4 weeks
(W) post-Tam for further analysis (Figure 3E). Immunostaining for tdT, β-catenin, GS, and E-
CAD on liver sections revealed similar levels of β-catenin and GS expression in the 3D mutant
liver compared to the control liver (Figure 3F). However, expression of β-catenin and GS almost
disappeared in the 4W mutant liver tdT$^+$ region compared with the control (Figure 3G),
suggesting efficient deletion of *Ctnnb1* that participates in regulating Wnt signaling downstream
gene target *Glul*, which encodes GS. Western blotting against β-catenin and qRT-PCR against
Ctnnb1 of sorted tdT$^+$ hepatocytes at 4W post-Tam revealed the significant *Ctnnb1* knock-out
(Figure 3H-K). Additionally, Wnt downstream target genes including *Glul*, *Axin2*, *Cyp1a2*,
Cyp2e1, *Oat*, *Tcf7*, *Lect2*, *Tbx3*, *Slc1a2*, and *Rhbg* were significantly reduced in the mutant

sorted tdT$^+$ hepatocytes than the control group (Figure 3L). Taken together, these data

demonstrate that DreER-induced mCre enables efficient intersectional genetic manipulation in

the cells of interest.

R26-loxCre-tdT is efficiently recombined by CreER

Having successfully constructed roxCre for Dre-induced mCre expression, we iterated the

system to enable mCre induction by CreER, a more widely available tool for biomedical research.

We replaced the rox sites of roxCre1 with loxP sites by inserting a *loxP-Stop-loxP* cassette into

Cre sequence at the same locus as roxCre1(Figure 4A). We then generated the *R26-loxCre-tdT*

mouse line, in which mCre and tdT would be expressed simultaneously after *Stop* removal,

allowing tdT as a surrogate marker for mCre detection in the same cell (Figure 4C). We then

designed two experiments to verify the baseline leakiness of Cre expression (Figure 4B) and to

examine if mCre/tdT could be expressed after *Stop* removal (Figure 4C), respectively, by *R26-*

loxCre-tdT;R26-tdT (Strategy 1) and by injecting *AAV8-Cre* virus into *R26-loxCre-tdT* mice

(Strategy 2). Immunostaining for tdT on tissue sections revealed no tdT expression in Strategy 1,

demonstrating minimal to negligible leakiness at the baseline. The robust tdT expression in the

liver in Strategy 2 showed mCre/tdT was allowed to express after *Stop* removal by AAV2/8-

hTBG-Cre (Figure 4D). The FACS results confirmed that no leakiness in *R26-loxCre-tdT*

(Figure S7A).

We then validate whether the new design can improve the labeling efficiency compared to the

conventional approach by crossing endothelial cell-specific *Cdh5-CreER* with the conventional

reporter *R26-tdT* (Strategy 3) and the new reporter *R26-loxCre-tdT* mice (strategy 4) under the

same protocol of Tam treatment (Figure 4E). Notably, immunostaining and FACS results

showed that *R26-loxCre-tdT* significantly limited the leakiness of adult *Cdh5-CreER* compared

to *R26-tdT* (Figure S7B). For the labeling efficiency comparison, immunostaining for tdT and

VE-Cad on tissue sections revealed no difference in the percentage of VE-Cad$^+$ endothelial cells

expressing tdT between strategies 3 and 4 (Figure 4F- G), suggesting that recombination of the

R26-loxCre-tdT allele by CreER was comparable to *R26-tdT*, which is an easy-to-recombine

reporter mouse line. Therefore, these data indicate that *R26-loxCre-tdT* can be an easily used

reporter, enabling efficient recombination to generate mCre/tdT in CreER-expressing cells

without spontaneous leakiness.

R26-loxCre-tdT enables CreER to recombine _R26-Confetti_ efficiently

We next examined whether mCre released from loxCre could also display high recombination

efficiency for the inert-to-recombine allele _R26-Confetti_. We generated the triple knock-in _Cdh5-_

CreER;R26-loxCre-tdT;R26-Confetti mouse line in which mCre was released for confetti

recombination upon Tam treatment, and _Cdh5-CreER;R26-Confetti_ was used as the control

(Figure 5A). Both mouse strains were treated with Tam at 8 weeks of age and their tissue

samples were collected for analysis at one week post-Tam (Figure 5B). As tdT expression is also

detected in _R26-loxCre-tdT_ after the first recombination by CreER, we only compared the YFP

and mCFP fluorescence signals of _R26-Confetti_ (Figure 5A). Wholemount-fluorescence imaging

showed markedly more YFP and mCFP signals in the retina of the _Cdh5-CreER;R26-loxCre-_

tdT;R26-Confetti mice, compared to _Cdh5-CreER;R26-Confetti_ mice (Figure 5C).

Immunostaining on frozen sections revealed a significantly higher percentage of endothelial cells

expressing YFP/mCFP in various vascularized organs of the _Cdh5-CreER;R26-loxCre-tdT;R26-_

Confetti mice, compared to _Cdh5-CreER;R26-Confetti_ mice (Figure 5D). Notably, the results

showed that tdT$^+$ endothelial cells highly simultaneously expressed YFP/mCFP in _Cdh5-_

CreER;R26-loxCre-tdT;R26-Confetti mice (Figure 5D). Thus, these data demonstrate that the

R26-loxCre-tdT line could be used as an adaptor to enhance CreER-mediated recombination with

a simultaneous expression of tdT as an accurate indicator.

R26-loxCre-tdT adaptor ensures efficient recombination

To evaluate the specificity and efficiency of the loxCre strategy, we generated the _Alb-_

CreER;R26-loxCre-tdT;R26-Confetti mouse line, in which _Alb-CreER_ recombines _R26-loxCre-_

tdT allele in hepatocytes upon Tam treatment to release mCre/tdT and mCre is expected to

subsequently recombine the _R26-Confetti_ allele (Figure S7C, the right panel). As a control, we

crossed the _Alb-CreER;R26-Confetti_ mouse with an alternative version of tdT reporter, _R26-tdT2_

mice[32], in which a longe _Stop_ cassette was inserted between two loxP sites to reduce the false-

positive tracing results (Figure S7C, the left panel). Moreover, we also crossed _Alb-CreER;R26-_

Confetti with _iSuRe-Cre_ (_R26-loxP-NphiM-Stop-loxP-tdT-Int-Cre_)[18], in which the first

recombination releases tdT and Cre expression, which is expected to subsequently recombine

R26-Confetti (Figure S7C, the middle panel). The *iSuRe-Cre* is a published and widely utilized

inducible reporter-Cre mouse line, which has been reported to improve recombination activity in

tdT$^+$ cells[18].

Since most hepatocytes are multinucleated and polyploidy, a single hepatocyte would carry

multiple copies of the *R26-Confetti* alleles, yielding two or more fluorescence reporters.

Therefore, we used tdT as the readout on *Alb-CreER* mediated recombination of these three

following reporters: *R26-tdT2*, *iSuRe-Cre*, and *R26-loxCre-tdT*; and used YFP/mCFP as readouts

on *R26-Confetti* (Figure S7E). We injected one dosage of Tam into *Alb-CreER;R26-tdT2;R26-*

Confetti, *Alb-CreER;iSuRe-Cre;R26-Confetti*, and *Alb-CreER;R26-loxCre-tdT;R26-Confetti*

mice at 7 weeks old and collected their livers for analysis at one week post-Tam (Figure S7D).

The fluorescence images presented that the percentage of hepatocytes expressing tdT was highest

in *R26-loxCre-tdT* (91.76 ± 1.77%), followed by *R26-tdT2* (35.24 ± 1.66%), and least in *iSuRe-*

Cre (1.40 ± 0.13%) (Figure S7E-F), indicating the orders of recombination efficiency for these

alleles were *R26-loxCre-tdT* > *R26-tdT2* > *iSuRe-Cre*. Furthermore, we compared the

recombination efficiency on *R26-Confetti* allele and found that the percentage of hepatocytes

expressing YFP/mCFP was 1.08 ± 0.23% by *Alb-CreER;R26-tdT2*, 19.42± 3.71% by *Alb-*

CreER;iSuRe-Cre; and 100.00 ± 0.00% by *Alb-CreER;R26-loxCre-tdT* (Figure S7G). Together,

these findings collectively indicate that the mCre, when released from the *R26-loxCre-tdT*

construct, exhibits a significantly enhanced recombination efficiency. This improvement is

notable when compared to both the conventional CreER and previously reported *iSuRe-Cre*[18].

R26-loxCre-tdT ensures gene deletion in tdT$^+$ cells

To investigate whether *R26-loxCre-tdT* enables *Alb-CreER* to more efficiently delete genes, we

crossed *Alb-CreER;R26-loxCre-tdT* with *Ctnnb1$^{flox/flox}$* to generate *Alb-CreER;R26-loxCre-*

tdT;Ctnnb1$^{flox/flox}$ mice for evaluating the efficiency of releasing mCre and also mCre-mediated

Ctnnb1 deletion in tdT$^+$ hepatocytes (Figure 6A). We also crossed *Alb-CreER;R26-tdT2* with

Ctnnb1$^{flox/flox}$ to generate *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$* mice to evaluate recombination

efficiency of both *R26-tdT2* and *Ctnnb1$^{flox/flox}$* alleles (Figure 6A). Additionally, *Alb-CreER;R26-*

loxCre-tdT;Ctnnb1$^{flox/+}$ mice were used as a control for heterozygous gene deletion (Figure 6B).

Firstly, the FACS and immunostaining data showed no leakiness in *Alb-CreER;R26-loxCre-tdT*

(Figure S8A). Based on the no leaky background, we treated *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$*

and *Alb-CreER;R26-loxCre-tdT*;*Ctnnb1$^{flox/+}$* mice with five dosages of Tam, and treated *Alb-*

CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$ mice only with one dosage of Tam, and collected

hepatocytes for analyzing *Ctnnb1* gene deletion at 4 weeks after Tam treatment (Figure 6B).

Immunostaining data showed that β-catenin, GS, and E-CAD were significantly reduced in *Alb-*

CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$ mice but were readily detectable in both *Alb-CreER;R26-*

tdT2;Ctnnb1$^{flox/flox}$ and *Alb-CreER;R26-loxCre-tdT*;*Ctnnb1$^{flox/+}$* mice (Figure 6C). Western

blotting of β-catenin and qRT-PCR of *Ctnnb1* of isolated hepatocytes revealed the significant

deletion of *Ctnnb1* in the *Alb-CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$* mice compared with those in

control groups (Figure 6D-F). Additionally, Wnt downstream or regulated genes, including *Glul*,

Axin2, *Cyp1a2*, *Cyp2e1*, *Oat*, *Tcf7*, *Lect2*, *Tbx3*, *Slc1a2*, and *Rhbg,* were significantly reduced in

hepatocytes derived from the *Alb-CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$* mice, compared with

those from the *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$* and *Alb-CreER;R26-loxCre-tdT*;*Ctnnb1$^{flox/+}$*

mice, respectively (Figure 6G).

We next elucidate whether the significant superiority of *Alb-CreER;R26-loxCre-*

tdT;Ctnnb1$^{flox/flox}$ in the above data is attributed to low Tam dosage, which may reduce Tam

toxicity and side effects. Initially, we treated *Alb-CreER;R26-tdT;Ctnnb1$^{flox/flox}$* mice with one

dosage of Tam and subsequently sorted out tdT$^+$ hepatocytes for qRT-PCR against *Ctnnb1* 4

weeks after Tam treatment. The results indicated that *Alb-CreER* with a single dose of Tam,

without the assistance of *R26-loxCre-tdT*, was insufficient to knock out *Ctnnb1* (Figure S8B-D).

Next, we conducted an experiment involving a proportional gradient of the Tam dosages, in

which we gave five dosages, one dosage, one-fifth of one dosage, one-twenty-fifth of one dosage,

and one-hundred-twenty-fifth of one dosage of Tam to *Alb-CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$*

mice, respectively. We analyzed samples 3 weeks after Tam's treatment. The FACS results

showed that five groups of the proportional gradient of the Tam dosages labeled a gradient of

tdT$^+$ cells: 94.33±0.76%, 90.14±0.41%, 57.07±0.51%, 27.30±2.35%, and 1.27±0.54% (Figure

S8E), respectively. We sorted out tdT$^+$ hepatocytes from each group for qRT-PCR against

Ctnnb1. Our finding revealed that varying doses of Tam treatment primarily affect the target cell

ratio but do not influence the high recombination efficiency of mCre at loxP sites within the

Ctnnb1 allele (Figure S8E-F). Additionally, the immunostaining analysis revealed a noticeable

trend, in which the proportion of GS$^+$ or E-CAD$^+$ hepatocytes was in decreased tendency with

the increase of target tdT$^+$ hepatocytes (Figure S8F-G). This observation can be attributed to the

significant knockout of *Ctnnb1* in tdT$^+$ cells (Figure S8E-F).

Together, the combined results lead to the conclusion that *R26-loxCre-tdT* significantly enhances

the recombination efficiency of CreER, facilitating the targeting of genes that are challenging to

manipulate by traditional CreER-loxP recombination. Furthermore, the tdT component in *R26-*

loxCre-tdT enables precise tracing of the modified cell lineage, which accurately investigates the

roles of genes in various biological processes.

Discussion

In this study, we demonstrated that the *rox-Stop-rox* insert placed in front of *Cre* does not

effectively prevent the leakiness of Cre in the Rosa26 allele. To address this issue, we modified

Cre by inserting the *RSR* cassette into its coding region, thereby dividing Cre into its N- and C-

terminal segments. Our study showed that the developed roxCre not only eliminates leakage but

also enables the release of mCre for targeted gene manipulation in specific cell subpopulations

through the promotion of DreER. According to the construction of roxCre, we advanced to

develop *R26-loxCre-tdT*, which significantly enhances the efficiency of inducible Cre-mediated

gene manipulation, particularly when employing less effective CreER drivers for gene functional

exploration.

Assuming successful gene deletion in the cell type of interest depends on efficient cell labeling

by a lineage-specific reporter mediated by the same Cre is unrigorous, as recombination

efficiency varies significantly among different mouse lines. Our data clearly showed the

differences in recombination efficiency of different alleles (e.g. *R26-tdT*, *R26-tdT2*, and *R26-*

Confetti), even located in the same genomic locus (e.g. Rosa26) mediated by the same CreER

under the same Tam treatment protocol (Figure 6C, Figure S7E-G, and Figure S8D). However,

R26-loxCre-tdT can be efficient for recombination mediated by CreER and release mCre in

targeted cells, thus enabling efficient and accurate manipulating of cells of interest.

Compared to the previously reported transgenic mice line, *iSuRe-Cre*[18], *R26-loxCre-tdT* shows

more stable and robust enhancement in the efficiency of CreER recombination (Figure S7E-G,

Figure 6), avoiding the toxic effects resulting from high-dose of Tam. This advancement

significantly improves the reliability of genetic manipulation and cell fate lineage tracing,

making *R26-loxCre-tdT* a valuable tool for researchers investigating cellular functions and

behaviors. The research group of the iSuRe-Cre system recently reported a new system, named

iSuRe-HadCre, which demonstrated to be effective in deleting floxed genes while avoiding the

toxicity of constitutive Cre[33]. Further studies can be carried out to compare the recombination

enhancement between *R26-loxCre-tdT* and *iSuRe-HadCre*. To achieve robust and rapid Cre

expression while minimizing the toxic effects associated with constitutive Cre, further

enhancements are required for both loxCre and roxCre systems.

Wnt-β-catenin plays an important role in liver development, zonation, homeostasis, and

diseases[30]. The administration of five doses of Tam in *Alb-CreER* mice did not impact

downstream protein GS expression due to not knocking out *Ctnnb1* completely. However, a

single dose of Tam could effectively induce a significant knockout of *Ctnnb1* through *Alb-*

CreER;R26-loxCre-tdT (Figure 6C). This indicates that *R26-loxCre-tdT* can assist CreER in

addressing challenging recombined loxP sits. Notably, *R26-loxCre-tdT* can release mCre by

CreER-mediated activation and report tdT expression across all cells that express the

recombinase. Therefore, this system does not produce a gradient of reporter fluorescence

reflective of the promoter activity of CreER. Consequently, it's advisable not to use *R26-loxCre-*

tdT with a nonspecific and unstable CreER mouse line, which would amplify the CreER mouse

line's weakness and result in unanticipated nonspecific cell-targeting.

Furthermore, the strategy of roxCre further enhances the precision for cell fate mapping with

genetic deletion simultaneously in Dre^+Cre^+ cells through intersectional genetic targeting. For

example, we have showcased the strength of roxCre for genetic manipulation in peri-central

hepatocytes, resulting in efficient gene knockout in tdT^+ hepatocytes specifically (Figure 3).

Additional roxCre mouse lines driven by different promoters can serve as valuable tools for

detailed investigations into specific cell subpopulations. By using roxCre, studies can achieve

precise spatial and temporal control of gene expression, facilitating targeted analysis that

contributes to the knowledge of developmental biology, disease mechanisms, and potential

therapeutic interventions.

Previous studies have used Cre-loxP to simultaneously over-express gene and fluorescence

reporter in a cell for functional genetic mosaic (ifgMosaic) analysis[34]. However, ifgMosaic could

not take advantage of the abundant resources from existing available floxed mouse lines for loss-

of-function study. Mosaic mutant analysis with spatial and temporal control of recombination

(MASTR) utilized FlpoER for tissue-specific expression of GFPCre to delete floxed genes[16].

Compared with FlpoER lines, CreER is a more broadly used recombinase that most laboratories

use in gene deletion experiments. A recent study using dual recombinase-mediated cassette

exchange (MADR) also permits stable labeling of mutant cells expressing transgenic elements

from defined chromosomal loci[35]. However, the use of viruses and electroporation, albeit

convenient and rapid, are inefficient to specifically target any type of cells for *in vivo* studies.

The powerful mosaic analysis with double markers (MADM) enables simultaneous lineage

tracing of a pair of mutant and control sibling cells with distinct fluorescence reporters, allowing

precise mosaic analysis of gene function in any cell[36, 37]. Nevertheless, the MADM cassette has

to be combined with the mutant null allele, which is not readily available for synchronizing

reporter expression and genetic modification in cells of interest. The ensured gene deletion with

tdT reporter by *R26-loxCre-tdT* could be coupled with present CreER and loxP alleles for

potential mosaic analysis. This could be achieved by adjusting Tam at a lower dosage so that the

recombination efficiency to release mCre could be low in cells of interest for sparse tdT labeling.

We believe that *R26-loxCre-tdT* could be useful for robust, efficient, and temporal gene deletion

at a population level and mosaic analysis at a single-cell level with improvement in the future.

In conclusion, we developed two novel genetic tools, roxCre and loxCre, permitting efficient

recombination to release mCre, which greatly enhanced the effectiveness of subsequent gene

deletion in tdT[+] cells, thus facilitating further detailed characterization of these cells both *in situ*

and *ex vivo*. These tools are particularly beneficial when utilizing less effective DreER or CreER

drivers to better elucidate the roles of specific genes in biological processes, facilitating more

robust studies.

Materials and Methods

Mice

Experiments using mice (Mus musculus) were carried out with the study protocols (SIBCB-S374-1702-001-C1) approved by the Institutional Animal Care and Use Committee of Center for Excellence in Molecular Cell Science (CEMCS), Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences. The *R26-DreER*[38], *Alb-CreER*[26], *R26-GFP*[29], *R26-tdT*[11], *R26-tdT2*[32], *R26-RSR-tdT*[39], *R26-Confetti*[12], *iSuRe-Cre*[18] and *Ctnnb1-flox*[31] mouse lines were used as previously described. New knock-in mouse lines *R26-RSR-Cre*, *R26-RSR-Cre2*, *R26-R-reverseCre-R*, *Cdh5-CreER*, *Alb-roxCre1-tdT*, *Alb-roxCre7-GFP*, *Cdh5-roxCre4-tdT*, *Cdh5-roxCre10-GFP*, *Cyp2e1-DreER*, and *R26-loxCre-tdT* were generated by homologous recombination using CRISPR/Cas9 technology. These new mouse lines were generated by the Shanghai Model Organisms Center, Inc. (SMOC). These mice were bred in a C57BL6/ICR mixed background. All mice were housed at the laboratory Animal center of the Center for Excellence in Molecular Cell Science in a Specific Pathogen Free facility with individually ventilated cages. The room has controlled temperature (20–25 °C), humidity (30–70%), and light (12 hours light-dark cycle). For the determination of the embryonic period of sampling, the day on which the vaginal plug was examined in female mice was recorded as E0.5. As the sex is not relevant to the topic of this study, male and female mice ranging in age from E13.5 to 12W (week) were allocated and mixed into the experimental groups in this study. No data in mouse experiments were excluded.

Genomic PCR

Genomic DNA was prepared from the mouse toes or embryonic tails. Tissues were precipitated by centrifugation at maximum speed for 1 min at room temperature. After that, tissues were lysed by lysis buffer (100 mM Tris-HCl, 5 mM EDTA, 0.2% SDS, 200 mM NaCl, and 100 μg/mL Proteinase K) at 55 °C overnight. About 750μL pure ethanol was added to the lysis mixture and mixed thoroughly, followed by centrifugation at maximum speed for 5 min at room temperature to collect the DNA precipitation. Then the supernatant was discarded and the mixture was dried at 55 °C for 1 hour. About 250μL double-distilled H_2O was added to dissolve the DNA. The genomic PCR primer pairs were designed for the mutant alleles spanning both

endogenous genomic fragments and insert fragments. The genetic constructs and genotyping of

the new knock-in lines can be reviewed in Figure S9.

***In vitro* screening of mCre for efficient recombination**

To test if the remaining rox sequence after Dre-rox recombination would impact Cre activity, the

pCDNA3.1[40] (Invitrogen, V79020) was used to express 12 types of modified Cre (mCre). For

testing the Cre activity, 500ng pCDNA3.1-mCre plasmids, or pCDNA3.1 (negative control), or

pHR-CMV-nlsCRE (positive control, Addgene, 12265) were mixed with 500ng pCAG-loxp-

stop-loxp-ZsGreen (Addgene, 51269) plasmid. *In-vitro* transfection experiment protocol is

performed according to that described previously[41]. For cell culture medium, DMEM

(ThermoFisher, 11965092) was supplemented with 10% fetal bovine serum (Gibco, 10099141c)

for preparing fresh complete culture medium. Poly-D-Lysine (ThermoFisher, A3890401)

precoated coverslips (Biosharp, BS-14-RC) or 10 cm dish (ThermoFisher, 150466) overnight to

dry, and sterile water was used to flush it for three times. A cryogenic vial of QBI293 was put in

a 37°C water bath, then thawed liquid contents into a 50 mL conical tube (Corning,430829)

prefilled with 5 mL prewarmed fresh complete culture medium. After centrifuging at $125 \times g$ for

5 min, the supernatant was discarded and cells were resuspended in 10 mL complete medium.

One-third of the cells were plated in a 10 cm coated dish and incubated in cultures at 37°C. On

the second day, the culture medium was discarded, and the cell layer was briefly rinsed with

prewarmed PBS (Gibco, C10010500BT). After that, PBS was removed and 1 mL 0.25% Trypsin

solution (with EDTA, Gibco, 25200072) was added. The cells were incubated at 37°C for 3 min.,

and a 6mL complete growth medium was added to aspirate cells by gently pipetting. The

medium-containing cells were transferred to a 50 mL conical tube and centrifuged at $125 \times g$ for

5 min. The supernatant was discarded, and cells were resuspended in 5.2 mL complete culture

medium. 24 well plates were placed into coated coverslips and added 1mL complete culture

medium. 200µL cells were added to every experimental well and incubated cultures at 37°C for

9 hours to grow up to ~80%. The new prewarmed fresh complete culture medium replaced the

old for 1mL per well. Lipofectamine™ 3000 Transfection Reagent (Thermo Fisher, L3000015)

was used for plasmid transfection. Samples were incubated in cultures at 37°C for 24 hours.

Samples were washed by PBS once and mounted on slides by VECTASHIELD®· Antifade

Mounting Medium with DAPI (Vector, H-1200). Images were acquired using an Olympus microscope (Olympus, BX53).

Tamoxifen injection and sample collection

For induction of CreER or DreER, tamoxifen (Sigma, T5648) was dissolved in corn oil and was administered to mice by oral gavage. *Cyp2e1-DreER;Alb-roxCre1-tdT;Ctnnb1^{flox/+}* and *Cyp2e1-DreER;Alb-roxCre1-tdT;Ctnnb1^{flox/flox}* mice were treated with 0.05 mg Tam per gram mouse body weight (0.05 mg/G) for one dose. For comparing tdT$^+$ cells, *Cyp2e1-DreER;R26-RSR-tdT* and *Cyp2e1-DreER;Alb-roxCre1-tdT* mice were treated with 0.05 mg/G Tam. *R26-DreER;Alb-roxCre1-tdT;R26-Confetti*, *R26-DreER;Cdh5-roxCre4-tdT;R26-Confetti*, *R26-DreER;Alb-roxCre7-GFP;R26-Confetti*, *R26-DreER;Cdh5-roxCre10-GFP;R26-Confetti*, *R26-DreER; R26-DreER;Alb-roxCre7-GFP;R26-GFP*, *Alb-CreER;R26-Confetti*, *Cdh5-CreER;R26-Confetti*, *Alb-CreER;R26-tdT2;R26-Confetti*, *Alb-CreER;iSuRe-Cre;R26-Confetti*, *Alb-CreER;R26-loxCre-tdT;R26-Confetti*, *Alb-CreER;R26-loxCre-tdT;Ctnnb1^{flox/flox}*, and *Alb-CreER;R26-tdT;Ctnnb1^{flox/flox}* mice were treated with 0.2 mg/G Tam for one dose. *Cdh5-CreER;R26-tdT*, *Cdh5-CreER;R26-loxCre-tdT*, *Cdh5-CreER;R26-Confetti*, *Cdh5-CreER;R26-loxCre-tdT;R26-Confetti*, *Alb-CreER;R26-tdT2;Ctnnb1^{flox/flox}*, and *Alb-CreER;R26-loxCre-tdT;Ctnnb1^{flox/+}* mice were treated with 0.2 mg/G Tam one dose per day for 5 days. *R26-RSR-Cre2;R26-GFP* and *R26-R-reverseCre-R;R26-tdT* mice were sacrificed at 8 weeks old for analysis. For the proportional gradient of the Tam dosages, five dosages group was given 0.2 mg/G Tam for constitutive 5 days, one-fifth of one dosage group was given 0.04 mg/G Tam for one dose, one-twenty-fifth of one dosage group was given 0.008 mg/G Tam for one dose, and one-hundred-twenty-fifth of one dosage group was given 0.0016 mg/G Tam for one dose. Mice, both males and females, at the age of 8–12 weeks were used for experiments with similar-aged mice for both control and experimental groups.

Whole-mount imaging and sectioning

The tissue samples were fixed with 4% paraformaldehyde (PFA) (Sigma, P6148-500g, wt/vol in PBS) for 1 or 2 hours (stomach, intestine, colon) at 4°C, followed by washing with PBS three times. The fixed tissues were placed in an agarose-filled petri dish for bright-field and fluorescence imaging by a Zeiss stereoscopic microscope (AxioZoom V16). For cryo-sections,

544   tissues were sectioned to slides of 10-μm thickness after dehydration by 30% sucrose

545   (Sinopharm, H-10021463, wt/vol in PBS) overnight and pre-embedding with OCT (Scigen, 4586)

546   at 4 °C for 1 hour.

Immunostaining

Immunostaining was performed as previously described[42]. 0.2% PBST was prepared with 0.2%

(vol/vol) Triton X-100(Sigma, T9284) dissolved in PBS. Tissue sections were blocked with 2.5%

normal donkey serum and 0.1% 4'6-diamidino-2-phenylindole (DAPI, Invitrogen, D21490)

dissolved in 0.2% PBST for 30 mins after washing with PBS three times. The tissue sections

were incubated with primary antibody diluted in 0.2% PBST at 4°C overnight. On the next day,

sections were incubated with secondary antibodies diluted in 0.2% PBST at room temperature

for 30 min, followed by PBS washing for three times. The slides were washed with PBS for three

times. The slides were mounted with a mounting medium (Vector Lab). For weak signals, the

endogenous peroxidase activity was quenched before blocking. Horseradish peroxidase or biotin-

conjugated secondary antibodies and a Tyramide signal amplification kit (PerkinElmer) were

used after incubating the primary antibodies. For primary antibodies of murine origin, mouse

immunoglobulins were blocked with an anti-mouse Fab antibody (Jackson, 715-007-003, 1:100).

The included primary antibodies are listed as follows: tdT (Rockland, 600-401-379, 1:500; or

Rockland, 200-101-379, 1:500), GFP (Invitrogen, A11122, 1:500), GFP (Rockland, 600-101-

215M, 1:500), GFP (Nacalai, 04404-84; 1:500), GS (Abcam, ab49873, 1:1000), β-catenin (BD

Pharmingen, 610153, 1:200), E-CAD (R&D, AF748, 1:500), Ep-CAM (Abcam, ab92382, 1:500),

VE-Cad (R&D, AF1002, 1:100), and CYP2E1(Abcam, ab28146, 1:100). The corresponding

secondary antibodies (JIR or Abcam) were diluted according to the instructions. Images were

captured by using a Nikon confocal (Nikon A1 FLIM) or an Olympus confocal (FV3000), and

captured images were analyzed by Image J2 (version 2.9.0/1.54f) and Photoline (version 23.02)

software. We collected five random fields from each liver section for quantification. Mutant and

control sections were processed at the same time to avoid potential batch differences during

staining. Imaging of all immunostained slides was performed under the same exposure and

contrast conditions using the same confocal microscope.

Imaging of Confetti fluorescence reporters

All the fluorescence images of *R26-confetti* were taken by a Leica confocal (Leica SP8 will). All

the sections were collected freshly and blocked with 0.2% PBST containing DAPI for 30 mins

after washing with PBS three times. The parameters of excitement light and emission light for all

channels are set as follows: DAPI (ex.405nm, em.415-450nm), CFP (ex.458nm, em.463-481nm),

GFP (ex.488nm, em.495-511nm), YFP (ex.520nm, em.527-545nm), tdT (ex.546nm, em.555-

590nm), markers for antibody staining (ex.647nm, em.657-700nm). Captured images were

analyzed by Image J2 (version 2.9.0/1.54f) and Photoline (version 23.02) software.

Hepatocyte dissociation

Mouse primary hepatocytes were isolated by the two-step collagenase perfusion method, which

was modified from a previous protocol[43]. Briefly, mice were anesthetized and the liver was

exposed through an incision in the lower abdomen. The inferior vena cava and portal vein were

also exposed. A needle was inserted into the inferior vena cava and secured with a hemostatic

clamp around the needle. The portal vein was cut immediately when the mouse liver was

perfused with a perfusion medium buffer (containing 0.5 mM EGTA) for 5 minutes using a

peristaltic pump. Then, the liver was perfused with medium containing collagenase type I (150

U/ml; Gibco, 17100-017) for 2-5 min to adequately digest the liver. After the gallbladder was

removed, the liver was dissected with cold DMEM to free the hepatic cells. Then the cell

suspension was passed through a 70-μm cell strainer (BD Biosciences, 352350) and centrifuged

at 50 x g for 3 min at 4 °C. The supernatant was removed, and cells were resuspended in Percoll

(GE Healthcare, 17-0891-01)/DMEM/10 x PBS (Sangon, B548117, diluted in 1:1) (1:1:0.1)

mixture and centrifuged at 300 x g for 5 min at 4 °C. After the supernatant was removed, cells

were dissected with cold DMEM and centrifuged at 50 x g for 3 min at 4 °C. Purified

hepatocytes were collected for FACS analysis, qRT-PCR or Western Blot analysis.

Flow cytometric analysis and isolation of tdT[+] hepatocytes

Cells were centrifuged at 50 x g for 3 min at 4 °C and dissected with relevant suspended solution.

The suspended solution is 0.1% Dnase I (Worthington, LS002139, diluted by DMEM) mixed

with 0.1% DAPI. tdT[+] hepatocytes were sorted by FACS Aria SORP machine and Sony MA900.

Hepatocytes were collected by DMEM and then centrifuged at 50 x g for 3 min at 4 °C.

Hepatic Non-parenchymal cell dissociation

The liver was diced into 1mm pieces and then transferred into 10 mL digestion buffer (0.05g/100 mL collagenase type IV (Worthington, LS004188), 2mg/mL collagenase type I, 5u/mL Dispase (Corning, 354235), and 1%Dnase I in PBS). The sample was incubated in a 37°C incubator shaker at 220 rotations per minute for 30 min, with mixing performed three times during the incubation. Following this, 0.5 mL FBS was added to the digestion mixture. The resulting cell suspension was passed through a 40-μm cell strainer (BD Biosciences, 352340), and transferred to a new tube containing 30 mL cold PBS after being centrifuged at 50 x g for 3 min at 4 °C. The supernatant was discarded after 700 x g for 5 min at 4 °C. To lyse erythrocytes, RBC lysis buffer was added, and the mixture was kept for at room temperature 5 min. The lysis was halted by adding 20 mL cold PBS, followed by another centrifugation at 700 x g for 5 min at 4 °C. The supernatant was discarded, and the pellet was resuspended in the suspended solution.

Total RNA extraction and qRT-PCR

Total RNA was extracted from hepatocytes isolated from indicated mice as previously described[44]. Cells were lysed with Trizol (Invitrogen, 15596018), and total RNA was extracted according to the manufacturer's instructions. For each sample, 1 μg of total RNA was reversely transcribed into cDNA using the Prime Script RT kit (Takara, RR047A). The SYBR Green qPCR master mix (Thermo Fisher Scientific, 4367659) was used and quantitative RT-PCR was performed on QuantStudio 6 Real-Time PCR System (Thermo Fisher Scientific). *Gapdh* was used as the internal control. Sequences of all primers would be provided upon request.

Western blot

All samples were lysed in RIPA lysis buffer (Beyotime, P0013B) containing protease inhibitors (Roche, 11836153001) for 30 min on ice, and then centrifuged at 15,000 x g for 15 min to collect the supernatant. All samples were separated with 30μL for testing the protein concentration by Pierce[TM] BCA Protein Assay kits (Thermo Scientific[TM],23227). The remaining samples were mixed with 5 x loading buffer (Beyotime, p0015L) and boiled at 100 °C for 5 min. 40μg samples were added to every well in precast gradient gels (Beyotime, P0469M) with 1x running buffer (Epizyme, PS105S, should be diluted into 1x). After running, samples were transferred onto Immobilon® PVDF membranes (Millipore, IPVH00010). After blocking in blocking buffer (Epizyme, PS108P), the membranes were incubated with primary antibodies (diluted by primary

antibody dilution buffer (Epzyme, PS114)) overnight at 4 °C, then washed for three times and

incubated with HRP-conjugated secondary antibodies (diluted by 1xTBST (Epzyme, PS103S,

should be diluted into 1x)). Samples were incubated with chemiluminescent HRP substrate

(Millipore, WBKLS0500), and related signals were detected by MiniChemi 610 Plus (Biogp).

The following antibodies were used: β-catenin (BD Biosciences, 610153, 1: 5000), GAPDH

(Proteintech, 60004-1-IG, 1:2000), β-actin (Epzyme, LF202,1:1000), HRP-donkey-a-mouse (JIR,

715-035-150, 1:5000), and HRP-goat a rabbit IgG (JIR, 111-035-047, 1:5000).

AAV injection

The AAVs used in this study were purchased from Taitool Biotechnologies company (Shanghai,

China). Briefly, AAVs were produced with cis-plasmids containing the full TBG promoter,

which is specifically active in hepatocytes, and Cre expression is under the control of TBG

promoter, replication incompetent AAV2/8-TBG-Cre virus (AAV8-TBG-Cre, S0657-8-H5) was

packaged and purified before application to mice. Mice were injected intraperitoneally at $1x\ 10^{11}$

genome copies of the virus per mouse. The above newly generated virus as well as targeting

plasmids will be provided upon request.

Statistical analysis

Each pot in every graph represented one individual mouse. The quantification of each mouse

based on the fluorescence images was performed by counting the average data of five different

sections. Statistical analyses were performed using GraphPad Prism (Version 9.5.1). All the

continuous variables were expressed as means ± standard error of the mean (SEM). One-way

ANOVA was used to detect statistical significance between three experimental groups. The

statistical difference between the two experimental groups was determined using the unpaired

student's *t*-test. The probability (*P*) values of less than 0.05 were considered statistically

significant. The equation, Log(agonist) vs. response – Variable slope (four parameters) in Prism,

was adapted for the dose-response curve for the proportional gradient of the Tam dosages

experiment analysis, with a confidence interval of 95%.

Acknowledgments

This study was supported by the National Key Research & Development Program of China
(2024YFA1803302, 2023YFA1800700, 2022YFA1104200, 2023YFA1801300,
2020YFA0803202), National Natural Science Foundation of China (82088101, 32370897,
32100648, 32370783, 32100592), CAS Project for Young Scientists in Basic Research (YSBR-
012), Shanghai Pilot Program for Basic Research-CAS, Shanghai Branch (JCYJ-SHFY-2021-0),
Research Funds of Hangzhou Institute for Advanced Study (2022ZZ01015, B04006C01600515),
Shanghai Municipal Science and Technology Major Project, and the New Cornerstone Science
Foundation through the New Cornerstone Investigator Program and the XPLORER PRIZE, and
CAS-Croucher Funding Scheme for Joint Laboratories.
677

678 **Author contributions**
679 M.S., J.L., and B.Z. designed the study, analyzed the data, and wrote the manuscript; X.L., K.L.,
680 L.H., W.P., W.W., S.Z., and H.Z. bred the mice and performed experiments; K.O.L. edited the
681 manuscript and provided intellectual input; B.Z. conceived, supervised, and organized the study.
682

683 **Declaration of interests**
684 The authors declare no competing interests.
685

686 **Data availability**
687 All data included in this study are included in the manuscript. The raw data are available upon
688 request.

References

1. Gu, H., Zou, Y. R., and Rajewsky, K. (1993). Independent control of immunoglobulin switch recombination at individual switch regions evidenced through Cre-loxP-mediated gene targeting. Cell *73*, 1155-1164.

2. Gu, H., Marth, J. D., Orban, P. C., Mossmann, H., and Rajewsky, K. (1994). Deletion of a DNA polymerase beta gene segment in T cells using cell type-specific gene targeting. Science *265*, 103-106. https://doi.org/10.1126/science.8016642.

3. Nagy, A. (2000). Cre recombinase: the universal reagent for genome tailoring. Genesis *26*, 99-109.

4. Sternberg, N., Hamilton, D., Austin, S., Yarmolinsky, M., and Hoess, R. (1981). Site-specific recombination and its role in the life cycle of bacteriophage P1. Cold Spring Harb Symp Quant Biol *45 Pt 1*, 297-309.

5. Sauer, B., and Henderson, N. (1988). Site-specific DNA recombination in mammalian cells by the Cre recombinase of bacteriophage P1. Proc Natl Acad Sci U S A *85*, 5166-5170.

6. Orban, P. C., Chui, D., and Marth, J. D. (1992). Tissue- and site-specific DNA recombination in transgenic mice. Proc Natl Acad Sci U S A *89*, 6861-6865.

7. Lakso, M., Sauer, B., Mosinger, B. J., Lee, E. J., Manning, R. W., Yu, S. H., Mulder, K. L., and Westphal, H. (1992). Targeted oncogene activation by site-specific recombination in transgenic mice. Proc Natl Acad Sci U S A *89*, 6232-6236.

8. Feil, R., Wagner, J., Metzger, D., and Chambon, P. (1997). Regulation of Cre recombinase activity by mutated estrogen receptor ligand-binding domains. Biochem Biophys Res Commun *237*, 752-757. https://doi.org/10.1006/bbrc.1997.7124.

9. Tian, X., Pu, W. T., and Zhou, B. (2015). Cellular Origin and Developmental Program of Coronary Angiogenesis. Circ Res *116*, 515-530. https://doi.org/10.1161/CIRCRESAHA.116.305097.

10. Tian, X., He, L., Liu, K., Pu, W., Zhao, H., Li, Y., Liu, X., Tang, M., Sun, R., Fei, J., et al. (2020). Generation of a self-cleaved inducible Cre recombinase for efficient temporal genetic manipulation. EMBO J *39*, e102675. https://doi.org/10.15252/embj.2019102675.

11. Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter, R. D., Hawrylycz, M. J., Jones, A. R., et al. (2010). A robust and high-

throughput Cre reporting and characterization system for the whole mouse brain. Nat
Neurosci *13*, 133-140. https://doi.org/10.1038/nn.2467.

12. Snippert, H. J., van der Flier, L. G., Sato, T., van Es, J. H., van den Born, M., Kroon-
Veenboer, C., Barker, N., Klein, A. M., van Rheenen, J., Simons, B. D., et al. (2010).
Intestinal crypt homeostasis results from neutral competition between symmetrically
dividing Lgr5 stem cells. Cell *143*, 134-144. https://doi.org/10.1016/j.cell.2010.09.016.

13. Becher, B., Waisman, A., and Lu, L. F. (2018). Conditional Gene-Targeting in Mice:
Problems   and   Solutions.   Immunity   *48*,   835-836.
https://doi.org/10.1016/j.immuni.2018.05.002.

14. Brash, J. T., Bolton, R. L., Rashbrook, V. S., Denti, L., Kubota, Y., and Ruhrberg, C.
(2020). Tamoxifen-Activated CreERT Impairs Retinal Angiogenesis Independently of
Gene Deletion. Circ Res *127*, 849-850. https://doi.org/10.1161/CIRCRESAHA.120.317025.

15. Rashbrook, V. S., Brash, J. T., and Ruhrberg, C. (2022). Cre toxicity in mouse models of
cardiovascular   physiology   and   disease.   Nat   Cardiovasc   Res   *1*,   806-816.
https://doi.org/10.1038/s44161-022-00125-6.

16. Lao, Z., Raju, G. P., Bai, C. B., and Joyner, A. L. (2012). MASTR: a technique for mosaic
mutant analysis with spatial and temporal control of recombination using conditional
floxed alleles in mice. Cell Rep *2*, 386-396. https://doi.org/10.1016/j.celrep.2012.07.004.

17. Wang, Q., Lin, Y. Y., Zhang, B., Wu, J., Roy, S., Ratiu, J. J., Xu, Y., Dai, M., Hale, L. P.,
Xiong, Y., et al. (2020). A mosaic analysis system with Cre or Tomato expression in the
mouse.   Proc   Natl   Acad   Sci   U   S   A   *117*,   28212-28220.
https://doi.org/10.1073/pnas.2014308117.

18. Fernández-Chacón, M., Casquero-García, V., Luo, W., Francesca Lunella, F., Ferreira
Rocha, S., Del Olmo-Cabrera, S., and Benedito, R. (2019). iSuRe-Cre is a genetic tool to
reliably induce and report Cre-dependent genetic modifications. Nat Commun *10*, 2262.
https://doi.org/10.1038/s41467-019-10239-4.

19. Liu, Q., Yang, R., Huang, X., Zhang, H., He, L., Zhang, L., Tian, X., Nie, Y., Hu, S., Yan,
Y., et al. (2016). Genetic lineage tracing identifies in situ Kit-expressing cardiomyocytes.
Cell Res *26*, 119-130. https://doi.org/10.1038/cr.2015.143.

20.   Gribben, C., Sancho, R., and Behrens, A. (2023). Response to Magenheim et al.: Ductal
749   Ngn3-expressing progenitors contribute to adult beta cell neogenesis in the pancreas. Cell
750   Stem Cell *30*, 345-347. https://doi.org/10.1016/j.stem.2023.03.002.

751   21.   Furuyama, K., Kawaguchi, Y., Akiyama, H., Horiguchi, M., Kodama, S., Kuhara, T.,
752   Hosokawa, S., Elbahrawy, A., Soeda, T., Koizumi, M., et al. (2011). Continuous cell
753   supply from a Sox9-expressing progenitor zone in adult liver, exocrine pancreas and
754   intestine. Nat Genet *43*, 34-41. https://doi.org/10.1038/ng.722.

755   22.   Xiao, X., Chen, Z., Shiota, C., Prasadan, K., Guo, P., El-Gohary, Y., Paredes, J., Welsh, C.,
756   Wiersch, J., and Gittes, G. (2013). No evidence for beta cell neogenesis in murine adult
757   pancreas. J Clin Invest *123*, 2207-2217. https://doi.org/10.1172/JCI66323.

758   23.   Tarlow, B. D., Finegold, M. J., and Grompe, M. (2014). Clonal tracing of Sox9+ liver
759   progenitors   in   mouse   oval   cell   injury.   Hepatology   *60*,   278-289.
760   https://doi.org/10.1002/hep.27084.

761   24.   He, L., Nguyen, N. B., Ardehali, R., and Zhou, B. (2020). Heart Regeneration by
762   Endogenous Stem Cells and Cardiomyocyte Proliferation: Controversy, Fallacy, and
763   Progress.   Circulation   *142*,   275-291.
764   https://doi.org/10.1161/CIRCULATIONAHA.119.045566.

765   25.   Liu, Q., Liu, K., Cui, G., Huang, X., Yao, S., Guo, W., Qin, Z., Li, Y., Yang, R., Pu, W., et
766   al. (2019). Lung regeneration by multipotent stem cells residing at the bronchioalveolar-
767   duct junction. Nat Genet *51*, 728-738. https://doi.org/10.1038/s41588-019-0346-6.

768   26.   He, L., Li, Y., Li, Y., Pu, W., Huang, X., Tian, X., Wang, Y., Zhang, H., Liu, Q., Zhang, L.,
769   et al. (2017). Enhancing the precision of genetic lineage tracing using dual recombinases.
770   Nat Med *23*, 1488-1498. https://doi.org/10.1038/nm.4437.

771   27.   Anastassiadis, K., Fu, J., Patsch, C., Hu, S., Weidlich, S., Duerschke, K., Buchholz, F.,
772   Edenhofer, F., and Stewart, A. F. (2009). Dre recombinase, like Cre, is a highly efficient
773   site-specific recombinase in E. coli, mammalian cells and mice. Dis Model Mech *2*, 508-
774   515. https://doi.org/10.1242/dmm.003087.

775   28.   Han, M., Liu, Z., Liu, L., Huang, X., Wang, H., Pu, W., Wang, E., Liu, X., Li, Y., He, L.,
776   et al. (2023). Dual genetic tracing reveals a unique fibroblast subpopulation modulating
777   cardiac fibrosis. Nat Genet *55*, 665-678. https://doi.org/10.1038/s41588-023-01337-7.

778   29.   Zhang, H., Pu, W., Li, G., Huang, X., He, L., Tian, X., Liu, Q., Zhang, L., Wu, S. M.,
779   Sucov, H. M., et al. (2016). Endocardium Minimally Contributes to Coronary Endothelium
780   in the Embryonic Ventricular Free Walls. Circ Res *118*, 1880-1893.
781   https://doi.org/10.1161/CIRCRESAHA.116.308749.

782   30.   Perugorria, M. J., Olaizola, P., Labiano, I., Esparza-Baquer, A., Marzioni, M., Marin, J. J.
783   G., Bujanda, L., and Banales, J. M. (2019). Wnt-β-catenin signalling in liver development,
784   health and disease. Nat Rev Gastroenterol Hepatol *16*, 121-136.
785   https://doi.org/10.1038/s41575-018-0075-9.

786   31.   Huelsken, J., Vogel, R., Erdmann, B., Cotsarelis, G., and Birchmeier, W. (2001). beta-
787   Catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. Cell
788   *105*, 533-545.

789   32.   Liu, K., Tang, M., Jin, H., Liu, Q., He, L., Zhu, H., Liu, X., Han, X., Li, Y., Zhang, L., et al.
790   (2020). Triple-cell lineage tracing by a dual reporter on a single allele. J Biol Chem *295*,
791   690-700. https://doi.org/10.1074/jbc.RA119.011349.

792   33.   Garcia-Gonzalez, I., Rocha, S. F., Hamidi, A., Garcia-Ortega, L., Regano, A., Sanchez-
793   Muñoz, M. S., Lytvyn, M., Garcia-Cabero, A., Roig-Soucase, S., Schoofs, H., et al. (2024).
794   iSuRe-HadCre is an essential tool for effective conditional genetics. Nucleic Acids Res *52*,
795   e56. https://doi.org/10.1093/nar/gkae472.

796   34.   Pontes-Quero, S., Heredia, L., Casquero-García, V., Fernández-Chacón, M., Luo, W.,
797   Hermoso, A., Bansal, M., Garcia-Gonzalez, I., Sanchez-Muñoz, M. S., Perea, J. R., et al.
798   (2017). Dual ifgMosaic: A Versatile Method for Multispectral and Combinatorial Mosaic
799   Gene-Function Analysis. Cell *170*, 800-814.e18. https://doi.org/10.1016/j.cell.2017.07.031.

800   35.   Kim, G. B., Rincon Fernandez Pacheco, D., Saxon, D., Yang, A., Sabet, S., Dutra-Clarke,
801   M., Levy, R., Watkins, A., Park, H., Abbasi Akhtar, A., et al. (2019). Rapid Generation of
802   Somatic Mouse Mosaics with Locus-Specific, Stably Integrated Transgenic Elements. Cell
803   *179*, 251-267.e24. https://doi.org/10.1016/j.cell.2019.08.013.

804   36.   Zong, H., Espinosa, J. S., Su, H. H., Muzumdar, M. D., and Luo, L. (2005). Mosaic
805   analysis with double markers in mice. Cell *121*, 479-492.
806   https://doi.org/10.1016/j.cell.2005.02.012.

807   37.   Liu, C., Sage, J. C., Miller, M. R., Verhaak, R. G., Hippenmeyer, S., Vogel, H., Foreman,
808   O., Bronson, R. T., Nishiyama, A., Luo, L., et al. (2011). Mosaic analysis with double

markers reveals tumor cell of origin in glioma. Cell *146*, 209-221. https://doi.org/10.1016/j.cell.2011.06.014.

38. Li, Y., He, L., Huang, X., Issa Bhaloo, S., Zhao, H., Zhang, S., Pu, W., Tian, X., Li, Y., Liu, Q., et al. (2018). Genetic Lineage Tracing of Non-Myocyte Population by Dual Recombinases. Circulation *138*, 793-805. https://doi.org/10.1161/CIRCULATIONAHA.118.034250.

39. Zhang, H., Pu, W., Liu, Q., He, L., Huang, X., Tian, X., Zhang, L., Nie, Y., Hu, S., Lui, K. O., et al. (2016). Endocardium Contributes to Cardiac Fat. Circ Res *118*, 254-265. https://doi.org/10.1161/CIRCRESAHA.115.307202.

40. Paxinou, E., Chen, Q., Weisse, M., Giasson, B. I., Norris, E. H., Rueter, S. M., Trojanowski, J. Q., Lee, V. M., and Ischiropoulos, H. (2001). Induction of alpha-synuclein aggregation by intracellular nitrative insult. J Neurosci *21*, 8053-8061. https://doi.org/10.1523/JNEUROSCI.21-20-08053.2001.

41. Jiang, P., Ren, L., Zhi, L., Hu, X., and Xiao, R. P. (2021). Protocol for cell preparation and gene delivery in HEK293T and C2C12 cells. STAR Protoc *2*, 100497. https://doi.org/10.1016/j.xpro.2021.100497.

42. Pu, W., Zhu, H., Zhang, M., Pikiolek, M., Ercan, C., Li, J., Huang, X., Han, X., Zhang, Z., Lv, Z., et al. (2023). Bipotent transitional liver progenitor cells contribute to liver regeneration. Nat Genet *55*, 651-664. https://doi.org/10.1038/s41588-023-01335-9.

43. Charni-Natan, M., and Goldstein, I. (2020). Protocol for Primary Mouse Hepatocyte Isolation. STAR Protoc *1*, 100086. https://doi.org/10.1016/j.xpro.2020.100086.

44. Pu, W., Zhang, H., Huang, X., Tian, X., He, L., Wang, Y., Zhang, L., Liu, Q., Li, Y., Li, Y., et al. (2016). Mfsd2a+ hepatocytes repopulate the liver during injury and regeneration. Nat Commun *7*, 13369. https://doi.org/10.1038/ncomms13369.

Figure legends

Figure 1. Design of roxCre for DreER-induced mCre expression

(A) A schematic showing genetic labeling and/or gene knockout in CreER-expressing cells.

(B, D, F) Strategies for DreER-induced Cre expression.

(C) Crossing with *R26-tdT* mice, *R26-RSR-Cre* exhibits leakiness in E13.5 embryos. Scale bars, yellow 1mm. Each image is representative of 5 individual biological samples.

(E) Examination of Cre leakiness by whole-mount fluorescence images of organs collected from *R26-RSR-Cre2;R26-GFP* adult mice. Scale bars, yellow 1mm. Each image is representative of 5 individual biological samples.

(G) Examination of Cre leakiness by whole-mount fluorescence images of organs collected from *R26-R-reverseCre-R;R26-tdT* adult mice. Scale bars, yellow 1mm. Each image is representative of 5 individual biological samples.

(H) A schematic showing the design for roxCre and mCre.

(I) A schematic showing mCre1 to mCre12 by insertion of rox into *Cre* cDNA at 12 different loci.

(J) Examination of mCre for recombination efficiency by the *loxP-Stop-loxP-GFP* reporter.

(K) Immunofluorescence images of cells stained with GFP on cells. Scale bars, white, 100 µm. Each image is representative of 5 individual biological samples.

(L) Quantification of the percentage of cells expressing GFP in each group. Data are the means ± SEM; n = 5.

Figure 2. DreER-induced mCre robustly recombines inert alleles

(A) A schematic showing the experimental design to test the recombination efficiency of mCre1 and mCre7 on the *R26-Confetti* allele. Tam-induced DreER-rox recombination leads to mCre1/tdT or mCre7/GFP expression in hepatocytes in strategy 1 or 2, respectively. Strategy 3 uses conventional *Alb-CreER* as control. YFP and mCFP signals are used for the examination of recombination on the *R26-Confetti* allele.

(B) Fluorescence images of YFP and mCFP on liver sections collected from mice in strategy 1-3. Strategy 1 and 3 exhibit tdT or GFP in hepatocytes after DreER-rox recombination, respectively. Scale bars, 100 µm. Each image is representative of 5 individual biological samples.

(C) Quantification of the percentage of hepatocytes (Heps) expressing either YFP and/or mCFP

in strategy 4-6. Data are the means ± SEM; n = 11 mice for each strategy; ****P < 0.0001 by

one-way ANOVA.

(D) A schematic showing the experimental strategy to test the recombination efficiency of

mCre4 and mCre10 on *R26-Confetti* allele. Tam-induced recombination leads to mCre4/tdT or

mCre10/GFP expression in endothelial cells (ECs) in strategy 4 or 5, respectively. Strategy 6

uses conventional *Cdh5-CreER* as control. YFP and mCFP signals are used for the examination

of recombination on *R26-Confetti* allele.

(E) Fluorescence images of YFP and mCFP on intestinal sections collected from mice in strategy

4-6. Strategy 4 and 5 exhibit tdT or GFP in ECs after DreER-rox recombination, respectively.

Scale bars, 100 µm. Each image is representative of 5 individual biological samples.

(F) Quantification of the percentage of ECs expressing either YFP and/or mCFP in strategy 4-6.

Data are the means ± SEM; n = 11 mice for each strategy; ****P < 0.0001 by one-way ANOVA.

Figure 3. Dre-induced mCre efficiently deletes genes in specific cell subpopulation.

(A) A schematic showing the experimental design for Dre-induced mCre expression and the

subsequent gene deletion.

(B) A schematic showing the experimental strategy.

(C and D) qRT-PCR analysis of the relative expression of *Ctnnb1* (C) and *Glul* (D) in the sorted

tdT$^+$ hepatocytes. Data are the means ± SEM; n = 5; ****P < 0.0001 by student's *t*-test.

(E) A schematic showing the experimental strategy.

(F) Immunostaining for tdT, β-Catenin, GS, and E-CAD on liver sections collected on the day 3

post-Tam. PV, portal vein; CV, central vein. Scale bars, yellow 1mm; white 100 µm. Each image

is representative of 5 individual biological samples.

(G) Immunostaining for tdT, β-Catenin, GS, and E-CAD on liver sections collected at week 4

post-Tam. Scale bars, yellow 1mm; white 100 µm. Each image is representative of 5 individual

biological samples.

(H) A schematic showing the experimental strategy.

(I) Western blotting of β-Catenin and GAPDH in sorted tdT$^+$ cells.

(J) Quantification of the relative expression of β-Catenin protein. Data are the means ± SEM; n =

3. **P < 0.01 by student's *t* test.

(K) qRT-PCR analysis of the relative expression of *Ctnnb1* in sorted tdT$^+$ cells. Data are the

means ± SEM; n = 6. ****$P < 0.0001$ by student's *t* test.

(L) qRT-PCR analysis of the relative expression of *Glul*, *Axin2*, *Cyp1a2*, *Cyp2e1*, *Oat*, *Tcf7*,

Lect2, *Tbx3*, *Slc1a2,* and *Rhbg* in sorted tdT$^+$ cells. Data are the means ± SEM; n = 6. ***$P <$

0.001, ****$P < 0.0001$ by student's *t* test.

Figure 4. *R26-loxCre-tdT* is efficiently recombined by CreER

(A) A schematic showing the knock-in strategy for the generation of the *R26-loxCre-tdT* allele.

In this line, tdT is used to denote the mCre expression after the removal of *Stop*.

(B) A schematic showing experimental strategy 1 to examine the leakiness of *R26-loxCre-tdT*

mice.

(C) A schematic showing experimental strategy 2 to test mCre/tdT expression by AAV8-TBG-

Cre.

(D) Immunostaining for tdT on tissue sections shows tdT expression in the liver and pancreas

when recombination was initiated by AAV8-TBG-Cre. Scale bars, white 100 µm. Each image is

representative of 5 individual biological samples.

(E) A schematic showing the experimental strategies 3 and 4 for comparing the CreER-mediated

first recombination efficiency between *Cdh5-CreER;R26-tdT* and *Cdh5-CreER;R26-loxCre-tdT*

mice.

(F) Immunostaining for tdT and VE-Cad on liver sections collected on day 7 post-Tam. Scale

bars, white 100 µm. Each image is representative of 5 individual biological samples.

(G) Quantification of the percentage of VE-Cad$^+$ ECs expressing tdT. Data are the means ± SEM;

n= 6. ****$P < 0.0001$ by student's *t*-test.

Figure 5. *R26-loxCre-tdT* enables CreER to recombine *R26-Confetti* efficiently

(A) Schematics showing the experimental design. In *Cdh5-CreER;R26-loxCre-tdT;R26-Confetti*

mice, Tam-induced CreER-loxp recombination switches *R26-loxCre-tdT* into *R26-mCre-tdT*

allele with simultaneous tdT labeling and expression of mCre, which subsequently targets *R26-*

Confetti (right panel). The conventional *Cdh5-CreER;R26-Confetti* mice are used as control.

(B) A schematic showing the experimental strategy.

(C) Whole-mount YFP and mCFP fluorescence images of retina collected from two mice groups.

Scale bars, white 100 µm. Each image is representative of 5 individual biological samples.

(D) Immunofluorescence images of YFP, mCFP, and VE-Cad on tissue sections show

significantly more YFP$^+$ and/or mCFP$^+$ ECs in the *Cdh5-CreER;R26-loxCre-tdT;R26-Confetti*

mice compared with that of *Cdh5-CreER;R26-Confetti* mice (left panel). The right panel shows

the quantification of ECs expressing YFP and/or mCFP. Data are the means ± SEM; n = 5.

****$P < 0.0001$ by student's *t*-test. Scale bars, white 100 µm. Each image is representative of 5

individual biological samples.

Figure 6. *R26-loxCre-tdT* enables *Alb-CreER* to efficiently knockout genes in hepatocytes

(A) Schematics showing the experimental designs for *Ctnnb1* gene knockout using either *Alb-*

CreER or *Alb-CreER;R26-loxCre-tdT* mice.

(B) A schematic showing the experimental strategy. *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$* and *Alb-*

CreER;R26-loxCre-tdT;Ctnnb1$^{flox/+}$ mice were injected with Tam for 5 times, while *Alb-*

CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$ mice were injected with Tam for one time.

(C) Immunostaining for tdT, β-Catenin, GS, and E-CAD on liver sections. Scale bars, yellow

1mm; white 100 µm. Each image is representative of 5 individual biological samples.

(D) Western blotting of β-Catenin and β-Actin expression in hepatocytes, n = 3.

(E) Quantification of β-Catenin expression. Data are the means ± SEM; n = 5.

(F) qRT-PCR analysis of the relative expression of *Ctnnb1* in the hepatocytes.

(G) qRT-PCR analysis of the relative expression of *Glul*, *Axin2*, *Cyp2e1*, *Oat*, *Tcf7*, *Lect2*, *Tbx3*,

and *Slc1a2* in hepatocytes. Data are the means ± SEM; n = 7; ns, non-significant; *$P < 0.05$, **P

< 0.01, ***$P < 0.001$, ****$P < 0.0001$ by one-way ANOVA.

947
948
949

Figure 1

Figure2

Figure3

A *R26-loxCre-tdT* knock-in strategy

B Strategy 1

C Strategy 2

D Strategy 1 / Strategy 2

E Comparison of recombination efficiency

F Strategy 3 / Strategy 4

G Strategy 3 / Strategy 4

Figure4

Figure5

Figure6