The robust, high-throughput, and temporally regulated roxCre and loxCre reporting

systems for genetic modifications *in vivo*

Mengyang Shi[1,5], Jie Li[1,5], Xiuxiu Liu[1], Kuo Liu[2], Wenjuan Pu[1], Wendong Weng[1], Shaohua

Zhang[1], Huan Zhao[1], Kathy O. Lui[3], Bin Zhou[1,2,4,6,*]

[1]New Cornerstone Science Laboratory, State Key Laboratory of Cell Biology, CAS Center for

Excellence in Molecular Cell Science, Shanghai Institute of Biochemistry and Cell Biology,

Chinese Academy of Sciences, University of Chinese Academy of Sciences, Shanghai, China

[2]Key Laboratory of Systems Health Science of Zhejiang Province, School of Life Science,

Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou,

Chinace

[3]Department of Chemical Pathology; and Li Ka Shing Institute of Health Sciences, The Chinese

University of Hong Kong, Prince of Wales Hospital, Hong Kong, China

[4]School of Life Science and Technology, ShanghaiTech University, Shanghai, China

[5]These authors contributed equally

[6]Lead contact

[*]Correspondence: zhoubin@sibs.ac.cn

Abstract

Cre-loxP technology, a cornerstone in fate mapping and *in vivo* gene function studies, faces

challenges in achieving precise and efficient conditional mutagenesis through inducible systems.

In this study, we introduce two innovative genetic tools designed to overcome these limitations.

The first, loxCre, facilitates conditional gene targeting by allowing any CreER line to induce Cre

expression with significantly enhanced efficiency. The second, roxCre, enables DreER-mediated

Cre release, paving the way for intersectional genetic manipulation that permits both increased

precision and efficiency. Both tools incorporate a fluorescent reporter for genetic lineage tracing,

revealing efficient gene knockout in cells marked by the reporter simultaneously. These

strategies hold great potential for precise and efficient exploration of lineage-specific gene

functions, marking a significant advancement in genetic research methodologies.

Introduction

Genetic modification such as gene knockout in a precise spatiotemporal manner forms the basis

for understanding cellular and molecular mechanisms in multiple biological processes[1, 2]. The

powerful Cre-loxP recombination system is the most widely used approach employed to

knockout or over-express specific genes *in vivo*[3]. Cre recombinase, derived from bacteriophage

P1 gene, targets short palindrome DNA sequence loxP (34bp) and recombines two loxP-flanked

sequences for deletion[4, 5]. Application of the Cre-loxP system in mouse genetics has

revolutionized gene functional analysis and significantly advanced our understanding of many

developmental and pathophysiological processes at the molecular level[6, 7]. To date, most of the

mouse genes have been successfully targeted by two loxP sites (floxed allele), which could be

recombined by cell-type specific Cre lines. For many *in vivo* genetic studies, a tissue-specific

gene knockout approach has been widely used as a gold standard to dissect gene function in a

particular cell type during a biological process.

While constitutively active Cre is robust and efficient in recombination for gene deletion, Cre

lacks temporal control when the promoter driving Cre is active from the embryonic to the adult

stage. Therefore, a temporally active Cre is needed to avoid early mortality as a result of the

ablation of an indispensable gene at a key developmental stage. To overcome the temporal

limitation of Cre, CreER, the fusion of Cre with a mutated estrogen receptor, is generated to

enable its activation only after tamoxifen (Tam) treatment, and interaction of ER with Tam

induces nuclear translocation of CreER from the cytoplasm[8]. As a result, the activity of inducible

CreER could be controlled by Tam treatment, thus restricting CreER activity at a specific time

window and permitting gene functional analysis at a higher spatiotemporal resolution than Cre[9].

As the loxP-flanked sequences and their locations in genome or chromatin influence its

accessibility for recombination, the recombination efficiency of CreER varies significantly

among different floxed alleles, ranging from easy- to inert-to-recombine alleles[10]. For example,

CreER has high efficiency in targeting the easy-to-recombine alleles such as generic reporter

R26-tdT[11], but exhibits low efficiency on some other reporters such as *R26-Confetti*[12], or may not

excise some floxed gene alleles that are inert or resistant for targeting, leading to failure of gene

deletion in some tdT$^+$ cells and causing false-positive tracing fate (Figure 1A). In other words,

tdT expression may not always necessarily denote gene deletion in the labeled cells. The notion

that the pattern of Cre-expression based on a reporter (e.g. *R26-tdT*) represents the pattern of

Cre-mediated recombination is less rigorous[13]. It is therefore not correct to assume the successful

deletion of the gene of interest based on the deletion of another gene (or reporter) by the same

CreER mouse line.

One way to increase the recombination efficiency of CreER is to treat mice with many times of

Tam, as the high dosage of Tam in theory could increase the chances of CreER-mediated

recombination. However, the increased dosage of Tam treatment is toxic and could lead to many

side effects on the phenotype, creating confounding effects on the study[14, 15]. To increase the

inducible recombination efficiency of floxed alleles, Lao et al. reported a mosaic mutant analysis

with spatial and temporal control of recombination (MASTR), in which initial FlpoER-mediated

recombination induces GFPCre expression, that subsequently recombines floxed alleles

effectively in GFP$^+$ cells[16]. Similarly, a Flp-induced mosaic analysis system with Cre or Tomato

(MASCOT) has been reported to express Cre and reporter simultaneously in a cell[17]. While

MASTR and MASCOT enable spatiotemporal labeling of mosaic mutant cells utilizing the

current resources of floxed mouse lines, it may not be suitable for effective gene knockout at a

tissue/population level due to low recombination efficiency initiated by Flp recombinase. In

order to improve the inducible recombination efficiency, Tian et al. have also reported a self-

cleaved inducible CreER (sCreER) that switches inducible CreER into a constitutively active Cre,

effectively recombining floxed allele for gene manipulation[10]. However, self-cleaved inducible

Cre mice have to be newly generated each time for targeting different promoters, which is time-

consuming and the system is not compatible with current resources largely based on CreER

mouse lines. A recent study has reported an iSuRe-Cre strategy to induce and report Cre-

dependent genetic modifications utilizing conventional CreER tools[18]. The reporter expression

reflects gene knockout in cells by *iSuRe-Cre*. However, two limitations hinder its widespread

applications. First, without any induction, the *iSuRe-Cre* transgene is leaky in some tissues such

as heart and skeletal muscle. Second, the initial recombination for releasing the *iSuRe-Cre* allele

induced by CreER is far from efficient in some tissues compared with easy-to-recombine alleles

such as *R26-tdT*[18], limiting its usage for efficient gene knockout at a tissue level. Thus, a new

method is needed to ensure gene knockout in cells as effectively and efficiently as recombination

on easy-to-recombine alleles, thus allowing efficient gene deletion at a population level.

Conventional Cre-loxP system uses tissue-specific promoter to drive Cre, thus the precision of

genetic targeting solely depends on promoter activity. Now it is known that many promoters are

not as specific as previously reported[19−21]. The ectopic or unwanted Cre expression in other cell

types may result in unspecific cell labeling or gene deletion, leading to many confounding issues

or controversies in multiple fields of research[22−24]. In addition, some cell types cannot be clearly

distinguished by one marker from another, and thus could not be specifically and genetically

targeted by only one gene promoter-driven recombinase[25]. To circumvent this limit, dual

recombinases-mediated genetic targeting utilizes two promoters to independently drive Cre and

Dre[26]. Dre is another bacteriophage recombinase that targets rox sites and is orthogonal to the

Cre-loxP system[27]. While cells of interest could be labeled by specific reporters that are

responsive to dual recombinases, gene deletion requires a final readout on singular Cre

recombinase for targeting floxed gene alleles.

In this study, we developed two genetic strategies: roxCre and loxCre, in which the *rox-Stop-rox*

(RSR) and *loxP-Stop-loxP* cassettes are respectively inserted into the *Cre* coding region, such

that Cre transcription and translation are not properly carried out before the removal of the *Stop*

cassette. We did not detect any spontaneous leakiness by roxCre or loxCre. We found that

CreER-induced loxCre effectively deletes genes in reporter-labeled cells, ensuring efficient gene

knockout for the evaluation of lineage-specific gene function. In addition, DreER-induced
roxCre efficiently deletes genes in cells for both specific and efficient intersectional genetic
studies. We expect that these tools would overcome the issues of inefficient and non-specific
genetic modifications, enhancing our ability to more precisely manipulate genes in a particular
cell lineage for a better understanding of gene functions in multiple biomedical fields.

Results

Design of roxCre for DreER-induced mCre expression

A sequential genetic approach has been recently reported to delete genes[28], which uses Dre-
induced CreER expression by removing the *loxP-Stop-loxP* sequence ahead. As aforementioned,
the recombination carried out by the released CreER for excising floxed gene alleles may not be
as efficient as reporters (e.g. *R26-tdT*). A possible strategy to control Cre activity is to engineer a
rox-flanked *Stop* sequence ahead of its DNA (*R26-RSR-Cre*) that will be removed after Dre-rox
recombination (Figure 1B). However, such a strategy is not ideal, as the upstream transcriptional
Stop sequence might not completely prevent *Cre* transcription, leading to the leakiness issue
associated with the *iSuRe-Cre* strategy[18], for instance (Figure 1C). We then inserted a longer
Stop sequence or reversed the direction of the *Cre* DNA by generating *R26-RSR-Cre2* and *R26-*
R-reverseCre-R lines, respectively, that still displayed heavy leakiness as Cre was independently
activated for recombination without being crossed with a Dre mouse line (Figure 1D-G). To
enable efficient dual recombinases-mediated gene knockout, we need to design a strategy for
DreER-induced expression of a constitutively active Cre, and the system should not exhibit
leakiness.

We first generated roxCre, in which an RSR cassette was inserted into the coding region of *Cre*,
splitting *Cre* into N- and C-terminal segments (Figure 1H). After removal of RSR by Dre-rox
recombination, Cre is recombined containing one remaining rox site within its coding sequence,
hereafter termed as modified Cre (mCre, Figure 1H). To ensure that mCre has the same
recombination efficiency as a conventional Cre, we inserted a rox sequence at 12 different sites
of *Cre* individually, most of which were located between helix domains (Figure 1I). As
insertions of additional amino acids of rox would change the original sequence of Cre,
potentially leading to reduced activity, we screened for their recombination efficiency by

transfecting cells with 12 versions of mCre (mCre1 to mCre12) and the responding GFP reporter

(Figure 1J). We found that mCre1, 2, 3, 4, 6, 7, 10, and 11 displayed a comparable efficiency as

the conventional Cre; whereas mCre5, 8, 9, and 12 exhibited impaired recombination efficiency

(Figure 1K, L). These data demonstrate that at least some of the mCre generated in our study

were as efficient as the conventional Cre *in vitro*.

Next, we sought to screen for the most robust version of mCre *in vivo*. We generated four roxCre

knock-in mice based on mCre1, 4, 7, and 10 as determined from the above *in vitro* screening

(Figure S1-S2). In these roxCre lines, RSR inserted into *Cre* was removed by Dre-rox

recombination resulting in the generation of mCre (Figure 1J). We then used hepatocyte- and

endothelial cell-specific promoters albumin (Alb) and VE-cadherin (Cdh5) to drive roxCre, and

generate *Alb-roxCre1-tdT*, *Cdh5-roxCre4-tdT*, *Alb-roxCre7-GFP*, and *Cdh5-roxCre10-GFP*

knock-in mice, respectively (Figure S1-S2). No fluorescence signal was detected in *Alb-*

roxCre1-tdT, *Cdh5-roxCre4-tdT*, *Alb-roxCre7-GFP*, and *Cdh5-roxCre10-GFP* knock-in mice

(Figure S1-S2), demonstrating no reporter leakiness without Dre-rox recombination. These mice

were crossed with *R26-GFP*[29] or *R26-tdT*[11] reporters. Nor did we detect leakiness as a result of

any Cre activity when they were crossed with *R26-tdT* or *R26-GFP* reporter (Figure S1-S2).

These data demonstrate that roxCre was functionally efficient yet non-leaky.

DreER-induced mCre robustly recombines inert alleles

The mCre system was originally designed for inducible gene knockout to study biological

functions. To more rigorously evaluate the recombination effectiveness among different versions

of mCre, we used an inducible DreER to temporally and specifically release mCre in hepatocytes

or endothelial cells. Although mCre efficiently labeled virtually all targeted cells (Figure S3A-E),

a strong Cre is not needed to drive easy-to-recombine alleles such as *R26-tdT* reporter in this

case. To identify a strong mCre, we attempted to target one of the most inert alleles for

recombination, *R26-Confetti*[12], which is often used as reporters for clonal analysis due to its

sparse labeling with rare recombination events[10, 12]. In the Cre-loxp recombination principle, Cre

can remove the sequence between two loxp sites that have the same transcription direction and

turn over the sequence in the middle of two loxp sites that have the opposite transcription

direction. As the recombination efficiency of CreER is limited, *R26-Confetti* only can be recombined into YFP, or nGFP (nuclear GFP), or mCFP (membrane CFP), or RFP (Figure S3F).

We first crossed *R26-DreER;R26-Confetti* mice with *Alb-roxCre1-tdT* (group 1) and *Alb-roxCre7-GFP* (group 2) mice, and Tam-induced Dre-rox recombination yielded *Alb-mCre1-tdT* and *Alb-mCre7-GFP* mice, respectively (Figure 2A). In this system, the constitutively active mCre would completely remove the sequence between two loxp sites that have the same transcription direction, and recombine *R26-Confetti* into two sets of reporter pairs: YFP-nGFP pair and mCFP-RFP pair, which could be detected in a single hepatocyte due to the poly-nuclear or polyploidy feature of hepatocytes (Figure 2A). In each pair, reporters can equally express, as mCre is strong enough to turn over the sequence in the middle of two loxp sites that have the opposite transcription direction (Figure S3G). Therefore, we used YFP/mCFP to detect mCre recombination efficiency and tdT/RFP or GFP to trace its endogenous reporter activity that was driven by Alb promoter in *Alb-mCre1-tdT* and *Alb-mCre7-GFP* mice, respectively (Figure 2A). Besides, *Alb-CreER*[26]*;R26-Confetti* was used as a control under the same Tam treatment (group 3, Figure 2A).

In fact, DreER-rox recombination in hepatocytes is not 100%. Therefore, we quantified YFP and mCFP expression in hepatocytes with DreER-rox recombination to evaluate the strength of mCre activity. We found sparse YFP^+ or $mCFP^+$ hepatocytes ($0.39 \pm 0.04\%$) in *Alb-CreER;R26-Confetti* mice. On the other hand, $99.94 \pm 0.02\%$ of tdT^+ hepatocytes (DreER recombined) were YFP^+ or $mCFP^+$ in *R26-DreER;Alb-roxCre1-tdT;R26-Confetti* mice; and $76.83 \pm 3.20\%$ of GFP^+ hepatocytes (DreER recombined) were YFP^+ or $mCFP^+$ in *R26-DreER;Alb-roxCre7-GFP;R26-Confetti* mice (Figure 2B, C). Similarly, we found that $39.81 \pm 2.61\%$ of tdT^+ endothelial cells and $74.15 \pm 3.64\%$ of GFP^+ endothelial cells in small intestine were YFP^+ or $mCFP^+$ in *R26-DreER;Cdh5-roxCre4-tdT;R26-Confetti* and *R26-DreER;Cdh5-roxCre10-GFP;R26-Confetti* mice, respectively, compared with sparse labeling in *Cdh5-CreER;R26-Confetti* mice (Figure 2D-F). Similar results were also observed in other tissues and organs (Figure S4). These data demonstrate that inducible DreER controlled roxCre activation, and mCre1 was the most efficient recombinase assisting the recombination of inert alleles *in vivo*.

Dre-induced mCre efficiently deletes floxed genes

Next, we examined the efficiency of gene knockout by Dre-induced mCre. We generated the *Alb-roxCre1-tdT;Ctnnb1$^{flox/flox}$* mouse line to examine if Dre-induced mCre could efficiently delete *Ctnnb1* gene that encodes β-catenin (Ctnnb1^{flox})[30]. To enhance the precision and specificity of targeted cells, we employed intersectional genetic targeting using dual recombinases, and generated the *Cyp2e1-DreER* mouse line that labeled peri-central hepatocytes and also renal epithelial cells (Figure S5). Crossing *Cyp2e1-DreER* with hepatocyte-specific *Alb-roxCre1-tdT* mice achieved genetic targeting of peri-central hepatocytes exclusively, circumventing issues associated with ectopic targeting of renal cells by *Cyp2e1-DreER* line (Figure S5) or unwanted targeting of peri-portal hepatocytes by *Alb-CreER* line[26].

We generated *Cyp2e1-DreER;Alb-roxCre1-tdT;Ctnnb1$^{flox/flox}$* triple knock-in mice (mutant), in which Tam-induced DreER-rox recombination yielded mCre that subsequently targeted floxed Ctnnb1 allele (Figure 3A). We treated mutant mice and their littermate control *Cyp2e1-DreER;Alb-roxCre1-tdT;Ctnnb1$^{flox/+}$* mice with Tam at 8 weeks of age, and purified tdT$^+$ hepatocytes for analysis at 3 days after Tam treatment (Figure 3B). qRT-PCR analysis showed significantly reduced *Ctnnb1* and *Glul* in tdT$^+$ hepatocytes of the mutant group, compared to that of the control group (Figure 3C, D). We then collected liver samples at 3 days (D) and 4 weeks (W) post-Tam for further analysis (Figure 3E). Immunostaining for tdT, β-catenin, GS, and E-CAD on liver sections revealed similar levels of β-catenin and GS expression in the 3D mutant liver than the control liver (Figure 3F). However, expression of β-catenin and GS almost disappeared in the 4W mutant sample compared with the control (Figure 3G), suggesting efficient deletion of *Ctnnb1* and down-regulation of Wnt signaling downstream gene target GS. Western blotting against β-catenin and qRT-PCR against *Ctnnb1* of purified tdT$^+$ hepatocytes at 4W post-Tam revealed a near complete loss of Ctnnb1 at both the protein and mRNA levels (Figure 3H-K). Additionally, Wnt downstream target genes including *Glul*, *Axin2*, *Cyp1a2*, *Cyp2e1*, *Oat*, *Tcf7*, *Lect2*, *Tbx3*, *Slc1a2*, and *Rhbg* were significantly reduced in the mutant than the control group (Figure 3L). Taken together, these data demonstrate that DreER-induced mCre enabled efficient intersectional genetic manipulation in the cells of interest.

***R26-loxCre-tdT* is efficiently recombined by CreER**

Having successfully established roxCre1 for Dre-induced mCre expression, we iterated the

system to enable mCre induction by CreER, a more widely available tool for biomedical research.

We replaced the rox sites of roxCre1 with loxP sites by inserting a *loxP-Stop-loxP* cassette into

Cre cDNA at the same locus as roxCre1(Figure 4A). We then generated a new *R26-loxCre-tdT*

mouse line, in which mCre and tdT would be expressed simultaneously after *Stop* removal,

allowing tdT as a surrogate marker for mCre detection in the same cell (Figure 4C). We then

designed two experiments to verify the baseline leakiness of Cre expression (Figure 4B) and to

examine if mCre/tdT could be expressed after *Stop* removal (Figure 4C), respectively, by *R26-*

loxCre-tdT;R26-tdT (Strategy 1) and by injecting *AAV8-Cre* virus into *R26-loxCre-tdT* mice

(Strategy 2). Immunostaining for tdT on tissue sections revealed no tdT expression in Strategy 1

demonstrating minimal to negligible leakiness at the baseline. The robust tdT expression in the

liver in Strategy 2 showed mCre/tdT was allowed to express after *Stop* removal by AAV2/8-

hTBG-Cre(Figure 4D).

We then confirm if the new design can improve the labeling efficiency compared to the

conventional approach by crossing endothelial cell-specific *Cdh5-CreER* with the conventional

reporter *R26-tdT* (Strategy 3) and the new reporter *R26-roxCre-tdT* mice (strategy 4) under the

same protocol of Tam treatment (Figure 4E). Immunostaining for tdT and VE-Cad on tissue

sections revealed no difference in the percentage of VE-Cad$^+$ endothelial cells expressing tdT

between strategies 3 and 4 (Figure 4F- G), suggesting that recombination of the *R26-roxCre-tdT*

allele by CreER was as efficient and easy as *R26-tdT* which is an easy-to-recombine reporter

mouse line. Therefore, *R26-roxCre-tdT* can be served as an easy-to-recombine reporter, enabling

efficient recombination to generate mCre/tdT in CreER-expressing cells, without leakiness.

***R26-loxCre-tdT* adaptor ensures efficient recombination**

We next examined whether mCre released from loxCre could also display high recombination

efficiency for the inert-to-recombine allele *R26-Confetti*. We generated the triple knock-in *Cdh5-*

CreER;R26-loxCre-tdT;R26-Confetti mouse line in which mCre was released for confetti

recombination upon Tam treatment, and *Cdh5-CreER;R26-Confetti* was used as a control (Figure

5A). Both mouse strains were treated with Tam at 8 weeks of age and their tissue samples were

collected for analysis at one week post-Tam (Figure 5B). As tdT expression is also detected in

R26-loxCre-tdT after the first recombination by CreER, we only compared the YFP and mCFP

fluorescence signals from *R26-Confetti* (Figure 5A). Wholemount-fluorescence imaging showed

significantly more YFP and mCFP signals in the retina of the *Cdh5-CreER;R26-loxCre-tdT;R26-*

Confetti mice, compared with that of *Cdh5-CreER;R26-Confetti* mice (Figure 5C).

Immunostaining on frozen sections revealed significantly higher percentage of endothelial cells

expressing YFP/mCFP in various vascularized organs of the *Cdh5-2a-CreER;R26-loxCre-*

tdT;R26-Confetti mice, compared with that of the *Cdh5-2a-CreER;R26-Confetti* mice (Figure

5D). Of note, virtually all tdT$^+$ endothelial cells simultaneously expressed YFP/mCFP (Figure

5D). These data demonstrate that the *R26-loxCre-tdT* line could be used as an adaptor to enhance

CreER-mediated recombination with a simultaneous expression of tdT as an accurate indicator.

R26-loxCre-tdT ensures gene deletion in tdT$^+$ cells

To examine if *R26-roxCre-tdT* enables *Alb-CreER* to more efficiently delete genes, we crossed

Alb-CreER;R26-roxCre-tdT   with   *Ctnnb1$^{flox/flox}$*   to   generate   *Alb-CreER;R26-roxCre-*

tdT;Ctnnb1$^{flox/flox}$ mice for evaluating the efficiency of releasing mCre and also mCre-mediated

Ctnnb1 deletion in tdT$^+$ hepatocytes (Figure 6A). We also crossed *Alb-CreER;R26-tdT2* with

Ctnnb1$^{flox/flox}$ to generate *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$* mice to evaluate recombination

efficiency of both *R26-tdT2* and *Ctnnb1$^{flox/flox}$* alleles (Figure 6A). Additionally, *Alb-CreER;R26-*

roxCre-tdT;Ctnnb1$^{flox/+}$ mice were used as a control for heterozygous gene deletion (Figure 6B).

We treated *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$* and *Alb-CreER;R26-roxCre-tdT;Ctnnb1$^{flox/+}$* mice

with five dosages of Tam, and treated *Alb-CreER;R26-roxCre-tdT;Ctnnb1$^{flox/flox}$* mice only with

one dosage of Tam, and collected tissues for analysis to determine *Ctnnb1* gene deletion at 4

weeks after Tam treatment (Figure 6B). Immunostaining data showed that β-catenin, GS, E-

CAD, and Cyp2e1 were significantly reduced in *Alb-CreER;R26-roxCre-tdT;Ctnnb1$^{flox/flox}$* mice,

but were readily detectable in both *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$* and *Alb-CreER;R26-*

roxCre-tdT;Ctnnb1$^{flox/+}$ mice (Figure 6C, D). Of note, a small number of hepatocytes continued

to express Cyp2e1 in *Alb-CreER;R26-roxCre-tdT;Ctnnb1$^{flox/flox}$* mice (Figure 6D). These

Cyp2e1-expressing hepatocytes were unanimously tdT$^−$, and not a single tdT$^+$ hepatocyte

expressed Cyp2e1 (Figure 6D), suggesting that these Cype2e1$^+$tdT$^−$ hepatocytes were likely not

targeted by *Alb-CreER* for the recombination to release mCre. Western blotting of β-catenin and

qRT-PCR of *Ctnnb1* of isolated hepatocytes revealed almost complete deletion of Ctnnb1 in the

Alb-CreER;R26-tdT2;Ctnnb1^{flox/flox} mice compared with those in control groups (Figure 6E-G).

Additionally, those Wnt downstream or regulated genes including *Glul*, *Axin2*, *Cyp1a2*, *Cyp2e1*,

Oat, *Tcf7*, *Lect2*, *Tbx3*, *Slc1a2*, and *Rhbg* were significantly reduced in hepatocytes derived from

the *Alb-CreER;R26-tdT2;Ctnnb1^{flox/flox}* mice, compared with those from the *Alb-CreER;R26-*

tdT2;Ctnnb1^{flox/flox} and *Alb-CreER;R26-roxCre-tdT;Ctnnb1^{flox/+}* mice, respectively (Figure 6H).

Discussion

In this study, we respectively generated two genetic tools based on roxCre and loxCre to enhance

the effectiveness of Cre-mediated gene deletion even under the control of a less efficient DreER

or CreER driver. The *R26-loxCre-tdT* line permitted efficient recombination to release mCre,

which greatly enhanced the effectiveness of subsequent gene deletion in tdT$^+$ cells. The inclusion

of a fluorescent reporter with roxCre or loxCre allele allowed us to readily detect cells with gene

deletion, thus facilitating further detailed characterization of these cells both *in situ* and *ex vivo*.

Furthermore, the strategy of roxCre further enhances the precision for cell fate mapping with

genetic deletion simultaneously in Dre$^+$Cre$^+$ cells through intersectional genetic targeting. For

example, we have showcased the power of roxCre for genetic manipulation in peri-central

hepatocytes, resulting in efficient gene knockout in tdT$^+$ hepatocytes specifically. Additionally,

we have demonstrated the efficient deletion of *Wls* and *Ctnnb1* genes in tdT$^+$ endothelial cells

and hepatocytes, respectively, by combining *R26-loxCre-tdT* adaptors with CreER lines.

Altogether, we have demonstrated that the *R26-loxCre-tdT* mouse line could be widely

applicable to any CreER-bearing mouse for enhancing its effectiveness in genetic modification

with simultaneous cell lineage tracing in multiple biomedical research fields.

Coupling gene deletion with reporter activity can greatly facilitate the accuracy of studying gene

function in targeted cells by *in situ* investigation with or without specific cell isolation for *ex vivo*

experiments. Assuming successful gene deletion in the cell type of interest because of efficient

cell labeling by a lineage-specific reporter mediated by the same Cre is incorrect, as

recombination efficiency varies significantly among different mouse lines. In fact, many factors

can potentially impact Cre-loxP recombination efficiency. Concerning the recombinase enzyme,

the type of recombinase, the strength of the promoter that drives the recombinase, the expression

level of the recombinase, and the translocation efficiency of an inducible recombinase such as

CreER from the cytoplasm to the nucleus after Tam induction could all influence the
recombination efficiency. For the targeted DNA sites, the genomic loci where targeted sites are
placed, loxP-flanked sequence length, and even the inserted DNA sequence *per se* could also
influence the recombination efficiency. We clearly observed the differences in recombination
efficiency of different alleles (e.g. *R26-tdT*, *R26-tdT2*, and *R26-Confetti*), even located in the
same genomic locus (e.g. Rosa26) mediated by the same CreER under the same Tam treatment
protocol, as evidenced by high efficiency in *R26-tdT*, medium efficiency in *R26-tdT2*, and low
efficiency in *R26-Confetti*. Importantly, *R26-loxCre-tdT* was found to be efficient for
recombination mediated by CreER, which was comparable to that of the *R26-tdT* allele, thus
enabling efficient labeling of cells of interest. Furthermore, *R26-loxCre-tdT* also enabled CreER
to efficiently ablate genes in any cells that had been genetically traced with tdT, synchronizing
genetic modification and lineage tracing in the labeled cells.

Despite the high demand in the quest for efficient genetic modification in genetically labeled
cells is significant, no efficient tool has been available to date. A previous study attempted to
generate an adaptor for CreER to efficiently delete genes in reporter-labeled cells by a transgenic
iSuRe-Cre[18]. However, the leakiness of *iSuRe-Cre* on the Rosa26 locus was too high in the male
germline. Therefore, a transgenic mouse line was reconstructed subjected to *iSuRe-Cre* mediated
recombination, which still displays consistently leaky recombination in some tissues such as
cardiac and skeletal muscle[18]. The leakiness was likely due to the inability of the *Stop* cassette to
completely block the transcription of *Cre* in *iSuRe-Cre*. Consistently, we also found leakiness as
a result of *Cre* transcription even in the presence of *Stop* ahead in the *R26-Stop-Cre* mouse line
(Figure 1C). These leaky events, despite being at different levels, demonstrate that transcription
of recombinases such as Cre could not be easily inhibited. Instead of blocking its transcription by
Stop, we inserted *Stop* in between the N- and C-terminus of *Cre*, thus preventing Cre expression
completely at the translational level. We found that the removal of *Stop* rendered mCre, which
efficiently recombined different types of alleles examined in our study. Additionally, we also
observed inefficient recombination in hepatocyte-specific *Alb-CreER;iSuRe -Cre* mice,
rendering this line less effective in conditional gene deletion in hepatocytes at a population level.
In contrast, *R26-loxCre-tdT* was an easy-to-recombine allele for *Alb-CreER* mediated
recombination, allowing tdT labeling and gene deletion simultaneously in the majority of

hepatocytes. Therefore, our *R26-loxCre-tdT* line has circumvented the leakiness issue associated

with *iSuRe-Cre*, exhibited as an easy-to-recombine allele for CreER targeting, and synchronized

fate mapping with efficient genetic modification robustly, efficiently and temporally in the same

cells.

The efficiency of CreER-loxP recombination depends not only on the location of the loxP-

containing alleles but also on the strength of CreER driven by an active promoter. For instance,

R26-loxCre-tdT was an easy-to-recombine allele compared with other tested R26 alleles, but the

recombination efficiency to release mCre for subsequent recombination depends on the strength

of the selected CreER lines for genetic targeting. It is known that different CreER mouse lines

vary significantly for recombination, largely due to the differences in expression levels of CreER

and Tam induction efficacy. Therefore, we should select a relatively stronger CreER line for

efficient recombination on *R26-loxCre-tdT* to release mCre for gene deletion at a population

level. In addition to bulk gene deletion, the *R26-loxCre-tdT* also offers an alternative way for

mosaic analysis of gene function by sparse labeling. Previous studies have used Cre-loxP to

over-express gene and fluorescence reporter simultaneously in a cell for functional genetic

mosaic (ifgMosaic) analysis[31]. While ifgMosaic could over-express some dominantly negative

genes for functional study, this system could not take advantage of rich resources from many

conventionally available floxed mouse lines for loss-of-function study. Mosaic mutant analysis

with spatial and temporal control of recombination (MASTR) utilized FlpoER for tissue-specific

expression of GFPCre to delete floxed genes[16]. Compared with FlpoER lines, CreER is a more

broadly used recombinase that most laboratories use in gene deletion experiments. A recent

study using dual recombinase-mediated cassette exchange (MADR) also permits stable labeling

of mutant cells expressing transgenic elements from defined chromosomal loci[32]. However, the

use of viruses and electroporation, albeit convenient and rapid, are inefficient to specifically

target any type of cells for *in vivo* studies. The powerful mosaic analysis with double markers

(MADM) enables simultaneous lineage tracing of a pair of mutant and control sibling cells with

distinct fluorescence reporters, allowing precise mosaic analysis of gene function in any cell[33, 34].

Nevertheless, the MADM cassette has to be combined with the mutant null allele, which is not

readily available for synchronizing reporter expression and genetic modification in cells of

interest. The ensured gene deletion with tdT reporter by *R26-roxCre-tdT* could be coupled with

any present CreER and loxP alleles for potential mosaic analysis. This could be achieved by

adjusting Tam at a lower dosage so that the recombination efficiency to release mCre could be

low in cells of interest for sparse tdT labeling. We believe that *R26-loxCre-tdT* could be useful

for robust, efficient, and temporal gene deletion at a population level and mosaic analysis at a

single-cell level with improvement in the future.

In this study, we deleted the Wnt signaling gene *Ctnnb1* in hepatocytes to present how roxCre

and loxCre respectively enable efficient recombination. The impact of deleting *Ctnnb1* in a

specific hepatocyte lineage on homeostasis, development, and injury recovery requires further

investigation. Besides, roxCre and loxCre should be involved in more gene functional studies, in

which knocking out genes continuously leads to embryonic death due to their crucial role in

development.Both loxCre and roxCre incorporate a fluorescent reporter tdT for genetic lineage

tracing, revealing efficient gene knockout in cells marked by the reporter simultaneously. In

other words, one reporter tdT can only show cells with gene knockout. One direction to improve

the current *R26-loxCre-tdT* for mosaic analysis in the future is to generate a second distinct

surrogate reporter for wild-type cells as an internal control during mosaic analysis, similar to that

in the MADM system.

Materials and Methods

Mice

Experiments using mice (Mus musculus) were carried out with the study protocols (SIBCB-S374-1702-001-C1) approved by the Institutional Animal Care and Use Committee of Center for Excellence in Molecular Cell Science (CEMCS), Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences. The *R26-DreER*[35], *Alb-CreER*[26], *R26-GFP*[29], *R26-tdT*[11], *R26-tdT2*[36], *R26-RSR-tdT*[37], *R26-Confetti*[12],and *Ctnnb1-flox*[30] mouse lines were used as previously described. New knock-in mouse lines *R26-RSR-Cre*, *R26-RSR-Cre2*, *R26-R-reverseCre-R*, *Cdh5-CreER*, *Alb-roxCre1-tdT*, *Alb-roxCre7-GFP*, *Cdh5-roxCre4-tdT*, *Cdh5-roxCre10-GFP*, *Cyp2e1-DreER*, and *R26-loxCre-tdT* were generated by homologous recombination using CRISPR/Cas9 technology. These new mouse lines were generated by the Shanghai Model Organisms Center, Inc. (SMOC). These mice were bred in a C57BL6/ICR mixed background. All mice were housed at the laboratory Animal center of the Center for Excellence in Molecular Cell Science in a Specific Pathogen Free facility with individually ventilated cages. The room has controlled temperature (20–25 °C), humidity (30–70%), and light (12 hours light-dark cycle). For the determination of the embryonic period of sampling, the day on which the vaginal plug was examined in female mice was recorded as E0.5. As the sex is not relevant to the topic of this study, male and female mice ranging in age from E13.5 to 12W (week) were allocated and mixed into the experimental groups in this study. No data in mouse experiments were excluded.

Genomic PCR

Genomic DNA was prepared from the mouse toes or embryonic tails. Tissues were precipitated by centrifugation at maximum speed for 1 min at room temperature. After that, tissues were lysed by lysis buffer (100 mM Tris-HCl, 5 mM EDTA, 0.2% SDS, 200 mM NaCl, and 100 µg/mL Proteinase K) at 55 °C overnight. About 750µL pure ethanol was added to the lysis mixture and mixed thoroughly, followed by centrifugation at maximum speed for 5 min at room temperature to collect the DNA precipitation. Then the supernatant was discarded and the mixture was dried at 55 °C for 1 hour. About 100–200µL double-distilled H_2O was added to dissolve the DNA. The genomic PCR primer pairs were designed for the mutant alleles spanning

both endogenous genomic fragments and insert fragments. All the new genomic PCR primer

sequences would be provided upon request.

***In vitro* screening of mCre for efficient recombination**

To test if the remaining rox sequence after Dre-rox recombination would impact Cre activity, the

pCDNA3.1[38] (Invitrogen, V79020) was used to express 12 types of modified Cre (mCre). For

testing the Cre activity, 500ng pCDNA3.1-mCre plasmids, or pCDNA3.1 (negative control), or

pHR-CMV-nlsCRE (positive control, Invitrogen, 12265) were mixed with 500ng pCAG-loxp-

stop-loxp-ZsGreen (Addgene, 51269) plasmid. *In-vitro* transfection experiment protocol is

performed according to that described previously[39]. For cell culture medium, DMEM

(ThermoFisher, 11965092) was supplemented with 10% fetal bovine serum (fetal bovine serum,

10099141c) for preparing fresh complete culture medium. Poly-D-Lysine (ThermoFisher,

A3890401) precoated coverslips (biosharp, BS-14-RC) or 10 cm dish (ThermoFisher, 150466)

overnight to dry, and sterile water was used to flush it for three times. A cryogenic vial of

QBI293 was put in a 37°C water bath, then thawed liquid contents into a 50 mL conical tube

(Corning,430829) prefilled with 5 mL prewarmed fresh complete culture medium. After

centrifuging at $125 \times g$ for 5 min, the supernatant was discarded and cells were resuspended in

10 mL complete medium. One-third of the cells were plated in a 10 cm coated dish and

incubated in cultures at 37°C. On the second day, the culture medium was discarded, and the cell

layer was briefly rinsed with prewarmed PBS (Gibco, C10010500BT). After that, PBS was

removed and 1 mL 0.25% Trypsin solution (with EDTA, Gibco, 25200072) was added. The cells

were incubated at 37°C for 3 min., and a 6mL complete growth medium was added to aspirate

cells by gently pipetting. The medium-containing cells were transferred to a 50 mL conical tube

and centrifuged at $125 \times g$ for 5 min. The supernatant was discarded, and cells were resuspended

in 5.2 mL complete culture medium. 24 well plates were placed into coated coverslips and added

1mL complete culture medium. 200µL cells were added to every experimental well and

incubated cultures at 37°C for 9 hours to grow up to ~80%. The new prewarmed fresh complete

culture medium replaced the old for 1mL per well. Lipofectamine™ 3000 Transfection Reagent

(Thermo Fisher, L3000015) was used for plasmid transfection. Samples were incubated in

cultures at 37°C for 24 hours. Samples were washed by PBS once and mounted on slides by

VECTASHIELD$^{®.}$ Antifade Mounting Medium with DAPI (Vector, H-1200). Images were

acquired using an Olympus microscope (Olympus, BX53).

Tamoxifen injection and sample collection

For induction of CreER or DreER, tamoxifen (Sigma, T5648) was dissolved in corn oil and was

administered to mice by oral gavage. *Cyp2e1-DreER;Alb-roxCre1-tdT;Ctnnb1$^{flox/+}$* and *Cyp2e1-*

DreER;Alb-roxCre1-tdT;Ctnnb1$^{flox/flox}$ mice were treated with 0.05 mg Tam per gram mouse

body weight for one dose. For comparing tdT$^+$ cells, *Cyp2e1-DreER;R26-RSR-tdT* and *Cyp2e1-*

DreER;Alb-roxCre1-tdT mice were treated with 0.05 mg Tam per gram mouse body weight.

R26-DreER;Alb-roxCre1-tdT;R26-Confetti, *R26-DreER;Cdh5-roxCre4-tdT;R26-Confetti*, *R26-*

DreER;Alb-roxCre7-GFP;R26-Confetti, *R26-DreER;Cdh5-roxCre10-GFP;R26-Confetti*, *R26-*

DreER; R26-DreER;Alb-roxCre7-GFP;R26-GFP, *Alb-CreER;R26-Confetti*, *Cdh5-CreER;R26-*

Confetti, *Alb-CreER;R26-tdT2;R26-Confetti*, *Alb-CreER;iSuRe-Cre;R26-Confetti*, *Alb-*

CreER;R26-loxCre-tdT;R26-Confetti, *Alb-CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$*, and *Cyp2e1-*

DreER;R26-RSR-tdT mice were treated with 0.2 mg Tam per gram mouse body weight for one

dose. *Cdh5-CreER;R26-tdT*, *Cdh5-CreER;R26-loxCre-tdT*, *Cdh5-CreER;R26-Confetti*, *Cdh5-*

CreER;R26-loxCre-tdT;R26-Confetti, *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$*, and *Alb-CreER;R26-*

loxCre-tdT;Ctnnb1$^{flox/+}$ mice were treated with 0.2 mg Tam per gram mouse body weight one

dose per day for 5 days. *R26-RSR-Cre2;R26-GFP* and *R26-R-reverseCre-R;R26-tdT* mice were

sacrificed at 8 weeks old for analysis. Mice, both males and females, at the age of 8–12 weeks

were used for experiments with similar-aged mice for both control and experimental groups.

Whole-mount imaging and sectioning

The tissue samples were fixed with 4% paraformaldehyde (PFA) (Sigma, P6148-500g, wt/vol in

PBS) for 1 or 2 hours (stomach, intestine, colon) at 4°C, followed by washing with PBS three

times. The fixed tissues were placed in an agarose-filled petri dish for bright-field and

fluorescence imaging by a Zeiss stereoscopic microscope (AxioZoom V16). For cryo-sections,

tissues were sectioned to slides of 10-μm thickness after dehydration by 30% sucrose

(Sinopharm, H-10021463, wt/vol in PBS) overnight and pre-embedding with OCT (Sakura) at

4 °C for 1 hour.

Immunostaining

Immunostaining was performed as previously described[40]. 0.2% PBST was prepared with 0.2% (vol/vol) Triton X-100(Sigma, T9284) dissolved in PBS. Tissue sections were blocked with 2.5% normal donkey serum and 0.1% 4'6-diamidino-2-phenylindole (DAPI, Vector lab, D21490) dissolved in 0.2% PBST for 30 mins after washing with PBS three times. The tissue sections were incubated with primary antibody diluted in 0.2% PBST at 4°C overnight. On the next day, sections were incubated with secondary antibodies diluted in 0.2% PBST at room temperature for 30 min, followed by PBS washing for three times. The slides were washed with PBS for three times. The slides were mounted with a mounting medium (Vector Lab). For weak signals, the endogenous peroxidase activity was quenched before blocking. Horseradish peroxidase or biotin-conjugated secondary antibodies and a Tyramide signal amplification kit (PerkinElmer) were used after incubating the primary antibodies. For primary antibodies of murine origin, mouse immunoglobulins were blocked with an anti-mouse Fab antibody (Jackson, 715-007-003, 1:100). The included primary antibodies are listed as follows: tdT (Rockland, 600-401-379, 1:500; or Rockland, 200-101-379, 1:500), GFP (Invitrogen, A11122, 1:500), GFP (Rockland, 600-101-215M, 1:500), GFP (Nacalai, 04404-84; 1:500), GS (Abcam, ab49873, 1:1000), β-catenin (BD Pharmingen, 610153, 1:200), E-CAD (R&D, AF748, 1:500), Ep-CAM (Abcam, ab92382, 1:500), VE-Cad (R&D, AF1002, 1:100), and CYP2E1(Abcam, ab28146, 1:100). The corresponding secondary antibodies (JIR or Abcam) were diluted according to the instructions. Images were captured by using a Nikon confocal (Nikon A1 FLIM) or an Olympus confocal (FV3000), and captured images were analyzed by Image J2 (version 2.9.0/1.54f) and Photoline (version 23.02) software. We collected five random fields from each liver section for quantification. Mutant and control sections were processed at the same time to avoid potential batch differences during staining. Imaging of all immunostained slides was performed under the same exposure and contrast conditions using the same confocal microscope.

Imaging of Confetti fluorescence reporters

All the fluorescence images of *R26-confetti* were taken by a Leica confocal (Leica SP8 will). All the sections were collected freshly and blocked with 0.2% PBST containing DAPI for 30 mins after washing with PBS three times. The parameters of excitement light and emission light for all channels are set as follows: DAPI (ex.405nm, em.415-450nm), CFP (ex.458nm, em.463-481nm),

GFP (ex.488nm, em.495-511nm), YFP (ex.520nm, em.527-545nm), tdT (ex.546nm, em.555-590nm), markers for antibody staining (ex.647nm, em.657-700nm). Captured images were analyzed by Image J2 (version 2.9.0/1.54f) and Photoline (version 23.02) software.

Hepatocyte dissociation

Mouse primary hepatocytes were isolated by the two-step collagenase perfusion method, which was modified from a previous protocol[41]. Briefly, mice were anesthetized and the liver was exposed through an incision in the lower abdomen. The inferior vena cava and portal vein were also exposed. A needle was inserted into the inferior vena cava and secured with a hemostatic clamp around the needle. The portal vein was cut immediately when the mouse liver was perfused with a perfusion medium buffer (containing 0.5 mM EGTA) for 5 minutes using a peristaltic pump. Then, the liver was perfused with medium containing collagenase type I (150 U/ml; Gibco, 17100-017) for 2-5 min to adequately digest the liver. After the gallbladder was removed, the liver was dissected with cold DMEM to free the hepatic cells. Then the cell suspension was passed through a 70-μm cell strainer (BD Biosciences, 352350) and centrifuged at 50 x g for 3 min at 4 °C. The supernatant was removed, and cells were resuspended in Percoll (GE Healthcare, 17-0891-01)/DMEM/10 x PBS (Sangon, B548117, diluted in 1:1) (1:1:0.1) mixture and centrifuged at 300 x g for 5 min at 4 °C. After the supernatant was removed, cells were dissected with cold DMEM and centrifuged at 50 x g for 3 min at 4 °C. Purified hepatocytes were collected for FACS analysis, qRT-PCR or Western Blot analysis.

Flow cytometric analysis and isolation of tdT$^+$ hepatocytes

Cells were stained with the following antibodies: anti-mouse CD45 APC (eBioscience, 17-0451-82, 1:400), anti-mouse CD140a APC (eBioscience, 17-1401-81, 1:100), anti-mouse CD31 APC (eBioscience, 17-0311-82, 1:40). All antibodies were diluted in 1~2mL DMEM (according to cells number) and cells were incubated with antibodies for 30 min in the dark at 4 °C. After staining, 10mL DMEM was added to stop staining, and the cell suspension was passed through a 70-μm cell strainer. Cells were centrifuged at 50 x g for 3 min at 4 °C and dissected with relevant suspended solution. For no DAPI control group, their suspended solution is 0.1% Dnase I (Worthington, LS002139, diluted by DMEM), otherwise, the suspended solution is 0.1% Dnase I mixed with 0.1% DAPI. Hepatocytes were sorted by FACS Aria SORP machine. Cell viability

was assessed with DAPI staining. APC$^-$tdT$^+$ hepatocytes were collected for subsequent

experiments.

Total RNA extraction and qRT-PCR

Total RNA was extracted from the liver of indicated mice or hepatocytes isolated from indicated

mice as previously described[42]. Cells or homogenized tissues were lysed with Trizol (Invitrogen,

15596018), and total RNA was extracted according to the manufacturer's instructions. For each

sample, 1 μg of total RNA was reversely transcribed into cDNA using the Prime Script RT kit

(Takara, RR047A). The SYBR Green qPCR master mix (Thermo Fisher Scientific, 4367659)

was used and quantitative RT-PCR was performed on QuantStudio 6 Real-Time PCR System

(Thermo Fisher Scientific). *Gapdh* was used as the internal control. Sequences of all primers

would be provided upon request.

Western blot

Liver tissues were collected at the indicated stages. All samples were lysed in RIPA lysis buffer

(Beyotime, P0013B) containing protease inhibitors (Roche, 11836153001) for 30 min on ice, and

then centrifuged at 15,000 x g for 15 min to collect the supernatant. All samples were separated

with 30μL for testing the protein concentration by Pierce^{TM} BCA Protein Assay kits (Thermo

Scientific^{TM},23227). The remaining samples were mixed with 5 x loading buffer (Beyotime,

p0015L) and boiled at 100 °C for 5 min. 40μg samples were added to every well in precast

gradient gels (Beyotime, P0469M) with 1x running buffer (Epizyme, PS105S, should be diluted

into 1x). After running, samples were transferred onto Immobilon® PVDF membranes

(Millipore, IPVH00010). After blocking in blocking buffer (Epizyme, PS108P), the membranes

were incubated with primary antibodies (diluted by primary antibody dilution buffer (Epzyme,

PS114)) overnight at 4 °C, then washed for three times and incubated with HRP-conjugated

secondary antibodies (diluted by 1xTBST (Epzyme, PS103S, should be diluted into 1x)).

Samples were incubated with chemiluminescent HRP substrate (Thermo Fisher Scientific,

WBKLS0500), and related signals were detected by MiniChemi 610 Plus (Biogp). The following

antibodies were used: β-catenin (BD Pharmingen, 610153, 1: 5000), GAPDH (Proteintech,

60004-1-IG, 1:2000), β-actin (Epzyme, LF202,1:1000), HRP-donkey-a-mouse (JIR, 715-035-

150, 1:5000), and HRP-goat a rabbit IgG (JIR, 111-035-047, 1:5000).

AAV injection

The AAVs used in this study were purchased from Taitool Biotechnologies company (Shanghai, China). Briefly, AAVs were produced with cis-plasmids containing the full TBG promoter, which is specifically active in hepatocytes, and Cre expression is under the control of TBG promoter, replication incompetent AAV2/8-TBG-Cre virus (AAV8-TBG-Cre, S0657-8-H5) was packaged and purified before application to mice. Mice were injected intraperitoneally at $1x\ 10^{11}$ genome copies of the virus per mouse. The above newly generated virus as well as targeting plasmids will be provided upon request.

Statistical analysis

Statistical analyses were performed using GraphPad Prism (Version 9.5.1). All the continuous variables were expressed as means ± standard error of the mean (SEM). One-way ANOVA was used to detect statistical significance between three experimental groups. The statistical difference between the two experimental groups was determined using the unpaired student's *t*-test. The probability (*P*) values of less than 0.05 were considered statistically significant.

Acknowledgments

This study was supported by the National Key Research & Development Program of China (2023YFA1800700, 2022YFA1104200, 2023YFA1801300, 2020YFA0803202), National Natural Science Foundation of China (82088101, 32370897, 32100648, 32370783, 32100592), CAS Project for Young Scientists in Basic Research (YSBR-012), Shanghai Pilot Program for Basic Research-CAS, Shanghai Branch (JCYJ-SHFY-2021-0), Research Funds of Hangzhou Institute for Advanced Study (2022ZZ01015, B04006C01600515), Shanghai Municipal Science and Technology Major Project, and the New Cornerstone Science Foundation through the New Cornerstone Investigator Program and the XPLORER PRIZE.

Author contributions

M.S., J.L., and B.Z. designed the study, analyzed the data, and wrote the manuscript; X.L., K.L., W.P., W.W., S.Z., and H.Z. bred the mice and performed experiments; K.O.L. edited the manuscript and provided intellectual input; B.Z. conceived, supervised, and organized the study.

Declaration of interests

The authors declare no competing interests.

Data availability

All data included in this study are included in the manuscript. The raw data are available upon request.

References

1. Gu, H., Zou, Y. R., and Rajewsky, K. (1993). Independent control of immunoglobulin switch recombination at individual switch regions evidenced through Cre-loxP-mediated gene targeting. Cell *73*, 1155-1164.

2. Gu, H., Marth, J. D., Orban, P. C., Mossmann, H., and Rajewsky, K. (1994). Deletion of a DNA polymerase beta gene segment in T cells using cell type-specific gene targeting. Science *265*, 103-106. https://doi.org/10.1126/science.8016642.

3. Nagy, A. (2000). Cre recombinase: the universal reagent for genome tailoring. Genesis *26*, 99-109.

4. Sternberg, N., Hamilton, D., Austin, S., Yarmolinsky, M., and Hoess, R. (1981). Site-specific recombination and its role in the life cycle of bacteriophage P1. Cold Spring Harb Symp Quant Biol *45 Pt 1*, 297-309.

5. Sauer, B., and Henderson, N. (1988). Site-specific DNA recombination in mammalian cells by the Cre recombinase of bacteriophage P1. Proc Natl Acad Sci U S A *85*, 5166-5170.

6. Orban, P. C., Chui, D., and Marth, J. D. (1992). Tissue- and site-specific DNA recombination in transgenic mice. Proc Natl Acad Sci U S A *89*, 6861-6865.

7. Lakso, M., Sauer, B., Mosinger, B. J., Lee, E. J., Manning, R. W., Yu, S. H., Mulder, K. L., and Westphal, H. (1992). Targeted oncogene activation by site-specific recombination in transgenic mice. Proc Natl Acad Sci U S A *89*, 6232-6236.

8. Feil, R., Wagner, J., Metzger, D., and Chambon, P. (1997). Regulation of Cre recombinase activity by mutated estrogen receptor ligand-binding domains. Biochem Biophys Res Commun *237*, 752-757. https://doi.org/10.1006/bbrc.1997.7124.

9. Tian, X., Pu, W. T., and Zhou, B. (2015). Cellular Origin and Developmental Program of Coronary Angiogenesis. Circ Res *116*, 515-530. https://doi.org/10.1161/CIRCRESAHA.116.305097.

10. Tian, X., He, L., Liu, K., Pu, W., Zhao, H., Li, Y., Liu, X., Tang, M., Sun, R., Fei, J., et al. (2020). Generation of a self-cleaved inducible Cre recombinase for efficient temporal genetic manipulation. EMBO J *39*, e102675. https://doi.org/10.15252/embj.2019102675.

11. Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter, R. D., Hawrylycz, M. J., Jones, A. R., et al. (2010). A robust and high-

throughput Cre reporting and characterization system for the whole mouse brain. Nat

Neurosci *13*, 133-140. https://doi.org/10.1038/nn.2467.

12. Snippert, H. J., van der Flier, L. G., Sato, T., van Es, J. H., van den Born, M., Kroon-

Veenboer, C., Barker, N., Klein, A. M., van Rheenen, J., Simons, B. D., et al. (2010).

Intestinal crypt homeostasis results from neutral competition between symmetrically

dividing Lgr5 stem cells. Cell *143*, 134-144. https://doi.org/10.1016/j.cell.2010.09.016.

13. Becher, B., Waisman, A., and Lu, L. F. (2018). Conditional Gene-Targeting in Mice:

Problems   and   Solutions.   Immunity   *48*,   835-836.

https://doi.org/10.1016/j.immuni.2018.05.002.

14. Brash, J. T., Bolton, R. L., Rashbrook, V. S., Denti, L., Kubota, Y., and Ruhrberg, C.

(2020). Tamoxifen-Activated CreERT Impairs Retinal Angiogenesis Independently of

Gene Deletion. Circ Res *127*, 849-850. https://doi.org/10.1161/CIRCRESAHA.120.317025.

15. Rashbrook, V. S., Brash, J. T., and Ruhrberg, C. (2022). Cre toxicity in mouse models of

cardiovascular   physiology   and   disease.   Nat   Cardiovasc   Res   *1*,   806-816.

https://doi.org/10.1038/s44161-022-00125-6.

16. Lao, Z., Raju, G. P., Bai, C. B., and Joyner, A. L. (2012). MASTR: a technique for mosaic

mutant analysis with spatial and temporal control of recombination using conditional

floxed alleles in mice. Cell Rep *2*, 386-396. https://doi.org/10.1016/j.celrep.2012.07.004.

17. Wang, Q., Lin, Y. Y., Zhang, B., Wu, J., Roy, S., Ratiu, J. J., Xu, Y., Dai, M., Hale, L. P.,

Xiong, Y., et al. (2020). A mosaic analysis system with Cre or Tomato expression in the

mouse.   Proc   Natl   Acad   Sci   U   S   A   *117*,   28212-28220.

https://doi.org/10.1073/pnas.2014308117.

18. Fernández-Chacón, M., Casquero-García, V., Luo, W., Francesca Lunella, F., Ferreira

Rocha, S., Del Olmo-Cabrera, S., and Benedito, R. (2019). iSuRe-Cre is a genetic tool to

reliably induce and report Cre-dependent genetic modifications. Nat Commun *10*, 2262.

https://doi.org/10.1038/s41467-019-10239-4.

19. Liu, Q., Yang, R., Huang, X., Zhang, H., He, L., Zhang, L., Tian, X., Nie, Y., Hu, S., Yan,

Y., et al. (2016). Genetic lineage tracing identifies in situ Kit-expressing cardiomyocytes.

Cell Res *26*, 119-130. https://doi.org/10.1038/cr.2015.143.

20. Gribben, C., Sancho, R., and Behrens, A. (2023). Response to Magenheim et al.: Ductal
Ngn3-expressing progenitors contribute to adult beta cell neogenesis in the pancreas. Cell
Stem Cell *30*, 345-347. https://doi.org/10.1016/j.stem.2023.03.002.

21. Furuyama, K., Kawaguchi, Y., Akiyama, H., Horiguchi, M., Kodama, S., Kuhara, T.,
Hosokawa, S., Elbahrawy, A., Soeda, T., Koizumi, M., et al. (2011). Continuous cell
supply from a Sox9-expressing progenitor zone in adult liver, exocrine pancreas and
intestine. Nat Genet *43*, 34-41. https://doi.org/10.1038/ng.722.

22. Xiao, X., Chen, Z., Shiota, C., Prasadan, K., Guo, P., El-Gohary, Y., Paredes, J., Welsh, C.,
Wiersch, J., and Gittes, G. (2013). No evidence for beta cell neogenesis in murine adult
pancreas. J Clin Invest *123*, 2207-2217. https://doi.org/10.1172/JCI66323.

23. Tarlow, B. D., Finegold, M. J., and Grompe, M. (2014). Clonal tracing of Sox9+ liver
progenitors in mouse oval cell injury. Hepatology *60*, 278-289.
https://doi.org/10.1002/hep.27084.

24. He, L., Nguyen, N. B., Ardehali, R., and Zhou, B. (2020). Heart Regeneration by
Endogenous Stem Cells and Cardiomyocyte Proliferation: Controversy, Fallacy, and
Progress. Circulation *142*, 275-291.
https://doi.org/10.1161/CIRCULATIONAHA.119.045566.

25. Liu, Q., Liu, K., Cui, G., Huang, X., Yao, S., Guo, W., Qin, Z., Li, Y., Yang, R., Pu, W., et
al. (2019). Lung regeneration by multipotent stem cells residing at the bronchioalveolar-
duct junction. Nat Genet *51*, 728-738. https://doi.org/10.1038/s41588-019-0346-6.

26. He, L., Li, Y., Li, Y., Pu, W., Huang, X., Tian, X., Wang, Y., Zhang, H., Liu, Q., Zhang, L.,
et al. (2017). Enhancing the precision of genetic lineage tracing using dual recombinases.
Nat Med *23*, 1488-1498. https://doi.org/10.1038/nm.4437.

27. Anastassiadis, K., Fu, J., Patsch, C., Hu, S., Weidlich, S., Duerschke, K., Buchholz, F.,
Edenhofer, F., and Stewart, A. F. (2009). Dre recombinase, like Cre, is a highly efficient
site-specific recombinase in E. coli, mammalian cells and mice. Dis Model Mech *2*, 508-
515. https://doi.org/10.1242/dmm.003087.

28. Han, M., Liu, Z., Liu, L., Huang, X., Wang, H., Pu, W., Wang, E., Liu, X., Li, Y., He, L.,
et al. (2023). Dual genetic tracing reveals a unique fibroblast subpopulation modulating
cardiac fibrosis. Nat Genet *55*, 665-678. https://doi.org/10.1038/s41588-023-01337-7.

29.   Zhang, H., Pu, W., Li, G., Huang, X., He, L., Tian, X., Liu, Q., Zhang, L., Wu, S. M.,
719   Sucov, H. M., et al. (2016). Endocardium Minimally Contributes to Coronary Endothelium
720   in the Embryonic Ventricular Free Walls. Circ Res *118*, 1880-1893.
721   https://doi.org/10.1161/CIRCRESAHA.116.308749.

722   30.   Huelsken, J., Vogel, R., Erdmann, B., Cotsarelis, G., and Birchmeier, W. (2001). beta-
723   Catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. Cell
724   *105*, 533-545.

725   31.   Pontes-Quero, S., Heredia, L., Casquero-García, V., Fernández-Chacón, M., Luo, W.,
726   Hermoso, A., Bansal, M., Garcia-Gonzalez, I., Sanchez-Muñoz, M. S., Perea, J. R., et al.
727   (2017). Dual ifgMosaic: A Versatile Method for Multispectral and Combinatorial Mosaic
728   Gene-Function Analysis. Cell *170*, 800-814.e18. https://doi.org/10.1016/j.cell.2017.07.031.

729   32.   Kim, G. B., Rincon Fernandez Pacheco, D., Saxon, D., Yang, A., Sabet, S., Dutra-Clarke,
730   M., Levy, R., Watkins, A., Park, H., Abbasi Akhtar, A., et al. (2019). Rapid Generation of
731   Somatic Mouse Mosaics with Locus-Specific, Stably Integrated Transgenic Elements. Cell
732   *179*, 251-267.e24. https://doi.org/10.1016/j.cell.2019.08.013.

733   33.   Zong, H., Espinosa, J. S., Su, H. H., Muzumdar, M. D., and Luo, L. (2005). Mosaic
734   analysis with double markers in mice. Cell *121*, 479-492.
735   https://doi.org/10.1016/j.cell.2005.02.012.

736   34.   Liu, C., Sage, J. C., Miller, M. R., Verhaak, R. G., Hippenmeyer, S., Vogel, H., Foreman,
737   O., Bronson, R. T., Nishiyama, A., Luo, L., et al. (2011). Mosaic analysis with double
738   markers reveals tumor cell of origin in glioma. Cell *146*, 209-221.
739   https://doi.org/10.1016/j.cell.2011.06.014.

740   35.   Li, Y., He, L., Huang, X., Issa Bhaloo, S., Zhao, H., Zhang, S., Pu, W., Tian, X., Li, Y., Liu,
741   Q., et al. (2018). Genetic Lineage Tracing of Non-Myocyte Population by Dual
742   Recombinases. Circulation *138*, 793-805.
743   https://doi.org/10.1161/CIRCULATIONAHA.118.034250.

744   36.   Liu, K., Tang, M., Jin, H., Liu, Q., He, L., Zhu, H., Liu, X., Han, X., Li, Y., Zhang, L., et al.
745   (2020). Triple-cell lineage tracing by a dual reporter on a single allele. J Biol Chem *295*,
746   690-700. https://doi.org/10.1074/jbc.RA119.011349.

747 37. Zhang, H., Pu, W., Liu, Q., He, L., Huang, X., Tian, X., Zhang, L., Nie, Y., Hu, S., Lui, K.
748    O., et al. (2016). Endocardium Contributes to Cardiac Fat. Circ Res *118*, 254-265.
749    https://doi.org/10.1161/CIRCRESAHA.115.307202.

750 38. Paxinou, E., Chen, Q., Weisse, M., Giasson, B. I., Norris, E. H., Rueter, S. M.,
751    Trojanowski, J. Q., Lee, V. M., and Ischiropoulos, H. (2001). Induction of alpha-synuclein
752    aggregation by intracellular nitrative insult. J Neurosci *21*, 8053-8061.
753    https://doi.org/10.1523/JNEUROSCI.21-20-08053.2001.

754 39. Jiang, P., Ren, L., Zhi, L., Hu, X., and Xiao, R. P. (2021). Protocol for cell preparation and
755    gene delivery in HEK293T and C2C12 cells. STAR Protoc *2*, 100497.
756    https://doi.org/10.1016/j.xpro.2021.100497.

757 40. Pu, W., Zhu, H., Zhang, M., Pikiolek, M., Ercan, C., Li, J., Huang, X., Han, X., Zhang, Z.,
758    Lv, Z., et al. (2023). Bipotent transitional liver progenitor cells contribute to liver
759    regeneration. Nat Genet *55*, 651-664. https://doi.org/10.1038/s41588-023-01335-9.

760 41. Charni-Natan, M., and Goldstein, I. (2020). Protocol for Primary Mouse Hepatocyte
761    Isolation. STAR Protoc *1*, 100086. https://doi.org/10.1016/j.xpro.2020.100086.

762 42. Pu, W., Zhang, H., Huang, X., Tian, X., He, L., Wang, Y., Zhang, L., Liu, Q., Li, Y., Li, Y.,
763    et al. (2016). Mfsd2a+ hepatocytes repopulate the liver during injury and regeneration. Nat
764    Commun *7*, 13369. https://doi.org/10.1038/ncomms13369.

Figure legends

Figure 1. Design of roxCre for DreER-induced mCre expression

(A) A schematic showing genetic labeling and/or gene knockout in CreER-expressing cells.

(B, D, F) Strategies for DreER-induced Cre expression.

(C) Crossing with R26-tdT mice, R26-RSR-Cre exhibits leakiness in E13.5 embryos. Scale bars, yellow 1mm. Each image is representative of 5 individual biological samples.

(E) Examination of Cre leakiness by whole-mount fluorescence images of organs collected from R26-RSR-Cre2;R26-26-GFP adult mice. Scale bars, yellow 1mm. Each image is representative of 5 individual biological samples.

(G) Examination of Cre leakiness by whole-mount fluorescence images of organs collected from R26-R-reverseCre-R;R26-tdT adult mice. Scale bars, yellow 1mm. Each image is representative of 5 individual biological samples.

(H) A schematic showing the design for roxCre and mCre.

(I) A schematic showing mCre1 to mCre12 by insertion of rox into Cre cDNA at 12 different loci. Two dual-recombinase designs for inducible Cre expression once tried before.

(J) Examination of mCre for recombination efficiency by the *loxP-Stop-loxP-GFP* reporter.

(K) Immunofluorescence images of cells stained with GFP on cells. Scale bars, white, 100 µm. Each image is representative of 5 individual biological samples.

(L) Quantification of the percentage of cells expressing GFP in each group. Data are the means ± SEM; n = 5.

Figure 2. DreER-induced mCre robustly recombines inert alleles

(A) A schematic showing the experimental design to test the recombination efficiency of mCre1 and mCre7 on the *R26-Confetti* allele. Tam-induced DreER-rox recombination leads to mCre1/tdT or mCre7/GFP expression in hepatocytes in strategy 1 or 2, respectively. Strategy 3 uses conventional *Alb-CreER* as control. YFP and mCFP signals are used for the examination of recombination on the *R26-Confetti* allele.

(B) Fluorescence images of YFP and mCFP on liver sections collected from mice in strategy 1-3. Strategy 1 and 3 exhibit tdT or GFP in hepatocytes after DreER-rox recombination, respectively. Scale bars, 100 µm. Each image is representative of 5 individual biological samples.

(C) Quantification of the percentage of hepatocytes (Heps) expressing either YFP and/or mCFP

in strategy 4-6. Data are the means ± SEM; n = 11 mice for each strategy; ****$P < 0.0001$ by

one-way ANOVA.

(D) A schematic showing the experimental strategy to test the recombination efficiency of

mCre4 and mCre10 on *R26-Confetti* allele. Tam-induced recombination leads to mCre4/tdT or

mCre10/GFP expression in endothelial cells (ECs) in strategy 4 or 5, respectively. Strategy 6

uses conventional *Cdh5-CreER* as control. YFP and mCFP signals are used for the examination

of recombination on *R26-Confetti* allele.

(E) Fluorescence images of YFP and mCFP on intestinal sections collected from mice in strategy

4-6. Strategy 4 and 5 exhibit tdT or GFP in ECs after DreER-rox recombination, respectively.

Scale bars, 100 µm. Each image is representative of 5 individual biological samples.

(F) Quantification of the percentage of ECs expressing either YFP and/or mCFP in strategy 4-6.

Data are the means ± SEM; n = 11 mice for each strategy; ****$P < 0.0001$ by one-way ANOVA.

Figure 3. Dre-induced mCre efficiently deletes genes

(A) A schematic showing the experimental design for Dre-induced mCre expression and the

subsequent gene deletion.

(B) A schematic showing the experimental strategy.

(C and D) qRT-PCR analysis of the relative expression of *Ctnnb1* (C) and Glul (D) in the tdT$^+$

hepatocytes. Data are the means ± SEM; n = 5; ****$P < 0.0001$ by student's *t*-test.

(E) A schematic showing the experimental strategy.

(F) Immunostaining for tdT, β-Catenin, GS, and E-CAD on liver sections collected on the day 3

post-Tam. PV, portal vein; CV, central vein. Scale bars, yellow 1mm; white 100 µm. Each image

is representative of 5 individual biological samples.

(G) Immunostaining for tdT, β-Catenin, GS, and E-CAD on liver sections collected at week 4

post-Tam. Scale bars, yellow 1mm; white 100 µm. Each image is representative of 5 individual

biological samples.

(H) A schematic showing the experimental strategy.

(I) Western blotting of β-Catenin and GAPDH in tdT$^+$ cells.

(J) Quantification of the relative expression of β-Catenin protein. Data are the means ± SEM; n =

3. **$P < 0.01$ by student's *t* test.

(K) qRT-PCR analysis of the relative expression of *Ctnnb1* in the tdT$^+$ cells. Data are the means

± SEM; n = 6. ****$P < 0.0001$ by student's *t* test.

(L) qRT-PCR analysis of the relative expression of *Glul*, *Axin2*, *Cyp1a2*, *Cyp2e1*, *Oat*, *Tcf7*,

Lect2, *Tbx3*, *Slc1a2,* and *Rhbg* in the tdT$^+$ cells. Data are the means ± SEM; n = 6. ***$P < 0.001$,

****$P < 0.0001$ by student's *t* test.

Figure 4. *R26-loxCre-tdT* is efficiently recombined by CreER

(A) A schematic showing the knock-in strategy for the generation of the *R26-loxCre-tdT* allele.

In this line, tdT is used to denote the mCre expression after the removal of *Stop*.

(B) A schematic showing the experimental strategy 1 to examine the leakiness of *R26-loxCre-*

tdT mice.

(C) A schematic showing the experimental strategy 2 to test mCre/tdT expression by AAV8-

TBG-Cre.

(D) Immunostaining for tdT on tissue sections shows tdT expression in the liver and pancreas

when recombination was intiated by AAV8-TBG-Cre. Scale bars, white 100 µm. Each image is

representative of 5 individual biological samples.

(E) A schematic showing the experimental strategies 3 and 4 for comparing the CreER-mediated

first recombination efficiency between *Cdh5-CreER;R26-tdT* and *Cdh5-CreER;R26-loxCre-tdT*

mice.

(F) Immunostaining for tdT and VE-Cad on liver sections collected on day 7 post-Tam. Scale

bars, white 100 µm. Each image is representative of 5 individual biological samples.

(G) Quantification of the percentage of VE-Cad$^+$ ECs expressing tdT. Data are the means ± SEM;

n= 6. ****$P < 0.0001$ by student's *t*-test.

Figure 5. *R26-loxCre-tdT* enables CreER to recombine *R26-Confetti* efficiently

(A) Schematics showing the experimental design. In *Cdh5-CreER;R26-loxCre-tdT;R26-Confetti*

mice, Tam-induced CreER-loxp recombination switches *R26-loxCre-tdT* into *R26-mCre-tdT*

allele with simultaneous tdT labeling and expression of mCre, which subsequently targets *R26-*

Confetti (right panel). The conventional *Cdh5-CreER;R26-Confetti* mice are used as control.

(B) A schematic showing the experimental strategy.

(C) Whole-mount YFP and mCFP fluorescence images of retina collected from two mice groups.

Scale bars, white 100 µm. Each image is representative of 5 individual biological samples.

(D) Immunofluorescence images of YFP, mCFP, and VE-Cad on tissue sections show

significantly more YFP$^+$ and/or mCFP$^+$ ECs in the *Cdh5-CreER;R26-loxCre-tdT;R26-Confetti*

mice compared with that of *Cdh5-CreER;R26-Confetti* mice (left panel). The right panel shows

the quantification of ECs expressing YFP and/or mCFP. Data are the means ± SEM; n = 5.

****$P < 0.0001$ by student's *t*-test. Scale bars, white 100 µm. Each image is representative of 5

individual biological samples.

Figure 6. *R26-loxCre-tdT* enables *Alb-CreER* to efficiently knockout genes in hepatocytes

(A) Schematics showing the experimental designs for Ctnnb1 gene knockout using either *Alb-*

CreER or *Alb-CreER;R26-loxCre-tdT* mice.

(B) A schematic showing the experimental strategy. *Alb-CreER;R26-tdT2;Ctnnb1$^{flox/flox}$* and *Alb-*

CreER;R26-loxCre-tdT;Ctnnb1$^{flox/+}$ mice were injected with Tam for 5 times, while *Alb-*

CreER;R26-loxCre-tdT;Ctnnb1$^{flox/flox}$ mice were injected with Tam for one time.

(C) Immunostaining for tdT, β-Catenin, GS, and E-CAD on liver sections. Scale bars, yellow

1mm; white 100 µm. Each image is representative of 5 individual biological samples.

(D) Immunostaining for tdT and CYP2E1 on liver sections. Scale bars, yellow 1mm; white 100

µm. Each image is representative of 5 individual biological samples.

(E) Western blotting of β-Catenin and β-Actin expression in hepatocytes, n = 3.

(F) Quantification of β-Catenin expression. Data are the means ± SEM; n = 5.

(G) qRT-PCR analysis of the relative expression of *Ctnnb1* in the hepatocytes.

(H) qRT-PCR analysis of the relative expression of *Glul*, *Axin2*, *Cyp2e1*, *Oat*, *Tcf7*, *Lect2*, *Tbx3*,

and *Slc1a2* in hepatocytes. Data are the means ± SEM; n = 7; ns, non-significant; *$P < 0.05$, **P

< 0.01, ***$P < 0.001$, ****$P < 0.0001$ by one-way ANOVA.

Figure1

Figure2

Figure3

A *R26-loxCre-tdT* knock-in strategy

B Strategy 1

C Strategy 2

D Strategy 1 | Strategy 2

E Comparison of recombination efficiency

Cdh5-CreER

F Strategy 3 | Strategy 4

G Strategy 3 | Strategy 4

Figure4

Figure5

Figure6