Rudhira-mediated microtubule stability controls TGFβ signaling during mouse vascular

development.

Divyesh Joshi[1#], Preeti Jindal[1#], Ronak Shetty[1] and Maneesha S. Inamdar[1,2*]

[1]Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore, India;. [2]Institute for Stem

Cell Science and Regenerative Medicine (inStem), Bangalore, India.

# Equal contribution

*Author for correspondence

Email: inamdar@jncasr.ac.in

Tel: +91-080-2208-2818

ORCID: 0000-0002-8243-2821

Running title: Rudhira bridges microtubule stability and TGFβ signaling.

Keywords: Rudhira; BCAS3; Microtubule cytoskeleton stability; Smad2/3; TGFβ signaling

Abstract

The Transforming Growth Factor β (TGFβ) signaling pathway is critical for survival, proliferation, and cell migration, and is tightly regulated during cardiovascular development. Smads, key effectors of TGFβ signaling, are sequestered by microtubules (MTs) and need to be released for pathway function. Independently, TGFβ signaling also stabilizes MTs. Molecular details and the *in vivo* relevance of this cross-regulation remain unclear, understanding which is important in complex biological processes such as cardiovascular development. Here, we use *rudhira/Breast Carcinoma Amplified Sequence 3 (BCAS3)*, a MT-associated, endothelium-restricted and developmentally essential proto-oncogene, as a pivot to decipher cellular mechanisms in bridging TGFβ signaling and MT stability. We show that Rudhira regulates TGFβ signaling *in vivo,* during mouse cardiovascular development, and in endothelial cells in culture. Rudhira associates with MTs and is essential for the activation and release of Smad2/3 from MTs. Consequently, Rudhira depletion attenuates Smad2/3-dependent TGFβ signaling thereby impairing cell migration. Interestingly, Rudhira is also a transcriptional target of Smad2/3-dependent TGFβ signaling essential for TGFβ-induced MT stability. Our study identifies an immediate early physical role and a slower, transcription-dependent role for Rudhira in cytoskeleton- TGFβ signaling crosstalk. These two phases of control could facilitate temporally- and spatially restricted targeting of the cytoskeleton and/or TGFβ signaling in vascular development and disease.

Significance statement

The developmental TGFβ pathway is essential for cell migration, cell-cell communication, adhesion, apoptosis, and matrix remodeling. Dysregulation of TGFβ signaling leads to aberrant vascular patterning and angiogenesis during mouse embryogenesis. Pathway activation involves phosphorylation and nuclear transport of Smads. Microtubules sequester Smads in the cytoplasm, thereby inhibiting TGFβ signaling. Conversely, TGFβ signaling stabilizes microtubules. However, the molecular components involved, and biological relevance of this cross-regulation remain unclear. We show that the oncoprotein Rudhira/BCAS3 facilitates Smad-MT dissociation upon ligand-mediated TGFβ receptor activation. Interestingly, Smad-dependent TGFβ signaling activation enhances rudhira transcription, essential for microtubule stabilization during cardiovascular development. This dual regulation of TGFβ signaling and microtubule stability by Rudhira allows sustained temporal control essential for development, and its dysregulation has pathological outcomes.

Introduction

Cytoskeletal organization and growth factor signaling are mutually dependent and coordinate multiple cellular processes [1]. The multifunctional Transforming growth factor β (TGFβ) pathway regulates cell proliferation, differentiation, survival and migration and is sensitive to cytoskeletal rearrangements [1-3]. Depletion of TGFβ pathway components causes developmental abnormalities often leading to embryonic death [4]. Ligand binding activates the TGFβ receptor, causing receptor-mediated phosphorylation, followed by nuclear translocation of Smad proteins, resulting in transcriptional regulation of target gene expression [5]. Smads, the effectors of the TGFβ pathway, are competitively bound by several proteins and are a hub for regulation of TGFβ signaling and its intersecting pathways [6]. MTs bind to and sequester Smad2/3 and Smad4, thereby negatively regulating TGFβ signaling, which is relieved upon TGFβ stimulation [7]. Conversely, TGFβ signaling stabilizes MTs [8, 9] through an as yet unclear mechanism, suggestive of a feedback loop.

During development, TGFβ signaling is activated in endothelial cells (ECs) of the remodeling angiogenic endothelium and is critical for vascular development [10]. MT stability is also essential for angiogenic sprouting in ECs [11]. MTs can be stabilized in multiple ways within the cell, initially by RhoA-mDia actin bundling and thereafter by interaction with multi-protein complexes. Serum components, TGFβ and Lysophosphatidic acid (LPA), have been shown to stabilize MTs [8, 12]. While LPA stabilizes MTs within 30 min post-stimulation and generically [12], the effect of TGFβ signaling takes at least 2 hours [8] but is more restricted to and essential for endothelial cell function *in vivo*. Slow and restricted induction of MT stability by TGFβ as compared to the more generic and rapid function of LPA suggests a requirement for new transcription/translation which may be developmentally important. However, the molecular mechanisms and biological relevance of cytoskeletal interactions with TGFβ signaling remain elusive. Characterization of tissue-restricted regulators of the ubiquitous MT cytoskeleton may aid in defining its specialized context- dependent functions.

In this study, we address the role of the MT-interacting protein Rudhira in connecting TGFβ signaling and MT stability. Rudhira associates with MTs and intermediate filaments (IFs), stabilizes MTs and

promotes directional cell migration [11, 13]. *Rudhira* knockout in mouse causes mid-gestation

lethality due to aberrant cardiovascular patterning [14]. Transcriptome analysis showed that

several processes were deregulated in the *rudhira* knockout, especially, a large number of TGFβ

signaling pathway components [14]. This suggests that Rudhira may regulate TGFβ signaling for

vascular development.

Here, we show that Rudhira-depleted cells have reduced Smad2/3 activation during mouse

embryonic development, indicating that Rudhira is a positive regulator of TGFβ signaling. We

demonstrate that Rudhira facilitates TGFβ-mediated release of Smad2/3 from MTs. Further, TGFβ

signaling stabilizes MTs by inducing transcription, in a Rudhira-dependent manner. Thus, Rudhira

bridges TGFβ signaling and MT stability, required for cardiovascular development. Ectopic and

aberrant Rudhira expression is seen in several carcinomas and positively correlates with tumor

metastatic potential [15-17]. Hence our study will aid understanding of TGFβ signaling and suggest

novel ways to regulate it in development and disease.

Results

Rudhira depletion attenuates Smad2/3-dependent TGFβ signaling *in vivo* and *in vitro*

Rudhira is an essential gene, critical for mouse cardio-vascular development [14]. Ubiquitous

(*rudh*$^{-/-}$) or endothelial (*rudh*^{CKO}) knockout of *rudhira* results in angiogenic defects and mid-

gestation embryonic lethality [14]. Transcripts of several components of the TGFβ pathway are

deregulated upon Rudhira depletion and the knockout phenocopies mutants of several TGFβ

pathway components, whose primary phenotype is in the cardiovascular system [10]. Hence, we

characterized the role of Rudhira in TGFβ signaling. Analysis of the molecular interaction, reaction

and relation networks by KEGG pathway mapping indicated that multiple components of the

TGFβ pathway were deregulated in the absence of Rudhira, affecting several processes important

for cardiovascular development (Fig. 1A). To validate this, we used non-silencing (NS) control and

rudhira knockdown (KD) saphenous vein endothelial cell lines (SVEC) as described before (Fig

S1A, B) [14]. To test the pathway regulation at the level of receptors, we analysed transcript and

basal as well as active protein levels. While we observed some reduction in transcript levels of

tgfbrI and *tgfbrII* (Fig. S2A), the basal and active protein levels of TGFβRI remained unchanged in
KD cells (Fig. S2B). Interestingly, a constitutively active form of TGFβRI (TGFβRIT204D), expressed
under the CMV promoter, was also suppressed in Rudhira KD (overexpression achieved in NS ~200
folds while in KD ~80 folds) (Fig. S2C). TGFBR1 protein has a short half-life, while the mRNA is
relatively stable [18]. Reduced amount of relatively stable RNA may be sufficient to produce
adequate protein levels. These increased levels of active TGFβRI in KD were unable to restore
Smad2 phosphorylation, unlike NS controls, which showed a clear increase (Fig. 1B). While
important to note that Rudhira or its loss may regulate *tgfbr* transcript levels, these results support
a receptor-independent role of Rudhira in regulating TGFβ pathway.

To probe further into the molecular mechanism by which Rudhira controls TGFβ signaling, we used
rudhira knockout mice and the KD endothelial cell line. Upon TGFβ-stimulation ECs in culture showed
reduced Smad2 activation in the KD as compared to NS control (Fig. 1C). Further, qRT-PCR analysis
showed that transcript levels of Smad2/3 targets, namely *mmp9* and *pai1*, did not increase
significantly in KD, whereas NS control showed >2-fold increase for both (Fig. 1D). Expectedly, partial
restoration of Rudhira level in KD cells restored Smad2 activation and target gene expression (Fig.
1C, D). The canonical TGFβ pathway also responds to BMP4 activation. However, upon stimulation
with BMP4 there was no significant difference in the levels of pSmad1/5/8 between NS and KD (Fig.
S2D). These data indicate that Rudhira is required for TGFβ-dependent Smad2 phosphorylation and
pathway activation.

Endothelial depletion of Rudhira (*rudh^{CKO}*) is sufficient to cause vascular defects [14] and functional
TGFβ signaling is essential for vascular development [19]. Expectedly, *rudh^{CKO}* embryos showed a
concomitant loss of pSmad2/3 in the embryonic endocardium (Fig. 1E, E'), without affecting the
total Smad2/3 level (Fig. S2E, E'). Thus, Rudhira regulates TGFβ signaling in a Smad2/3-dependent
manner *in vivo*, which is essential for angiogenesis and endothelial cell migration during cardio-
vascular development.

Rudhira is required for TGFβ mediated cell migration

As Rudhira is essential for cell migration, we tested the effect of altered TGFβ pathway activity in
Rudhira KD endothelial cells (SVEC) in a transwell migration assay. We also validated our results in
HEK293 cells as these allow a high efficiency of transfection and Rudhira overexpression, for

migration assays. TGFβ addition did not enhance migration in *rudhira* knockdown (KD) cells as

compared to controls (NS and rescued KD) (Fig. 2A). In addition, ectopic Rudhira expression in

HEK293 cells caused increase in migration rate as compared to the control, consistent with our

earlier study [13] (Fig. 2B,C).

TGFβ supplementation of Rudhira overexpressing cells dramatically enhanced the rate of cell

migration (~2.6 fold), suggesting a direct correlation between Rudhira levels and pathway

function (Fig. 1C, D). Serum contains TGFβ in a concentration range of 10-40 ng/ml [20] and

we used 5% FBS with or without additional TGFβ in the transwell migration assay, since purified

TGFβ alone is unable to induce migration. The ratio of increase in migration rates with or without

exogenous TGFβ addition was similar in both the control and Rudhira-overexpressing cells (~1.5

fold). Interestingly, TGFβ pathway-inhibition by SB431542 (SB) restored the effect of Rudhira

overexpression (enhancement in cell migration rate) to control levels (Fig. 2B, S3A) suggesting

that Rudhira function is regulated by the TGFβ pathway downstream of receptor activation. In

addition to the multiple molecular pathways that function in concert to control cell migration [21],

our data support a role for Rudhira in TGFβ-dependent cell migration.

Rudhira regulates TGFβ signaling by promoting the release of Smad2/3 from MTs

Since Rudhira localizes to the cytoskeletal components, namely microtubules and intermediate

filaments [13], we hypothesized that it regulates TGFβ signaling at the cytoskeleton. However, we

could not detect any interaction between Rudhira and Smad2/3 under the conditions tested (Fig.

3A), suggesting that the effect on the pathway may be indirect, possibly through MTs. TGFβ

signaling is controlled at multiple levels, including cytoskeletal regulation at MTs, which interact

with and sequester Smad2/3 thereby inhibiting the TGFβ pathway [7]. TGFβ stimulation releases

Smads from MT sequestration in multiple cell types including endothelial cells, facilitating its

activation and nuclear translocation [7]. Expectedly, TGFβ stimulation in ECs (confirmed by increase

in pSmad2 levels) reduced Smad-MT binding as seen by co-immunoprecipitation for β-Tubulin

and immunoblotting for Smad2 (Fig. 3B).

To assess whether Rudhira is required for TGFβ-dependent release of Smad from MTs, we tested

for β-Tubulin and Smad interaction upon TGFβ stimulation in NS and KD cells by performing *in situ*

Proximity Ligation assay (PLA) [22]. While NS cells showed reduced Smad-MT interaction, KD cells

retained Smad-MT interaction even upon TGFβ stimulation in multiple KD cell lines (Fig. 3C, Fig. S3B).

Interestingly, TGFβ stimulation resulted in an increase in Rudhira-MT association in wild type

cells, detected by colocalization (Fig. 3D) and confirmed by PLA (Fig. 3E). Thus TGFβ-stimulated

Rudhira-MT association makes MTs unavailable for Smad binding. However, MTs are abundant and

present as subsets, differing in their Tubulin isotype and isoform composition, post translational

modifications (PTMs) and stability [23], which may also govern their interactions with other

molecules. It is unlikely that Rudhira binds to all MTs present. Also, it is likely that only a subset of

MTs may bind and sequester Smads.

Rudhira is a transcriptional target of TGFβ signaling and is essential for MT stability

The TGFβ pathway is auto-regulatory, due to which several pathway components are induced

or inhibited [24, 25]. The _rudhira_ promoter harbors putative Smad-binding elements (SBEs) (Fig. S4A).

It also harbors putative binding sites of major TGFβ pathway regulated transcription factors (TFBS)

(Fig. S4B, Table S2). Expectedly, TGFβ stimulation increased _rudhira_ mRNA and protein levels (Fig.

4A, B, C). Knockdown of _smad2_ or _smad3_ resulted in significant reduction of Rudhira levels,

indicating that _rudhira_ is a Smad2/3-dependent TGFβ target (Fig. 4D). Blocking transcription

with Actinomycin D (AcD) or translation using Cycloheximide (CHX) prevented increase in _rudhira_

levels upon TGFβ stimulation, confirming that _rudhira_ is a transcriptional target of the Smad2/3-

dependent TGFβ pathway (Fig. 5A, S4C). However, additional investigation is required to confirm

whether _rudhira_ is a direct target of Smad2/3-dependent TGFβ pathway.

Rudhira and TGFβ signaling, both can induce MT stabilization [8, 11]. Although the mechanism

is unclear, PTMs and/or transcription may control TGFβ-induced MT stability [8]. Since _in vivo_

studies indicated that _rudhira_ deletion severely affects the TGFβ pathway [14], we tested whether

TGFβ- induced transcription is required for MT stability. TGFβ stimulation results in a slow

enhancement of MT stability over a period of several hours, allowing sufficient time for

transcriptional regulation [8]. Interestingly, we find that blocking transcription or translation,

prevented TGFβ-dependent MT stabilization, as detected by Glu-tubulin levels (Fig. 5A, S4C),

correlating well with Rudhira levels and validated by increased sensitivity to Nocodazole-mediated

MT depolymerization (Fig. 5B). These data show that TGFβ induced MT stability is transcription-

dependent.

TGFβ signaling stabilizes MTs in a Rudhira-dependent manner

Since *rudhira* transcription is dependent on TGFβ signaling and Rudhira is required for MT stability,

we further tested whether TGFβ-induced MT stability is Rudhira-dependent. Serum starvation

resulted in low detectable levels of Glu-tubulin in NS and KD cells (Fig. 5C). Stimulation with

TGFβ or serum led to increased Glu-tubulin level in NS but not in KD cells showing that TGFβ-

induced MT stability is dependent on Rudhira (Fig. 5C, D). To test whether Rudhira stabilizes MTs

in vivo also, we analyzed the effect of loss of Rudhira on MT stability in the developing

endocardium. Interestingly, along with reduced pSmads levels observed earlier (Fig. 1E, E'), we found

that Glu-tubulin positive stable MTs were fewer in *rudh^CKO* embryonic endocardium (Fig. 5D),

supporting a role of Rudhira in regulation of TGFβ signaling and MT stability during cardiovascular

development. Combined, our study suggests that Rudhira primarily affects MTs to modulate

their Smad-binding property and downstream TGFβ signaling. Consequently, Smad2/3-dependent

TGFβ signaling induces *rudhira* transcription leading to MT stabilization essential for cardiovascular

remodeling during development (Fig. 5E).

Discussion

Growth factor signaling and cytoskeletal remodeling are mutually dependent processes that require complex context-dependent molecular crosstalk [1]. Specifically, the developmentally essential, tissue-restricted TGFβ signaling pathway, regulates and is regulated by the MT cytoskeleton. However, the underlying molecular mechanism and *in vivo* relevance of this cross-regulation remain elusive. In this study, we probed the role of Rudhira in the cytoskeletal control of TGFβ signaling and, in turn, its effect on MT properties. Our study identifies a dual role of Rudhira during vascular development- an early role in MT-mediated regulation of Smad2/3 and a late role as a transcriptional target of TGFβ signaling required for MT stability.

While several processes important for cardiovascular development were deregulated in *rudhira^-/-* embryos, the TGFβ pathway was primarily affected [14]. TGFβ signaling is a complex pathway and is regulated at multiple levels by various feedback loops. One of the complicated yet delicate feedback circuits is the regulation of receptor levels and activity [25]. Rudhira depletion reduces transcript level without altering basal and active protein level of TGFBRI. This disparity in the transcript and protein level may arise from the fact that while the *tgfbr* transcript is very stable, the protein is quite unstable, and lower transcript levels may still be sufficient for generating adequate protein levels. Such a phenomenon is not unusual, particularly for membrane proteins, where mRNA levels may not correlate with protein levels. Rudhira, in a manner not yet understood, may also be a part of a broader feedback pathway as it further negatively regulates levels of TGFβ pathway inhibitors such as Smurfs and Smad7 [14], suggesting that additional regulatory mechanisms may operate.

TGFβ signaling and Rudhira promote cell migration in a synergistic and inter-dependent manner. Mechanistically, Rudhira promotes TGFβ-dependent release of Smads from MTs likely due to increased Rudhira-MT association. This could allow pathway activation and result in enhanced cell migration. Additional, confounding factors downstream of the autoregulatory TGFβ pathway or pathway targets, as described above, may also contribute to Rudhira dependent cell migration in response to TGFβ. Converse regulation of Rudhira-MT and Smad-MT interaction is indicative of probable competition between Smads and Rudhira for MT binding, which merits further investigation. Binding of Rudhira to MTs may also alter them in a way that makes MTs unavailable for Smad binding. Since TGFβ stimulation is known to stabilize microtubules, we hypothesize that

TGFβ stimulation increases Rudhira binding to stable microtubules. While unclear molecular events
involved in release of Smads from MTs, and MT stabilization by Rudhira make it difficult to assign
causal roles, this study supports the hypothesis that dissociation of Smads from MTs is essential for
Smad activation. In addition, though MTs are ubiquitously and abundantly present, our data
suggest that only a few MTs participate in Smad sequestration. This is in concordance to our
earlier study which shows that Rudhira associates preferentially with stable MTs [11].

The TGFβ signaling pathway is autoregulatory and many of its components are also targets of the
pathway [24]. The *rudhira* promoter harbors Smad Binding Elements (SBEs). We show that Rudhira,
in addition to being a regulator of the TGFβ pathway, is also a transcriptional target. However,
further analysis is required to identify the specific mechanisms by which Smad-dependent, TGFβ-
induced, *rudhira* transcription is regulated during development. In contrast to this positive feedback,
TGFβ pathway is also negatively self-regulated, allowing for return to basal state amenable for
reactivation. This self-suppression could possibly involve increased turnover of receptors and
Smad2/3, and increased expression of inhibitory Smads, that may recover responsiveness to TGFβ
stimulation. Additionally, in the context of MT-dependent Smad2/3 inhibition, the still short turnover
time of stable MTs (several minutes to hours) may also promote quick return to resting state. These
interesting possibilities can now be tested.

Rudhira expression is tightly regulated and restricted to the remodeling endothelium
during development [14]. Mechanistically, Rudhira promotes MT stability for angiogenic
remodeling [11]. Independent studies show that TGFβ signaling, and MT reorganization are
essential for cardiovascular development [26-28]. Deletion of *smad2/3* or *tubulins* leads to
embryonic lethality with severely defective embryonic and extra-embryonic vasculature [29].
However, the role of MT stability for angiogenic sprouting *in vivo* and the underlying molecular
mechanisms remain elusive. Our study suggests that endothelial cytoskeleton remodeling by
Rudhira and activation of TGFβ signaling during cardiovascular development are inter-dependent
and cross-regulated. TGFβ signaling may lead to restricted induction of MT stability by inducing
the expression of MT-stabilizing developmentally regulated proteins like Rudhira. Additional
cytoskeletal components such as Vimentin that interacts with Rudhira, and is also a target of TGFβ
signaling, are likely to be involved [11, 13]. Our analysis will help gain better insight into cellular
processes such as migration, which require maintenance of cytoskeletal stability and active

signaling for extended periods. An added advantage will be the ability to decipher deregulated
migration such as in tumor metastasis. While targeting MT-Smad interaction would allow
regulation of the TGFβ pathway, the ubiquitous nature of these molecules is likely to result
in widespread and possibly undesirable effects of therapeutic intervention. Rudhira expression,
being more restricted, could provide a suitable target to regulate developmental and pathological
TGFβ signaling.

Materials and methods

Animal experiments

Animal experiments were performed as mentioned earlier [14]. All animal experimental protocols
were approved by the Institutional Animal Ethics Committee (IAEC) of JNCASR (Project number
MSI006. All animals were maintained, and experiments were performed according to the guidelines
of the animal ethics committee of JNCASR. *Rudhira* floxed mice and conditional knockout mice (*Tek-*
Cre mediated endothelial deletion, *rudhira^{flox/flox}; TekCre^+*, abbreviated to *rudh^{CKO}*) were validated by
genotyping.

Cell culture, transfections, and small molecule treatments

Cell line sources and generation is reported elsewhere [14]. In brief, *rudhira* shRNA vectors (715,
716), and scrambled (non- silencing) control vector (TR30015) (Origene, USA) were microporated
into SVEC (mouse endothelial cell line) and selected for stable line generation. Control cell line is
referred to as 'NS' and *rudhira* knockdown as 'KD'. For Rudhira overexpression and rescue line, the
plasmid constructs used were GFP-2A-GFP and Rudh-2A-GFP which are described elsewhere [13].
Constitutively active TGFβRI (pcDNA3-ALK5 T204D, 80877) was purchased from addgene for
transfection of SVEC NS and KD cells. Control and shRNA vectors for s*mad2* or *smad3* knockdown)
were purchased from ShRNA Resource Centre (Department of Microbiology and Cell Biology, Indian
Institute of Science, Bangalore, India) and were from the SIGMA Library of shRNAs for Human gene
products. The shRNAs were transfected in HEK293T cells and selected in 1 µg/ml Puromycin for 7

days. Details of the oligonucleotide sequence of the shRNAs are provided in Supplementary Table S3.

HEK293, HEK293T cells were transfected using Calcium Phosphate method while SVEC cells were

transfected using Lipofectamine 2000 (ThermoFisher Scientific, USA). All small molecules were from

Sigma Chemical Co., USA, and treatments were performed in DMEM, unless otherwise indicated.

Cells were washed three times in PBS and serum starved for 12h. Thereafter, cells were kept

untreated or treated with 10 ng/ml of TGFβ (Fig. 1C) or BMP4 (Fig. S2D) for 2 h and then taken for

immunoblotting. For gene expression analysis, cells were induced with 10 ng/ml of TGFβ for 24 to 48

h (Figs. 1D, 4A, B, C, S3C). SB431542 was used at a concentration of 10 μM, as indicated (Fig. 2A, B

and S3A). For Transwell migration assays, small molecule dilutions were prepared in 5% FBS

containing DMEM (Fig. 2A, B). For interaction studies cells were serum starved for 12h and thereafter

kept untreated or treated with 10 ng/ml of TGFβ or 10% FBS for 2 h and then taken for PLA,

immunostaining or western blotting, as desired (Fig. 3, S3A, B). For TGFβ-dependent *rudhira*

transcription and MT stability assays, cells were serum starved for 48h in serum-free medium (SFM)

containing 5 mg/ml BSA and 20mM HEPES (pH 7.4) in DMEM. Thereafter, various treatments were

performed for 7h (Fig. 5A, B, C, S4C, D). TGFβ (10 ng/ml), Actinomycin D (AcD, 10 μM), Cycloheximide

(50 μg/ml) were diluted in SFM.

Quantitative RT-PCR (qRT-PCR)

RNA was isolated using TRIzol reagent (Invitrogen). Reverse transcription was performed using 2 μg

of DNase treated RNA and Superscript II (Invitrogen, Carlsbad, CA) according to manufacturer's

instructions. Quantitative RT-PCR (qRT-PCR) was carried out using EvaGreen (BIO-RAD, CA) in Biorad-

CFX96 Thermal cycler (BIO-RAD, CA). Details of primers used are provided in Supplementary Table

S1.

Immunostaining, Immunohistochemistry, Fluorescence Microscopy and Analysis

Yolk sac or embryos dissected at E10.5 were fixed in 4% paraformaldehyde and processed for

Cryosectioning (embryos) and immunostaining using standard procedures. Control and knockout

embryos were identified by embryonic tail genotyping. Cells were fixed in 4% paraformaldehyde at

room temperature or 100% methanol at -20 ˚C and processed for immunostaining using standard

procedures [30]. Primary antibodies used were against PECAM1 (CD31) (BD Biosciences, USA),

Rudhira, Smad2/3, pSmad2, pSmad2/3 (Cell Signaling Technology), β-Tubulin (DSHB, Iowa), Glu-

Tubulin (Abcam), GFP (Invitrogen). Secondary antibodies were conjugated with Alexa-Fluor 488 or

Alexa-Fluor 568 or Alexa-Fluor 633 (Molecular Probes). Bright field and phase contrast microscopy

were performed using an inverted (IX70, Olympus) microscope. Confocal microscope, LSM 880 with

airy scan, Zeiss, was used for fluorescence microscopy. All images in a set were adjusted equally for

brightness and contrast using Adobe Photoshop CS2, where required.

Immunoblot analysis and co-immunoprecipitation

50 µg whole cell lysate was used for western blot analysis by standard protocols. Blots were cut into strips and incubated with primary antibodies as indicated: Smad2/3, pSmad2, Smad1/5/8, pSmad1/5/8 (Cell Signaling Technology), GAPDH (Sigma Chemical Co., USA), Glu-Tubulin, Ubiquitin, TGFβRI, pTGFβRI (Abcam, USA), β- Tubulin (DSHB, Iowa), Rudhira (Bethyl Labs, USA). HRP conjugated secondary antibodies against appropriate species were used and signal developed by using Clarity Western ECL substrate (Biorad, USA). For co-immunoprecipitation, 500 µg whole cell lysate and 2 µg of the desired antibody were used. Complexes were captured on Protein G-Agarose beads and analyzed by immunoblotting.

Transwell migration assay

The assays and quantitation were carried out as mentioned earlier [13]. Briefly, 24 h after transfection with desired plasmid vectors, cells were serum-starved for 12 h and 20000 HEK cells or 40000 SVEC were plated onto the upper chamber of the transwell filter inserts with 8 µm pore size, 24-well format (Costar, USA). 10% FBS containing medium (with desired small molecules) was added to the lower chamber to serve as a chemo-attractant. After 24 h, cells were fixed in 4% paraformaldehyde for 10 min at room temperature. Cells on the top of the filter were removed using a cotton swab. Cells that had migrated to the bottom were fixed and stained with 0.5% crystal violet for 10 min at room temperature. The dye was extracted in methanol and absorbance measured spectrophotometrically at 570 nm.

***In situ* Proximity Ligation Assay (PLA or Duolink assay)**

In situ PLA (Proximity Ligation Assay) reaction was performed on SVEC cell lines. The cells were cultured, fixed, permeabilised and stained with primary antibodies for Smad2/3, β-Tubulin or Rudhira, as desired. Thereafter, the protocol for PLA as recommended by manufacturer (Duolink, USA) was followed. Post PLA, nuclei were counterstained with DAPI.

Rudhira promoter *in silico* analysis

Smad-binding elements or motifs (SBEs) and other transcription factor binding sites (TFBS) were

identified from experimental studies, available in JASPAR database (https://jaspar.genereg.net/).

PSCAN software (http://159.149.160.88/pscan) was used for motif scanning in mouse and human

rudhira/BCAS3 gene promoters (1 kb sequence upstream of the transcription start site) using

matrices obtained from JASPAR bioinformatics tool, as indicated. (Supplementary Table S2).

Quantification and Statistical analyses

Statistical analyses were performed using One Way ANOVA in the Data Analysis package in Microsoft

Excel, and two-way ANOVA and one-sample t-test in GraphPad Prism. $p < 0.05$ was considered

significant.

Acknowledgements

We thank Aksah Sam for maintaining mouse stocks; Developmental Studies Hybridoma Bank,

University of Iowa, USA for some antibodies; JNCASR Imaging facility, JNCASR Animal Facility, NCBS

Animal facility for access and Inamdar laboratory members for fruitful discussions. We are thankful

to Arghakusum Das for repeating the experiment for Figure 4A. This work was funded by grants

from the Department of Biotechnology, Government of India (Sanction no.

BT/PR11246/BRB/10/644/2008 dated 29.09.2009), JC Bose fellowship (Grant no. JCB/2019/000020),

Department of Science and Technology, Government of India, the Wellcome Trust, UK

(094879/B/10/Z) and intramural funds from Jawaharlal Nehru Centre for Advanced Scientific

Research, India.

Author Contributions

MSI conceived of the project and directed the work. DJ, PJ, RS, MSI performed and analyzed

all the experiments. DJ, PJ, MSI wrote the manuscript. Authors have no competing interests.

References

1. Tzima, E., *Role of small GTPases in endothelial cytoskeletal dynamics and the shear stress response.* Circ Res, 2006. **98**(2): p. 176-85.
2. Guo, X. and X.F. Wang, *Signaling cross-talk between TGF-beta/BMP and other pathways.* Cell Res, 2009. **19**(1): p. 71-88.
3. Moustakas, A. and C.H. Heldin, *Dynamic control of TGF-beta signaling and its links to the cytoskeleton.* FEBS Lett, 2008. **582**(14): p. 2051-65.
4. Lebrin, F., et al., *TGF-beta receptor function in the endothelium.* Cardiovasc Res, 2005. **65**(3): p. 599-608.
5. Hata, A. and Y.G. Chen, *TGF-beta Signaling from Receptors to Smads.* Cold Spring Harb Perspect Biol, 2016. **8**(9).
6. Massague, J., *TGFbeta in Cancer.* Cell, 2008. **134**(2): p. 215-30.
7. Dong, C., et al., *Microtubule binding to Smads may regulate TGF beta activity.* Mol Cell, 2000. **5**(1): p. 27-34.
8. Gundersen, G.G., I. Kim, and C.J. Chapin, *Induction of stable microtubules in 3T3 fibroblasts by TGF-beta and serum.* J Cell Sci, 1994. **107 (Pt 3)**: p. 645-59.
9. You, E., et al., *Dynein-mediated nuclear translocation of yes-associated protein through microtubule acetylation controls fibroblast activation.* Cell Mol Life Sci, 2020. **77**(20): p. 4143-4161.
10. Goumans, M.J., Z. Liu, and P. ten Dijke, *TGF-beta signaling in vascular biology and dysfunction.* Cell Res, 2009. **19**(1): p. 116-27.
11. Joshi, D. and M.S. Inamdar, *Rudhira/BCAS3 couples microtubules and intermediate filaments to promote cell migration for angiogenic remodeling.* Mol Biol Cell, 2019. **30**(12): p. 1437-1450.
12. Cook, T.A., T. Nagasaki, and G.G. Gundersen, *Rho guanosine triphosphatase mediates the selective stabilization of microtubules induced by lysophosphatidic acid.* J Cell Biol, 1998. **141**(1): p. 175-85.
13. Jain, M., et al., *Rudhira/BCAS3 is a cytoskeletal protein that controls Cdc42 activation and directional cell migration during angiogenesis.* Exp Cell Res, 2012. **318**(6): p. 753-67.
14. Shetty, R., et al., *Rudhira/BCAS3 is essential for mouse development and cardiovascular patterning.* Sci Rep, 2018. **8**(1): p. 5632.
15. Siva, K., et al., *Human BCAS3 expression in embryonic stem cells and vascular precursors suggests a role in human embryogenesis and tumor angiogenesis.* PLoS One, 2007. **2**(11): p. e1202.
16. Barlund, M., et al., *Cloning of BCAS3 (17q23) and BCAS4 (20q13) genes that undergo amplification, overexpression, and fusion in breast cancer.* Genes Chromosomes Cancer, 2002. **35**(4): p. 311-7.
17. Gururaj, A.E., et al., *MTA1, a transcriptional activator of breast cancer amplified sequence 3.* Proc Natl Acad Sci U S A, 2006. **103**(17): p. 6670-5.
18. Vizan, P., et al., *Controlling long-term signaling: receptor dynamics determine attenuation and refractory behavior of the TGF-beta pathway.* Sci Signal, 2013. **6**(305): p. ra106.
19. Goumans, M.J. and C. Mummery, *Functional analysis of the TGFbeta receptor/Smad pathway through gene ablation in mice.* Int J Dev Biol, 2000. **44**(3): p. 253-65.
20. Nockowski, P., et al., *Serum concentrations of transforming growth factor beta 1 in patients with psoriasis vulgaris.* Acta Dermatovenerol Croat, 2004. **12**(1): p. 2-6.
21. Devreotes, P. and A.R. Horwitz, *Signaling networks that regulate cell migration.* Cold Spring Harb Perspect Biol, 2015. **7**(8): p. a005959.
22. Bagchi, S., R. Fredriksson, and A. Wallen-Mackenzie, *In Situ Proximity Ligation Assay (PLA).* Methods Mol Biol, 2015. **1318**: p. 149-59.
23. Janke, C., *The tubulin code: molecular components, readout mechanisms, and functions.* J Cell Biol, 2014. **206**(4): p. 461-72.

24. Miyazono, K., *Positive and negative regulation of TGF-beta signaling.* J Cell Sci, 2000. **113 (Pt 7)**: p.
1101-9.
25. Yan, X., X. Xiong, and Y.G. Chen, *Feedback regulation of TGF-beta signaling.* Acta Biochim Biophys Sin
(Shanghai), 2018. **50**(1): p. 37-50.
26. Ferrari, G., et al., *Transforming growth factor-beta 1 (TGF-beta1) induces angiogenesis through*
vascular endothelial growth factor (VEGF)-mediated apoptosis. J Cell Physiol, 2009. **219**(2): p. 449-58.
27. Martin, M., et al., *Control of endothelial cell polarity and sprouting angiogenesis by non-centrosomal*
microtubules. Elife, 2018. **7**.
28. Bayless, K.J. and G.A. Johnson, *Role of the cytoskeleton in formation and maintenance of angiogenic*
sprouts. J Vasc Res, 2011. **48**(5): p. 369-85.
29. Itoh, F., et al., *Smad2/Smad3 in endothelium is indispensable for vascular stability via S1PR1 and N-*
cadherin expressions. Blood, 2012. **119**(22): p. 5320-8.
30. Schlaeger, T.M., et al., *Vascular endothelial cell lineage-specific promoter in transgenic mice.*
Development, 1995. **121**(4): p. 1089-98.
476

477   **Figures**

Figure 1

Fig. 1. Rudhira depletion deregulates developmental endothelial TGFβ signaling. (A) KEGG pathway
map indicating TGFβ pathway genes from KEGG database that are deregulated upon *rudhira*
depletion in mouse embryonic yolk sac, based on [14]. Genes were mapped based on fold change
in *rudhira*$^{-/-}$ yolk sacs in comparison to control to understand pathway regulation. Green:
downregulated, Red: upregulated. Blue circles indicate affected genes or processes known to be
affected upon Rudhira depletion, from earlier studies. (B) NS and KD SVEC cells, transfected with
constitutively active TGFβRI for 24h, were analyzed for Smad2 phosphorylation by Immunoblotting.
Graph shows the quantitation of pSmad2/Smad2 levels (N= 3 independent experiments). (C, D)
Control (NS), *rudhira* knockdown (KD) and rescued KD (KD + Rudhira-2A-GFP) cells were kept
untreated or treated with TGFβ and used for various assays, as indicated. (C) Immunoblot for
Smad2/3 and pSmad2. Graph shows the quantitation of pSmad2/Smad2 levels (N= 3 independent
experiments). (D) qRT-PCR analysis of *mmp9* and *pai1*, known Smad2/3 targets in ECs (N= 3
independent experiments). (E, E') Immunostaining for phosphorylated Smad2/3 in control and
rudhira conditional knockout *(rudh^{CKO})* embryos at E10.5. Boxed region in the embryonic heart in (E)
marking the endocardium and the myocardium are magnified in (E') (N= 3 embryos for each
genotype). White dotted line indicates the endocardium. Graph shows the quantitation of pSmad2/3
levels (N= 3 independent experiments). Statistical analysis was performed using two-way ANOVA
.(A, B, C, D) and student's t-test (E). Error bars indicate standard error of mean (SEM). *p<0.05,
**p<0.01, ***p<0.001. Scale bar: (E) 100 μm; (E') 20 μm.

Figure 2

Fig. 2. Rudhira functions downstream of TGFβ receptor activation to promote TGFβ-dependent cell migration. (A) Control (NS), *rudhira* knockdown (KD) and rescued KD (KD + Rudhira-2A-GFP) cells were analyzed for migration rates with or without TGFβ using a transwell migration assay. Graph shows the extent of cell migration as measured by crystal violet absorbance (N= 3 independent experiments). (B, C) HEK293 cells transfected with vector alone or Rudhira were analyzed for migration rates using a transwell migration assay in various conditions, as indicated. Graph shows the extent of cell migration as measured by crystal violet absorbance (N= 3 independent experiments). Statistical analysis was performed using one-way ANOVA. Error bars indicate SEM. Scale bar: (A, B) 100 μm. *p<0.05, **p<0.01, ***p<0.001.

Figure 3

Fig. 3. Rudhira is essential for the release of Smads from microtubules for TGFβ pathway

activation. (A) Analysis of interaction between Rudhira, Smad2/3 and MTs by co-

immunoprecipitation and immunoblotting, as indicated (N= 3 independent experiments). (B) Loss of

Smad-MT interaction on TGFβ stimulation in SVEC cells (N= 3 independent experiments). (C) NS and

KD cells were analyzed for MT and Smad2/3 interaction by PLA for Tubulin and Smad2/3 with or

without TGFβ treatment. PLA dots represent Smad2/3 and β-tubulin interaction. Graph shows

the quantitation of PLA dots per cell (quantitation from >80 cells for two knockdown cell lines

from N= 3 independent experiments; also see Fig. S3). (D, E) Association of Rudhira and MTs was

analyzed by immunostaining (D) or Proximity Ligation Assay (PLA) (E) for Rudhira and Tubulin with

or without TGFβ stimulation in SVEC. PLA dots represent Rudhira and β-tubulin interaction. Graph

(D) shows the quantitation of colocalization percentage (quantitation from ~40 cells from 9

images each with or without TGFβ respectively from N= 3 independent experiments) and graph

(E) shows the quantitation of PLA dots per cell (quantitation from 86 or 68 cells from 9 images

each with or without TGFβ respectively from N= 3 independent experiments). Statistical

analysis was performed using one-way ANOVA. Error bars indicate SEM. Scale bar: (C) 20 μm; (D, E)

10 μm. *p<0.05, **p<0.01, ***p<0.001.

Figure 4

Fig. 4. Rudhira is a Smad2/3-dependent target of TGFβ signaling. (A-C) TGFβ stimulation followed by qRT-PCR (A), immunoblot (B) or immunostaining (C) analysis (quantitation from 34 cells in each condition) of Rudhira levels in SVEC cells. Graphs in (B) and (C) show the quantitation of Rudhira levels with or without TGFβ (N= 3 independent experiments). (D) Analysis of Rudhira levels upon *smad2* or *smad3* knockdown in HEK293T cells by immunoblotting (N= 3 independent experiments). Statistical analysis was performed using one-way ANOVA. Error bars indicate SEM.

533   Scale bar: (C) 10 μm. *p<0.05, **p<0.01, ***p<0.001.

Figure 5

Fig. 5. TGFβ-dependent *rudhira* transcription stabilizes MTs. (A) Analysis of Rudhira levels and

MT stability (marked by Glu-Tubulin) upon treatments as indicated, by immunoblotting. Graphs

show the quantitation of Rudhira or Glu-Tub levels (N= 3 independent experiments). (B) Analysis of

MT resistance to Nocodazole mediated depolymerization at indicated dosage and time (N= 30 cells).

(C) MT stability in serum starved NS or KD SVEC cells kept untreated or treated with TGFβ was

analyzed by immunoblot for Glu-Tubulin. Graph shows the quantitation of Glu-Tub levels (N= 3

independent experiments). (D) Immunostaining for Glu-Tubulin in control and *rudhira* conditional

knockout *(rudh^{CKO})* embryos at E10.5. Endocardium in the heart is marked by PECAM1. Boxed region

in the embryonic heart is magnified in the insets. Red arrowheads in the insets mark the PECAM1

positive cells (N= 4 embryos for each genotype). Graph shows the quantitation of Glu-Tub

fluorescence intensity in the endocardium. (E) Model depicting cellular and molecular response to

TGFβ addition. Rudhira has an early and a late effect on MTs. Previously known information
about Rudhira is shown in blue. Events identified in this study are shown in red or pink. Parallel
lines indicate protein- protein interaction. Solid lines indicate direct effects. Dotted lines indicate
effects that may be direct or indirect. Arrows indicate positive regulation. Bar-headed line
indicates inhibition. Statistical analysis was performed using one-way ANOVA. Error bars indicate
SEM. Scale bar: (B) 10 μm; (D) 100 μm. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Rudhira-mediated microtubule stability controls TGFβ signaling during mouse vascular development.

Divyesh Joshi[1#], Preeti Jindal[1#], Ronak Shetty[1] and Maneesha S. Inamdar[1,2*]

[1]Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore, India;[2]Institute for Stem Cell Science and Regenerative Medicine (inStem), Bangalore, India.

Supplementary Figures and Figure legends

Figure S1

Fig. S1 (related to Fig. 1). Cell line validation. (A, B) Validation of Rudhira knockdown and restoration in SVEC cells by immunoblotting. Statistical analysis was performed using one-sample t-test. Error bars indicate standard error of mean (SEM). *p<0.05, **p<0.01,***p<0.001.

Figure S2

Fig. S2 (related to Fig. 1). Rudhira acts downstream to TGFβ receptors. (A) qRT-PCR analysis of *tgfβrI* and *tgfβrII* in Rudhira knockdown endothelial cells (N= 3 independent experiments). (B) TGFβRI phosphorylation was analyzed using immunoblot technique in SVEC NS and KD. Graph shows the quantitation of pTGFβRI/TGFβRI levels (N= 3 independent experiments). (C) NS and KD SVEC cells, transfected with constitutively active TGFβRI for 24h, were analyzed for total TGFβRI level by qRT-PCR. (D) SVEC control (NS) and knockdown (KD) cells were stimulated by BMP4 and used for immunoblotting of pSmad1/5/8. Graph shows the quantitation of pSmad1/5/8 by Smad1/5/8 levels (N= 3 independent experiments). (E, E') Immunostaining for total Smad2/3 in control and rudhira conditional knockout (rudhCKO) embryos at E10.5. Boxed region in the embryonic heart in (E) marking the endocardium and the myocardium are magnified in (E') (N= 3 embryos for each genotype). White dotted line indicates the endocardium. Graph shows the quantitation of Smad2/3 level (N= 3 independent experiments). Statistical analysis was performed using one-sample t-test. Error bars indicate standard error of mean (SEM). *p<0.05, **p<0.01,***p<0.001, ****p<0.0001. Scale bar: (E) 100 μm; (E') 20 μm.

Figure S3

Fig. S3 (related to Fig. 1, 2 and 3). Rudhira specifically regulates TGFβ-dependent Smad2/3 activation. (A) SVEC cells treated with SB431542 (10μM) for 2h and analyzed for Smad2 phosphorylation by Immunoblotting. (B) NS and two KD SVEC lines (KD1 and KD2) were analyzed for MT-SMAD interaction by PLA for β-Tubulin and Smad2/3 With or without TGFβ treatment (N= 3 independent experiments). (C) qRT-PCR analysis of negative regulators of TGFβ pathway (N=3 independent experiments). Statistical analysis was performed Using One way ANOVA. Error bars indicate standard error of mean (SEM). Scale bar (B): 20 μm. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure S4

Fig. S4 (related to Figs. 4 and 5). TGFβ induces *rudhira* transcription for MT stability. (A) Smad-binding element (SBE) matrix (SMAD3, ID MA0795.1; Smad4, ID MA1153.1) and predicted binding sites in mouse (in red) and human (in blue) *rudhira* promoters, from JASPAR and PSCAN bioinformatics tool. (B) Transcription factor binding sites (TFBS) available from JASPAR and PSCAN bioinformatics tool.(C) Analysis of Rudhira levels and MT stability (marked by Glu-Tubulin) upon treatments as indicated by immunoblotting (N= 3 independent experiments). (D) Analysis of MT stability in serum starved WT and KD cells kept untreated or treated with TGFβ or serum (N= 3 independent experiments).

50 **Supplementary Tables**

51 **Supplementary Table S1. Primers used for qRT-PCR analysis.**

Gene	Forward primer (5' to 3')	Reverse primer (5' to 3')
rudhira	TCCTACATGGAGAGCGTCG	GGAGGCTCATTTCCAGTGC
smurf2	AGACTGGTGTGAGCACATGG	CTTCCTGTTGCGGTATTGCG
smad6	GGGTGAATTCTCAGACGCC	GGTCGTACACCGCATAGAGG
smad7	CGGAAGTCAAGAGGCTGTGT	GACAGCCTGCAGTTGGTTTG
pai1	CTGTGCCCATGATGGCTCAG	ATGAACATGCTGAGGGTGTC
mmp9	GTCCAGACCAAGGGTACAGC	ATACAGCGGGTACATGAGCG
gapdh	GGTGAGGCCGGTGCTGAG	TGATGTCATCATACTTGGCAGG

55   **Supplementary Table S2. Predicted Transcription factor binding sites in mouse and human**

56   *rudhira/BCAS3* promoters, obtained from JASPAR bioinformatics tool.

Name	Position (from TSS)	Score	Background score	p-value	Matrix ID (JASPAR)	Strand	Predicted sequence
MAX::MYC	-428	0.959466	0.823	0.00252271	MA0059.1	-	AAGCACGTGGA
MYC	-429	0.959143	0.827	0.00590903	MA0147.3	+	GTCCACGTGCTT
TWIST1	-651	0.988341	0.854	0.00194672	MA1123.2	+	TTTCCAGATGTTC
HIF1A	-427	0.952744	0.849	0.0315803	MA1106.1	+	CCACGTGCTT
FOXP1	-386	0.992493	0.876	0.0124319	MA0481.3	+	ATGTAAACACA
FOXF2	-515	0.932307	0.826	0.0101525	MA0030.1	+	GGAAGGTAAACACA
MAX::MYC	-431	0.965445	0.823	0.00084454	MA0059.1	-	GAGCACGTGGA
MYC	-432	0.97587	0.831	0.00370719	MA0147.3	+	GTCCACGTGCTC
TWIST1	-412	0.88988	0.849	0.19157	MA1123.2	+	GAACCAGCTGTCG
HIF1A	-430	0.963428	0.856	0.0274206	MA1106.1	+	CCACGTGCTC
FOXP1	-389	0.989613	0.864	0.0168508	MA0481.3	+	ATGTAAACAGA
FOXF2	-518	0.932307	0.817	0.0116588	MA0030.1	+	GGAAGGTAAACACA

60   **Supplementary Table S3. The oligonucleotide sequence of the *smad2* and *smad3* shRNAs.**

Gene target	Oligonucleotide sequence in the shRNA (target sequence in red)
Smad2 sh#1	CCGGCGATTAGATGAGCTTGAGAAACTCGAGTTTCTCAAGCTCATCTAATCGTTTTTG
Smad2 sh#2	CCGGCCTAAGTGATAGTGCAATCTTCTCGAGAAGATTGCACTATCACTTAGGTTTTTG
Smad3 sh#1	CCGGGCCTCAGTGACAGCGCTATTTCTCGAGAAATAGCGCTGTCACTGAGGCTTTTTG
Smad3 sh#2	CCGGGAGCCTGGTCAAGAAACTCAACTCGAGTTGAGTTTCTTGACCAGGCTCTTTTTG

62