

1 **The role of extra-striate areas in conscious motor behavior: a**
2 **registered report with Fast-Optical Imaging**

3 Elisabetta Colombari^a, Giorgia Parisi^a, Sonia Mele^a, Chiara Mazzi^a and Silvia
4 Savazzi^a

5
6 ^a Perception and Awareness (PandA) Laboratory, Department of Neuroscience, Biomedicine and
7 Movement Sciences, University of Verona, Strada le Grazie 8, Verona, Italy

8
9

10 **Abstract**

11 Disclosing the brain areas responsible for the emergence of visual awareness and their timing of
12 activation represents one of the major challenges in consciousness research. In particular, isolating
13 the neural processes strictly related to consciousness from concurrent neural dynamics either related
14 to prerequisites or post-perceptual processing has long engaged consciousness research. In this
15 framework, the present study aims at unravelling the spatio-temporal dynamics underlying conscious
16 vision by adopting a distinctive experimental design in which both awareness and motor response are
17 manipulated, allowing the segregation of neural activity strictly related to awareness from response-
18 related mechanisms. To this aim, we will employ a GO/NOGO detection task, in which participants
19 will respond or withhold responding according to the experimental condition. Critically, during the
20 performance of the task, participants' brain activity will be recorded by means of Event-Related
21 Optical Signal (EROS) technique, which provides accurate information about brain functions both
22 from the temporal and spatial point of view, simultaneously. The combination of this experimental
23 design with EROS recording will enable us to pinpoint the neural correlates underlying conscious
24 vision and to disentangle them from processes related to the response. In addition, by coupling
25 conventional EROS analysis with Granger Causality analysis, we will be able to clarify the potential
26 interplay between consciousness-related extra-striate areas and response-related motor areas.

27 **1. Introduction**

28 Consciousness, namely the set of subjective experiences we have when we are awake, is one of the
29 most intriguing topics debated in neuroscience research. In particular, the search for its neural
30 correlates (NCC) has permeated the literature in recent decades. In broad strokes, one of the most
31 widely used approaches to assess such NCCs involves contrasting brain activity occurring when a
32 visual stimulus enters consciousness with brain activity occurring when the same stimulus does not
33 reach awareness. This renowned paradigm is known as *contrastive analysis* (Aru *et al.*, 2012) and
34 has been frequently combined with electrophysiological recording or functional neuroimaging,
35 leading to numerous and dissimilar results (Förster *et al.*, 2020). Indeed, the interpretations of spatio-
36 temporal dynamics underlying conscious vision are among the most disparate. ERP studies propose
37 two possible electrophysiological markers as correlates of visual awareness: an earlier occipito-
38 temporal negative deflection (i.e., Visual Awareness Negativity – VAN) detectable 200 ms after the
39 presentation of the stimulus, and a later positivity (i.e., Late Positivity – LP) widespread over centro-
40 parietal regions, peaking 300-500 ms after the stimulus onset (Koivisto & Revonsuo, 2010). However,
41 the electrophysiological signature/s characterizing conscious vision has still to be elucidated. This
42 may be attributed to one of the main limitations of the contrastive analysis, which is represented by
43 its ineffectiveness in dissociating the true NCC (i.e., the set of neural correlates necessary and
44 sufficient to enable consciousness) from concurrent neural dynamics either related to prerequisites or
45 post-perceptual processing (Aru *et al.*, 2012). In most prior studies aiming at identifying such NCCs,
46 participants were asked to make judgments about their experience. However, such an operation could
47 lead to confounding neural processes related to the task, not strictly to awareness per se.
48 For this reason, in an effort to disentangle the proper correlates of consciousness from neural activity
49 related to the response, no-report paradigms have been employed. In this framework, no-report
50 paradigms, where participants are not requested to perform any tasks or to provide any judgments
51 about their perceptual experience, represent an advantageous tool to dissociate the neural processes
52 strictly related to consciousness from subsequent processes related to the required response (Tsuchiya
53 *et al.*, 2015; Hatamimajoumerd *et al.*, 2022). Studies employing this kind of paradigm with different
54 techniques such as EEG and fMRI concluded that LP is highly modulated by several different
55 cognitive processes occurring at later stages of processing (Mazzi *et al.*, 2020; Schlossmacher *et al.*,
56 2020; Dembski *et al.*, 2021; Kronemer *et al.*, 2022), as well as by the task relevance of the stimulus
57 (Makeig & Jung, 2000; Pitts *et al.*, 2014; Shafto & Pitts, 2015; Schelonka *et al.*, 2017; Dellert *et al.*,
58 2021; Hense *et al.*, 2024). By contrast, the role of response requirements, as well as that of attention,
59 on the VAN are still debated as different studies have reported both positive (e.g., Bola &
60 Doradzińska, 2021; Dellert *et al.*, 2021; Doradzińska & Bola, 2024) and negative (e.g., Koivisto *et*

61 *al.*, 2006; Cohen *et al.*, 2020; Dellert *et al.*, 2022; Ciupińska *et al.*, 2024) results. Interestingly, in a
62 study published in 2016 by Koivisto and colleagues (Koivisto *et al.*, 2016), authors successfully
63 dissociated ERP correlates of visual awareness from those related to post-perceptual mechanisms,
64 disclosing that VAN was not modulated by response requirements. The authors adopted a particular
65 partial-report paradigm in which participants were sometimes asked to provide a report by pressing a
66 response button when they were aware of the stimulus and sometimes to withhold responding in case
67 of awareness. They found that, while the amplitude of LP was modulated by the response (i.e., it was
68 greater in trials where participants were asked to respond in case of awareness, compared to the Aware
69 condition where they were asked to withhold responding), VAN did not change depending on task
70 requirements. This allowed Koivisto and colleagues to advocate for an early onset of visual
71 awareness: the phenomenal content of a visual experience, indeed, takes place before LP, more
72 specifically in the temporal window of VAN.

73 Several pieces of evidence are consistent in considering VAN as the electrophysiological signature
74 of phenomenal consciousness (Koivisto *et al.*, 2008; Railo *et al.*, 2015), while the localization of its
75 neural generator still remains open. In this regard, previous MEG source localization studies (Vanni
76 *et al.*, 1996; Liu *et al.*, 2012) identified the Lateral Occipital Complex (LOC), an extra-striate visual
77 areas traditionally associated with objects recognition, as the generator of VAN. Moreover, previous
78 no-report studies using both EEG and fMRI (Dellert *et al.*, 2021; Kronemer *et al.*, 2022) have also
79 found awareness effects in LOC and linked it to VAN. The same result was achieved in a recent work
80 aimed at unravelling the spatio-temporal dynamics underlying conscious vision (Colombari *et al.*,
81 2024). In such study, participants were asked to perform a discrimination task on the orientation of a
82 tilted Gabor patch while their brain activity was recorded first with EEG and then with Fast Optical
83 Imaging. This allowed authors to identify the exact temporal window of VAN and LP and then, by
84 taking advantage of the peculiarity of Fast Optical Imaging of achieving both temporal and spatial
85 accurate information (Gratton & Corballis, 1995; Gratton & Fabiani, 2010; Baniqued *et al.*, 2013), to
86 investigate the spatio-temporal unfolding of brain activity occurring in these predetermined time
87 windows. Authors contrasted activity of Aware trials (i.e., trials in which participants reported to
88 perceive the orientation of the stimulus) with activity of Unaware ones and observed a sustained
89 activation of LOC in the VAN temporal window, consistently with the above-mentioned MEG
90 studies. More interestingly, they observed that, only when the stimulus crossed the threshold of
91 consciousness, activity in extra-striate visual areas triggered subsequent activation of motor areas,
92 although motor response was required in both Aware and Unaware conditions. Authors tried to
93 interpret this unexpected finding by ascribing it to the selection of the correct response, that could be
94 provided in the Aware trials only where participants consciously perceived the stimulus. Indeed, in

95 Aware trials participants had to press a specific button on the response box (to provide the correct
96 answer about the orientation of the Gabor patch), while when the stimulus was unseen (i.e., Unaware
97 trials) they had to respond randomly, by pressing indifferently one of the two response buttons.
98 However, the employed experimental paradigm did not allow the authors to thoroughly investigate
99 this issue. Thus, in order to clarify the interplay between extra-striate areas and motor regions in
100 awareness, in the present study we will adopt a go/no-go detection task (similar to that adopted by
101 Koivisto *et al.*, 2016), while recording participants' brain activity by means of Fast Optical Imaging.
102 Specifically, Event-Related Optical Signal (EROS) technique will be employed. This technique, by
103 shedding near-infrared light through the brain tissues, is able to detect changes in the light scattering
104 properties that are known to be directly related to neural activity, thus providing accurate information
105 about brain functions both from the temporal and spatial point of view, simultaneously (Gratton *et*
106 *al.*, 1997; Gratton & Fabiani, 1998, 2001). Critically, the study will adopt a distinctive paradigm
107 manipulating both awareness and response. The latter, indeed, will be provided sometimes in the
108 Aware condition (condition Aware-GO/Unaware-NOGO) and sometimes in the Unaware one
109 (condition Aware-NOGO/Unaware-GO). This double manipulation will enable us to unravel the
110 spatio-temporal unfolding of awareness-related activity, by disentangling neural activity related to
111 awareness from response-related mechanisms. Indeed, in the present study, we can investigate the
112 NCCs both when the motor response is required and when no task is performed, thus allowing to
113 isolate consciousness effects from the effects related to the task. Importantly, the experimental
114 paradigm adopted will enable us to elucidate the interplay between extra-striate visual areas and
115 motor areas. Indeed, in addition to conventional EROS analyses, we will perform Granger Causality
116 analysis, in order to disclose the relationship existing among the investigated areas. In broad strokes,
117 Granger analysis allows to move beyond the classical identification of cortical activation provided by
118 EROS analysis by disclosing functional circuits underpinning the investigated brain function (Seth *et*
119 *al.*, 2015). When coupled with EROS, Granger Causality analysis represents a powerful tool to
120 highlight predictive relationship between activations in the investigated regions of interest (ROI) at
121 different time-points (Parisi *et al.*, 2020).

122 Based on previous literature suggesting that VAN is independent from subjective report (Koivisto *et*
123 *al.*, 2016; Ye *et al.*, 2024) and LOC represents the cortical generator or VAN (Liu *et al.*, 2012;
124 Colombari *et al.*, 2024), we expect Aware trials to elicit early greater activation of LOC,
125 independently of the response requirement. Moreover, by combining EROS conventional analysis
126 with Granger Causality analysis, and manipulating both awareness and motor response, we aim to
127 highlight potential interplay between consciousness-related extra-striate areas and response-related
128 motor areas both when the motor response is required and when it has to be inhibited.

129 **2. Methods**

130 **2.1 Ethics Information**

131 The study is approved by the local Ethics Committee (Prog.171CESC) and it will be conducted in
132 accordance with the principles laid down in the 2013 Declaration of Helsinki ~~and~~. Participants will
133 be recruited from the University of Verona community, by means of printed flyers displayed on notice
134 boards at different University of Verona sites and through advertisements on social media. Each
135 participant will be fully informed about the modalities of the study before taking part in the
136 experiment and written informed consent will be signed. In addition, participants will receive
137 compensation for their participation and will be debriefed after the conclusion of the experiment.

138 **2.2 Participants**

139 We will recruit healthy adults, right-handed (as assessed by means of the standard handedness
140 inventory *Edinburgh Handedness Questionnaire*; Oldfield, 1971) and aged between 18 and 50 years
141 old. All of them will have to report normal or corrected-to-normal vision, no history of neurological
142 or psychiatric disorders and no contraindications to MRI. The study will be conducted at the Panda
143 lab of the University of Verona (Italy).

144 *2.2.1 Sample size estimation*

145 The estimate of the sample size for the current study is based on our previous EROS study (Colombari
146 et al., 2024), in which a similar paradigm was adopted, and similar analyses on similar ROIs were
147 performed. Specifically, EROS data from the ROI of LOC were extracted, and significant time-points
148 were averaged within participants so to have one value for each of them. Then, a one-sample t-test
149 was performed ($t(23) = 2.99$, $p = .006$, Cohen's $d = .611$), and the resulting Cohen's d was employed
150 to compute the sample size estimation for the current study. Specifically, the estimated sample size
151 for research questions Q1 (i.e., "Can we replicate Colombari et al., 2024 findings showing that LOC
152 is an NCC?") and Q2 (i.e., "Is the activity in LOC independent from the response?") was calculated
153 with G-Power software (v. 3.1.9.7), with a power of 90% and a level of significance of 2%. The
154 estimated sample size resulted in 32 participants (critical $t = 2.143$; actual power = 0.900). Considering
155 that the estimated sample size for this study ($n=32$) is more than double the typical sample size of
156 EROS studies present in literature, the same estimated sample size seems to be also adequate to
157 answer research questions Q3 (i.e., "Does consciousness modulate the activation of motor areas in a
158 detection task?") and Q4 (i.e., "Does consciousness modulate the activation of motor areas in
159 ABSENCE of motor response?"). For a review of the existing EROS literature, see Supplementary
160 Table 1 at

161 https://osf.io/ebfu3/?view_only=9ec2e6bf32ba4a8bb8b858639ec40a59) from which emerges that,
162 on average, EROS studies employ experimental samples composed of about 13 participants (mean
163 12.944; SD 7.008).

164 *2.2.2 Exclusion Criteria*

165 As better specified in section 2.3, before getting involved in the study, participants will undergo a
166 perceptual threshold assessment, in order to identify the proper stimulus to be employed in the main
167 experiment. To be enrolled in the study, participants will have to successfully complete this session.
168 The criterion used is that one of the stimuli presented during the threshold assessment will have to
169 be acknowledged as perceived a minimum of 25%, a maximum of 75%, or closest to the 50% of the
170 times (i.e., at perceptual threshold level). If no stimulus results at the threshold level, the participant
171 should not be enrolled in the study.

172 In addition, participants who will not complete all the experimental sessions, as well as participants
173 reporting a level of Awareness superior to 75% or inferior to 25% at the end of the experiment will
174 be excluded from analyses. This is to maintain comparable the number of trials in the two
175 experimental conditions (i.e., Aware and Unaware) and to ensure a reliable EROS activity (because
176 of its relatively low signal-to-noise ratio, EROS needs a high number of trials per condition, in
177 order to compute statistics). Moreover, participants whose behavioral performance will be affected
178 by biases related to the behavioral response (as assessed by catch trial analysis, explained more in
179 detail below) will be excluded from the analyses (see below –*Section 2.8.1 Behavioral data* for
180 more detailed information on the analysis of catch trials). Finally, participants whose EROS signal
181 could not be detected properly during the experiment (for example because of too dark hair or
182 technical issues) will not ~~also~~ be included in the analyses as well. In particular, the opacity value
183 (i.e., the product of the scattering and absorption coefficients) will be estimated for each participant.
184 Based on this value, it is possible to judge the quality of the signal for each participant,
185 independently from the experimental condition. Opacity values of all participants will be averaged
186 together providing the absorption coefficient to be used when running statistical analysis.
187 Participants whose opacity value is equal to 0 or exceeds three standard deviations of the mean will
188 be excluded from statistical analyses.

189 Importantly, each participant who will be excluded due to the previously mentioned exclusion criteria,
190 will be replaced with the recruitment of another participant. Thus, the number of participants to be
191 recruited will be increased to reach a total of 32 analyzed subjects, as specified in section 2.2.1.

192 **2.3 Stimuli**

193 Stimuli will be created by means of a custom-made Matlab script (version R2022b; the MathWorks,
194 Inc., Natick, MA) and resized by means of Photoshop (Adobe Photoshop CC, v2014.0.0). As shown
195 in Figure 1, they will be gray circles (.85 .85 .85 RGB), presented on a black background, with 8 radii
196 equally distanced one from another. One radius (the first one, clockwise) can be slightly thicker than
197 the others (critical trials) or not (catch trials). The thickness of the radius for critical stimuli will be
198 individually assessed for each participant on the basis of a subjective perceptual threshold assessment
199 that will be held before the main experiment.

200 Both in the perceptual threshold assessment and in the main experiment, the stimulus will be
201 presented in the lower right quadrant of the screen, specifically at an eccentricity of 3.5° from the
202 fixation cross along the vertical meridian and of 2° along the horizontal one. This is to allow a left-
203 lateralized EROS montage, as a full-head montage is not achievable in our lab due to technical
204 constraints (i.e., insufficient probes). Moreover, since EROS technique is sensitive to depth, a right-
205 lateralized stimulus ensures that it elicits activity in the left portion of the primary visual cortex, which
206 is known to be anatomically closer to the skull compared to the right one (Zhao *et al.*, 2022), thus
207 ensuring a better penetration of near-infrared light through brain tissues.

208 **2.4 Perceptual Threshold Assessment**

209 Before starting the experiment, participants will undergo a perceptual threshold assessment, with the
210 aim of identifying, for each participant, the level of thickness of the critical radius so that it results to
211 be perceived as thicker 50% of the times. To this aim, stimuli with different levels of radius thickness
212 will be randomly presented and the subjective perceptual threshold will be measured using the method
213 of constant stimuli. Specifically, 9 levels of radius thickness will be presented. The range of stimuli
214 to be used in the perceptual threshold assessment will be selected based on the results of a pilot
215 experiment in which participants were asked to perform the same task employed in the perceptual
216 threshold assessment while presented with a wider range of radius thickness. This will allow us to
217 identify a smaller range of optimal stimuli to be presented thus excluding a range of stimuli whose
218 thickness was almost never or always reported by participants. Each level of radius thickness will be
219 presented 5 times per block, for a total of 8 blocks. Thus, all the stimuli, as well as the catch stimulus,
220 will be presented 40 times each. Participants will be asked to press the spacebar as soon as they detect
221 the stimulus with a thicker radius. The stimulus identified as perceived a minimum of 25%, a
222 maximum of 75%, and closest to 50% of the times at the end of the subjective perceptual threshold
223 assessment will be used in the experimental task, together with the catch. The perceptual threshold
224 assessment, as well as the main experiment, will be conducted in a dimly illuminated room and

225 participants will be sitting in front of a 17 in. LCD monitor (resolution 1920x1080, refresh rate of
226 144 Hz) placed at a viewing distance of 57 cm. Their head will be held in place by means of an
227 adaptable chin rest so that eyes are aligned with the center of the screen. Both the perceptual threshold
228 assessment and the main experiment will be programmed and administered using E-Prime 3.0
229 software (E-Prime Psychology Software Tools Inc., Pittsburgh, PA, USA). Before starting the
230 perceptual threshold assessment, participants will undergo a fixation training (Leung *et al.*, 2009), in
231 order to ensure they will maintain their gaze on the central fixation cross correctly.

232 **2.5 Experimental Procedure**

233 The experiment will be composed of two identical sessions lasting approximately 3 hours each
234 performed on different days. The first session will be preceded by the assessment of the subjective
235 perceptual threshold, which, in turn, will last around 20 minutes. The two experimental sessions will
236 be identical except for the EROS montages, specifically devised to obtain better coverage of the brain
237 areas of interest. The order of the montages will be counterbalanced across participants, as well as
238 the order of conditions (see below for more detailed information).

239 The task will be a two-conditions go/no-go detection task, similar to that adopted by Koivisto *et al.*,
240 2016, in which participants have to respond in different ways according to the experimental condition
241 (Table 1). In condition “Aware-GO”, they will be asked to press the spacebar on the keyboard as soon
242 as they perceive the thicker radius, and withhold responding when they do not perceive any difference
243 among radii. On the contrary, in condition “Aware-NOGO”, participants will be asked to withhold
244 responding when they perceive a thicker radius, and press the response button when they do not
245 perceive any difference. Each trial will begin with the presentation of a central fixation cross,
246 followed 500 ms later by a sound (1000Hz) presented for 100 ms, notifying participants of the
247 subsequent onset of the stimulus. After a random interval ranging from 500 to 600 ms, the stimulus
248 will be presented for 100 ms in the lower right quadrant of the screen. After that, participants will be
249 asked to respond according to the experimental condition. Each experimental session will be
250 composed of 24 blocks: 12 blocks for condition Aware-GO/Unaware-NOGO and 12 blocks for
251 condition Aware-NOGO/Unaware-GO, counterbalanced across participants according to the order
252 depicted in Table 1. Each block will consist of 50 critical trials and 15 catch trials. The whole
253 experiment will be composed of 48 blocks per participant, for a total of 2400 critical trials and 720
254 catch trials per participant.

255

256

Awareness

		yes	no
		Aware-NOGO	Unaware-NOGO
Response	no	Aware-NOGO	Unaware-NOGO
	yes	Aware-GO	Unaware-GO

257 **Table 1. Experimental conditions.** Both Awareness and Response are manipulated: Awareness is
 258 experimentally manipulated by employing a threshold stimulus, so that sometimes it is consciously perceived
 259 (Aware) and sometimes not (Unaware). Response is manipulated by the task: in condition GO participants are
 260 asked to respond by pressing a key, while in condition NOGO they are asked to withhold responding. The
 261 combination of these two manipulations gives rise to the 4 experimental conditions depicted in the table.

262

Participants	Day 1		Day 2	
	EROS montage 1	Task	EROS montage 2	Task
1	A	GNNG	B	NGGN
2	B	GNNG	A	NGGN
3	A	NGGN	B	GNNG
4	B	NGGN	A	GNNG

263 **Table 2. Counterbalancing of montages and task conditions across participants.** Both EROS montages
 264 and task conditions (G = Aware-GO/Unaware-NOGO; N = Aware-NOGO/Unaware-GO) will be
 265 counterbalanced across participants. In the column “Task”, each letter represents 6 blocks of task. Thus, each
 266 day, participants will perform 12 blocks per condition, for a total of 24 blocks of task per day.

267 2.6 Optical Recording

268 Three synchronized Imagent frequency domain systems (ISS, Inc., Champaign, IL) will be used to
 269 record continuous fast optical data throughout experimental sessions. Each system is equipped with
 270 4 photo-multiplier tubes detectors, for a total of 12 detectors. Near-infrared light (830 nm) will be
 271 delivered from 48 laser diodes on participants’ scalp and it will be modulated at 110 MHz. Each of
 272 12 detectors will receive light from sets of 16 light emitters, multiplexed every 25.6 ms, resulting in
 273 a sampling rate of 39.0625 Hz.

274 To avoid cross-talk between channels, the array of source-detector pairs (i.e., the montage) will be
 275 created by means of a specific program (NOMAD, Near-Infrared Optode Montage Automated
 276 Design) implemented in Matlab, useful to place sources and detectors at optimal distances. In this
 277 experiment, we will set the minimal distance to 17.5 mm and the maximum distance to 50 mm, in
 278 order to ensure an extensive coverage of the brain regions of interest both from the spatial and the
 279 depth point of view. The distance between the source and the detector of a channel, in fact, determines

280 the depth of the light pathway (Gratton *et al.*, 2000), thus corresponding to the depth of the
281 investigation: namely, longer channels can investigate deeper layers and shorter channels can
282 examine shallower regions.

283 Both light emitters and detectors will be placed on participants head using a custom-built helmet. To
284 minimize interferences, before placing the optical fibers on the head, the hair will be carefully moved
285 with cotton buds, so that the fibers can reach the scalp directly. In order to better adhere to the head
286 of the participant, we will employ two helmets of different sizes: one 55-56 cm large, and one 57-58
287 cm large. For each helmet, we will develop two different montages, so that to provide a dense
288 coverage of the regions of interest (i.e., the left occipital, temporal and parietal cortices, see Figure
289 2). Each montage will consist of the combination of 12 detectors and 48 light emitters, resulting in a
290 total of 192 channels per montage. As mentioned before, each montage will be recorded in a separate
291 session, and the order will be counterbalanced across participants.

292 At the end of each EROS session, the scalp location of each source and detector will be digitized in
293 relation to four fiducial points (i.e., nasion, inion and pre-auricular points) with a neuro-navigation
294 software (SofTaxis, E.M.S., Bologna, Italy) combined with a 3D optical digitizer (Polaris Vicra, NDI,
295 Waterloo, Canada). Afterwards, the digitized scalp locations will be co-registered with each
296 participant's individual MRI, using a dedicated software package (OCP, Optimized Co-registration
297 Package, Matlab code developed by Chiarelli and colleagues (Chiarelli *et al.*, 2015).

298 For this reason, participants will undergo a structural MRI at the Azienda Ospedaliera Universitaria
299 Integrata of Verona (AOUI).

300 **2.7 MRI Acquisition**

301 Participants' individual structural MRI will be acquired by means of a 3 Tesla Philips Ingenia scanner
302 with a 32-channel head RF receive coils. A whole brain high-resolution 3D T1-weighted image (T1w)
303 Turbo-field echo image (1mm-isotropic TE/TR=3.8/8.4 ms, TI=1050 ms) will be acquired.

304 The T1w field of view (240 x 240 x 180 mm) will be large enough to allow for the ears and the entire
305 scalp to be fully included in the image to facilitate later and accurate co-registration with functional
306 data.

307 **2.8 Data Analysis**

308 *2.8.1 Behavioral data*

309 Raw data will be processed by means of custom scripts created on Matlab (the MathWorks, Inc.,
310 Natick, MA). Data will be divided into the 4 experimental conditions (i.e., Aware-GO, Unaware-
311 NOGO, Aware-NOGO, Unaware-GO). For each participant, trials with reaction times lower than 150
312 ms and higher than 3 standard deviations from the mean will be excluded from the analysis. Data will

313 be successively analyzed using Jamovi (version 2.3.28): first, the percentage of Aware and Unaware
314 trials will be calculated, in order to assess that a sufficient amount of trials is present for each
315 condition. Participants presenting more than 75% or less than 25% of Awareness will be discarded
316 from the sample. This is because, in that case, the number of Unaware (or Aware) trials would be
317 insufficient for statistical EROS analysis. EROS technique, indeed, although having a high
318 localization power from both the spatial and temporal point of view, has a relatively low signal-to-
319 noise ratio. For this reason, a high number of trials is required for statistical analysis. Subsequently,
320 reaction times (RTs) will be analyzed for the “GO” conditions, thus paired sample t-tests (two-tailed)
321 will be applied to compare the mean RTs between Aware-GO and Unaware-GO conditions. Finally,
322 to verify that participants are performing the task accurately and that there are no biases related to the
323 response, catch trials will be analyzed. As mentioned above, catch trials are those trials in which all
324 the radii of the stimulus are equally thick, thus no differences in the stimulus are present. In case of
325 catch trials, the participants’ task will be different according to the condition: in the Aware-GO
326 condition, they are expected to withhold responding, while in the Aware-NOGO condition, they are
327 expected to respond. Thus, catch trials will be analyzed separately for the two conditions (GO and
328 NOGO) by means of a paired sample t-test (two-tailed), in order to ensure that the behavioral
329 performance follows the above-mentioned trend. Paired sample t-tests (two-tailed) will indeed be
330 performed to test whether catch trials performance is significantly different from critical trials.

331 2.8.2 EROS data

332 Pre-processing of continuous phase delay (i.e., time-of-flight) data will be computed by means of a
333 dedicated in-house software, P-POD (Pre-Processing of Optical Data, run in Matlab, version
334 R2013b). Thus, raw data will be normalized (i.e., corrected for phase wrapping and de-trended to
335 remove low-frequency drifts), demeaned and filtered by means of a 6th order Butterworth band-pass
336 filter which allows frequencies between 0.5 Hz and 15 Hz. Pulse artifact will be removed by using a
337 regression algorithm (GRATTON *et al.*, 1995). After that, data will be averaged separately for each
338 subject, condition, and channel and segmented into epochs time-locked to the onset of the stimulus.
339 Each epoch will comprise a period from 486 ms before the stimulus onset to 998 ms following the
340 stimulus onset, resulting in an epoch lasting 1484 ms. Subsequently, statistical analyses will be
341 computed with an in-house software package (Opt-3d; (Gratton, 2000)), which provides statistical
342 spatial maps of fast optical data.


343 To perform statistics, data from channels whose diffusion paths intersect a given voxel will be
344 combined (Wolf *et al.*, 2014). Phase delay data will be spatially filtered with an 8-mm Gaussian
345 kernel and baseline corrected using a 204 ms time-window preceding the stimulus onset. Within each

346 ROI, t-Statistics will be calculated at group level, converted into Z-scores and corrected for multiple
347 comparisons using random field theory (Worsley *et al.*, 1995; Kiebel *et al.*, 1999). Then, Z-scores
348 will be weighted and orthogonally projected onto the surface of an MNI template brain, according to
349 the physical homogenous model (Arridge & Schweiger, 1995; Gratton, 2000).

350 In order to investigate the neural dynamics related to conscious vision and to disentangle the role of
351 the motor areas, the following contrasts between conditions will be computed: 1) Aware-GO versus
352 Unaware-GO and 2) Aware-NOGO versus Unaware-NOGO. These contrasts allow to investigate the
353 research questions the proposed study aims at answering (see Section 3 for a detailed description of
354 the planned analysis). Importantly, both frequentist and Bayesian statistics (with default priors) will
355 be computed, to test both positive and negative effects.

356 Moreover, Granger Causality analysis will be computed. Granger Causality analysis allows to explore
357 the predictive interactions between different brain areas at different time-points. Specifically, this
358 approach requires a region of interest (ROI) to be used as a “seed” and investigating whether the
359 activity of this seed predicts activity in the other ROIs at a later time-lag, by deriving statistical maps
360 from t-statistics computation (then transformed into z scores) for each lag.


361 Statistical functional analysis will be computed within specific predetermined regions of interest
362 (ROIs) and time intervals. ROIs will be defined by a 2-dimensional box-shaped structure, covering
363 an area of 20x20 millimeters. Critical ROIs will be selected on the basis of the results obtained in the
364 above-mentioned experiment (Colombari *et al.*, 2024) and they will be located in the occipital and in
365 the left parietal and temporal lobes, specifically over the primary visual cortex (V1, Brodmann Area
366 17), the left lateral occipital cortex (LOC, Brodmann Area 19), the left supplementary motor area
367 (SMA, Brodmann Area 6), the left premotor area (PM, Brodmann Area 6) and the left primary motor
368 cortex (M1, Brodmann Area 4). Statistical analysis will be computed within specific temporal
369 windows of interest selected on the basis of the results obtained by Colombari *et al.*, 2024. This is to
370 reduce the risk of false positives, as Opt3d does not offer the possibility to correct data for multiple
371 comparisons in the temporal domain. The specific time windows tested for each hypothesis are listed
372 in Table 3.



373

374 **Figure 1. Trial procedure and stimuli:** A) Experimental procedure: the trial begins with a fixation cross
375 persisting at the center of the screen for 500 ms. After that, an acoustic tone lasting 100 ms will be presented,
376 followed by a random interval ranging from 500 to 600 ms. Then, the stimulus will be presented for 100 ms
377 and participants will be asked to respond according to the experimental condition (i.e., Aware-GO or Aware-
378 NOGO). B) Example of stimuli: on the left is shown the catch stimulus, with all the radii equally thick; on the
379 right is depicted the critical stimulus, with the first radius, clockwise, thicker than the others.

380



381

382 **Figure 2. Covered area.** The gray area represents the area covered by the EROS montages (combined
383 together) from the sagittal, axial and coronal point of view.

3. Study design

Question	Hypothesis	Sampling Plan	Analysis Plan	Rationale for deciding the sensitivity of the test for confirming or disconfirming the hp	Interpretation given different outcomes
<p>Q1: Can we replicate Colombari et al., 2024 findings showing that LOC is an NCC?</p>	<p>H1: We hypothesize to replicate Colombari et al., 2024 results: greater activity in LOC in an early temporal window (i.e., 150-350 ms post stimulus onset) is observed when contrasting Aware and Unaware trials in the condition in which the response is required (i.e., GO condition)</p> <p>Expected outcome: LOC aware-GO>LOC unaware-GO, as measured by EROS activity</p>	<p>Since the present research question aims at replicating the results of Colombari et al., 2024, sample size estimation is based on those EROS data. Sample size calculation was thus performed with G-Power software (v. 3.1.9.7), with a power of 90% and a significance level of 2%, resulting in 32 participants.</p>	<p>A1: The goal is to replicate the results of Colombari et al., 2024, in which the manual response was required for both Aware and Unaware conditions. Here, in order to perform the same analysis, early LOC activity in Aware-GO and Unaware-GO trials will be compared by using a paired-sample one-tailed t-test, computed with the EROS dedicated analysis software “Opt3d”.</p> <p>In addition to frequentist statistics, Bayesian statistics (with default prior) will be computed to test potential null effects.</p> <p>Contrast to be computed: AWARE GO VS UNAWARE GO</p> <p>ROI to be tested: LOC</p>	<p>Sample size estimation is based on previous EROS data (Colombari et al., 2024), on which similar analyses were performed.</p>	<p>O1.1: A significant t-test within the interval of interest will be interpreted as a successful replication of previous findings, supporting the involvement of LOC in NCC.</p> <p>O1.2: The absence of this effect will not confirm the hypothesis, suggesting that LOC is not involved in the conscious detection of a stimulus property.</p>

			Time interval of interest: 150-350ms after stimulus onset		
Q2: Is the activity in LOC independent from the response?	H2: We hypothesize that LOC activity is independent from response requirement: when contrasting activity elicited by Aware-NOGO trials with activity elicited by Unaware-NOGO trials, we expect to find the same activation of LOC found in the Aware-GO vs Unaware-GO contrast. Expected outcome: LOC aware-NOGO>LOC unaware-NOGO, as measured by EROS activity (LOC aware-GO>LOC unaware-GO)=(LOC aware-NOGO>LOC unaware-NOGO)	Since research question Q2 involves the same analyses of research question Q1 (but for the NOGO condition), the sampling plan for Q2 is the same as for Q1.	A2.1: A paired-sample one-tailed t-test will be computed in order to compare early activity in LOC in the NOGO condition. Thus, activity in Aware-NOGO and Unaware-NOGO trials will be contrasted. Both frequentist and Bayesian statistics (with default prior) will be computed. Contrast to be computed: AWARE NOGO VS UNAWARE NOGO ROI to be tested: LOC Time interval of interest: 150-350ms after stimulus onset	As above	O2.1.1: A significant t-test in the time window of interest will suggest that LOC activity is independent from response, since its activity is observed even when no response is required (NOGO conditions). O2.1.2: If greater activity in LOC in the time window of interest is not observed, then it means that LOC activity is somehow related to the motor response.
			A2.2: The interaction effect between awareness and motor response will be tested by means of a paired-sample one-tailed		O2.2.1: Significant interaction effect will suggest that activity in LOC depends from response requirement O2.2.2: The absence of a difference between the two effects will suggest

			<p>t-test computed between contrast Aware-GO VS Unaware-GO and contrast Aware-NOGO VS Unaware-NOGO</p> <p>In addition to frequentist statistics, Bayesian statistics (with default prior) will be computed to test potential null effects</p> <p>Contrast to be computed:</p> <p>(AWARE GO VS UNAWARE GO) - (AWARE NOGO VS UNAWARE NOGO)</p> <p>ROI to be tested: LOC</p> <p>Time interval of interest: 150-350ms after stimulus onset</p>		that motor response does not affect awareness-related activity in LOC
<p>Q3: Does consciousness modulate activation of motor areas in a detection task?</p>	<p>H3: When a motor response is required, consciousness modulates activation of motor areas (MA), as activity in motor areas is triggered by LOC (Colombari et al., 2024)</p> <p>Expected outcome:</p>	<p>Considering that the estimated sample size for this study (n= 32) is more than twice the typical sample size of EROS studies present in literature (see Supplementary Table 1, where a systematic review of existing EROS literature revealing that the typical sample size used is</p>	<p>A3.1 A paired-sample one-tailed t-test will be computed in order to compare early activity in Motor Areas in the GO condition. Thus, activity in Aware-GO and Unaware-GO trials will be contrasted.</p> <p>In addition to frequentist statistics, Bayesian</p>	As above	<p>O3.1.1: A statistically significant difference between the two conditions will suggest that, even in a detection task, response related motor activity is stronger in the Aware condition compared to the Unaware one.</p> <p>In Colombari et al., 2024 this difference was observed. Importantly, in this previous study a <i>discrimination</i> task was employed and participants were asked to provide two different</p>

	<p>MA aware-GO>MA unaware-GO, as measured by EROS activity</p> <p>LOC activity predicts MA activity (investigated by means of Granger Causality Analysis)</p>	<p>13 participants is depicted), the same estimated sample size of Q1 and Q2 seems to be also adequate to answer research questions Q3 and Q4.</p>	<p>statistics (with default prior) will be computed to test potential null effects</p> <p>Contrast to be computed:</p> <p>AWARE GO VS UNAWARE GO</p> <p>ROI to be tested:</p> <p>Motor areas</p> <p>Time interval of interest:</p> <p>Based on mean RTs, with a time window of ± 1.5 sd around the mean</p>		<p>responses in case of Awareness (intentional) or Unawareness (random). Instead, in this study participants are asked to perform a <i>detection</i> task, in which the motor behavior made to provide the response, when required, is the same for both Aware and Unaware condition and thus no response selection is required.</p> <p>O3.1.2: If no difference between the tested conditions is observed, it will suggest that in a detection task there is no difference in the motor activity related to the response.</p>
			<p>A3.2: In order to further investigate the flow of activity occurring in the investigated brain areas, Granger Causality Analysis will be performed. In the present study, we will perform Granger analysis on the “Aware-GO VS Unaware-GO” contrast, since we are interested in investigating whether activity in motor areas is predicted by previous activity in LOC, when a motor response is required (i.e., in the GO condition). Thus, LOC will be used as seed ROI</p>		<p>O3.2.1: Significant predictive interactions between LOC and motor areas will suggest that, when the stimulus enters consciousness, awareness-related activity in LOC predicts subsequent activity in motor areas. This (expected) outcome will suggest that consciousness modulates subsequent response-related motor activity, by directly triggering activation of motor areas, as observed in Colombari et al., 2024</p> <p>O3.2.2: If no significant interactions between LOC and MA will be highlighted, then it would mean that activity in motor areas is not predicted by LOC. Specifically, it could be surmised that in a <i>detection</i> task, consciousness does not modulate activation of motor areas, as observed in Colombari et al., 2024, where a</p>

			<p>and later activity in motor areas will be investigated.</p> <p>In addition to frequentist statistics, Bayesian statistics (with default prior) will be computed to test potential null effects</p> <p>Contrast to be computed:</p> <p>AWARE GO VS UNAWARE GO</p> <p>ROI to be tested: LOC (as seed ROI) Motor areas as predicted areas</p> <p>Time interval of interest: LOC: 150-350 ms after the stimulus onset (within this interval, about ± 4 time points around the peak value -according to the peak waveform- will be selected)</p> <p>MA: based on mean RTs, with a time window of ± 1.5 sd around the mean</p>		<p><i>discrimination</i> task was employed. The difference in the two tasks, indeed, consists in the type of motor response required: in the case of the discrimination task, the participant is asked to press one button or another according to the response. Conversely, in a detection task, the participant has to press a key when the target stimulus is detected. Thus, no selection of the response is needed. This difference could play a role in the relationship between consciousness and motor areas.</p>
<p>Q4: Does consciousness modulate activation of motor areas in ABSENCE of</p>	<p>H4: Consciousness modulates activation of motor areas, even if the motor response is not required</p>	<p>As Q3</p>	<p>A4.1: A paired-sample one-tailed t-test will be computed in order to compare activity in Motor Areas in the NOGO condition. Thus, activity</p>	<p>As above</p>	<p>O4.1.1: A statistically significant t-test will suggest that, when a motor response is not provided, the inhibition required to withhold responding is stronger when the visual characteristic of the stimulus is consciously</p>

<p>motor response?</p>	<p>Expected outcome: MA aware-NOGO>MA unaware-NOGO</p> <p>LOC predicts MA (investigated by means of Granger Causality Analysis)</p>		<p>in Aware-NOGO and Unaware-NOGO trials will be contrasted.</p> <p>In addition to frequentist statistics, Bayesian statistics (with default prior) will be computed to test potential null effects</p> <p>Contrast to be computed:</p> <p>AWARE NOGO VS UNAWARE NOGO</p> <p>ROI to be tested: Motor areas</p> <p>Time interval of interest: Based on mean RTs, with a time window of ± 1.5 sd around the mean</p>		<p>perceived, compared to when no difference is perceived.</p> <p>O4.1.2: If no difference between the tested conditions is observed, this will suggest that i) no inhibition is required to withhold responding, both in the Aware and Unaware condition, or ii) the inhibition is equally strong for the two conditions.</p>
			<p>A4.2: With the aim of investigating the flow of activity occurring in the investigated brain areas also in the condition where no response is required, Granger Analysis will be performed on the “Aware-NOGO VS Unaware-NOGO” contrast. This will allow to investigate whether activity in motor areas is triggered by previous activity in LOC,</p>		<p>O4.2.1: If significant predictive interactions between LOC and motor areas will be observed, then consciousness modulates subsequent activity in motor areas also in absence of a motor response. This could be due to inhibition of the response processes.</p> <p>O4.2.2: If no significant interactions between LOC and MA will be highlighted, then LOC does not predict activity in motor areas in absence of motor response.</p>

			<p>even when a motor response is not required. Thus, LOC will be used as seed ROI and later activity in motor areas will be investigated.</p> <p>In addition to frequentist statistics, Bayesian statistics (with default prior) will be computed to test potential null effects</p> <p>Contrast to be computed:</p> <p>AWARE NOGO VS UNAWARE NOGO</p> <p>ROI to be tested: LOC (as seed ROI)</p> <p>Motor areas as predicted areas</p> <p>Time interval of interest: LOC: 150-350 ms after the stimulus onset (within this interval, about ± 4 time points around the peak value -according to the peak waveform- will be selected)</p> <p>MA: Based on mean RTs, with a time window of ± 1.5 sd around the mean</p>		
--	--	--	--	--	--

386 **4. Pilot study**

387 In order to test the experimental paradigm, we pilot-tested the task.

388 A total of 10 right-handed participants (5 females and 5 males; mean age \pm standard deviation: 21
389 years \pm 1.0) took part in the pilot study. They all reported normal or corrected-to-normal vision and
390 no history of neurological or psychiatric disorders. All of them provided written informed consent
391 before starting the experiment.

392 After the first session, two participants dropped out the experiment, hence data from 8 participants
393 were included in the statistical analyses. Moreover, in order to maintain an equal number of trials in
394 both the conditions (i.e., Aware and Unaware), the percentage of Aware and Unaware trials was
395 calculated and data from participants reporting a proportion of awareness equal or superior to 80%
396 (i.e., 3 participants) were discarded from subsequent analysis. For this pilot study, we decided to raise
397 the awareness threshold of acceptance to 80% (instead of 75%, that will be used in the experiment)
398 in order to be more inclusive, given the low number of participants.

399 Thus, in total, data from 5 participants were included in the behavioral and functional analyses.


400 **4.1 Preliminary Results**

401 *4.1.1 Behavioral results*

402 Raw data were processed by means of scripts created on Matlab (version R2017b; the MathWorks,
403 Inc., Natick, MA). According to the participants' responses, trials were sorted into the four
404 experimental conditions (i.e., Aware-GO, Unaware-NOGO, Aware-NOGO and Unaware-GO).
405 Aware trials were those trials in which the participant reported to perceive the thicker radius, while
406 Unaware trials were those trials in which participants could not perceive that the radius was thicker.
407 As specified in Section 2.8, trials with RTs lower than 150 ms or higher than 3SD from the mean
408 were removed. After removal, we had on average 830.6 trials for the Aware-GO condition, 389.2 for
409 the Unaware-NOGO condition, 738.8 trials for the Aware-NOGO condition and 491.4 for the
410 Unaware-GO.

411 Subsequently, once assessed the normality of RTs and Awareness distributions (Shapiro-Wilk test.
412 RTs distribution: $W=0.824$, $p=0.125$; Awareness distribution: $W=0.817$, $p=0.112$), the percentage of
413 Awareness for the two conditions was calculated: in the GO condition, Aware trials represented on
414 average 68.02% of the trials, while in the NOGO condition, Aware trials constituted the 59.82% of
415 the trials. Paired sample (two-tailed) t-test performed with Jamovi (version 2.3.28) highlighted that
416 there was no significant difference between the two conditions ($t_{(4)} = 1.88$, $p = .134$, Cohen's $d =$
417 $.839$). Similarly, mean RTs for Aware and Unaware trials in the GO condition were contrasted and
418 the statistical analysis (Paired sample two-tailed t-test) revealed that mean RTs for the Aware

419 condition (628.530 ms) and the Unaware condition (675.317 ms) were not statistically different ($t(4)$
420 = - 1.77, $p = .152$, Cohen's $d = -.791$). The behavioral results are depicted in Figure 3.
421 Moreover, in order to verify that the employed paradigm works as planned and that participants
422 performed the task accurately, analysis on catch trials was performed as described in Section 2.8.1
423 *Behavioral data*. As specified above, catch trials were those trials in which all the radii of the stimulus
424 are equally thick. Hence, in those cases, participants should report not to see the thicker radius. As
425 expected, they correctly reported not seeing the thicker radius on average the 96.47% of times
426 ($sd=2.49$) in the Aware GO condition and the 98.36% ($sd=1.89$) in the Aware NOGO condition.
427 Paired sample (two-tailed) t-test revealed no significant difference between the two conditions.
428



429
430 **Figure 3. Behavioral results.** The percentage of Awareness was calculated for both “GO” and “NOGO”
431 conditions (on the left). Mean reaction times were calculated for Aware and Unaware trials only for the “GO”
432 condition (on the right). No significant differences were observed. Error bars represent SEM and gray dots
433 represent individual data points showing the data distribution.


434 4.1.2 EROS results

435 EROS data were pre-processed with a dedicated in-house software, P-POD (Pre-Processing of Optical
436 Data, run in Matlab, version R2013b), as described in Section 2.8. Subsequently, we computed
437 statistical analyses on pre-processed data by means of the dedicated in-house software package Opt-
438 3d.

439 For this pilot study, participants' individual structural MR images could not be acquired, so an
440 estimated MR-based head model was individually created using the Softaxic Optic system (Softaxic,
441 E.M.S., Bologna, Italy) combined with a 3D optical digitizer (Polaris Vicra, NDI, Waterloo, Canada).
442 EROS data were thus co-registered with the estimated MRI using a specific procedure performed in
443 OCP software package (as specified above). Finally, co-registered data were transformed into MNI
444 space for subsequent analyses.

445 For both GO and NOGO conditions, Aware and Unaware trials were contrasted. As shown in Figure
446 4, the Aware-GO vs Unaware-GO contrast replicated the results obtained by Colombari et al., 2024.

447 In this contrast, indeed, we compared conditions in which the motor response was required, thus
448 replicating the task carried out in the previously mentioned experiment. Also in this case, Aware trials
449 elicited a sustained activation of LOC (230 and 332 ms after the stimulus onset), followed by the
450 recurrent activation of the primary visual cortex (V1) and the motor areas (MA) at later stages of
451 stimulus processing.
452 Similarly, contrasting Aware and Unaware trials in the condition where the motor response was not
453 required (i.e., the NOGO condition), greater activation of LOC was elicited in a timing comparable
454 to that of the contrast just mentioned above (i.e., 307 ms after the stimulus presentation). Interestingly,
455 also in this case awareness-related processing elicited activity in the motor areas, 563 ms after the
456 stimulus onset, despite in this condition no response was required, possibly suggesting an inhibition
457 to respond for the NOGO trials.



458 **Figure 4. EROS results.** Statistical parametric maps of the z-score difference computed contrasting Aware
459 and Unaware trials in the GO (upper panel) and NOGO condition (lower panel). Each map represents a 25.6
460 ms interval.
461

462 4.2 Preliminary Discussion

463 The aim of the present pilot study was to assess whether the task and the experimental procedure were
464 suitable to investigate the study's research questions.

465 As described in Section 4.1, the pilot study successfully replicated the trend of activations observed
466 by Colombari et al., 2024, suggesting that the proposed study proves to be feasible in terms of
467 methodology. For the sake of clarity, it is important to point out that the preliminary results reported
468 here do not reach the statistical level of significance. This outcome was expected as data from only 5
469 participants were included in the analysis. For the same reason, we decided not to perform Granger
470 Causality analysis as for this kind of analysis results from 5 participants would have been

471 uninformative. Nevertheless, it was possible to observe that the proposed task could elicit a pattern
472 of activation similar to that observed by Colombari et al., 2024, suggesting that the experimental
473 paradigm proposed to investigate the research questions is suitable.

474

475 **Data availability**

476 Upon acceptance of the Stage 2 registered report, we will share all raw and processed anonymized
477 data as well as study materials publicly available as open data. Pilot raw and processed data can be
478 found on this link: https://osf.io/ebfu3/?view_only=9ec2e6bf32ba4a8bb8b858639ec40a59

479 **Code availability**

480 All analysis codes will be made publicly available upon acceptance of the Stage 2 registered report.

481 **Acknowledgments**

482 The present project is supported by the grant program "Funding Consciousness Research with
483 Registered Reports" and Fondazione Cassa di Risparmio di Verona, Vicenza, Belluno e Ancona
484 "Ricerca scientifica d'eccellenza 2018", "Emergence of Consciousness: From neural dynamics to
485 complex conscious behaviour" grant no. 2018.0861; SS is supported by #NEXTGENERATIONEU
486 (NGEU) and funded by the Ministry of University and Research (MUR), National Recovery and
487 Resilience Plan (NRRP), project MNESYS (PE0000006)—"A Multiscale integrated approach to the
488 study of the nervous system in health and disease" (DN. 1553 11.10.2022), CM is supported by MIUR
489 D.M. 737/2021—"Neural correlates of perceptual awareness: from neural architecture to the
490 preservation of conscious vision in brain tumor patients".

491 **Author contributions**

492 **EC** Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data
493 Curation, Writing - Original Draft, Visualization, Funding Acquisition; **GP** Methodology, Formal
494 Analysis, Investigation, Data Curation, Writing - Review & Editing; **SM** Methodology, Investigation,
495 Writing - Review & Editing **CM** Methodology, Software, Data Curation, Writing - Review & Editing,
496 Supervision; **SS** Conceptualization, Methodology, Resources, Writing - Review & Editing,
497 Supervision, Project administration, Funding acquisition.

498 **Competing interests**

499 The authors declare no competing interests.

500 **References**

- 501 Arridge, S.R. & Schweiger, M. (1995) &title>Sensitivity to prior knowledge in optical
502 tomographic reconstruction&/title> In Chance, B. & Alfano, R.R. (eds), *Optical*
503 *Tomography, Photon Migration, and Spectroscopy of Tissue and Model Media: Theory,*
504 *Human Studies, and Instrumentation*. SPIE, pp. 378–388.
- 505 Aru, J., Bachmann, T., Singer, W., & Melloni, L. (2012) Distilling the neural correlates of
506 consciousness. *Neurosci. Biobehav. Rev.*, **36**, 737–746.
- 507 Baniqued, Pauline L.; Low, Kathy A.; Fabiani, Monica; Gratton, G. (2013) Frontoparietal Traffic
508 Signals: A Fast Optical Imaging Study of Preparatory Dynamics in Response Mode Switching.
509 *J. Cogn. Neurosci.* 25 887–902.,.
- 510 Bola, M. & Doradzińska, Ł. (2021) Perceptual Awareness Negativity—Does It Reflect Awareness
511 or Attention? *Front. Hum. Neurosci.*, **15**, 1–4.
- 512 Chiarelli, A.M., Maclin, E.L., Low, K.A., Fabiani, M., & Gratton, G. (2015) Comparison of
513 procedures for co-registering scalp-recording locations to anatomical magnetic resonance
514 images. *J. Biomed. Opt.*, **20**, 016009.
- 515 Ciupińska, K., Orłowska, W., Zębrowski, A., Łępa, L., Koculak, M., Bola, M., & Wierzchon, M.
516 (2024) The influence of spatial and temporal attention on visual awareness—a behavioral and
517 ERP study. *Cereb. Cortex*, **34**.
- 518 Cohen, M.A., Ortego, K., Kyroudis, A., & Pitts, M. (2020) Distinguishing the Neural Correlates of
519 Perceptual Awareness and Postperceptual Processing. *J. Neurosci.*, **40**, 4925–4935.
- 520 Colombari, E., Parisi, G., Tafuro, A., Mele, S., Mazzi, C., & Savazzi, S. (2024) Beyond primary
521 visual cortex: The leading role of lateral occipital complex in early conscious visual
522 processing. *Neuroimage*, **298**, 120805.
- 523 Dellert, T., Krebs, S., Bruchmann, M., Schindler, S., Peters, A., & Straube, T. (2022) Neural
524 correlates of consciousness in an attentional blink paradigm with uncertain target relevance.
525 *Neuroimage*, **264**, 119679.
- 526 Dellert, T., Müller-Bardorff, M., Schlossmacher, I., Pitts, M., Hofmann, D., Bruchmann, M., &
527 Straube, T. (2021) Dissociating the neural correlates of consciousness and task relevance in
528 face perception using simultaneous EEG-fMRI. *J. Neurosci.*, **41**, 7864–7875.

- 529 Dembski, C., Koch, C., & Pitts, M. (2021) Perceptual awareness negativity: a physiological
530 correlate of sensory consciousness. *Trends Cogn. Sci.*, **25**, 660–670.
- 531 Doradzińska, Ł. & Bola, M. (2024) Early Electrophysiological Correlates of Perceptual
532 Consciousness Are Affected by Both Exogenous and Endogenous Attention. *J. Cogn.*
533 *Neurosci.*, **36**, 1297–1324.
- 534 Förster, J., Koivisto, M., & Revonsuo, A. (2020) ERP and MEG correlates of visual consciousness:
535 The second decade. *Conscious. Cogn.*, **80**.
- 536 Gratton, G. (2000) “Opt-cont” and “opt-3D”: A software suite for the analysis and 3D
537 reconstruction of the event-related optical signal (EROS). *Psychophysiology*,.
- 538 Gratton, G. & Corballis, P.M. (1995) Removing the heart from the brain: Compensation for the
539 pulse artifact in the photon migration signal. *Psychophysiology*, **32**, 292–299.
- 540 GRATTON, G., CORBALLIS, P.M., CHO, E., FABIANI, M., & HOOD, D.C. (1995) Shades of
541 gray matter: Noninvasive optical images of human brain reponses during visual stimulation.
542 *Psychophysiology*,.
- 543 Gratton, G. & Fabiani, M. (1998) Dynamic brain imaging: Event-related optical signal (EROS)
544 measures of the time course and localization of cognitive-related activity. *Psychon. Bull. Rev.*,
545 **5**, 535–563.
- 546 Gratton, G. & Fabiani, M. (2001) Shedding light on brain function: The event-related optical signal.
547 *Trends Cogn. Sci.*, **5**, 357–363.
- 548 Gratton, G. & Fabiani, M. (2003) The event-related optical signal (EROS) in visual cortex:
549 Replicability, consistency, localization, and resolution. *Psychophysiology*, **40**, 561–571.
- 550 Gratton, G. & Fabiani, M. (2010) Fast optical imaging of human brain function. *Front. Hum.*
551 *Neurosci.*, **4**, 1–9.
- 552 Gratton, G., Fabiani, M., Corballis, P.M., Hood, D.C., Goodman-Wood, M.R., Hirsch, J., Kim, K.,
553 Friedman, D., & Gratton, E. (1997) Fast and localized event-related optical signals(EROS) in
554 the human occipital cortex: Comparisons with the visual evoked potential and fMRI.
555 *Neuroimage*, **6**, 168–180.
- 556 Gratton, G., Goodman-Wood, M.R., & Fabiani, M. (2001) Comparison of neuronal and
557 hemodynamic measures of the brain response to visual stimulation: An optical imaging study.
558 *Hum. Brain Mapp.*, **13**, 13–25.

- 559 Gratton, G., Low, K.A., Maclin, E.L., Brumback, C.R., Gordon, B., & Fabiani, M. (2006) Time
560 course of activation of human occipital cortex measured with the event-related optical signal
561 (EROS). *Opt. InfoBase Conf. Pap.*, 5–8.
- 562 Gratton, G., Sarno, A., Maclin, E., Corballis, P.M., & Fabiani, M. (2000) Toward noninvasive 3-D
563 imaging of the time course of cortical activity: Investigation of the depth of the event-related
564 optical signal. *Neuroimage*, **11**, 491–504.
- 565 Hatamimajoumerd, E., Ratan Murty, N.A., Pitts, M., & Cohen, M.A. (2022) Decoding perceptual
566 awareness across the brain with a no-report fMRI masking paradigm. *Curr. Biol.*, **32**, 4139-
567 4149.e4.
- 568 Hense, A., Peters, A., Bruchmann, M., Dellert, T., & Straube, T. (2024) Electrophysiological
569 correlates of sustained conscious perception. *Sci. Rep.*, **14**, 1–11.
- 570 Kiebel, S.J., Poline, J.B., Friston, K.J., Holmes, A.P., & Worsley, K.J. (1999) Robust smoothness
571 estimation in statistical parametric maps using standardized residuals from the general linear
572 model. *Neuroimage*, **10**, 756–766.
- 573 Knight, R.S., Chen, T., Gratton, G., Fabiani, M., Savazzi, S., Mazzi, C., & Beck, D.M. (2024)
574 Neuropsychologia Bypassing input to V1 in visual awareness : A TMS-EROS investigation.
575 *Neuropsychologia*, **198**, 108864.
- 576 Koivisto, M., Lähteenmäki, M., Sørensen, T.A., Vangkilde, S., Overgaard, M., & Revonsuo, A.
577 (2008) The earliest electrophysiological correlate of visual awareness? *Brain Cogn.*, **66**, 91–
578 103.
- 579 Koivisto, M. & Revonsuo, A. (2010) Event-related brain potential correlates of visual awareness.
580 *Neurosci. Biobehav. Rev.*, **34**, 922–934.
- 581 Koivisto, M., Revonsuo, A., & Lehtonen, M. (2006) Independence of visual awareness from the
582 scope of attention: An electrophysiological study. *Cereb. Cortex*, **16**, 415–424.
- 583 Koivisto, M., Salminen-Vaparanta, N., Grassini, S., & Revonsuo, A. (2016) Subjective visual
584 awareness emerges prior to P3. *Eur. J. Neurosci.*, **43**, 1601–1611.
- 585 Kronemer, S.I., Aksen, M., Ding, J.Z., Ryu, J.H., Xin, Q., Ding, Z., Prince, J.S., Kwon, H., Khalaf,
586 A., Forman, S., Jin, D.S., Wang, K., Chen, K., Hu, C., Agarwal, A., Saberski, E., Mohammad,
587 S., Wafa, A., Morgan, O.P., Wu, J., Christison-lagay, K.L., Hasulak, N., Morrell, M., Urban,
588 A., Constable, R.T., & Pitts, M. (2022) Human visual consciousness involves large scale

- 589 cortical and subcortical networks independent of task report and eye movement activity.
- 590 Leung, P., Franconeri, S., Grabowecky, M., & Suzuki, S. (2009) Rapid eye-fixation training
591 without eyetracking **16**, 491–496.
- 592 Liu, Y., Paradis, A.L., Yahia-Cherif, L., & Tallon-Baudry, C. (2012) Activity in the lateral occipital
593 cortex between 200 and 300 ms distinguishes between physically identical seen and unseen
594 stimuli. *Front. Hum. Neurosci.*, **6**, 1–9.
- 595 Low, K.A., Leaver, E., Kramer, A.F., Fabiani, M., & Gratton, G. (2006) Fast optical imaging of
596 frontal cortex during active and passive oddball tasks. *Psychophysiology*, **43**, 127–136.
- 597 Makeig, S. & Jung, T. (2000) Independent Component Analysis of Simulated ERP Data. *Brain*, 1–
598 24.
- 599 Mazzi, C., Mazzeo, G., & Savazzi, S. (2020) Late Positivity Does Not Meet the Criteria to be
600 Considered a Proper Neural Correlate of Perceptual Awareness. *Front. Syst. Neurosci.*, **14**, 1–
601 14.
- 602 Medvedev, A. V., Kainerstorfer, J., Borisov, S. V., Barbour, R.L., & VanMeter, J. (2008) Event-
603 related fast optical signal in a rapid object recognition task: Improving detection by the
604 independent component analysis. *Brain Res.*, **1236**, 145–158.
- 605 Medvedev, A. V., Kainerstorfer, J.M., Borisov, S. V., Gandjbakhche, A.H., & VanMeter, J. (2010)
606 “Seeing” electroencephalogram through the skull: imaging prefrontal cortex with fast optical
607 signal. *J. Biomed. Opt.*, **15**, 061702.
- 608 Oldfield, R. (1971) The assessment and analysis of handedness: the Edinburgh Inventory.
609 *Neuropsychologia*,.
- 610 Parisi, G., Mazzi, C., Colombari, E., Chiarelli, A.M., Metzger, B.A., Marzi, C.A., & Savazzi, S.
611 (2020) Spatiotemporal dynamics of attentional orienting and reorienting revealed by fast
612 optical imaging in occipital and parietal cortices. *Neuroimage*, **222**, 117244.
- 613 Pitts, M.A., Padwal, J., Fennelly, D., Martínez, A., & Hillyard, S.A. (2014) Gamma band activity
614 and the P3 reflect post-perceptual processes, not visual awareness. *Neuroimage*, **101**, 337–350.
- 615 Proulx, N., Samadani, A.A., & Chau, T. (2018) Quantifying fast optical signal and event-related
616 potential relationships during a visual oddball task. *Neuroimage*, **178**, 119–128.
- 617 Railo, H., Revonsuo, A., & Koivisto, M. (2015) Behavioral and electrophysiological evidence for
618 fast emergence of visual consciousness. *Neurosci. Conscious.*, **2015**, 1–12.

- 619 Schelonka, K., Grauly, C., Canseco-gonzalez, E., & Pitts, M.A. (2017) ERP signatures of
620 conscious and unconscious word and letter perception in an inattentional blindness paradigm.
621 *Conscious. Cogn.*, **54**, 56–71.
- 622 Schlossmacher, I., Dellert, T., Pitts, M., Bruchmann, M., & Straube, T. (2020) Differential effects
623 of awareness and task relevance on early and late ERPs in a no-report visual oddball paradigm.
624 *J. Neurosci.*, **40**, 2906–2913.
- 625 Seth, A.K., Barrett, A.B., & Barnett, L. (2015) Granger causality analysis in neuroscience and
626 neuroimaging. *J. Neurosci.*, **35**, 3293–3297.
- 627 Shafto, J.P. & Pitts, M.A. (2015) Neural signatures of conscious face perception in an inattentional
628 blindness paradigm. *J. Neurosci.*, **35**, 10940–10948.
- 629 Toscano, J.C., Anderson, N.D., Fabiani, M., Gratton, G., & Garnsey, S.M. (2018) The time-course
630 of cortical responses to speech revealed by fast optical imaging. *Brain Lang.*, **184**, 32–42.
- 631 Tse, C.Y. & Penney, T.B. (2007) Preattentive change detection using the event-related optical
632 signal. *IEEE Eng. Med. Biol. Mag.*, **26**, 52–58.
- 633 Tse, C.Y., Shum, Y.H., Xiao, X.Z., & Wang, Y. (2021) Fronto-occipital mismatch responses in pre-
634 attentive detection of visual changes: Implication on a generic brain network underlying
635 Mismatch Negativity (MMN). *Neuroimage*, **244**, 118633.
- 636 Tsuchiya, N., Wilke, M., Frässle, S., & Lamme, V.A.F. (2015) No-Report Paradigms: Extracting
637 the True Neural Correlates of Consciousness. *Trends Cogn. Sci.*, **19**, 757–770.
- 638 Vanni, S., Revonsuo, A., Saarinen, J., & Hari, R. (1996) Visual awareness of objects correlates with
639 activity of right occipital cortex. *Neuroreport*, **8**, 183–186.
- 640 Wolf, M., Wolf, U., Choi, J.H., Toronov, V., Paunescu, L.A., Michalos, A., & Gratton, E. (2003)
641 Fast cerebral functional signal in the 100-ms range detected in the visual cortex by frequency-
642 domain near-infrared spectrophotometry. *Psychophysiology*, **40**, 521–528.
- 643 Wolf, U., Wolf, M., Toronov, V., Michalos, A., Paunescu, L.A., & Gratton, E. (2014) Detecting
644 cerebral functional slow and fast signals by frequency-domain near-infrared spectroscopy
645 using two different sensors. In *Biomedical Optical Spectroscopy and Diagnostics (2000)*,
646 *Paper TuF10*. The Optical Society, p. TuF10.
- 647 Worsley, K.J., Poline, J.B., Vandal, A.C., & Friston, K.J. (1995) Tests for distributed, nonfocal
648 brain activations. *Neuroimage*, **2**, 183–194.

- 649 Ye, M., Wang, A., Liang, H., & Liu, X. (2024) Late positivity correlates with subjective reports:
650 Evidence from the low-frequency and high-frequency reporting tasks. *Neuroscience*,.
- 651 Zhao, L., Matloff, W., Shi, Y., Cabeen, R.P., & Toga, A.W. (2022) Mapping Complex Brain
652 Torque Components and Their Genetic Architecture and Phenomic Associations in 24 , 112
653 Individuals. *Biol. Psychiatry*, **91**, 753–768.
- 654