RAG suppresses group 2 innate lymphoid cells

Aaron M. Ver Heul[1], Madison Mack[2], Lydia Zamidar[3,4,5,6], Masato Tamari[3,4,5,6,], Ting-Lin Yang[8], Anna M. Trier[8], Do-Hyun Kim[9,10], Hannah Janzen-Meza[1], Steven J. Van Dyken[9], Chyi-Song Hsieh[11], Jenny M. Karo[12], Joseph C. Sun[12], Brian S. Kim[3,4,5,6,7,*]

[1]Division of Allergy and Immunology, Department of Medicine, Washington University School of Medicine, St. Louis, MO 63130, USA

[2]Immunology & Inflammation Research Therapeutic Area, Sanofi, Cambridge, MA 02141, USA

[3]Kimberly and Eric J. Waldman Department of Dermatology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[4]Mark Lebwohl Center for Neuroinflammation and Sensation, Icahn School of Medicine at Mount Sinai, New York, NY 10019, USA

[5]Marc and Jennifer Lipschultz Precision Immunology Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[6]Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[7]Allen Discovery Center for Neuroimmune Interactions, Icahn School of Medicine at Mount Sinai 10019

[8]Division of Dermatology, Department of Medicine, Washington University School of Medicine, St. Louis, MO 63130, USA

[9]Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, MO 63130, USA

[10]Department of Life Science, College of Natural Sciences, Hanyang University, Seoul 04763, Republic of Korea

[11]Division of Rheumatology, Department of Medicine, Washington University School of Medicine, St. Louis, MO, USA.

[12]Immunology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA; Immunology and Microbial Pathogenesis Program, Graduate School of Medical Sciences, Weill Cornell Medical College, New York, NY 10065, USA.

*Correspondence: Brian S. Kim, MD, MTR, Icahn School of Medicine at Mount Sinai, 1425 Madison Avenue, New York, NY, 10029. E-mail: brian.kim3@mountsinai.org

ABSTRACT

Antigen specificity is the central trait distinguishing adaptive from innate immune function. Assembly of antigen-specific T cell and B cell receptors occurs through V(D)J recombination mediated by the Recombinase Activating Gene endonucleases RAG1 and RAG2 (collectively called RAG). In the absence of RAG, mature T and B cells do not develop and thus RAG is critically associated with adaptive immune function. In addition to adaptive T helper 2 (Th2) cells, group 2 innate lymphoid cells (ILC2s) contribute to type 2 immune responses by producing cytokines like Interleukin-5 (IL-5) and IL-13. Although it has been reported that RAG expression modulates the function of innate natural killer (NK) cells, whether other innate immune cells such as ILC2s are affected by RAG remains unclear. We find that in RAG-deficient mice, ILC2 populations expand and produce increased IL-5 and IL-13 at steady state and contribute to increased inflammation in atopic dermatitis (AD)-like disease. Further, we show that RAG modulates ILC2 function in a cell-intrinsic manner independent of the absence or presence of adaptive T and B lymphocytes. Lastly, employing multiomic single cell analyses of RAG1 lineage-traced cells, we identify key transcriptional and epigenomic ILC2 functional programs that are suppressed by a history of RAG expression. Collectively, our data reveal a novel role for RAG in modulating innate type 2 immunity through suppression of ILC2s.

INTRODUCTION

Atopic disorders such as atopic dermatitis (AD), asthma, and food allergy are associated with T helper type 2 (Th2) cell responses, elevated production of the type 2 cytokines interleukin(IL)-4, IL-5, and IL-13, and induction of immunoglobulin(Ig)E[1–4].Classically, this allergic inflammatory cascade is believed to originate with antigenic stimulation of T cell receptors on adaptive T cells, which in turn results in the production of IgE from B and plasma cells capable of binding the same antigen. Indeed, presence of antigen-specific IgE reactivity is a hallmark across atopic disorders.[5] Thus, for decades, antigen-specific adaptive Th2 cell responses have been the primary focus of investigation in the pathogenesis of atopic diseases. However, recent studies indicate that innate immune cells are sufficient to not only drive allergic pathology, but also amplify adaptive Th2 cell responses[6–9]. These studies suggest that innate immune mechanisms may play a larger role in driving atopic inflammation than previously recognized.

Innate lymphoid cells (ILCs), while lacking antigen receptors generated by recombination activating gene proteins RAG1 and RAG2 (collectively called RAG), are the innate counterparts of T cells. For example, ILC2s mirror adaptive Th2 cells in their developmental requirements, cytokine profiles, and effector functions[10]. Unlike classical T cells, ILC2s are concentrated at barrier surfaces to rapidly respond to microbial and environmental stimuli. ILC2s are key mediators of inflammatory skin conditions like AD[11–13]. Indeed, in murine models of AD-like disease, type 2 skin inflammation can still occur despite the absence of adaptive T cells, but is further reduced after depletion of ILC2s[12,13]. Furthermore, recent studies have shown that ILC2s harbor non-redundant functions in the presence of the adaptive immune system in the setting of anti-helminth immunity[14,15]. These findings suggest that ILC2 dysfunction may also uniquely contribute to the pathogenesis of atopic diseases, independent of adaptive immunity. However, the cell-intrinsic mechanisms that drive ILC2 dysregulation remain poorly understood.

ILC2s were originally discovered due to their capacity to orchestrate multiple allergic

pathologies in immunocompromised mice, most notably in RAG-deficient mice that lack T and B

cells[16–22]. These discoveries fundamentally redefined our understanding of allergic diseases and

placed a major focus on ILC2s as potential drivers of human allergic disease. However, despite

ILC2s not requiring RAG expression for their development, fate mapping studies in mice have

demonstrated that up to 60% of ILC2s have historically expressed RAG1 during development[23,24].

Although previous work has described roles of RAG beyond antigen receptor recombination in

developing T and B cells[25–27] and NK cells[24], how this developmental expression of RAG impacts

ILC2s remains unclear.

By directly comparing RAG-deficient and RAG-sufficient mice, we unexpectedly found

enhanced AD-like disease in RAG-deficient mice, despite the lack of adaptive lymphocytes to

contribute to AD-like inflammation. Using splenocyte replenishment and bone marrow chimeras,

we show that RAG suppresses ILC2 activation and expansion in a cell-intrinsic manner.

Employing a RAG1-lineage reporter mouse line, we performed simultaneous single-cell multiomic

RNA and ATAC sequencing to show that RAG fate-mapped ILC2s display unique transcriptional

and epigenomic alterations consistent with the suppression of effector cytokine production.

Collectively, our studies reveal evolutionarily conserved regulatory functions of RAG within innate

lymphocytes, extending beyond the generation of antigen receptors in adaptive lymphocytes.

RESULTS

94
RAG deficiency leads to expansion and activation of ILC2s

AD-like disease can be elicited in the skin of mice with repeated application of the topical vitamin

D analog calcipotriol (MC903)[28]. Although it has been previously demonstrated that MC903 can

induce AD-like disease in RAG-deficient mice that lack T and B cells, in part via ILC2

activation[12,13], the relative contributions of ILC2s and the adaptive lymphocyte compartment have

not been rigorously evaluated. We hypothesized that the presence of Th2 cells, in addition to

ILC2s, would lead to enhanced AD-like disease in an additive fashion. In testing this, we evaluated

both RAG1-sufficient wild-type (WT) mice and RAG1-deficient $Rag1^{-/-}$ mice in the setting of AD-

like disease (**Fig. 1A**). Unexpectedly, we observed that $Rag1^{-/-}$ mice developed increased ear skin

thickness (**Fig. 1B**) and increased absolute numbers and proportion of ILC2s in the skin-draining

lymph nodes (sdLNs) compared to control WT mice (**Fig. 1C,D; S1A,B**). Furthermore, a larger

proportion of ILC2s from $Rag1^{-/-}$ mice exhibited production of both IL-5 (**Fig. 1E, S1C**) and IL-13

(**Fig. 1F, S1C**). Our findings indicated that RAG1 deficiency results in paradoxically worse AD-

like disease in association with enhanced ILC2 expansion and activation.

To determine whether this phenomenon was specific to AD-like pathological conditions,

we next examined the sdLNs in $Rag1^{-/-}$ and lymphocyte-sufficient $Rag^{+/-}$ littermate control mice in

the absence of disease (**Fig. 1G**). We found that the absolute number and frequency of ILC2s

was increased at steady state in $Rag1^{-/-}$ sdLNs (**Fig. 1H,I**) and that a higher proportion of these

ILC2s produced both IL-5 (**Fig. 1J**) and IL-13 (**Fig. 1K**) compared to WT controls following ex vivo

stimulation and intracellular cytokine staining. The RAG recombinase requires both RAG1 and

RAG2 components to successfully rearrange a functional antigen receptor in adaptive

lymphocytes[29]. Thus, to test whether our findings are specific to RAG1, or related to function of

the overall RAG complex, we similarly examined the steady-state profile of ILC2s in $Rag2^{-/-}$ mice

(**Fig. S2A**). Deficiency of RAG2 led to an expansion of ILC2s in the sdLNs (**Fig. S2B,C**) and

increased proportions of ILC2s expressing IL-5 (**Fig. S2D**) and IL-13 (**Fig. S2E**) similar to $Rag1^{-/-}$

mice. Collectively, these findings suggest that the RAG recombinase modulates ILC2 function at

steady state and during type 2 inflammation. However, whether the hyperactive ILC2 phenotype

is due to a cell-intrinsic process or simply due to the absence of T and B cells was unclear.

ILC2 suppression by RAG is cell intrinsic

Given the importance of the adaptive lymphocyte compartment in shaping the secondary lymphoid organ repertoire, we next wanted to examine whether the presence of adaptive lymphocytes could restore ILC2 homeostasis in RAG-deficient mice. To test this, we created splenocyte chimera mice by reconstituting both $Rag1^{-/-}$ and control WT mice with splenocytes containing T and B cells from WT donor mice (**Fig. 2A**). We first assessed the overall level of immune reconstitution in the recipient mice and found fully restored proportions of CD4$^+$ (**Fig. S3A**) and CD8$^+$ (**Fig. S3B**) T cells in the spleens of recipient $Rag1^{-/-}$ mice, although B cell numbers remained significantly lower than in WT mice (**Fig. S3C**). Upon induction of AD-like disease, we found that the $Rag1^{-/-}$ mice still exhibited increased ear skin thickness (**Fig. 2B**), enhanced expansion of ILC2s (**Fig. 2C,D**), and increased proportions of ILC2s expressing IL-5 (**Fig. 2E**) and IL-13 (**Fig. 2F**) in the sdLNs. Interestingly, we found significantly higher proportions of eosinophils in the spleens of $Rag1^{-/-}$ recipient mice (**Fig. S3D**), possibly reflecting the increased IL-5 production we observed from ILC2s. These findings indicate that the mere introduction of exogenous T and B cells is not sufficient to suppress ILC2 dysregulation in the setting of RAG deficiency.

To further test whether this phenotype is mediated by cell-intrinsic RAG expression, we next generated mixed bone marrow (BM) chimeras. We harvested BM from congenic CD90.1$^+$ WT and CD90.2$^+$ $Rag1^{-/-}$ donor mice on a CD45.2$^+$ background and reconstituted sub-lethally irradiated CD45.1$^+$ congenic WT recipients with a 50:50 mixture of WT:$Rag1^{-/-}$ BM (**Fig. 2G**). After confirming reconstitution of donor immune cells in the sdLN (**Fig. S4A,D**), we examined the frequency and activity of ILC2s in the sdLNs based on whether they originated from WT (CD90.1$^+$) or $Rag1^{-/-}$ (CD90.2$^+$) donors (**Fig. S4B,C**). Strikingly, of the total donor ILC2s, the majority were derived from $Rag1^{-/-}$ donors (**Fig. 2F,G**). This was not due to differences in overall donor

148 reconstitution, since measuring all Lin⁻ cells revealed WT donor cells outnumbered those from

149 *Rag1*^(-/-) donors (**Fig. S4E**). Of the total IL-5- (**Fig. 2H,I,K**) and IL-13-expressing (**Fig. 2 H,J,L**) ILCs,

150 the majority were also derived from *Rag1*^(-/-) donors. Taken together, these data suggest that cell-

151 intrinsic RAG activity in ILC2s may limit their capacity to expand and become activated.

A history of RAG expression marks a subpopulation of ILC2s in the skin draining lymph node

In contrast to resting naïve T cells, ILC2s resemble activated Th2 cells at steady state

based on their transcriptomic and epigenomic profiles[30,31]. While both T cells and ILC2s exhibit

historical RAG expression[23], they do not actively express the protein in their mature state[32]. Taken

together, these findings provoke the hypothesis that ILC2s are regulated by RAG early in

development to imprint alterations that influence their activity as mature cells. To distinguish ILC2s

as either having a history of RAG expression or not, we utilized a RAG lineage tracing system,

whereby a *Rag1*^(Cre) mouse was crossed to a reporter mouse expressing tandem dimer red

fluorescent protein (tdRFP) in a Cre-dependent manner from the *Rosa26* locus (**Fig. 3A**)[24,33]. This

system allowed us to compare RAG-experienced (RAG^(exp)) and RAG-naïve (RAG^(naïve)) lymphoid

cells, including ILC2s, simultaneously originating from the same immunocompetent host, thus

removing confounders inherent in knockout and chimera experiments. Analysis of sdLN from the

reporter mice revealed that nearly all CD4⁺ T cells, CD8⁺ T cells, and B220⁺ B cells expressed

tdRFP (positive history of Cre expression from the *Rag1* locus), consistent with the known

requirement of RAG expression for their development (**Fig. 3B-D,G**). We also examined NK cells,

since certain subsets of NK cells are known to express RAG during their development[24], and we

observed that 60% of NK cells were tdRFP⁺, similar to previous findings (**Fig. 3E,G**)[24]. In the ILC2

population of the sdLN, around 50% were tdRFP⁺ (**Fig. 3F,G**), similar to proportions of RAG fate-

mapped ILC2s previously observed in the fat[24] and lung[23,24]. These findings demonstrate that

there are heterogeneous populations of ILC2s marked by differential tdRFP[+] fate mapping.

Importantly, this provided us with the possibility to profile these different ILC2 subsets based on

Rag1[Cre]-activated expression of tdRFP.

176 Multiomic profiling enhances the detection of rare tissue-specific ILC2s

Transient RAG expression early in lymphoid development leads to well-characterized,

durable effects on B and T cell development and function mainly through successful genomic

rearrangement of antigen receptors. Yet, our data indicate that RAG expression also imprints

phenotypic changes on ILC2s, which can develop independently of functional antigen receptors,

provoking the hypothesis that RAG expression may affect broader epigenomic and transcriptional

programs. Furthermore, our data indicate that the impact of RAG on ILC2 function has

implications for AD-like skin inflammation, suggesting a persistent effect that modulates

phenotypes of type 2 inflammation.

To test these hypotheses, we performed combined single-nuclei RNA sequencing

(snRNA-seq) and ATAC sequencing (snATAC-seq) of sdLN cells from RAG fate-mapped mice at

steady state and in the setting of AD-like disease (**Fig. S5A,B**). Because fluorescence-activated

cell sorting (FACS) can cause physical stress, cell loss, and contamination, which can introduce

unwanted perturbations in target cells, instead we utilized gentle initial negative selection by

magnetic activated cell sorting (MACS) to remove most B and T cells and monocytes prior to

sequencing (**Fig. S5A**). This allowed us to enrich for innate immune cell populations prior to

sequencing. Further, we used the gene encoding tdRFP as a barcode to differentiate between

RAG[exp] (RAG fate map-positive) and RAG[naïve] (RAG fate map-negative) ILC2s at the single-cell

level (**Fig. 5A**). The multiomic data was analyzed using recently developed pipelines in Cell

Ranger, Seurat[34–36], and Signac[37], and sequenced cells were further filtered computationally to

enrich for ILCs, as in previous studies (see methods)[38].

In addition to gene expression (GEX) information derived from snRNA-seq (**Fig. 4A**), we

calculated a "gene activity" (GA) score based on chromatin accessibility at gene loci (**Fig. 4B**)

from the corresponding snATAC-seq dataset[37]. Clustering the cells with each data subset alone

and in combination using weighted nearest neighbor (WNN) analysis, we identified six clusters

(**Fig. 4C**) that demonstrated consistent differences in cellular markers based on both metrics of

GEX and GA (**Fig. 4D,E; Table S1**). Additionally, top markers for each cluster clearly

differentiated each cell type (**Fig. 4F**). Despite successful ILC2 enrichment via MACS depletion

for lineage markers and computational filtering (see methods), our data set included non-ILC2

populations determined by gene expression to be T cells, dendritic cells, B cells, and NK cells

(**Fig. 4A,F; Table S1**), which allowed for broader multidimensional comparisons while studying

this ILC2-enriched data set.

To further complement the GEX and GA assays, we utilized another method of detecting

cell-specific marker genes, whereby chromatin regions that are differentially accessible (DA), or

open, in each cluster could be linked by their physical proximity to specific genes (**Fig. 4G, S6**;

Table S2; see methods). Comparing the top 100 GEX, GA, and DA markers in the ILC2

population, we identified a multiomic ILC2 signature of 235 unique genes (**Fig. 4H; Table S3,4**).

While there was some overlap between each respective set, the multiomic approach enabled

more extensive identification of an ILC2 gene program than through either snRNA-seq or

snATAC-seq alone. The analysis revealed a variety of canonical ILC2-associated genes specific

to the ILC2 cluster (**Fig. 4I, S6,7**) including the ILC2-activating cell surface receptors *Icos*[39,40],

Il2ra[41,42], *Il18r1*[38,43,44], *Nmur*[14,15,45–47], and *Il1rl1* (encoding the receptor for IL-33)[11,12,48,49], as well

as transcription factors such as *Gata3*[19,50–53], *Bcl11b*[54–57], *Maf*[19,58,59], *Ets1*[60], and *Rora*[38,51,61–63], all

previously shown to be important for ILC2 development and/or function.

Expression of some secreted proteins can be difficult to capture in droplet-based snRNA-

seq experiments due to low transcript levels and relatively shallow sequencing depth. With the

addition of the complementary GA and DA assays from snATAC-seq, our analysis identified *Il5*

(**Fig. 4I, S6B**), a canonical ILC2 cytokine, in the DA assay, while in the GA assay we found

Bmp7[64,65], which has been shown to be secreted by ILC2s to influence browning of adipose tissue.

Additionally, we identified the secreted chemokine *Ccl1* as an ILC2 marker gene[66–69], which along

with its cognate receptor *Ccr8* (also an ILC2 marker in our analysis)[44,67,70], participates in a feed-

forward circuit to drive ILC2 recruitment and expansion[71]. Thus, our findings demonstrate how

genetic barcoding, combining transcriptomic and epigenomic analyses, and cross-validation

across many published studies can yield new insights while providing internal control measures

to elevate the rigor, robustness, and confidence of identifying gene signatures of rare populations

such as ILC2s at the single-cell level.

ILC2s with a history of RAG expression are epigenomically suppressed

As noted above, barcoding the ILC2s afforded the opportunity to transcriptionally and

epigenomically profile ILC2s under identical developmental conditions by dividing the ILC2 cluster

into RAG^{exp} and RAG^{naïve} populations. We hypothesized that RAG^{exp} ILC2s would have a distinct

transcriptional profile compared to ILC2s without any history of RAG expression. To test this, we

calculated differentially expressed genes (DEGs) for the ILC2 cluster by *Rag1* fate-map status.

Genes with higher expression in RAG^{exp} cells relative to RAG^{naïve} cells had positive fold change

values, and vice versa, with genes relatively increased in RAG^{naïve} cells having negative values

(**Fig. S8A, Table S5**). Using gene set enrichment analysis (GSEA)[72,73] on the ranked list of DEGs,

we found that gene sets generally representing lymphocyte activation and differentiation were

suppressed in RAG^{exp} ILC2s compared to RAG^{naïve} ILC2s (**Fig. S8B,C, Table S6**), consistent with

our previous observations that ILC2s are expanded and more activated in RAG-deficient mice

relative to WT mice.

We next employed newly described methodologies[37,74] that quantify associations between

open chromatin peaks and the expression of nearby genes to describe the functional regulomes

of both RAG[exp] and RAG[naïve] ILC2s (**Fig. 5B**). In this analysis, each ATAC peak can be linked to

multiple genes, and each gene to multiple peaks, generating a list of "gene-to-peak links" or GPLs

(see methods). For each gene, we interpreted the number of corresponding GPLs as a

quantitative representation of the regulome activity for that gene. Considering RAG[exp] and

RAG[naïve] cells as two separate populations, we generated two lists of GPLs (**Table S7**) defining

functional regulomes for each population. We focused our analysis on the functional regulomes

of ILC2s by filtering the GPL lists for the 235 unique ILC2 genes identified in our multiomic gene

set (**Fig. 4H; Table S8**), then calculated the difference in GPLs between RAG[exp] and RAG[naïve]

cells for each gene and sorted them; genes displaying greater numbers of GPLs in the RAG[naïve]

population are at the top, and genes with more GPLs in the RAG[exp] population are at the bottom

(**Fig. 5C; Table S9**).

We found that the ILC2 marker genes segregating toward the top of this list, corresponding

to enhanced epigenomic activity in the RAG[naïve] cells, tended to be genes previously identified to

play positive roles in development, expansion, and activation of lymphoid cells. These included

transcriptional regulators such as *Tox*[75–78], *Rora*[38,51,61–63], *Maf*[19,58,59], and *Gata3*[19,50–52], which are

involved in early differentiation of both ILCs and lymphocytes (**Fig. S9A**)*.* Indeed, epigenetic

activation of the *Gata3* locus is recognized to play a critical role in development of both ILC2s[79]

and Th2 cells[80,81]. Additionally, surface receptors known to drive ILC2 activation upon stimulation

including *Il18r1*[38,43,44], *Il1rl1*[21,38,43,82,83], and *Icos*[39,40] had increased functional regulome activity in

RAG[naïve] ILC2s. In contrast, genes with more GPLs in the RAG[exp] ILC2s tended to be associated

with suppressive functions. For example, *Ndfip1* (**Fig. S9B**) encodes a regulatory protein that

enhances activity of the ubiquitin ligase ITCH to negatively regulate inflammation[84,85] and has

been associated with asthma risk in GWAS studies[86]. *Dusp1* partially mediates glucocorticoid

effects through its ability to negatively regulate inflammation[87,88], is associated with eczema by

GWAS[89], and has recently been shown to mark an anti-inflammatory set of ILCs[69]. Last, *Asxl1*

encodes a tumor suppressor that inhibits clonal hematopoiesis through its epigenomic regulatory

effects in both mice and humans[90–93]. Collectively, our GPL analysis stratifies the ILC2 gene

signature based on RAG experience, where genes associated with ILC2 expansion and activation

are poised in RAG^{naïve} cells, while genes associated with suppressive effects are poised in RAG^{exp}

cells.

We expanded our multiomic analysis to infer information about transcription factor (TF)

activity from open chromatin regions in our snATAC data. We used the chromVAR package[94],

which finds known TF binding motifs in open chromatin regions in each cell, to identify TF motifs

enriched in each cell cluster (**Fig. S10A,B**, **Table S10**). The enriched motifs were consistent with

known functional roles of associated TFs in each cell type. For example, in the NK cell cluster we

found enriched motifs recognized by the TFs EOMES and T-bet (encoded by *Eomes* and *Tbx21*,

respectively), which are critical for development of NK cells[95]. A limitation of this analysis is that

while TF motif accessibility can be inferred from open chromatin in snATAC data, which TFs are

bound to the identified accessible sites is not known. We reasoned that complementary gene

expression information from our multiomic data could mitigate this limitation in part by comparing

accessibility of TF binding motifs to expression levels of corresponding TFs (**Fig. S10C**). Indeed,

motifs for both RORα and RORγ (encoded by *Rora* and *Rorg*, respectively), which share a

common DNA binding 5'-AGGTCA-3' half site, have similar calculated accessibilities in both the

ILC2 and NK cell clusters. Yet only *Rora* is expressed at appreciable levels, and only in ILC2s,

consistent with its critical role in ILC2 development[61,62]. In contrast, ILC2 development is not

dependent on *Rorg* expression, and neither RORα nor RORγ plays a major role in NK cells. Taken

together, this analysis confirms the known role of *Rora* in ILC2s and highlights how matched

multiomic chromatin accessibility and gene expression data can clarify ambiguities inherent in TF

enrichment analyses.

The broad effects of RAG expression on ILC2 transcriptional regulomes we observed (**Fig.**

5C) led us to hypothesize that distinct cohorts of TFs may contribute to the differences observed

between RAG[naïve] and RAG[exp] ILC2s. To test this hypothesis, we analyzed the open chromatin

regions in GPLs unique to each RAG fate mapped ILC2 population using the FindMotifs function

in Signac[37], which returns a ranked list of enriched motifs corrected for background presence of

each motif in all cells. In both RAG[naïve] and RAG[exp] ILC2s, we identified enriched TF motifs (**Fig.**

S10D, Table S11) that are GC rich regions recognized by a large family of C2H2 zinc finger TFs,

particularly the Krüppel-like factors (KLFs), which are well-established as key regulators of

lymphocyte development[96,97]. Given the strong sequence similarities of the identified TF motifs,

we turned to the matched gene expression data to clarify which TFs may be available to engage

the accessible binding sites. Of the eleven unique TFs identified, only six were detected in the

gene expression assay (**Fig. S10E**). We observed much higher expression of *Klf2*, *Klf6*, and *Klf12*

in Rag[exp] ILC2s compared to RAG[naïve] ILC2s (**Fig. S10E**). Notably, all three of these TFs have

been associated with reduced cellular proliferation and/or activation[98–100]. *Klf2* expression plays a

key role in T cell quiescence[101,102], and both *Klf2* and *Klf6* were recently identified as markers of

"quiescent-like" skin resident ILCs[69]. In contrast, although detected in a smaller fraction of cells in

our data, we found *Klf7* expression was higher in RAG[naïve] ILC2s compared to Rag[exp] ILC2s (**Fig.**

S10E). Increased expression of *Klf7* has been shown to enhance survival of early thymocytes

and is a predictor of poor outcome in acute lymphoblastic leukemia[103]. Collectively, these findings

link the relatively activated or suppressed epigenomic and transcriptomic states of RAG[naïve] and

Rag[exp] ILC2s, respectively, to distinct cohorts of homeostatic TFs.

A history of RAG expression modulates ILC2 epigenomes at steady state and in AD-like
inflammation

Although our GPL and TF analyses revealed a suppressive effect of RAG expression on

ILC2 gene programs, we did not account for the additional variable of disease state in the initial

analysis. To test whether RAG expression promotes a suppressive epigenomic program in ILC2s

that is durable in the setting of inflammation, we first recalculated GPLs after splitting our dataset

by both history of RAG expression (naïve vs. experienced) and disease (steady state vs. AD-like

disease) to yield four lists of GPLs (**Fig 6A; Table S12**). When we examined the intersection, or

overlap, of peaks from ILC2 GPLs (**Table S13**), several notable patterns emerged (**Fig. 6B**). First,

the largest set of peaks was shared by all RAG[naïve] cells (gray bar), regardless of disease state,

with the next largest peak sets belonging to either steady state or AD-like disease in the RAG[naïve]

cells. Second, there was a large set of peaks shared by all RAG[exp] cells (red bar). Third, the

intersections corresponding to disease states (steady state – yellow, AD-like disease – dark red),

had relatively few unique peaks. These findings suggest that early exposure to RAG expression

plays a larger role in modulating the epigenomic signature of the ILC2 gene program than

exposure to disease. To confirm that the patterns we observed represent a specific effect of RAG

expression on the ILC2 gene program, we performed the same analysis on GPL peaks for all

genes in the dataset (**Fig. S11**). In contrast to the ILC2 gene set, the majority of GPL peaks for

all genes was shared among all cell populations, consistent with epigenomic regulation of most

genes being minimally affected by either RAG expression or AD-like disease. Last, in the ILC2

gene set analysis, we noted a set of poised peaks shared by all RAG[naïve] cells and RAG[exp] cells

in the setting of AD-like disease, but not with RAG[exp] cells at steady state (blue bar, **Fig. 6B**). We

reasoned that this condition might capture some genomic loci that are suppressed by a history of

RAG expression at steady state but are induced during inflammation.

Thus, we next quantified and sorted these GPLs to generate a list of genes with the most

peaks "induced" during AD-like disease (**Fig. 6C, Table S14**). Among the identified genes, we

selected *Rora* (**Fig. 6C**) and *Ccr6* (**Fig. 6D**) to examine more closely for evidence of epigenomic

activation in AD-like disease, given the role of these genes in ILC2 expansion[61,62] and homing to

sites of inflammation[44,104], respectively. For both genes, we observed more widespread open

chromatin over the genomic region in the RAG[naïve] cells compared to the RAG[exp] cells, but this

difference was partially abolished by increased open chromatin in AD-like disease in the RAG^{exp}

cells. Taken together, our analysis reveals that a history of RAG expression selectively modulates

activity of ILC2 gene programs across both steady state and during AD-like inflammation, while

some programs are more evident at steady state given the uniquely poised nature of ILC2s.

RAG suppresses the Th2 locus

Our functional data demonstrate a role for RAG expression in regulating ILC2

development and activation, including limiting proportions of IL5^{+} and IL-13^{+} ILC2s at steady state

and in AD-like disease. Prior work identified epigenomic priming in ILC2s early in development at

the Th2 locus (comprised of the *Il4*, *Il13*, *Rad50*, and *Il5* gene loci) to enable rapid transcriptional

responses during inflammation[30]. Thus, we hypothesized that RAG promotes the functional

observations in ILC2s by suppressing the establishment of an active regulome at the Th2 locus.

To test this hypothesis, we analyzed the Th2 locus in our multiomic data in greater detail. Using

a similar strategy to our analysis of ILC2 marker genes, we calculated the number of GPLs in the

RAG^{exp} and RAG^{naïve} cells, respectively, for the genes in the Th2 locus. (**Fig. 7A, Table S7**). We

found many GPLs associated with the four Th2 locus genes, including significant crosstalk

between these genes, similar to previous observations (**Fig. 7A**)[105–107]. Importantly, we identified

fewer GPLs in the RAG^{exp} cells, particularly for the *Il5* and *Il13* loci (**Fig. 7A**). As in our analysis

of the ILC2 marker GPLs, we quantified the differences based on RAG fate mapping and found

that all genes in this locus had increased GPLs in RAG^{naïve} cells relative to RAG^{exp} cells (**Fig. 6B;**

Table S15). We also applied the same analysis strategy that identified TFs potentially mediating

observed differences between RAG^{exp} and RAG^{naïve} ILC2s (**Fig. S10D,E**) specifically to the four

Th2 locus genes. Given the limited size of genomic regions (and thus open chromatin peaks)

analyzed in the Th2 locus compared to all ILC2 genes, we found overall fewer enriched motifs.

Strikingly, significant enrichment of TF motifs was only present in unique peaks from RAG^{naïve}

ILC2s, while no TF motifs met the cutoff in RAG^{exp} cells (**Fig S12A, Table S16**). These motifs

primarily contained the canonical 5'-(A/T)GATA(A/G)-3' binding site recognized by the GATA

family of zinc finger TFs[108] . When we compared enriched motifs in open chromatin to gene

expression of the corresponding TFs, only *Gata3* was expressed at appreciable levels (**Fig**

S12B). Critically, *Gata3* expression was higher in RAG[naïve] compared to RAG[exp] cells, consistent

with our previous analyses of the ILC2 gene regulomes (**Fig. 5C**). Collectively, our data confirm

the established role of GATA3 in mediating activation of the Th2 locus[50] and are consistent with

a role for RAG expression in suppressing the type 2 regulome at the Th2 locus.

379   We next considered the additional effects of AD-like inflammation on the Th2 epigenomic

regulome using the same approach we used to analyze the ILC2 gene set in **Fig. 6**. Again, we

found the largest set of peaks was shared by the RAG[naïve] cells, regardless of disease state, with

the next largest peak sets belonging to either steady state or AD-like disease in the RAG[naïve] cells

(**Fig. 7C**). Furthermore, there was a large proportion of peaks shared by both RAG[exp] cells,

consistent with a major contribution of a history of RAG expression to epigenomic modulation of

the Th2 regulome. To quantify the potential effect of AD-like inflammation on reversing RAG-

mediated suppression of Th2 locus genes, we mapped the 14 peaks shared by RAG[naive] cells and

RAG[exp] cells in the setting of AD-like inflammation (i.e. only suppressed in RAG[naïve] cells at steady

state) back to the Th2 genes via their respective GPLs (**Fig. 7C - blue bar, Table S17**).

Interestingly, *Il13*, which was not identified as a top ILC2 marker in our earlier analyses, had the

highest number of linked peaks associated with potential induction in AD-like disease (**Fig. 7D**).

When we examined the *Il13* locus in the ILC2 cluster more closely, we found more widespread

open chromatin in the RAG[naïve] cells compared to the RAG[exp] cells (**Fig. 7E**). However, in the AD-

like disease sample, the RAG[exp] cells displayed increased open chromatin relative to the steady

state, consistent with induction in the setting of inflammation, like our earlier findings for ILC2

genes such as *Ccr6* (**Fig. 6E**). Taken together, our functional data and multiomic analyses

demonstrate a role for RAG expression in modulating genes critical for ILC2 development and

function, including the key type 2 cytokines expressed from the Th2 locus.

DISCUSSION

400   RAG recombinases evolved nearly 500 million years ago from endogenous transposons,

crucially enabling antigen receptor rearrangement and emergence of the adaptive immune cell

lineages present in all modern vertebrates[29,109]. Indeed, RAG deficiency leads to a complete lack

of B and T lymphocytes, manifesting clinically as severe combined immunodeficiency (SCID)[110–

112]. However, fate mapping studies have shown that multiple mature immune cell populations

other than adaptive B and T lymphocytes have a history of RAG expression[23,24,33,113,114]. More

recent studies by Karo et al found that RAG expression during NK cell development influences

multiple cellular functions including antitumor cytotoxicity, cell proliferation, and survival[24]. Yet

whether RAG modulates cellular functions of innate immune cell populations other than NK cells

remains poorly understood. Here, using RAG-deficient mice, RAG fate mapping mice, and

multiomic analyses, we report that RAG suppresses developmental and effector functions of

ILC2s.

412   Our functional data in RAG-deficient mice demonstrate that populations of ILC2s

producing the type 2 cytokines IL-5 and IL-13 preferentially expand in the absence of a history of

RAG expression. This implies a specific role for RAG in developmental repression of ILC2s.

Building on this, our multiomic RAG fate mapping analyses of ILC2 gene programs demonstrate

extensive epigenomic differences between RAG[exp] and RAG[naïve] cells. We found RAG-associated

epigenomic suppression at multiple functional levels, including cell surface receptors, key

transcription factors, and the Th2 locus encoding the type 2 cytokine genes *Il5* and *Il13*. Although

RAG is only transiently expressed early in lymphoid development[115], our data demonstrate that

RAG expression can imprint durable effects on ILC2 gene programs to restrain their function.

Our observations imply that RAG expression may mark a developmentally distinct

population of ILC2s. In adaptive lymphocytes, RAG expression in T cells is restricted to their time

in the thymus. However, ILC2 populations have been observed in the thymus, provoking the

hypothesis that thymic ILC2s may be uniquely high in expression of RAG[63,116]. Prior studies by

Schneider et al have identified ILC2 populations in adult tissues that variably derive from

expansion of fetal, postnatal, and adult populations[68]. Yet how RAG expression in ILC2s may be

restricted spatially or temporally remains unknown. The mouse strains used in the fate mapping

studies by Schneider et al would be incompatible with our RAG fate mapping mice. Thus, novel

mouse strains enabling intersectional genetics to trace ILC2 ontogeny (e.g. CreER/lox for

temporally restricted fate mapping and FlpO/frt for RAG fate mapping[117]) are needed to more

precisely determine when and where RAG expression occurs during ILC2 development. Beyond

steady state ontogeny, our data suggest a history of RAG expression also imprints suppressed

proliferative and type 2 inflammatory functions on ILC2s in the setting of AD-like disease.

It is increasingly recognized that expression of effector molecules for both ILCs and their

counterpart adaptive lymphocytes (e.g. IL-13 from ILC2s and Th2 cells) is governed by finely

tuned transcriptomic and epigenomic regulomes[30,118–120]. ILCs tend to adopt these regulomes

earlier in their development than T cells, and these "poised" regulatory elements are thought to

underlie the ability of tissue-resident ILCs to rapidly respond to stimuli. In contrast, the regulomes

of naïve T cells remain relatively inactive until stimulation. Given that T cells are uniformly RAG-

experienced, our data provoke the hypothesis that RAG^{exp} ILC2s adopt a phenotype closer to that

of naive T cells and may require stronger stimuli than RAG^{naïve} ILC2s to become activated. Indeed,

our analyses found the RAG-associated suppressive programs could be overcome in the setting

of AD-like inflammation. Thus, sufficient RAG expression may mediate key events underlying the

establishment and maintenance of functional regulomes not only in ILCs, but also T cells. How

RAG might affect these changes, and whether they are independent of its enzymatic activity

and/or antigen receptor recombination, remains to be elucidated.

Clinically, a link between enhanced type 2 immune activity and RAG dysfunction is well-

established. Omenn Syndrome (OS) is a form of SCID characterized by exaggerated type 2

immune activation and typically arises in the setting of hypomorphic RAG gene mutations.

Impaired antigen receptor rearrangement, with rare "leaky" recombination events, leads to

expansion of autoreactive oligoclonal T cells, eosinophilia, and markedly elevated IgE[121–125].

Similar phenotypes have been observed in mice harboring RAG mutations analogous to those

found in human patients with OS[126,127]. Notwithstanding these findings, the mechanisms

underlying the propensity of oligoclonal T cells with hypomorphic RAG activity to preferentially

develop into the Th2 subtype are unclear. Prior studies have found a role for regulatory T cells in

controlling type 2 skewing of transferred T cells in RAG-deficient hosts, potentially explaining

similar observations in patients with OS[128]. Our data provide an additional mechanism by which

RAG dysfunction may lead to OS through loss of cell-intrinsic RAG-mediated suppression of type

2 cellular programs. Additionally, increased type 2 cytokine production from RAG-deficient ILC2s

may, in *trans*, enhance expansion of the oligoclonal Th2 cell populations, IgE induction, and

eosinophilia observed in RAG-deficient states like OS. However, whether other immune cell types

with RAG dysfunction, such as ILCs, contribute to the pathogenesis of OS in humans has not

been investigated.

Lymphoid acquisition of RAG activity may represent a newer evolutionary mechanism that

fine tunes ancient innate immune cell programs in addition to enabling development of relatively

newer antigen-specific adaptive immune cell populations. Independent of antigen receptor

diversity, loss of this function may offer an explanation as to why oligoclonal T cells tend to expand

and skew towards a Th2 cell phenotype in the setting of hypomorphic RAG function as in OS[128].

Further studies are needed to define whether the suppressive effects of RAG expression operate

similarly in T and B cells. Although we demonstrate that this phenomenon is observed in ILC2s,

whether hypomorphic RAG expression in bona fide Th2 cells not only results in oligoclonality but

also loss of suppression of the Th2 locus independently of antigen receptor rearrangement

remains an outstanding question. Indeed, during development of gene therapy strategies for

RAG-deficient SCID, lower doses of wild type RAG transgene expression have been associated

with development of OS-like conditions in transplanted RAG-deficient mice[129–132].

A major limitation of our study is a focus on cutaneous type 2 inflammation, which

stemmed from our initial observations in the MC903 mouse model of AD-like disease. Further,

given the scarcity of skin-resident ILC2 populations, key functional investigations in our study

such as cytokine production and multiomic sequencing were limited to the sdLN, as in prior

studies[13,133]. However, ILC2s are recognized to have highly tissue-specific functions that extend

much beyond inflammation to other processes including regeneration and metabolism. In addition

to Il-5 and IL-13, ILC2s can produce other effector molecules such as acetylcholine, IL-9,

methionine-enkephalin peptides, and amphiregulin, which modulate tissue responses across

numerous organs[134–141]. Considering that the complexity of ILC2 biology may result in markedly

divergent responses to RAG expression in other tissues and disease models, we thus restricted

our initial studies to the skin, where we had strong molecular, cellular, and phenotypic outcomes.

An implication of our findings in the skin is that RAG expression may modulate a variety of ILC2

functions in other tissues. Broader surveys of how RAG impacts ILC2 development and function

in different tissues and disease states remains an exciting area of inquiry.

While we focused our multiomic analyses on ILC2s, it is likely that RAG may impact other

ILC populations. For example, hyperactivation of intestinal ILC3s has been observed in $Rag1^{-/-}$

mice secondary to persistent phosphorylation of Signal Transducer and Activator of Transcription

3 (STAT3). Adoptive transfer of T regulatory cells rescued this phenotype, providing a cell-

extrinsic mechanism for the observation of hyperactivated ILC3s in the setting of RAG

deficiency[142]. However, our data supporting a cell-intrinsic role for RAG in ILC2s may offer

additional mechanistic insight into the prior observations in ILC3s. We found that the regulome of

Jak2, which encodes JAK2, an upstream activator of STAT3, was more activated in RAG[naïve]

ILC2s (**Fig. 5C**). Additionally, the regulome for *Dusp1*, which encodes dual specificity

phosphatase 1 (DUSP1), was more activated in RAG[exp] ILC2s (**Fig. 5C**). While not implicated in

directly dephosphorylating STAT3, a recent study found DUSP1 overexpression negatively

regulated the JAK2/STAT3 pathway[143]. Notably, recent transcriptional profiling of skin ILCs

identified a potential mechanism for skin ILC populations to transition to an ILC3-like phenotype[69],

but how this process is regulated remains poorly understood. Taken together, our data provoke

compelling new hypotheses about cell-intrinsic functions of RAG that may be complementary,

rather than contradictory, to prior observations in gut and skin ILC populations. Additionally, our

studies provide a rationale to design novel reagents to enable more comprehensive studies on

the role of RAG in multiple innate immune cell populations across different tissues and disease

models.

Our observations are also limited by lack of a defined mechanism for how RAG expression

imprints durable epigenomic and transcriptomic changes in ILC2s. The mechanisms by which

RAG mediates VDJ recombination are well-defined, from the biochemical details of DNA-binding

to the epigenomic accessibility of antigen receptor loci and timing of RAG expression[29,144–146].

Notwithstanding genomic stress[24] or potential RAG dose effects[129–132], how RAG expression

might modulate broad developmental and functional lymphoid programs other than V(D)J

recombination remains unclear. The RAG complex can bind both DNA and modified histones and

has been observed to occupy thousands of sites across the genome[27]. Thus, RAG may directly

influence open chromatin states or obscure transcription factor binding sites to alter ILC2

development and function. Notably, RAG preferentially binds near transcription start sites of open

chromatin in mouse thymocytes and pre-B cells, although corresponding effects on gene

expression were not observed[27]. Although canonical recombination sites are concentrated in the

antigen receptor loci, cryptic recombination sites in other regions may be deleted or rearranged

by RAG activity, altering transcriptional regulation of associated genes[27]. In contrast to developing

B and T lymphocytes, the precise timing and location of RAG expression in ILC2s is not known.

Thus, combined with the relative scarcity of ILC2s, conventional methods of chromatin

immunoprecipitation to identify potential epigenomic regulatory mechanisms mediated by RAG

expression may not be feasible in ILC2s or other rare cell populations. Instead, newer

technologies such as self-reporting transposons[147] could be adapted to trace the genomic

footprint of RAG in cells at various stages of development and in various tissues independent of

the constraint of concurrent RAG expression. Finally, through its E3 ubiquitin ligase activity[29],

RAG may influence immune signaling pathways independently of transcription altogether. Given

that direct targeting of RAG would lead to unacceptable side effects, elucidating the mechanisms

by which RAG imprints phenotypic changes beyond antigen receptor rearrangement is a critical

next step in translating these findings to potential new therapies.

Our studies expand prior work implicating RAG in critical immune functions beyond

antigen receptor rearrangement that is exclusive to adaptive lymphocytes. Further, we provide

additional insights into why patients with OS exhibit atopic syndromes in the setting of adaptive

lymphocyte deficiency. Future studies into mechanisms underlying these findings may lead to

new therapeutic avenues for disorders such as atopic dermatitis, food allergy, and asthma.

ACKNOWLEDGEMENTS

We thank all members of the Kim lab for helpful comments and discussion. This work is

supported by the Allen Discovery Center program, a Paul G. Allen Frontiers Group advised

program of the Paul G. Allen Family Foundation, the Doris Duke Charitable Foundation, LEO

Pharma, the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS)

(AR070116, AR077007, and AR080392), and the National Institute of Allergy and Infectious

Disease (NIAID) (AI167933 and AI167047) of the National Institutes of Health (NIH). A.M.V. is

supported by NIAMS (1K08AR080219). M.T. is supported by the Japanese Society of Allergology

(JSA) International Scholarship. A.M.T. is supported by the NIAID (AI007163 and AI154912). We

thank the Genome Technology Access Center at the McDonnell Genome Institute at Washington

University School of Medicine for help with genomic analysis. The Center is partially supported

by NCI Cancer Center Support Grant #P30 CA91842 to the Siteman Cancer Center and by

ICTS/CTSA Grant# UL1TR002345 from the National Center for Research Resources (NCRR), a

component of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research.

This publication is solely the responsibility of the authors and does not necessarily represent the

official view of NCRR or NIH.

AUTHOR CONTRIBUTIONS

Conceptualization: A.M.V., M.M., L.Z., M.T., T-L.Y., and B.S.K.

Methodology: A.M.V., M.M., M.T., L.Z., T-L.Y., and B.S.K.

Validation: A.M.V., M.M., L.Z., M.T., and T-L.Y.

Formal Analysis: A.M.V. and M.M.

Investigation: A.M.V., M.M., L.Z., M.T., T-L.Y., D-H.K., and H.J-M.

Resources: S.V.D. and B.S.K.

Writing – Original Draft: A.M.V.

Writing – Review & Editing: All authors

Supervision, B.S.K.

Funding Acquisition, B.S.K.

DECLARATION OF INTERESTS

FIGURE TITLES AND LEGENDS

Figure 1. RAG deficiency leads to expansion and activation of ILC2s during inflammation

and at steady state.

A) Experimental schematic of AD-like disease. WT (Control) B6 mice or *Rag1*$^{-/-}$ mice treated

topically to the inner surface of each ear with 2 nmol MC903 in 10 μL ethanol vehicle daily for 7

days develop AD-like inflammation. **B)** Ear thickness measured daily in AD-like inflammation.

Data representative of at least 2 independent experiments, 5 mice/group. ** P < 0.01 by 2-way

ANOVA with Sidak's multiple comparisons test, day 7. **C)** Total number ILC2s normalized to 10^5

live cells and **D**) ILC2 proportion of CD90$^+$, Lin$^-$ cells (Lin$^-$ defined as CD3$^-$, CD5$^-$, CD11b$^-$, CD11c$^-$

, CD19$^-$, NK1.1$^-$, and FcεR1$^-$) determined to be ILC2s (IL-33R$^+$) in skin-draining lymph nodes

(sdLN) from WT or *Rag1*$^{-/-}$ mice with AD-like ear inflammation. Percent ILC2 from sdLN in AD-like

disease following PMA/iono stimulation positive for **E)** IL-5 or **F)** IL-13 staining. **G)** Schematic of

steady state analysis of sdLN from WT (Control) or *Rag1*$^{-/-}$ mice. **H)** Total number ILC2s

normalized to 10^5 live cells and **I**) ILC2 proportion of steady state sdLN CD90$^+$, Lin$^-$ cells

determined to be ILC2s as in **(C,D)**. Percent ILC2 from sdLN in steady state following PMA/iono

stimulation positive for **J)** IL-5 or **K)** IL-13 staining. **C-F; H-K)** Data representative of at least 2

independent experiments, 4-5 mice/group * P < 0.05, ** P < 0.01, *** P < 0.001 by two-tailed

Welch's t test. All data represented as mean with standard deviation.

Figure S1. ILC2 and IL-5/IL-13 gating.

Gating for **A)** CD45$^+$, CD90$^+$, Lin$^-$ cells (Lin$^-$ defined as CD3$^-$, CD5$^-$, CD11b$^-$, CD11c$^-$, CD19$^-$,

NK1.1$^-$, and FcεR1$^-$), then gating on **B)** ILC2 (IL-33R$^+$ Lin$^-$) corresponding to **Fig. 1C**, with

subsequent gating of **C)** IL-5$^+$ and IL-13$^+$ ILC2, corresponding to **Fig. 1D-E.**

Figure S2. Expansion and activation of ILC2s in RAG2 deficiency compared to littermates.

A) Schematic of steady state analysis of WT B6 (Control) mice or *Rag2$^{-/-}$* mice. **B)** Proportion of CD90$^+$ Lin$^-$ cells (Lin- defined as CD3$^-$, CD5$^-$, CD11b$^-$, CD11c$^-$, CD19$^-$, NK1.1$^-$, and FcεR1$^-$) determined to be ILC2s (IL-33R$^+$) in sdLN at steady state from WT or *Rag2$^{-/-}$* mice. Percent ILC2 from sdLN at steady state following PMA/iono stimulation positive for **C)** IL-5 or **D)** IL-13 staining. Data representative of 2 independent experiments, 2-3 mice per group. * P <0.05, ** P < 0.01 by two-tailed Welch's t test. All data represented as mean with standard deviation.

Figure 2. Homeostatic expansion and activation of RAG-deficient ILC2s is cell intrinsic

A) Experimental schematic of AD-like disease in splenocyte chimera experiment. WT B6 or *Rag1$^{-/-}$* mice received WT splenocytes and developed AD-like inflammation after subsequent topical treatment with 2 nmol MC903 in 10 μL ethanol vehicle to each ear daily for 10 days. **B)** Ear thickness measured daily in AD-like inflammation. Data representative of 2 independent experiments, 4-5 mice per group. **** P <0.0001 by 2-way ANOVA with Sidak's multiple comparisons test, day 10. **C)** Total number ILC2s normalized to 10^5 live cells and **D)** ILC2 proportion of CD90$^+$, Lin$^-$ cells (Lin$^-$ defined as CD3$^-$, CD5$^-$, CD11b$^-$, CD11c$^-$, CD19$^-$, NK1.1$^-$, and FcεR1$^-$) determined to be ILC2s (IL-33R$^+$). Percent ILC2 from sdLN in splenocyte chimera mice with AD-like disease after PMA/iono stimulation positive for **E)** IL-5 or **F)** IL-13 staining. **G)** Schematic of bone marrow chimera experiment. Equal quantities of bone marrow cells from *Rag1$^{-/-}$* (CD45.2, CD90.2 - orange) and WT (CD45.2, CD90.1 - blue) C57Bl/6J donor mice were used to reconstitute the immune systems of irradiated recipient WT (CD45.1 - black) C57Bl/6J mice. **H)** Proportion of donor (CD45.2$^+$) ILC2 defined as in (C) in sdLN by donor source (CD90.1$^+$ - WT, CD90.2$^-$ - *Rag1$^{-/-}$*). Proportion of Lin$^-$ ILCs by donor source positive for **I)** IL-5 and **J)** IL-13 following PMA/iono stimulation and cytokine staining. **C-F)** Data representative of at least 2 independent experiments, 4-5 mice per group. ** P < 0.01, **** P < 0.0001 by two-tailed Welch's t test. **H-J)**

Data representative of at least 2 independent experiments, 4-5 mice per group. ** P < 0.01 by

two-tailed ratio means paired t test. All data represented as mean with standard deviation.

Figure S3. Confirmation of splenocyte reconstitution in splenocyte chimera mice.

Proportion of CD45$^+$ splenocytes from splenocyte chimera mice (related to **Fig. 2A-E**) determined

to be **A)** CD4$^+$ T cells (CD4$^+$, CD8$^-$, CD19$^-$), **B)** CD8$^+$ T cells (CD4$^-$, CD8$^+$, CD19$^-$), **C)** B cells (CD4$^-$

, CD8$^-$, CD19$^+$), and **D)** Eosinophils (SiglecF$^+$, CD4$^-$, CD8$^-$).

Figure S4. Donor cell reconstitution and gating in sdLN of WT:*Rag1$^{-/-}$* bone marrow chimera

mice.

A) Gating of live host (CD45.1$^+$) and donor (CD45.2$^+$) cells then **B)** Gating on donor cells by

genotype (CD90.1$^+$ = WT [blue]; CD90.2$^+$ = Rag1$^{-/-}$ [orange]) in Lin$^-$ population then **C**) Gating on

Lin$^-$ IL-33R$^+$ ILC2s in the skin draining lymph node (sdLN). **D)** Host/donor CD45$^+$ cell reconstitution

in sdLN of WT:*Rag1$^{-/-}$* bone marrow chimera mice. **E)** quantification of CD45.2$^+$ Lin$^-$ donor cells

by genotype in sdLN of WT:*Rag1$^{-/-}$* bone marrow chimera mice. **F**) total numbers of ILC2s

normalized to 10^5 live cells and **G**) ILC2 proportion of Lin- cells in the sdLN. **H**) Gating for IL-5 and

IL-13 after in vitro stimulation and intracellular cytokine staining of ILC2s from sdLN. Quantification

of total positive cells normalized to 10^5 live cells for **I**) IL-5 and **J**) IL-13 and proportion of ILC2

positive for **K**) IL-5 and **L**) IL-13. **E,G,I-L:** * P < 0.05, ** P <0.01 by ratio means paired t test. **D,F:**

** P <0.01, *** P < 0.001, **** P < 0.0001, by RM one-way ANOVA test with Geisser-Greenhouse

correction. All data represented as mean with standard deviation. Related to **Fig. 2G-J.**

Figure 3. A history of RAG expression marks a population of ILC2s in the sdLN.

A) Schematic of RAG fate mapping in the lymphoid cell compartment using reporter mice

expressing Cre-inducible tandem dimer red fluorescent protein (tdRFP) from the *Rosa26* locus

crossed to mice expressing Cre recombinase from the *Rag1* locus. **B-F)** Histograms of tdRFP

signal in CD45$^+$ sdLN cells by cell type for **B)** CD4$^+$ T cells (B220$^-$, CD3$^+$, CD4$^+$), **C)** CD8$^+$ T cells

(B220$^-$, CD3$^+$, CD8$^+$), **D)** B cells (MHCII$^+$, B220$^+$), **E)** NK cells (B220$^-$, CD3$^-$, CD4$^-$, CD8$^-$, CD49b$^+$,

NK1.1$^+$), **F)** ILC2s (B220$^-$, CD3$^-$, CD4$^-$, CD8$^-$, CD49b$^-$, NK1.1$^-$, CD11b$^-$, CD11c$^-$, SiglecF$^-$, CD90$^+$,

KLRG1$^+$ or ICOS$^+$ or IL-33R$^+$), **F)** quantification of tdRFP$^+$ proportion of each cell type.

Figure S5. sdLN multiome experiment.

A) *Rag1*^{Cre}::*Rosa26*$^{LSL-tdRFP}$ reporter mice were given topical treatments with 2 nmol MC903

dissolved in ethanol vehicle or with ethanol vehicle alone to each ear daily for 7 days. Harvested

sdLN processed using Magnetic Activated Cell Sorting (MACS) led to depletion of cells

expressing the CD3, CD19, and CD11b lineage markers and remaining cells were further

processed in the 10X Multiome pipeline, generating both single cell RNA-sequencing and single

cell ATAC-sequencing data for each cell. **B)** Ear thickness measured daily in the AD-like disease

multiome experiment. Data representative of one experiment, with 4 mice per group pooled for

sequencing. **** P <0.0001 by 2-way ANOVA with Sidak's multiple comparisons test, day 7. All

data represented as mean with standard deviation.

Figure 4. Multiomic analysis of ILC2s through single nuclei sequencing of the sdLN.

A) Schematic of the gene expression (GEX) assay derived from snRNA-seq data. **B)** Schematic

of the gene activity (GA) assay, representing estimated transcription scores derived from

snATAC-seq data using Signac. **C)** UMAP visualizations of independent analyses of RNA-seq

and ATAC-seq data for 2034 sdLN cells after dimensional reduction and clustering combined

using weighted nearest neighbor (WNN) analysis in Seurat. Cluster identities are color coded

consistently throughout the following panels. Heatmaps of **D)** top 25 GEX marker genes and **E)**

top 25 GA marker genes identified for each cluster. See **Table S1** for full lists of genes. **F)** Dotplots

comparing selected marker genes for each cluster between the GEX and GA assays, with

emphasis on known cell type-specific markers. **G)** Schematic of differentially accessible (DA)

chromatin assay, which finds the nearest gene to any peak calculated to be differentially open in

a particular cell cluster. See **Table S2** for full lists of top 25 DA cluster markers. **H)** Overlap of top

100 markers for the ILC2 cluster from the GEX, GA, and DA assays. See **Table S3** for top 100

DA peaks and distances to nearest genes and **Table S4** for full list of top 100 ILC2 markers. **I)**

Selected genes from the ILC2 gene set for each assay individually and for overlaps.

Figure S6. ILC2 marker genes identified in the differentially accessible open chromatin

assay.

Differentially accessible (DA) open chromatin peaks identified for the ILC2 cluster are highlighted

in gray and shown next to the closest gene for **A)** Neuromedin U receptor 1 (*Nmur1*) and **B)** IL-5

(*Il5*). See **Table S3** for top 100 DA peaks and distances to nearest genes for the ILC2 cluster.

Figure S7. Dotplots of selected ILC2 marker genes.

Dotplots comparing selected marker genes from the multiomic ILC2 gene set (**Figure 4I, Table**

S4) for each cluster between the GEX and GA assays, with genes highlighted by color

corresponding to individual assays or overlap of assays in which they were identified. *Rora* was

not detected in the GA assay.

Figure 5. A history of RAG expression imprints transcriptomic and epigenomic modulation

of ILC2 gene programs.

A) Schematic of transcriptional RAG fate mapping. Sequenced cells from the RAG fate map

mouse (see Figure 3A) transcribe tdRFP only after Cre is expressed from the *Rag1* locus. Cells

were assigned as either having a history of RAG expression (RAG^{exp} - tomato red) or not (RAG^{naïve}

- dark gray) based on detection of tdRFP transcript in the RNA-seq data. **B)** Schematic of mapping

gene to peak links (GPLs). The LinkPeaks function of Signac (see methods) calculates significant

correlations between open chromatin at defined peaks (teal bars) and nearby gene expression.

These links represent inferred epigenomic-transcriptomic regulation, or "regulomes" based on the

correlated snRNA- and snATAC-sequencing data. After calculating GPLs separately for each

population (gray for RAG^{naïve} and tomato red for RAG^{exp}), GPLs found in only one group, but not

the other, can be identified (teal boxes). The difference in GPLs based on RAG experience for

any given gene (e.g. Gene X) can be visualized on a bar graph, with the number of GPLs for

RAG^{naïve} (gray - left) and RAG^{exp} (red - right) plotted, with the difference overlaid as a black bar.

C) GPLs calculated as in (B) for the multiomic ILC2 gene set identified in **Figure 4H** and **Table**

S4). All identified GPLs are listed in **Table S7**, while ILC2 GPLs are listed in **Table S8**. Genes are

sorted from more links identified in the RAG^{naïve} population at top to more links identified in the

RAG^{exp} population at bottom. Select genes are labeled. Full ranked list by difference in GPLs is

available in **Table S9**.

Figure S8. Gene set enrichment analysis of differentially expressed genes in ILC2s.

A) Volcano plot of differentially expressed genes (DEGs) by RAG fate map for the ILC2 cluster.

A ranked list (**Table S5**) was constructed for all DEGs with log$_2$(fold change) >0.1 for gene set

enrichment analysis (GSEA). **B)** Dotplot of GSEA result calculated using ClusterProfiler and the

gene ontology (GO) biological process (BP) database (see methods). Full results in **Table S6**. **C)**

GSEA plot of the GO BP "positive regulation of immune system process" gene set. RAG^{naïve}, RAG

fate map negative; RAG^{exp}, RAG fate map positive.

Figure S9. Mapping gene to peak links in select ILC2 genes.

Gene to peak links (GPLs) mapped for the RAG^{naïve} and RAG^{exp} states as depicted in **Figure 5B**

for **A)** GATA binding protein 3 (*Gata3*) and **B)** Nedd4 family interacting protein 1 (*Ndfip1*). Only

GPLs that fit in the coverage window are shown. Select peaks (teal bars) present in one state,

but not the other, are highlighted in teal boxes. Full gene names not shown on figure in (A) are

*9230102O04Rik and **4930412O13Rik and in (B) #Gm42690.

Figure S10. Multiomic transcription factor analysis of ILC2s

 A) Schematic of assigning transcription factor (TF) motif enrichment in differentially open chromatin to cell clusters using chromVAR. **B**) Heatmap of top 5 TF motif activity scores for each cluster. **C**) Dotplots comparing expression levels of selected TFs in the gene expression (GEX) assay with the chromVAR activity score of the corresponding TF motif. The TFs and corresponding motifs for *Rora* and *Rorg* in the ILC2 cluster are highlighted by boxes. An expanded list of cluster TF motif markers identified using chromVAR is in **Table S10**. **D**) Analysis of TF motifs enriched in ILC2 gene to peak links (GPLs) unique to RAG^{naïve} and RAG^{exp} populations determined using the FindMotifs function in Signac. The top 6 TF motifs for each population are shown and are ranked by the -log$_{10}$ transformed false discovery rate (FDR - Bonferroni corrected p values). An expanded list is in **Table S11**. **E**) Dotplot of gene expression for TFs corresponding to the top enriched TF motifs identified in ILC2 GPLs from (D). TF genes that were not detected in the GEX assay are labeled N.D.

Figure 6. A history of RAG expression broadly influences ILC2 genes at steady state and in AD-like inflammation.

A) Schematic of the process to determine contribution of RAG fate map and disease states to GPLs for subsequent intersection analyses. GPLs were first calculated for all indictated cells, regardless of disease state or fate map. Cells were then split, first by RAG fate map (RAG^{exp} and RAG^{naïve}), and again by disease state (SS - steady state, AD - AD-like inflammation). GPLs were recalculated for each split sample and matched back to the original set of total GPLs. **B)** UpSet plot visualizing intersections of peaks identified from ILC2 GPLs for split samples. Each row represents one of the four sets, and each column corresponds to an intersection of one or more sets (see methods). See **Table S12** for full list of GPLs for all genes. **Table S13** lists total and ILC2 peaks used for intersection analyses in each of the four sets. Columns identifying key

intersections are color coded by the corresponding RAG fate map or treatment groups. The blue

column indicates the intersection of peaks from RAG^{naïve} cells and peaks induced by AD-like

disease in RAG^{exp} cells. **C)** Top genes with the most AD-like disease-induced peaks. Peaks from

the intersection between RAG^{naïve} cells and inflamed RAG^{exp} cells were identified in corresponding

GPLs, and genes were ranked by number of linked peaks identified. See **Table S14** for full list of

ranked genes and associated GPLs. Open chromatin in the ILC2 cell cluster split by disease state

and then by RAG fate map for the genomic loci of **C)** *Ccr6* and **D)** *Rora*.

Figure S11. Gene to peak link analysis by RAG fate map and disease for all detected genes.

A) UpSet plot of overlaps in peaks identified from GPLs of all genes split by both RAG fate map

(RAG^{exp} and RAG^{naïve}) and disease state (SS - steady state, AD - AD-like inflammation). Each row

corresponds to one of the four sets, and each column corresponds to an intersection of one or

more sets (see methods). See **Table S13** for full list of peaks from GPLs for all genes in each set.

Columns identifying key intersections are color coded by the corresponding RAG fate map or

disease groups. The blue column indicates the intersection of peaks from RAG^{naïve} cells and

peaks induced by AD-like disease in RAG^{exp} cells.

Figure 7. RAG suppresses the Th2 locus.

A) Coverage plot of the Th2 genomic locus. Open chromatin in the ILC2 cluster for each *Rag1*

fate-mapped state is shown on top, and corresponding peaks (teal) and gene to peaks links

(GPLs) are shown below for the RAG^{naïve} sample (gray) and the RAG^{exp} sample (tomato red).

Only GPLs that fit in the coverage window are shown. **B)** All GPLs identified in each fate map

state for the Th2 locus genes *Il4*, *Il13*, *Rad50*, and *Il5*. See **Table S15** for full list of Th2 GPLs.

The number of GPLs for each gene is shown on the left in gray for RAG^{naïve} and on the right in

tomato red for RAG^{exp}. The difference is superimposed in black, and genes are sorted from more

GPLs identified in RAG^{naïve} at top to more links identified in RAG^{exp} at bottom. **C)** UpSet plot of

intersections of peaks identified from Th2 locus GPLs. GPLs were recalculated, this time for

samples separately by both RAG fate map status (RAG^{exp} and RAG^{naïve}) and disease (SS - steady

state, AD - AD-like inflammation). Each row represents one of the four sets of peaks, and each

column corresponds to an intersection of one or more sets (see methods). See **Table S13** for full

list of peaks from GPLs for all genes, including Th2 genes, in each of the four sets. Columns

identifying key intersections are color coded by the corresponding RAG fate map or disease

groups. The blue column indicates the intersection of peaks from RAG^{naïve} cells and peaks

induced in AD-like disease in RAG^{exp} cells. (**D)** Th2 genes sorted by number of AD-like disease-

induced peaks. Peaks induced by AD-like disease were identified in corresponding GPLs, and

genes were ranked by frequency of links to induced peaks (representation in identified GPLs).

See **Table S17** for full list of ranked Th2 locus genes and associated GPLs. **E)** Open chromatin

tracks, split by disease (beige box – steady state; maroon box – AD-like disease) and by RAG

fate map (RAG^{naïve} - gray, RAG^{exp} - red) for *Il13*.

Figure S12. Multiomic transcription factor analysis of Th2 locus

A) Analysis of TF motifs enriched in Th2 locus subset of ILC2 gene to peak links (GPLs) unique

to RAG^{naïve} and RAG^{exp} cell populations determined using the FindMotifs function in Signac. The

top 6 TF motifs for each population are shown and are ranked by the -log$_{10}$ transformed false

discovery rate (FDR - Bonferroni corrected p values). No TFs met the -log10(FDR) minimum cutoff

value of 1.5 in the RAG^{exp} cell population. The full list of enriched motifs is in **Table S16**. **B**) Dotplot

of gene expression for TFs corresponding to the top enriched TF motifs identified in Th2 locus

GPLs from (A). TF genes that were not detected in the GEX assay are labeled N.D.

MATERIALS AND METHODS

Animal studies

Wild-type (WT) C57BL/6J and WT congenic strains (CD90.1[+], CD45.1[+]), *Rag1*[-/-], and *Rag2*[-/-] mice were initially purchased from the Jackson Laboratory and bred in house. The RAG fate-mapping strain *Rag1*[Cre]::*Rosa26*[LSL-tdRFP] was originally created in the lab of Paul Kincade[33] and bred in house. All mice were housed in specific-pathogen-free condition and environmentally controlled animal faculty with a 12-hour light-dark cycle and given unrestricted access to food and water at Icahn School of Medicine at Mount Sinai or Washington University School of Medicine in St. Louis. All animal protocols and experiments were approved by the Institutional Animal Care and Use Committee (IACUC) at Icahn School of Medicine at Mount Sinai or Washington University School of Medicine in St. Louis. Experiments were performed on independent cohorts of male and female mice. For induction of AD-like disease, 8- to 12-week-old mice were treated with 2 nmol calcipotriol (MC903, Tocris Bioscience) in 10 µL of 100% ethanol (EtOH) vehicle, or vehicle alone, on the bilateral ear skin daily for 7-10 days. Body weight and ear thickness were measured daily with a digital scale and analog caliper by the same investigator. For tissue harvest, animals were euthanized by CO_2 inhalation.

Flow cytometry

Cervical skin draining lymph nodes (sdLN) were removed from the mice and immediately homogenized manually through a 100 µm cell strainer (Fisher Scientific) into a 50 mL tube with the end of a plunger from a 3 mL syringe. The strainer was washed with wash medium (2% vol/vol FBS/PBS) and the strained cells were centrifuged at 400g for 5 minutes at 4°C. Lymph node cell samples were stained with Zombie NIR viability dye (Biolegend; 1:500) to exclude dead cells, followed by Fc-receptor blocking and cell-surface staining with specific antibodies. The cells were analyzed using either LSR Fortessa[TM] (BD) or Cytek® Aurora (CYTEK) flow cytometers. Data was

833   obtained using either FACSDiva^{TM} (BD) or SpectroFlo® (CYTEK) software and was further

834   analyzed using FlowJo^{TM}.

Lymphocyte stimulations

After tissue harvest, ILC stimulations were performed by incubating 0.5-1x10^6 cells for 4

hours at 37°C in stimulation media (DMEM with 5% fetal bovine serum, 1%

penicillin/streptomycin, 2 mM L-glutamine, 50 ng/mL Phorbol 12-myristate 13-acetate (PMA), 100

ng/mL ionomycin, 5 ug/mL Brefeldin A (BFA), 2 uM monensin). T cell stimulations were performed

by first coating a 96-well plate with 5 µg/mL anti-mouse CD3 (Biolegend) in 50 µL/well PBS

overnight the day before tissue harvest. The following day, 0.5-1x10^6 cells were resuspended in

50 µL/well T cell stimulation media (5 µg/mL anti-mouse CD28 (Biolegend), 5 ug/mL BFA, 2 µM

monensin) and incubated for 20 minutes hours at 37 C. The cells were then transferred to the

anti-mouse CD3 coated plate and incubated for 4 hours at 37 C. After stimulation, cells were

washed in wash medium, fixed, and stained for surface and intracellular markers as described for

unstimulated cells.

Splenocyte chimeras

Spleens were harvested from donor WT B6 mice and immediately homogenized manually

through a 100 µm cell strainer (Fisher Scientific) into a 50 mL tube with the end of a plunger from

a 3 mL syringe. The strainer was washed with wash medium (2% vol/vol FBS/PBS) and the

strained cells were centrifuged at 400g for 5 minutes at 4°C followed by treatment with RBC lysis

buffer for two minutes and two wash steps using 2 volumes of wash medium. Cells were counted,

and 5 million splenocytes were injected intraperitoneally into each recipient mouse. Experiments

were performed 4 weeks following splenocyte add-back to allow immune reconstitution.

Bone marrow chimeras

Recipient mice were provided with antibiotic water, consisting of 5 mL of Sulfatrim

(sulfamethoxazole/trimethoprim) added into 200 mL of drinking water, for one week starting from

the day prior to irradiation (day -1). On day 0, recipient mice were irradiated with 950 cGy using

the X-RAD 320 (Precision X-Ray). BM was harvested from donor mice femurs and tibias and

treated with RBC lysis buffer (Sigma-Aldrich) for two minutes. BM cells were transferred into a 15

mL conical tube through a 70 μm cell strainer (Fisher Scientific) and the cell strainer and cells

were washed with 2% (vol/vol) FBS/PBS. The concentration of living cells was determined using

a Cellometer Auto 2000 (Nexcelom Bioscience) with ViaStain^{TM} AOPI Staining Solution

(Nexcelom Bioscience). Recipient mice received the same number of cells, at 1 x 10^7 live bone

marrow cells per mouse, through retroorbital injection within 24-hour after irradiation. Recipients

were given 8 weeks for immune reconstitution after BM transplantation before experimental use.

Cryopreserving sdLN cells for sequencing

Rag1^{Cre}::*Rosa26*$^{LSL-tdRFP}$ mice were treated with 2 nmol calcipotriol (MC903, Tocris

Bioscience) in 10 μL of 100% ethanol (EtOH) vehicle, or vehicle alone, on the bilateral ear skin

daily for 7 days to induce AD-like inflammation. The next day, cervical sdLN were harvested and

immediately homogenized manually through a 100 μm cell strainer (Fisher Scientific) into a 50

mL tube with the end of a plunger from a 3 mL syringe. The strainer was washed with wash

medium (2% vol/vol FBS/PBS) and the strained cells were centrifuged at 400g for 5 minutes at

4°C. Next, cells were incubated with biotinylated antibodies (anti-mouse CD3e, CD19, CD11b;

1:300; Biolegend) in 100 μL of wash buffer for 20 minutes at 4°C, followed by two washes in 2

volumes of wash buffer. Next, no more than 10^7 cells were incubated with Streptavidin

MicroBeads (Miltenyi) in 500 μL separation buffer (0.5% w/v BSA in PBS; BSA and PBS from

Sigma) at 4°C for 20 minutes, then added to LD columns (Miltenyi) pre-equilibrated with

separation buffer and loaded in a QuadroMACS™ Separator (Miltenyi) for negative cell selection.

Remaining cells were eluted in 1 mL separation buffer and cells were centrifuged at 400g for 5

minutes at 4°C, followed by resuspension in freezing buffer (10% DMSO, Invitrogen; 20% FBS in

DMEM, Sigma) and slow freezing to -80°C in a CoolCell™ LX (Corning) device.

Processing cryopreserved cells for multiome

Cryopreserved sdLN cells were processed as recommended by the 10X Genomics

DemonstratedProtocol_NucleiIsolation_ATAC_GEX_Sequencing_RevC_(CG000365)

instructions for primary cells without any modification to the protocol. Briefly, cells were thawed in

a 37°C water bath followed by dilution into media (RPMI + 15% FBS, Sigma) and centrifugation

at 400g for 5 minutes at 4°C. For each final sample (EtOH vehicle- or MC903-treated), cells were

pooled from sample from 3 individual mice. Cells were resuspended in PBS + 0.04% BSA (Sigma)

and passed through a 40 µm Flomi strainer (Bel-art) followed by determination of cell

concentration using the using Cellometer Auto 2000 (Nexcelom Bioscience) with ViaStain™ AOPI

Staining Solution (Nexcelom Bioscience). Cells were centrifuged 5 minutes at 4°C and

supernatant removed. Lysis Buffer (Tris HCl base with 0.1% Tween-20, 0.1% NP-40, 0.01%

digitonin, 1 mM DTT, and 1 U/µL Protectors RNase inhibitor, Sigma; full recipe in 10X protocol)

was added, cells mixed 10x, and incubated on ice for 3 minutes. Nuclei from lysed cells were

centrifuged at 400g for 5 minutes at 4°C and washed in 1 mL Wash Buffer (Lysis Buffer, but

without NP-40 or digitonin). The wash step was repeated two more times. Nuclei concentration

was determined as for cell concentration using the Cellometer and ViaStain™ solution. The AOPI

staining indicated 97-99% lysis efficiency of the cells. We manually confirmed nuclei count using

a Bright-Line™ hemacytometer (Hausser Scientific™). Nuclei were centrifuged at 400g for 5

minutes and resuspended in a volume of 1X Nuclei Buffer (10X Genomics) to yield roughly 4,000

nuclei/µL. We then immediately proceeded to the 10X Chromium Next GEM Single Cell Multiome

ATAC + Gene Expression pipeline.

Multiome library construction and sequencing

Multiome 3v3.1 GEX and ATAC libraries were prepared as recommended by 10X

Genomics protocol Chromium_NextGEM_Multiome_ATAC_GEX_User_Guide_RevD

((CG000338). For sample preparation on the 10x Genomics platform, the Chromium Next GEM

Single Cell Multiome ATAC + Gene Expression Reagent Bundle, 16 rxns PN-

1000283, Chromium Next GEM Chip J Single Cell Kit, 48 rxns PN-1000234, Single Index Kit N

Set A, 96 rxns PN-1000212 (ATAC), Dual Index Kit TT Set A, 96 rxns PN-1000215 (3v3.1 GEX),

were used. The concentration of each library was accurately determined through qPCR utilizing

the KAPA library Quantification Kit according to the manufacturer's protocol (KAPA

Biosystems/Roche) to produce cluster counts appropriate for the Illumina NovaSeq6000

instrument. GEX libraries were pooled and run over 0.05 of a NovaSeq6000 S4 flow cell using

the XP workflow and running a 28x10x10x150 sequencing recipe in accordance with

manufacturer's protocol. Target coverage was 500M reads per sample. ATAC libraries were

pooled and run over 0.167 of a NovaSeq6000 S1 flow cell using the XP workflow and running a

51x8x16x51 sequencing recipe in accordance with manufacturer's protocol. Target coverage

was 250M reads per sample.

Multiomic data analysis

The cellranger-arc-2.0.0 (10X Genomics) pipeline was used to generate FASTQ files,

gene expression matrices, and ATAC fragment tables for each sample, followed by aggregation

using the aggr function. Default settings were utilized, with the exception that we incorporated a

custom reference with the sequence for tdRFP (see **Supplemental file S1**) added to the default

mouse reference sequence provided by cellranger (refdata-cellranger-arc-mm10-2020-A-2.0.0).

Correction for ambient RNA was performed using SoupX[148] with clustering information provided

by the default cellranger outputs. Doublets were removed using Scrublet[149] with default settings.

Corrected data was then processed using Signac[37] and Seurat[34–36]. ATAC-seq peaks were

identified using MACS2[150] through the CallPeaks function in Signac. Per-cell quality control

metrics were computed using the TSSEnrichment and NucleosomeSignal functions, and cells

retained with a nucleosome signal score < 1.5, TSS enrichment score > 1, total RNA counts <

15,000 and > 1,000, total ATAC counts < 75,000 and > 100, percent mitochondrial reads < 5%,

and percent ribosomal genes detected <10%. After these filtering steps, 10,304 cells remained.

Cells were further filtered by their expression of lineage defining markers similar to the negative

selection step during sample processing. Cells with detectable transcripts for *Cd3d*, *Cd3e*, *Cd3g*,

Cd4, *Cd19*, *Cd8a*, and *Itgam* were removed. This left 2,034 remaining cells for further analysis.

The SCTransform function of Seurat was used to normalize RNA counts. We performed

integration of the two samples using the RNA assay to correct for batch effects and treatment in

the initial clustering using the default parameters for the functions SelectIntegrationFeatures,

FindIntegrationAnchors, and IntegrateData. The integrated data was used for PCA (25

dimensions) and UMAP reduction for the RNA assay alone. With default parameters in Signac,

we used TFIDF to normalize ATAC peaks and latent semantic indexing (LSI) to reduce the

dimensionality of the ATAC data. We constructed a UMAP of the ATAC data alone using the LSI

reduction (dimensions 2-25). To construct a joint graph and UMAP using equal weighting from

the RNA and ATAC assays, we used the FindMultiModalNeighbors function of Seurat/Signac

using default parameters (RNA dimensions 1-25, ATAC dimensions 2-25). We used a resolution

of 0.1 to identify clusters with the FindClusters function in Seurat/Signac. Cell types were assigned

based on manual curation of marker genes. Initially, 7 clusters were identified, but two highly

similar lymphocyte clusters were merged for a total of 6 cell types.

The inferred Gene Activity (GA) assay from the ATAC-seq data was calculated using

default parameters of the GeneActivity function in Signac. FindAllMarkers was used to identify

top markers by cluster for both RNA gene expression data (GEX) and GA, with setting

adjustments including min.pct = 0.20 and logfc.threshold = 0.25. The differentially accessible (DA)

open chromatin assay was calculated in Signac with the FindMarkers function on the ATAC-seq

peaks assay (called using MACS2 as above). The differential test used was 'LR' (logistical

regression, as suggested for snRNA-seq[151]). The total number of ATAC fragments was used as

a latent variable to mitigate effect of differential sequencing depth. Given the sparsity of the data,

the min.pct parameter was set to 0.02. After identifying the top differentially accessible peaks for

each cluster, the gene closest to each peak was determined using the ClosestFeature function in

Signac. Results were filtered for genes within 10^5 base pairs of the corresponding peak. The

filtered gene lists were used for the "DA" assay as markers of each cluster (top 25) and an

expanded list for the ILC2 cluster (top 100). Venn diagrams were calculated using

BioVenn/BioVennR[152].

Gene set enrichment analysis was performed and visualized using ClusterProfiler[153]. For

GSEA on steady state ILC2 DEGs between fate mapped states, we opted to use more permissive

filtering parameters instead of default parameters. We created the ranked list of DEGs using the

FindMarkers function in Seurat with min.pct = 0.1 and logfc.threshold = 0.1. The DEG list from

the GEX assay was used to generate the GSEA results. The DEG list from the GA assay did not

yield any significant GSEA results. The ClusterProfiler function gseGO was used to analyze the

ranked DEG list using the paramters minGSSize = 50, maxGSSize = 500, pvalueCutoff = 0.05.

A motif matrix was constructed from the ATAC data Granges using the "CORE" collection

and "vertebrates" taxonomy group from the JASPAR2022 position weight matrix set and the

mm10 reference genome. Per cell transcription factor motif activity was calculated with

chromVar[94] using the motif matrix and MACS2 called peaks. Transcription factor motifs were

identified in differentially accessible chromatin using the FindMotifs function in Signac.

The correlation coefficients, or gene to peak links (GPLs), between gene expression and

accessibility of each peak were calculated for all peaks within 10^6 base pairs of the transcription

start sites for all detected genes using the LinkPeaks function of Signac with min.cells = 2. GPLs

were filtered by gene for the curated ILC2 and Th2 gene sets. Since multiple genes can be linked

to one peak by GPL analysis, finding intersections of GPLs in set analysis would result in counting

some epigenomic regions multiple times. Thus, for set analysis, we eliminated GPLs with

redundant peaks. Then, we used each list of non-redundant peaks as input sets to generate

UpSet plots and lists of intersecting peaks between states (Rag1 fate map positive or negative;

AD-like disease or steady state) using UpSetR[154]. Coverage plots of the single cell multiomic data,

including open chromatin, peaks, and links (GPLs), were plotted using the CoveragePlot function

in Signac.

Data and code availability

Sequencing data have been deposited at GEO and accession numbers are listed in the

key resources table. Aggregated data are supplied in the supplemental file. All data reported in

this paper will be shared by the lead contact upon request. This paper does not report original

code. Any additional information required to reanalyze the data reported in this paper is available

from the lead contact upon request.

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
anti-mouse CD11b	eBioscience	Cat#: 45-0112-82; RRID: AB_953558
anti-mouse CD11c	eBioscience	Cat#: 45-0114-82; RRID: AB_925727
anti-mouse CD19	eBioscience	Cat#: 45-0193-82; RRID: AB_1106999
anti-mouse CD3e	eBioscience	Cat#: 45-0031-82; RRID: AB_1107000
anti-mouse CD5	eBioscience	Cat#: 45-0051-82; RRID: AB_914334
anti-mouse NK1.1	eBioscience	Cat#: 45-5941-82; RRID: AB_914361
anti-mouse FcεR1α	eBioscience	Cat#: 11-5898-82; RRID: AB_465308
anti-mouse IL-13	eBioscience	Cat#: 48-7136-42; RRID: AB_2784729
anti-mouse CD45.1	BioLegend	Cat#: 110706; RRID: AB_313495
anti-mouse CD45.2	BioLegend	Cat#: 109808; RRID: AB_313445
anti-mouse ICOS	BioLegend	Cat#: 313506; RRID: AB_416330
anti-mouse CD62L	BioLegend	Cat#: 104428; RRID: AB_830799
anti-mouse CD69	BioLegend	Cat#: 104530; RRID: AB_2563062
anti-mouse KLRG1	BioLegend	Cat#: 138424; RRID: AB_2564051
anti-mouse IL-5	BioLegend	Cat#: 504306; RRID: AB_315330
anti-mouse IL-13	BioLegend	Cat#: 503826; RRID: AB_2650897
anti-mouse IL-17A	BioLegend	Cat#: 506927; RRID: AB_11126144
anti-mouse IFNγ	BioLegend	Cat#: 505806; RRID: AB_315400
anti-mouse CD90.2	BioLegend	Cat#: 105328; RRID: AB_10613293
anti-mouse IL-33Rα	BioLegend	Cat#: 145308; RRID: AB_2565569
anti-mouse IL-33Rα	BioLegend	Cat#: 145327; RRID: AB_2565569
anti-mouse CD8a	BioLegend	Cat#: 100762; RRID: AB_2564027
anti-mouse CD25	BioLegend	Cat#: 102016; RRID: AB_312865
anti-mouse CD45.2	BioLegend	Cat#: 109806; RRID: AB_313443
anti-mouse CD90.2	BioLegend	Cat#: 109830; RRID: AB_1186098
anti-mouse CD4	BioLegend	Cat#: 100449; RRID: AB_2564587
anti-mouse Gata3	BioLegend	Cat#: 653814; RRID: AB_2563221
anti-mouse CD25	BioLegend	Cat#: 102036; RRID: AB_2563059
anti-mouse CD90.1	BioLegend	Cat#: 202537; RRID: AB_2562644
anti-mouse CD3e	BioLegend	Cat#: 155608; RRID: AB_2750434
anti-mouse CD3e	BioLegend	Cat#: 100339; RRID: AB_11150783
anti-mouse CD28	BioLegend	Cat#: 102115; RRID: AB_11150408
anti-mouse CD3	BioLegend	Cat#: 100243; RRID: AB_2563946
anti-mouse CD19	BioLegend	Cat#: 115503; RRID: AB_313638
anti-mouse CD11b	BioLegend	Cat#: 101203; RRID: AB_312786
anti-mouse IL-4	BioLegend	Cat#: 504109; RRID: AB_493320
anti-mouse CD117 (c-Kit)	BioLegend	Cat#: 105838; RRID: AB_2616739
anti-mouse CD49b	Invitrogen	Cat#: 17-5971-82; RRID: AB_469485
anti-mouse CD45R/B220	BioLegend	Cat#: 103275; RRID: AB_2860602
anti-mouse I-A/I-E (MHCII)	BioLegend	Cat#: 107622; RRID: AB_493727
anti-mouse Ly-6A/E (Sca-1)	BioLegend	Cat#: 122512; RRID: AB_756197
anti-mouse F4/80	BioLegend	Cat#: 123112; RRID: AB_893482

anti-mouse SiglecF	BD Biosciences	Cat#: 562757; RRID: AB_2687994
anti-mouse TCR γ/δ	eBioscience	Cat#: 48-5711-82; RRID: AB_2574071
anti-mouse Ly-6G	eBioscience	Cat#: 62-9668-82; RRID: AB_2762763
anti-mouse CD16/CD32	Bio X Cell	Cat#: BE0307; RRID: AB_1107647
streptavidin	BioLegend	Cat#: 405204
streptavidin	BioLegend	Cat#: 405207
Chemicals, peptides, and recombinant proteins		
Phorbol 12-myristate 13-acetate (PMA)	Sigma	P1585
Ionomycin	Sigma	I0634
Monensin	Biolegend	420701
Brefeldin A Solution	Biolegend	420601
DMEM	Sigma	D6429
RPMI-1640 Medium	Sigma	R8758
Bovine serum albumin (BSA)	Sigma	A2058
Penicillin-Streptomycin	Gibco	15140122
L-glutamine solution - 200 mM	Corning	MT25005CI
Fetal bovine serum	Sigma	2442
Calcipotriol (MC903)	Tocris Biosciences	2700
Phosphate Buffered Saline	Sigma	806552
DMSO	Invitrogen	D12345
Nuclei Buffer (20X)	10x Genomics	2000153/2000207
Digitonin	ThermoFisher	BN2006
Trizma hydrochloride solution pH 7.4	Sigma	T2194
Sodium Chloride Solution 5 M	Sigma	59222C
Magnesium Chloride Solution 1M	Sigma	M1028
Nonidet P40 Substitute	Sigma	74385
Protector RNase inhibitor	Sigma	3335402001
DTT	Sigma	646563
Flowmi Cell Strainer 40 μm	Bel-Art	H13680-0040
Tween 20	Bio-Rad	1662404
ZombieNIR	Biolegend	423106
ZombieUV	Biolegend	423107
Experimental models: Organisms/strains		
B6 WT	Jackson Laboratory	C57BL/6J; Cat# 000664
Rag1[-/-]	Jackson Laboratory	B6.129S7-Rag1tm1Mom/J; Cat# 002216
Rag2[-/-]	Jackson Laboratory	B6.Cg-Rag2tm1.1Cgn/J; Cat# 002014
B6 CD45.1	Jackson Laboratory	B6.SJL-Ptprca Pepcb/BoyJ; Cat# 008450
B6 CD90.1	Jackson Laboratory	B6.PL-Thy1a/CyJ; Cat# 000406
Rag1 fate-map mice	[24,33]	

Software and algorithms		
BD FACSDiva (v8.0)	BD Life Sciences	https://www.bdbiosciences.com/en-us/products/software/instrument-software/bd-facsdiva-software
FlowJo™ (v10.8)	BD Life Sciences	https://www.flowjo.com/
SpectroFlo® (v3.0)	CYTEK	https://cytekbio.com/pages/spectro-flo
Prism 9	GraphPad Software	https://www.graphpad.com/scientific-software/prism/
R (v4.2.2)	R core	https://www.r-project.org/
Seurat (v4.2.0)	[35]	https://github.com/satijalab/seurat
Signac (v1.8.0)	[37]	https://github.com/stuart-lab/signac
chromVAR (v1.18.0)	[94]	https://github.com/GreenleafLab/chromVAR
JASPAR2022 (v0.99.7)		https://github.com/da-bar/JASPAR2022
SoupX (v1.6.1)	[148]	https://github.com/constantAmateur/SoupX
clusterProfiler (v4.4.4)	[153]	https://github.com/YuLab-SMU/clusterProfiler
singleCellTK (v2.6.0)	[155]	https://github.com/compbiomed/singleCellTK
biomaRt (v2.52.0)	[156]	https://github.com/grimbough/biomaRt
EnsDb.Mmusculus.v79 (v2.99.0)	[157]	https://bioconductor.org/packages/release/data/annotation/html/EnsDb.Mmusculus.v79.html
BSgenome.Mmusculus.UCSC.mm10 (v1.4.3)	[158]	https://bioconductor.org/packages/release/data/annotation/html/BSgenome.Mmusculus.UCSC.mm10.html
ggplot2 (v3.3.6)	[159]	https://github.com/tidyverse/ggplot2
viridis (v0.6.2)	[160]	https://github.com/sjmgarnier/viridis
TFBSTools (v1.34.0)	[161]	https://bioconductor.org/packages/release/bioc/html/TFBSTools.html
motifmatchr (v1.18.0)	[162]	https://github.com/GreenleafLab/motifmatchr
BioVenn/BioVennR	[152]	https://www.biovenn.nl/
UpSetR	[154]	http://gehlenborglab.org/research/projects/upsetr/
Cell Ranger	[163,164]	https://support.10xgenomics.com/single-cell-multiome-atac-gex/software/overview/welcome
Python (v3.7)	Python Software Foundation	https://www.python.org/
Deposited data		
scRNA-seq dataset	This paper	GEO: GSE192597

scATAC-seq dataset	This paper	GEO: [GSE192597](GSE192597)
Other		
Streptavidin MicroBeads	Miltenyi	130-048-101
LD Columns	Miltenyi	130-042-901
QuadroMACS™ Separator	Miltenyi	130-090-976
CoolCell™ LX	Corning	CLS432001
Bright-Line™ hemacytometer	Hausser Scientific	3110V
Cytek® Aurora	CYTEK	N/A
LSR FortessaTM	BD Life Sciences	N/A

REFERENCES

1. Ständer, S. Atopic Dermatitis. *New Engl J Med* **384,** 1136–1143 (2021).

2. Hammad, H. & Lambrecht, B. N. The basic immunology of asthma. *Cell* **184,** 1469–1485 (2021).

3. Yu, W., Freeland, D. M. H. & Nadeau, K. C. Food allergy: immune mechanisms, diagnosis and immunotherapy. *Nat Rev Immunol* **16,** 751–765 (2016).

4. Tordesillas, L., Berin, M. C. & Sampson, H. A. Immunology of Food Allergy. *Immunity* **47,** 32–50 (2017).

5. Ring, J. Terminology of Allergic Phenomena. *Chem Immunol Allergy* **100,** 46–52 (2014).

6. Halim, T. Y. F., Hwang, Y. Y., Scanlon, S. T., Zaghouani, H., Garbi, N., Fallon, P. G. & McKenzie, A. N. J. Group 2 innate lymphoid cells license dendritic cells to potentiate memory TH2 cell responses. *Nat Immunol* **17,** 57–64 (2016).

7. Halim, T. Y. F., Steer, C. A., Mathä, L., Gold, M. J., Martinez-Gonzalez, I., McNagny, K. M., McKenzie, A. N. J. & Takei, F. Group 2 Innate Lymphoid Cells Are Critical for the Initiation of Adaptive T Helper 2 Cell-Mediated Allergic Lung Inflammation. *Immunity* **40,** 425–435 (2014).

8. Sokol, C. L., Chu, N.-Q., Yu, S., Nish, S. A., Laufer, T. M. & Medzhitov, R. Basophils function as antigen-presenting cells for an allergen-induced T helper type 2 response. *Nat. Immunol.* **10,** 713–720 (2009).

9. Perrigoue, J. G., Saenz, S. A., Siracusa, M. C., Allenspach, E. J., Taylor, B. C., Giacomin, P. R., Nair, M. G., Du, Y., Zaph, C., Rooijen, N. van, Comeau, M. R., Pearce, E. J., Laufer, T. M. & Artis, D. MHC class II–dependent basophil–CD4+ T cell interactions promote TH2 cytokine–dependent immunity. *Nat Immunol* **10,** 697–705 (2009).

10. Vivier, E., Artis, D., Colonna, M., Diefenbach, A., Santo, J. P. D., Eberl, G., Koyasu, S., Locksley, R. M., McKenzie, A. N. J., Mebius, R. E., Powrie, F. & Spits, H. Innate Lymphoid Cells: 10 Years On. *Cell* **174,** 1054–1066 (2018).

11. Imai, Y., Yasuda, K., Sakaguchi, Y., Haneda, T., Mizutani, H., Yoshimoto, T., Nakanishi, K. & Yamanishi, K. Skin-specific expression of IL-33 activates group 2 innate lymphoid cells and elicits atopic dermatitis-like inflammation in mice. *P Natl Acad Sci Usa* **110,** 13921–6 (2013).

12. Salimi, M., Barlow, J. L., Saunders, S. P., Xue, L., Gutowska-Owsiak, D., Wang, X., Huang, L.-C., Johnson, D., Scanlon, S. T., McKenzie, A. N. J., Fallon, P. G. & Ogg, G. S. A role for IL-25 and IL-33–driven type-2 innate lymphoid cells in atopic dermatitis. *J Exp Med* **210,** 2939–2950 (2013).

13. Kim, B. S., Siracusa, M. C., Saenz, S. A., Noti, M., Monticelli, L. A., Sonnenberg, G. F.,
Hepworth, M. R., Voorhees, A. S. V., Comeau, M. R. & Artis, D. TSLP elicits IL-33-
independent innate lymphoid cell responses to promote skin inflammation. *Sci Transl Med* **5,**
170ra16 (2013).

14. Tsou, A. M., Yano, H., Parkhurst, C. N., Mahlakõiv, T., Chu, C., Zhang, W., He, Z., Jarick,
K. J., Zhong, C., Putzel, G. G., Hatazaki, M., Consortium, J. I. L. C. B., Longman, R.,
Sonnenberg, G., Scherl, E., Lukin, D., Battat, R., Sockolow, R., Ciecierega, T., Solomon, A.,
Barfield, E., Chien, K., Ferreira, J., Williams, J., Khan, S., Chong, P. S., Mozumder, S., Chou,
L., Zhou, W., Ahmed, A., Joseph, A. M., Lorenz, I. C., Andrew, D., Balderes, P., Klose, C. S.
N., Lira, S. A. & Artis, D. Neuropeptide regulation of non-redundant ILC2 responses at barrier
surfaces. *Nature* **611,** 787–793 (2022).

15. Jarick, K. J., Topczewska, P. M., Jakob, M. O., Yano, H., Arifuzzaman, M., Gao, X.,
Boulekou, S., Stokic-Trtica, V., Leclère, P. S., Preußer, A., Rompe, Z. A., Stamm, A., Tsou, A.
M., Chu, C., Heinrich, F. R., Guerra, G. M., Durek, P., Ivanov, A., Beule, D., Helfrich, S., Duerr,
C. U., Kühl, A. A., Stehle, C., Romagnani, C., Mashreghi, M.-F., Diefenbach, A., Artis, D. &
Klose, C. S. N. Non-redundant functions of group 2 innate lymphoid cells. *Nature* **611,** 794–800
(2022).

16. Fort, M. M., Cheung, J., Yen, D., Li, J., Zurawski, S. M., Lo, S., Menon, S., Clifford, T.,
Hunte, B., Lesley, R., Muchamuel, T., Hurst, S. D., Zurawski, G., Leach, M. W., Gorman, D. M.
& Rennick, D. M. IL-25 Induces IL-4, IL-5, and IL-13 and Th2-Associated Pathologies In Vivo.
Immunity **15,** 985–995 (2001).

17. Hurst, S. D., Muchamuel, T., Gorman, D. M., Gilbert, J. M., Clifford, T., Kwan, S., Menon,
S., Seymour, B., Jackson, C., Kung, T. T., Brieland, J. K., Zurawski, S. M., Chapman, R. W.,
Zurawski, G. & Coffman, R. L. New IL-17 Family Members Promote Th1 or Th2 Responses in
the Lung: In Vivo Function of the Novel Cytokine IL-25. *J Immunol* **169,** 443–453 (2002).

18. Fallon, P. G., Ballantyne, S. J., Mangan, N. E., Barlow, J. L., Dasvarma, A., Hewett, D. R.,
McIlgorm, A., Jolin, H. E. & McKenzie, A. N. J. Identification of an interleukin (IL)-25–
dependent cell population that provides IL-4, IL-5, and IL-13 at the onset of helminth expulsion.
J Exp Medicine **203,** 1105–1116 (2006).

19. Moro, K., Yamada, T., Tanabe, M., Takeuchi, T., Ikawa, T., Kawamoto, H., Furusawa, J.,
Ohtani, M., Fujii, H. & Koyasu, S. Innate production of TH2 cytokines by adipose tissue-
associated c-Kit+Sca-1+ lymphoid cells. *Nature* **463,** 540–544 (2010).

20. Saenz, S. A., Siracusa, M. C., Perrigoue, J. G., Spencer, S. P., Jr, J. F. U., Tocker, J. E.,
Budelsky, A. L., Kleinschek, M. A., Kastelein, R. A., Kambayashi, T., Bhandoola, A. & Artis,
D. IL25 elicits a multipotent progenitor cell population that promotes TH2 cytokine responses.
Nature **464,** 1362–1366 (2010).

21. Neill, D. R., Wong, S. H., Bellosi, A., Flynn, R. J., Daly, M., Langford, T. K. A., Bucks, C.,
Kane, C. M., Fallon, P. G., Pannell, R., Jolin, H. E. & McKenzie, A. N. J. Nuocytes represent a
new innate effector leukocyte that mediates type-2 immunity. *Nature* **464,** 1367–1370 (2010).

22. Price, A. E., Liang, H.-E., Sullivan, B. M., Reinhardt, R. L., Eisley, C. J., Erle, D. J. &
Locksley, R. M. Systemically dispersed innate IL-13-expressing cells in type 2 immunity. *P Natl*
Acad Sci Usa **107,** 11489–94 (2010).

23. Yang, Q., Saenz, S. A., Zlotoff, D. A., Artis, D. & Bhandoola, A. Cutting Edge: Natural
Helper Cells Derive from Lymphoid Progenitors. *J Immunol* **187,** 5505–5509 (2011).

24. Karo, J. M., Schatz, D. G. & Sun, J. C. The RAG Recombinase Dictates Functional
Heterogeneity and Cellular Fitness in Natural Killer Cells. *Cell* **159,** 94–107 (2014).

25. Bredemeyer, A. L., Helmink, B. A., Innes, C. L., Calderon, B., McGinnis, L. M., Mahowald,
G. K., Gapud, E. J., Walker, L. M., Collins, J. B., Weaver, B. K., Mandik-Nayak, L., Schreiber,
R. D., Allen, P. M., May, M. J., Paules, R. S., Bassing, C. H. & Sleckman, B. P. DNA double-
strand breaks activate a multi-functional genetic program in developing lymphocytes. *Nature*
456, 819–823 (2008).

26. Bednarski, J. J., Nickless, A., Bhattacharya, D., Amin, R. H., Schlissel, M. S. & Sleckman,
B. P. RAG-induced DNA double-strand breaks signal through Pim2 to promote pre–B cell
survival and limit proliferation. *J. Exp. Med.* **209,** 11–17 (2012).

27. Teng, G., Maman, Y., Resch, W., Kim, M., Yamane, A., Qian, J., Kieffer-Kwon, K.-R.,
Mandal, M., Ji, Y., Meffre, E., Clark, M. R., Cowell, L. G., Casellas, R. & Schatz, D. G. RAG
Represents a Widespread Threat to the Lymphocyte Genome. *Cell* **162,** 751–765 (2015).

28. Li, M., Hener, P., Zhang, Z., Kato, S., Metzger, D. & Chambon, P. Topical vitamin D3 and
low-calcemic analogs induce thymic stromal lymphopoietin in mouse keratinocytes and trigger
an atopic dermatitis. *Proc National Acad Sci* **103,** 11736–11741 (2006).

29. Liu, C., Zhang, Y., Liu, C. C. & Schatz, D. G. Structural insights into the evolution of the
RAG recombinase. *Nat Rev Immunol* 1–18 (2021). doi:10.1038/s41577-021-00628-6

30. Shih, H.-Y., Sciumè, G., Mikami, Y., Guo, L., Sun, H.-W., Brooks, S. R., Urban, J. F., Davis,
F. P., Kanno, Y. & O'Shea, J. J. Developmental Acquisition of Regulomes Underlies Innate
Lymphoid Cell Functionality. *Cell* **165,** 1120–1133 (2016).

31. Dyken, S. J. V., Nussbaum, J. C., Lee, J., Molofsky, A. B., Liang, H.-E., Pollack, J. L., Gate,
R. E., Haliburton, G. E., Ye, C. J., Marson, A., Erle, D. J. & Locksley, R. M. A tissue checkpoint
regulates type 2 immunity. *Nat Immunol* **17,** 1381–1387 (2016).

32. Turka, L. A., Schatz, D. G., Oettinger, M. A., Chun, J. J. M., Gorka, C., Lee, K.,
McCormack, W. T. & Thompson, C. B. Thymocyte Expression of RAG-1 and RAG-2:
Termination by T Cell Receptor Cross-Linking. *Science* **253,** 778–781 (1991).

33. Welner, R. S., Esplin, B. L., Garrett, K. P., Pelayo, R., Luche, H., Fehling, H. J. & Kincade,
P. W. Asynchronous RAG-1 Expression during B Lymphopoiesis. *J Immunol* **183,** 7768–7777
(2009).

34. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y.,
Stoeckius, M., Smibert, P. & Satija, R. Comprehensive Integration of Single-Cell Data. *Cell* **177,**
1888-1902.e21 (2019).

35. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., Lee, M. J.,
Wilk, A. J., Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain,
J., Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., Rogers, A. J., McElrath, J. M., Blish, C.
A., Gottardo, R., Smibert, P. & Satija, R. Integrated analysis of multimodal single-cell data. *Cell*
(2021). doi:10.1016/j.cell.2021.04.048

36. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36,**
411–420 (2018).

37. Stuart, T., Srivastava, A., Madad, S., Lareau, C. A. & Satija, R. Single-cell chromatin state
analysis with Signac. *Nat Methods* **18,** 1333–1341 (2021).

38. Ghaedi, M., Shen, Z. Y., Orangi, M., Martinez-Gonzalez, I., Wei, L., Lu, X., Das, A.,
Heravi-Moussavi, A., Marra, M. A., Bhandoola, A. & Takei, F. Single-cell analysis of RORα
tracer mouse lung reveals ILC progenitors and effector ILC2 subsets. *J Exp Med* **217,** e20182293
(2019).

39. Maazi, H., Patel, N., Sankaranarayanan, I., Suzuki, Y., Rigas, D., Soroosh, P., Freeman, G.
J., Sharpe, A. H. & Akbari, O. ICOS:ICOS-Ligand Interaction Is Required for Type 2 Innate
Lymphoid Cell Function, Homeostasis, and Induction of Airway Hyperreactivity. *Immunity* **42,**
538–551 (2015).

40. Paclik, D., Stehle, C., Lahmann, A., Hutloff, A. & Romagnani, C. ICOS regulates the pool of
group 2 innate lymphoid cells under homeostatic and inflammatory conditions in mice. *Eur J*
Immunol **45,** 2766–2772 (2015).

41. Roediger, B., Kyle, R., Yip, K. H., Sumaria, N., Guy, T. V., Kim, B. S., Mitchell, A. J., Tay,
S. S., Jain, R., Forbes-Blom, E., Chen, X., Tong, P. L., Bolton, H. A., Artis, D., Paul, W. E.,
Groth, B. F. de S., Grimbaldeston, M. A., Gros, G. L. & Weninger, W. Cutaneous
immunosurveillance and regulation of inflammation by group 2 innate lymphoid cells. *Nat*
Immunol **14,** 564–573 (2013).

42. Roediger, B., Kyle, R., Tay, S. S., Mitchell, A. J., Bolton, H. A., Guy, T. V., Tan, S.-Y.,
Forbes-Blom, E., Tong, P. L., Köller, Y., Shklovskaya, E., Iwashima, M., McCoy, K. D., Gros,
G. L., Groth, B. F. de S. & Weninger, W. IL-2 is a critical regulator of group 2 innate lymphoid
cell function during pulmonary inflammation. *J Allergy Clin Immun* **136,** 1653-1663.e7 (2015).

43. Zeis, P., Lian, M., Fan, X., Herman, J. S., Hernandez, D. C., Gentek, R., Elias, S., Symowski,
C., Knöpper, K., Peltokangas, N., Friedrich, C., Doucet-Ladeveze, R., Kabat, A. M., Locksley,
R. M., Voehringer, D., Bajenoff, M., Rudensky, A. Y., Romagnani, C., Grün, D. & Gasteiger, G.
In Situ Maturation and Tissue Adaptation of Type 2 Innate Lymphoid Cell Progenitors. *Immunity*
53, 775-792.e9 (2020).

44. Ricardo-Gonzalez, R. R., Dyken, S. J. V., Schneider, C., Lee, J., Nussbaum, J. C., Liang, H.-
E., Vaka, D., Eckalbar, W. L., Molofsky, A. B., Erle, D. J. & Locksley, R. M. Tissue signals
imprint ILC2 identity with anticipatory function. *Nat Immunol* **19,** 1093–1099 (2018).

45. Cardoso, V., Chesne, J., Ribeiro, H., Garcia-Cassani, B., Carvalho, T., Bouchery, T., Shah,
K., Barbosa-Morais, N. L., Harris, N. & Veiga-Fernandes, H. Neuronal regulation of type 2
innate lymphoid cells via neuromedin U. *Nature* **549,** 277–281 (2014).

46. Klose, C. S. N., Mahlakõiv, T., Moeller, J. B., Rankin, L. C., Flamar, A.-L., Kabata, H.,
Monticelli, L. A., Moriyama, S., Putzel, G. G., Rakhilin, N., Shen, X., Kostenis, E., König, G.
M., Senda, T., Carpenter, D., Farber, D. L. & Artis, D. The neuropeptide neuromedin U
stimulates innate lymphoid cells and type 2 inflammation. *Nature* **549,** 282–286 (2017).

47. Wallrapp, A., Riesenfeld, S. J., Burkett, P. R., Abdulnour, R.-E. E., Nyman, J., Dionne, D.,
Hofree, M., Cuoco, M. S., Rodman, C., Farouq, D., Haas, B. J., Tickle, T. L., Trombetta, J. J.,
Baral, P., Klose, C. S. N., Mahlakõiv, T., Artis, D., Rozenblatt-Rosen, O., Chiu, I. M., Levy, B.
D., Kowalczyk, M. S., Regev, A. & Kuchroo, V. K. The neuropeptide NMU amplifies ILC2-
driven allergic lung inflammation. *Nature* **549,** 351–356 (2017).

48. Mjösberg, J. M., Trifari, S., Crellin, N. K., Peters, C. P., Drunen, C. M. van, Piet, B.,
Fokkens, W. J., Cupedo, T. & Spits, H. Human IL-25- and IL-33-responsive type 2 innate
lymphoid cells are defined by expression of CRTH2 and CD161. *Nat Immunol* **12,** 1055–1062
(2011).

49. Hung, L.-Y., Lewkowich, I. P., Dawson, L. A., Downey, J., Yang, Y., Smith, D. E. &
Herbert, D. R. IL-33 drives biphasic IL-13 production for noncanonical Type 2 immunity against
hookworms. *Proc National Acad Sci* **110,** 282–287 (2013).

50. Mjösberg, J., Bernink, J., Golebski, K., Karrich, J. J., Peters, C. P., Blom, B., te Velde, A. A.,
Fokkens, W. J., van Drunen, C. M. & Spits, H. The Transcription Factor GATA3 Is Essential for
the Function of Human Type 2 Innate Lymphoid Cells. *Immunity* **37,** 649–659 (2012).

51. Hoyler, T., Klose, C. S. N., Souabni, A., Turqueti-Neves, A., Pfeifer, D., Rawlins, E. L.,
Voehringer, D., Busslinger, M. & Diefenbach, A. The Transcription Factor GATA-3 Controls
Cell Fate and Maintenance of Type 2 Innate Lymphoid Cells. *Immunity* **37,** 634–648 (2012).

52. Wolterink, R. G. J. K., Serafini, N., Nimwegen, M. van, Vosshenrich, C. A. J., Bruijn, M. J.
W. de, Pereira, D. F., Fernandes, H. V., Hendriks, R. W. & Santo, J. P. D. Essential, dose-
dependent role for the transcription factor Gata3 in the development of IL-5+ and IL-13+ type 2
innate lymphoid cells. *Proc National Acad Sci* **110,** 10240–10245 (2013).

53. Pokrovskii, M., Hall, J. A., Ochayon, D. E., Yi, R., Chaimowitz, N. S., Seelamneni, H.,
Carriero, N., Watters, A., Waggoner, S. N., Littman, D. R., Bonneau, R. & Miraldi, E. R.
Characterization of Transcriptional Regulatory Networks that Promote and Restrict Identities and
Functions of Intestinal Innate Lymphoid Cells. *Immunity* **51,** 185-197.e6 (2019).

54. Califano, D., Cho, J. J., Uddin, M. N., Lorentsen, K. J., Yang, Q., Bhandoola, A., Li, H. &
Avram, D. Transcription Factor Bcl11b Controls Identity and Function of Mature Type 2 Innate
Lymphoid Cells. *Immunity* **43,** 354–368 (2015).

55. Walker, J. A., Oliphant, C. J., Englezakis, A., Yu, Y., Clare, S., Rodewald, H.-R., Belz, G.,
Liu, P., Fallon, P. G. & McKenzie, A. N. J. Bcl11b is essential for group 2 innate lymphoid cell
development. *J Exp Med* **212,** 875–882 (2015).

56. Yu, Y., Wang, C., Clare, S., Wang, J., Lee, S.-C., Brandt, C., Burke, S., Lu, L., He, D.,
Jenkins, N. A., Copeland, N. G., Dougan, G. & Liu, P. The transcription factor Bcl11b is
specifically expressed in group 2 innate lymphoid cells and is essential for their development. *J*
Exp Med **212,** 865–874 (2015).

57. Hosokawa, H., Romero-Wolf, M., Yang, Q., Motomura, Y., Levanon, D., Groner, Y., Moro,
K., Tanaka, T. & Rothenberg, E. V. Cell type–specific actions of Bcl11b in early T-lineage and
group 2 innate lymphoid cells. *J Exp Med* **217,** e20190972 (2019).

58. Björklund, Å. K., Forkel, M., Picelli, S., Konya, V., Theorell, J., Friberg, D., Sandberg, R. &
Mjösberg, J. The heterogeneity of human CD127+ innate lymphoid cells revealed by single-cell
RNA sequencing. *Nat Immunol* **17,** 451–460 (2016).

59. Trabanelli, S., Ercolano, G., Wyss, T., Gomez-Cadena, A., Falquet, M., Cropp, D., Imbratta,
C., Leblond, M. M., Salvestrini, V., Curti, A., Adotevi, O., Jandus, C. & Verdeil, G. c-Maf
enforces cytokine production and promotes memory-like responses in mouse and human type 2
innate lymphoid cells. *Embo J* **41,** e109300 (2022).

60. Zook, E. C., Ramirez, K., Guo, X., Voort, G. van der, Sigvardsson, M., Svensson, E. C., Fu,
Y.-X. & Kee, B. L. The ETS1 transcription factor is required for the development and cytokine-
induced expansion of ILC2. *J Exp Med* **213,** 687–696 (2016).

61. Wong, S. H., Walker, J. A., Jolin, H. E., Drynan, L. F., Hams, E., Camelo, A., Barlow, J. L.,
Neill, D. R., Panova, V., Koch, U., Radtke, F., Hardman, C. S., Hwang, Y. Y., Fallon, P. G. &
McKenzie, A. N. J. Transcription factor RORα is critical for nuocyte development. *Nat Immunol*
13, 229–236 (2012).

62. Halim, T. Y. F., MacLaren, A., Romanish, M. T., Gold, M. J., McNagny, K. M. & Takei, F.
Retinoic-Acid-Receptor-Related Orphan Nuclear Receptor Alpha Is Required for Natural Helper
Cell Development and Allergic Inflammation. *Immunity* **37,** 463–474 (2012).

63. Ferreira, A. C. F., Szeto, A. C. H., Heycock, M. W. D., Clark, P. A., Walker, J. A., Crisp, A.,
Barlow, J. L., Kitching, S., Lim, A., Gogoi, M., Berks, R., Daly, M., Jolin, H. E. & McKenzie,

A. N. J. RORα is a critical checkpoint for T cell and ILC2 commitment in the embryonic thymus. *Nat Immunol* **22,** 166–178 (2021).

64. Miyajima, Y., Ealey, K. N., Motomura, Y., Mochizuki, M., Takeno, N., Yanagita, M., Economides, A. N., Nakayama, M., Koseki, H. & Moro, K. Effects of BMP7 produced by group 2 innate lymphoid cells on adipogenesis. *Int Immunol* **32,** 407–419 (2020).

65. Chen, W.-Y., Wu, Y.-H., Tsai, T.-H., Li, R.-F., Lai, A. C.-Y., Li, L.-C., Yang, J.-L. & Chang, Y.-J. Group 2 innate lymphoid cells contribute to IL-33-mediated alleviation of cardiac fibrosis. *Theranostics* **11,** 2594–2611 (2021).

66. Xu, H., Ding, J., Porter, C. B. M., Wallrapp, A., Tabaka, M., Ma, S., Fu, S., Guo, X., Riesenfeld, S. J., Su, C., Dionne, D., Nguyen, L. T., Lefkovith, A., Ashenberg, O., Burkett, P. R., Shi, H. N., Rozenblatt-Rosen, O., Graham, D. B., Kuchroo, V. K., Regev, A. & Xavier, R. J. Transcriptional Atlas of Intestinal Immune Cells Reveals that Neuropeptide α-CGRP Modulates Group 2 Innate Lymphoid Cell Responses. *Immunity* **51,** 696-708.e9 (2019).

67. Vivier, E., Pavert, S. A. van de, Cooper, M. D. & Belz, G. T. The evolution of innate lymphoid cells. *Nat Immunol* **17,** 790–794 (2016).

68. Schneider, C., Lee, J., Koga, S., Ricardo-Gonzalez, R. R., Nussbaum, J. C., Smith, L. K., Villeda, S. A., Liang, H.-E. & Locksley, R. M. Tissue-Resident Group 2 Innate Lymphoid Cells Differentiate by Layered Ontogeny and In Situ Perinatal Priming. *Immunity* **50,** 1425-1438.e5 (2019).

69. Bielecki, P., Riesenfeld, S. J., Hütter, J.-C., Triglia, E. T., Kowalczyk, M. S., Ricardo-Gonzalez, R. R., Lian, M., Vesely, M. C. A., Kroehling, L., Xu, H., Slyper, M., Muus, C., Ludwig, L. S., Christian, E., Tao, L., Kedaigle, A. J., Steach, H. R., York, A. G., Skadow, M. H., Yaghoubi, P., Dionne, D., Jarret, A., McGee, H. M., Porter, C. B. M., Licona-Limón, P., Bailis, W., Jackson, R., Gagliani, N., Gasteiger, G., Locksley, R. M., Regev, A. & Flavell, R. A. Skin-resident innate lymphoid cells converge on a pathogenic effector state. *Nature* **592,** 128–132 (2021).

70. Sun, Z., Han, S., Zhu, X., Sato, A. & Islam, S. A. CCR8 Signaling Regulates Skin ILC2s to Promote Chronic Atopic Dermatitis. *Ssrn Electron J* (2022). doi:10.2139/ssrn.4135728

71. Knipfer, L., Schulz-Kuhnt, A., Kindermann, M., Greif, V., Symowski, C., Voehringer, D., Neurath, M. F., Atreya, I. & Wirtz, S. A CCL1/CCR8-dependent feed-forward mechanism drives ILC2 functions in type 2–mediated inflammation. *J Exp Med* **216,** 2763–2777 (2019).

72. Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., Paulovich, A., Pomeroy, S. L., Golub, T. R., Lander, E. S. & Mesirov, J. P. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc National Acad Sci* **102,** 15545–15550 (2005).

73. Mootha, V. K., Lindgren, C. M., Eriksson, K.-F., Subramanian, A., Sihag, S., Lehar, J.,
Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., Houstis, N., Daly, M. J., Patterson, N.,
Mesirov, J. P., Golub, T. R., Tamayo, P., Spiegelman, B., Lander, E. S., Hirschhorn, J. N.,
Altshuler, D. & Groop, L. C. PGC-1α-responsive genes involved in oxidative phosphorylation
are coordinately downregulated in human diabetes. *Nat Genet* **34,** 267–273 (2003).

74. Ma, S., Zhang, B., LaFave, L. M., Earl, A. S., Chiang, Z., Hu, Y., Ding, J., Brack, A., Kartha,
V. K., Tay, T., Law, T., Lareau, C., Hsu, Y.-C., Regev, A. & Buenrostro, J. D. Chromatin
Potential Identified by Shared Single-Cell Profiling of RNA and Chromatin. *Cell* **183,** 1103-
1116.e20 (2020).

75. Constantinides, M. G., McDonald, B. D., Verhoef, P. A. & Bendelac, A. A committed
precursor to innate lymphoid cells. *Nature* **508,** 397–401 (2014).

76. Seehus, C. R., Aliahmad, P., Torre, B. de la, Iliev, I. D., Spurka, L., Funari, V. A. & Kaye, J.
Innate lymphoid cell development requires TOX-dependent generation of a common ILC
progenitor. *Nat Immunol* **16,** 599–608 (2015).

77. Aliahmad, P., Torre, B. de la & Kaye, J. Shared dependence on the DNA-binding factor TOX
for the development of lymphoid tissue–inducer cell and NK cell lineages. *Nat Immunol* **11,**
945–952 (2010).

78. Aliahmad, P. & Kaye, J. Development of all CD4 T lineages requires nuclear factor TOX. *J*
Exp Medicine **205,** 245–256 (2008).

79. Kasal, D. N., Liang, Z., Hollinger, M. K., O'Leary, C. Y., Lisicka, W., Sperling, A. I. &
Bendelac, A. A Gata3 enhancer necessary for ILC2 development and function. *Proc National*
Acad Sci **118,** e2106311118 (2021).

80. Wei, L., Vahedi, G., Sun, H.-W., Watford, W. T., Takatori, H., Ramos, H. L., Takahashi, H.,
Liang, J., Gutierrez-Cruz, G., Zang, C., Peng, W., O'Shea, J. J. & Kanno, Y. Discrete Roles of
STAT4 and STAT6 Transcription Factors in Tuning Epigenetic Modifications and Transcription
during T Helper Cell Differentiation. *Immunity* **32,** 840–851 (2010).

81. Onodera, A., Yamashita, M., Endo, Y., Kuwahara, M., Tofukuji, S., Hosokawa, H., Kanai,
A., Suzuki, Y. & Nakayama, T. STAT6-mediated displacement of polycomb by trithorax
complex establishes long-term maintenance of GATA3 expression in T helper type 2 cells. *J Exp*
Med **207,** 2493–2506 (2010).

82. Barlow, J. L., Peel, S., Fox, J., Panova, V., Hardman, C. S., Camelo, A., Bucks, C., Wu, X.,
Kane, C. M., Neill, D. R., Flynn, R. J., Sayers, I., Hall, I. P. & McKenzie, A. N. J. IL-33 is more
potent than IL-25 in provoking IL-13–producing nuocytes (type 2 innate lymphoid cells) and
airway contraction. *J Allergy Clin Immun* **132,** 933–941 (2013).

83. Molofsky, A. B., Van Gool, F., Liang, H.-E., Van Dyken, S. J., Nussbaum, J. C., Lee, J.,
Bluestone, J. A. & Locksley, R. M. Interleukin-33 and Interferon-γ Counter-Regulate Group 2
Innate Lymphoid Cell Activation during Immune Perturbation. *Immunity* **43,** 161–174 (2015).

84. Oliver, P. M., Cao, X., Worthen, G. S., Shi, P., Briones, N., MacLeod, M., White, J., Kirby,
P., Kappler, J., Marrack, P. & Yang, B. Ndfip1 Protein Promotes the Function of Itch Ubiquitin
Ligase to Prevent T Cell Activation and T Helper 2 Cell-Mediated Inflammation. *Immunity* **25,**
929–940 (2006).

85. Altin, J. A., Daley, S. R., Howitt, J., Rickards, H. J., Batkin, A. K., Horikawa, K., Prasad, S.
J., Nelms, K. A., Kumar, S., Wu, L. C., Tan, S.-S., Cook, M. C. & Goodnow, C. C. Ndfip1
mediates peripheral tolerance to self and exogenous antigen by inducing cell cycle exit in
responding CD4+ T cells. *Proc National Acad Sci* **111,** 2067–2074 (2014).

86. Ferreira, M. A., Matheson, M. C., Duffy, D. L., Marks, G. B., Hui, J., Souëf, P. L., Danoy,
P., Baltic, S., Nyholt, D. R., Jenkins, M., Hayden, C., Willemsen, G., Ang, W., Kuokkanen, M.,
Beilby, J., Cheah, F., Geus, E. J. de, Ramasamy, A., Vedantam, S., Salomaa, V., Madden, P. A.,
Heath, A. C., Hopper, J. L., Visscher, P. M., Musk, B., Leeder, S. R., Jarvelin, M.-R., Pennell,
C., Boomsma, D. I., Hirschhorn, J. N., Walters, H., Martin, N. G., James, A., Jones, G.,
Abramson, M. J., Robertson, C. F., Dharmage, S. C., Brown, M. A., Montgomery, G. W.,
Thompson, P. J. & Consortium, for the A. A. G. Identification of IL6R and chromosome 11q13.5
as risk loci for asthma. *Lancet* **378,** 1006–1014 (2011).

87. Chi, H., Barry, S. P., Roth, R. J., Wu, J. J., Jones, E. A., Bennett, A. M. & Flavell, R. A.
Dynamic regulation of pro- and anti-inflammatory cytokines by MAPK phosphatase 1 (MKP-1)
in innate immune responses. *Proc National Acad Sci* **103,** 2274–2279 (2006).

88. Clark, A. R., Martins, J. R. S. & Tchen, C. R. Role of Dual Specificity Phosphatases in
Biological Responses to Glucocorticoids*. *J Biol Chem* **283,** 25765–25769 (2008).

89. Grosche, S., Marenholz, I., Esparza-Gordillo, J., Arnau-Soler, A., Pairo-Castineira, E.,
Rüschendorf, F., Ahluwalia, T. S., Almqvist, C., Arnold, A., (AAGC), A. A. G. C., Baurecht, H.,
Bisgaard, H., Bønnelykke, K., Brown, S. J., Bustamante, M., Curtin, J. A., Custovic, A.,
Dharmage, S. C., Esplugues, A., Falchi, M., Fernandez-Orth, D., Ferreira, M. A. R., Franke, A.,
Gerdes, S., Gieger, C., Hakonarson, H., Holt, P. G., Homuth, G., Hubner, N., Hysi, P. G.,
Jarvelin, M.-R., Karlsson, R., Koppelman, G. H., Lau, S., Lutz, M., Magnusson, P. K. E., Marks,
G. B., Müller-Nurasyid, M., Nöthen, M. M., Paternoster, L., Pennell, C. E., Peters, A., Rawlik,
K., Robertson, C. F., Rodriguez, E., Sebert, S., Simpson, A., Sleiman, P. M. A., Standl, M.,
Stölzl, D., Strauch, K., Szwajda, A., Tenesa, A., Thompson, P. J., Ullemar, V., Visconti, A.,
Vonk, J. M., Wang, C. A., Weidinger, S., Wielscher, M., Worth, C. L., Xu, C.-J. & Lee, Y.-A.
Rare variant analysis in eczema identifies exonic variants in DUSP1, NOTCH4 and SLC9A4.
Nat Commun **12,** 6618 (2021).

90. Xie, M., Lu, C., Wang, J., McLellan, M. D., Johnson, K. J., Wendl, M. C., McMichael, J. F.,
Schmidt, H. K., Yellapantula, V., Miller, C. A., Ozenberger, B. A., Welch, J. S., Link, D. C.,
Walter, M. J., Mardis, E. R., Dipersio, J. F., Chen, F., Wilson, R. K., Ley, T. J. & Ding, L. Age-

related mutations associated with clonal hematopoietic expansion and malignancies. *Nat Med* **20,**
1472–1478 (2014).

91. Giulio, G., K., K. A., E., H. R., Johan, L., A., R. S., F., B. S., Kimberly, C., Eran, M., M., N.
B., Menachem, F., M., P. S., Oscar, S., Mikael, L., Martin, H., Sören, L., B., G. S., L., M. J., S.,
L. E., F., S. P., Pamela, S., Henrik, G., M., H. C. & A., M. S. Clonal Hematopoiesis and Blood-
Cancer Risk Inferred from Blood DNA Sequence. *New Engl J Med* **371,** 2477–2487 (2014).

92. Siddhartha, J., Pierre, F., Jason, F., Alisa, M., V., G. P., G., M. B., Coleman, L. R., H., M. C.,
Noel, B., Alejandro, C., M., H. J., Vladislav, M., C., K. F., J., K. M., Brian, H., Leena, K., A., K.
H., Claes, L., Gad, G., Adolfo, C., F., B. B., Stacey, G., Sekar, K., M., S. H., I., M. M., Michael,
B., Jaakko, T., Christopher, H., Leif, G., Gil, A., G., W. J., Donna, N., David, A. & L., E. B.
Age-Related Clonal Hematopoiesis Associated with Adverse Outcomes. *New Engl J Med* **371,**
2488–2498 (2014).

93. Nagase, R., Inoue, D., Pastore, A., Fujino, T., Hou, H.-A., Yamasaki, N., Goyama, S., Saika,
M., Kanai, A., Sera, Y., Horikawa, S., Ota, Y., Asada, S., Hayashi, Y., Kawabata, K. C., Takeda,
R., Tien, H.-F., Honda, H., Abdel-Wahab, O. & Kitamura, T. Expression of mutant Asxl1
perturbs hematopoiesis and promotes susceptibility to leukemic transformation. *J Exp Med* **215,**
1729–1747 (2018).

94. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat Methods* **14,**
975–978 (2017).

95. Bando, J. K. & Colonna, M. Innate lymphoid cell function in the context of adaptive
immunity. *Nat Immunol* **17,** 783–789 (2016).

96. Hart, G. T., Hogquist, K. A. & Jameson, S. C. Krüppel-like Factors in Lymphocyte Biology.
J Immunol **188,** 521–526 (2012).

97. Cao, Z., Sun, X., Icli, B., Wara, A. K. & Feinberg, M. W. Role of Krüppel-like factors in
leukocyte development, function, and disease. *Blood* **116,** 4404–4414 (2010).

98. Weinreich, M. A., Takada, K., Skon, C., Reiner, S. L., Jameson, S. C. & Hogquist, K. A.
KLF2 Transcription-Factor Deficiency in T Cells Results in Unrestrained Cytokine Production
and Upregulation of Bystander Chemokine Receptors. *Immunity* **31,** 122–130 (2009).

99. Godin-Heymann, N., Brabetz, S., Murillo, M. M., Saponaro, M., Santos, C. R., Lobley, A.,
East, P., Chakravarty, P., Matthews, N., Kelly, G., Jordan, S., Castellano, E. & Downward, J.
Tumour-suppression function of KLF12 through regulation of anoikis. *Oncogene* **35,** 3324–3334
(2016).

100. Narla, G., Heath, K. E., Reeves, H. L., Li, D., Giono, L. E., Kimmelman, A. C., Glucksman,
M. J., Narla, J., Eng, F. J., Chan, A. M., Ferrari, A. C., Martignetti, J. A. & Friedman, S. L.

KLF6, a Candidate Tumor Suppressor Gene Mutated in Prostate Cancer. *Science* **294,** 2563–2566 (2001).

101. Kuo, C. T., Veselits, M. L. & Leiden, J. M. LKLF: A Transcriptional Regulator of Single-Positive T Cell Quiescence and Survival. *Science* **277,** 1986–1990 (1997).

102. Buckley, A. F., Kuo, C. T. & Leiden, J. M. Transcription factor LKLF is sufficient to program T cell quiescence via a c-Myc–dependent pathway. *Nat Immunol* **2,** 698–704 (2001).

103. Schuettpelz, L. G., Gopalan, P. K., Giuste, F. O., Romine, M. P., Os, R. van & Link, D. C. Kruppel-like factor 7 overexpression suppresses hematopoietic stem and progenitor cell function. *Blood* **120,** 2981–2989 (2012).

104. Kobayashi, T., Voisin, B., Kim, D. Y., Kennedy, E. A., Jo, J.-H., Shih, H.-Y., Truong, A., Doebel, T., Sakamoto, K., Cui, C.-Y., Schlessinger, D., Moro, K., Nakae, S., Horiuchi, K., Zhu, J., Leonard, W. J., Kong, H. H. & Nagao, K. Homeostatic Control of Sebaceous Glands by Innate Lymphoid Cells Regulates Commensal Bacteria Equilibrium. *Cell* **176,** 982-997.e16 (2019).

105. Lee, G. R., Fields, P. E., Griffin, T. J. & Flavell, R. A. Regulation of the Th2 Cytokine Locus by a Locus Control Region. *Immunity* **19,** 145–153 (2003).

106. Loots, G. G., Locksley, R. M., Blankespoor, C. M., Wang, Z. E., Miller, W., Rubin, E. M. & Frazer, K. A. Identification of a Coordinate Regulator of Interleukins 4, 13, and 5 by Cross-Species Sequence Comparisons. *Science* **288,** 136–140 (2000).

107. Fields, P. E., Lee, G. R., Kim, S. T., Bartsevich, V. V. & Flavell, R. A. Th2-Specific Chromatin Remodeling and Enhancer Activity in the Th2 Cytokine Locus Control Region. *Immunity* **21,** 865–876 (2004).

108. Lentjes, M. H., Niessen, H. E., Akiyama, Y., Bruïne, A. P. de, Melotte, V. & Engeland, M. van. The emerging role of GATA transcription factors in development and disease. *Expert Rev Mol Med* **18,** e3 (2016).

109. Zhang, Y., Cheng, T. C., Huang, G., Lu, Q., Surleac, M. D., Mandell, J. D., Pontarotti, P., Petrescu, A. J., Xu, A., Xiong, Y. & Schatz, D. G. Transposon molecular domestication and the evolution of the RAG recombinase. *Nature* **569,** 79–84 (2019).

110. Buckley, R. H. Molecular Defects in Human Severe Combined Immunodeficiency and Approaches to Immune Reconstitution. *Annu Rev Immunol* **22,** 625–655 (2004).

111. Mombaerts, P., Iacomini, J., Johnson, R. S., Herrup, K., Tonegawa, S. & Papaioannou, V. E. RAG-1-deficient mice have no mature B and T lymphocytes. *Cell* **68,** 869–877 (1992).

112. Shinkai, Y., Rathbun, 0Gary, Lam, K.-P., Oltz, E. M., Stewart, V., Mendelsohn, M.,
Charron, J., Datta, M., Young, F., Stall, A. M. & Alt, F. W. RAG-2-deficient mice lack mature
lymphocytes owing to inability to initiate V(D)J rearrangement. *Cell* **68,** 855–867 (1992).

113. Igarashi, H., Gregory, S. C., Yokota, T., Sakaguchi, N. & Kincade, P. W. Transcription
from the RAG1 Locus Marks the Earliest Lymphocyte Progenitors in Bone Marrow. *Immunity*
17, 117–130 (2002).

114. Pelayo, R., Hirose, J., Huang, J., Garrett, K. P., Delogu, A., Busslinger, M. & Kincade, P.
W. Derivation of 2 categories of plasmacytoid dendritic cells in murine bone marrow. *Blood* **105,**
4407–4415 (2005).

115. Wilson, A., Held, W. & MacDonald, H. R. Two waves of recombinase gene expression in
developing thymocytes. *J Exp Medicine* **179,** 1355–1360 (1994).

116. Gentek, R., Munneke, J. M., Helbig, C., Blom, B., Hazenberg, M. D., Spits, H. & Amsen,
D. Modulation of Signal Strength Switches Notch from an Inducer of T Cells to an Inducer of
ILC2. *Front Immunol* **4,** 334 (2013).

117. Rodríguez, C. I., Buchholz, F., Galloway, J., Sequerra, R., Kasper, J., Ayala, R., Stewart, A.
F. & Dymecki, S. M. High-efficiency deleter mice show that FLPe is an alternative to Cre-loxP.
Nat. Genet. **25,** 139–140 (2000).

118. Zhu, J., Yamane, H. & Paul, W. E. Differentiation of Effector CD4 T Cell Populations*.
Annu Rev Immunol **28,** 445–489 (2010).

119. Koues, O. I., Collins, P. L., Cella, M., Robinette, M. L., Porter, S. I., Pyfrom, S. C., Payton,
J. E., Colonna, M. & Oltz, E. M. Distinct Gene Regulatory Pathways for Human Innate versus
Adaptive Lymphoid Cells. *Cell* **165,** 1134–1146 (2016).

120. Agarwal, S. & Rao, A. Modulation of Chromatin Structure Regulates Cytokine Gene
Expression during T Cell Differentiation. *Immunity* **9,** 765–775 (1998).

121. Omenn, G. S. Familial Reticuloendotheliosis with Eosinophilia. *New Engl J Medicine* **273,**
427–432 (1965).

122. Villa, A., Notarangelo, L. D. & Roifman, C. M. Omenn syndrome: inflammation in leaky
severe combined immunodeficiency. *J Allergy Clin Immun* **122,** 1082–1086 (2008).

123. Villa, A., Santagata, S., Bozzi, F., Giliani, S., Frattini, A., Imberti, L., Gatta, L. B., Ochs, H.
D., Schwarz, K., Notarangelo, L. D., Vezzoni, P. & Spanopoulou, E. Partial V(D)J
recombination activity leads to Omenn syndrome. *Cell* **93,** 885–896 (1998).

124. Villa, A., Sobacchi, C., Notarangelo, L. D., Bozzi, F., Abinun, M., Abrahamsen, T. G.,
Arkwright, P. D., Baniyash, M., Brooks, E. G., Conley, M. E., Cortes, P., Duse, M., Fasth, A.,
Filipovich, A. M., Infante, A. J., Jones, A., Mazzolari, E., Muller, S. M., Pasic, S., Rechavi, G.,

Sacco, M. G., Santagata, S., Schroeder, M. L., Seger, R., Strina, D., Ugazio, A., Väliaho, J., Vihinen, M., Vogler, L. B., Ochs, H., Vezzoni, P., Friedrich, W. & Schwarz, K. V(D)J recombination defects in lymphocytes due to RAG mutations: severe immunodeficiency with a spectrum of clinical presentations. *Blood* **97,** 81–88 (2001).

125. Wada, T., Toma, T., Okamoto, H., Kasahara, Y., Koizumi, S., Agematsu, K., Kimura, H., Shimada, A., Hayashi, Y., Kato, M. & Yachie, A. Oligoclonal expansion of T lymphocytes with multiple second-site mutations leads to Omenn syndrome in a patient with RAG1-deficient severe combined immunodeficiency. *Blood* **106,** 2099–2101 (2005).

126. Marrella, V., Poliani, P. L., Casati, A., Rucci, F., Frascoli, L., Gougeon, M.-L., Lemercier, B., Bosticardo, M., Ravanini, M., Battaglia, M., Roncarolo, M. G., Cavazzana-Calvo, M., Facchetti, F., Notarangelo, L. D., Vezzoni, P., Grassi, F. & Villa, A. A hypomorphic R229Q Rag2 mouse mutant recapitulates human Omenn syndrome. *J Clin Invest* **117,** 1260–1269 (2007).

127. Khiong, K., Murakami, M., Kitabayashi, C., Ueda, N., Sawa, S., Sakamoto, A., Kotzin, B. L., Rozzo, S. J., Ishihara, K., Verella-Garcia, M., Kappler, J., Marrack, P. & Hirano, T. Homeostatically proliferating CD4+ T cells are involved in the pathogenesis of an Omenn syndrome murine model. *J Clin Invest* **117,** 1270–1281 (2007).

128. Milner, J. D., Ward, J. M., Keane-Myers, A. & Paul, W. E. Lymphopenic mice reconstituted with limited repertoire T cells develop severe, multiorgan, Th2-associated inflammatory disease. *Proc. Natl. Acad. Sci.* **104,** 576–581 (2007).

129. Pike-Overzet, K., Rodijk, M., Ng, Y.-Y., Baert, M. R. M., Lagresle-Peyrou, C., Schambach, A., Zhang, F., Hoeben, R. C., Hacein-Bey-Abina, S., Lankester, A. C., Bredius, R. G. M., Driessen, G. J. A., Thrasher, A. J., Baum, C., Cavazzana-Calvo, M., Dongen, J. J. M. van & Staal, F. J. T. Correction of murine Rag1 deficiency by self-inactivating lentiviral vector-mediated gene transfer. *Leukemia* **25,** 1471–1483 (2011).

130. Pike-Overzet, K., Baum, C., Bredius, R. G. M., Cavazzana, M., Driessen, G.-J., Fibbe, W. E., Gaspar, H. B., Hoeben, R. C., Lagresle-Peyrou, C., Lankester, A., Meij, P., Schambach, A., Thrasher, A., Dongen, J. J. M. V., Zwaginga, J.-J. & Staal, F. J. T. Successful RAG1-SCID gene therapy depends on the level of RAG1 expression. *J Allergy Clin Immun* **134,** 242–243 (2014).

131. Til, N. P. van, Cortes, P., Danos, O., Cassani, B., Poliani, P. L., Villa, A. & Wagemaker, G. Reply. *J Allergy Clin Immun* **134,** 243–244 (2014).

132. Til, N. P. van, Sarwari, R., Visser, T. P., Hauer, J., Lagresle-Peyrou, C., Velden, G. van der, Malshetty, V., Cortes, P., Jollet, A., Danos, O., Cassani, B., Zhang, F., Thrasher, A. J., Fontana, E., Poliani, P. L., Cavazzana, M., Verstegen, M. M. A., Villa, A. & Wagemaker, G. Recombination-activating gene 1 (Rag1)–deficient mice with severe combined immunodeficiency treated with lentiviral gene therapy demonstrate autoimmune Omenn-like syndrome. *J Allergy Clin Immun* **133,** 1116–1123 (2014).

133. Tamari, M., Bel, K. L. D., Heul, A. M. V., Zamidar, L., Orimo, K., Hoshi, M., Trier, A. M., Yano, H., Yang, T.-L., Biggs, C. M., Motomura, K., Shibuya, R., Yu, C. D., Xie, Z., Iriki, H., Wang, Z., Auyeung, K., Damle, G., Demircioglu, D., Gregory, J. K., Hasson, D., Dai, J., Chang, R. B., Morita, H., Matsumoto, K., Jain, S., Dyken, S. V., Milner, J. D., Bogunovic, D., Hu, H., Artis, D., Turvey, S. E. & Kim, B. S. Sensory neurons promote immune homeostasis in the lung. *Cell* **187,** 44-61.e17 (2024).

134. Monticelli, L. A., Sonnenberg, G. F., Abt, M. C., Alenghat, T., Ziegler, C. G. K., Doering, T. A., Angelosanto, J. M., Laidlaw, B. J., Yang, C. Y., Sathaliyawala, T., Kubota, M., Turner, D., Diamond, J. M., Goldrath, A. W., Farber, D. L., Collman, R. G., Wherry, E. J. & Artis, D. Innate lymphoid cells promote lung tissue homeostasis following acute influenza virus infection. *Nat. Immunol.* **12,** 1045–1054 (2011).

135. Brestoff, J. R., Kim, B. S., Saenz, S. A., Stine, R. R., Monticelli, L. A., Sonnenberg, G. F., Thome, J. J., Farber, D. L., Lutfy, K., Seale, P. & Artis, D. Group 2 innate lymphoid cells promote beiging of white adipose tissue and limit obesity. *Nature* **519,** 242–246 (2015).

136. Monticelli, L. A., Osborne, L. C., Noti, M., Tran, S. V., Zaiss, D. M. W. & Artis, D. IL-33 promotes an innate immune pathway of intestinal tissue protection dependent on amphiregulin–EGFR interactions. *Proc National Acad Sci* **112,** 10762–10767 (2015).

137. Mohapatra, A., Dyken, S. J. V., Schneider, C., Nussbaum, J. C., Liang, H.-E. & Locksley, R. M. Group 2 innate lymphoid cells utilize the IRF4-IL-9 module to coordinate epithelial cell maintenance of lung homeostasis. *Mucosal Immunol.* **9,** 275–286 (2016).

138. Turner, J.-E., Morrison, P. J., Wilhelm, C., Wilson, M., Ahlfors, H., Renauld, J.-C., Panzer, U., Helmby, H. & Stockinger, B. IL-9–mediated survival of type 2 innate lymphoid cells promotes damage control in helminth-induced lung inflammation. *J. Exp. Med.* **210,** 2951–2965 (2013).

139. Wang, S., Xia, P., Chen, Y., Qu, Y., Xiong, Z., Ye, B., Du, Y., Tian, Y., Yin, Z., Xu, Z. & Fan, Z. Regulatory Innate Lymphoid Cells Control Innate Intestinal Inflammation. *Cell* **171,** 201-216.e18 (2017).

140. Bando, J. K., Gilfillan, S., Luccia, B. D., Fachi, J. L., Sécca, C., Cella, M. & Colonna, M. ILC2s are the predominant source of intestinal ILC-derived IL-10. *J. Exp. Med.* **217,** e20191520 (2019).

141. Roberts, L. B., Schnoeller, C., Berkachy, R., Darby, M., Pillaye, J., Oudhoff, M. J., Parmar, N., Mackowiak, C., Sedda, D., Quesniaux, V., Ryffel, B., Vaux, R., Gounaris, K., Berrard, S., Withers, D. R., Horsnell, W. G. C. & Selkirk, M. E. Acetylcholine production by group 2 innate lymphoid cells promotes mucosal immunity to helminths. *Sci. Immunol.* **6,** (2021).

142. Mao, K., Baptista, A. P., Tamoutounour, S., Zhuang, L., Bouladoux, N., Martins, A. J., Huang, Y., Gerner, M. Y., Belkaid, Y. & Germain, R. N. Innate and adaptive lymphocytes sequentially shape the gut microbiota and lipid metabolism. *Nature* **554,** 255–259 (2018).

143. Chen, H., Ren, S., Wan, H., Wei, W., Luo, Y. & Cai, M. DUSP1 regulates the
JAK2/STAT3 signaling pathway through targeting miR-21 in cervical cancer cells. *Cell. Mol.*
Biol. **69,** 40–44 (2023).

144. Schatz, D. G. & Swanson, P. C. V(D)J Recombination: Mechanisms of Initiation. *Annu Rev*
Genet **45,** 167–202 (2011).

145. Kuo, T. C. & Schlissel, M. S. Mechanisms controlling expression of the RAG locus during
lymphocyte development. *Curr Opin Immunol* **21,** 173–178 (2009).

146. Desiderio, S. Temporal and spatial regulatory functions of the V(D)J recombinase. *Semin*
Immunol **22,** 362–369 (2010).

147. Moudgil, A., Wilkinson, M. N., Chen, X., He, J., Cammack, A. J., Vasek, M. J., Lagunas,
T., Qi, Z., Lalli, M. A., Guo, C., Morris, S. A., Dougherty, J. D. & Mitra, R. D. Self-Reporting
Transposons Enable Simultaneous Readout of Gene Expression and Transcription Factor
Binding in Single Cells. *Cell* **182,** 992-1008.e21 (2020).

148. Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet-
based single-cell RNA sequencing data. *Gigascience* **9,** giaa151 (2020).

149. Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell
Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8,** 281-291.e9 (2019).

150. Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nusbaum,
C., Myers, R. M., Brown, M., Li, W. & Liu, X. S. Model-based Analysis of ChIP-Seq (MACS).
Genome Biol **9,** R137 (2008).

151. Ntranos, V., Yi, L., Melsted, P. & Pachter, L. A discriminative learning approach to
differential expression analysis for single-cell RNA-seq. *Nat Methods* **16,** 163–166 (2019).

152. Hulsen, T., Vlieg, J. de & Alkema, W. BioVenn – a web application for the comparison and
visualization of biological lists using area-proportional Venn diagrams. *Bmc Genomics* **9,** 488
(2008).

153. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L.,
Fu, X., Liu, S., Bo, X. & Yu, G. clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. *Innovation* **2,** 100141 (2021).

154. Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of
intersecting sets and their properties. *Bioinformatics* **33,** 2938–2940 (2017).

155. Wang, Y., Sarfraz, I., Hong, R., Koga, Y., Alabdullatif, S., Jenkins, D., Akavoor, V., Cao,
X., Bandyadka, S., Leshchyk, A., Faits, T., Khan, M. M., Wang, Z., Johnson, W. E. & Campbell,
J. D. singleCellTK: Comprehensive and Interactive Analysis of Single Cell   RNA-Seq Data.
(2022). at <https://www.camplab.net/sctk/>

156. Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the integration
of genomic datasets with the R/Bioconductor package biomaRt. *Nat Protoc* **4,** 1184–1191
(2009).

157. Rainer, J. EnsDb.Mmusculus.v79: Ensembl based annotation package. (2017).

158. TBD, T. BSgenome.Mmusculus.UCSC.mm10: Full genome sequences for Mus musculus
(UCSC version mm10, based on GRCm38.p6). (2021).

159. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. (2016). at
<https://ggplot2.tidyverse.org>

160. Garnier, Simon, Ross, Noam, Rudis, Robert, Camargo, Pedro, A., Sciaini, Marco, Scherer
& Cédric. viridis - Colorblind-Friendly Color Maps for R. (2021). doi:10.5281/zenodo.4679424

161. Tan, G. & Lenhard, B. TFBSTools: an R/bioconductor package for transcription factor
binding site analysis. *Bioinformatics* **32,** 1555–1556 (2016).

162. Schep, A. motifmatchr: Fast Motif Matching in R. (2022).

163. Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S.
B., Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L.,
Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W.,
Hindson, C. M., Bharadwaj, R., Wong, A., Ness, K. D., Beppu, L. W., Deeg, H. J., McFarland,
C., Loeb, K. R., Valente, W. J., Ericson, N. G., Stevens, E. A., Radich, J. P., Mikkelsen, T. S.,
Hindson, B. J. & Bielas, J. H. Massively parallel digital transcriptional profiling of single cells.
Nat Commun **8,** 14049 (2017).

164. Satpathy, A. T., Granja, J. M., Yost, K. E., Qi, Y., Meschi, F., McDermott, G. P., Olsen, B.
N., Mumbach, M. R., Pierce, S. E., Corces, M. R., Shah, P., Bell, J. C., Jhutty, D., Nemec, C. M.,
Wang, J., Wang, L., Yin, Y., Giresi, P. G., Chang, A. L. S., Zheng, G. X. Y., Greenleaf, W. J. &
Chang, H. Y. Massively parallel single-cell chromatin landscapes of human immune cell
development and intratumoral T cell exhaustion. *Nat Biotechnol* **37,** 925–936 (2019).

1551

Figure 1

Figure S1

Figure S2

A

Steady state model

sdLN

B
ILC2

C
ILC2

D
IL-5⁺

E
IL-13⁺

Figure 2

Figure S3

Figure S4

Figure 3

A RAG fate mapping in lymphoid differentiation

sdLN

B CD4+ T cells

C CD8+ T cells

D B cells

E NK cells

F ILC2s

G Percent tdRFP+

Figure S5

A

Schematic of multiome experiment

B

Ear thickness

Figure 4

Figure S6

Selected ILC2 marker genes

Figure 5

A Fate assignment schematic

RAG fate mapped cells → Isolated nuclei → tdRFP transcripts

RAG^naïve   RAG^exp

B Mapping gene to peak links (GPLs)

Open chromatin

RAG^naïve
RAG^exp

Peaks

Gene X

GPLs

4 GPLs for Gene X in RAG^naïve
2 GPLs for Gene X in RAG^exp

4   2   0   2   4

Regulome for Gene X more active in RAG^naïve cells

C GPLs for ILC2 gene set

n = 235 genes

More links in RAG^naïve

Difference between RAG fate-mapped states (black bars)

More links in RAG^exp

Tox (33)
Jak2 (22)
Gata3 (19)
Il18r1 (16)
Rora (16)
Il1rl1 (15)
Ccr8 (11)
Icos (9)

Fosb (6)
Ndfip1 (6)
Asxl1 (11)
Junb (11)
Dusp1 (20)

80   40   0   40   80
Number of GPLs

A

ILC2 DEGs by RAG fate map

B

GSEA of RAG^exp vs RAG^naïve ILC2 upregulated genes

C

GO BP: positive regulation of immune system process

Figure S9

Figure S10

A

chromVAR TF motif
enrichment from snATAC-seq

B

Top 5 cluster
chromVAR enriched TF motifs

C

Cell type-specific TFs

D

Top enriched TF motifs in
ILC2 GPLs by RAG fate map

E

Expression of top TFs
from ILC2 GPLs

Figure 6

A Schematic of GPL assignment by RAG fate map and disease

B Intersection of ILC2 GPL peaks by RAG fate map and disease

C Top genes with AD-like disease-induced peaks

D *Rora*

E *Ccr6*

Figure S11

Intersection of all GPL peaks by RAG fate map and disease

A

Th2 locus in ILC2s

Open chromatin (0-11)

RAG^naïve

RAG^exp

Il4   *Il13*   *Rad50*   *Il5*

Peaks

GPLs

5kb

B

Th2 locus GPLs

RAG^naïve   RAG^exp

Il13 (26)
Il4 (23)
Il5 (19)
Rad50 (5)

50 25 0 25 50
Number of GPLs

C

Intersection of Th2 locus GPL peaks by RAG fate map and disease

Intersection size

25 19 17 16 14 10 9 8 8 4 3 2 1 1

exp SS
exp AD
naïve SS
naïve AD

80 40 0
Set size

D

Top Th2 peaks induced in AD-like inflammation

Il13 (7)
Il4 (6)
Il5 (5)
Rad50 (5)

0 4 8

E

Il13

Steady state

AD-like disease

RAG^naïve   RAG^exp

500 bp

A

Top enriched TF motifs in Th2 locus
GPLs by RAG fate map

RORC
GATA4
GATA5
HOXC8
GATA2
GATA3

none
>1.5 cutoff

Enriched in RAGnaïve   3   2   1   0   Enriched in RAGexp

-log$_{10}$(FDR)

B

Expression of top TFs
from ILC2 GPLs

Gata3
Gata2 --- N.D. ---
Hxc8 --- N.D. ---
Gata5 --- N.D. ---
Gata4
Rorc

RAGnaïve   RAGexp

Avg.
expression
z-score

0.4
0
-0.4

Percent
expressed

100
75
50
25
0