1 **Title:** Transcriptome analysis reveals role for WRKY70 in early *N*-hydroxy-pipecolic acid
2 signaling
3
4

5 **Authors:** Jessica Foret[1], Jung-Gun Kim[1], Elizabeth S. Sattely[2,3] and Mary Beth Mudgett[1,*]
6
7
8 **Affiliations:**
9 [1]Department of Biology, Stanford University, Stanford, CA 94305, USA
10 [2]Department of Chemical Engineering, Stanford University, Stanford CA 94305, USA
11 [3]Howard Hughes Medical Institute, Stanford University, Stanford CA, 94305, USA
12
13
14 ***Corresponding Author:** Professor Mary Beth Mudgett (mudgett@stanford.edu);
15 Stanford University, 371 Jane Stanford Way, 228A Gilbert Biosciences, Stanford, CA
16 94305-5020

17
18
19 **Short running head:** Early transcriptional changes in response to NHP
20
21

Abstract   (250 words)

N-hydroxy-pipecolic acid (NHP) is a mobile metabolite essential for inducing and
amplifying systemic acquired resistance (SAR) following pathogen attack. Early phases
of NHP signaling leading to immunity have remained elusive. Here we report the early
transcriptional changes mediated by NHP and the role salicylic acid (SA) plays during this
response. We show that distinct waves of expression within minutes to hours of NHP
treatment include increased expression of WRKY transcription factors as the primary
transcriptional response, followed by the induction of WRKY-regulated defense genes as
the secondary response. The majority of genes induced by NHP within minutes were SA-
dependent, whereas those induced within hours were SA-independent. These data
suggest that NHP induces the primary transcriptional response in a low SA environment
and new SA biosynthesis is dispensable for induction of the secondary transcriptional
response. We demonstrate that WRKY70 is required for the induced expression of a set
of genes defining some of the secondary transcriptional response, SAR protection, and
NHP-dependent enhancement of ROS production in response to flagellin treatment.
Taken together, our study highlights the key genes and pathways defining early NHP
responses and a role for WRKY70 in the regulation of NHP-dependent transcription.

Introduction

Systemic acquired resistance (SAR) is a whole-plant defense response triggered by an initial pathogen attack. During the establishment of SAR, mobile signals are sent from infected tissue to uninfected tissue, initiating the priming of immune responses and enhancing immunity against secondary pathogen infections (Fu and Dong, 2013; Tian and Zhang, 2019; Kachroo and Kachroo, 2020). To achieve a primed state, plants execute substantial transcriptional reprogramming of several hundred to a few thousand genes, resulting in the increased accumulation of defense hormone salicylic acid (SA), sensing and signaling components for pathogen detection, and production of antimicrobial compounds (Jaskiewicz et al., 2011; Conrath et al., 2015; Bernsdorff et al., 2016). In addition, several small molecules produced by plants during infection help to orchestrate this defense priming, including SA and the bioactive metabolite *N*-hydroxy-pipecolic acid (NHP) (Chen et al., 2018; Hartmann et al., 2018; Klessig et al., 2018).

NHP is a mobile metabolite that can move from one leaf to another to mediate long-distance cellular communication (Chen et al., 2018; Mohnike et al., 2021; Yildiz et al., 2021). In *Arabidopsis thaliana* (Arabidopsis), the enzyme FLAVIN-DEPENDENT MONOOXYGENASE 1 (FMO1) catalyzes the *N*-hydroxylation of lysine-derived pipecolic acid (Pip) in response to pathogen infection to produce NHP (Chen et al., 2018; Hartmann et al., 2018). Notably, *fmo1* null mutants are impaired in local disease resistance during pathogen infection and fail to establish SAR (Chen et al., 2018; Hartmann et al., 2018). Treatment of *fmo1* mutant plants with exogenous NHP is sufficient to induce systemic resistance against bacterial and oomycete pathogens (Chen et al., 2018; Hartmann et al., 2018). Furthermore, NHP treatment of a single leaf is sufficient to initiate and amplify defense signaling throughout the plant, resulting in the transcriptional activation of a large subset of SAR-associated genes within 24 h of treatment (Yildiz et al., 2021; Yildiz et al., 2023), including genes encoding the SA biosynthetic enzyme ISOCHORISMATE SYNTHASE 1/SA-INDUCTION DEFICIENT 2 (ICS1/SID2), the Pip biosynthetic enzymes AGD2-LIKE DEFENSE RESPONSE PROTEIN 1 (ALD1) and SAR-DEFICIENT 4 (SARD4), as well as its own biosynthetic

enzyme FMO1 (Chen et al., 2018). Collectively, these findings highlight the importance of

NHP in initiating immune signaling to establish local and systemic plant defenses.

Induction of *ICS1* transcription by NHP signaling results in the accumulation of SA in both

local and systemic tissues (Chen et al., 2018; Hartmann et al., 2018). Elevation of SA in

turn amplifies SAR-associated transcriptional changes (Bernsdorff et al., 2016). Signaling

downstream of SA accumulation is largely driven by the action of the SA receptor and

transcriptional coactivator NONEXPRESSER OF PR GENES 1 (NPR1), which interacts

with a number of TGA transcription factors (TFs) from the basic leucine zipper (bZIP)

family to alter gene expression (Zhang et al., 1999; Wang et al., 2006; Wu et al., 2012).

Members of the biotic and abiotic stress-response WRKY TF family are among the known

targets of the NPR1/TGA interaction (Thibaud-Nissen et al., 2006; Wang et al., 2006).

WRKY TFs have been identified as key regulators of NPR1-dependent and NPR1-

independent SAR responses. Additionally, some WRKY members play important roles in

feedback regulation of transcription (Pandey and Somssich, 2009; Fu and Dong, 2013).

For example, WRKY54 and WRKY70 modulate SA biosynthesis through repression of

ICS1 expression (Wang et al., 2006).

The interplay between SA and NHP appears to be interdependent and synergistic with

both metabolites being able to upregulate one another's biosynthetic pathways (Chen et

al., 2018; Hartmann et al., 2018; Sun et al., 2020). Notably, elevated endogenous levels

of Pip, the precursor of NHP, primes the expression of both NHP and SA biosynthetic

genes under low concentrations of SA by stabilizing NPR1 levels (Kim et al., 2020). This

points to an intriguing dynamic relationship between SA and NHP over the course of SAR

activation. Evidence suggests NHP accumulates to detectable levels in systemic tissues

roughly 24 h before SA accumulation is detected (Hartmann and Zeier, 2019). In such a

case, NHP would need to drive systemic signaling for a substantial period of time before

SA could further amplify SAR responses. It is currently unclear how the levels of SA in

leaves impact early NHP signaling. Furthermore, little is known about the very early phase

of NHP signaling in the tissues where it is first produced or in the distal tissues once NHP

arrives there. Key players of NHP signaling and the progression of NHP-dependent
transcriptional reprogramming require further investigation.

Presently, the majority of genome-wide transcriptomic studies of SAR have been carried
out on the timescale of 1-2 days post induction; this includes pharmacological studies
using Pip and NHP treatment (Bernsdorff et al., 2016; Hartmann et al., 2018; Yildiz et al.,
2021; Yildiz et al., 2023). These data have revealed that NHP ultimately activates SAR in
untreated tissue, but do not shed light on early signaling events that lead to the observed
SAR phenotype. The immediate consequences of NHP accumulation on gene expression
have not been described. Studying very early transcriptional changes, within minutes to
hours of elicitor treatment, has been an effective strategy elucidating key components of
signaling pathways for SA, jasmonic acid (JA), and brassinosteroid (Friedrichsen et al.,
2002; Chini et al., 2007; Ding et al., 2018). Thus, in an effort to better understand NHP's
primary action in signal transduction, we determined the genome-wide transcriptional
responses in Arabidopsis from 15 minutes up to 6 hours after treatment with exogenous
NHP.

Here we report the early NHP transcriptional profile and the role SA plays during this
response. We present evidence indicating that increased expression of WRKY TFs define
the primary transcriptional responses mediated by NHP within minutes (i.e. 15 to 30
minutes) post-treatment, and WRKY TFs act as executors of the secondary
transcriptional responses within hours (i.e. 3 to 6 hours). The majority of genes induced
by NHP within minutes were SA-dependent, whereas those induced within hours were
SA-independent. These data suggest that NHP induces the primary transcriptional
response in a low SA environment and new SA biosynthesis is dispensable for induction
of the secondary transcriptional response. We also show that WRKY70 is required for
NHP induced transcription of a set of genes defining secondary transcriptional changes
as well as NHP enhancement of reactive oxygen species (ROS) and SAR. Taken
together, our study highlights the key genes and pathways defining early NHP responses
and a role for WRKY70 in the regulation of NHP-dependent transcription.

Results

NHP induces expression of defense-associated genes within minutes

Previous studies in Arabidopsis have demonstrated that exogenous application of NHP to a leaf (defined as "local tissue") is sufficient to initiate and amplify defense signaling in untreated leaves (defined as "distal tissue"), including the transcriptional activation of SAR marker genes (Chen et al., 2018; Hartmann et al., 2018; Nair et al., 2021; Yildiz et al., 2021). Notably, these studies focused on gene expression changes in distal tissues at late time points (24 to 48 h) following NHP elicitation. However, it was unknown which gene expression changes occur immediately in response to NHP treatment. We hypothesized that once present in the tissue, either via endogenous synthesis or exogenous application, NHP initiates signaling that leads to transcriptional changes on a timescale of minutes to several hours, which are essential for execution of NHP-dependent immune responses.

In order to test this hypothesis, we set out to identify NHP responsive transcriptional markers and determine the earliest time frame when NHP could induce these genes. We first questioned if NHP could induce the expression of classical SAR-associated marker genes (i.e., *FMO1*, *ICS1*, and *PATHOGENESIS-RELATED PROTEIN 1* (*PR1*)) within several hours of treatment. Studies have shown *FMO1*, *ICS1*, and *PR1* are induced between 8 to 48 h following NHP treatment (Chen et al., 2018; Nair et al., 2021). Therefore, we began by testing the response to NHP within 24 h of treatment. The leaves of 4.5-week-old Arabidopsis Col-0 (wild type) plants were infiltrated with 1 mM NHP and then treated leaves were collected for mRNA isolation at 2, 6 and 24 h. Quantitative real-time PCR (qRT-PCR) analysis showed significantly increased transcript abundance of all three genes in NHP treated leaves at 24 h but not at 2 and 6 h compared to water (mock) treated leaves (Fig. 1A). These data indicate that *FMO1*, *ICS1,* and *PR1* are responsive to NHP treatment but are likely not primary targets of NHP transcriptional activation.

Next, we tested the ability of NHP to induce expression of *WRKY38* and *UGT76B1*, two genes known to respond to SA within 2 h of treatment (Blanco et al., 2009). *WRKY38*

encodes a defense-associated TF and *UGT76B1* encodes a UDP-dependent
glycosyltransferase shown to glycosylate SA and NHP (Eulgem et al., 2000; Bauer et al.,
2021; Cai et al., 2021; Holmes et al., 2021; Mohnike et al., 2021). We found a significant
increase of mRNA abundance for both *WRKY38* and *UGT76B1* in NHP treated leaves at
2 h compared to mock treated leaves (Fig. 1A). *WRKY38* transcript levels were also
elevated at 6 and 24 h after treatment with NHP (Fig. 1A). These data indicate that both
WRKY38 and *UGT76B1* are early NHP responsive transcripts.
178
We used the *WRKY38* gene as a marker to define the earliest time when NHP-induced
transcriptional changes could be measured. To this end, we implemented a
hydroponically grown seedling assay to synchronize chemical treatment and increase the
number of plants sampled, while minimizing mechanical stress. Wild type seedlings were
grown in liquid medium in six-well plates with 15-18 seeds per well. The seedlings were
treated with mock, 1 mM Pip, or 1 mM NHP at 10-days-post germination and then
collected at 15 min, 30 min, 3 h, and 6 h post treatment to isolate mRNA (Fig. 1B).
WRKY38 transcript abundance was significantly elevated in Arabidopsis seedlings
treated with NHP as early as 15 min, peaking in abundance at 30 min, when compared
to those treated with mock or Pip (Fig. 1C). This trend continued through 3 h following
NHP treatment. The NHP precursor Pip did not elicit an increase in *WRKY38* mRNA
within 3 h but showed a slight trend of increased abundance at the 6 h time point (Fig.
1C). These results indicate that exogenous NHP, but not Pip, is sufficient to elevate
WRKY38 transcript abundance within 15 minutes of application.
193
Since *WRKY38* expression is known to be induced by SA treatment (Blanco et al., 2009),
we questioned if NHP-induced expression of *WRKY38* is dependent on SA derived from
the ICS1/SID2 biosynthetic pathway by utilizing the well characterized *sid2-2* mutant that
does not accumulate SA upon pathogen infection (Wildermuth et al., 2001). Wild type and
sid2-2 seedlings were grown hydroponically, treated with mock, 1 mM Pip, or 1 mM NHP,
and then collected 3 h after treatment. NHP treatment increased *WRKY38* mRNA
abundance in the *sid2-2* mutant relative to mock treatment, similar to wild type seedlings
(Supplemental Fig. S1). These findings indicate that NHP is sufficient to alter *WRKY38*

transcript abundance in the absence of SA accumulation, defining *WRKY38* as an early

NHP induced, SA-independent transcript.

The early NHP-responsive transcriptome reveals distinct waves of expression

We next questioned how the presence of NHP impacts whole-genome transcriptional

changes within minutes to hours of treatment in a SA-dependent and -independent

manner. To address this, Arabidopsis wild type and *sid2-2* mutant seedlings were grown

in the same hydroponic design as previously described for qRT-PCR experiments (Fig.

1B). The *sid2-2* mutant was analyzed in order to identify transcriptional changes

dependent on NHP-induced SA biosynthesis (Chen et al., 2018). Each pool of seedlings

was treated with mock or 0.5 mM NHP for 15 min, 30 min, 3 h and 6 h. A reduced

concentration of NHP was used as we found that 0.5 mM NHP induced similar *WRKY38*

mRNA accumulation as 1 mM NHP (Supplemental Fig. S2). The abundance of mRNA in

the treated seedlings was assessed for each condition via RNA-sequencing (RNA-seq).

After general quality control and mapping to the Arabidopsis TAIR10 genome, genes

differentially expressed by NHP were determined relative to mock treatment for each time

point and genotype using DESeq2 (Love et al., 2014). Genes were considered

differentially expressed if they returned an adjusted *P*-value below 0.05 ($P_{adj} < 0.05$).

In wild type seedlings, a total of 2,079 genes were differentially expressed in response to

NHP for at least one of the measured time points. To explore the set of early NHP-

responsive genes, we first selected for genes showing a robust fold change over mock

by applying the cutoff of $\log_2$(fold change, FC) > 1 for upregulated genes and $\log_2$(FC) <

-1 for downregulated genes. This identified 352 genes that were differentially expressed

in response to NHP for at least one time point, with 163 genes up and 189 genes down

(Supplemental Table S3). Next, the upregulated genes were hierarchically clustered

based on their $\log_2$(FC) values over the indicated time points and four well-defined gene

clusters were identified (Fig. 2A). For each group of genes in the defined clusters, the

$\log_2$(FC) values were then averaged by time point, revealing dynamic patterns of

expression. Two of the clusters were upregulated within minutes (designated as 'early

transient' and 'early strong') and two were upregulated by several hours (designated as
'late weak' and 'late strong') (Fig. 2B, Table 1, Table 2, Supplemental Table S4).

The 'early transient' cluster includes 42 genes and is defined by elevated transcript
abundance 15-30 min after NHP treatment that did not persist by 3 h. Gene ontology (GO)
analysis for this cluster showed strong enrichment for biological processes associated
with responses to jasmonic acid (JA), wounding, and oxidative stress. Some of these
genes are associated with the regulation of defense as well as defense responses to fungi
and bacteria (Fig. 2C). Among the most highly expressed genes in this cluster are JA-
responsive genes, which play key roles in the regulation of wounding responses, including
CML39 which encodes a calmodulin-like protein, *JAZ10* which encodes a jasmonate-zim-
domain protein, and *MYC2* which encodes a jasmonate-inducible transcription factor
(Table 1, Table 2). Taken together, the early yet unsustained induction of these genes
within 30 min suggests a burst of JA- and wound-responsive gene expression that acts
as a primary wave of NHP signaling.

The 'early strong' cluster includes 10 genes and is defined by a high average $\log_2$(FC) 15
min after NHP treatment that was maintained for all subsequent time points. Of the 10
genes, two (AT1G06475 and AT5G40980) were strongly upregulated by NHP at 30 min
and no other time points. These two genes were removed from the average $\log_2$(FC)
analysis and subsequent analyses because they did not behave like the other genes in
this cluster (Fig. 2B). The early strong cluster is notably enriched in biological processes
involved in the regulation of transcription, with over half the genes belonging to the WRKY
TF family, including four group III *WRKY* genes (*WRKY38*, *WRKY54*, *WRKY62* and
WRKY70) and two group II-c genes (*WRKY51* and *WRKY59*) (Eulgem et al., 2000; Kalde
et al., 2003); (Table 1, Table2, Supplemental Table S4). Additional GO terms enriched in
the early strong cluster includes regulation of defense and regulation of hormonal
signaling pathways such as SA, JA, brassinosteroid, and ethylene (Fig. 2C). The early
and sustained induction of these genes, particularly the *WRKYs*, suggests they are
primary transcriptional target genes of NHP signaling.

The 'late weak' cluster includes the largest number of genes (96) that averaged a $\log_2$(FC)
around 1 (Fig. 2B). GO term analysis of the cluster revealed significant enrichment of
biological processes involved in response to bacteria and oomycetes, as well as SA-
mediated signaling and regulation of SAR (Fig. 2C). This indicates NHP activation of SAR
and pathogen defense genes can be detected as early as 3-6 h in treated tissues.
Notably, genes involved in the regulation and biosynthesis of NHP were observed in this
wave of transcription, including *SARD1* (*SAR DEFICIENT 1*), *NIMIN-1* (*NIM-*
INTERACTING 1), *ALD1,* and *UGT76B1* (Table 1, Table 2, Supplemental Table S4)
(Song et al., 2004; Zhang et al., 2010a; Holmes et al., 2021).

The 'late strong' cluster consists of 15 genes with $\log_2$(FC) values averaging between 3.5
- 4 (Fig. 2B). Response to SA was the only significantly enriched GO term for this cluster
(Fig. 2C). Despite having a role in the response to SA, the majority of genes classified as
late strong were upregulated by NHP in both wild type and *sid2-2* (Fig. 3A, Table 1),
indicating that accumulation of these transcripts was at least partially SA-independent. Of
the 14 SA-independent genes identified in this cluster, only five have been found to be
upregulated 24 h after NHP treatment in an SA-independent manner (Yildiz et al., 2021).
Thus, our analysis has uncovered additional NHP-responsive genes that operate
independently of SA as early as 6 h post NHP signaling.

We also performed a similar clustering analysis on the 189 NHP-downregulated
transcripts and identified three gene clusters (Supplemental Fig. S3A, B). In contrast to
the upregulated gene clusters, the downregulated gene clusters had less distinct patterns
of expression and averaged $\log_2$(FC) values around -1 to -1.5. Two clusters were
downregulated within 15-30 min and persisted 3-6 h hours after NHP treatment. One of
these showed weaker downregulation and was classified as an 'early weak' cluster. The
other showed stronger downregulation and was classified as an 'early strong' cluster. The
third did not show decreased transcript abundance until 3-6 h after treatment and thus
was labeled as a 'late' cluster (Supplemental Fig. S3B). All three clusters were enriched
for biological processes involving growth and development such as root morphogenesis,
plant epidermis development, plant-type cell wall organization, and root hair cell

differentiation (Supplemental Fig. S3C). Given the NHP precursor Pip is known to inhibit

root growth of seedlings (Wang et al., 2018), it is likely that NHP also inhibits root growth.

These clusters shed new light on the genes that may be associated with altered plant

growth and development in response to elevated levels of NHP.

SA biosynthesis is dispensable for the early induction of a majority of NHP

upregulated genes

Interrogation into the late strong cluster suggested NHP may be sufficient to activate a

subset of SA-responsive genes in the absence of NHP-induced SA biosynthesis (Fig. 2C,

Fig. 3A, 'late strong'). This was an intriguing possibility given NHP likely accumulates in

distal tissues prior to SA accumulation following a local tissue infection (Bernsdorff et al.,

2016; Hartmann et al., 2018; Hartmann and Zeier, 2019). We therefore hypothesized that

NHP would need to drive transcriptional reprogramming as SA levels rise from low to

high. To investigate this, we assessed the impact of SA biosynthesis on NHP-induced

gene expression from 15 min to 6 h. For each time point of the wild type gene clusters

described in Fig. 2, the percentage of genes upregulated in both wild type and the *sid2-2*

mutant was determined. This group of genes was defined as 'SA-independent'. Genes

upregulated only in wild type and not in *sid2-2* at the defined time point were conversely

defined as 'SA-dependent'.

Notably, the majority of genes upregulated by NHP at 15 min in the early transient cluster

were SA-dependent. For example, 87% of the genes (27 out of 31) were upregulated at

15 min in wild type but not in *sid2-2* seedlings (Fig. 3A) and similar trends were seen at

30 min. By contrast, 100% of the genes (8 out of 8) in the early strong cluster upregulated

3-6 h after NHP treatment were SA-independent (Fig. 3A). 69% of the genes (36 out of

52) in the late weak cluster were SA-independent at 3h, which dropped to 53% (37 out of

70 genes) at 6 h (Fig. 3A).

A similar pattern of gene expression was observed when we analyzed the total number

of NHP upregulated genes reaching a $\log_2(FC) > 1$ without gene clustering (Fig. 3B). The

majority of genes induced by NHP between 15-30 min were SA-dependent (i.e., 36 genes

up in wild type and 9 up in both wild type and *sid2-2* at 15 min) and the majority of genes

induced by NHP between 3-6 h were SA-independent (i.e., 20 genes up in wild type and

55 up in both wild type and *sid2-2* at 3 h) (Fig. 3B). Taken together, this analysis highlights

an important temporal role of SA on NHP induced transcription. Our data suggest that

new SA biosynthesis is required for the induction of NHP-responsive genes within

minutes of exposure to NHP, whereas SA biosynthesis is dispensable for NHP-induced

expression several hours later.

We also analyzed the downregulated genes using the same approach and found a slight

trend of SA-dependence for these genes following NHP treatment. For example, the early

weak and late clusters defined in Supplemental Fig. S3 showed that a majority of genes

are downregulated in wild type but not the *sid2-2* mutant at each time point (Supplemental

Fig. S4A). By contrast, the genes of the early strong cluster were less dependent on SA

biosynthesis, such that by the 6 h time point only 15% of genes (8 out of 44) were

downregulated in wild type and not in *sid2-2* seedlings (Supplemental Fig. S4A).

Independent of clustering, greater than 60% of the downregulated genes were found to

be SA-dependent at 15 min and 3 h and approximately 40% were SA-dependent at 30

min and 6 h (Supplemental Fig. S4B). Taken together, these findings suggest a stronger

dependence on SA biosynthesis for NHP-dependent downregulation of gene expression

over the time courses studied.

SA biosynthesis antagonizes NHP-elicited gene expression at 3 to 6 h

Looking at the genes upregulated only in the *sid2-2* mutant and not wild type, we

discovered an unexpectedly large number of transcripts that increased in abundance

upon NHP treatment at 3 and 6 h post treatment (Fig. 3B). This includes 151 genes at 3

h and 98 genes at 6 h that were significantly upregulated by NHP in the *sid2-2* mutant but

not in wild type (Fig. 3B). GO-term enrichment analysis of these genes showed significant

enrichment for a number of biotic and abiotic stress responses, such as defense response

to bacteria and fungi, SAR, response to abscisic acid, and response to hypoxia (Fig. 3C).

These data suggest that SA may play an important role in modulating the transcription of

a suite of genes, through an unknown mechanism, once NHP-mediated transcriptional
reprogramming has been initiated.

TGA and WRKY transcription factor *cis*-regulatory elements are enriched in the
358
promoters of early NHP upregulated genes

To identify TFs that might directly control the expression of early NHP-responsive genes,
we analyzed the promoter regions of all genes upregulated after NHP treatment in both
wild type and *sid2-2* mutant seedlings to identify putative TF bindings sites (i.e., *cis*-
regulatory elements, CREs). A less stringent cutoff of $\log_2(FC) > 0$ and $P_{adj} < 0.05$ was
used for the upregulated genes to capture putative transcriptional regulators of all genes
significantly elevated in transcript abundance, regardless of the magnitude of expression
change. Following a modified workflow detailed by Bjornson et al., 2021, the NHP
upregulated genes were then separated by the time points in which they were
differentially expressed. Enrichment of transcription factor CREs was then determined for
each time point using the DAPseq database (O'Malley et al., 2016) (Fig. 4A).
Supplemental Fig. 5 shows all the CREs found to be significantly enriched in the
promoters of NHP-upregulated genes in both wild type and *sid2-2* seedings between 15
min and 6 h following NHP treatment.

We found that CREs for bZIP family members, specifically TGA TFs, were enriched in the
promoters of genes upregulated by NHP as early as 15 min. For some TGAs (e.g., TGA2,
TGA5, and TGA7), the enrichment of their TF binding sites was observed for all time
points (Fig. 4B). Enrichment of TGA binding sites in the promoters of early NHP
upregulated genes is consistent with previous work demonstrating TGA2/5/6 and TGA1/4
are required for the induction of NHP responsive genes at 24-48 h post NHP treatment
(Nair et al., 2021; Yildiz et al., 2023). TGA TFs are known to interact with the
transcriptional coactivator NPR1 to drive SAR- and SA-induced transcriptional changes
(Zhang et al., 1999). The enrichment of TGA binding sites in the promoters of NHP
upregulated genes as early as 15 min implicates TGAs and NPR1 among the primary
transcriptional regulators of NHP signaling. Furthermore, enrichment of TGA CREs was
not observed in *sid2-2* seedlings until 30 min and was also not significantly enriched at 3

h (Fig. 4B). These findings suggest that SA biosynthesis and likely elevated SA levels are

required for rapid and stable induction of some TGA-regulated genes induced by NHP

signaling.

Moreover, we discovered that a large number of WRKY CREs were highly enriched in the

promoter regions of NHP upregulated genes at 3-6 h in both wild type and *sid2-2*

seedlings (Fig. 4C). This correlates with the majority of genes identified in the NHP

upregulated early strong cluster belonging to the WRKY TF family (Fig. 2, Table 1). This

finding suggests a model where NHP induces the expression of *WRKY* genes as a

primary transcriptional response, followed by WRKY-regulated gene expression changes

that act as a secondary wave of transcriptional control.

It is striking that the confidence in WRKY CRE enrichment was greater (i.e., lower *P* value)

at 3 and 6 h in the promoters of genes upregulated in the *sid2-2* mutant compared to

those of wild type (Fig. 4C). These data suggest that a substantial number of NHP

upregulated genes are directly regulated by WRKY TFs, and such regulation is influenced

by SA levels. A similar pattern emerged for members of the NAC TF family, where NAC

CREs were enriched in the promoters of NHP upregulated genes at 3 h in *sid2-2* but not

in the promoters of genes upregulated in wild type (Supplemental Fig. S5). Both WRKY

and NAC TFs are involved in the regulation of a number of biotic and abiotic stress

responses (Eulgem et al., 2000; Mohanta et al., 2020). Notably, the set of genes

upregulated by NHP at 3-6 h only in the *sid2-2* mutant but not in wild type were enriched

in biological processes associated with biotic and abiotic stress responses (Fig. 3B, C).

Taken together, these data suggest SA may antagonize the transcription of a subset of

NHP responsive genes by regulating the activities of WRKY and NAC TFs.

The same CRE enrichment analysis was also carried out on all NHP downregulated

genes, regardless of the magnitude of fold change ($\log_2$(FC) < 0 and P_{adj} < 0.05). We

found multiple WRKY CREs enriched in the promoter regions of NHP downregulated

genes at 15 min, 3 h, and 6 h, in wild type seedlings (Supplemental Fig. S6), further

supporting a role for WRKY TFs in relaying NHP-induced transcriptional changes.

Finally, we found the CREs of Homeobox gene family members to be abundant in the

promoters of NHP downregulated genes spanning from 15 min through 6 h, with the

greatest confidence of enrichment found in the promoters of genes downregulated by

NHP at 3-6 h in an SA-dependent manner (Supplemental Fig. S6). Among these was

ATHB5, a homeodomain leucine zipper (HDZip) protein linked to the abscisic acid (ABA)-

driven repression of germination and root growth in seedlings (Johannesson et al., 2003).

Given that NHP precursor Pip inhibits seedling root growth, we now hypothesize that NHP

may drive the repression of root growth and development genes (Supplemental Fig. 3C)

through the action of HDZip Homeobox proteins like ATHB5.

Early NHP-responsive gene *WRKY70* is required for NHP-elicited SAR

To further dissect the NHP response pathway, we carried out a reverse genetic screen to

identify Arabidopsis mutants with compromised NHP-elicited SAR. Given WRKY CREs

were the predominant regulatory elements identified in genes responsive to NHP 3-6 h

following treatment (Fig. 4C, Supplemental Fig. S6), we selected the six *WRKY*s

(*WRKY38, WRKY51, WRKY54, WRKY59, WRKY62, WRKY70)* from the early strong

NHP upregulated cluster (Figure 2B, Table 1, Table 2) for mutant analysis.

To measure NHP-induced SAR, we obtained homozygous Arabidopsis mutants and

quantified bacterial growth in their leaves after treatment with exogenous NHP (Chen et

al., 2018). Specifically, three lower leaves of 4.5-week-old wild type and mutant plants

were infiltrated with water (mock) or 0.5 mM NHP. One day later, one distal upper leaf

was inoculated with a 1 x 10^5 CFU/mL suspension of the virulent bacterium *Pseudomonas*

syringae pathovar *maculicola* strain ES4326 (*Psm*). Bacterial growth of *Psm* was

quantified three days post infection (dpi) to determine if the mutants were altered in their

resistance to pathogen infection. Mutants were considered to exhibit NHP-elicited SAR if

bacterial titer was significantly lower in NHP treated plants compared to mock treated

plants. By contrast, mutants were considered insensitive or partially insensitive to NHP if

the bacterial titer of NHP treated mutants was higher than NHP treated wild type plants.

Of the six *wrky* mutants screened, five (*wrky38*, *wrky62*, *wrky54*, *wrky51*, and *wrky59*)
exhibited normal NHP-elicited SAR (Fig. 5B-E). Only one, *wrky70*-1 (T-DNA insertional
line SALK_025198) (Li et al., 2006), displayed compromised NHP-elicited SAR (Fig. 5A).
That is, NHP treatment did not decrease bacterial growth in distal leaves of *wrky70-1*
plants compared to mock treatment. We also observed that *wrky70-1* mutants treated
with mock contained less bacteria than wild type plants treated with mock, demonstrating
that basal resistance is enhanced in the *wrk70-1* mutant (Fig. 5A). Notably, distal leaves
of *wrky70-1* plants treated with NHP contained higher levels of bacteria than distal leaves
of wild type plants treated with NHP (Fig. 5A). The inability of *wrky70-1* plants to further
restrict bacterial growth following NHP treatment suggests that the mutant may be
insensitive to NHP or defective in NHP signaling. Taken together, these data indicate that
WRKY70 is required for full NHP-elicited systemic resistance.

We next questioned if systemic resistance to *Psm* was similarly compromised in *wrky70-
1* plants primed with the avirulent pathogen *P. syringae* pathovar *tomato* (*Pst*) strain
DC3000 carrying *avrRpt2* (*Pst avrRpt2*). Localized infection with *Pst avrRpt2* is known to
elicit a strong immune response that leads to SAR and protection against *Psm* in distal
leaves (Kohler et al., 2002). We found that the *Psm* titer in distal leaves reached similar
levels for *wrk70-1* plants treated with mock or *Pst avrRpt2* and this level was comparable
to *Psm* growth in wild type leaves of plants treated with *Pst avrRpt2* (Supplemental Fig.
S7A). These data indicate that *wrky70-1* mutants exhibit a basal-level of defense priming
and *Pst avrRpt2* primary infection does not further enhance this resistance.

We also examined pathogen growth in local *wrky70-1* leaves. We found that the titer of
Psm in infected *wrky70-1* leaves was lower than in *Psm* infected wild type leaves
(Supplemental Fig. S7B). This phenotype corroborates previous work demonstrating loss
of *WRKY70* expression enhances resistance to *Psm* infection (Zhou et al., 2018).

Expression of *SARD1* and *PR* genes is elevated in *wrky70-1* independent of NHP
Previous studies in Arabidopsis have demonstrated a role for WRKY70 in the
transcriptional regulation of several genes required for SAR signaling and disease

resistance, including *SARD1* and the *PATHOGENESIS-RELATED* genes *PR1*, *PR2*, and
PR5 (Li et al., 2004; Zhou et al., 2018; Liu et al., 2021). These genes were also shown to
be upregulated in wild type Arabidopsis plants 24 h after treatment with NHP (Yildiz et al.,
2021). To determine if WRKY70 is required for NHP induced expression of *SARD1*, *PR1*,
PR2, and *PR5* at 24 h, we quantified transcript abundance in wild type and *wrky70-1*
plants treated with water or 1 mM NHP. As expected, qRT-PCR showed increased
abundance of *SARD1*, *PR1*, *PR2*, and *PR5* mRNA in wild type plants treated with NHP
compared to water (Supplemental Fig. S7C). However, NHP treatment did not further
increase transcript levels in *wrky70-1* plants (Supplemental Fig. S7C). Notably, all four
genes showed elevated transcript abundance in water treated *wrky70-1* mutants
compared to water treated wild type. The elevated expression of *SARD1* and *PR1* in
water treated *wrky70-1* plants is consistent with previous studies demonstrating WRKY70
negatively regulates these two genes in the absence of a pathogen (Zhou et al., 2018; Liu
et al., 2021).
493
WRKY70 is required for NHP induction of genes in the late weak and late strong
clusters
Given *wrky70-1* mutants exhibited enhanced basal resistance but were still unable to
achieve the same level of resistance as NHP treated wild type plants (Figure 5A), we
hypothesized that a subset of early NHP-responsive genes containing putative WRKY70
CREs in their promoters may be dependent on WRKY70 function for full, wild type gene
expression levels following NHP treatment. To investigate this, we selected three genes
from the late weak and late strong clusters (Fig. 2, Table 1), specifically *BDA1*,
PROSCOOP4, and *CML10,* and examined their transcript accumulation in response to 1
mM NHP over a time course of 0 to 12 h. The promoters of these genes contain one or
more WRKY70 CREs (i.e., W-box (TTGACY), WT-box (YGACTTTT), and WRKY70
DAPseq motif; (Rushton et al., 2010; Machens et al., 2014; O'Malley et al., 2016)) within
2000 base pairs upstream of their respective putative transcriptional start site (Fig. 6A).
507
We found that *BDA1* (*bian da*; "becoming big" in Chinese), a gene encoding an ankyrin-
repeat transmembrane protein (Yang et al., 2012), exhibited the strongest dependence

on WRKY70 for NHP-induced expression. *BDA1* transcripts were significantly increased
at 6, 9, and 12 h post NHP treatment in wild type leaves but not *wrk70-1* leaves (Fig. 6B).
Transcript levels for *PROSCOOP4*, a gene encoding the precursor of secreted peptide
SCOOP4/STMP10 (Gully et al., 2019; Hou et al., 2021), were also significantly higher at
6 and 9 h post NHP treatment in wild type leaves compared to that observed for *wrk70-1*
plants (Fig. 6C). Unlike *BDA1*, we found that *PROSCOOP4* transcripts were higher in
NHP-treated *wrk70-1* leaves at 6 and 9h compared to the 0h timepoint (Fig. 6C). These
data show that *wrk70-1* plants responded to NHP treatment, suggesting that they can
sense NHP but are impaired in NHP signaling. Similar trends were observed for *CML10*
(*CALMODULIN-LIKE 10/CaBP22*; (Luan et al., 2002)) at the 9 h timepoint (Fig. 6D). We
confirmed these results by analyzing a second *wrky70* allele, *wrky70-2* (Supplementary
Fig. S8).
522
523 Taken together, our findings reveal that *WRKY70* is required for the proper expression of
524 NHP responsive genes from the late weak and late strong clusters, highlighting a role for
525 WRKY70 in the transcriptional response to NHP several hours following treatment.
526
NHP pretreatment enhances flg22-elicited ROS production and WRKY70
contributes to the full response
We next questioned if WRKY70 regulates a specific branch of NHP defense signaling.
One such branch involves the response to common molecular features of microbes (e.g.,
bacterial flagellin or fungal chitin) known as microbe-associated molecular patterns
(MAMPs) and the response to plant-derived damage-associated molecular patterns
(DAMPs), a process collectively referred to as pattern-triggered immunity (PTI). The
WRKY70-dependent gene *BDA1* is known to mediate signaling in response to MAMP
detection and interacts with receptor-like protein SNC2 (Suppressor of NPR1,
Constitutive2) to relay MAMP-triggered defense responses (Yang et al., 2012).
Production of reactive oxygen species (ROS) is a MAMP-triggered defense response
which can act directly or indirectly as an antimicrobial agent and serves as a secondary
signal to activate further defense responses (Boller and Felix, 2009; Couto and Zipfel,
2016). Similarly, DAMPs, such as secreted peptides from the PROSCOOP family, are

known to elicit an increase in ROS production (Gully et al., 2019; Hou et al., 2021;

Rhodes et al., 2021). Incidentally, *CML10* encodes a calmodulin-like protein known to

interact with phosphomannomutase (PMM) in order to modulate ascorbic acid synthesis

and cellular homeostasis of ROS (Cho et al., 2016).

We thus investigated how NHP alters MAMP-triggered ROS production using bacterial

flagellin as the elicitor in the presence and absence of NHP pretreatment and asked if

WRKY70 is required for the response. Leaf discs of wild type and *wrky70-1* plants were

floated on solutions of water, NHP, and SA. SA was included as a positive control, as it

is known to enhance the MAMP-triggered ROS burst (Yi et al., 2014). After 24 h, leaf discs

were treated with a 100 nM solution of flagellin peptide (flg22) and then ROS was

measured using a luminol-based assay. We found ROS levels from mock treated *wrky70-*

1 plants trended lower than water treated wild type, suggesting WRKY70 is required for

maximal ROS accumulation (Fig. 7, upper panel). NHP treatment increased ROS

generation in both wild type and *wrky70-1* tissue exposed to flg22 when compared to

water treated tissues. The average ROS production in NHP treated *wrky70-1* leaf discs

was lower than NHP treated wild type leaf discs (Fig. 7, lower panel). Similar results were

found when analyzing *wrky70-2* plants (Supplementary Fig. S9). Collectively, these data

indicate that NHP is able to enhance the flg22-elicited ROS burst, and this enhancement

requires WRKY70 for full ROS generation.

Discussion

NHP has emerged as an essential signaling molecule required for the establishment and

amplification of SAR in Arabidopsis and crop plants (Chen et al., 2018; Holmes et al.,

2019), though the early signaling events mediated by NHP had yet to be described. Our

study provides the first insights into the primary transcriptional responses to NHP, as well

as the role that NHP-induced SA biosynthesis plays during this response. We also

discovered that TGA and WRKY CREs are enriched in the promoters of early NHP

induced genes, providing a framework for dissecting primary and secondary

transcriptional responses. Further, we show that early transcriptional expression of

WRKY70, within minutes of NHP treatment, mediates secondary transcriptional
reprogramming that is required for NHP signaling, ROS production, and SAR.

Early short-lived NHP transcriptional responses associate with general stress
In the early response to NHP, we identified a primary wave of NHP signaling that was
short lived and activated transcriptional responses mirroring wounding/DAMP and JA
signaling in a partially SA-dependent manner. Of the pathways activated by NHP, we
found a considerable enrichment of known JA-response genes, including *MYC2* and
JAZ10 (Fig. 2C, Table 1, Table 2). Given the established antagonism between JA and SA
signaling pathways (Pieterse et al., 2012), it was surprising to find the majority of genes
in the early transient cluster were SA-dependent (Fig. 3A). These findings suggest that
part of the early NHP transcriptional response depends on synergism between JA
signaling components and pre-infection levels of SA. Our data support a model where
general stress response signatures are activated by NHP treatment in an early pulse of
gene expression that acts synergistically with SA to modulate activation of defense
pathways.

Previous work has described a strong, systemic JA response in Arabidopsis leaves
infected with an avirulent *Pst* strain after 4 h, suggesting JA signaling pathways are
involved in SAR development (Truman et al., 2007). The JA response detected by
Truman *et al*. in the systemic leaves of avirulent *Pst*-treated plants after 4 h may be
reflective of the early and transient response to NHP observed in our transcriptional
analysis. These JA- and early NHP-responsive marker genes may prove useful for
determining optimal time points in testing the systemic translocation of NHP. Furthermore,
although genetic studies contest the requirement for JA biosynthesis and signaling in SAR
(Attaran et al., 2009), it remains a possibility that differences in primary pathogen
treatment or secondary pathogen infection may shift the requirement for JA signaling in
SAR establishment and warrant further study.

Early strong NHP responses activate a primary wave of *WRKY* transcription

We discovered another primary wave of early NHP signaling that was partially SA-independent and exhibited strong induction of *WRKY* gene transcription within minutes of NHP treatment, persisting several hours after treatment. All six *WRKY* TF genes in the early stable upregulated gene cluster were activated by NHP treatment in the *sid2-2* mutant, and although some showed delays in the timing of induction in *sid2-2* seedlings, all six were strongly upregulated by 3 h (Table 1, Table 2). These findings are intriguing given that the expression of these *WRKY* TFs is known to be induced by SA and SA-analogs in an NPR1-dependent manner (Kalde et al., 2003; Wang et al., 2006). The ability of NHP to induce an early wave of *WRKY* expression, independent of elevated SA, suggests NHP and not SA is the primary signal driving this early response.

The early and strong induction of *WRKY* transcription, including two sets of homologs belonging to WRKY group III (*WRKY38* and *WRKY62* as well as *WRKY54* and *WRKY70*), suggested WRKY TFs may play a role in driving the secondary transcriptional response to NHP. Members of WRKY group III are known to play both positive and negative regulatory roles in defense responses (Kalde et al., 2003; Pandey and Somssich, 2009). Consistent with this, we found CREs of WRKY TFs to be abundant in the promoters of genes, both up and downregulated, several hours after NHP treatment (Fig. 4C, Supplemental Fig. 5). For example, the CRE bound by WRKY70, as defined by the DAPseq Plant Cistrome Database (O'Malley et al., 2016) , was enriched in the promoters of genes upregulated by NHP at 3 and 6 h (Fig. 4C). Other canonical motifs bound by WRKY70, such as the W-box and WT-box (Rushton et al., 2010; Machens et al., 2014), were also found in the promoters of genes upregulated several hours after NHP treatment (Fig. 6A). Taken together, these findings link the early NHP-driven induction of *WRKY* expression to the subsequent up and downregulation of WRKY-targeted genes 3-6 h after initial NHP treatment. Based on these findings, we hypothesize a model of early NHP signaling wherein activation of *WRKY* transcription is a primary response to NHP signaling that is followed by a secondary wave of expression driven through WRKY regulation.

While not the focus of this study, it is noteworthy to mention that TGA TFs are also
associated with the primary wave of NHP mediated transcription. CREs of TGAs were
abundant in the promoters of early NHP upregulated genes across all time points
sampled, particularly those in the early strong cluster (Fig. 4B). These findings are in
agreement with recent studies demonstrating TGAs are redundantly required for NHP-
induction of SAR and SAR marker genes (Nair et al., 2021; Yildiz et al., 2023). Our data
provide further evidence highlighting the importance of TGAs in NHP-driven signaling and
suggest that a NPR1/TGA complex, or a key interactor of this complex, directly relays the
NHP signal.

Late weak and late strong genes define the secondary wave of transcription
Finally, our RNA-seq analysis also revealed a secondary wave of NHP signaling defined
by late weak and late strong upregulated genes. These genes are associated with
pathogen defense, SAR, and SA responses, and showed features of SA-dependent and
SA-independent transcriptional induction (Fig. 2C, Fig. 3A). Upregulation of SAR-
associated genes by NHP is consistent with previous studies (Bernsdorff et al., 2016;
Yildiz et al., 2021; Yildiz et al., 2023). Our study reveals that transcriptional activation of
these SAR-associated genes occurs as early as 3-6 h following detection of the NHP
signal. Interestingly, 24 genes from the late weak cluster and 4 genes from the late strong
cluster are SA-independent within 6 h of NHP treatment but SA-dependent 24 h post NHP
(Yildiz et al., 2021). Among this group are key SA and NHP biosynthetic and regulatory
genes including *ALD1*, *UGT76B1*, *SARD1* and *NPR3*. Together, these data suggest NHP
is sufficient to drive gene expression over a short time frame but requires SA
accumulation for sustained expression, thus highlighting the importance of interrogation
into early NHP transcriptional changes. Further elucidation of a timeframe for early NHP
responses should prove useful in studying other aspects of NHP biology, such as the
biosynthesis and translocation of NHP and the identification of NHP sensors and their
respective downstream signaling pathways.

Primary NHP-responsive gene *WRKY70* mediates secondary transcriptional changes required for NHP-elicited SAR, signaling, and ROS production

We found *WRKY70* expression is induced within 15 min of NHP treatment and is required for full NHP-elicited SAR, as NHP did not enhance resistance in distal leaves of the *wrky70-1* mutant and could not achieve the same level of resistance observed in NHP-treated wild type plants (Fig. 5A). WRKY70 is known to repress many pathogen-inducible genes in the absence of infection, but is also required for full activation of pathogen-inducible genes during infection (Zhou et al., 2018). We observed enhanced resistance to *Psm* infection and enhanced transcript abundance of defense genes *SARD1*, *PR1*, *PR2*, and *PR5* after mock treatment (Supplement Fig. S7), consistent with a repressive role for WRKY70. However, NHP treatment did not enhance resistance nor increase transcript abundance of these genes in *wrky70-1* (Fig. 5A, Supplement Fig. S7). Furthermore, expression of the NHP-upregulated early strong gene *BDA1* was completely abolished in *wrky70* mutant lines (Fig. 6B, Supplemental Fig. S8), demonstrating early expression of *WRKY70* is required to drive and modulate a subset of the secondary transcriptional responses of NHP signaling.

Evidence supports a SA- and NPR1-independent defense pathway driven by WRKY70 (Shah et al., 2001; Zhang et al., 2010b). We found that NHP was sufficient to induce expression of *WRKY70* in the *sid2-2* mutant within 30 min of treatment (Table 1), suggesting SA biosynthesis is dispensable for the early transcriptional activation of *WRKY70*. Likewise, WRKY70-regulated genes *BDA1* and *CML10* are induced within 3-6 h of NHP signaling partially independent of SA (Table1). Notably, 24 h after NHP treatment, *BDA1* and *CML10* are also upregulated in both *sid2-1* and *npr1-3* mutants (Yildiz et al., 2021). It is therefore possible that NHP signaling activates a SA- and NPR1-independent defense pathway through the action of WRKY70.

Studies into the MAMP-signaling pathway functioning through the receptor-like protein SNC2 suggest a role for WRKY70 and BDA1 in PTI responses such as ROS production (Yang et al., 2012). While flg22-elicited ROS production was not significantly different

between mock treated wild type and *wrky70* mutants, the enhanced production of flg22-
elicited ROS following NHP and SA pretreatment was compromised in *wrky70* mutants
(Fig. 7, Supplemental Fig. S9). Our findings indicate WRKY70 positively contributes to
the NHP and SA signaling pathways that prime PTI. Further, our discovery that NHP
primes plants for enhanced production of flg22-elicited ROS is consistent with reports of
NHP priming Arabidopsis for enhanced metabolic and transcriptional responses following
flg22 treatment (Löwe et al., 2023). Our work links WRKY70-dependent signaling
pathways to this NHP-primed PTI response.

Conclusion
Our work describes the early transcriptional response to NHP and highlights key genes
and signaling pathways that may contribute to the physiological processes regulated by
this important bioactive molecule. It also provides a foundation for better understanding
the transcriptional regulators, particularly the role of WRKY70, and gene networks that
define early NHP signal transduction.

Material and methods
Plant materials and growth conditions
Arabidopsis thaliana (Arabidopsis) lines used in this study were in the Col-0 ecotype (wild
type) with mutant lines as follows: *sid2-2* (Wildermuth et al., 2001) and T-DNA insertion
lines *wrky70-1* (SALK_025198) and *wrky70-2* (GABI_324D11) (Li et al., 2006), *wrky38*
(WiscDsLox489-492C21), *wrky54* (SALK_017254), and *wrky59* (SALK_039436) (Wang
et al., 2006), *wrky51* (SALK_022198) (Yan et al., 2018), and *wrky62* (SM_3_38820) (Kim
et al., 2008). All mutant lines were obtained from the Arabidopsis Biological Resource
Center (ABRC). The presence of T-DNA insertions were confirmed with the genotyping
primers listed in Supplemental Table S1. Plants were grown on soil in a controlled growth
chamber at 80% humidity and 22°C on a 10 h light/14 h dark cycle for gene expression
and bacterial growth assays. For seedlings grown hydroponically, seeds were surface
sterilized and stratified at 4°C in darkness for 3 days. Seeds were placed into sterile six-
well plates (15-18 seeds per well) with half-strength Murashige and Skoog (MS) medium,
0.5% (w/v) sucrose, 0.05% (w/v) MES at pH 5.7. Plates were covered with a lid, sealed

with micropore tape (3M), and placed in a chamber on a 10 h light/14 h dark cycle. Liquid

medium was agitated twice daily to facilitate gas exchange and after five days of growth,

spent medium was replaced with fresh MS.

Bacterial strains and growth conditions

P. syringae pv. *maculicola* strain ES4326 (*Psm*) and *P. syringae* pv. *tomato* DC3000 (*Pst*

DC3000) carrying pVSP61 + *avrRpt2* (*Pst avrRpt2*) were used in this study. Growth

conditions for all bacterial strains are as previously described (Holmes et al., 2021).

Chemical stocks

Two sources of NHP were used in this study, one synthesized by Elizabeth Sattely's lab

as previously described (Chen et al., 2018) and the second obtained from Med Chem

Express. Gene expression and NHP-elicited SAR assays showed similar results for both

sources of NHP. Pipecolic acid stocks were made from L -pipecolic acid (Oakwood

Chemical). Salicylic acid stocks were made from sodium salicylate (Mallinckrodt).

Chemical treatment of plants for qRT-PCR assays

For gene expression by qRT-PCR of 4.5-week-old, soil-grown Arabidopsis, three leaves

were infiltrated with sterile water or NHP dissolved in water at the indicated

concentrations. The three treated leaves were collected and flash frozen at the indicated

time points. For the time course of gene expression in *wrky70-1 and wrky70-2* mutants,

three leaves were treated with sterile water or 1 mM NHP. One leaf from the same plant

was sampled at 6, 9, and 12 h and pooled with a second plant of the same condition for

one replicate. Samples collected at 0 h were untreated before collecting.

For gene expression in Arabidopsis seedling-based assays, plants were grown as

described for seedling hydroponics and treated at 10-days-post-germination. Before

addition of treatment, all liquid medium was removed from plate wells and replaced with

fresh MS medium, seedlings were allowed to recover under lights for at least 1 h, followed

by application of 2x solutions of Pip, NHP, or mock in MS medium. The 15-18 seedlings

within a single well were pooled and treated as a single replicate, with three replicates

sampled per condition. Following treatment, seedlings were blotted dry, and flash frozen
at the indicated time points.

RNA isolation and qRT-PCR

Total RNA was isolated from leaves and seedlings using TRIsure™ reagent (Meridian
Bioscience) according to the manufacturer's instructions. 5 µg of RNA was used to
synthesize cDNA by oligo dT (New England Biolabs) and reverse transcriptase (Thermo
Fisher), followed by dilution of cDNA. The qRT-PCR reactions were performed on a MJ
Opticon 2 (Bio-Rad) using Green Taq DNA polymerase (GenScript) with EvaGreen Dye
(Biotium) for amplification and detection. Three technical replicates were performed per
sample and three to four biological replicates were included for each condition.
Expression values were normalized to the reference gene *UBC21* (AT5G25760) for
relative expression determined using $2^{-\Delta Ct}$ and normalized to both *UBC21* and mock
treated samples for fold change using $2^{-\Delta\Delta Ct}$. Genes assayed include *FMO1*
(AT1G19250), *ICS1*/*SID2* (AT1G74710), *UGT76B1* (AT3G11340), *WRKY38*
(AT5G22570), *SARD1* (AT1G73805), *PR1* (AT2G14610), *PR2* (AT3G57260), *PR5*
(AT1G75040), *BDA1* (AT5G54610), *PROSCOOP4* (AT5G44568), and *CML10*
(AT2G41090). Primers used in these experiments can be found in Supplemental Table
S1.

RNA-seq treatment and library preparation

Seedlings used for mRNA isolation were grown hydroponically as described above. Six-
well plates contained equal numbers of Col-0 and *sid2-2* seedlings, with one genotype
per plate and 15 seedlings per well. At 10-days-post-germination, all liquid medium was
removed from each well and fresh MS medium was added to the seedlings. Seedlings
were placed back into the growth chamber for an hour to acclimate, followed by treatment
with 2x NHP (final concentration 0.5 mM) or mock (MS medium). Three wells of pooled
seedlings were collected for each genotype, treatment, and time (15 min, 30 min, 3 h,
and 6 h) combinations, blotted dry, and flash frozen for mRNA isolation.

Total RNA was extracted from pooled seedlings using TRIsure™ reagent (Meridian
Bioscience) according to the manufacturer's instructions. Total RNA was then divided
with 5 µg used for cDNA synthesis and marker gene expression to assess quality of
treatment, as described for qRT-PCR experiments, and 5 µg was used for cDNA library
preparation. The cDNA library was prepared using NEBNext® Ultra™ II RNA Library Prep
Kit for Illumina® with NEBNext® Poly(A) mRNA Magnetic Isolation Module and then
multiplexed via PCR amplification using NEBNext® Multiplex Oligos for Illumina®
according to the manufacturer's instructions (New England Biolabs). Library
concentration was quantified with Qubit™ dsDNA HS Assay Kit (Thermo Fisher Scientific)
followed by measurement of quality and average cDNA length using the DNA 1000 Series
II chip on a 2100 Bioanalyzer (Agilent Technologies). The multiplexed cDNA libraries
were then pooled (two sets of 36 libraries) and sequenced using both lanes of the
NovaSeq 6000 (Illumina) S1 flow cell with run type 100 bp paired-end reads at the
Genome Sequencing Service Center by the Stanford Center for Genomics and
Personalized Medicine, supported by the grant award NIH S10OD025212.
797

RNA-seq data analysis
Read data was initially assessed using FastQC. CLC Genomics Workbench version
11.0.2 (QIAGEN) was used for trimming, discarding reads trimmed below 20 bp, and then
mapped to the Arabidopsis TAIR10 genome with the auto-detect paired distances option
deselected. Differential expression analysis was performed using DESeq2 (Love et al.,
2014) with mock treated samples compared to NHP treated samples for each respective
time point and genotype. Genes were considered differentially expressed if they returned
an adjusted P-value below 0.05 ($P_{adj} < 0.05$). The complete list of early NHP-responsive
transcripts is provided in Supplemental Table S3 and the raw sequencing data deposited
in the Gene Expression Omnibus (GEO) under accession code GSE263276.
808

Exploratory analysis of differentially expressed genes
To define patterns of early expression in response to NHP treatment, differentially
expressed genes were considered upregulated in wild type if they had a $\log_2(FC) > 1$ at
any time point and downregulated with a $\log_2(FC) < -1$ at any time point. The sets of up

and downregulated genes were then hierarchically clustered by $\log_2$(FC) over the four
timepoints using the hclust function and dendextend packages in R, with Euclidean
distances and Ward.D2 linkage (Galili, 2015). For each defined cluster in wild type, the
$\log_2$(FC) values were averaged at each time point to visualize broad trends in expression.
817
818 The PANTHER classification system accessed via TAIR was used for gene ontology
819 enrichment analysis (https://www.arabidopsis.org/tools/go_term_enrichment.jsp; (Mi et
820 al., 2021)). Version 17.0 of the PANTHER overrepresentation test was used (Fisher's
821 exact test with FDR correction) and matched to the GO database 2021_03 release.
822

Promoter and *cis*-regulatory element enrichment analysis

Araport11 sequences 1000 bp upstream of the TSS of NHP-responsive genes were
downloaded from TAIR (https://www.arabidopsis.org/tools/bulk/sequences/index.jsp) for
each time point and genotype grouping of the NHP upregulated and downregulated
genes. A less stringent cut-off of $\log_2$(FC) > 0 (upregulated) or $\log_2$(FC) < 0
(downregulated) with a $P_{adj} < 0.05$ was used to select for NHP-responsive genes. Motif
identification and enrichment of *cis*-regulatory elements (CREs) was performed using
SEA, part of the MEME suite (Bailey et al., 2015; Bailey and Grant, 2021) and compared
to the published library of TF binding sites found within the DAPseq database (O'Malley
et al., 2016). Only CREs from the DAP-seq library were used in this study, omitting CREs
from the ampDAP-seq library (where DNA modifications have been removed). Following
the workflow detailed by Bjornson et al. (2021), relative enrichment was determined
against the same three sets of randomly sampled genes detected in this RNA-seq
experiment for all genotypes and time points tested. The identified motif was considered
enriched if it returned a significant P-value (< 0.05) relative to at least two of the three
randomly sampled gene sets. P-values for significantly enriched TF-binding motifs were
then averaged for visualization.
840

Measurement of bacterial growth in plants

Chemical treatment of leaves for NHP-elicited SAR assays were performed as described
in (Chen et al., 2018) with minor alterations. At 4.5-weeks-old, three lower leaves of wild

type and mutant Arabidopsis plants were infiltrated with sterile water, 0.5 mM NHP or 1

mM NHP. After 24 h, one untreated distal leaf of each plant was inoculated with a 1 x 10^5

CFU/mL suspension of *Psm*. The distal infected leaf was always separated by one leaf

from the original three treated (i.e., primary treatment in leaf no. 7-9 then infected leaf is

no. 11). Inoculated plants were covered with a dome to increase humidity and 3 dpi the

titer of *Psm* in the distal leaves was quantified. Leaf discs were homogenized in 1 mL of

10 mM $MgCl_2$, plated in triplicate in a dilution series on nutrient yeast glycerol medium

supplemented with 1.5% wt/vol agar (NYGA) with rifampicin (100 µg/mL), incubated at 28

°C for 1.5 d, and bacterial colonies counted. For each individual experiment, 4-12 plants

were tested per condition.

Pathogen priming of leaves for SAR assays were performed as previously described

(Chen et al., 2018). Three lower leaves of wild type and mutant Arabidopsis plants (4.5-

weeks-old) were infiltrated with 10 mM $MgCl_2$ or a 5 x 10^6 CFU/mL suspension of *Pst*

avrRpt2 in 10 mM $MgCl_2$. After 48 h, one distal untreated leaf of each plant was inoculated

with a 1 x 10^5 CFU/mL suspension of *Psm* and plants were covered with a dome to

increase humidity. At 3 dpi, the titer of *Psm* in the upper leaves was quantified as

described above for NHP-elicited SAR. For each individual experiment, 8 plants were

tested per condition and experiments repeated twice with similar results.

For bacterial growth measurements in unprimed leaves of Arabidopsis wild type and

mutant plants, a suspension of 1 x 10^5 CFU/mL *Psm* was infiltrated into two young leaves

per plant. Plants were placed under a dome and 3 dpi the two infected leaves were pooled

and bacterial titer quantified for 12 plants per genotype.

Oxidative burst assay

Four leaf discs (4 mm diameter) from 5-week-old Arabidopsis plants (n = 12) were

incubated on water, 500 µM NHP, or 100 µM SA in a 96-well plate (one leaf disc per well)

for 24 h. After pretreatment, ROS was measured by addition of flg22 (100 nM) in a solution

of 20 mg/mL horseradish peroxidase and 200 µM Luminol (Sigma) (Gómez-Gómez et al.,

1999), followed by immediate measurement of luminescence in a Synergy H1 Microplate
Reader (Biotek). Relative luminescence units (RLU) are reported.

Statistical analyses

Two-tailed Student's t tests were performed to determine statistically significant
differences between the different conditions for measurements of bacterial growth using
the stat_compare_means function of the ggpubr package (Kassambara, 2023). P values
< 0.05 were considered statistically significant and all significance levels are indicated in
the figure legends (*P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001). Statistical
significance of qRT-PCR assays was determined using Mann-Whitney U test or one-way
ANOVA with post hoc Sidak test using the ggpubr and lsmeans R packages, respectively
(Lenth, 2016).

Acknowledgements

We thank the Mudgett laboratory for feedback on the research and manuscript. We also
thank Zhiyong Wang, Dominique Bergmann, Sharon Long, and Virginia Walbot for
instrument use and intellectual discussion. This work was supported by National Science
Foundation IOS-2026368 (to M.B.M. and E.S.S) and National Science Foundation
Graduate Research Fellowship DGE-1656518 (to J.F.).

Author contributions

J.F., J.-G.K., E.S.S. and M.B.M contributed to the study design; J.F. and J.-G.K.
performed research; J.F., J.-G.K., E.S.S. and M.B.M analyzed data; J.F., and M.B.M.
wrote the manuscript; J.F., J.-G.K., E.S.S. and M.B.M. reviewed and revised the
manuscript for accuracy and final approval.

References

Attaran E, Zeier TE, Griebel T, Zeier J (2009) Methyl salicylate production and
jasmonate signaling are not essential for systemic acquired resistance in *Arabidopsis*.
Plant Cell **21**: 954–971

Bailey TL, Grant CE (2021) SEA: Simple Enrichment Analysis of motifs. BioRxiv. doi:
10.1101/2021.08.23.457422

Bailey TL, Johnson J, Grant CE, Noble WS (2015) The MEME Suite. Nucleic Acids
Res **43**: W39-49

Bartsch M, Gobbato E, Bednarek P, Debey S, Schultze JL, Bautor J, Parker JE
(2006) Salicylic acid-independent ENHANCED DISEASE SUSCEPTIBILITY1
signaling in Arabidopsis immunity and cell death is regulated by the monooxygenase
FMO1 and the Nudix hydrolase NUDT7. Plant Cell **18**: 1038–1051

Bauer S, Mekonnen DW, Hartmann M, Yildiz I, Janowski R, Lange B, Geist B, Zeier
J, Schäffner AR (2021) UGT76B1, a promiscuous hub of small molecule-based
immune signaling, glucosylates N-hydroxypipecolic acid, and balances plant
immunity. Plant Cell **33**: 714–734

Bernsdorff F, Döring A-C, Gruner K, Schuck S, Bräutigam A, Zeier J (2016)
Pipecolic Acid Orchestrates Plant Systemic Acquired Resistance and Defense
Priming via Salicylic Acid-Dependent and -Independent Pathways. Plant Cell **28**:
102–129

Bjornson M, Pimprikar P, Nürnberger T, Zipfel C (2021) The transcriptional
landscape of *Arabidopsis thaliana* pattern-triggered immunity. Nat Plants **7**: 579–586

Blanco F, Salinas P, Cecchini NM, Jordana X, Van Hummelen P, Alvarez ME,
Holuigue L (2009) Early genomic responses to salicylic acid in Arabidopsis. Plant
Mol Biol **70**: 79–102

Boller T, Felix G (2009) A renaissance of elicitors: perception of microbe-associated
molecular patterns and danger signals by pattern-recognition receptors. Annu Rev
Plant Biol **60**: 379–406

Cai J, Jozwiak A, Holoidovsky L, Meijler MM, Meir S, Rogachev I, Aharoni A (2021)
Glycosylation of N-hydroxy-pipecolic acid equilibrates between systemic acquired
resistance response and plant growth. Mol Plant **14**: 440–455

Chen Y-C, Holmes EC, Rajniak J, Kim J-G, Tang S, Fischer CR, Mudgett MB,
Sattely ES (2018) *N*-hydroxy-pipecolic acid is a mobile metabolite that induces
systemic disease resistance in *Arabidopsis.* Proc Natl Acad Sci USA **115**: E4920–
E4929

Chini A, Fonseca S, Fernández G, Adie B, Chico JM, Lorenzo O, García-Casado G,
López-Vidriero I, Lozano FM, Ponce MR, et al (2007) The JAZ family of repressors
is the missing link in jasmonate signalling. Nature **448**: 666–671

Cho K-M, Nguyen HTK, Kim SY, Shin JS, Cho DH, Hong SB, Shin JS, Ok SH (2016)
CML10, a variant of calmodulin, modulates ascorbic acid synthesis. New Phytol **209**:
664–678

Conrath U, Beckers GJM, Langenbach CJG, Jaskiewicz MR (2015) Priming for
enhanced defense. Annu Rev Phytopathol **53**: 97–119

Couto D, Zipfel C (2016) Regulation of pattern recognition receptor signalling in plants.
Nat Rev Immunol **16**: 537–552

Ding Y, Sun T, Ao K, Peng Y, Zhang Y, Li X, Zhang Y (2018) Opposite Roles of
Salicylic Acid Receptors NPR1 and NPR3/NPR4 in Transcriptional Regulation of
Plant Immunity. Cell **173**: 1454-1467.e15

Eulgem T, Rushton PJ, Robatzek S, Somssich IE (2000) The WRKY superfamily of
plant transcription factors. Trends Plant Sci **5**: 199–206

Friedrichsen DM, Nemhauser J, Muramitsu T, Maloof JN, Alonso J, Ecker JR,
Furuya M, Chory J (2002) Three redundant brassinosteroid early response genes
encode putative bHLH transcription factors required for normal growth. Genetics **162**:
1445–1456

Fu ZQ, Dong X (2013) Systemic acquired resistance: turning local infection into global
defense. Annu Rev Plant Biol **64**: 839–863

Galili T (2015) dendextend: an R package for visualizing, adjusting and comparing
trees of hierarchical clustering. Bioinformatics **31**: 3718–3720

Gómez-Gómez L, Felix G, Boller T (1999) A single locus determines sensitivity to
bacterial flagellin in Arabidopsis thaliana. Plant J **18**: 277–284

Gully K, Pelletier S, Guillou M-C, Ferrand M, Aligon S, Pokotylo I, Perrin A, Vergne
E, Fagard M, Ruelland E, et al (2019) The SCOOP12 peptide regulates defense
response and root elongation in *Arabidopsis thaliana*. J Exp Bot **70**: 1349–1365

Hartmann M, Zeier J (2019) N-hydroxypipecolic acid and salicylic acid: a metabolic duo
for systemic acquired resistance. Curr Opin Plant Biol **50**: 44–57

Hartmann M, Zeier T, Bernsdorff F, Reichel-Deland V, Kim D, Hohmann M,
Scholten N, Schuck S, Bräutigam A, Hölzel T, et al (2018) Flavin Monooxygenase-
Generated N-Hydroxypipecolic Acid Is a Critical Element of Plant Systemic Immunity.
Cell **173**: 456-469.e16

Holmes EC, Chen Y-C, Mudgett MB, Sattely ES (2021) Arabidopsis UGT76B1
glycosylates *N*-hydroxy-pipecolic acid and inactivates systemic acquired resistance in
tomato. Plant Cell **33**: 750–765

Holmes EC, Chen Y-C, Sattely ES, Mudgett MB (2019) An engineered pathway for *N*-
hydroxy-pipecolic acid synthesis enhances systemic acquired resistance in tomato.
Sci Signal. 12:eaay3066

Hou S, Liu D, Huang S, Luo D, Liu Z, Xiang Q, Wang P, Mu R, Han Z, Chen S, et al
(2021) The Arabidopsis MIK2 receptor elicits immunity by sensing a conserved
signature from phytocytokines and microbes. Nat Commun **12**: 5494

Jaskiewicz M, Conrath U, Peterhänsel C (2011) Chromatin modification acts as a
memory for systemic acquired resistance in the plant stress response. EMBO Rep **12**:
50–55

Johannesson H, Wang Y, Hanson J, Engström P (2003) The *Arabidopsis thaliana*
homeobox gene *ATHB5* is a potential regulator of abscisic acid responsiveness in
developing seedlings. Plant Mol Biol **51**: 719–729

Kachroo A, Kachroo P (2020) Mobile signals in systemic acquired resistance. Curr
Opin Plant Biol **58**: 41–47

Kalde M, Barth M, Somssich IE, Lippok B (2003) Members of the *Arabidopsis* WRKY
group III transcription factors are part of different plant defense signaling pathways.
Mol Plant Microbe Interact **16**: 295–305

Kassambara A (2023) *ggpubr: 'ggplot2' Based Publication Ready Plots*. R package
version 0.6.0. https://rpkgs.datanovia.com/ggpubr/

Kim K-C, Lai Z, Fan B, Chen Z (2008) *Arabidopsis* WRKY38 and WRKY62
transcription factors interact with histone deacetylase 19 in basal defense. Plant Cell
20: 2357–2371

Kim Y, Gilmour SJ, Chao L, Park S, Thomashow MF (2020) *Arabidopsis* CAMTA
transcription factors regulate pipecolic acid biosynthesis and priming of immunity
genes. Mol Plant **13**: 157–168

Klessig DF, Choi HW, Dempsey DA (2018) Systemic acquired resistance and salicylic
acid: past, present, and future. Mol Plant Microbe Interact **31**: 871–888

Kohler A, Schwindling S, Conrath U (2002) Benzothiadiazole-Induced Priming for
Potentiated Responses to Pathogen Infection, Wounding, and Infiltration of Water into
Leaves Requires the *NPR1*/*NIM1* Gene in Arabidopsis. Plant Physiol **128**: 1046–1056

Lenth RV (2016) Least-Squares Means: the *R* package lsmeans. J Stat Softw **69**: 1–33

Liu H, Liu B, Lou S, Bi H, Tang H, Tong S, Song Y, Chen N, Zhang H, Jiang Y, et al
(2021) CHYR1 ubiquitinates the phosphorylated WRKY70 for degradation to balance
immunity in *Arabidopsis thaliana*. New Phytol **230**: 1095–1109

Li J, Brader G, Kariola T, Palva ET (2006) WRKY70 modulates the selection of
signaling pathways in plant defense. Plant J **46**: 477–491

Li J, Brader G, Palva ET (2004) The WRKY70 transcription factor: a node of
convergence for jasmonate-mediated and salicylate-mediated signals in plant
defense. Plant Cell **16**: 319–331

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol **15**: 550

Löwe M, Jürgens K, Zeier T, Hartmann M, Gruner K, Müller S, Yildiz I, Perrar M,
Zeier J (2023) N-hydroxypipecolic acid primes plants for enhanced microbial pattern-
induced responses. Front Plant Sci **14**: 1217771

Luan S, Kudla J, Rodriguez-Concepcion M, Yalovsky S, Gruissem W (2002)
Calmodulins and calcineurin B-like proteins: calcium sensors for specific signal
response coupling in plants. Plant Cell **14 Suppl**: S389-400

Machens F, Becker M, Umrath F, Hehl R (2014) Identification of a novel type of
WRKY transcription factor binding site in elicitor-responsive *cis*-sequences from
Arabidopsis thaliana. Plant Mol Biol **84**: 371–385

Mariani L, Weinand K, Vedenko A, Barrera LA, Bulyk ML (2017) Identification of
Human Lineage-Specific Transcriptional Coregulators Enabled by a Glossary of
Binding Modules and Tunable Genomic Backgrounds. Cell Syst **5**: 187-201.e7

Mishina TE, Zeier J (2006) The Arabidopsis Flavin-Dependent Monooxygenase FMO1
Is an Essential Component of Biologically Induced Systemic Acquired Resistance.
Plant Physiol **141**: 1666–1675

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD
(2021) PANTHER version 16: a revised family classification, tree-based classification
tool, enhancer regions and extensive API. Nucleic Acids Res **49**: D394–D403

Mohanta TK, Yadav D, Khan A, Hashem A, Tabassum B, Khan AL, Abd Allah EF,
Al-Harrasi A (2020) Genomics, molecular and evolutionary perspective of NAC
transcription factors. PLoS ONE **15**: e0231425

Mohnike L, Rekhter D, Huang W, Feussner K, Tian H, Herrfurth C, Zhang Y,
Feussner I (2021) The glycosyltransferase UGT76B1 modulates N-hydroxy-pipecolic
acid homeostasis and plant immunity. Plant Cell **33**: 735–749

Nair A, Goyal I, Voß E, Mrozek P, Prajapati S, Thurow C, Tietze L, Tittmann K, Gatz
C (2021) N-hydroxypipecolic acid-induced transcription requires the salicylic acid
signaling pathway at basal SA levels. Plant Physiol **187**: 2803–2819

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M,
Gallavotti A, Ecker JR (2016) Cistrome and Epicistrome Features Shape the
Regulatory DNA Landscape. Cell **165**: 1280–1292

Pandey SP, Somssich IE (2009) The Role of WRKY Transcription Factors in Plant
Immunity. Plant Physiol **150**: 1648–1655

Pieterse CMJ, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SCM (2012)
Hormonal modulation of plant immunity. Annu Rev Cell Dev Biol **28**: 489–521

Rhodes J, Yang H, Moussu S, Boutrot F, Santiago J, Zipfel C (2021) Perception of a
divergent family of phytocytokines by the Arabidopsis receptor kinase MIK2. Nat
Commun **12**: 705

Rushton PJ, Somssich IE, Ringler P, Shen QJ (2010) WRKY transcription factors.
Trends Plant Sci **15**: 247–258

Shah J, Kachroo P, Nandi A, Klessig DF (2001) A recessive mutation in the
Arabidopsis SSI2 gene confers SA- and *NPR1*-independent expression of *PR* genes
and resistance against bacterial and oomycete pathogens. Plant J **25**: 563–574

Song JT, Lu H, McDowell JM, Greenberg JT (2004) A key role for ALD1 in activation
of local and systemic defenses in *Arabidopsis*. Plant J **40**: 200–212

Sun T, Huang J, Xu Y, Verma V, Jing B, Sun Y, Ruiz Orduna A, Tian H, Huang X,
Xia S, et al (2020) Redundant CAMTA Transcription Factors Negatively Regulate the
Biosynthesis of Salicylic Acid and *N*-Hydroxypipecolic Acid by Modulating the
Expression of *SARD1* and *CBP60g*. Mol Plant **13**: 144–156

Thibaud-Nissen F, Wu H, Richmond T, Redman JC, Johnson C, Green R, Arias J,
Town CD (2006) Development of Arabidopsis whole-genome microarrays and their
application to the discovery of binding sites for the TGA2 transcription factor in
salicylic acid-treated plants. Plant J **47**: 152–162

Tian H, Zhang Y (2019) The emergence of a mobile signal for systemic acquired
resistance. Plant Cell **31**: 1414–1415

Truman W, Bennett MH, Kubigsteltig I, Turnbull C, Grant M (2007) *Arabidopsis*
systemic immunity uses conserved defense signaling pathways and is mediated by
jasmonates. Proc Natl Acad Sci USA **104**: 1075–1080

Wang D, Amornsiripanitch N, Dong X (2006) A genomic approach to identify
regulatory nodes in the transcriptional network of systemic acquired resistance in
plants. PLoS Pathog **2**: e123

Wang Y, Schuck S, Wu J, Yang P, Döring A-C, Zeier J, Tsuda K (2018) A MPK3/6-
WRKY33-ALD1-Pipecolic Acid Regulatory Loop Contributes to Systemic Acquired
Resistance. Plant Cell **30**: 2480–2494

Wildermuth MC, Dewdney J, Wu G, Ausubel FM (2001) Isochorismate synthase is
required to synthesize salicylic acid for plant defence. Nature **414**: 562–565

Wu Y, Zhang D, Chu JY, Boyle P, Wang Y, Brindle ID, De Luca V, Després C (2012)
The *Arabidopsis* NPR1 Protein Is a Receptor for the Plant Defense Hormone Salicylic
Acid. Cell Rep **1**: 639–647

Yang Y, Zhang Y, Ding P, Johnson K, Li X, Zhang Y (2012) The ankyrin-repeat
transmembrane protein BDA1 functions downstream of the receptor-like protein
SNC2 to regulate plant immunity. Plant Physiol **159**: 1857–1865

Yan C, Fan M, Yang M, Zhao J, Zhang W, Su Y, Xiao L, Deng H, Xie D (2018) Injury
Activates Ca2+/Calmodulin-Dependent Phosphorylation of JAV1-JAZ8-WRKY51
Complex for Jasmonate Biosynthesis. Mol Cell **70**: 136-149.e7

Yi SY, Shirasu K, Moon JS, Lee S-G, Kwon S-Y (2014) The activated SA and JA
signaling pathways have an influence on flg22-triggered oxidative burst and callose
deposition. PLoS ONE **9**: e88951

Yildiz I, Gross M, Moser D, Petzsch P, Köhrer K, Zeier J (2023) N-hydroxypipecolic
acid induces systemic acquired resistance and transcriptional reprogramming via
TGA transcription factors. Plant Cell Environ. doi: 10.1111/pce.14572

Yildiz I, Mantz M, Hartmann M, Zeier T, Kessel J, Thurow C, Gatz C, Petzsch P,
Köhrer K, Zeier J (2021) The mobile SAR signal N-hydroxypipecolic acid induces
NPR1-dependent transcriptional reprogramming and immune priming. Plant Physiol
186: 1679–1705

Zhang Y, Fan W, Kinkema M, Li X, Dong X (1999) Interaction of NPR1 with basic
leucine zipper protein transcription factors that bind sequences required for salicylic
acid induction of the *PR-1* gene. Proc Natl Acad Sci USA **96**: 6523–6528

Zhang Y, Xu S, Ding P, Wang D, Cheng YT, He J, Gao M, Xu F, Li Y, Zhu Z, et al
(2010a) Control of salicylic acid synthesis and systemic acquired resistance by two
members of a plant-specific family of transcription factors. Proc Natl Acad Sci USA
107: 18220–18225

Zhang Y, Yang Y, Fang B, Gannon P, Ding P, Li X, Zhang Y (2010b) *Arabidopsis*
snc2-1D activates receptor-like protein-mediated immunity transduced through
WRKY70. Plant Cell **22**: 3153–3163

Zhou M, Lu Y, Bethke G, Harrison BT, Hatsugai N, Katagiri F, Glazebrook J (2018)
WRKY70 prevents axenic activation of plant immunity by direct repression of *SARD1*.
New Phytol **217**: 700–712

 Tables

Table 1. SA-independent/partially independent NHP upregulated genes by cluster.

AGI	Name	Annotation	Col-0 15m	30m	3h	6h	sid2-2 15m	30m	3h	6h
Early transient										
AT3G44870	FAMT-L	FARNESOIC ACID METHYL TRANSFERASE-LIKE	4.3	4.4	-	-	2.3	2.7	-	-
AT2G44840	ERF13	Ethylene-responsive transcription factor	3.3	2.1	-	-	1.4	-	-	-
AT1G76640	CML39	CALMODULIN LIKE 39	2.7	2.7	-	-	1.7	-	-	-
AT5G13220	JAZ10	JASMONATE-ZIM-DOMAIN PROTEIN 10	1.6	1.1	-	0.6	1.2	1.0	-	-
AT1G80840	WRKY40	WRKY Transcription Factor; Group II-a	1.8	1.1	-	0.5	-	1.4	1.4	0.7
AT3G17690	CNGC19	Cyclic nucleotide-gated ion channel	1.6	1.6	-	-	-	1.2	-	-
AT4G21840	MSRB8	Methionine sulfoxide reductase B8	3.1	-	-	1.5	-	2.2	-	-
Early strong										
AT5G22570	WRKY38	WRKY Transcription Factor; Group III	5.8	5.2	4.7	3.6	3.9	6.2	5.8	2.9
AT5G01900	WRKY62	WRKY Transcription Factor; Group III	3.9	5.0	4.4	2.6	-	5.8	4.1	4.4
AT1G28480	GRXC9	GRX480/ROXY19, glutaredoxin family	3.2	2.2	2.7	1.8	3.2	3.1	2.3	2.1
AT2G40750	WRKY54	WRKY Transcription Factor; Group III	2.7	3.7	2.8	2.9	3.0	3.6	3.5	3.0
AT5G64810	WRKY51	WRKY Transcription Factor; Group II-c	2.6	3.4	3.3	3.0	1.7	3.2	3.2	3.5
AT3G56400	WRKY70	WRKY Transcription Factor; Group III	1.6	1.6	2.4	2.1	-	1.3	2.3	1.8
AT3G25882	NIMIN-2	NIM1-INTERACTING 2	1.4	1.5	1.8	1.6	1.2	1.1	1.0	1.6
Late weak										
AT3G60470	-	Transmembrane protein, putative (DUF247)	-	-	-	3.3	-	-	-	5.1
AT2G13810	ALD1	AGD2-LIKE DEFENSE RESPONSE PROTEIN 1	-	-	-	2.3	-	-	-	6.6
AT1G01680	PUB54	Plant U-box type E3 ubiquitin ligase	-	-	-	2.2	-	-	-	1.4
AT3G12220	SCPL16	Serine carboxypeptidase-like	-	-	1.1	2.0	-	-	-	2.0
AT1G78340	GSTU22	Glutathione S-transferase U22	-	-	2.0	2.0	-	-	1.6	1.7
AT5G03350	LLP	Lectin-like protein	-	-	2.1	1.8	-	-	2.1	1.9
AT5G24530	DMR6	DOWNY MILDEW RESISTANCE 6	0.7	1.2	1.6	1.7	0.5	1.6	2.0	1.7
AT2G41090	CML10	Calmodulin-like protein	-	-	1.1	1.7	-	-	0.3	1.2
AT4G14400	ACD6	ACCELERATED CELL DEATH 6	-	-	2.1	1.6	-	-	1.4	1.4
AT2G15490	UGT73B4	UDP-DEPENDENT GLYCOSYLTRANSFERASE 73B4	-	0.7	1.2	1.5	-	0.9	2.7	1.5
AT4G08040	ACS11	1-aminocyclopropane-1-carboxylate synthase 11	-	1.0	1.5	1.5	-	-	1.0	1.2
AT1G21250	WAK1	CELL WALL-ASSOCIATED KINASE 1	0.7	-	1.8	1.5	-	-	1.7	2.0
AT1G17170	GSTU24	Glutathione S-transferase U24	-	-	-	1.4	-	-	1.6	1.5
AT3G11340	UGT76B1	UDP-DEPENDENT GLYCOSYLTRANSFERASE 76B1	-	1.0	0.9	1.1	-	0.7	1.1	1.0
AT1G73805	SARD1	SAR DEFICIENT 1	1.0	0.7	1.5	1.0	-	-	1.3	1.3

Late strong

AGI	Symbol	Description								
AT3G12230	SCPL14	Serine carboxypeptidase-like	-	-	5.9	6.9	-	-	5.4	4.6
AT1G15610	-	Transmembrane protein	-	-	5.5	5.9	-	-	-	5.1
AT3G28510	-	AAA-type ATPase family protein	-	-	3.2	5.6	-	-	-	4.9
AT2G26400	ARD	ACIREDUCTONE DIOXYGENASE 3	-	2.2	4.4	5.0	-	-	5.1	5.8
AT4G19750	-	Glycosyl hydrolase family with chitinase insertion domain	-	-	3.9	4.8	-	-	6.7	5.3
AT1G19960	-	Putative uncharacterized protein	1.2	1.6	4.0	4.0	-	2.9	5.5	4.4
AT4G10500	DLO1	DMR6-LIKE OXYGENASE 1	-	1.4	3.1	3.8	-	-	3.4	4.2
AT5G41280	CRRSP57	Receptor-like protein kinase-related family protein	-	-	2.6	3.2	-	-	3.2	2.2
AT2G14560	LURP1	late upregulated in response to *Hyaloperonospora parasitica*	-	-	3.0	3.1	-	-	-	4.5
AT1G02230	NAC004	NAC DOMAIN CONTAINING PROTEIN 4	-	2.8	3.8	3.0	-	2.5	3.5	4.3
AT5G54610	BDA1	Ankyrin repeat protein family	-	-	2.2	3.0	-	-	3.0	6.4

Selection of genes upregulated by NHP in Col-0 and *sid2-2* seedlings. Clusters defined for wild type expression are noted (i.e., early transient, early strong, late weak, late strong). Indicated $\log_2$ fold change (FC) values determined relative to mock treatment for each genotype and time point. Genes are categorized as SA-independent/partially independent if significant upregulation was observed in *sid2-2* within 15-30 min for the early clusters and within 3-6 h for the late clusters. Not all NHP upregulated genes are shown in this table, the complete list can be found in Supplemental Table S3.

1111

Table 2. SA-dependent NHP upregulated genes by cluster.

			Log$_2$(FC)							
			Col-0				sid2-2			
AGI	Name	Annotation	15m	30m	3h	6h	15m	30m	3h	6h
Early transient										
AT1G32970	SBT3.2	Subtilisin-like protease	2.8	3.2	-	-	-	-	-	-
AT1G21240	WAK3	Wall-associated receptor kinase	2.5	-	-	1.4	-	-	-	2.0
AT5G44430	PDF1.2C	PLANT DEFENSIN 1.2C	2.4	-	-	-	-	-	-	-
AT2G14610	PR1	PATHOGENESIS-RELATED GENE 1	2.3	-	-	-	-	-	-	-
AT4G34410	ERF109	Ethylene-responsive transcription factor	2.1	-	-	-	-	-	-	-
AT2G32140	-	Transmembrane receptor	2.1	-	-	-	-	-	2.0	-
AT5G44420	PDF1.2A	PLANT DEFENSIN 1.2	2.0	-	-	-	-	-	-	-
AT3G14260	-	LURP-one-like protein (DUF567)	1.8	-	-	-	-	-	-	-
AT2G26020	PDF1.2B	PLANT DEFENSIN 1.2B	1.6	-	-	-	-	-	-	-
AT5G64905	PROPEP3	ELICITOR PEPTIDE 3 PRECURSOR	1.5	-	-	-	-	-	-	-
AT1G32640	MYC2	JASMONATE INSENSITIVE 1	1.4	-	-	-	-	-	-	-
AT2G18660	EGC2	PLANT NATRIURETIC PEPTIDE A	1.4	-	-	-	-	-	-	-
Early strong										
AT2G21900	WRKY59	WRKY Transcription Factor; Group II-c	1.9	2.4	2.3	2.3	-	-	2.2	2.7
Late weak										
AT2G28850	CYP710A3	Cytochrome P450 710A3	-	-	2.4	2.1	-	-	-	-
AT1G15630	-	Transmembrane protein	-	-	1.9	1.4	-	-	-	-
AT5G44568	PROSCOOP4	SERINE RICH ENDOGENOUS PEPTIDE	-	-	1.3	1.3	-	-	0.7	1.0
AT1G18830	SEC31A	Protein transport protein SEC31 homolog A	-	-	-	2.8	-	-	-	-
AT2G45550	CYP76C4	Cytochrome P450 76C4	-	-	-	2.6	-	-	-	-
AT1G10155	ATPP2-A10	PHLOEM PROTEIN 2-A10	-	-	-	2.3	-	-	-	-
AT1G02450	NIMIN-1	NIM1-INTERACTING 1	-	-	-	2.3	-	-	-	-
AT3G45860	CRK4	Cysteine-rich receptor-like protein kinase	-	-	-	2.1	-	-	-	-
AT5G51500	PME60	Pectin methylesterase inhibitor	-	-	-	1.7	-	-	-	-
AT2G29350	SAG13	SENESCENCE-ASSOCIATED GENE 13	-	-	-	1.3	-	-	-	-
Late strong										
AT3G22231	PCC1	PATHOGEN AND CIRCADIAN CONTROLLED 1	-	-	3.8	3.9	-	-	-	-

Selection of genes upregulated by NHP in Col-0 and *sid2-2* seedlings. Clusters defined for wild type (Col-0) expression are noted. Indicated log$_2$ fold change (FC) values were determined relative to mock treatment for each genotype and time point. Genes are categorized as SA-dependent if significant upregulation was not observed in *sid2-2* within 15-30 min for the early clusters and within 3-6 h for the late clusters. Not all NHP upregulated genes are shown in this table, the complete list can be found in Supplemental Table S3.

1112

1113

1114　**Figure Legends**

1115

Figure 1 NHP induces *WRKY38* expression at early time points.

A, Transcript abundance of SAR marker genes *FMO1*, *ICS1*, and *PR1* and early SA-

responsive marker genes *WRKY38* and *UGT76B1* in 4.5-week-old wild type (Col-0)

Arabidopsis plants. Three leaves were infiltrated with sterile water (mock) or 1 mM NHP

and collected after 2, 6, and 24 h for mRNA isolation. Transcript abundance was

measured via quantitative real-time PCR (qRT-PCR) and normalized relative to *UBC21*

(Ubiquitin-Conjugating Enzyme 21; At5g25760). Fold change was determined relative to

mock ($2^{-\Delta\Delta Ct}$). Asterisks indicate a significant difference between mock and NHP per time

point (n = 4; Mann-Whitney *U* test; *P < 0.05, ns = not significant). B, Design of seedling-

based assay. Arabidopsis seedlings (15-18 per well) were hydroponically grown, treated

with MS medium (mock), 1 mM Pip, or 1 mM NHP in MS medium then collected at 15

min, 30 min, 3 h, and 6 h for qRT-PCR analysis of *WRKY38* transcript abundance in wild

type. C, Expression of *WRKY38* in seedlings from the experiment described in B.

Transcript abundance was normalized relative to *UBC21* and mock treated samples for

each condition ($2^{-\Delta\Delta Ct}$). Asterisks indicate a significant difference between mock and Pip

or NHP at each time point (n = 4-6; Mann-Whitney *U* test; **$P <$ 0.01, ns = not significant).

1132

1133

Figure 2 Profile of early NHP-upregulated genes in wild type seedlings.

A, Heatmap of genes upregulated ($\log_2$(FC) > 1, P_{adj} < 0.05) in response to NHP treatment

in wild type seedlings. One biological replicate consists of 15 pooled seedlings per

condition (n = 3). Euclidean distances determined from the $\log_2$(FC) values were

hierarchically clustered across the indicated time points (15 min, 30 min, 3 h and 6 h),

resulting in four distinct clusters of gene expression. B, Average $\log_2$(FC) of all genes

within each cluster defined in (A) showing the expression trends across the four time

points. Note, the two genes expressed only at 30 min, and no other time, were removed

from the 'early strong' cluster average. Error bars represent the ± SEM. C, Biological

processes of significantly enriched (FDR < 0.05) Gene Ontology (GO) terms for the genes

within each cluster are shown. Bars represent the -log(FDR) of each significantly enriched

GO term. Bars were omitted if the term was not significantly enriched for the defined

cluster. The most significantly enriched GO term for each cluster is indicated in bold.

1147

1148

Figure 3 Comparison of NHP upregulated genes in wild type and *sid2-2* seedlings.

A, Percent of NHP upregulated genes expressed only in wild type (Col-0, SA-dependent)

and in both wild type and the *sid2-2* mutant (SA-independent) for the wild type gene

expression clusters defined in Figure 2. Total number of genes in each condition is

indicated. B, Total number of NHP upregulated genes ($\log_2$(FC) > 1, P_{adj} < 0.05) per time

point unique to wild type (Col-0), shared between wild type and *sid2-2*, and unique to

sid2-2. The number of upregulated genes in each grouping is indicated. C, Biological

processes of significantly enriched (gray bar) or depleted (white bar) (FDR < 0.05) GO

terms for the set of genes upregulated 3 and 6 h after NHP treatment that were

upregulated only in the *sid2-2* mutant and not wild type. Bars represent the negative

log(FDR) of each significantly enriched GO term, bars were omitted if the term was not

significantly enriched or depleted for the defined time point.

1161

Figure 4 Presence of TGA/bZIP and WRKY TF *cis*-regulatory elements (CREs) in the promoters of NHP-upregulated genes in wild type (Col-0) and *sid2-2* seedlings.

A, Schematic of the TF binding motif (i.e., CRE) enrichment analysis performed. All genes with increased transcript abundance upon NHP treatment ($\log_2$(FC) > 0 and P_{adj} < 0.05) were grouped by time point for the indicated genotype, these groups are represented as "test promoter group" in the diagram. CREs of known Arabidopsis TFs from the DAPseq database (O'Malley et al., 2016) were identified in the promoters (1 kb upstream of the transcriptional start site) of each group of promoters. Enrichment of identified CREs was determined relative to CREs found in promoters pulled from a random sampling of genes detected in this RNA-seq experiment. Schematic design is modified from Mariani et al., 2017. Enrichment analysis of TGA/bZIP (B) and WRKY (C) CREs. The name of the TF

1173   known to bind the enriched CRE is listed on the right and the TF family name is indicated

1174   above (i.e., bZIP or WRKY). The -$\log_2$(P value) of the enrichment analysis is indicated by

1175   the scale bar with black indicating no significant enrichment of the CRE and purple to

1176   yellow denoting an enriched CRE.

1177

1178

Figure 5 *WRKY70* is involved in NHP-elicited SAR

Bacterial growth in the distal leaves of mock and NHP treated wild type (Col-0) and *wrky* mutant plants. *WRKY* genes for mutant analysis were selected from the early stable cluster defined in Figure 2 and include *WRKY70* (A), *WRKY38* and *WRKY62* (B), *WRKY54* (C), *WRKY51* (D), and *WRKY59* (E). Three lower (1°) leaves were infiltrated with water (mock), 0.5 mM NHP, or 1 mM NHP (only for *wrky59*) and 1 d later one upper, distal leaf was inoculated with a 1 x 10^5 CFU/mL suspension of *Psm*, followed by quantification of bacterial titer 3 dpi (n = 4-8). Asterisks indicate significant differences in bacterial titer (two-tailed t-test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns = not significant).

1189

1190

Figure 6 Loss of WRKY70 function impacts upregulation of early NHP-responsive genes
A, Diagram of putative WRKY70 CREs, including W-box, WT-box, and WRKY70 (W70)
DAPseq motifs in the promoters (-2 Kb upstream of the transcriptional start site; TSS) of
NHP-responsive genes *BDA1*, *PROSCOOP4*, and *CML10*. Note W70 DAPseq sites
adjacent to a W- or WT-box overlap. Expression of *BDA1* (B), *PROSCOOP4* (C), and
CML10 (D) in *wrky70-1* mutant plants. Three leaves of 4.5-week-old wild type (Col-0) and
*wrky70-*1 mutant plants were infiltrated with water or 1 mM NHP. Samples collected at 0
h were untreated before collection. Transcript abundance was determined relative to
UBC21 for each condition (2$^{-\Delta Ct}$). Asterisks indicate a significant difference between wild

1200   type and *wrky70-1* at each time point (two-tailed t-test; *$P < 0.05$, **$P < 0.01$, ns = not

1201   significant).

1202

1203

Figure 7 NHP enhances ROS production and requires WRKY70 for full ROS levels.

Flg22-elicited ROS quantification in *wrky70-1* plants. Four leaf discs from 4.5 to 5-week-

old wild type and *wrky70-1* plants were pretreated by floating leaf discs on water, 0.1 mM

SA, or 0.5 mM NHP for 24 h before treating with 100 nM flg22 in horseradish peroxidase

and Luminol. Top, traces of the average relative luminescence units (RLU) over the

indicated run time. Each point represents the average of 6 plants (n = 24) from two

experiments. Bottom, total quantification of RLUs from top plot. For each leaf disc, RLUs

from each time point were summed by condition over the run time and averaged with

error bars representing the standard error of the mean. Statistical analysis was performed

using a one-way ANOVA and post hoc Sidak test, different letters indicated statistical

differences between means with $P < 0.05$.

Parsed Citations

Attaran E, Zeier TE, Griebel T, Zeier J (2009) Methyl salicylate production and jasmonate signaling are not essential for systemic acquired resistance in Arabidopsis. Plant Cell 21: 954–971
  Google Scholar: Author Only Title Only Author and Title

Bailey TL, Grant CE (2021) SEA: Simple Enrichment Analysis of motifs. BioRxiv. doi: 10.1101/2021.08.23.457422
  Google Scholar: Author Only Title Only Author and Title

Bailey TL, Johnson J, Grant CE, Noble WS (2015) The MEME Suite. Nucleic Acids Res 43: W39-49
  Google Scholar: Author Only Title Only Author and Title

Bartsch M, Gobbato E, Bednarek P, Debey S, Schultze JL, Bautor J, Parker JE (2006) Salicylic acid-independent ENHANCED DISEASE SUSCEPTIBILITY1 signaling in Arabidopsis immunity and cell death is regulated by the monooxygenase FMO1 and the Nudix hydrolase NUDT7. Plant Cell 18: 1038–1051
  Google Scholar: Author Only Title Only Author and Title

Bauer S, Mekonnen DW, Hartmann M, Yildiz I, Janowski R, Lange B, Geist B, Zeier J, Schäffner AR (2021) UGT76B1, a promiscuous hub of small molecule-based immune signaling, glucosylates N-hydroxypipecolic acid, and balances plant immunity. Plant Cell 33: 714–734
  Google Scholar: Author Only Title Only Author and Title

Bernsdorff F, Döring A-C, Gruner K, Schuck S, Bräutigam A, Zeier J (2016) Pipecolic Acid Orchestrates Plant Systemic Acquired Resistance and Defense Priming via Salicylic Acid-Dependent and -Independent Pathways. Plant Cell 28: 102–129
  Google Scholar: Author Only Title Only Author and Title

Bjornson M, Pimprikar P, Nürnberger T, Zipfel C (2021) The transcriptional landscape of Arabidopsis thaliana pattern-triggered immunity. Nat Plants 7: 579–586
  Google Scholar: Author Only Title Only Author and Title

Blanco F, Salinas P, Cecchini NM, Jordana X, Van Hummelen P, Alvarez ME, Holuigue L (2009) Early genomic responses to salicylic acid in Arabidopsis. Plant Mol Biol 70: 79–102
  Google Scholar: Author Only Title Only Author and Title

Boller T, Felix G (2009) A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. Annu Rev Plant Biol 60: 379–406
  Google Scholar: Author Only Title Only Author and Title

Cai J, Jozwiak A, Holoidovsky L, Meijler MM, Meir S, Rogachev I, Aharoni A (2021) Glycosylation of N-hydroxy-pipecolic acid equilibrates between systemic acquired resistance response and plant growth. Mol Plant 14: 440–455
  Google Scholar: Author Only Title Only Author and Title

Chen Y-C, Holmes EC, Rajniak J, Kim J-G, Tang S, Fischer CR, Mudgett MB, Sattely ES (2018) N-hydroxy-pipecolic acid is a mobile metabolite that induces systemic disease resistance in Arabidopsis. Proc Natl Acad Sci USA 115: E4920–E4929
  Google Scholar: Author Only Title Only Author and Title

Chini A, Fonseca S, Fernández G, Adie B, Chico JM, Lorenzo O, García-Casado G, López-Vidriero I, Lozano FM, Ponce MR, et al (2007) The JAZ family of repressors is the missing link in jasmonate signalling. Nature 448: 666–671
  Google Scholar: Author Only Title Only Author and Title

Cho K-M, Nguyen HTK, Kim SY, Shin JS, Cho DH, Hong SB, Shin JS, Ok SH (2016) CML10, a variant of calmodulin, modulates ascorbic acid synthesis. New Phytol 209: 664–678
  Google Scholar: Author Only Title Only Author and Title

Conrath U, Beckers GJM, Langenbach CJG, Jaskiewicz MR (2015) Priming for enhanced defense. Annu Rev Phytopathol 53: 97–119
  Google Scholar: Author Only Title Only Author and Title

Couto D, Zipfel C (2016) Regulation of pattern recognition receptor signalling in plants. Nat Rev Immunol 16: 537–552
  Google Scholar: Author Only Title Only Author and Title

Ding Y, Sun T, Ao K, Peng Y, Zhang Y, Li X, Zhang Y (2018) Opposite Roles of Salicylic Acid Receptors NPR1 and NPR3/NPR4 in Transcriptional Regulation of Plant Immunity. Cell 173: 1454-1467.e15
  Google Scholar: Author Only Title Only Author and Title

Eulgem T, Rushton PJ, Robatzek S, Somssich IE (2000) The WRKY superfamily of plant transcription factors. Trends Plant Sci 5: 199–206
  Google Scholar: Author Only Title Only Author and Title

Friedrichsen DM, Nemhauser J, Muramitsu T, Maloof JN, Alonso J, Ecker JR, Furuya M, Chory J (2002) Three redundant

brassinosteroid early response genes encode putative bHLH transcription factors required for normal growth. Genetics 162: 1445–1456

Google Scholar: Author Only Title Only Author and Title

Fu ZQ, Dong X (2013) Systemic acquired resistance: turning local infection into global defense. Annu Rev Plant Biol 64: 839–863

Google Scholar: Author Only Title Only Author and Title

Galili T (2015) dendextend: an R package for visualizing, adjusting and comparing trees of hierarchical clustering. Bioinformatics 31: 3718–3720

Google Scholar: Author Only Title Only Author and Title

Gómez-Gómez L, Felix G, Boller T (1999) A single locus determines sensitivity to bacterial flagellin in Arabidopsis thaliana. Plant J 18: 277–284

Google Scholar: Author Only Title Only Author and Title

Gully K, Pelletier S, Guillou M-C, Ferrand M, Aligon S, Pokotylo I, Perrin A, Vergne E, Fagard M, Ruelland E, et al (2019) The SCOOP12 peptide regulates defense response and root elongation in Arabidopsis thaliana. J Exp Bot 70: 1349–1365

Google Scholar: Author Only Title Only Author and Title

Hartmann M, Zeier J (2019) N-hydroxypipecolic acid and salicylic acid: a metabolic duo for systemic acquired resistance. Curr Opin Plant Biol 50: 44–57

Google Scholar: Author Only Title Only Author and Title

Hartmann M, Zeier T, Bernsdorff F, Reichel-Deland V, Kim D, Hohmann M, Scholten N, Schuck S, Bräutigam A, Hölzel T, et al (2018) Flavin Monooxygenase-Generated N-Hydroxypipecolic Acid Is a Critical Element of Plant Systemic Immunity. Cell 173: 456-469.e16

Google Scholar: Author Only Title Only Author and Title

Holmes EC, Chen Y-C, Mudgett MB, Sattely ES (2021) Arabidopsis UGT76B1 glycosylates N-hydroxy-pipecolic acid and inactivates systemic acquired resistance in tomato. Plant Cell 33: 750–765

Google Scholar: Author Only Title Only Author and Title

Holmes EC, Chen Y-C, Sattely ES, Mudgett MB (2019) An engineered pathway for N-hydroxy-pipecolic acid synthesis enhances systemic acquired resistance in tomato. Sci Signal. 12:eaay3066

Google Scholar: Author Only Title Only Author and Title

Hou S, Liu D, Huang S, Luo D, Liu Z, Xiang Q, Wang P, Mu R, Han Z, Chen S, et al (2021) The Arabidopsis MIK2 receptor elicits immunity by sensing a conserved signature from phytocytokines and microbes. Nat Commun 12: 5494

Google Scholar: Author Only Title Only Author and Title

Jaskiewicz M, Conrath U, Peterhänsel C (2011) Chromatin modification acts as a memory for systemic acquired resistance in the plant stress response. EMBO Rep 12: 50–55

Google Scholar: Author Only Title Only Author and Title

Johannesson H, Wang Y, Hanson J, Engström P (2003) The Arabidopsis thaliana homeobox gene ATHB5 is a potential regulator of abscisic acid responsiveness in developing seedlings. Plant Mol Biol 51: 719–729

Google Scholar: Author Only Title Only Author and Title

Kachroo A, Kachroo P (2020) Mobile signals in systemic acquired resistance. Curr Opin Plant Biol 58: 41–47

Google Scholar: Author Only Title Only Author and Title

Kalde M, Barth M, Somssich IE, Lippok B (2003) Members of the Arabidopsis WRKY group III transcription factors are part of different plant defense signaling pathways. Mol Plant Microbe Interact 16: 295–305

Google Scholar: Author Only Title Only Author and Title

Kassambara A (2023) ggpubr: 'ggplot2' Based Publication Ready Plots. R package version 0.6.0. https://rpkgs.datanovia.com/ggpubr/

Google Scholar: Author Only Title Only Author and Title

Kim K-C, Lai Z, Fan B, Chen Z (2008) Arabidopsis WRKY38 and WRKY62 transcription factors interact with histone deacetylase 19 in basal defense. Plant Cell 20: 2357–2371

Google Scholar: Author Only Title Only Author and Title

Kim Y, Gilmour SJ, Chao L, Park S, Thomashow MF (2020) Arabidopsis CAMTA transcription factors regulate pipecolic acid biosynthesis and priming of immunity genes. Mol Plant 13: 157–168

Google Scholar: Author Only Title Only Author and Title

Klessig DF, Choi HW, Dempsey DA (2018) Systemic acquired resistance and salicylic acid: past, present, and future. Mol Plant Microbe Interact 31: 871–888

Google Scholar: Author Only Title Only Author and Title

Kohler A, Schwindling S, Conrath U (2002) Benzothiadiazole-Induced Priming for Potentiated Responses to Pathogen Infection, Wounding, and Infiltration of Water into Leaves Requires the NPR1/NIM1 Gene in Arabidopsis. Plant Physiol 128: 1046–1056
  Google Scholar: Author Only Title Only Author and Title

Lenth RV (2016) Least-Squares Means: the R package lsmeans. J Stat Softw 69: 1–33
  Google Scholar: Author Only Title Only Author and Title

Liu H, Liu B, Lou S, Bi H, Tang H, Tong S, Song Y, Chen N, Zhang H, Jiang Y, et al (2021) CHYR1 ubiquitinates the phosphorylated WRKY70 for degradation to balance immunity in Arabidopsis thaliana. New Phytol 230: 1095–1109
  Google Scholar: Author Only Title Only Author and Title

Li J, Brader G, Kariola T, Palva ET (2006) WRKY70 modulates the selection of signaling pathways in plant defense. Plant J 46: 477–491
  Google Scholar: Author Only Title Only Author and Title

Li J, Brader G, Palva ET (2004) The WRKY70 transcription factor: a node of convergence for jasmonate-mediated and salicylate-mediated signals in plant defense. Plant Cell 16: 319–331
  Google Scholar: Author Only Title Only Author and Title

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15: 550
  Google Scholar: Author Only Title Only Author and Title

Löwe M, Jürgens K, Zeier T, Hartmann M, Gruner K, Müller S, Yildiz I, Perrar M, Zeier J (2023) N-hydroxypipecolic acid primes plants for enhanced microbial pattern-induced responses. Front Plant Sci 14: 1217771
  Google Scholar: Author Only Title Only Author and Title

Luan S, Kudla J, Rodriguez-Concepcion M, Yalovsky S, Gruissem W (2002) Calmodulins and calcineurin B-like proteins: calcium sensors for specific signal response coupling in plants. Plant Cell 14 Suppl: S389-400
  Google Scholar: Author Only Title Only Author and Title

Machens F, Becker M, Umrath F, Hehl R (2014) Identification of a novel type of WRKY transcription factor binding site in elicitor-responsive cis-sequences from Arabidopsis thaliana. Plant Mol Biol 84: 371–385
  Google Scholar: Author Only Title Only Author and Title

Mariani L, Weinand K, Vedenko A, Barrera LA, Bulyk ML (2017) Identification of Human Lineage-Specific Transcriptional Coregulators Enabled by a Glossary of Binding Modules and Tunable Genomic Backgrounds. Cell Syst 5: 187-201.e7
  Google Scholar: Author Only Title Only Author and Title

Mishina TE, Zeier J (2006) The Arabidopsis Flavin-Dependent Monooxygenase FMO1 Is an Essential Component of Biologically Induced Systemic Acquired Resistance. Plant Physiol 141: 1666–1675
  Google Scholar: Author Only Title Only Author and Title

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD (2021) PANTHER version 16: a revised family classification, tree-based classification tool, enhancer regions and extensive API. Nucleic Acids Res 49: D394–D403
  Google Scholar: Author Only Title Only Author and Title

Mohanta TK, Yadav D, Khan A, Hashem A, Tabassum B, Khan AL, Abd Allah EF, Al-Harrasi A (2020) Genomics, molecular and evolutionary perspective of NAC transcription factors. PLoS ONE 15: e0231425
  Google Scholar: Author Only Title Only Author and Title

Mohnike L, Rekhter D, Huang W, Feussner K, Tian H, Herrfurth C, Zhang Y, Feussner I (2021) The glycosyltransferase UGT76B1 modulates N-hydroxy-pipecolic acid homeostasis and plant immunity. Plant Cell 33: 735–749
  Google Scholar: Author Only Title Only Author and Title

Nair A, Goyal I, Voß E, Mrozek P, Prajapati S, Thurow C, Tietze L, Tittmann K, Gatz C (2021) N-hydroxypipecolic acid-induced transcription requires the salicylic acid signaling pathway at basal SA levels. Plant Physiol 187: 2803–2819
  Google Scholar: Author Only Title Only Author and Title

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR (2016) Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. Cell 165: 1280–1292
  Google Scholar: Author Only Title Only Author and Title

Pandey SP, Somssich IE (2009) The Role of WRKY Transcription Factors in Plant Immunity. Plant Physiol 150: 1648–1655
  Google Scholar: Author Only Title Only Author and Title

Pieterse CMJ, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SCM (2012) Hormonal modulation of plant immunity. Annu Rev Cell Dev Biol 28: 489–521
  Google Scholar: Author Only Title Only Author and Title

Rhodes J, Yang H, Moussu S, Boutrot F, Santiago J, Zipfel C (2021) Perception of a divergent family of phytocytokines by the

Arabidopsis receptor kinase MIK2. Nat Commun 12: 705

Google Scholar: Author Only Title Only Author and Title

Rushton PJ, Somssich IE, Ringler P, Shen QJ (2010) WRKY transcription factors. Trends Plant Sci 15: 247–258

Google Scholar: Author Only Title Only Author and Title

Shah J, Kachroo P, Nandi A, Klessig DF (2001) A recessive mutation in the Arabidopsis SSI2 gene confers SA- and NPR1-independent expression of PR genes and resistance against bacterial and oomycete pathogens. Plant J 25: 563–574

Google Scholar: Author Only Title Only Author and Title

Song JT, Lu H, McDowell JM, Greenberg JT (2004) A key role for ALD1 in activation of local and systemic defenses in Arabidopsis. Plant J 40: 200–212

Google Scholar: Author Only Title Only Author and Title

Sun T, Huang J, Xu Y, Verma V, Jing B, Sun Y, Ruiz Orduna A, Tian H, Huang X, Xia S, et al (2020) Redundant CAMTA Transcription Factors Negatively Regulate the Biosynthesis of Salicylic Acid and N-Hydroxypipecolic Acid by Modulating the Expression of SARD1 and CBP60g. Mol Plant 13: 144–156

Google Scholar: Author Only Title Only Author and Title

Thibaud-Nissen F, Wu H, Richmond T, Redman JC, Johnson C, Green R, Arias J, Town CD (2006) Development of Arabidopsis whole-genome microarrays and their application to the discovery of binding sites for the TGA2 transcription factor in salicylic acid-treated plants. Plant J 47: 152–162

Google Scholar: Author Only Title Only Author and Title

Tian H, Zhang Y (2019) The emergence of a mobile signal for systemic acquired resistance. Plant Cell 31: 1414–1415

Google Scholar: Author Only Title Only Author and Title

Truman W, Bennett MH, Kubigsteltig I, Turnbull C, Grant M (2007) Arabidopsis systemic immunity uses conserved defense signaling pathways and is mediated by jasmonates. Proc Natl Acad Sci USA 104: 1075–1080

Google Scholar: Author Only Title Only Author and Title

Wang D, Amornsiripanitch N, Dong X (2006) A genomic approach to identify regulatory nodes in the transcriptional network of systemic acquired resistance in plants. PLoS Pathog 2: e123

Google Scholar: Author Only Title Only Author and Title

Wang Y, Schuck S, Wu J, Yang P, Döring A-C, Zeier J, Tsuda K (2018) A MPK3/6-WRKY33-ALD1-Pipecolic Acid Regulatory Loop Contributes to Systemic Acquired Resistance. Plant Cell 30: 2480–2494

Google Scholar: Author Only Title Only Author and Title

Wildermuth MC, Dewdney J, Wu G, Ausubel FM (2001) Isochorismate synthase is required to synthesize salicylic acid for plant defence. Nature 414: 562–565

Google Scholar: Author Only Title Only Author and Title

Wu Y, Zhang D, Chu JY, Boyle P, Wang Y, Brindle ID, De Luca V, Després C (2012) The Arabidopsis NPR1 Protein Is a Receptor for the Plant Defense Hormone Salicylic Acid. Cell Rep 1: 639–647

Google Scholar: Author Only Title Only Author and Title

Yang Y, Zhang Y, Ding P, Johnson K, Li X, Zhang Y (2012) The ankyrin-repeat transmembrane protein BDA1 functions downstream of the receptor-like protein SNC2 to regulate plant immunity. Plant Physiol 159: 1857–1865

Google Scholar: Author Only Title Only Author and Title

Yan C, Fan M, Yang M, Zhao J, Zhang W, Su Y, Xiao L, Deng H, Xie D (2018) Injury Activates Ca2+/Calmodulin-Dependent Phosphorylation of JAV1-JAZ8-WRKY51 Complex for Jasmonate Biosynthesis. Mol Cell 70: 136-149.e7

Google Scholar: Author Only Title Only Author and Title

Yi SY, Shirasu K, Moon JS, Lee S-G, Kwon S-Y (2014) The activated SA and JA signaling pathways have an influence on flg22-triggered oxidative burst and callose deposition. PLoS ONE 9: e88951

Google Scholar: Author Only Title Only Author and Title

Yildiz I, Gross M, Moser D, Petzsch P, Köhrer K, Zeier J (2023) N-hydroxypipecolic acid induces systemic acquired resistance and transcriptional reprogramming via TGA transcription factors. Plant Cell Environ. doi: 10.1111/pce.14572

Google Scholar: Author Only Title Only Author and Title

Yildiz I, Mantz M, Hartmann M, Zeier T, Kessel J, Thurow C, Gatz C, Petzsch P, Köhrer K, Zeier J (2021) The mobile SAR signal N-hydroxypipecolic acid induces NPR1-dependent transcriptional reprogramming and immune priming. Plant Physiol 186: 1679–1705

Google Scholar: Author Only Title Only Author and Title

Zhang Y, Fan W, Kinkema M, Li X, Dong X (1999) Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the PR-1 gene. Proc Natl Acad Sci USA 96: 6523–6528

Google Scholar: Author Only Title Only Author and Title

Zhang Y, Xu S, Ding P, Wang D, Cheng YT, He J, Gao M, Xu F, Li Y, Zhu Z, et al (2010a) Control of salicylic acid synthesis and systemic acquired resistance by two members of a plant-specific family of transcription factors. Proc Natl Acad Sci USA 107: 18220–18225

Google Scholar: Author Only Title Only Author and Title

Zhang Y, Yang Y, Fang B, Gannon P, Ding P, Li X, Zhang Y (2010b) Arabidopsis snc2-1D activates receptor-like protein-mediated immunity transduced through WRKY70. Plant Cell 22: 3153–3163

Google Scholar: Author Only Title Only Author and Title

Zhou M, Lu Y, Bethke G, Harrison BT, Hatsugai N, Katagiri F, Glazebrook J (2018) WRKY70 prevents axenic activation of plant immunity by direct repression of SARD1. New Phytol 217: 700–712

Google Scholar: Author Only Title Only Author and Title