1   **Title:** Transcriptome analysis reveals role for WRKY70 in early *N*-hydroxy-pipecolic acid
2   signaling
3
4

5   **Authors:** Jessica Foret[1], Jung-Gun Kim[1], Elizabeth S. Sattely[2,3] and Mary Beth
6   Mudgett[1,*]
7
8
9   **Affiliations:**
10   [1]Department of Biology, Stanford University, Stanford, CA 94305, USA
11   [2]Department of Chemical Engineering, Stanford University, Stanford CA 94305, USA
12   [3]Howard Hughes Medical Institute, Stanford University, Stanford CA, 94305, USA
13
14
15   ***Corresponding Author:** Professor Mary Beth Mudgett ([mudgett@stanford.edu)](mailto:mudgett@stanford.edu);
16   Stanford University, 371 Jane Stanford Way, 228A Gilbert Biosciences, Stanford, CA
17   94305-5020

18
19
20   **Short running head:** Early transcriptional changes in response to NHP
21
22
23   **Key words:** N-hydroxy-pipecolic acid, RNA-sequencing, WRKY transcription factors,
24   salicylic acid, systemic acquired resistance, Arabidopsis thaliana, WRKY70

Abstract   (250 words)

N-hydroxy-pipecolic acid (NHP) is a mobile metabolite essential for inducing and amplifying systemic acquired resistance (SAR) following pathogen attack. Early phases of NHP signaling leading to immunity have remained elusive. Here we report the early transcriptional changes mediated by NHP and the role salicylic acid (SA) plays during this response. We show that distinct waves of expression within minutes to hours of NHP treatment include increased expression of WRKY transcription factors as the primary transcriptional response, followed by the induction of WRKY-regulated defense genes as the secondary response. The majority of genes induced by NHP within minutes were SA-dependent, whereas those induced within hours were SA-independent. These data suggest that NHP induces the primary transcriptional response in a low SA environment and new SA biosynthesis is dispensable for induction of the secondary transcriptional response. We demonstrate that WRKY70 is required for the induced expression of a set of genes defining some of the secondary transcriptional response, SAR protection, and NHP-dependent enhancement of ROS production in response to flagellin treatment. Taken together, our study highlights the key genes and pathways defining early NHP responses and a role for WRKY70 in the regulation of NHP-dependent transcription.

Introduction

Systemic acquired resistance (SAR) is a whole-plant defense response triggered by an initial pathogen attack. During the establishment of SAR, mobile signals are sent from infected tissue to uninfected tissue, initiating the priming of immune responses and enhancing immunity against secondary pathogen infections (Fu and Dong, 2013; Tian and Zhang, 2019; Kachroo and Kachroo, 2020). To achieve a primed state, plants execute substantial transcriptional reprogramming of several hundred to a few thousand genes, resulting in the increased accumulation of defense hormone salicylic acid (SA), sensing and signaling components for pathogen detection, and production of antimicrobial compounds (Jaskiewicz et al., 2011; Conrath et al., 2015; Bernsdorff et al., 2016). In addition, several small molecules produced by plants during infection help to orchestrate this defense priming, including SA and the bioactive metabolite *N*-hydroxy-pipecolic acid (NHP) (Chen et al., 2018; Hartmann et al., 2018; Klessig et al., 2018).

NHP is a mobile metabolite that can move from one leaf to another to mediate long-distance cellular communication (Chen et al., 2018; Mohnike et al., 2021; Yildiz et al., 2021). In *Arabidopsis thaliana* (Arabidopsis), the enzyme FLAVIN-DEPENDENT MONOOXYGENASE 1 (FMO1) catalyzes the *N*-hydroxylation of lysine-derived pipecolic acid (Pip) in response to pathogen infection to produce NHP (Chen et al., 2018; Hartmann et al., 2018). Notably, *fmo1* null mutants are impaired in local disease resistance during pathogen infection and fail to establish SAR (Chen et al., 2018; Hartmann et al., 2018). Treatment of *fmo1* mutant plants with exogenous NHP is sufficient to induce systemic resistance against bacterial and oomycete pathogens (Chen et al., 2018; Hartmann et al., 2018). Furthermore, NHP treatment of a single leaf is sufficient to initiate and amplify defense signaling throughout the plant, resulting in the transcriptional activation of a large subset of SAR-associated genes within 24 h of treatment (Yildiz et al., 2021; Yildiz et al., 2023), including genes encoding the SA biosynthetic enzyme ISOCHORISMATE SYNTHASE 1/SA-INDUCTION DEFICIENT 2 (ICS1/SID2), the Pip biosynthetic enzymes AGD2-LIKE DEFENSE RESPONSE PROTEIN 1 (ALD1) and SAR-DEFICIENT 4 (SARD4), as well as its own biosynthetic

enzyme FMO1 (Chen et al., 2018). Collectively, these findings highlight the importance

of NHP in initiating immune signaling to establish local and systemic plant defenses.

Induction of *ICS1* transcription by NHP signaling results in the accumulation of SA in

both local and systemic tissues (Chen et al., 2018; Hartmann et al., 2018). Elevation of

SA in turn amplifies SAR-associated transcriptional changes (Bernsdorff et al., 2016).

Signaling downstream of SA accumulation is largely driven by the action of the SA

receptor and transcriptional coactivator NONEXPRESSER OF PR GENES 1 (NPR1),

which interacts with a number of TGA transcription factors (TFs) from the basic leucine

zipper (bZIP) family to alter gene expression (Zhang et al., 1999; Wang et al., 2006; Wu

et al., 2012). Members of the biotic and abiotic stress-response WRKY TF family are

among the known targets of the NPR1/TGA interaction (Thibaud Nissen et al., 2006;

Wang et al., 2006). WRKY TFs have been identified as key regulators of NPR1-

dependent and NPR1-independent SAR responses. Additionally, some WRKY

members play important roles in feedback regulation of transcription (Pandey and

Somssich, 2009; Fu and Dong, 2013). For example, WRKY54 and WRKY70 modulate

SA biosynthesis through repression of *ICS1* expression (Wang et al., 2006).

The interplay between SA and NHP appears to be interdependent and synergistic with

both metabolites being able to upregulate one another's biosynthetic pathways (Chen et

al., 2018; Hartmann et al., 2018; Sun et al., 2020). Notably, elevated endogenous levels

of Pip, the precursor of NHP, primes the expression of both NHP and SA biosynthetic

genes under low concentrations of SA by stabilizing NPR1 levels (Kim et al., 2020). This

points to an intriguing dynamic relationship between SA and NHP over the course of

SAR activation. Evidence suggests NHP accumulates to detectable levels in systemic

tissues roughly 24 h before SA accumulation is detected (Hartmann and Zeier, 2019). In

such a case, NHP would need to drive systemic signaling for a substantial period of

time before SA could further amplify SAR responses. It is currently unclear how the

levels of SA in leaves impact early NHP signaling. Furthermore, little is known about the

very early phase of NHP signaling in the tissues where it is first produced or in the distal

tissues once NHP arrives there. Key players of NHP signaling and the progression of

NHP-dependent transcriptional reprogramming require further investigation.

Presently, the majority of genome-wide transcriptomic studies of SAR have been carried

out on the timescale of 1-2 days post induction; this includes pharmacological studies

using Pip and NHP treatment (Bernsdorff et al., 2016; Hartmann et al., 2018; Yildiz et al.,

2021; Yildiz et al., 2023). These data have revealed that NHP ultimately activates SAR

in untreated tissue, but do not shed light on early signaling events that lead to the

observed SAR phenotype. The immediate consequences of NHP accumulation on gene

expression have not been described. Studying very early transcriptional changes, within

minutes to hours of elicitor treatment, has been an effective strategy elucidating key

components of signaling pathways for SA, jasmonic acid (JA), and brassinosteroid

(Friedrichsen et al., 2002; Chini et al., 2007; Ding et al., 2018). Thus, in an effort to better

understand NHP's primary action in signal transduction, we determined the genome-

wide transcriptional responses in Arabidopsis from 15 minutes up to 6 hours after

treatment with exogenous NHP.

Here we report the early NHP transcriptional profile and the role SA plays during this

response. We present evidence indicating that increased expression of WRKY TFs

define the primary transcriptional responses mediated by NHP within minutes (i.e. 15 to

30 minutes) post-treatment, and WRKY TFs act as executors of the secondary

transcriptional responses within hours (i.e. 3 to 6 hours). The majority of genes induced

by NHP within minutes were SA-dependent, whereas those induced within hours were

SA-independent. These data suggest that NHP induces the primary transcriptional

response in a low SA environment and new SA biosynthesis is dispensable for induction

of the secondary transcriptional response. We also show that WRKY70 is required for

NHP induced transcription of a set of genes defining secondary transcriptional changes

as well as NHP enhancement of reactive oxygen species (ROS) and SAR. Taken

together, our study highlights the key genes and pathways defining early NHP

responses and a role for WRKY70 in the regulation of NHP-dependent transcription.

Results

NHP induces expression of defense-associated genes within minutes

Previous studies in Arabidopsis have demonstrated that exogenous application of NHP
to a leaf (defined as "local tissue") is sufficient to initiate and amplify defense signaling
in untreated leaves (defined as "distal tissue"), including the transcriptional activation of
SAR marker genes (Chen et al., 2018; Hartmann et al., 2018; Nair et al., 2021; Yildiz et
al., 2021). Notably, these studies focused on gene expression changes in distal tissues
at late time points (24 to 48 h) following NHP elicitation. However, it was unknown which
gene expression changes occur immediately in response to NHP treatment. We
hypothesized that once present in the tissue, either via endogenous synthesis or
exogenous application, NHP initiates signaling that leads to transcriptional changes on
a timescale of minutes to several hours, which are essential for execution of NHP-
dependent immune responses.

In order to test this hypothesis, we set out to identify NHP responsive transcriptional
markers and determine the earliest time frame when NHP could induce these genes.
We first questioned if NHP could induce the expression of classical SAR-associated
marker genes (i.e., *FMO1*, *ICS1*, and *PATHOGENESIS-RELATED PROTEIN 1* (*PR1*))
within several hours of treatment. Studies have shown *FMO1*, *ICS1*, and *PR1* are
induced between 8 to 48 h following NHP treatment (Chen et al., 2018; Nair et al.,
2021). Therefore, we began by testing the response to NHP within 24 h of treatment.
The leaves of 4.5-week-old Arabidopsis Col-0 (wild type) plants were infiltrated with 1
mM NHP and then treated leaves were collected for mRNA isolation at 2, 6 and 24 h.
Quantitative real-time PCR (qRT-PCR) analysis showed significantly increased
transcript abundance of all three genes in NHP treated leaves at 24 h but not at 2 and 6
h compared to water (mock) treated leaves (Fig. 1A). These data indicate that *FMO1*,
ICS1, and *PR1* are responsive to NHP treatment but are likely not primary targets of
NHP transcriptional activation.

Next, we tested the ability of NHP to induce expression of *WRKY38* and *UGT76B1*, two
genes known to respond to SA within 2 h of treatment (Blanco et al., 2009). *WRKY38*
encodes a defense-associated TF and *UGT76B1* encodes a UDP-dependent
glycosyltransferase shown to glycosylate SA and NHP (Eulgem et al., 2000; Bauer et
al., 2021; Cai et al., 2021; Holmes et al., 2021; Mohnike et al., 2021).   We   found   a
significant increase of mRNA abundance for both *WRKY38* and *UGT76B1* in NHP
treated leaves at 2 h compared to mock treated leaves (Fig. 1A). *WRKY38* transcript
levels were also elevated at 6 and 24 h after treatment with NHP (Fig. 1A). These data
indicate that both *WRKY38* and *UGT76B1* are early NHP responsive transcripts.

We used the *WRKY38* gene as a marker to define the earliest time when NHP-induced
transcriptional changes could be measured. To this end, we implemented a
hydroponically grown seedling assay to synchronize chemical treatment and increase
the number of plants sampled, while minimizing mechanical stress. Wild type seedlings
were grown in liquid medium in six-well plates with 15-18 seeds per well. The seedlings
were treated with mock, 1 mM Pip, or 1 mM NHP at 10-days-post germination and then
collected at 15 min, 30 min, 3 h, and 6 h post treatment to isolate mRNA (Fig. 1B).
WRKY38 transcript abundance was significantly elevated in Arabidopsis seedlings
treated with NHP as early as 15 min, peaking in abundance at 30 min, when compared
to those treated with mock or Pip (Fig. 1C). This trend continued through 3 h following
NHP treatment. The NHP precursor Pip did not elicit an increase in *WRKY38* mRNA
within 3 h but showed a slight trend of increased abundance at the 6 h time point (Fig.
1C). These results indicate that exogenous NHP, but not Pip, is sufficient to elevate
WRKY38 transcript abundance within 15 minutes of application.

Since *WRKY38* expression is known to be induced by SA treatment (Blanco et al.,
2009), we questioned if NHP-induced expression of *WRKY38* is dependent on SA
derived from the ICS1/SID2 biosynthetic pathway by utilizing the well characterized
sid2-2 mutant that does not accumulate SA upon pathogen infection (Wildermuth et al.,
2001). Wild type and *sid2-2* seedlings were grown hydroponically, treated with mock, 1
mM Pip, or 1 mM NHP, and then collected 3 h after treatment. NHP treatment increased

WRKY38 mRNA abundance in the *sid2-2* mutant relative to mock treatment, similar to

wild type seedlings (Supplemental Fig. S1). These findings indicate that NHP is

sufficient to alter *WRKY38* transcript abundance in the absence of SA accumulation,

defining *WRKY38* as an early NHP induced, SA-independent transcript.

The early NHP-responsive transcriptome reveals distinct waves of expression

We next questioned how the presence of NHP impacts whole-genome transcriptional

changes within minutes to hours of treatment in a SA-dependent and -independent

manner. To address this, Arabidopsis wild type and *sid2-2* mutant seedlings were grown

in the same hydroponic design as previously described for qRT-PCR experiments (Fig.

1B). The *sid2-2* mutant was analyzed in order to identify transcriptional changes

dependent on NHP-induced SA biosynthesis (Chen et al., 2018). Each pool of seedlings

was treated with mock or 0.5 mM NHP for 15 min, 30 min, 3 h and 6 h. A reduced

concentration of NHP was used as we found that 0.5 mM NHP induced similar *WRKY38*

mRNA accumulation as 1 mM NHP (Supplemental Fig. S2). The abundance of mRNA

in the treated seedlings was assessed for each condition via RNA-sequencing (RNA-

seq). After general quality control and mapping to the Arabidopsis TAIR10 genome,

genes differentially expressed by NHP were determined relative to mock treatment for

each time point and genotype using DESeq2 (Love et al., 2014). Genes were

considered differentially expressed if they returned an adjusted *P*-value below 0.05 (P_{adj}

< 0.05).

In wild type seedlings, a total of 2,079 genes were differentially expressed in response

to NHP for at least one of the measured time points. To explore the set of early NHP-

responsive genes, we first selected for genes showing a robust fold change over mock

by applying the cutoff of $\log_2$(fold change, FC) > 1 for upregulated genes and $\log_2$(FC) <

-1 for downregulated genes. This identified 352 genes that were differentially expressed

in response to NHP for at least one time point, with 163 genes up and 189 genes down

(Supplemental Table S3). Next, the upregulated genes were hierarchically clustered

based on their $\log_2$(FC) values over the indicated time points and four well-defined gene

clusters were identified (Fig. 2A). For each group of genes in the defined clusters, the

log$_2$(FC) values were then averaged by time point, revealing dynamic patterns of
expression. Two of the clusters were upregulated within minutes (designated as 'early
transient' and 'early strong') and two were upregulated by several hours (designated as
'late weak' and 'late strong') (Fig. 2B, Table 1, Table 2, Supplemental Table S4).

The 'early transient' cluster includes 42 genes and is defined by elevated transcript
abundance 15-30 min after NHP treatment that did not persist by 3 h. Gene ontology
(GO) analysis for this cluster showed strong enrichment for biological processes
associated with responses to jasmonic acid (JA), wounding, and oxidative stress. Some
of these genes are associated with the regulation of defense as well as defense
responses to fungi and bacteria (Fig. 2C). Among the most highly expressed genes in
this cluster are JA-responsive genes, which play key roles in the regulation of wounding
responses, including *CML39* which encodes a calmodulin-like protein, *JAZ10* which
encodes a jasmonate-zim-domain protein, and *MYC2* which encodes a jasmonate-
inducible transcription factor (Table 1, Table 2). Taken together, the early yet
unsustained induction of these genes within 30 min suggests a burst of JA- and wound-
responsive gene expression that acts as a primary wave of NHP signaling.

The 'early strong' cluster includes 10 genes and is defined by a high average log$_2$(FC)
15 min after NHP treatment that was maintained for all subsequent time points. Of the
10 genes, two (AT1G06475 and AT5G40980) were strongly upregulated by NHP at 30
min and no other time points. These two genes were removed from the average
log$_2$(FC) analysis and subsequent analyses because they did not behave like the other
genes in this cluster (Fig. 2B). The early strong cluster is notably enriched in biological
processes involved in the regulation of transcription, with over half the genes belonging
to the WRKY TF family, including four group III *WRKY* genes (*WRKY38*, *WRKY54*,
WRKY62 and *WRKY70*) and two group II-c genes (*WRKY51* and *WRKY59*) (Eulgem et
al., 2000; Kalde et al., 2003); (Table 1, Table2, Supplemental Table S4). Additional GO
terms enriched in the early strong cluster includes regulation of defense and regulation
of hormonal signaling pathways such as SA, JA, brassinosteroid, and ethylene (Fig.

2C). The early and sustained induction of these genes, particularly the *WRKYs*, suggests they are primary transcriptional target genes of NHP signaling.

The 'late weak' cluster includes the largest number of genes (96) that averaged a $\log_2$(FC) around 1 (Fig. 2B). GO term analysis of the cluster revealed significant enrichment of biological processes involved in response to bacteria and oomycetes, as well as SA-mediated signaling and regulation of SAR (Fig. 2C). This indicates NHP activation of SAR and pathogen defense genes can be detected as early as 3-6 h in treated tissues. Notably, genes involved in the regulation and biosynthesis of NHP were observed in this wave of transcription, including *SARD1* (*SAR DEFICIENT 1*), *NIMIN-1* (*NIM-INTERACTING 1*), *ALD1,* and *UGT76B1* (Table 1, Table 2, Supplemental Table S4) (Song et al., 2004; Zhang et al., 2010a; Holmes et al., 2021).

The 'late strong' cluster consists of 15 genes with $\log_2$(FC) values averaging between 3.5 - 4 (Fig. 2B). Response to SA was the only significantly enriched GO term for this cluster (Fig. 2C). Despite having a role in the response to SA, the majority of genes classified as late strong were upregulated by NHP in both wild type and *sid2-2* (Fig. 3A, Table 1), indicating that accumulation of these transcripts was at least partially SA-independent. Of the 14 SA-independent genes identified in this cluster, only five have been found to be upregulated 24 h after NHP treatment in an SA-independent manner (Yildiz et al., 2021). Thus, our analysis has uncovered additional NHP-responsive genes that operate independently of SA as early as 6 h post NHP signaling.

We also performed a similar clustering analysis on the 189 NHP-downregulated transcripts and identified three gene clusters (Supplemental Fig. S3A, B). In contrast to the upregulated gene clusters, the downregulated gene clusters had less distinct patterns of expression and averaged $\log_2$(FC) values around -1 to -1.5. Two clusters were downregulated within 15-30 min and persisted 3-6 h hours after NHP treatment. One of these showed weaker downregulation and was classified as an 'early weak' cluster. The other showed stronger downregulation and was classified as an 'early strong' cluster. The third did not show decreased transcript abundance until 3-6 h after

treatment and thus was labeled as a 'late' cluster (Supplemental Fig. S3B). All three
clusters were enriched for biological processes involving growth and development such
as root morphogenesis, plant epidermis development, plant-type cell wall organization,
and root hair cell differentiation (Supplemental Fig. S3C). Given the NHP precursor Pip
is known to inhibit root growth of seedlings (Wang et al., 2018), it is likely that NHP also
inhibits root growth. These clusters shed new light on the genes that may be associated
with altered plant growth and development in response to elevated levels of NHP.

SA biosynthesis is dispensable for the early induction of a majority of NHP
upregulated genes
Interrogation into the late strong cluster suggested NHP may be sufficient to activate a
subset of SA-responsive genes in the absence of NHP-induced SA biosynthesis (Fig.
2C, Fig. 3A, 'late strong'). This was an intriguing possibility given NHP likely
accumulates in distal tissues prior to SA accumulation following a local tissue infection
(Bernsdorff et al., 2016; Hartmann et al., 2018; Hartmann and Zeier, 2019). We therefore
hypothesized that NHP would need to drive transcriptional reprogramming as SA levels
rise from low to high. To investigate this, we assessed the impact of SA biosynthesis on
NHP-induced gene expression from 15 min to 6 h. For each time point of the wild type
gene clusters described in Fig. 2, the percentage of genes upregulated in both wild type
and the *sid2-2* mutant was determined. This group of genes was defined as 'SA-
independent'. Genes upregulated only in wild type and not in *sid2-2* at the defined time
point were conversely defined as 'SA-dependent'.

Notably, the majority of genes upregulated by NHP at 15 min in the early transient
cluster were SA-dependent. For example, 87% of the genes (27 out of 31) were
upregulated at 15 min in wild type but not in *sid2-2* seedlings (Fig. 3A) and similar
trends were seen at 30 min. By contrast, 100% of the genes (8 out of 8) in the early
strong cluster upregulated 3-6 h after NHP treatment were SA-independent (Fig. 3A).
69% of the genes (36 out of 52) in the late weak cluster were SA-independent at 3h,
which dropped to 53% (37 out of 70 genes) at 6 h (Fig. 3A).

A similar pattern of gene expression was observed when we analyzed the total number
of NHP upregulated genes reaching a $\log_2$(FC) > 1 without gene clustering (Fig. 3B).
The majority of genes induced by NHP between 15-30 min were SA-dependent (i.e., 36
genes up in wild type and 9 up in both wild type and *sid2-2* at 15 min) and the majority
of genes induced by NHP between 3-6 h were SA-independent (i.e., 20 genes up in wild
type and 55 up in both wild type and *sid2-2* at 3 h) (Fig. 3B). Taken together, this
analysis highlights an important temporal role of SA on NHP induced transcription. Our
data suggest that new SA biosynthesis is required for the induction of NHP-responsive
genes within minutes of exposure to NHP, whereas SA biosynthesis is dispensable for
NHP-induced expression several hours later.

We also analyzed the downregulated genes using the same approach and found a
slight trend of SA-dependence for these genes following NHP treatment. For example,
the early weak and late clusters defined in Supplemental Fig. S3 showed that a majority
of genes are downregulated in wild type but not the *sid2-2* mutant at each time point
(Supplemental Fig. S4A). By contrast, the genes of the early strong cluster were less
dependent on SA biosynthesis, such that by the 6 h time point only 15% of genes (8 out
of 44) were downregulated in wild type and not in *sid2-2* seedlings (Supplemental Fig.
S4A). Independent of clustering, greater than 60% of the downregulated genes were
found to be SA-dependent at 15 min and 3 h and approximately 40% were SA-
dependent at 30 min and 6 h (Supplemental Fig. S4B). Taken together, these findings
suggest a stronger dependence on SA biosynthesis for NHP-dependent downregulation
of gene expression over the time courses studied.

SA biosynthesis antagonizes NHP-elicited gene expression at 3 to 6 h
Looking at the genes upregulated only in the *sid2-2* mutant and not wild type, we
discovered an unexpectedly large number of transcripts that increased in abundance
upon NHP treatment at 3 and 6 h post treatment (Fig. 3B). This includes 151 genes at 3
h and 98 genes at 6 h that were significantly upregulated by NHP in the *sid2-2* mutant
but not in wild type (Fig. 3B). GO-term enrichment analysis of these genes showed
significant enrichment for a number of biotic and abiotic stress responses, such as

defense response to bacteria and fungi, SAR, response to abscisic acid, and response

to hypoxia (Fig. 3C). These data suggest that SA may play an important role in

modulating the transcription of a suite of genes, through an unknown mechanism, once

NHP-mediated transcriptional reprogramming has been initiated.

TGA and WRKY transcription factor *cis*-regulatory elements are enriched in the

promoters of early NHP upregulated genes

To identify TFs that might directly control the expression of early NHP-responsive

genes, we analyzed the promoter regions of all genes upregulated after NHP treatment

in both wild type and *sid2-2* mutant seedlings to identify putative TF bindings sites (i.e.,

cis-regulatory elements, CREs). A less stringent cutoff of $\log_2$(FC) > 0 and P_{adj} < 0.05

was used for the upregulated genes to capture putative transcriptional regulators of all

genes significantly elevated in transcript abundance, regardless of the magnitude of

expression change. Following a modified workflow detailed by Bjornson et al., 2021, the

NHP upregulated genes were then separated by the time points in which they were

differentially expressed. Enrichment of transcription factor CREs was then determined

for each time point using the DAPseq database (O'Malley et al., 2016) (Fig. 4A).

Supplemental Fig. 5 shows all the CREs found to be significantly enriched in the

promoters of NHP-upregulated genes in both wild type and *sid2-2* seedings between 15

min and 6 h following NHP treatment.

We found that CREs for bZIP family members, specifically TGA TFs, were enriched in

the promoters of genes upregulated by NHP as early as 15 min. For some TGAs (e.g.,

TGA2, TGA5, and TGA7), the enrichment of their TF binding sites was observed for all

time points (Fig. 4B). Enrichment of TGA binding sites in the promoters of early NHP

upregulated genes is consistent with previous work demonstrating TGA2/5/6 and

TGA1/4 are required for the induction of NHP responsive genes at 24-48 h post NHP

treatment (Nair et al., 2021; Yildiz et al., 2023). TGA TFs are known to interact with the

transcriptional coactivator NPR1 to drive SAR- and SA-induced transcriptional changes

(Zhang et al., 1999). The enrichment of TGA binding sites in the promoters of NHP

upregulated genes as early as 15 min implicates TGAs and NPR1 among the primary

transcriptional regulators of NHP signaling. Furthermore, enrichment of TGA CREs was

not observed in *sid2-2* seedlings until 30 min and was also not significantly enriched at

3 h (Fig. 4B). These findings suggest that SA biosynthesis and likely elevated SA levels

are required for rapid and stable induction of some TGA-regulated genes induced by

NHP signaling.

Moreover, we discovered that a large number of WRKY CREs were highly enriched in

the promoter regions of NHP upregulated genes at 3-6 h in both wild type and *sid2-2*

seedlings (Fig. 4C). This correlates with the majority of genes identified in the NHP

upregulated early strong cluster belonging to the WRKY TF family (Fig. 2, Table 1). This

finding suggests a model where NHP induces the expression of *WRKY* genes as a

primary transcriptional response, followed by WRKY-regulated gene expression

changes that act as a secondary wave of transcriptional control.

It is striking that the confidence in WRKY CRE enrichment was greater (i.e., lower *P*

value) at 3 and 6 h in the promoters of genes upregulated in the *sid2-2* mutant

compared to those of wild type (Fig. 4C). These data suggest that a substantial number

of NHP upregulated genes are directly regulated by WRKY TFs, and such regulation is

influenced by SA levels. A similar pattern emerged for members of the NAC TF family,

where NAC CREs were enriched in the promoters of NHP upregulated genes at 3 h in

sid2-2 but not in the promoters of genes upregulated in wild type (Supplemental Fig.

S5). Both WRKY and NAC TFs are involved in the regulation of a number of biotic and

abiotic stress responses (Eulgem et al., 2000; Mohanta et al., 2020). Notably, the set of

genes upregulated by NHP at 3-6 h only in the *sid2-2* mutant but not in wild type were

enriched in biological processes associated with biotic and abiotic stress responses

(Fig. 3B, C). Taken together, these data suggest SA may antagonize the transcription of

a subset of NHP responsive genes by regulating the activities of WRKY and NAC TFs.

The same CRE enrichment analysis was also carried out on all NHP downregulated

genes, regardless of the magnitude of fold change ($\log_2$(FC) < 0 and P_{adj} < 0.05). We

found multiple WRKY CREs enriched in the promoter regions of NHP downregulated

genes at 15 min, 3 h, and 6 h, in wild type seedlings (Supplemental Fig. S6), further

supporting a role for WRKY TFs in relaying NHP-induced transcriptional changes.

Finally, we found the CREs of Homeobox gene family members to be abundant in the

promoters of NHP downregulated genes spanning from 15 min through 6 h, with the

greatest confidence of enrichment found in the promoters of genes downregulated by

NHP at 3-6 h in an SA-dependent manner (Supplemental Fig. S6). Among these was

ATHB5, a homeodomain leucine zipper (HDZip) protein linked to the abscisic acid

(ABA)-driven repression of germination and root growth in seedlings (Johannesson et

al., 2003). Given that NHP precursor Pip inhibits seedling root growth, we now

hypothesize that NHP may drive the repression of root growth and development genes

(Supplemental Fig. 3C) through the action of HDZip Homeobox proteins like ATHB5.

Early NHP-responsive gene *WRKY70* is required for NHP-elicited SAR

To further dissect the NHP response pathway, we carried out a reverse genetic screen

to identify Arabidopsis mutants with compromised NHP-elicited SAR. Given WRKY

CREs were the predominant regulatory elements identified in genes responsive to NHP

3-6 h following treatment (Fig. 4C, Supplemental Fig. S6), we selected the six *WRKY*s

(*WRKY38, WRKY51, WRKY54, WRKY59, WRKY62, WRKY70)* from the early strong

NHP upregulated cluster (Figure 2B, Table 1, Table 2) for mutant analysis.

To measure NHP-induced SAR, we obtained homozygous Arabidopsis mutants and

quantified bacterial growth in their leaves after treatment with exogenous NHP (Chen et

al., 2018). Specifically, three lower leaves of 4.5-week-old wild type and mutant plants

were infiltrated with water (mock) or 0.5 mM NHP. One day later, one distal upper leaf

was inoculated with a 1×10^5 CFU/mL suspension of the virulent bacterium

Pseudomonas syringae pathovar *maculicola* strain ES4326 (*Psm*). Bacterial growth of

Psm was quantified three days post infection (dpi) to determine if the mutants were

altered in their resistance to pathogen infection. Mutants were considered to exhibit

NHP-elicited SAR if bacterial titer was significantly lower in NHP treated plants

compared to mock treated plants. By contrast, mutants were considered insensitive or

partially insensitive to NHP if the bacterial titer of NHP treated mutants was higher than

NHP treated wild type plants.

Of the six *wrky* mutants screened, five (*wrky38*, *wrky62*, *wrky54*, *wrky51*, and *wrky59*)

exhibited normal NHP-elicited SAR (Fig. 5B-E). Only one, *wrky70*-1 (T-DNA insertional

line SALK_025198) (Li et al., 2006), displayed compromised NHP-elicited SAR (Fig.

5A). That is, NHP treatment did not decrease bacterial growth in distal leaves of

wrky70-1 plants compared to mock treatment. We also observed that *wrky70-1* mutants

treated with mock contained less bacteria than wild type plants treated with mock,

demonstrating that basal resistance is enhanced in the *wrk70-1* mutant (Fig. 5A).

Notably, distal leaves of *wrky70-1* plants treated with NHP contained higher levels of

bacteria than distal leaves of wild type plants treated with NHP (Fig. 5A). The inability of

wrky70-1 plants to further restrict bacterial growth following NHP treatment suggests

that the mutant may be insensitive to NHP or defective in NHP signaling. Taken

together, these data indicate that *WRKY70* is required for full NHP-elicited systemic

resistance.

We next questioned if systemic resistance to *Psm* was similarly compromised in

wrky70-1 plants primed with the avirulent pathogen *P*. *syringae* pathovar *tomato* (*Pst*)

strain DC3000 carrying *avrRpt2* (*Pst avrRpt2*). Localized infection with *Pst avrRpt2* is

known to elicit a strong immune response that leads to SAR and protection against *Psm*

in distal leaves (Kohler et al., 2002). We found that the *Psm* titer in distal leaves reached

similar levels for *wrk70-1* plants treated with mock or *Pst avrRpt2* and this level was

comparable to *Psm* growth in wild type leaves of plants treated with *Pst avrRpt2*

(Supplemental Fig. S7A). These data indicate that *wrky70-1* mutants exhibit a basal-

level of defense priming and *Pst avrRpt2* primary infection does not further enhance this

resistance.

We also examined pathogen growth in local *wrky70-1* leaves. We found that the titer of

Psm in infected *wrky70-1* leaves was lower than in *Psm* infected wild type leaves

(Supplemental Fig. S7B). This phenotype corroborates previous work demonstrating

loss of *WRKY70* expression enhances resistance to *Psm* infection (Zhou et al., 2018).

Expression of *SARD1* and *PR* genes is elevated in *wrky70-1* independent of NHP

Previous studies in Arabidopsis have demonstrated a role for WRKY70 in the

transcriptional regulation of several genes required for SAR signaling and disease

resistance, including *SARD1* and the *PATHOGENESIS-RELATED* genes *PR1*, *PR2*,

and *PR5* (Li et al., 2004; Zhou et al., 2018; Liu et al., 2021). These genes were also

shown to be upregulated in wild type Arabidopsis plants 24 h after treatment with NHP

(Yildiz et al., 2021). To determine if WRKY70 is required for NHP induced expression of

SARD1, *PR1*, *PR2*, and *PR5* at 24 h, we quantified transcript abundance in wild type

and *wrky70-1* plants treated with water or 1 mM NHP. As expected, qRT-PCR showed

increased abundance of *SARD1*, *PR1*, *PR2*, and *PR5* mRNA in wild type plants treated

with NHP compared to water (Supplemental Fig. S7C). However, NHP treatment did not

further increase transcript levels in *wrky70-1* plants (Supplemental Fig. S7C). Notably,

all four genes showed elevated transcript abundance in water treated *wrky70-1* mutants

compared to water treated wild type. The elevated expression of *SARD1* and *PR1* in

water treated *wrky70-1* plants is consistent with previous studies demonstrating

WRKY70 negatively regulates these two genes in the absence of a pathogen (Zhou et

al., 2018; Liu et al., 2021).

WRKY70 is required for NHP induction of genes in the late weak and late strong

clusters

Given *wrky70-1* mutants exhibited enhanced basal resistance but were still unable to

achieve the same level of resistance as NHP treated wild type plants (Figure 5A), we

hypothesized that a subset of early NHP-responsive genes containing putative

WRKY70 CREs in their promoters may be dependent on WRKY70 function for full, wild

type gene expression levels following NHP treatment. To investigate this, we selected

three genes from the late weak and late strong clusters (Fig. 2, Table 1), specifically

BDA1, *PROSCOOP4*, and *CML10,* and examined their transcript accumulation in

response to 1 mM NHP over a time course of 0 to 12 h. The promoters of these genes

contain one or more WRKY70 CREs (i.e., W-box (TTGACY), WT-box (YGACTTTT), and WRKY70 DAPseq motif; (Rushton et al., 2010; Machens et al., 2014; O'Malley et al., 2016)) within 2000 base pairs upstream of their respective putative transcriptional start site (Fig. 6A).

We found that *BDA1* (*bian da*; "becoming big" in Chinese), a gene encoding an ankyrin-repeat transmembrane protein (Yang et al., 2012), exhibited the strongest dependence on WRKY70 for NHP-induced expression. *BDA1* transcripts were significantly increased at 6, 9, and 12 h post NHP treatment in wild type leaves but not *wrk70-1* leaves (Fig. 6B). Transcript levels for *PROSCOOP4*, a gene encoding the precursor of secreted peptide SCOOP4/STMP10 (Gully et al., 2019; Hou et al., 2021), were also significantly higher at 6 and 9 h post NHP treatment in wild type leaves compared to that observed for *wrk70-1* plants (Fig. 6C). Unlike *BDA1*, we found that *PROSCOOP4* transcripts were higher in NHP-treated *wrk70-1* leaves at 6 and 9h compared to the 0h timepoint (Fig. 6C). These data show that *wrk70-1* plants responded to NHP treatment, suggesting that they can sense NHP but are impaired in NHP signaling. Similar trends were observed for *CML10* (*CALMODULIN-LIKE 10*/*CaBP22*; (Luan et al., 2002)) at the 9 h timepoint (Fig. 6D). We confirmed these results by analyzing a second *wrky70* allele, *wrky70-2* (Supplementary Fig. S8).

Taken together, our findings reveal that *WRKY70* is required for the proper expression of NHP responsive genes from the late weak and late strong clusters, highlighting a role for WRKY70 in the transcriptional response to NHP several hours following treatment.

NHP pretreatment enhances flg22-elicited ROS production and WRKY70 contributes to the full response

We next questioned if WRKY70 regulates a specific branch of NHP defense signaling. One such branch involves the response to common molecular features of microbes (e.g., bacterial flagellin or fungal chitin) known as microbe-associated molecular patterns (MAMPs) and the response to plant-derived damage-associated molecular patterns (DAMPs), a process collectively referred to as pattern-triggered immunity (PTI).

The WRKY70-dependent gene *BDA1* is known to mediate signaling in response to
MAMP detection and interacts with receptor-like protein SNC2 (Suppressor of NPR1,
Constitutive2) to relay MAMP-triggered defense responses (Yang et al., 2012).
Production of reactive oxygen species (ROS) is a MAMP-triggered defense response
which can act directly or indirectly as an antimicrobial agent and serves as a secondary
signal to activate further defense responses (Boller and Felix, 2009; Couto and Zipfel,
2016). Similarly, DAMPs, such as secreted peptides from the PROSCOOP family, are
known to elicit an increase in ROS production (Gully et al., 2019; Hou et al., 2021;
Rhodes et al., 2021). Incidentally, *CML10* encodes a calmodulin-like protein known to
interact with phosphomannomutase (PMM) in order to modulate ascorbic acid synthesis
and cellular homeostasis of ROS (Cho et al., 2016).

We thus investigated how NHP alters MAMP-triggered ROS production using bacterial
flagellin as the elicitor in the presence and absence of NHP pretreatment and asked if
WRKY70 is required for the response. Leaf discs of wild type and *wrky70-1* plants were
floated on solutions of water, NHP, and SA. SA was included as a positive control, as it
is known to enhance the MAMP-triggered ROS burst (Yi et al., 2014). After 24 h, leaf
discs were treated with a 100 nM solution of flagellin peptide (flg22) and then ROS was
measured using a luminol-based assay. We found ROS levels from mock treated
wrky70-1 plants trended lower than water treated wild type, suggesting WRKY70 is
required for maximal ROS accumulation (Fig. 7, upper panel). NHP treatment increased
ROS generation in both wild type and *wrky70-1* tissue exposed to flg22 when compared
to water treated tissues. The average ROS production in NHP treated *wrky70-1* leaf
discs was lower than NHP treated wild type leaf discs (Fig. 7, lower panel). Similar
results were found when analyzing *wrky70-2* plants (Supplementary Fig. S9).
Collectively, these data indicate that NHP is able to enhance the flg22-elicited ROS
burst, and this enhancement requires WRKY70 for full ROS generation.

Discussion
NHP has emerged as an essential signaling molecule required for the establishment
and amplification of SAR in Arabidopsis and crop plants (Chen et al., 2018; Holmes et

al., 2019), though the early signaling events mediated by NHP had yet to be described.
Our study provides the first insights into the primary transcriptional responses to NHP,
as well as the role that NHP-induced SA biosynthesis plays during this response. We
also discovered that TGA and WRKY CREs are enriched in the promoters of early NHP
induced genes, providing a framework for dissecting primary and secondary
transcriptional responses. Further, we show that early transcriptional expression of
WRKY70, within minutes of NHP treatment, mediates secondary transcriptional
reprogramming that is required for NHP signaling, ROS production, and SAR.

Early short-lived NHP transcriptional responses associate with general stress
In the early response to NHP, we identified a primary wave of NHP signaling that was
short lived and activated transcriptional responses mirroring wounding/DAMP and JA
signaling in a partially SA-dependent manner. Of the pathways activated by NHP, we
found a considerable enrichment of known JA-response genes, including *MYC2* and
JAZ10 (Fig. 2C, Table 1, Table 2). Given the established antagonism between JA and
SA signaling pathways (Pieterse et al., 2012), it was surprising to find the majority of
genes in the early transient cluster were SA-dependent (Fig. 3A). These findings
suggest that part of the early NHP transcriptional response depends on synergism
between JA signaling components and pre-infection levels of SA. Our data support a
model where general stress response signatures are activated by NHP treatment in an
early pulse of gene expression that acts synergistically with SA to modulate activation of
defense pathways.

Previous work has described a strong, systemic JA response in Arabidopsis leaves
infected with an avirulent *Pst* strain after 4 h, suggesting JA signaling pathways are
involved in SAR development (Truman et al., 2007). The JA response detected by
Truman *et al*. in the systemic leaves of avirulent *Pst*-treated plants after 4 h may be
reflective of the early and transient response to NHP observed in our transcriptional
analysis. These JA- and early NHP-responsive marker genes may prove useful for
determining optimal time points in testing the systemic translocation of NHP.
Furthermore, although genetic studies contest the requirement for JA biosynthesis and

signaling in SAR (Attaran et al., 2009), it remains a possibility that differences in primary

pathogen treatment or secondary pathogen infection may shift the requirement for JA

signaling in SAR establishment and warrant further study.

Early strong NHP responses activate a primary wave of *WRKY* transcription

We discovered another primary wave of early NHP signaling that was partially SA-

independent and exhibited strong induction of *WRKY* gene transcription within minutes

of NHP treatment, persisting several hours after treatment. All six *WRKY* TF genes in

the early stable upregulated gene cluster were activated by NHP treatment in the *sid2-2*

mutant, and although some showed delays in the timing of induction in *sid2-2* seedlings,

all six were strongly upregulated by 3 h (Table 1, Table 2). These findings are intriguing

given that the expression of these *WRKY* TFs is known to be induced by SA and SA-

analogs in an NPR1-dependent manner (Kalde et al., 2003; Wang et al., 2006). The

ability of NHP to induce an early wave of *WRKY* expression, independent of elevated

SA, suggests NHP and not SA is the primary signal driving this early response.

The early and strong induction of *WRKY* transcription, including two sets of homologs

belonging to WRKY group III (*WRKY38* and *WRKY62* as well as *WRKY54* and

WRKY70), suggested WRKY TFs may play a role in driving the secondary

transcriptional response to NHP. Members of WRKY group III are known to play both

positive and negative regulatory roles in defense responses (Kalde et al., 2003; Pandey

and Somssich, 2009). Consistent with this, we found CREs of WRKY TFs to be

abundant in the promoters of genes, both up and downregulated, several hours after

NHP treatment (Fig. 4C, Supplemental Fig. 5). For example, the CRE bound by

WRKY70, as defined by the DAPseq Plant Cistrome Database (O'Malley et al., 2016) ,

was enriched in the promoters of genes upregulated by NHP at 3 and 6 h (Fig. 4C).

Other canonical motifs bound by WRKY70, such as the W-box and WT-box (Rushton et

al., 2010; Machens et al., 2014), were also found in the promoters of genes upregulated

several hours after NHP treatment (Fig. 6A). Taken together, these findings link the

early NHP-driven induction of *WRKY* expression to the subsequent up and

downregulation of WRKY-targeted genes 3-6 h after initial NHP treatment. Based on

these findings, we hypothesize a model of early NHP signaling wherein activation of
WRKY transcription is a primary response to NHP signaling that is followed by a
secondary wave of expression driven through WRKY regulation.

While not the focus of this study, it is noteworthy to mention that TGA TFs are also
associated with the primary wave of NHP mediated transcription. CREs of TGAs were
abundant in the promoters of early NHP upregulated genes across all time points
sampled, particularly those in the early strong cluster (Fig. 4B). These findings are in
agreement with recent studies demonstrating TGAs are redundantly required for NHP-
induction of SAR and SAR marker genes (Nair et al., 2021; Yildiz et al., 2023). Our data
provide further evidence highlighting the importance of TGAs in NHP-driven signaling
and suggest that a NPR1/TGA complex, or a key interactor of this complex, directly
relays the NHP signal.

Late weak and late strong genes define the secondary wave of transcription
Finally, our RNA-seq analysis also revealed a secondary wave of NHP signaling defined
by late weak and late strong upregulated genes. These genes are associated with
pathogen defense, SAR, and SA responses, and showed features of SA-dependent and
SA-independent transcriptional induction (Fig. 2C, Fig. 3A). Upregulation of SAR-
associated genes by NHP is consistent with previous studies (Bernsdorff et al., 2016;
Yildiz et al., 2021; Yildiz et al., 2023). Our study reveals that transcriptional activation of
these SAR-associated genes occurs as early as 3-6 h following detection of the NHP
signal. Interestingly, 24 genes from the late weak cluster and 4 genes from the late
strong cluster are SA-independent within 6 h of NHP treatment but SA-dependent 24 h
post NHP (Yildiz et al., 2021). Among this group are key SA and NHP biosynthetic and
regulatory genes including *ALD1*, *UGT76B1*, *SARD1* and *NPR3*. Together, these data
suggest NHP is sufficient to drive gene expression over a short time frame but requires
SA accumulation for sustained expression, thus highlighting the importance of
interrogation into early NHP transcriptional changes. Further elucidation of a timeframe
for early NHP responses should prove useful in studying other aspects of NHP biology,

such as the biosynthesis and translocation of NHP and the identification of NHP

sensors and their respective downstream signaling pathways.

Primary NHP-responsive gene *WRKY70* mediates secondary transcriptional

changes required for NHP-elicited SAR, signaling, and ROS production

We found *WRKY70* expression is induced within 15 min of NHP treatment and is

required for full NHP-elicited SAR, as NHP did not enhance resistance in distal leaves

of the *wrky70-1* mutant and could not achieve the same level of resistance observed in

NHP-treated wild type plants (Fig. 5A). WRKY70 is known to repress many pathogen-

inducible genes in the absence of infection, but is also required for full activation of

pathogen-inducible genes during infection (Zhou et al., 2018). We observed enhanced

resistance to *Psm* infection and enhanced transcript abundance of defense genes

SARD1, *PR1*, *PR2*, and *PR5* after mock treatment (Supplement Fig. S7), consistent

with a repressive role for WRKY70. However, NHP treatment did not enhance

resistance nor increase transcript abundance of these genes in *wrky70-1* (Fig. 5A,

Supplement Fig. S7). Furthermore, expression of the NHP-upregulated early strong

gene *BDA1* was completely abolished in *wrky70* mutant lines (Fig. 6B, Supplemental

Fig. S8), demonstrating early expression of *WRKY70* is required to drive and modulate

a subset of the secondary transcriptional responses of NHP signaling.

Evidence supports a SA- and NPR1-independent defense pathway driven by WRKY70

(Shah et al., 2001; Zhang et al., 2010b). We found that NHP was sufficient to induce

expression of *WRKY70* in the *sid2-2* mutant within 30 min of treatment (Table 1),

suggesting SA biosynthesis is dispensable for the early transcriptional activation of

WRKY70. Likewise, WRKY70-regulated genes *BDA1* and *CML10* are induced within 3-

6 h of NHP signaling partially independent of SA (Table1). Notably, 24 h after NHP

treatment, *BDA1* and *CML10* are also upregulated in both *sid2-1* and *npr1-3* mutants

(Yildiz et al., 2021). It is therefore possible that NHP signaling activates a SA- and

NPR1-independent defense pathway through the action of WRKY70.

695 Studies into the MAMP-signaling pathway functioning through the receptor-like protein

696 SNC2 suggest a role for WRKY70 and BDA1 in PTI responses such as ROS production

697 (Yang et al., 2012). While flg22-elicited ROS production was not significantly different

698 between mock treated wild type and *wrky70* mutants, the enhanced production of flg22-

699 elicited ROS following NHP and SA pretreatment was compromised in *wrky70* mutants

700 (Fig. 7, Supplemental Fig. S9). Our findings indicate WRKY70 positively contributes to

701 the NHP and SA signaling pathways that prime PTI. Further, our discovery that NHP

702 primes plants for enhanced production of flg22-elicited ROS is consistent with reports of

703 NHP priming Arabidopsis for enhanced metabolic and transcriptional responses

704 following flg22 treatment (Löwe et al., 2023). Our work links WRKY70-dependent

705 signaling pathways to this NHP-primed PTI response.

707 **Conclusion**

708 Our work describes the early transcriptional response to NHP and highlights key genes

709 and signaling pathways that may contribute to the physiological processes regulated by

710 this important bioactive molecule. It also provides a foundation for better understanding

711 the transcriptional regulators, particularly the role of WRKY70, and gene networks that

712 define early NHP signal transduction.

714 **Material and methods**

715 **Plant materials and growth conditions**

716 *Arabidopsis thaliana* (Arabidopsis) lines used in this study were in the Col-0 ecotype

717 (wild type) with mutant lines as follows: *sid2-2* (Wildermuth et al., 2001) and T-DNA

718 insertion lines *wrky70-1* (SALK_025198) and *wrky70-2* (GABI_324D11) (Li et al., 2006),

719 *wrky38* (WiscDsLox489-492C21), *wrky54* (SALK_017254), and *wrky59* (SALK_039436)

720 (Wang et al., 2006), *wrky51* (SALK_022198) (Yan et al., 2018), and *wrky62*

721 (SM_3_38820) (Kim et al., 2008). All mutant lines were obtained from the Arabidopsis

722 Biological Resource Center (ABRC). The presence of T-DNA insertions were confirmed

723 with the genotyping primers listed in Supplemental Table S1. Plants were grown on soil

724 in a controlled growth chamber at 80% humidity and 22°C on a 10 h light/14 h dark

cycle for gene expression and bacterial growth assays. For seedlings grown
hydroponically, seeds were surface sterilized and stratified at 4°C in darkness for 3
days. Seeds were placed into sterile six-well plates (15-18 seeds per well) with half-
strength Murashige and Skoog (MS) medium, 0.5% (w/v) sucrose, 0.05% (w/v) MES at
pH 5.7. Plates were covered with a lid, sealed with micropore tape (3M), and placed in a
chamber on a 10 h light/14 h dark cycle. Liquid medium was agitated twice daily to
facilitate gas exchange and after five days of growth, spent medium was replaced with
fresh MS.
733

734 **Bacterial strains and growth conditions**
735 *P. syringae* pv. *maculicola* strain ES4326 (*Psm*) and *P. syringae* pv. *tomato* DC3000
736 (*Pst* DC3000) carrying pVSP61 + *avrRpt2* (*Pst avrRpt2*) were used in this study. Growth
737 conditions for all bacterial strains are as previously described (Holmes et al., 2021).
738

Chemical stocks
Two sources of NHP were used in this study, one synthesized by Elizabeth Sattely's lab
as previously described (Chen et al., 2018) and the second obtained from Med Chem
Express. Gene expression and NHP-elicited SAR assays showed similar results for
both sources of NHP. Pipecolic acid stocks were made from L -pipecolic acid
(Oakwood Chemical). Salicylic acid stocks were made from sodium salicylate
(Mallinckrodt).
746

Chemical treatment of plants for qRT-PCR assays
For gene expression by qRT-PCR of 4.5-week-old, soil-grown Arabidopsis, three leaves
were infiltrated with sterile water or NHP dissolved in water at the indicated
concentrations. The three treated leaves were collected and flash frozen at the
indicated time points. For the time course of gene expression in *wrky70-1 and wrky70-2*
mutants, three leaves were treated with sterile water or 1 mM NHP. One leaf from the
same plant was sampled at 6, 9, and 12 h and pooled with a second plant of the same
condition for one replicate. Samples collected at 0 h were untreated before collecting.
755

For gene expression in Arabidopsis seedling-based assays, plants were grown as
described for seedling hydroponics and treated at 10-days-post-germination. Before
addition of treatment, all liquid medium was removed from plate wells and replaced with
fresh MS medium, seedlings were allowed to recover under lights for at least 1 h,
followed by application of 2x solutions of Pip, NHP, or mock in MS medium. The 15-18
seedlings within a single well were pooled and treated as a single replicate, with three
replicates sampled per condition. Following treatment, seedlings were blotted dry, and
flash frozen at the indicated time points.

RNA isolation and qRT-PCR
Total RNA was isolated from leaves and seedlings using TRIsure™ reagent (Meridian
Bioscience) according to the manufacturer's instructions. 5 µg of RNA was used to
synthesize cDNA by oligo dT (New England Biolabs) and reverse transcriptase (Thermo
Fisher), followed by dilution of cDNA. The qRT-PCR reactions were performed on a MJ
Opticon 2 (Bio-Rad) using Green Taq DNA polymerase (GenScript) with EvaGreen Dye
(Biotium) for amplification and detection. Three technical replicates were performed per
sample and three to four biological replicates were included for each condition.
Expression values were normalized to the reference gene *UBC21* (AT5G25760) for
relative expression determined using $2^{-\Delta Ct}$ and normalized to both *UBC21* and mock
treated samples for fold change using $2^{-\Delta\Delta Ct}$. Genes assayed include *FMO1*
(AT1G19250), *ICS1*/*SID2* (AT1G74710), *UGT76B1* (AT3G11340), *WRKY38*
(AT5G22570), *SARD1* (AT1G73805), *PR1* (AT2G14610), *PR2* (AT3G57260), *PR5*
(AT1G75040), *BDA1* (AT5G54610), *PROSCOOP4* (AT5G44568), and *CML10*
(AT2G41090). Primers used in these experiments can be found in Supplemental Table
S1.

RNA-seq treatment and library preparation
Seedlings used for mRNA isolation were grown hydroponically as described above. Six-
well plates contained equal numbers of Col-0 and *sid2-2* seedlings, with one genotype
per plate and 15 seedlings per well. At 10-days-post-germination, all liquid medium was
removed from each well and fresh MS medium was added to the seedlings. Seedlings

were placed back into the growth chamber for an hour to acclimate, followed by

treatment with 2x NHP (final concentration 0.5 mM) or mock (MS medium). Three wells

of pooled seedlings were collected for each genotype, treatment, and time (15 min, 30

min, 3 h, and 6 h) combinations, blotted dry, and flash frozen for mRNA isolation.

Total RNA was extracted from pooled seedlings using TRIsure™ reagent (Meridian

Bioscience) according to the manufacturer's instructions. Total RNA was then divided

with 5 µg used for cDNA synthesis and marker gene expression to assess quality of

treatment, as described for qRT-PCR experiments, and 5 µg was used for cDNA library

preparation. The cDNA library was prepared using NEBNext® Ultra™ II RNA Library

Prep Kit for Illumina® with NEBNext® Poly(A) mRNA Magnetic Isolation Module and

then multiplexed via PCR amplification using NEBNext® Multiplex Oligos for Illumina®

according to the manufacturer's instructions (New England Biolabs). Library

concentration was quantified with Qubit™ dsDNA HS Assay Kit (Thermo Fisher

Scientific) followed by measurement of quality and average cDNA length using the DNA

1000 Series II chip on a 2100 Bioanalyzer (Agilent Technologies). The multiplexed

cDNA libraries were then pooled (two sets of 36 libraries) and sequenced using both

lanes of the NovaSeq 6000 (Illumina) S1 flow cell with run type 100 bp paired-end reads

at the Genome Sequencing Service Center by the Stanford Center for Genomics and

Personalized Medicine, supported by the grant award NIH S10OD025212.

RNA-seq data analysis

Read data was initially assessed using FastQC. CLC Genomics Workbench version

11.0.2 (QIAGEN) was used for trimming, discarding reads trimmed below 20 bp, and

then mapped to the Arabidopsis TAIR10 genome with the auto-detect paired distances

option deselected. Differential expression analysis was performed using DESeq2 (Love

et al., 2014) with mock treated samples compared to NHP treated samples for each

respective time point and genotype. Genes were considered differentially expressed if

they returned an adjusted *P*-value below 0.05 ($P_{adj} < 0.05$). The complete list of early

NHP-responsive transcripts is provided in Supplemental Table S3 and the raw

sequencing data deposited in the Gene Expression Omnibus (GEO) under accession
code GSE263276.

Exploratory analysis of differentially expressed genes

To define patterns of early expression in response to NHP treatment, differentially
expressed genes were considered upregulated in wild type if they had a $\log_2(FC) > 1$ at
any time point and downregulated with a $\log_2(FC) < -1$ at any time point. The sets of up
and downregulated genes were then hierarchically clustered by $\log_2(FC)$ over the four
timepoints using the hclust function and dendextend packages in R, with Euclidean
distances and Ward.D2 linkage (Galili, 2015). For each defined cluster in wild type, the
$\log_2(FC)$ values were averaged at each time point to visualize broad trends in
expression.

The PANTHER classification system accessed via TAIR was used for gene ontology
enrichment analysis (https://www.arabidopsis.org/tools/go_term_enrichment.jsp; (Mi et
al., 2021)). Version 17.0 of the PANTHER overrepresentation test was used (Fisher's
exact test with FDR correction) and matched to the GO database 2021_03 release.

Promoter and *cis*-regulatory element enrichment analysis

Araport11 sequences 1000 bp upstream of the TSS of NHP-responsive genes were
downloaded from TAIR (https://www.arabidopsis.org/tools/bulk/sequences/index.jsp) for
each time point and genotype grouping of the NHP upregulated and downregulated
genes. A less stringent cut-off of $\log_2(FC) > 0$ (upregulated) or $\log_2(FC) < 0$
(downregulated) with a $P_{adj} < 0.05$ was used to select for NHP-responsive genes. Motif
identification and enrichment of *cis*-regulatory elements (CREs) was performed using
SEA, part of the MEME suite (Bailey et al., 2015; Bailey and Grant, 2021) and compared
to the published library of TF binding sites found within the DAPseq database (O'Malley
et al., 2016). Only CREs from the DAP-seq library were used in this study, omitting
CREs from the ampDAP-seq library (where DNA modifications have been removed).
Following the workflow detailed by Bjornson et al. (2021), relative enrichment was
determined against the same three sets of randomly sampled genes detected in this

RNA-seq experiment for all genotypes and time points tested. The identified motif was

considered enriched if it returned a significant *P*-value (< 0.05) relative to at least two of

the three randomly sampled gene sets. *P*-values for significantly enriched TF-binding

motifs were then averaged for visualization.

Measurement of bacterial growth in plants

Chemical treatment of leaves for NHP-elicited SAR assays were performed as

described in (Chen et al., 2018) with minor alterations. At 4.5-weeks-old, three lower

leaves of wild type and mutant Arabidopsis plants were infiltrated with sterile water, 0.5

mM NHP or 1 mM NHP. After 24 h, one untreated distal leaf of each plant was

inoculated with a 1 x 10^5 CFU/mL suspension of *Psm*. The distal infected leaf was

always separated by one leaf from the original three treated (i.e., primary treatment in

leaf no. 7-9 then infected leaf is no. 11). Inoculated plants were covered with a dome to

increase humidity and 3 dpi the titer of *Psm* in the distal leaves was quantified. Leaf

discs were homogenized in 1 mL of 10 mM $MgCl_2$, plated in triplicate in a dilution series

on nutrient yeast glycerol medium supplemented with 1.5% wt/vol agar (NYGA) with

rifampicin (100 µg/mL), incubated at 28 °C for 1.5 d, and bacterial colonies counted. For

each individual experiment, 4-12 plants were tested per condition.

Pathogen priming of leaves for SAR assays were performed as previously described

(Chen et al., 2018). Three lower leaves of wild type and mutant Arabidopsis plants (4.5-

weeks-old) were infiltrated with 10 mM $MgCl_2$ or a 5 x 10^6 CFU/mL suspension of *Pst*

avrRpt2 in 10 mM $MgCl_2$. After 48 h, one distal untreated leaf of each plant was

inoculated with a 1 x 10^5 CFU/mL suspension of *Psm* and plants were covered with a

dome to increase humidity. At 3 dpi, the titer of *Psm* in the upper leaves was quantified

as described above for NHP-elicited SAR. For each individual experiment, 8 plants were

tested per condition and experiments repeated twice with similar results.

For bacterial growth measurements in unprimed leaves of Arabidopsis wild type and

mutant plants, a suspension of 1 x 10^5 CFU/mL *Psm* was infiltrated into two young

leaves per plant. Plants were placed under a dome and 3 dpi the two infected leaves
were pooled and bacterial titer quantified for 12 plants per genotype.

Oxidative burst assay

Four leaf discs (4 mm diameter) from 5-week-old Arabidopsis plants (n = 12) were
incubated on water, 500 μM NHP, or 100 μM SA in a 96-well plate (one leaf disc per
well) for 24 h. After pretreatment, ROS was measured by addition of flg22 (100 nM) in a
solution of 20 mg/mL horseradish peroxidase and 200 μM Luminol (Sigma)
(Gómez☐Gómez et al., 1999), followed by immediate measurement of luminescence in
a Synergy H1 Microplate Reader (Biotek). Relative luminescence units (RLU) are
reported.

Statistical analyses

Two-tailed Student's *t* tests were performed to determine statistically significant
differences between the different conditions for measurements of bacterial growth using
the stat_compare_means function of the ggpubr package (Kassambara, 2023). *P* values
< 0.05 were considered statistically significant and all significance levels are indicated in
the figure legends (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$). Statistical
significance of qRT-PCR assays was determined using Mann-Whitney *U* test or one-
way ANOVA with post hoc Sidak test using the ggpubr and lsmeans R packages,
respectively (Lenth, 2016).

Acknowledgements

We thank the Mudgett laboratory for feedback on the research and manuscript. We also
thank Zhiyong Wang, Dominique Bergmann, Sharon Long, and Virginia Walbot for
instrument use and intellectual discussion. This work was supported by National
Science Foundation IOS-2026368 (to M.B.M. and E.S.S) and National Science
Foundation Graduate Research Fellowship DGE-1656518 (to J.F.).

Author contributions

J.F., J.-G.K., E.S.S. and M.B.M contributed to the study design; J.F. and J.-G.K. performed research; J.F., J.-G.K., E.S.S. and M.B.M analyzed data; J.F., and M.B.M. wrote the manuscript; J.F., J.-G.K., E.S.S. and M.B.M. reviewed and revised the manuscript for accuracy and final approval.

References

Attaran E, Zeier TE, Griebel T, Zeier J (2009) Methyl salicylate production and jasmonate signaling are not essential for systemic acquired resistance in *Arabidopsis*. Plant Cell **21**: 954–971

Bailey TL, Grant CE (2021) SEA: Simple Enrichment Analysis of motifs. BioRxiv. doi: 10.1101/2021.08.23.457422

Bailey TL, Johnson J, Grant CE, Noble WS (2015) The MEME Suite. Nucleic Acids Res **43**: W39-49

Bartsch M, Gobbato E, Bednarek P, Debey S, Schultze JL, Bautor J, Parker JE (2006) Salicylic acid-independent ENHANCED DISEASE SUSCEPTIBILITY1 signaling in Arabidopsis immunity and cell death is regulated by the monooxygenase FMO1 and the Nudix hydrolase NUDT7. Plant Cell **18**: 1038–1051

Bauer S, Mekonnen DW, Hartmann M, Yildiz I, Janowski R, Lange B, Geist B, Zeier J, Schäffner AR (2021) UGT76B1, a promiscuous hub of small molecule-based immune signaling, glucosylates N-hydroxypipecolic acid, and balances plant immunity. Plant Cell **33**: 714–734

Bernsdorff F, Döring A-C, Gruner K, Schuck S, Bräutigam A, Zeier J (2016) Pipecolic Acid Orchestrates Plant Systemic Acquired Resistance and Defense Priming via Salicylic Acid-Dependent and -Independent Pathways. Plant Cell **28**: 102–129

Bjornson M, Pimprikar P, Nürnberger T, Zipfel C (2021) The transcriptional landscape of *Arabidopsis thaliana* pattern-triggered immunity. Nat Plants **7**: 579–586

Blanco F, Salinas P, Cecchini NM, Jordana X, Van Hummelen P, Alvarez ME, Holuigue L (2009) Early genomic responses to salicylic acid in Arabidopsis. Plant Mol Biol **70**: 79–102

Boller T, Felix G (2009) A renaissance of elicitors: perception of microbe-associated
molecular patterns and danger signals by pattern-recognition receptors. Annu Rev
Plant Biol **60**: 379–406

Cai J, Jozwiak A, Holoidovsky L, Meijler MM, Meir S, Rogachev I, Aharoni A (2021)
Glycosylation of N-hydroxy-pipecolic acid equilibrates between systemic acquired
resistance response and plant growth. Mol Plant **14**: 440–455

Chen Y-C, Holmes EC, Rajniak J, Kim J-G, Tang S, Fischer CR, Mudgett MB,
Sattely ES (2018) *N*-hydroxy-pipecolic acid is a mobile metabolite that induces
systemic disease resistance in *Arabidopsis.* Proc Natl Acad Sci USA **115**: E4920–
E4929

Chini A, Fonseca S, Fernández G, Adie B, Chico JM, Lorenzo O, García-Casado G,
López-Vidriero I, Lozano FM, Ponce MR, et al (2007) The JAZ family of repressors
is the missing link in jasmonate signalling. Nature **448**: 666–671

Cho K-M, Nguyen HTK, Kim SY, Shin JS, Cho DH, Hong SB, Shin JS, Ok SH (2016)
CML10, a variant of calmodulin, modulates ascorbic acid synthesis. New Phytol **209**:
664–678

Conrath U, Beckers GJM, Langenbach CJG, Jaskiewicz MR (2015) Priming for
enhanced defense. Annu Rev Phytopathol **53**: 97–119

Couto D, Zipfel C (2016) Regulation of pattern recognition receptor signalling in plants.
Nat Rev Immunol **16**: 537–552

Ding Y, Sun T, Ao K, Peng Y, Zhang Y, Li X, Zhang Y (2018) Opposite Roles of
Salicylic Acid Receptors NPR1 and NPR3/NPR4 in Transcriptional Regulation of
Plant Immunity. Cell **173**: 1454-1467.e15

Eulgem T, Rushton PJ, Robatzek S, Somssich IE (2000) The WRKY superfamily of
plant transcription factors. Trends Plant Sci **5**: 199–206

Friedrichsen DM, Nemhauser J, Muramitsu T, Maloof JN, Alonso J, Ecker JR,
Furuya M, Chory J (2002) Three redundant brassinosteroid early response genes
encode putative bHLH transcription factors required for normal growth. Genetics **162**:
1445–1456

Fu ZQ, Dong X (2013) Systemic acquired resistance: turning local infection into global
defense. Annu Rev Plant Biol **64**: 839–863

Galili T (2015) dendextend: an R package for visualizing, adjusting and comparing
trees of hierarchical clustering. Bioinformatics **31**: 3718–3720

Gómez-Gómez L, Felix G, Boller T (1999) A single locus determines sensitivity to
bacterial flagellin in Arabidopsis thaliana. Plant J **18**: 277–284

Gully K, Pelletier S, Guillou M-C, Ferrand M, Aligon S, Pokotylo I, Perrin A, Vergne
E, Fagard M, Ruelland E, et al (2019) The SCOOP12 peptide regulates defense
response and root elongation in *Arabidopsis thaliana*. J Exp Bot **70**: 1349–1365

Hartmann M, Zeier J (2019) N-hydroxypipecolic acid and salicylic acid: a metabolic duo
for systemic acquired resistance. Curr Opin Plant Biol **50**: 44–57

Hartmann M, Zeier T, Bernsdorff F, Reichel-Deland V, Kim D, Hohmann M,
Scholten N, Schuck S, Bräutigam A, Hölzel T, et al (2018) Flavin Monooxygenase-
Generated N-Hydroxypipecolic Acid Is a Critical Element of Plant Systemic Immunity.
Cell **173**: 456-469.e16

Holmes EC, Chen Y-C, Mudgett MB, Sattely ES (2021) Arabidopsis UGT76B1
glycosylates *N*-hydroxy-pipecolic acid and inactivates systemic acquired resistance in
tomato. Plant Cell **33**: 750–765

Holmes EC, Chen Y-C, Sattely ES, Mudgett MB (2019) An engineered pathway for *N*-
hydroxy-pipecolic acid synthesis enhances systemic acquired resistance in tomato.
Sci Signal. 12:eaay3066

Hou S, Liu D, Huang S, Luo D, Liu Z, Xiang Q, Wang P, Mu R, Han Z, Chen S, et al
(2021) The Arabidopsis MIK2 receptor elicits immunity by sensing a conserved
signature from phytocytokines and microbes. Nat Commun **12**: 5494

Jaskiewicz M, Conrath U, Peterhänsel C (2011) Chromatin modification acts as a
memory for systemic acquired resistance in the plant stress response. EMBO Rep **12**:
50–55

Johannesson H, Wang Y, Hanson J, Engström P (2003) The *Arabidopsis thaliana*
homeobox gene *ATHB5* is a potential regulator of abscisic acid responsiveness in
developing seedlings. Plant Mol Biol **51**: 719–729

Kachroo A, Kachroo P (2020) Mobile signals in systemic acquired resistance. Curr
Opin Plant Biol **58**: 41–47

Kalde M, Barth M, Somssich IE, Lippok B (2003) Members of the *Arabidopsis* WRKY
group III transcription factors are part of different plant defense signaling pathways.
Mol Plant Microbe Interact **16**: 295–305

Kassambara A (2023) *ggpubr: 'ggplot2' Based Publication Ready Plots*. R package
version 0.6.0. https://rpkgs.datanovia.com/ggpubr/

Kim K-C, Lai Z, Fan B, Chen Z (2008) *Arabidopsis* WRKY38 and WRKY62 transcription factors interact with histone deacetylase 19 in basal defense. Plant Cell **20**: 2357–2371

Kim Y, Gilmour SJ, Chao L, Park S, Thomashow MF (2020) *Arabidopsis* CAMTA transcription factors regulate pipecolic acid biosynthesis and priming of immunity genes. Mol Plant **13**: 157–168

Klessig DF, Choi HW, Dempsey DA (2018) Systemic acquired resistance and salicylic acid: past, present, and future. Mol Plant Microbe Interact **31**: 871–888

Kohler A, Schwindling S, Conrath U (2002) Benzothiadiazole-Induced Priming for Potentiated Responses to Pathogen Infection, Wounding, and Infiltration of Water into Leaves Requires the *NPR1*/*NIM1* Gene in Arabidopsis. Plant Physiol **128**: 1046–1056

Lenth RV (2016) Least-Squares Means: the *R* package lsmeans. J Stat Softw **69**: 1–33

Liu H, Liu B, Lou S, Bi H, Tang H, Tong S, Song Y, Chen N, Zhang H, Jiang Y, et al (2021) CHYR1 ubiquitinates the phosphorylated WRKY70 for degradation to balance immunity in *Arabidopsis thaliana*. New Phytol **230**: 1095–1109

Li J, Brader G, Kariola T, Palva ET (2006) WRKY70 modulates the selection of signaling pathways in plant defense. Plant J **46**: 477–491

Li J, Brader G, Palva ET (2004) The WRKY70 transcription factor: a node of convergence for jasmonate-mediated and salicylate-mediated signals in plant defense. Plant Cell **16**: 319–331

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol **15**: 550

Löwe M, Jürgens K, Zeier T, Hartmann M, Gruner K, Müller S, Yildiz I, Perrar M, Zeier J (2023) N-hydroxypipecolic acid primes plants for enhanced microbial pattern-induced responses. Front Plant Sci **14**: 1217771

Luan S, Kudla J, Rodriguez-Concepcion M, Yalovsky S, Gruissem W (2002) Calmodulins and calcineurin B-like proteins: calcium sensors for specific signal response coupling in plants. Plant Cell **14 Suppl**: S389-400

Machens F, Becker M, Umrath F, Hehl R (2014) Identification of a novel type of WRKY transcription factor binding site in elicitor-responsive *cis*-sequences from *Arabidopsis thaliana*. Plant Mol Biol **84**: 371–385

Mariani L, Weinand K, Vedenko A, Barrera LA, Bulyk ML (2017) Identification of Human Lineage-Specific Transcriptional Coregulators Enabled by a Glossary of Binding Modules and Tunable Genomic Backgrounds. Cell Syst **5**: 187-201.e7

Mishina TE, Zeier J (2006) The Arabidopsis Flavin-Dependent Monooxygenase FMO1
Is an Essential Component of Biologically Induced Systemic Acquired Resistance.
Plant Physiol **141**: 1666–1675

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD
(2021) PANTHER version 16: a revised family classification, tree-based classification
tool, enhancer regions and extensive API. Nucleic Acids Res **49**: D394–D403

Mohanta TK, Yadav D, Khan A, Hashem A, Tabassum B, Khan AL, Abd Allah EF,
Al-Harrasi A (2020) Genomics, molecular and evolutionary perspective of NAC
transcription factors. PLoS ONE **15**: e0231425

Mohnike L, Rekhter D, Huang W, Feussner K, Tian H, Herrfurth C, Zhang Y,
Feussner I (2021) The glycosyltransferase UGT76B1 modulates N-hydroxy-pipecolic
acid homeostasis and plant immunity. Plant Cell **33**: 735–749

Nair A, Goyal I, Voß E, Mrozek P, Prajapati S, Thurow C, Tietze L, Tittmann K, Gatz
C (2021) N-hydroxypipecolic acid-induced transcription requires the salicylic acid
signaling pathway at basal SA levels. Plant Physiol **187**: 2803–2819

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M,
Gallavotti A, Ecker JR (2016) Cistrome and Epicistrome Features Shape the
Regulatory DNA Landscape. Cell **165**: 1280–1292

Pandey SP, Somssich IE (2009) The Role of WRKY Transcription Factors in Plant
Immunity. Plant Physiol **150**: 1648–1655

Pieterse CMJ, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SCM (2012)
Hormonal modulation of plant immunity. Annu Rev Cell Dev Biol **28**: 489–521

Rhodes J, Yang H, Moussu S, Boutrot F, Santiago J, Zipfel C (2021) Perception of a
divergent family of phytocytokines by the Arabidopsis receptor kinase MIK2. Nat
Commun **12**: 705

Rushton PJ, Somssich IE, Ringler P, Shen QJ (2010) WRKY transcription factors.
Trends Plant Sci **15**: 247–258

Shah J, Kachroo P, Nandi A, Klessig DF (2001) A recessive mutation in the
Arabidopsis SSI2 gene confers SA- and *NPR1*-independent expression of *PR* genes
and resistance against bacterial and oomycete pathogens. Plant J **25**: 563–574

Song JT, Lu H, McDowell JM, Greenberg JT (2004) A key role for ALD1 in activation
of local and systemic defenses in *Arabidopsis*. Plant J **40**: 200–212

Sun T, Huang J, Xu Y, Verma V, Jing B, Sun Y, Ruiz Orduna A, Tian H, Huang X,
Xia S, et al (2020) Redundant CAMTA Transcription Factors Negatively Regulate the

Biosynthesis of Salicylic Acid and *N*-Hydroxypipecolic Acid by Modulating the Expression of *SARD1* and *CBP60g*. Mol Plant **13**: 144–156

Thibaud-Nissen F, Wu H, Richmond T, Redman JC, Johnson C, Green R, Arias J, Town CD (2006) Development of Arabidopsis whole-genome microarrays and their application to the discovery of binding sites for the TGA2 transcription factor in salicylic acid-treated plants. Plant J **47**: 152–162

Tian H, Zhang Y (2019) The emergence of a mobile signal for systemic acquired resistance. Plant Cell **31**: 1414–1415

Truman W, Bennett MH, Kubigsteltig I, Turnbull C, Grant M (2007) *Arabidopsis* systemic immunity uses conserved defense signaling pathways and is mediated by jasmonates. Proc Natl Acad Sci USA **104**: 1075–1080

Wang D, Amornsiripanitch N, Dong X (2006) A genomic approach to identify regulatory nodes in the transcriptional network of systemic acquired resistance in plants. PLoS Pathog **2**: e123

Wang Y, Schuck S, Wu J, Yang P, Döring A-C, Zeier J, Tsuda K (2018) A MPK3/6-WRKY33-ALD1-Pipecolic Acid Regulatory Loop Contributes to Systemic Acquired Resistance. Plant Cell **30**: 2480–2494

Wildermuth MC, Dewdney J, Wu G, Ausubel FM (2001) Isochorismate synthase is required to synthesize salicylic acid for plant defence. Nature **414**: 562–565

Wu Y, Zhang D, Chu JY, Boyle P, Wang Y, Brindle ID, De Luca V, Després C (2012) The *Arabidopsis* NPR1 Protein Is a Receptor for the Plant Defense Hormone Salicylic Acid. Cell Rep **1**: 639–647

Yang Y, Zhang Y, Ding P, Johnson K, Li X, Zhang Y (2012) The ankyrin-repeat transmembrane protein BDA1 functions downstream of the receptor-like protein SNC2 to regulate plant immunity. Plant Physiol **159**: 1857–1865

Yan C, Fan M, Yang M, Zhao J, Zhang W, Su Y, Xiao L, Deng H, Xie D (2018) Injury Activates Ca2+/Calmodulin-Dependent Phosphorylation of JAV1-JAZ8-WRKY51 Complex for Jasmonate Biosynthesis. Mol Cell **70**: 136-149.e7

Yi SY, Shirasu K, Moon JS, Lee S-G, Kwon S-Y (2014) The activated SA and JA signaling pathways have an influence on flg22-triggered oxidative burst and callose deposition. PLoS ONE **9**: e88951

Yildiz I, Gross M, Moser D, Petzsch P, Köhrer K, Zeier J (2023) N-hydroxypipecolic acid induces systemic acquired resistance and transcriptional reprogramming via TGA transcription factors. Plant Cell Environ. doi: 10.1111/pce.14572

Yildiz I, Mantz M, Hartmann M, Zeier T, Kessel J, Thurow C, Gatz C, Petzsch P,
Köhrer K, Zeier J (2021) The mobile SAR signal N-hydroxypipecolic acid induces
NPR1-dependent transcriptional reprogramming and immune priming. Plant Physiol
186: 1679–1705

Zhang Y, Fan W, Kinkema M, Li X, Dong X (1999) Interaction of NPR1 with basic
leucine zipper protein transcription factors that bind sequences required for salicylic
acid induction of the *PR-1* gene. Proc Natl Acad Sci USA **96**: 6523–6528

Zhang Y, Xu S, Ding P, Wang D, Cheng YT, He J, Gao M, Xu F, Li Y, Zhu Z, et al
(2010a) Control of salicylic acid synthesis and systemic acquired resistance by two
members of a plant-specific family of transcription factors. Proc Natl Acad Sci USA
107: 18220–18225

Zhang Y, Yang Y, Fang B, Gannon P, Ding P, Li X, Zhang Y (2010b) *Arabidopsis*
snc2-1D activates receptor-like protein-mediated immunity transduced through
WRKY70. Plant Cell **22**: 3153–3163

Zhou M, Lu Y, Bethke G, Harrison BT, Hatsugai N, Katagiri F, Glazebrook J
(2018) WRKY70 prevents axenic activation of plant immunity by direct repression of
SARD1. New Phytol **217**: 700–712

Tables

Table 1. SA-independent/partially independent NHP upregulated genes by cluster.

| | | | Log$_2$(FC) | | | | | | | |
| | | | Col-0 | | | | sid2-2 | | | |
AGI	Name	Annotation	15m	30m	3h	6h	15m	30m	3h	6h
Early transient										
AT3G44870	FAMT-L	FARNESOIC ACID METHYL TRANSFERASE-LIKE	4.3	4.4	-	-	2.3	2.7	-	-
AT2G44840	ERF13	Ethylene-responsive transcription factor	3.3	2.1	-	-	1.4	-	-	-
AT1G76640	CML39	CALMODULIN LIKE 39	2.7	2.7	-	-	1.7	-	-	-
AT5G13220	JAZ10	JASMONATE-ZIM-DOMAIN PROTEIN 10	1.6	1.1	-	0.6	1.2	1.0	-	-
AT1G80840	WRKY40	WRKY Transcription Factor; Group II-a	1.8	1.1	-	0.5	-	1.4	1.4	0.7
AT3G17690	CNGC19	Cyclic nucleotide-gated ion channel	1.6	1.6	-	-	-	1.2	-	-
AT4G21840	MSRB8	Methionine sulfoxide reductase B8	3.1	-	-	1.5	-	2.2	-	-
Early strong										
AT5G22570	WRKY38	WRKY Transcription Factor; Group III	5.8	5.2	4.7	3.6	3.9	6.2	5.8	2.9
AT5G01900	WRKY62	WRKY Transcription Factor; Group III	3.9	5.0	4.4	2.6	-	5.8	4.1	4.4
AT1G28480	GRXC9	GRX480/ROXY19, glutaredoxin family	3.2	2.2	2.7	1.8	3.2	3.1	2.3	2.1
AT2G40750	WRKY54	WRKY Transcription Factor; Group III	2.7	3.7	2.8	2.9	3.0	3.6	3.5	3.0
AT5G64810	WRKY51	WRKY Transcription Factor; Group II-c	2.6	3.4	3.3	3.0	1.7	3.2	3.2	3.5
AT3G56400	WRKY70	WRKY Transcription Factor; Group III	1.6	1.6	2.4	2.1	-	1.3	2.3	1.8
AT3G25882	NIMIN-2	NIM1-INTERACTING 2	1.4	1.5	1.8	1.6	1.2	1.1	1.0	1.6
Late weak										
AT3G60470	-	Transmembrane protein, putative (DUF247)	-	-	-	3.3	-	-	-	5.1
AT2G13810	ALD1	AGD2-LIKE DEFENSE RESPONSE PROTEIN 1	-	-	-	2.3	-	-	-	6.6
AT1G01680	PUB54	Plant U-box type E3 ubiquitin ligase	-	-	-	2.2	-	-	-	1.4
AT3G12220	SCPL16	Serine carboxypeptidase-like	-	-	1.1	2.0	-	-	-	2.0
AT1G78340	GSTU22	Glutathione S-transferase U22	-	-	2.0	2.0	-	-	1.6	1.7
AT5G03350	LLP	Lectin-like protein	-	-	2.1	1.8	-	-	2.1	1.9
AT5G24530	DMR6	DOWNY MILDEW RESISTANCE 6	0.7	1.2	1.6	1.7	0.5	1.6	2.0	1.7
AT2G41090	CML10	Calmodulin-like protein	-	-	1.1	1.7	-	-	0.3	1.2
AT4G14400	ACD6	ACCELERATED CELL DEATH 6	-	-	2.1	1.6	-	-	1.4	1.4
AT2G15490	UGT73B4	UDP-DEPENDENT GLYCOSYLTRANSFERASE 73B4	-	0.7	1.2	1.5	-	0.9	2.7	1.5
AT4G08040	ACS11	1-aminocyclopropane-1-carboxylate synthase 11	-	1.0	1.5	1.5	-	-	1.0	1.2
AT1G21250	WAK1	CELL WALL-ASSOCIATED KINASE 1	0.7	-	1.8	1.5	-	-	1.7	2.0
AT1G17170	GSTU24	Glutathione S-transferase U24	-	-	-	1.4	-	-	1.6	1.5
AT3G11340	UGT76B1	UDP-DEPENDENT GLYCOSYLTRANSFERASE 76B1	-	1.0	0.9	1.1	-	0.7	1.1	1.0
AT1G73805	SARD1	SAR DEFICIENT 1	1.0	0.7	1.5	1.0	-	-	1.3	1.3

Late strong

AT number	Gene	Description								
AT3G12230	SCPL14	Serine carboxypeptidase-like	-	-	5.9	6.9	-	-	5.4	4.6
AT1G15610	-	Transmembrane protein	-	-	5.5	5.9	-	-	-	5.1
AT3G28510	-	AAA-type ATPase family protein	-	-	3.2	5.6	-	-	-	4.9
AT2G26400	ARD	ACIREDUCTONE DIOXYGENASE 3	-	2.2	4.4	5.0	-	-	5.1	5.8
AT4G19750	-	Glycosyl hydrolase family with chitinase insertion domain	-	-	3.9	4.8	-	-	6.7	5.3
AT1G19960	-	Putative uncharacterized protein	1.2	1.6	4.0	4.0	-	2.9	5.5	4.4
AT4G10500	DLO1	DMR6-LIKE OXYGENASE 1	-	1.4	3.1	3.8	-	-	3.4	4.2
AT5G41280	CRRSP57	Receptor-like protein kinase-related family protein	-	-	2.6	3.2	-	-	3.2	2.2
AT2G14560	LURP1	late upregulated in response to *Hyaloperonospora parasitica*	-	-	3.0	3.1	-	-	-	4.5
AT1G02230	NAC004	NAC DOMAIN CONTAINING PROTEIN 4	-	2.8	3.8	3.0	-	2.5	3.5	4.3
AT5G54610	BDA1	Ankyrin repeat protein family	-	-	2.2	3.0	-	-	3.0	6.4

Selection of genes upregulated by NHP in Col-0 and *sid2-2* seedlings. Clusters defined for wild type expression are noted (i.e., early transient, early strong, late weak, late strong). Indicated log$_2$ fold change (FC) values determined relative to mock treatment for each genotype and time point. Genes are categorized as SA-independent/partially independent if significant upregulation was observed in *sid2-2* within 15-30 min for the early clusters and within 3-6 h for the late clusters. Not all NHP upregulated genes are shown in this table, the complete list can be found in Supplemental Table S3.

1124

Table 2. SA-dependent NHP upregulated genes by cluster.

AGI	Name	Annotation	Col-0 15m	30m	3h	6h	sid2-2 15m	30m	3h	6h
Early transient										
AT1G32970	SBT3.2	Subtilisin-like protease	2.8	3.2	-	-	-	-	-	-
AT1G21240	WAK3	Wall-associated receptor kinase	2.5	-	-	1.4	-	-	-	2.0
AT5G44430	PDF1.2C	PLANT DEFENSIN 1.2C	2.4	-	-	-	-	-	-	-
AT2G14610	PR1	PATHOGENESIS-RELATED GENE 1	2.3	-	-	-	-	-	-	-
AT4G34410	ERF109	Ethylene-responsive transcription factor	2.1	-	-	-	-	-	-	-
AT2G32140	-	Transmembrane receptor	2.1	-	-	-	-	-	2.0	-
AT5G44420	PDF1.2A	PLANT DEFENSIN 1.2	2.0	-	-	-	-	-	-	-
AT3G14260	-	LURP-one-like protein (DUF567)	1.8	-	-	-	-	-	-	-
AT2G26020	PDF1.2B	PLANT DEFENSIN 1.2B	1.6	-	-	-	-	-	-	-
AT5G64905	PROPEP3	ELICITOR PEPTIDE 3 PRECURSOR	1.5	-	-	-	-	-	-	-
AT1G32640	MYC2	JASMONATE INSENSITIVE 1	1.4	-	-	-	-	-	-	-
AT2G18660	EGC2	PLANT NATRIURETIC PEPTIDE A	1.4	-	-	-	-	-	-	-
Early strong										
AT2G21900	WRKY59	WRKY Transcription Factor; Group II-c	1.9	2.4	2.3	2.3	-	-	2.2	2.7
Late weak										
AT2G28850	CYP710A3	Cytochrome P450 710A3	-	-	2.4	2.1	-	-	-	-
AT1G15630	-	Transmembrane protein	-	-	1.9	1.4	-	-	-	-
AT5G44568	PROSCOOP4	SERINE RICH ENDOGENOUS PEPTIDE	-	-	1.3	1.3	-	-	0.7	1.0
AT1G18830	SEC31A	Protein transport protein SEC31 homolog A	-	-	-	2.8	-	-	-	-
AT2G45550	CYP76C4	Cytochrome P450 76C4	-	-	-	2.6	-	-	-	-
AT1G10155	ATPP2-A10	PHLOEM PROTEIN 2-A10	-	-	-	2.3	-	-	-	-
AT1G02450	NIMIN-1	NIM1-INTERACTING 1	-	-	-	2.3	-	-	-	-
AT3G45860	CRK4	Cysteine-rich receptor-like protein kinase	-	-	-	2.1	-	-	-	-
AT5G51500	PME60	Pectin methylesterase inhibitor	-	-	-	1.7	-	-	-	-
AT2G29350	SAG13	SENESCENCE-ASSOCIATED GENE 13	-	-	-	1.3	-	-	-	-
Late strong										
AT3G22231	PCC1	PATHOGEN AND CIRCADIAN CONTROLLED 1	-	-	3.8	3.9	-	-	-	-

Selection of genes upregulated by NHP in Col-0 and sid2-2 seedlings. Clusters defined for wild type (Col-0) expression are noted. Indicated $\log_2$ fold change (FC) values were determined relative to mock treatment for each genotype and time point. Genes are categorized as SA-dependent if significant upregulation was not observed in sid2-2 within 15-30 min for the early clusters and within 3-6 h for the late clusters. Not all NHP upregulated genes are shown in this table, the complete list can be found in Supplemental Table S3.

1125

 Figure Legends

Figure 1 NHP induces *WRKY38* expression at early time points.

A, Transcript abundance of SAR marker genes *FMO1*, *ICS1*, and *PR1* and early SA-

responsive marker genes *WRKY38* and *UGT76B1* in 4.5-week-old wild type (Col-0)

Arabidopsis plants. Three leaves were infiltrated with sterile water (mock) or 1 mM NHP

and collected after 2, 6, and 24 h for mRNA isolation. Transcript abundance was

measured via quantitative real-time PCR (qRT-PCR) and normalized relative to *UBC21*

(Ubiquitin-Conjugating Enzyme 21; At5g25760). Fold change was determined relative to

mock ($2^{-\Delta\Delta Ct}$). Asterisks indicate a significant difference between mock and NHP per

time point (n = 4; Mann-Whitney *U* test; *$P < 0.05$, ns = not significant). B, Design of

seedling-based assay. Arabidopsis seedlings (15-18 per well) were hydroponically

grown, treated with MS medium (mock), 1 mM Pip, or 1 mM NHP in MS medium then

collected at 15 min, 30 min, 3 h, and 6 h for qRT-PCR analysis of *WRKY38* transcript

abundance in wild type. C, Expression of *WRKY38* in seedlings from the experiment

described in B. Transcript abundance was normalized relative to *UBC21* and mock

treated samples for each condition ($2^{-\Delta\Delta Ct}$). Asterisks indicate a significant difference

between mock and Pip or NHP at each time point (n = 4-6; Mann-Whitney *U* test; *P <

0.01, ns = not significant).

1146

1147

Figure 2 Profile of early NHP-upregulated genes in wild type seedlings.

A, Heatmap of genes upregulated ($\log_2$(FC) > 1, P_{adj} < 0.05) in response to NHP
treatment in wild type seedlings. One biological replicate consists of 15 pooled
seedlings per condition (n = 3). Euclidean distances determined from the $\log_2$(FC)
values were hierarchically clustered across the indicated time points (15 min, 30 min, 3
h and 6 h), resulting in four distinct clusters of gene expression. B, Average $\log_2$(FC) of
all genes within each cluster defined in (A) showing the expression trends across the
four time points. Note, the two genes expressed only at 30 min, and no other time, were
removed from the 'early strong' cluster average. Error bars represent the ± SEM. C,
Biological processes of significantly enriched (FDR < 0.05) Gene Ontology (GO) terms
for the genes within each cluster are shown. Bars represent the -log(FDR) of each
significantly enriched GO term. Bars were omitted if the term was not significantly
enriched for the defined cluster. The most significantly enriched GO term for each
cluster is indicated in bold.
1162

1163

Figure 3 Comparison of NHP upregulated genes in wild type and *sid2-2* seedlings.

A, Percent of NHP upregulated genes expressed only in wild type (Col-0, SA-

dependent) and in both wild type and the *sid2-2* mutant (SA-independent) for the wild

type gene expression clusters defined in Figure 2. Total number of genes in each

condition is indicated. B, Total number of NHP upregulated genes ($\log_2$(FC) > 1, P_{adj} <

0.05) per time point unique to wild type (Col-0), shared between wild type and *sid2-2*,

and unique to *sid2-2*. The number of upregulated genes in each grouping is indicated.

C, Biological processes of significantly enriched (gray bar) or depleted (white bar) (FDR

< 0.05) GO terms for the set of genes upregulated 3 and 6 h after NHP treatment that

were upregulated only in the *sid2-2* mutant and not wild type. Bars represent the

negative log(FDR) of each significantly enriched GO term, bars were omitted if the term

was not significantly enriched or depleted for the defined time point.

1176

Figure 4 Presence of TGA/bZIP and WRKY TF *cis*-regulatory elements (CREs) in the promoters of NHP-upregulated genes in wild type (Col-0) and *sid2-2* seedlings.

A, Schematic of the TF binding motif (i.e., CRE) enrichment analysis performed. All genes with increased transcript abundance upon NHP treatment ($\log_2$(FC) > 0 and P_{adj} < 0.05) were grouped by time point for the indicated genotype, these groups are represented as "test promoter group" in the diagram. CREs of known Arabidopsis TFs from the DAPseq database (O'Malley et al., 2016) were identified in the promoters (1 kb upstream of the transcriptional start site) of each group of promoters. Enrichment of identified CREs was determined relative to CREs found in promoters pulled from a random sampling of genes detected in this RNA-seq experiment. Schematic design is modified from Mariani et al., 2017. Enrichment analysis of TGA/bZIP (B) and WRKY (C)

1188   CREs. The name of the TF known to bind the enriched CRE is listed on the right and

1189   the TF family name is indicated above (i.e., bZIP or WRKY). The -$\log_2$(P value) of the

1190   enrichment analysis is indicated by the scale bar with black indicating no significant

1191   enrichment of the CRE and purple to yellow denoting an enriched CRE.

1192

1193

Figure 5 *WRKY70* is involved in NHP-elicited SAR

Bacterial growth in the distal leaves of mock and NHP treated wild type (Col-0) and *wrky* mutant plants. *WRKY* genes for mutant analysis were selected from the early stable cluster defined in Figure 2 and include *WRKY70* (A), *WRKY38* and *WRKY62* (B), *WRKY54* (C), *WRKY51* (D), and *WRKY59* (E). Three lower (1°) leaves were infiltrated with water (mock), 0.5 mM NHP, or 1 mM NHP (only for *wrky59*) and 1 d later one upper, distal leaf was inoculated with a 1 x 10^5 CFU/mL suspension of *Psm*, followed by quantification of bacterial titer 3 dpi (n = 4-8). Asterisks indicate significant differences in bacterial titer (two-tailed t-test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns = not significant).

1204

1205

Figure 6 Loss of WRKY70 function impacts upregulation of early NHP-responsive genes

A, Diagram of putative WRKY70 CREs, including W-box, WT-box, and WRKY70 (W70) DAPseq motifs in the promoters (-2 Kb upstream of the transcriptional start site; TSS) of NHP-responsive genes *BDA1*, *PROSCOOP4*, and *CML10*. Note W70 DAPseq sites adjacent to a W- or WT-box overlap. Expression of *BDA1* (B), *PROSCOOP4* (C), and *CML10* (D) in *wrky70-1* mutant plants. Three leaves of 4.5-week-old wild type (Col-0) and *wrky70*-1 mutant plants were infiltrated with water or 1 mM NHP. Samples collected at 0 h were untreated before collection. Transcript abundance was determined relative to *UBC21* for each condition ($2^{-\triangle Ct}$). Asterisks indicate a significant difference between

1216   wild type and *wrky70-1* at each time point (two-tailed t-test; $*P < 0.05$, $**P < 0.01$, ns =

1217   not significant).

1218

1219

Figure 7 NHP enhances ROS production and requires WRKY70 for full ROS levels.
Flg22-elicited ROS quantification in *wrky70-1* plants. Four leaf discs from 4.5 to 5-week-
old wild type and *wrky70-1* plants were pretreated by floating leaf discs on water, 0.1
mM SA, or 0.5 mM NHP for 24 h before treating with 100 nM flg22 in horseradish
peroxidase and Luminol. Top, traces of the average relative luminescence units (RLU)
over the indicated run time. Each point represents the average of 6 plants (n = 24) from
two experiments. Bottom, total quantification of RLUs from top plot. For each leaf disc,
RLUs from each time point were summed by condition over the run time and averaged
with error bars representing the standard error of the mean. Statistical analysis was
performed using a one-way ANOVA and post hoc Sidak test, different letters indicated
statistical differences between means with $P < 0.05$.