Structural basis for molecular assembly of fucoxanthin chlorophyll *a/c*-binding proteins in a

diatom photosystem I supercomplex

Koji Kato[1,†], Yoshiki Nakajima[1,†], Jian Xing[2,†], Minoru Kumazawa[2], Haruya Ogawa[1], Jian-Ren

Shen[1,]*, Kentaro Ifuku[2,]*, Ryo Nagao[3,]*

[1]Research Institute for Interdisciplinary Science and Graduate School of Environmental, Life,

Natural Science and Technology, Okayama University, Okayama 700-8530, Japan

[2]Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

[3]Faculty of Agriculture, Shizuoka University, Shizuoka 422-8529, Japan

[†]These authors contributed equally to this work.

*Corresponding Authors:

Jian-Ren Shen, E-mail: shen@cc.okayama-u.ac.jp

Kentaro Ifuku, E-mail: ifuku.kentaro.2m@kyoto-u.ac.jp

Ryo Nagao, E-mail: nagry@shizuoka.ac.jp

Abstract

Photosynthetic organisms exhibit remarkable diversity in their light-harvesting complexes (LHCs).

LHCs are associated with photosystem I (PSI), forming a PSI-LHCI supercomplex. The number of

LHCI subunits, along with their protein sequences and pigment compositions, has been found to

differ greatly among the PSI-LHCI structures. However, the mechanisms by which LHCIs

recognize their specific binding sites within the PSI core remain unclear. In this study, we

determined the cryo-electron microscopy structure of a PSI supercomplex incorporating

fucoxanthin chlorophyll *a*/*c*-binding proteins (FCPs), designated as PSI-FCPI, isolated from the

diatom *Thalassiosira pseudonana* CCMP1335. Structural analysis of PSI-FCPI revealed five FCPI

subunits associated with a PSI monomer; these subunits were identified as RedCAP, Lhcr3, Lhcq10,

Lhcf10, and Lhcq8. Through structural and sequence analyses, we identified specific

protein-protein interactions at the interfaces between FCPI and PSI subunits, as well as among FCPI

subunits themselves. Comparative structural analyses of PSI-FCPI supercomplexes, combined with

phylogenetic analysis of FCPs from *T. pseudonana* and the diatom *Chaetoceros gracilis*, underscore

the evolutionary conservation of protein motifs crucial for the selective binding of individual FCPI

subunits. These findings provide significant insights into the molecular mechanisms underlying the

assembly and selective binding of FCPIs in diatoms.

Introduction

Oxygenic photosynthesis in cyanobacteria, algae, and land plants converts solar energy into chemical energy and releases molecular oxygen into the atmosphere (1). The conversion of light energy takes place within two multi-subunit membrane protein complexes, known as photosystem I (PSI) and photosystem II (PSII), which perform light harvesting, charge separation, and electron transfer reactions (2-5). To optimize light energy capture, numerous light-harvesting antenna subunits are associated with the periphery of the PSI and PSII core complexes, transferring excitation energy to the respective photosystem cores (1). These light-harvesting antennae exhibit significant diversity among photosynthetic organisms, both in protein sequences and pigment compositions, and can be broadly categorized into two major groups: membrane proteins and water-soluble proteins (1).

The membrane protein category primarily consists of the light-harvesting complex (LHC) protein superfamily (6, 7), which absorbs light energy through chlorophylls (Chls) and carotenoids (Cars). The number and types of Chls and Cars vary significantly among LHCs, which can be grouped into green and red lineages, leading to color diversity in photosynthetic organisms (8). The green lineage includes green algae and land plants, while the red lineage encompasses red algae, diatoms, haptophytes, cryptophytes, and dinoflagellates (8). LHCs specific to PSI (LHCIs) bind to a eukaryotic PSI monomer, forming a PSI-LHCI supercomplex (9, 10), the structures of which have been revealed by cryo-electron microscopy (cryo-EM) in various eukaryotes (9, 10). In the red lineage, the number of LHCIs and their protein sequences and pigment compositions exhibit considerable variation among the PSI-LHCI structures of red algae (11-14), a diatom (15, 16), a cryptophyte (17), and dinoflagellates (18, 19).

Recently, we demonstrated the conservation and diversity of LHCIs among red-lineage algae through structural and phylogenetic analyses of PSI-LHCI supercomplexes (14). This study revealed that while the binding sites of LHCIs to PSI were conserved to some extent among red-lineage algae, their evolutionary relationships were weak. It is known that LHCIs have similar overall protein structures across photosynthetic organisms, with particular similarity in their

three-transmembrane helices, regardless of whether they belong to the green or red lineages (9, 10).

However, individual LHCIs have altered their sequences and structures to adapt their respective

binding sites to the PSI cores during the assembly of PSI-LHCI supercomplexes. These

observations raise a critical question: how do LHCIs recognize their binding sites in the PSI core?

Diatoms are among the most essential phytoplankton in aquatic environments, playing a

crucial role in the global carbon cycle, supporting marine food webs, and contributing significantly

to nutrient cycling, thus ensuring the health and sustainability of marine ecosystems (20). Diatoms

possess unique LHCs known as fucoxanthin Chl *a*/*c*-binding proteins (FCPs), which differ in

pigment composition and amino acid sequences from the LHCs of land plants (21-23). Previous

studies have reported the isolation and structural characterization of PSI-FCPI supercomplexes

isolated from the diatom *Chaetoceros gracilis* (15, 16, 24-29). Kumazawa et al. showed significant

diversity in FCPs between *C. gracilis* and *Thalassiosira pseudonana*, with 46 and 44 FCPs

identified, respectively (30). These FCPs are categorized into multiple, closely related subgroups

(30), and their amino acid sequences are not entirely identical between the two diatoms.

Consequently, comparing FCPIs, including their amino acid residues and protein structures at

similar binding sites in PSI-FCPIs, may provide molecular insights into how FCPIs interact with

PSI. However, an overall structure of the *T. pseudonana* PSI-FCPI supercomplex has yet to be

solved.

In this study, we solved the structure of the PSI-FCPI supercomplex from *T. pseudonana*

CCMP1335 at a resolution of 2.30 Å by cryo-EM single-particle analysis. The structure reveals a

PSI-monomer core and five FCPI subunits. Structural and sequence comparisons highlight unique

protein-protein interactions between each FCPI subunit and PSI. Based on these findings, we

discuss the molecular assembly and selective binding mechanisms of FCPI subunits in diatom

species.

Results and discussion

Overall structure of the *T. pseudonana* PSI-FCPI supercomplex

The PSI-FCPI supercomplexes were purified from the diatom *T. pseudonana* CCMP1335 and

analyzed by biochemical and spectroscopic techniques (Fig. S1). Notably, the protein bands of

PSI-FCPI closely resembled those reported in a previous study (31). Cryo-EM images of the

PSI-FCPI supercomplex were obtained using a JEOL CRYO ARM 300 electron microscope

operated at 300 kV. The final cryo-EM map was determined at a resolution of 2.30 Å with a C1

symmetry (Fig. S2, S3, Table S1), based on the "gold standard" Fourier shell correlation (FSC) =

0.143 criterion (Fig. S3A).

The atomic model of PSI-FCPI was built based on the cryo-EM map obtained (see

Methods; Fig. S3, Table S1–S3). The structure reveals a monomeric PSI core associated with five

FCPI subunits (Fig. 1A, B). The five FCPI subunits were named FCPI-1 to 5 (Fig. 1A), following

the nomenclature of LHCI subunits in the PSI-LHCI structure of *Cyanidium caldarium* RK-1

(NIES-2137) (14). Specifically, the positions of FCPI-1 and FCPI-2 in the *T. pseudonana* PSI-FCPI

structure (Fig. 1A) correspond to those of LHCI-1 and LHCI-2 in the *C. caldarium* PSI-LHCI

structure. The PSI core comprises 94 Chls *a*, 18 *β*-carotenes (BCRs), 1 zeaxanthin (ZXT), 3

[4Fe-4S] clusters, 2 phylloquinones, and 6 lipid molecules, whereas the five FCPI subunits include

45 Chls *a*, 7 Chls *c*, 2 BCRs, 15 fucoxanthins (Fxs), 7 diadinoxanthins (Ddxs), and 3 lipid

molecules (Table S3).

Structure of the *T. pseudonana* PSI core

The PSI core contains 12 subunits, 11 of which are identified as PsaA, PsaB, PsaC, PsaD, PsaE,

PsaF, PsaI, PsaJ, PsaL, PsaM, and Psa29 (Fig. 1B). The remaining subunit could not be assigned

due to insufficient map resolution and was therefore modeled as polyalanines (Fig. S4A). This

unidentified subunit, designated as Unknown, occupies the same site as Psa28 in the *C. gracilis*

PSI-FCPI (15). The structural comparison reveals that Unknown closely resembles Psa28 in the *C.*

gracilis PSI-FCPI (Fig. S4B). Psa28, a novel subunit identified in the *C. gracilis* PSI-FCPI structure

(15), follows the previously established nomenclature rule (32). Historically, genes encoding PSI

proteins have been designated as *psaA*, *psaB*, and so forth. PsaZ was identified in the PSI cores of

Gloeobacter violaceus PCC 7421 (33, 34). Subsequent discoveries led to the designation of a new

subunit as Psa27, which was identified in the PSI cores of *Acaryochloris marina* MBIC11017

(35-37). Consequently, we designated this novel subunit as Psa28 (15). However, Xu et al. referred

to this subunit as PsaR in the PSI-FCPI structure of *C. gracilis* (16).

Psa29 is newly identified in the *T. pseudonana* PSI-FCPI structure using ModelAngelo

(38) and the NCBI database (https://www.ncbi.nlm.nih.gov/) (Fig. 2). The subunit corresponding to

Psa29 was also observed previously in the *C. gracilis* PSI-FCPI structures (15, 16), where it was

modeled as polyalanines and referred to as either Unknown1 (15) or PsaS (16). Psa29 exhibits a

unique structure distinct from the other PSI subunits in the *T. pseudonana* PSI-FCPI (Fig. 2A, Fig.

S4C) and engages in multiple interactions with PsaB, PsaC, PsaD, and PsaL at distances of 2.5–3.2

Å (Fig. 2B–G). Sequence analyses suggest that Psa29 has undergone evolutionary divergence

between Bacillariophyceae (diatoms) and Bolidophyceae, the latter of which is a sister group of

diatoms within Stramenopiles (Fig. 2H), although this subunit has not been found in other

organisms. The arrangement of PSI subunits in the *T. pseudonana* PSI-FCPI is virtually identical to

that in the *C. gracilis* PSI-FCPI structures already reported (15, 16). However, the functional and

physiological roles of Psa29 remain unclear at present. It is evident that Psa29 does not have any

pigments, quinones, or metal complexes, suggesting no contribution of Psa29 to electron transfer

reactions within PSI. Further mutagenesis studies will be necessary to investigate the role of Psa29

in diatom photosynthesis.

The number and arrangement of Chls and Cars within the PSI core in the *T. pseudonana*

PSI-FCPI structure (Fig. S4D, E) are largely similar to those in the *C. gracilis* PSI-FCPI structure

(15). However, Chl a102 of PsaI was found in the *T. pseudonana* PSI-FCPI structure but not in the

C. gracilis PSI-FCPI structure (15), whereas a844 of PsaA and BCR843 of PsaB were identified in

the *C. gracilis* PSI-FCPI structure (15) but not in the *T. pseudonana* PSI-FCPI structure. One of the

Car molecules in PsaJ was identified as ZXT103 in the *T. pseudonana* PSI-FCPI structure, while it

is BCR103 in the *C. gracilis* PSI-FCPI structure (15).

Structure of the *T. pseudonana* FCPIs

Kumazawa et al. classified 44 *Lhc* genes in *T. pseudonana*, designating them as *Lhcf*, *Lhcq*, *Lhcr*, *Lhcx*, *Lhcz*, and *CgLhcr9* homologs (30). Based on this classification, the five FCPI subunits in the PSI-FCPI structure were identified using five genes: *RedCAP*, *Lhcr3*, *Lhcq10*, *Lhcf10*, and *Lhcq8*, corresponding to FCPI-1 to 5, respectively (Fig. 1A). It is important to note that *RedCAP* is not included among the 44 *Lhc* genes (30) but is classified within the LHC protein superfamily (6, 7). For the assignment of each FCPI subunit, we focused on characteristic amino acid residues derived from their cryo-EM map, especially S61/V62/Q63 in FCPI-1; A70/R71/W72 in FCPI-2; Y64/R65/E66 in FCPI-3; M63/R64/Y65 in FCPI-4; and A62/R63/R64 in FCPI-5 (Fig. S5). The root mean square deviations (RMSDs) of the structures between FCPI-4 and the other four FCPIs range from 1.91 to 3.73 Å (Table S4).

Each FCPI subunit binds several Chl and Car molecules: 7 Chls *a*/1 Chl *c*/2 Fxs/3 Ddxs/2 BCRs in FCPI-1; 10 Chls *a*/1 Chl *c*/3 Fxs/1 Ddx in FCPI-2; 7 Chls *a*/3 Chls *c*/2 Fxs/2 Ddxs in FCPI-3; 11 Chls *a*/2 Chls *c*/4 Fxs in FCPI-4; and 10 Chls *a*/4 Fxs/1 Ddx in FCPI-5 (Fig. S6A–E, Table S3). The axial ligands of the central Mg atoms of Chls within each FCPI are primarily provided by the main and side chains of amino acid residues (Table S5). Potential excitation-energy-transfer pathways can be proposed based on the close physical interactions among Chls between FCPI-3 and PsaA, between FCPI-3 and PsaL, between FCPI-1 and PsaI, and between FCPI-2 and PsaB (Fig. S7).

Structural characteristics of RedCAP and its evolutionary implications

Among the FCPI subunits, only FCPI-1 contains two BCRs in addition to Fxs and Ddxs (Fig. S6A, F). This is the first report of BCR binding to FCPIs in diatoms. FCPI-1 is identified as RedCAP, a member of the LHC protein superfamily but distinct from the LHC protein family (6, 7); however, the functional and physiological roles of RedCAP remain unknown. FCPI-1 is positioned near PsaB, PsaI, and PsaL through protein-protein interactions with these subunits on both the stromal and lumenal sides (Fig. 3A). At the stromal side, I138 and S139 of FCPI-1 interact with K121, G122,

and F125 of PsaL (Fig. 3B), whereas at the lumenal side, multiple interactions occur between I109

of FCPI-1 and F5 of PsaI, between T105/L106/T108 of FCPI-1 and W92/P94/F96 of PsaB, and

between E102/W103 of FCPI-1 and S71/I73 of PsaL (Fig. 3C). The protein-protein interactions at

the lumenal side (Fig. 3C) appear to be caused by a loop structure of FCPI-1 from Q96 to T116

(pink in Fig. 3D), which is unique to FCPI-1 but absent in the other four FCPI subunits (pink in Fig.

3E). This loop structure is inserted into a cavity formed by PsaB, PsaI, and PsaL (Fig. 3C, D). These

findings indicate that the Q96–T116 loop of FCPI-1 specifically recognizes and binds to the cavity

provided by the PSI subunits.

RedCAP of *C. gracilis* (CgRedCAP) was not identified in the *C. gracilis* PSI-FCPI

structures (15, 16). As previously discussed (14), we proposed that CgRedCAP may bind to the *C.*

gracilis PSI core at a site similar to LHCI-1 in the red alga *C. caldarium* PSI-LHCI through

sequence analysis. This site corresponds to the FCPI-1 site in the PSI-FCPI of *T. pseudonana* in this

study. A sequence alignment between RedCAP of *T. pseudonana* (TpRedCAP) and CgRedCAP is

shown in Fig. S8A, exhibiting a 72% sequence similarity. CgRedCAP contains a protein motif,

Q106–I113 (QWGTLATI), corresponding to E102–I109 (EWGTLATI) in TpRedCAP (Fig. 3C).

These findings suggest the potential binding of CgRedCAP to PSI in *C. gracilis* at a position similar

to FCPI-1 in the *T. pseudonana* PSI-FCPI structure. However, it remains unclear (i) whether

CgRedCAP is indeed bound to the *C. gracilis* PSI-FCPI supercomplex and (ii) if a loop structure

corresponding to the Q96–T116 loop of TpRedCAP exists in CgRedCAP. Further structural studies

of the *C. gracilis* PSI-FCPI are required to elucidate the molecular assembly mechanism of diatom

RedCAPs.

RedCAPs have been found in the PSI-LHCI structures of the red alga *Porphyridium*

purpureum (13) and a PSI supercomplex with alloxanthin Chl *a/c*-binding proteins (PSI-ACPI) in

the cryptophyte *Chroomonas placoidea* (17), as summarized in our previous study (14). Both *P.*

purpureum RedCAP (PpRedCAP) and *C. placoidea* RedCAP (CpRedCAP) exhibit loop structures

similar to the Q96–T116 loop in TpRedCAP observed in the present study (Fig. S8B). Multiple

sequence alignments of TpRedCAP with PpRedCAP and CpRedCAP are shown in Fig. S8C,

revealing sequence similarities of 39% and 60%, respectively. PpRedCAP contains a protein motif of V105–L112 (VWGPLAQL), while CpRedCAP has a protein motif of Q117–A124 (QWGPLASA). These motifs correspond to E102–I109 (EWGTLATI) in TpRedCAP; however, the sequence conservation between TpRedCAP and PpRedCAP/CpRedCAP is lower than between TpRedCAP and CgRedCAP. Among the four RedCAPs, the amino acids Trp, Gly, Leu, and Ala are conserved in the protein motifs (xWGxLAxx), implying that this conserved loop structure contributes to the binding of RedCAP to PSI across the red-lineage algae.

Protein-protein interactions of the other FCPI subunits

FCPI-2 (Lhcr3) is positioned near PsaB and PsaM, engaging in protein-protein interactions with these subunits at distances of 3.0–4.3 Å at both the stromal and lumenal sides (Fig. 4). The amino acid residues I63/T65/D66/Y69/W134/Y138/D140 of FCPI-2 are associated with W153/L154/K159/F160/W166 of PsaB at the stromal side (Fig. 4B), while F116 and F120 of FCPI-2 interact with F5/I9/M12 of PsaM at the lumenal side (Fig. 4C). The amino acid sequences corresponding to I63–Y69, F116–F120, and W134–D140 in Lhcr3 are not conserved in the Lhcr subfamily, comprising Lhcr1, Lhcr4, Lhcr7, Lhcr11, Lhcr12, Lhcr14, Lhcr17, Lhcr18, Lhcr19, and Lhcr20, as reported by Kumazawa et al. (30) (Fig. S9).

FCPI-3 (Lhcq10) is positioned near PsaL, with protein-protein interactions at distances of 2.3–4.2 Å at the stromal side (Fig. 5A, B). The amino acid residues L126/I130/L142/Y146/W147/V148/W155 of FCPI-3 are associated with F4/K6/P20/S25/L26/L30 of PsaL (Fig. 5B). Given the homology between TpLhcq10 and CgLhcr9 (30), we compared the amino acid sequence of Lhcq10 with the Lhcq and Lhcr subfamilies in *T. pseudonana* (Fig. S10A, B). The sequence L126–W155 of Lhcq10 is not conserved in the Lhcq subfamily, comprising Lhcq1, Lhcq2, Lhcq3, Lhcq4, Lhcq5, Lhcq6, Lhcq7, Lhcq8, and Lhcq9 (Fig. S10A), nor in the Lhcr subfamily, comprising Lhcr1, Lhcr3, Lhcr4, Lhcr7, Lhcr11, Lhcr12, Lhcr14, Lhcr17, Lhcr18, Lhcr19, and Lhcr20, as reported by Kumazawa et al. (30) (Fig. S10B).

FCPI-4 (Lhcf10) is positioned near FCPI-5 through protein-protein interactions with it at

distances of 2.6–3.6 Å at the lumenal side (Fig. 5A, C). The amino acid residues Y196/P198/F199

of FCPI-4 interact with F82/F86/G87 of FCPI-5 (Fig. 5C). The amino acid sequence Y196–F199 of

Lhcf10 is not conserved in the Lhcf subfamily, comprising Lhcf1, Lhcf2, Lhcf3, Lhcf4, Lhcf5,

Lhcf6, Lhcf7, Lhcf8, Lhcf9, Lhcf11, and Lhcf12, as reported by Kumazawa et al. (30) (Fig. S11).

FCPI-5 (Lhcq8) is positioned near PsaL and FCPI-4 through protein-protein interactions at

distances of 2.6–4.1 Å at both the stromal and lumenal sides (Fig. 5A, C, D). The amino acid

residues P108/Q109/A112/I115 of FCPI-5 interact with F134/I137/S141 of PsaL at the lumenal side

(Fig. 5D). The interactions between FCPI-5 and FCPI-4 are shown in Fig. 5C. The amino acid

sequences F82–G87 and P107–I115 of Lhcq8 are not conserved in the Lhcq subfamily, comprising

Lhcq1, Lhcq2, Lhcq3, Lhcq4, Lhcq5, Lhcq6, Lhcq7, Lhcq9, and Lhcq10, as reported by Kumazawa

et al.(30) (Fig. S12A, B).

Molecular insights into the assembly of FCPIs in diatom PSI-FCPI supercomplexes

To evaluate the molecular assembly of FCPI subunits in the *T. pseudonana* PSI-FCPI structure, we

focused on protein-protein interactions based on their close proximities (Fig. 3–5) and the amino

acid residues in non-conserved regions among 44 FCPs (Fig. S9–S12). This approach is based on

the premise that selective associations of FCPIs with PSI require specific amino acid residues

unique to each FCPI. Protein-protein interactions among FCPI subunits, as well as between FCPI

and PSI subunits, occur at both the stromal and lumenal sides (Fig. 3–5), and are likely recognized

by unique amino acid residues of FCPIs that are not conserved in each LHC subfamily (Fig.

S9–S12). Thus, the binding and assembly of each FCPI subunit to PSI are likely determined by the

amino acid sequences within the loop regions of the 44 FCPs in *T. pseudonana*.

The diatom *C. gracilis* exhibits two distinct PSI-FCPI structures: one with 16 FCPI

subunits (15) and the other with 24 FCPI subunits (16). These structural variations arise from

changes in the antenna sizes of FCPIs within the *C. gracilis* PSI-FCPI supercomplexes, in response

to varying growth conditions, especially CO_2 concentrations and temperatures (39). Notably, the *C.*

gracilis PSI-FCPI structure contains five FCPI subunits located at the same binding sites as FCPI-1

to 5 in the *T. pseudonana* PSI-FCPI structure (Fig. 6A). A summary of the relationship between the

Lhc genes encoding FCPs, the distinct gene *RedCAP*, and the binding positions of FCPI-1 to 5 in *T.*

pseudonana and *C. gracilis* is shown in Fig. 6B. The gene nomenclature for the *C. gracilis* FCPIs

follows the conventions established by Kumazawa et al. (30), as discussed in our recent study (14).

258   Phylogenetic analysis clearly showed that at the FCPI-1, 2, 3, and 5 sites in the *T.*

pseudonana PSI-FCPI structure, TpRedCAP, TpLhcr3, TpLhcq10, and TpLhcq8 are orthologous to

CgRedCAP, CgLhcr1, CgLhcr9, and CgLhcq12, respectively (Fig. 6C). The characteristic protein

loops of TpRedCAP and CpRedCAP likely participate in interactions with PSI at the FCPI-1 site, as

noted above (Fig. S8). At the FCPI-2 site, comparative analyses revealed that the amino acid

residues facilitating interactions between TpLhcr3 and TpPsaB/TpPsaM closely parallel those

observed in the CgLhcr1-CgPsaB and CgLhcr1-CgPsaM pairs (Fig. S13). Similarly, a high degree

of   similarity   characterized   the   residues   involved   in   the   interaction   pairs   of

TpLhcq10-TpPsaL/CgLhcr9-CgPsaL   at   the   FCPI-3   site   (Fig.   S14A,   B),   as   well   as

TpLhcq8-TpPsaL/CgLhcq12-CgPsaL at the FCPI-5 site (Fig. S14C, D), underscoring the conserved

nature of these interactions. However, TpLhcf10 was not homologous to CgLhcf3 (Fig. 6C), despite

both being located at the FCPI-4 site in their respective PSI-FCPI structures (Fig. 6A). These

findings suggest that the two diatoms possess both a conserved mechanism of protein-protein

interactions across characteristic protein motifs between FCPI and PSI subunits, and a different

mechanism of interactions among FCPIs.

273   It is notable that the *C. gracilis* PSI-FCPI structure binds remarkably more FCPI subunits

than that of *T. pseudonana*, e.g., 16 or 24 subunits in *C. gracilis* as reported in the previous studies

(15, 16), versus 5 subunits in *T. pseudonana* in the present study. The reason for this difference

remains   unclear.   One   possibility   is   that   some   FCPI   subunits   are   released   during   detergent

solubilization in *T. pseudonana*, while they are retained in *C. gracilis*. Alternatively, the number of

FCPI subunits may be inherently lower in *T. pseudonana*, which may reflect adaptations to different

living environments. Further studies are needed to resolve this question.

Extension to molecular assembly of PSI-LHCI supercomplexes

The mechanisms of protein-protein interactions in diatom PSI-FCPI supercomplexes are likely developed by the specific binding of FCPs selected from 44 TpFCPs and 46 CgFCPs in addition to RedCAPs. Like a lock-and-key mechanism, one FCP cannot be substituted by another in forming the PSI-FCPI supercomplexes in the two diatoms; for example, TpLhcq10 binds specifically at the FCPI-3 site but not at the other sites such as FCPI-2. This selective binding mechanism of FCPIs may dictate the molecular assembly of PSI-FCPI. Importantly, the selective binding of FCPIs was identified for the first time by comparing the structures of PSI-FCPI supercomplexes and the amino acid sequences of FCPIs between the two diatom species. This approach can be extended to the LHC protein superfamily in the green and red lineages, enabling comparisons of protein structures and sequences of PSI-LHCI supercomplexes among closely related species. This, in turn, lays the foundation for elucidating the underlying mechanism of PSI-LHCI supercomplex assembly. Thus, this study will shed light on answering the evolutionary question of how LHCIs recognize their binding sites at PSI in photosynthetic organisms.

Methods

Cell growth and preparation of thylakoid membranes

The marine centric diatom *T. pseudonana* CCMP1335 was grown in artificial seawater supplemented with sodium metasilicate and KW21 (40) at 20°C under a photosynthetic photon flux density of 30 μmol photons m^{-2} s^{-1} provided by white LED, with bubbling of air containing 3% (v/v) CO_2. The cells were harvested by centrifugation, disrupted by agitation with glass beads (41), and the thylakoid membranes were pelleted by further centrifugation. The resulting thylakoid membranes were suspended in 50 mM Mes-NaOH (pH 6.5) buffer containing 1 M betaine and 1 mM EDTA.

Purification of the PSI-FCPI supercomplex

Thylakoid membranes were solubilized with 1% (w/v) n-dodecyl-β-D-maltoside (β-DDM) at a Chl

concentration of 0.5 mg mL^{-1} for 20 min on ice in the dark with gentle stirring. After centrifugation

at 162,000 × g for 20 min at 4°C, the supernatant was loaded onto a Q-Sepharose anion-exchange

column (1.6 cm inner diameter, 25 cm length) equilibrated with 20 mM Mes-NaOH (pH 6.5) buffer

containing 0.2 M trehalose, 5 mM CaCl$_2$, and 10 mM MgCl$_2$, and 0.03% β-DDM (buffer A). The

column was washed with buffer A until the eluate became colorless. Elution was performed at a

flow rate of 1.0 mL min^{-1} using a linear gradient of buffer A and buffer B (buffer A plus 500 mM

NaCl) with the following time and gradient: 0–600 min, 0–60% buffer B; 600–800 min, 60–100%

buffer B; 800–900 min, 100% buffer B. The PSI-FCPI-enriched fraction was eluted at 194–247 mM

NaCl, then collected and subsequently loaded onto a linear gradient containing 10–40% (w/v)

trehalose in 20 mM Mes-NaOH (pH 6.5) buffer containing 5 mM CaCl$_2$, 10 mM MgCl$_2$, 100 mM

NaCl, and 0.03% β-DDM. After centrifugation at 154,000 × g for 18 h at 4°C (P40ST rotor;

Hitachi), a green fraction (Fig. S1A) was collected and concentrated using a 150 kDa cut-off filter

(Apollo; Orbital Biosciences) at 4,000 × g. The concentrated samples were stored in liquid nitrogen

until use.

Biochemical and spectroscopic analyses of the PSI-FCPI supercomplex

The polypeptide bands of PSI-FCPI were analyzed by SDS- PAGE with 16% (w/v) acrylamide and

7.5 M urea, following the method of Ikeuchi and Inoue (42) (Fig. S1B). The PSI-FCPI

supercomplexes (4 µg of Chl) were solubilized in 3% lithium lauryl sulfate and 75 mM

dithiothreitol for 10 min at 60°C, and then loaded onto the gel. A standard molecular weight marker

(SP-0110; APRO Science) was used. The absorption spectrum of PSI-FCPI was measured at room

temperature using a UV-Vis spectrophotometer (UV-2450; Shimadzu) (Fig. S1C), and the

fluorescence emission spectrum of PSI-FCPI was measured at 77 K upon excitation at 430 nm

using a spectrofluorometer (RF-5300PC; Shimadzu) (Fig. S1D). The pigment composition of

PSI-FCPI was analyzed by HPLC following the method of Nagao et al. (43), and the elution profile

was monitored at 440 nm (Fig. S1E).

Cryo-EM data collection

A 3-µL aliquot of the *T. pseudonana* PSI-FCPI supercomplex (3.0 mg of Chl mL^{-1}) in 20 mM

MES-NaOH (pH 6.5) buffer containing 0.5 M betaine, 5 mM CaCl$_2$, 10 mM MgCl$_2$, and 0.03%

β-DDM was applied to Quantifoil R1.2/1.3 Cu 300 mesh grids in the chamber of FEI Vitrobot Mark

IV (Thermo Fisher Scientific). The grid was then blotted with filter paper for 4 sec at 4°C under

100% humidity and plunged into liquid ethane cooled by liquid nitrogen. The frozen grid was

transferred to a CRYO ARM 300 electron microscope (JEOL) equipped with a cold-field emission

gun operated at 300 kV. All image stacks were collected from 5 × 5 holes per stage adjustment to

the central hole and image shifts were applied to the surrounding holes while maintaining an axial

coma-free condition. The images were recorded using an in-column energy filter with a slit width of

20 eV at a nominal magnification of × 60,000 on a direct electron detector (Gatan K3, AMETEK).

The nominal defocus range was −1.8 to −1.2 µm, and the physical pixel size corresponded to 0.752

Å. Each image stack was exposed at a dose rate of 21.46 e$^-$Å^{-2}sec^{-1} for 2.33 sec in CDS mode, with

dose-fractionated 50 movie frames. A total of 8,950 image stacks were collected.

Cryo-EM image processing

The resultant movie frames were aligned and summed using MotionCor2 (44) to produce

dose-weighted images. The contrast transfer function (CTF) estimation was performed using

CTFFIND4 (45). All subsequent processes were carried out using RELION-4.0 (46). A total of

2,733,572 particles were automatically picked and subjected to reference-free 2D classification.

From these, 1,132,721 particles were selected from well-defined 2D classes and further processed

for 3D classification without imposing any symmetry. An initial model for the first 3D

classification was generated *de novo* from the 2D classification. A 240-Å spherical mask was used

during the 3D classification and refinement processes. As illustrated in Fig. S2C, the final PSI-FCPI

structure was reconstructed from 75,667 particles. The overall resolution of the cryo-EM map was

determined to be 2.30 Å, based on the gold-standard FSC curve with a cut-off value of 0.143 (Fig.

S3A) (47). Local resolutions were calculated using RELION (Fig. S3C).

Model building and refinement

Two types of the cryo-EM maps were employed for the model building of the PSI-FCPI

supercomplex: a postprocessed map and a denoised map generated using Topaz version 0.2.4 (48).

The postprocessed map was denoised using a trained model over 100 epochs using two half-maps.

Initial models of each subunit in the PSI-FCPI supercomplex were generated by ModelAngelo (38)

and subsequently inspected and manually adjusted against the maps with Coot (49). Each model

was built based on interpretable features from the density maps at a contour level of 2.5 σ in both

the denoised and postprocessed maps. For the assignment of Chls, Chls a and c were distinguished

by inspecting the density map corresponding to the phytol chain at the least level not to link the map

of Chls with that of noise. All Chls c were assigned as Chl c1 due to the inability to distinguish

between Chl c1 and Chl c2 at the present resolution. For the assignment of Cars, Fx and Ddx were

distinguished based on the density surrounding the head groups of Cars with the above threshold.

The PSI-FCPI structure was refined using phenix.real_space_refine (50) and Servalcat (51),

incorporating geometric restraints for protein-cofactor coordination. The final model was validated

with MolProbity (52), EMRinger (53), and Q-score (54). The statistics for all data collection and

structure refinement are summarized in Table S1, S2. All structural figures were prepared using

PyMOL (55), UCSF Chimera (56), and UCSF ChimeraX (57). Since the numbering of Chls, Cars,

and other cofactors in this paper differs from those in the PDB data, the corresponding relationships

are provided in Table S6–S8.

Phylogenetic analysis

Amino acid sequences were aligned using MAFFT L-INS-i v7.490 or MAFFT E-INS-i v7.520 (58).

The alignment was trimmed using ClipKit v1.4.1 with the smart-gap mode. Phylogenetic trees were

inferred using IQ-TREE 2 (59) with the model selected by ModelFinder (60). The trees were

visualized using iTOL v6 (61). Ultrafast bootstrap approximation was performed with 1000

replicates (62).

Data availability

Atomic coordinates, cryo-EM maps, and raw image data for the reported structure have been

deposited in the Protein Data Bank under an accession code 8XLS

[https://www.rcsb.org/structure/8XLS], in the Electron Microscopy Data Bank under an accession

code EMD-38457 [https://www.ebi.ac.uk/emdb/EMD-38457], and in the Electron Microscopy

Public Image Archive under an accession code EMPIAR-12142

[https://doi.org/10.6019/EMPIAR-12142], respectively.

References

1. R. E. Blankenship, *Molecular Mechanisms of Photosynthesis* (Wiley-Blackwell, Oxford, U.K., ed. 3rd, 2021).

2. J. H. Golbeck, Structure and function of photosystem I. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* **43**, 293-324 (1992).

3. K. Brettel, W. Leibl, Electron transfer in photosystem I. *Biochim. Biophys. Acta, Bioenerg.* **1507**, 100-114 (2001).

4. J.-R. Shen, The structure of photosystem II and the mechanism of water oxidation in photosynthesis. *Annu. Rev. Plant Biol.* **66**, 23-48 (2015).

5. D. Shevela, J. F. Kern, G. Govindjee, J. Messinger, Solar energy conversion by photosystem II: principles and structures. *Photosynth. Res.* **156**, 279-307 (2023).

6. J. Engelken, H. Brinkmann, I. Adamska, Taxonomic distribution and origins of the extended LHC (light-harvesting complex) antenna protein superfamily. *BMC Evol. Biol.* **10**, 233 (2010).

7. S. Sturm *et al.*, A novel type of light-harvesting antenna protein of red algal origin in algae with secondary plastids. *BMC Evol. Biol.* **13**, 159 (2013).

8. P. G. Falkowski *et al.*, The evolution of modern eukaryotic phytoplankton. *Science* **305**, 354-360 (2004).

9. M. Hippler, N. Nelson, The plasticity of photosystem I. *Plant Cell Physiol.* **62**, 1073-1081 (2021).

10. J.-R. Shen, "Structure, function, and variations of the photosystem I-antenna supercomplex from different photosynthetic organisms" in Macromolecular Protein Complexes IV. Subcellular Biochemistry, J. R. Harris, J. Marles-Wright, Eds. (Springer, Cham., 2022), vol. 99, pp. 351-377.

11. X. Pi *et al.*, Unique organization of photosystem I-light-harvesting supercomplex revealed by cryo-EM from a red alga. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 4423-4428 (2018).

12. M. Antoshvili, I. Caspy, M. Hippler, N. Nelson, Structure and function of photosystem I in *Cyanidioschyzon merolae*. *Photosynth. Res.* **139**, 499-508 (2019).

13. X. You *et al.*, In situ structure of the red algal phycobilisome-PSII-PSI-LHC megacomplex. *Nature* **616**, 199-206 (2023).

14. K. Kato *et al.*, The structure of PSI-LHCI from *Cyanidium caldarium* provides evolutionary insights into conservation and diversity of red-lineage LHCs. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2319658121 (2024).

15. R. Nagao *et al.*, Structural basis for assembly and function of a diatom photosystem I-light-harvesting supercomplex. *Nat. Commun.* **11**, 2481 (2020).

16. C. Xu *et al.*, Structural basis for energy transfer in a huge diatom PSI-FCPI supercomplex. *Nat. Commun.* **11**, 5081 (2020).

17. L.-S. Zhao *et al.*, Structural basis and evolution of the photosystem I-light-harvesting supercomplex of cryptophyte algae. *Plant Cell* **35**, 2449-2463 (2023).

18. X. Li *et al.*, Structures and organizations of PSI-AcpPCI supercomplexes from red tidal and coral symbiotic photosynthetic dinoflagellates. *Proc. Natl. Acad. Sci. U. S. A.* **121**,

437   e2315476121 (2024).

19. L.-S. Zhao *et al.*, Architecture of symbiotic dinoflagellate photosystem I-light-harvesting supercomplex in *Symbiodinium*. *Nat. Commun.* **15**, 2392 (2024).

20. C. B. Field, M. J. Behrenfeld, J. T. Randerson, P. Falkowski, Primary production of the biosphere: Integrating terrestrial and oceanic components. *Science* **281**, 237-240 (1998).

21. B. R. Green, D. G. Durnford, The chlorophyll-carotenoid proteins of oxygenic photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **47**, 685-714 (1996).

22. C. Büchel, Light harvesting complexes in chlorophyll *c*-containing algae. *Biochim. Biophys. Acta Bioenerg.* **1861**, 148027 (2020).

23. W. Wang, J.-R. Shen, "Structure, organization and function of light-harvesting complexes associated with photosystem II" in Photosynthesis: molecular approaches to solar energy conversion, J.-R. Shen, K. Satoh, S. I. Allakhverdiev, Eds. (Springer, Cham., 2021), pp. 163-194.

24. Y. Ikeda *et al.*, Photosystem I complexes associated with fucoxanthin-chlorophyll-binding proteins from a marine centric diatom, *Chaetoceros gracilis*. *Biochim. Biophys. Acta, Bioenerg.* **1777**, 351-361 (2008).

25. R. Nagao, K. Kagatani, Y. Ueno, J.-R. Shen, S. Akimoto, Ultrafast excitation energy dynamics in a diatom photosystem I-antenna complex: a femtosecond fluorescence upconversion study. *J. Phys. Chem. B* **123**, 2673-2678 (2019).

26. R. Nagao *et al.*, Biochemical characterization of photosystem I complexes having different subunit compositions of fucoxanthin chlorophyll *a*/*c*-binding proteins in the diatom *Chaetoceros gracilis*. *Photosynth. Res.* **140**, 141-149 (2019).

27. R. Nagao, M. Yokono, Y. Ueno, J.-R. Shen, S. Akimoto, Low-energy chlorophylls in fucoxanthin chlorophyll *a*/*c*-binding protein conduct excitation energy transfer to photosystem I in diatoms. *J. Phys. Chem. B* **123**, 66-70 (2019).

28. R. Nagao, M. Yokono, Y. Ueno, J.-R. Shen, S. Akimoto, pH-sensing machinery of excitation energy transfer in diatom PSI-FCPI complexes. *J. Phys. Chem. Lett.* **10**, 3531-3535 (2019).

29. R. Nagao, M. Yokono, Y. Ueno, J.-R. Shen, S. Akimoto, Excitation-energy transfer and quenching in diatom PSI-FCPI upon P700 cation formation. *J. Phys. Chem. B* **124**, 1481-1486 (2020).

30. M. Kumazawa *et al.*, Molecular phylogeny of fucoxanthin-chlorophyll *a*/*c* proteins from *Chaetoceros gracilis* and Lhcq/Lhcf diversity. *Physiol. Plant.* **174**, e13598 (2022).

31. Y. Ikeda *et al.*, Two types of fucoxanthin-chlorophyll-binding proteins I tightly bound to the photosystem I core complex in marine centric diatoms. *Biochim. Biophys. Acta* **1827**, 529-539 (2013).

32. Y. Kashino *et al.*, Proteomic analysis of a highly active photosystem II preparation from the cyanobacterium *Synechocystis* sp. PCC 6803 reveals the presence of novel polypeptides. *Biochemistry* **41**, 8004-8012 (2002).

33. H. Inoue *et al.*, Unique constitution of photosystem I with a novel subunit in the cyanobacterium *Gloeobacter violaceus* PCC 7421. *FEBS Lett.* **578**, 275-279 (2004).

477   34.   K. Kato *et al.*, Structural basis for the absence of low-energy chlorophylls in a photosystem I

478   trimer from *Gloeobacter violaceus*. *eLife* **11**, e73990 (2022).

479   35.   T. Tomo *et al.*, Characterization of highly purified photosystem I complexes from the

480   chlorophyll *d*-dominated cyanobacterium *Acaryochloris marina* MBIC 11017. *J. Biol. Chem.*

481   **283**, 18198-18209 (2008).

482   36.   T. Hamaguchi *et al.*, Structure of the far-red light utilizing photosystem I of *Acaryochloris*

483   *marina*. *Nat. Commun.* **12**, 2333 (2021).

484   37.   C. Xu *et al.*, A unique photosystem I reaction center from a chlorophyll *d*-containing

485   cyanobacterium *Acaryochloris marina*. *J. Integr. Plant Biol.* **63**, 1740-1752 (2021).

486   38.   K. Jamali *et al.*, Automated model building and protein identification in cryo-EM maps.

487   *Nature* 10.1038/s41586-024-07215-4 (2024).

488   39.   R. Nagao, Y. Ueno, S. Akimoto, J.-R. Shen, Effects of CO_2 and temperature on photosynthetic

489   performance in the diatom *Chaetoceros gracilis*. *Photosynth. Res.* **146**, 189-195 (2020).

490   40.   R. Nagao *et al.*, Isolation and characterization of oxygen-evolving thylakoid membranes and

491   Photosystem II particles from a marine diatom *Chaetoceros gracilis*. *Biochim. Biophys. Acta,*

492   *Bioenerg.* **1767**, 1353-1362 (2007).

493   41.   R. Nagao, M. Yamaguchi, S. Nakamura, H. Ueoka-Nakanishi, T. Noguchi, Genetically

494   introduced hydrogen bond interactions reveal an asymmetric charge distribution on the radical

495   cation of the special-pair chlorophyll P680. *J. Biol. Chem.* **292**, 7474-7486 (2017).

496   42.   M. Ikeuchi, Y. Inoue, A new photosystem II reaction center component (4.8 kDa protein)

497   encoded by chloroplast genome. *FEBS Lett.* **241**, 99-104 (1988).

498   43.   R. Nagao, M. Yokono, S. Akimoto, T. Tomo, High excitation energy quenching in fucoxanthin

499   chlorophyll *a*/*c*-binding protein complexes from the diatom *Chaetoceros gracilis*. *J. Phys.*

500   *Chem. B* **117**, 6888-6895 (2013).

501   44.   S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for improved

502   cryo-electron microscopy. *Nat. Methods* **14**, 331-332 (2017).

503   45.   J. A. Mindell, N. Grigorieff, Accurate determination of local defocus and specimen tilt in

504   electron microscopy. *J. Struct. Biol.* **142**, 334-347 (2003).

505   46.   D. Kimanius, L. Dong, G. Sharov, T. Nakane, S. H. W. Scheres, New tools for automated

506   cryo-EM single-particle analysis in RELION-4.0. *Biochem. J.* **478**, 4169-4185 (2021).

507   47.   N. Grigorieff, S. C. Harrison, Near-atomic resolution reconstructions of icosahedral viruses

508   from electron cryo-microscopy. *Curr. Opin. Struc. Biol.* **21**, 265-273 (2011).

509   48.   T. Bepler, K. Kelley, A. J. Noble, B. Berger, Topaz-Denoise: general deep denoising models

510   for cryoEM and cryoET. *Nat. Commun.* **11**, 5208 (2020).

511   49.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of *Coot*. *Acta*

512   *Crystallogr. D Biol. Crystallogr.* **66**, 486-501 (2010).

513   50.   P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular

514   structure solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213-221 (2010).

515   51.   K. Yamashita, C. M. Palmer, T. Burnley, G. N. Murshudov, Cryo-EM single-particle structure

516   refinement and map calculation using Servalcat. *Acta Crystallogr. D Struct. Biol.* **77**,

1282-1291 (2021).

52.   V. B. Chen *et al.*, MolProbity: all-atom structure validation for macromolecular

crystallography. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 12-21 (2010).

53.   B. A. Barad *et al.*, EMRinger: side chain-directed model and map validation for 3D

cryo-electron microscopy. *Nat. Methods* **12**, 943-946 (2015).

54.   G. Pintilie *et al.*, Measurement of atom resolvability in cryo-EM maps with *Q*-scores. *Nat.*

Methods **17**, 328-334 (2020).

55.   L. L. C. Schrödinger, The PyMOL Molecular Graphics System. Version 2.5.0.   (2021).

56.   E. F. Pettersen *et al.*, UCSF Chimera - A visualization system for exploratory research and

analysis. *J. Comput. Chem.* **25**, 1605-1612 (2004).

57.   E. F. Pettersen *et al.*, UCSF ChimeraX: structure visualization for researchers, educators, and

developers. *Protein Sci.* **30**, 70-82 (2021).

58.   K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772-780 (2013).

59.   B. Q. Minh *et al.*, IQ-TREE 2: new models and efficient methods for phylogenetic inference in

the genomic era. *Mol. Biol. Evol.* **37**, 1530-1534 (2020).

60.   S. Kalyaanamoorthy, B. Q. Minh, T. K. F. Wong, A. von Haeseler, L. S. Jermiin, ModelFinder:

fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587-589 (2017).

61.   I. Letunic, P. Bork, Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. *Nucleic Acids Res.* **49**, W293-W296 (2021).

62.   D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2: improving the

ultrafast bootstrap approximation. *Mol. Biol. Evol.* **35**, 518-522 (2018).

Acknowledgements

We thank Kumiyo Kato and Satoko Kakiuchi for their assistance in this study. The cells of *T.*

pseudonana CCMP1335 were given by Prof. Yusuke Matsuda, Kwansei Gakuin University, Japan.

Cryo-EM data was obtained using EM01CT and EM02CT of SPring-8 with the approval of the

Japan Synchrotron Radiation Research Institute (JASRI Proposal No. 2022B2728 (J.-R.S.) and No.

2023A2715 (Y.N.)). This work was supported by JSPS KAKENHI grant Nos. JP22KJ2017 (M.K.),

JP23K14211 (Y.N.), JP22H04916 (J.-R.S.), JP23H02347 (K.I.), and JP23H02423 (R.N.), Takeda

Science Foundation (R.N., K.K.), and Research Support Project for Life Science and Drug

Discovery (Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS))

from AMED under support No. 4176 (J.-R.S.).

Author Contributions

R.N. conceived the project; H.O. and R.N. prepared the PSI-FCPI supercomplexes and analyzed

their biochemical characteristics; J.X., M.K., and K.I. performed phylogenetic and sequence

analyses; Y.N. collected cryo-EM images; K.K. processed the cryo-EM data and reconstructed the

final cryo-EM map; K.K. built the structural model and refined the final model; K.K., Y.N., M.K.,

K.I., and R.N. drafted the original manuscript; J.-R.S. modified the manuscript; and R.N. wrote the

final manuscript, and all authors joined the discussion of the results.

Declaration of competing interest

The authors declare no conflict of interest.

Fig. 1 | Overall structure of the PSI-FCPI supercomplex from *T. pseudonana*.

Structures are viewed from the stromal side (left panels) and from the direction perpendicular to the membrane normal (right panels). Only protein structures are depicted, with cofactors omitted for clarity. The FCPI (**A**) and PSI-core (**B**) subunits are labeled and colored distinctly. The five FCPI subunits are labeled as FCPI-1 to 5 (red), with their corresponding gene products indicated in parentheses (black) in panel **A**.

Fig. 2 | Structure and diversity of Psa29.

(**A**) Structure of Psa29 depicted as cartoons. Psa29 was modeled from V47 to L178. (**B**) Cryo-EM

map of Psa29 and its surrounding environment, viewed from the stromal side. The red-squared

areas are enlarged in panels **C**–**G**. Yellow, PsaB; cyan, PsaC; blue, PsaD; magenta, PsaL; orange,

Psa29. (**C**–**G**) Protein-protein interactions of Psa29 with PsaB/PsaC (**C**), PsaC/PsaD (**D**),

PsaB/PsaD (**E**), PsaB/PsaD/PsaL (**F**), and PsaD (**G**). Interactions are indicated by dashed lines, and

the numbers are distances in Å. Amino acid residues participating in the interactions are labeled; for

example, A71/C indicates Ala71 of PsaC. B, PsaB; C, PsaC; D, PsaD, L, PsaL; 29, Psa29. (**H**)

Phylogenetic analysis of Psa29 in photosynthetic organisms. A maximum-likelihood tree of Psa29

proteins was inferred using IQ-TREE v2.2.2.7 with the WAG+F+G4 model and a trimmed

alignment of 22 sequences comprising 245 amino acid residues. Numbers at the nodes represent

ultrafast bootstrap support (%) (1000 replicates). The tree was mid-point rooted between diatoms

and Bolidophyceae Parmales. Psa29 of *T. pseudonana* CCMP1335 is indicated by a red underline.

Fig. 3 | Structural characteristics of FCPI-1 (RedCAP).

(**A**) Interactions of FCPI-1 with PsaB, PsaI, and PsaL viewed from the stromal (left) and lumenal

(right) sides. The areas encircled by black squares are enlarged in panels **B** and **C**. Yellow, PsaB;

magenta, PsaI; dark red, PsaL; red, FCPI-1. (**B**, **C**) Protein-protein interactions of FCPI-1 with PsaL

(**B**) and with PsaB/PsaI/PsaL (**C**). Interactions are indicated by dashed lines, and the numbers are

distances in Å. Amino acid residues involved in the interactions are labeled; for example, S139/1

indicates Ser139 of FCPI-1. B, PsaB; I, PsaI; L, PsaL; 1, FCPI-1. (**D**) Characteristic loop structure

from Q96 to T116 in FCPI-1, viewed from the lumenal side. Q96 and T116 are labeled with sticks,

and the Q96–T116 loop is colored pink. (**E**) Superpositions of FCPI-1 with FCPI-2, FCPI-3, FCPI-4,

and FCPI-5. Only proteins are depicted. Q96 and T116 in the Q96–T116 loop of FCPI-1 are shown

with sticks.

Fig. 4 | Structural characteristics of FCPI-2.

(**A**) Interactions of FCPI-2 with PsaB and PsaM viewed from the stromal (left) and lumenal (right)

sides. The areas encircled by black squares are enlarged in panels **B** and **C**. PSI subunits are colored

grey, and FCPI subunits are colored yellow. Protein-protein interactions are shown in different

colors: green, FCPI-2; cyan, PsaB; pink, PsaM. (**B**, **C**) Protein-protein interactions of FCPI-2 with

PsaB (**B**) and PsaM (**C**). Interactions are indicated by dashed lines, and the numbers represent

distances in Å. Amino acid residues involved in the interactions are labeled; for example, Y138/2

indicates Tyr138 of FCPI-2. B, PsaB; M, PsaM; 2, FCPI-2.

Fig. 5 | Structural characteristics of FCPI-3, 4, and 5.

(**A**) Interactions among FCPIs and between FCPIs and PsaL, viewed from the stromal (left) and

lumenal (right) sides. The areas encircled by black squares are enlarged in panels **B**–**D**. PSI subunits

are colored grey, and FCPI subunits are colored yellow. Protein-protein interactions are shown in

different colors: blue, FCPI-3; magenta, FCPI-4; orange, FCPI-5; purple, PsaL. (**B**–**D**)

Protein-protein interactions between FCPI-3 and PsaL (**B**), between FCPI-4 and FCPI-5 (**C**), and

between FCPI-5 and PsaL (**D**). Interactions are indicated by dashed lines, and the numbers represent

distances in Å. Amino acid residues involved in the interactions are labeled; for example, L126/3

indicates Leu126 of FCPI-3. L, PsaL; 3, FCPI-3; 4, FCPI-4; 5, FCPI-5.

Fig. 6. Comparisons of structures and sequences of FCPIs in the PSI-FCPI structures between *T. pseudonana* and *C. gracilis*.

(**A**) Superposition of the PSI-FCPI structures between *T. pseudonana* and *C. gracilis* (PDB: 6LY5). FCPI subunits from *T. pseudonana* and *C. gracilis* are colored red and cyan, respectively. The structures are viewed from the stromal side. The FCPI-1 to 5 sites are labeled. (**B**) Correlation of the names of FCPIs in the structures with their corresponding genes between *T. pseudonana* and *C. gracilis*. The FCPI genes are derived from Kumazawa et al. (30) and Kato et al. (14) for *C. gracilis*. (**C**) Phylogenetic analysis of FCPs and RedCAPs from *T. pseudonana* (Tp) and *C. gracilis* (Cg). In addition to the RedCAP family, 44 TpFCPs and 46 CgFCPs are grouped into five Lhc subfamilies

632 and CgLhcr9 homologs. Maroon, RedCAP family; magenta, Lhcq subfamily; red, Lhcz subfamily;

633 orange, Lhcr subfamily; brown, CgLhcr9 homologs; green, Lhcf subfamily; blue, Lhcx subfamily.

634 The FCPs and RedCAPs located at the FCPI-1 to 5 sites are labeled. The tree was inferred using

635 IQ-TREE 2 (59) with the Q.pfam + R4 model selected by ModelFinder (60). The light purple

636 circular symbols on the tree represent bootstrap support (%).