Association of diet and inflammation with the vaginal microbiota of pregnant

individuals with or without IBD

Daniela Vargas-Robles[1], Yan Rou Yap[1], Biplab Singha[1], Joyce Tien[2], Mallika Purandare[2], Mayra Rojas-

Correa[1], Camilla Madziar[1], Mellissa Picker[3], Tina Dumont[4]; Heidi Leftwich[4], Christine F. Frisard[5], Doyle

V. Ward[1], Inga Peter[3], Barbara Olendzki[5], Ana Maldonado-Contreras[1]*

[1] Department of Microbiology and Physiological Systems, Program of Microbiome Dynamics; [2] School of

Medicine; [5] Department of Population and Quantitative Health Science. University of Massachusetts

Chan Medical School, Worcester, MA, 01655

[3] Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York,

NY, 10029

[4] Department of Obstetrics and Gynecology, Division of Maternal-Fetal Medicine, University of

Massachusetts Chan Medical School, Worcester, MA, 01655

*Corresponding author: Ana Maldonado-Contreras.

Email: ana.maldonado@umassmed.edu

Telephone: +1 774-455-3697

Short title: Inflammation, diet, and the vaginal microbiota.

The authors report no conflict of interest.

Why was the study conducted?

An altered vaginal microbiota has been implicated in preterm birth. There is no research on the vaginal microbiome and the factors that influence it in pregnant individuals with Inflammatory Bowel Disease (IBD) at a higher risk of preterm delivery.

Key findings

Pregnant individuals with IBD exhibit a comparable vaginal microbiome to healthy pregnant individuals. However, pregnant individuals with IBD present a vaginal immune profile characterized by increased levels of Th17 pro-inflammatory cytokines. High dietary quality, and optimal consumption of vegetables and added sugars were associated with vaginal dominance by the beneficial *L. crispatus*.

What does this add to what is known?

Our results indicate that the vaginal immune environment and not the microbiome might explain poor pregnancy outcomes for individuals with IBD. Moreover, our study supports the importance of diet to favor *L. crispatus,* a bacterium associated with a lower risk of preterm birth.

Abstract

Background and aims: Vaginal dysbiosis has been associated with adverse pregnancy outcomes. Here, we characterized the vaginal microbiota of pregnant individuals with inflammatory bowel disease (IBD) and investigated whether gut or vaginal inflammation and diet influence the vaginal microbiota diversity of these individuals.

Study Design: We recruited 48 individuals in their third trimester of pregnancy (IBD=23 and HC=18). We characterized the vaginal microbiota by *16S rRNA* sequencing and the gut microbiota by shotgun sequencing. We measured fecal calprotectin in stool and pro-inflammatory cytokines in vaginal fluids. We determine dietary quality using validated 24-hour dietary recalls.

Results: Pregnant individuals with IBD exhibit higher levels of fecal calprotectin and increased expression of Th17 pro-inflammatory cytokines (i.e., IL-6, IL-8, IL-17) in the vaginal mucosa compared to healthy pregnant individuals. High fecal calprotectin correlated with high vaginal microbiota diversity. Also, IL-4 (reduced in IBD) was associated with vaginal microbial composition. Regardless of IBD status, pregnant individuals with healthier diets and particularly optimal servings of vegetables and sugars exhibited a vaginal microbiota dominated by *Lactobacillus crispatus*, a species associated with a lower risk of preterm birth and bacterial vaginosis.

Conclusion: Besides gut inflammation, pregnant individuals with IBD also exhibit a Th17 immune tone in the vaginal mucosa. The vaginal microbiota diversity or composition, particularly high in the beneficial *L. crispatus,* is positively associated with healthier diets, regardless of IBD status.

Keywords: Vaginal, microbiota, IBD, inflammation, pregnancy

Introduction

The vaginal microbiota composition of pregnant individuals with IBD is stable throughout pregnancy but has not been directly compared to a healthy cohort (1). Only one study has reported that pregnant individuals with IBD were more likely to have vaginal infections compared to healthy pregnant individuals (2). In general, healthy pregnancies are characterized by vaginal microbiotas with low bacterial diversity and dominant *Lactobacillus* species (3). For instance, pregnant individuals with a vaginal microbiota dominated by *L. crispatus* have a lower risk of preterm birth compared to individuals with low-lactobacilli abundance (4). Moreover, vaginal microbial communities dominated by *L. crispatus* are better protected against infections than those dominated by *L. iners* (5). Therefore, understanding *Lactobacillus* species dominance in individuals with IBD is important to predicting their risk of poor pregnancy outcomes. Our goal is to characterize and compare the vaginal microbiota of pregnant individuals with and without IBD using high-throughput microbiota sequencing.

Moreover, we sought to determine the role of environmental factors, such as diet and local inflammation on the vaginal microbiota composition. To our knowledge, there have been only a few studies evaluating the influence of diet on the vaginal microbiota of pregnant individuals using high throughput microbiota sequencing (6-8), yet none of the studies included all the relevant diet components from validated instruments aiming at measuring dietary intake. Additionally, the immune tone of the vaginal mucosa in pregnant women with IBD remains understudied. Hence, we will test whether diet influences the vaginal microbiota makeup thus diet can be potentially used as a strategy to revert vaginal dysbiosis during pregnancy, and whether an inflammatory environment in the gut and vagina are linked to vaginal microbiota profiles.

Materials and Methods

Recruitment: We conducted a case–control study nested into our ongoing MELODY Trial (9) including participants with 27^{th}–29^{th} weeks of gestation before any dietary intervention. Pregnant women with and without IBD were recruited nationwide under approved IRB protocol (IRB # H00016462) as previously described (10) (see supplementary methods). IBD disease activity was evaluated using validated scoring systems, namely the Harvey Bradshaw index (11) for participants with CD and the 6-point Mayo score (12) for participants with UC.

Sample collection: Vaginal and stool samples were self-collected using the OMNIgeneVAGINAL collection tube (DNA Genotek, Canada) or the ALPCO EasySampler kit (ALPCO, USA) following manufacturer instructions. Samples were received in the lab frozen 30h after sample collection.

Nucleic acid isolation: DNA from both vaginal and stool samples was isolated with Dneasy PowerSoil Pro kit (QIAGEN, Germany) and the RNA from vaginal samples was isolated with PowerMicrobiome kit (QIAGEN, Germany) following the manufacturer's protocol.

Cytokine expression: RNA from vaginal samples was subjected to qRT–PCRs (see supplementary methods). Oligonucleotides used to estimate cytokine expression are listed in **Table S1** (Integrated DNA Technology, USA).

Fecal calprotectin quantification was performed using the CalproLab ELISA ALP (Svar Life Sciences, Norway) according to the manufacturer's instructions. Total protein was quantified using the Pierce BCA Protein Assay kit (Thermo Fisher Scientific, USA). Calprotectin was normalized to initial stool weight (ng calprotectin/mg stool).

Vaginal microbiota sequencing and profiling was performed by *16S rRNA* sequencing of the V3-V4 hypervariable region as previously described (13). Sequencing libraries were sequenced on 600 cycles

using the MiSeq platform (Illumina, CA, USA). QIIME2 was used to process paired-end sequences. The DADA2 (14) algorithm was used for quality control and to obtain representative sequences (Amplicon Sequence Variant or ASV). We used a custom database (15, 16) for taxonomy classification. Only taxa with at least 0.10% abundance and present in a minimum of one sample were used for the analyses, as previously done (17, 18). **Table S2** describes the sequence counts included in the analyses.

Gut microbiota sequencing and profiling were performed using whole genome sequencing. Specifically, library generation and 150bp paired-end sequencing were carried out on the Illumina NextSeq 500 platform. KneadData (dec_v0.1, http://huttenhower.sph.harvard.edu/kneaddata) was used to eliminate human sequences and for quality control. MetaPhlAn4 database (vOct22)(19) was used for taxonomic assignment. **Table S2** describes the sequence counts included in the analyses.

Vaginal and gut microbiota diversity analyses were done in R, particularly the Phyloseq package (20). We performed data imputation for two individuals lacking BMI information (IBD=2), and four individuals (IBD=2, HC=2) without fecal calprotectin measurements using the median values for the IBD or HC group, respectively. All analyses were performed at the ASV level. Microbial alpha diversity was estimated using the Shannon and Simpson's Indexes (1-D) with rarefied sequencing data. Shannon and Simpson's Index were log-transformed to reach 'normality' of the residuals when necessary. To determine associations in alpha diversity we used linear regression models corrected by cofounders as age, body mass index (BMI), antibiotic use, and gestational diabetes. Results for the best-fitted model are reported.

We used Permutational Multivariate Analysis of Variance (PERMANOVA (21)) with adonis2 function to evaluate beta diversity (measured by Aitchison distance) of non-rarefied sequencing data.

We only assessed the associations between the vaginal microbiota and cytokines that were significantly different between IBD and HC. When assessing the association with dietary components, we only included

dietary components with no collinearity (Spearman correlation >|0.5|). One participant did not complete dietary recalls and thus was excluded from diet analyses.

Discriminant taxa analysis: Microbial taxa and their association with clinical variables were assessed using MaAsLin2's (22), also including the previously named confounding variables.

Community State Types (CST): Each vaginal sample was classified into CST as described before (23). To compare CST by discrete variables (health status) we use Fisher exact test (24); and by continuous variables (i.e., fecal calprotectin, cytokine expression, or diet score) Kruskal-Wallis test.

Dietary assessment: We conducted 24-hour dietary recalls (24HDRs) around the same time as vaginal/stool collection and estimated the Healthy Eating index-2015 (HEI-2015) as previously described by us (10) (see supplementary methods). Wilcoxon test was used to compare HEI-2015 and its dietary components by health status and CST.

Results

A total of 48 pregnant individuals were enrolled in the study: 23 with diagnosed IBD (n=18 CD, and n=5, UC) and 25 HC without IBD. Participants' demographics and clinical information are summarized in **Table 1**. Briefly, participants' mean age was 33.8 years, most had normal BMI (41.7%) or were overweight (37.5%), and most self-identified as White (93.8%). Only a few participants reported gestational diabetes or the use of antibiotics currently or previously during pregnancy. None of the clinical and demographic variables differed by health status (IBD vs. HC, **Table 1**) or IBD diagnosis (CD vs. UC, **Table S3**). More than 50% of the IBD participants were in self-reported remission (**Table 1**).

Vaginal microbiota diversity is associated with gut and vaginal inflammation, regardless of health status.

We first determined whether the vaginal microbiota diversity and composition differed by health status and found no significant differences (**Figure S1**). We also found no significant differences in the vaginal microbiota by IBD medications or when comparing individuals with IBD not on medication with the HC (**Figure S2**).

Given that most of the study participants were in remission or with mild disease we sought to further determine inflammatory markers that could influence the vaginal microbiota. We observed significantly higher levels of fecal calprotectin in IBD participants compared to their HC counterparts. Concomitantly, fecal calprotectin levels were positively associated with vaginal microbiota Simpson diversity but not with Shannon index (**Figure 1 and S1**) or for the beta diversity (**Figure S1**). Additionally, there were no significant associations between fecal calprotectin and vaginal microbiota diversity by health status (**Figure S1**).

We then assessed inflammation in the vaginal mucosae. We observed that pregnant individuals with IBD exhibited higher gene expression of Th17 pro-inflammatory cytokines, specifically IL-6, IL-8, and IL-17 than

HC. Conversely, expression of Th1 and Th2 cytokines IFN-γ and IL-4 respectively, was lower in pregnant

individuals with IBD compared to HC (**Figure 1**). We observe that the vaginal microbiota composition – not

its alpha diversity- correlated with the expression of IL-4 in the vaginal mucosa (**Figures 1 and S3**).

There were no specific bacteria associated with any of the inflammatory markers studied. We only identify

Dialister sp. and WAL 1855D (an uncultured bacterium from the *Tissierellaceae* family) increased in

pregnant individuals with high BMI (**Figure S4**).

Finally, we examined the gut microbiota of the pregnant individuals included in the study. There were no

significant differences in the gut microbiota diversity and composition by health status or fecal calprotectin

levels (**Figure S5**). However, we observed that *Collinsella* was lower in pregnant individuals with IBD

compared to HC (**Figure S5**).

Vaginal microbial diversity is associated with the consumption of vegetables and added sugars.

Pregnant individuals in this cohort had a HEI-2015 score of 63.8, comparable to the average of 63.0

reported by pregnant individuals in the USA (25, 26). There were no significant differences in the HEI-2015

score by health status (**Table S5**). Similarly, there were no significant differences in vaginal microbial

diversity or composition by HEI-2015 score (**Figure S6**).

We further investigated the associations between each dietary component included in the HEI-2015 score

and the vaginal microbiota diversity. Consumption of each dietary component was similar between

pregnant individuals with IBD and HC (**Table S5**). Whole fruit, fatty acid, and seafood/plant protein were

excluded from the analysis due to collinearity (Spearman score >|0.5|). We observed that high scores of

total vegetables (high vegetable intake) were predictive of a high vaginal microbiota diversity (**Figure 2**)

along with increasing abundance of *L. crispatus* (**Figure 3**). *L. iners* showed an opposite trend, although

not significant (**Figures 3 and S7**). We observed that vaginal microbial composition differed by added

sugar score, although no bacterial taxon was significantly associated with it (**Figure 2**).

Higher Lactobacillus crispatus *dominance (CST-I) is associated with higher dietary quality and*

vegetable consumption.

We determine the vaginal CST for each participant. More than 40% exhibited CST-I (*L. crispatus* dominance), followed by CST-III (*L. iners* dominance*)*, CST-II (*L. gasseri* dominance*)*, CST-IV (non-*Lactobacillus*-dominated), and CST-V (*L. jensenii* dominance. **Table S4**) (27). We did not observe differences in CSTs by health status, IBD diagnosis (CD vs. UC), levels of fecal calprotectin, or any of the vaginal cytokines assessed (**Table S3 and S4**).

Pregnant individuals with CST-I (*L. crispatus*-dominated, N=21) showed a higher HEI-2015 score, reflective of better dietary quality than CST-III individuals (*L. iners*-dominated, N=14. **Figure 3)**. Moreover, scoring for vegetable and added sugar intakes was significantly higher in individuals with CST-I than those with CST-III (**Figure 3**).

Discussion

Principal findings

The vaginal and gut microbiota composition of pregnant individuals with IBD was comparable to HC during the third trimester of pregnancy. Yet pregnant individuals with IBD presented gut inflammation linked to high vaginal microbial diversity and a vaginal pro-inflammatory immune tone. Pregnant individuals with higher dietary quality and optimal consumption of vegetables exhibited a vaginal microbiota profile dominated by the beneficial *L. crispatus*.

Results in the context of what is known.

Reduced levels of IFN-γ and IL-4 and elevated levels of IL-8 and IL-6, as seen in the pregnant individuals with IBD in this study, have been associated with an increased risk of preterm birth (28). To the best of our knowledge, our study is the first one to assess vaginal inflammatory markers in IBD pregnant patients and might explain the increased risk of preterm birth of pregnant individuals with IBD, even when in remission

or with mild disease (29, 30). Despite differences in the vaginal immune profile, the vaginal microbiota of

pregnant individuals with IBD was comparable to healthy controls. This result is unexpected as

inflammation relates to alteration in the microbiota in the vaginal mucosa (31, 32). Here, the vaginal

immune profile was assessed by gene expression (by RT-qPCR) and no protein/cytokine levels; thus,

further studies are required to accurately evaluate the immune tone in the vaginal mucosa.

We observe that gut inflammation in IBD individuals is positively associated with increased vaginal

microbial diversity. Of note, contrary to the gut microbiota, high microbial diversity in the vagina relates to

unhealthy states with increased risk for bacterial vaginosis (reviewed in (33)) and preterm birth (34, 35).

We then investigated whether diet impacts the vaginal microbiota. Pregnant individuals with higher dietary

quality exhibited a vaginal microbiota profile dominated by the beneficial *L. crispatus,* whereas the vaginal

microbiota profile of those with lower dietary quality was dominated by *L. iners*. For individual dietary

components, our findings showed that high vegetable consumption was associated with greater microbial

diversity (linked to vaginal dysbiosis (36)), similar to what previous studies have found for non-pregnant

vegetarians compared to non-vegetarians (37). However, our results also showed that, despite the higher

diversity, high vegetable intake resulted in a greater abundance of the beneficial *L. crispatus*. This indicates

the importance of not only considering a diversity index but identifying members of the vaginal microbiome

at the specie level to understand the potential implications of the microbiota in vaginal health.

Additionally, we found that lower added sugar intake resulted in decreased microbial alpha diversity and

increased *L. crispatus*. Concomitantly, high vegetable consumption and low intake of sweetened

beverages, has been also positively associated with abundance of *L. crispatus* in White and Black pregnant

women (8).

Importantly, *L. crispatus* is believed to offer the most protective benefits to the host compared to other

Lactobacillus species, with *L. iners* offering the least protective benefits (reviewed in (5)). *L. crispatus*

creates a highly acidic vaginal niche (pH<4.5) inhospitable to non-beneficial microbes, such as bacterial

vaginosis-related bacteria (38).

Clinical implications

We find that intestinal inflammation correlates with high vaginal microbiota diversity, indicative of unhealthy

states with increased risk for bacterial vaginosis and preterm birth. Thus, our results highlight the

importance of continuing therapy during pregnancy to reduce IBD-related intestinal inflammation.

Moreover, we found that diet can influence the dominance of a beneficial *L. crispatus* associated with

decreased risk of pre-term birth and bacterial vaginosis. Hence, emphasizing dietary quality during

pregnancy is a must, not only for the sustainment of pregnancy but to fuel a healthy vaginal microbiota.

Research implications

Our results highlight the need for a large prospective study that includes pregnant individuals with IBD

experiencing different disease activity (i.e., mild, moderate, severe). Future studies including dietary

interventions will unveil the role of diet as a strategy to support a healthy vaginal microbiome.

Strengths and limitations

Our study is constrained by a few key limitations. The modest sample size for both IBD and HC cohorts

curtails the statistical comparisons, particularly when several confounding variables, such as antibiotic use,

IBD medications, and gestational diabetes, are considered. This factor reduces the statistical power of the

study. Additionally, the IBD samples predominantly represent individuals in remission or with mild disease

with CD and only a few participants with UC, which narrows the scope of our conclusions to this specific

severity level of IBD and IBD diagnosis. Moreover, the ethnic/racial composition of our study sample, which

is mainly White, introduces a limitation since the vaginal microbiome is known to vary with race and

ethnicity (39). These limitations suggest the need for studies that include participants with severe disease,

an equal representation of IBD diagnosis (UC and CD), as well as individuals with diverse ethnic/racial

backgrounds.

Conclusions

Our results demonstrate that although the vaginal and gut microbiota of pregnant individuals with IBD and HC is similar in the third trimester of pregnancy, it varies depending on the immune tone of each mucosa. We show that pregnant individuals with IBD exhibit a pro-inflammatory cytokine profile that has been associated with an increased risk of pre-term birth. Finally, a high-quality diet with optimal intakes of vegetables and added sugars favors *L. crispatus* vaginal dominance.

Acknowledgments

We thank all the participants for their time and effort in collecting samples and responding to surveys. We also thank Rafael Lopez Martinez for contributing to designing scripts for analysis automatization. This work was supported by the Leona M. and Harry B. Helmsley Charitable Trust.

Data availability statement

The raw sequences and their associated metadata will be released at NCBI BioProject ID PRJNA915128 upon manuscript publication.

References

1. Hill JE, Pena-Sanchez JN, Fernando C, Freitas AC, Withana Gamage N, Fowler S. 2022. Composition and
Stability of the Vaginal Microbiota of Pregnant Women With Inflammatory Bowel Disease. Inflamm Bowel
Dis 28:905-911.

2. Rosta K, Mazzucato-Puchner A, Kiss H, Malik V, Mandl P, Petricevic L, Foessleitner P, Shafran I, Temsch
W, Farr A. 2021. Vaginal microbiota in pregnant women with inflammatory rheumatic and inflammatory
bowel disease: A matched case-control study. Mycoses 64:909-917.

3. Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Nikita L, Galuppi M, Lamont RF, Chaemsaithong
P, Miranda J. 2014. The composition and stability of the vaginal microbiota of normal pregnant women is
different from that of non-pregnant women. Microbiome 2:1-19.

4. Gudnadottir U, Debelius JW, Du J, Hugerth LW, Danielsson H, Schuppe-Koistinen I, Fransson E,
Brusselaers N. 2022. The vaginal microbiome and the risk of preterm birth: a systematic review and
network meta-analysis. Sci Rep 12:7926.

5. France M, Alizadeh M, Brown S, Ma B, Ravel J. 2022. Towards a deeper understanding of the vaginal
microbiota. Nat Microbiol 7:367-378.

6. Sun H, Yamada P, Paetow A, Chan M, Arslan A, Landberg R, Dominguez-Bello MG, Young BK. 2022. A
randomized controlled trial of the effects of whole grains versus refined grains diets on the microbiome in
pregnancy. Sci Rep 12:7509.

7. Jefferson KK, Parikh HI, Garcia EM, Edwards DJ, Serrano MG, Hewison M, Shary JR, Powell AM, Hollis
BW, Fettweis JM. 2019. Relationship between vitamin D status and the vaginal microbiome during
pregnancy. Journal of Perinatology 39:824-836.

8. Rosen EM, Martin CL, Siega-Riz AM, Dole N, Basta PV, Serrano M, Fettweis J, Wu M, Sun S, Thorp Jr JM.
2022. Is prenatal diet associated with the composition of the vaginal microbiome? Paediatric and Perinatal
Epidemiology 36:243-253.

9. Peter I, Maldonado-Contreras A, Eisele C, Frisard C, Simpson S, Nair N, Rendon A, Hawkins K, Cawley C,
Debebe A. 2020. A dietary intervention to improve the microbiome composition of pregnant women with
Crohn's disease and their offspring: the MELODY (Modulating Early Life Microbiome through Dietary
Intervention in Pregnancy) trial design. Contemporary clinical trials communications 18:100573.

10.   Olendzki BC, Hsiao BS, Weinstein K, Chen R, Frisard C, Madziar C, Picker M, Pauplis C, Maldonado-

353   Contreras A, Peter I. 2023. Dietary Intake of Pregnant Women with and without Inflammatory Bowel

354   Disease in the United States. Nutrients 15.

355   11.   Harvey RF, Bradshaw JM. 1980. A simple index of Crohn's-disease activity. Lancet 1:514.

356   12.   Lewis JD, Chuai S, Nessel L, Lichtenstein GR, Aberra FN, Ellenberg JH. 2008. Use of the noninvasive

357   components of the Mayo score to assess clinical response in ulcerative colitis. Inflamm Bowel Dis 14:1660-

358   6.

359   13.   Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a dual-index

360   sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina

361   sequencing platform. Appl Environ Microbiol 79:5112-20.

362   14.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016. DADA2: High-resolution

363   sample inference from Illumina amplicon data. Nat Methods 13:581-3.

364   15.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA, Gregory Caporaso J.

365   2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-

366   classifier plugin. Microbiome 6:1-17.

367   16.   Usyk M, Zolnik CP, Castle PE, Porras C, Herrero R, Gradissimo A, Gonzalez P, Safaeian M, Schiffman M,

368   Burk RD, Costa Rica HPVVTG. 2020. Cervicovaginal microbiome and natural history of HPV in a

369   longitudinal study. PLoS Pathog 16:e1008376.

370   17.   Reitmeier S, Hitch TC, Treichel N, Fikas N, Hausmann B, Ramer-Tait AE, Neuhaus K, Berry D, Haller D,

371   Lagkouvardos I. 2021. Handling of spurious sequences affects the outcome of high-throughput 16S rRNA

372   gene amplicon profiling. ISME Communications 1:1-12.

373   18.   Linz AM, Crary BC, Shade A, Owens S, Gilbert JA, Knight R, McMahon KD. 2017. Bacterial community

374   composition and dynamics spanning five years in freshwater bog lakes. MSphere 2:e00169-17.

375   19.   Blanco-Miguez A, Beghini F, Cumbo F, McIver LJ, Thompson KN, Zolfo M, Manghi P, Dubois L, Huang KD,

376   Thomas AM, Nickols WA, Piccinno G, Piperni E, Puncochar M, Valles-Colomer M, Tett A, Giordano F,

377   Davies R, Wolf J, Berry SE, Spector TD, Franzosa EA, Pasolli E, Asnicar F, Huttenhower C, Segata N.

378   2023. Extending and improving metagenomic taxonomic profiling with uncharacterized species using

379   MetaPhlAn 4. Nat Biotechnol doi:10.1038/s41587-023-01688-w.

380   20.   McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis and graphics
381   of microbiome census data. PloS one 8:e61217.
382   21.   Anderson MJ. 2014. Permutational multivariate analysis of variance (PERMANOVA). Wiley statsref:
383   statistics reference online:1-15.
384   22.   Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL, Weingart G, Ren B, Schwager
385   EH, Chatterjee S, Thompson KN, Wilkinson JE, Subramanian A, Lu Y, Waldron L, Paulson JN, Franzosa
386   EA, Bravo HC, Huttenhower C. 2021. Multivariable association discovery in population-scale meta-omics
387   studies. PLoS Comput Biol 17:e1009442.
388   23.   France MT, Ma B, Gajer P, Brown S, Humphrys MS, Holm JB, Waetjen LE, Brotman RM, Ravel J. 2020.
389   VALENCIA: a nearest centroid classification method for vaginal microbial communities based on
390   composition. Microbiome 8:166.
391   24.   Hervé M, Hervé MM. 2020. Package 'RVAideMemoire'. See https://CRANR-projectorg/package=
392   RVAideMemoire.
393   25.   Krebs-Smith SM, Pannucci TE, Subar AF, Kirkpatrick SI, Lerman JL, Tooze JA, Wilson MM, Reedy J.
394   2018. Update of the Healthy Eating Index: HEI-2015. J Acad Nutr Diet 118:1591-1602.
395   26.   U.S. Department of Agriculture FaNS, Center for Nutrition Policy and Promotion 2021. Average Healthy
396   Eating Index-2015 Scores for Non-Pregnant Non-Lactating, Pregnant and Lactating Women 20-44 Years.
397   What We Eat in America, NHANES 2013-2018.
398   27.   Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket
399   CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011. Vaginal microbiome of reproductive-age
400   women. Proc Natl Acad Sci U S A 108 Suppl 1:4680-7.
401   28.   Kumar M, Murugesan S, Singh P, Saadaoui M, Elhag DA, Terranegra A, Kabeer BSA, Marr AK, Kino T,
402   Brummaier T, McGready R, Nosten F, Chaussabel D, Al Khodor S. 2021. Vaginal Microbiota and Cytokine
403   Levels Predict Preterm Delivery in Asian Women. Front Cell Infect Microbiol 11:639665.
404   29.   Marild K, Soderling J, Stephansson O, Axelrad J, Halfvarson J, Group SS, Broms G, Marsal J, Olen O,
405   Ludvigsson JF. 2022. Histological remission in inflammatory bowel disease and risk of adverse pregnancy
406   outcomes: A nationwide study. EClinicalMedicine 53:101722.
407   30.   Baird DD, Narendranathan M, Sandler RS. 1990. Increased risk of preterm birth for women with
408   inflammatory bowel disease. Gastroenterology 99:987-94.

31. Masson L, Barnabas S, Deese J, Lennard K, Dabee S, Gamieldien H, Jaumdally SZ, Williamson AL, Little

F, Van Damme L, Ahmed K, Crucitti T, Abdellati S, Bekker LG, Gray G, Dietrich J, Jaspan H, Passmore JS.

2019. Inflammatory cytokine biomarkers of asymptomatic sexually transmitted infections and vaginal

dysbiosis: a multicentre validation study. Sex Transm Infect 95:5-12.

32. Campisciano G, Zanotta N, Licastro D, De Seta F, Comar M. 2018. In vivo microbiome and associated

immune markers: New insights into the pathogenesis of vaginal dysbiosis. Sci Rep 8:2307.

33. Chen X, Lu Y, Chen T, Li R. 2021. The female vaginal microbiome in health and bacterial vaginosis.

Frontiers in cellular and infection microbiology:271.

34. Hillier SL, Nugent RP, Eschenbach DA, Krohn MA, Gibbs RS, Martin DH, Cotch MF, Edelman R, Pastorek

JG, Rao AV. 1995. Association between bacterial vaginosis and preterm delivery of a low-birth-weight

infant. New England Journal of Medicine 333:1737-1742.

35. Donders G, Van Calsteren K, Bellen G, Reybrouck R, Van den Bosch T, Riphagen I, Van Lierde S. 2009.

Predictive value for preterm birth of abnormal vaginal flora, bacterial vaginosis and aerobic vaginitis during

the first trimester of pregnancy. BJOG: An International Journal of Obstetrics & Gynaecology 116:1315-

1324.

36. Srinivasan S, Hoffman NG, Morgan MT, Matsen FA, Fiedler TL, Hall RW, Ross FJ, McCoy CO, Bumgarner

R, Marrazzo JM, Fredricks DN. 2012. Bacterial communities in women with bacterial vaginosis: high

resolution phylogenetic analyses reveal relationships of microbiota to clinical criteria. PLoS One 7:e37818.

37. Song SD, Acharya KD, Zhu JE, Deveney CM, Walther-Antonio MRS, Tetel MJ, Chia N. 2020. Daily Vaginal

Microbiota Fluctuations Associated with Natural Hormonal Cycle, Contraceptives, Diet, and Exercise.

mSphere 5.

38. Ling Z, Kong J, Liu F, Zhu H, Chen X, Wang Y, Li L, Nelson KE, Xia Y, Xiang C. 2010. Molecular analysis

of the diversity of vaginal microbiota associated with bacterial vaginosis. BMC genomics 11:1-16.

39. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket

CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011. Vaginal microbiome of reproductive-age

women. Proc Natl Acad Sci U S A 108 Suppl 1:4680-7.

Figure 1. Variation in inflammatory markers and microbial diversity by health status. (A) Fecal calprotectin

levels of individuals with Inflammatory Bowel Disease (IBD) or Healthy Controls (HC; Wilcoxon test). (B)

439   Linear regression correlation between vaginal microbial alpha diversity and fecal calprotectin levels. (C)

440   Expression of cytokines on the vaginal mucosa of IBD and HC participants (Wilcoxon test. Asterisks denote

441   ** <5E-2, *** <5E-3, ****<5E-4). (D) Principal Coordinates Analysis (PCoA) of vaginal microbial beta

442   diversity by IL-4 levels (Aitchison distance and PERMANOVA).

457 **Figure 2**. HEI-2015 total vegetable and added sugar components are significantly associated with vaginal

458 microbiota diversity or composition, respectively. (A) Linear regression of microbial alpha diversity and

459 HEI-2015 total vegetable score. Gray shades in the graphs represent the 95% confidence interval. (B)

460 Principal Coordinates Analysis (PCoA) demonstrates significance for vaginal microbial composition by

461 HEI-2015 added sugar score (Aitchison distance and PERMANOVA).

Figure 3. HEI-20215 added sugar and total vegetable correlate with CST classification. Pregnant individuals with CST-I (*L. crispatus*- dominated profile) exhibit higher scores for (A) HEI-2015, (B) added sugar, and (C) total vegetables (Wilcoxon-test, adjusting for multiple comparisons). The arrows indicate the direction of consumption of each dietary component. (D) Total vegetable score positively correlates with the abundance of *L. crispatus* and negatively correlates with the abundance of *L. iners* (E, Spearman Correlation).

470 **Table 1**. Demographic and clinical variables for pregnant individuals with Inflammatory Bowel Disease

471 (IBD) or Healthy Controls (HC) recruited for the study between 2019 and 2022.

Demographics and clinical variables	IBD	HC	Overall	P value[&]
	(N=23)	(N=25)	(N=48)	
Age				*0.414*
Mean (SD)	33.3 (4.63)	34.4 (4.97)	33.8 (4.79)	
Median [Min, Max]	33.0 [22.0, 41.0]	36.0 [22.0, 42.0]	34.0 [22.0, 42.0]	
BMI categories [&&]				*0.179*
Underweight	1 (4.3%)	0 (0%)	1 (2.1%)	
Normal	12 (52.2%)	8 (32.0%)	20 (41.7%)	
Overweight	8 (34.8%)	10 (40.0%)	18 (37.5%)	
Obese	2 (8.7%)	7 (28.0%)	9 (18.8%)	
Race				*0.490*
White	23 (100%)	22 (88.0%)	45 (93.8%)	
Asian	0 (0%)	1 (4.0%)	1 (2.1%)	
Other	0 (0%)	2 (8.0%)	2 (4.2%)	
Gestational diabetes				*0.098*
Yes	3 (13.0%)	0 (0%)	3 (6.3%)	
No	16 (69.6%)	21 (84.0%)	37 (77.1%)	
Information unavailable	4 (17.4%)	4 (16.0%)	8 (16.7%)	
Use of antibiotic during all pregnancy				*0.468*
No	19 (82.6%)	22 (88.0%)	41 (85.4%)	
Yes	3 (13.0%)	3 (12.0%)	6 (12.5%)	
Information unavailable	1 (4.3%)	0 (0%)	1 (2.1%)	
IBD diagnosis				-
Crohn's disease	18 (78.3%)	0 (0%)	18 (37.5%)	
Ulcerative colitis	5 (21.7%)	0 (0%)	5 (10.4%)	
Healthy controls	0 (0%)	25 (100%)	25 (52.1%)	
IBD disease activity [&&&]				-
Mild disease	6 (26.1%)	0 (0%)	6 (12.5%)	
Remission	13 (56.5%)	0 (0%)	13 (27.1%)	
Information unavailable	4 (17.4%)	25 (100%)	29 (60.4%)	
Use of IBD medication				*1.70E-06*
No	9 (39.1%)	25 (100%)	34 (70.8%)	
Yes	14 (60.9%)	0 (0%)	14 (29.2%)	
Preterm				*0.106*
No	15 (65.2%)	17 (68.0%)	32 (66.7%)	
Yes	4 (17.4%)	0 (0%)	4 (8.3%)	
Information unavailable	4 (17.4%)	8 (32.0%)	12 (25.0%)	
Infant birth weight (g)				*0.151*
Mean (SD)	3150 (470)	3330 (662)	3240 (579)	
Median [Min, Max]	3230 [1810, 3710]	3290 [1080, 4520]	3230 [1080, 4520]	

[&]Fisher's exact test for categorical variables and Wilcoxon test for continuous variables.

[&&]BMIs categories correspond to the WHO's classifications: Underweight (<18.5), normal weight (18.5–24.9), overweight (≥25.0), and obese (≥30).

[&&&]Disease activity was estimated using the Harvey Bradshaw Index and the Mayo score for individuals with Crohn's Disease or Ulcerative colitis, respectively.

SUPPLEMENTARY MATERIAL

METHODS:

Recruitment: We conducted a case–control study nested into our ongoing MELODY Trial, which is a

prospective non-randomized diet intervention trial testing the effects of IBD anti-inflammatory diet during

the third trimester of pregnancy on maternal IBD activity and microbiome composition in mothers and their

babies(9). Written informed consent was obtained from all eligible participants. Eligible participants were

adult (18+ years of age) pregnant individuals with or without IBD, with less than 29 weeks of pregnancy,

carrying a singleton, with or without IBD diagnosis. We excluded pregnant individuals with scheduled C-

section or induction of vaginal delivery prior to week 37 at the time of enrollment, those with medical

conditions that required a special diet, or those unable to speak or understand English.

Cytokine expression: RNA from vaginal samples was reverse transcribed using iScript cDNA Synthesis

Kit (Bio-Rad, USA), and qRT–PCRs were performed using iTaq Universal SYBR Green-Supermix (Bio-

Rad, USA) in an Applied Biosystems ViiA7 Real-Time PCR machine (Thermo-Scientific, USA). Expression

of each cytokine was measured in triplicate and the mean was normalized by the expression of a

housekeeping gene (GAPDH). The $2^{(-\Delta\Delta Ct)}$ method was used to analyze the relative changes in gene

expression.

Dietary assessment: We conducted three, 24-hour dietary recalls (24HDRs) around the same time of

vaginal/stool collection. 24HDRs were performed using the USDA Automated Multiple Pass Method(40) in

conjunction with the University of Minnesota Nutrition Data for Research (NDSR) software (Current

version: NDS-R 2022)(41-44) as we have previously done(45-47). We estimated the Healthy Eating Index

2015 (HEI-2015) from the average of the three 24HDRs. The population ratio method(48) was used to

compute the individual scores of each of the 13 HEI-2015 components: total fruits, whole fruits, total

vegetables, greens and beans, dairy, total protein foods, seafood and plant proteins, fatty acids, refined

grains, whole grains, sodium, added sugars and saturated fats. Six food groups have values from 0 to 5; and seven from 0 to 10, for a total maximum score of 100 as previously described(25). Higher values represent better dietary quality. Wilcoxon test was used to compare HEI-2015 and individual HEI-2015' dietary components by health status and CST.

References for supplementary material section:

1. Hill JE, Pena-Sanchez JN, Fernando C, Freitas AC, Withana Gamage N, Fowler S. 2022. Composition and Stability of the Vaginal Microbiota of Pregnant Women With Inflammatory Bowel Disease. Inflamm Bowel Dis 28:905-911.

2. Rosta K, Mazzucato-Puchner A, Kiss H, Malik V, Mandl P, Petricevic L, Foessleitner P, Shafran I, Temsch W, Farr A. 2021. Vaginal microbiota in pregnant women with inflammatory rheumatic and inflammatory bowel disease: A matched case-control study. Mycoses 64:909-917.

3. Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Nikita L, Galuppi M, Lamont RF, Chaemsaithong P, Miranda J. 2014. The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. Microbiome 2:1-19.

4. Gudnadottir U, Debelius JW, Du J, Hugerth LW, Danielsson H, Schuppe-Koistinen I, Fransson E, Brusselaers N. 2022. The vaginal microbiome and the risk of preterm birth: a systematic review and network meta-analysis. Sci Rep 12:7926.

5. France M, Alizadeh M, Brown S, Ma B, Ravel J. 2022. Towards a deeper understanding of the vaginal microbiota. Nat Microbiol 7:367-378.

6. Sun H, Yamada P, Paetow A, Chan M, Arslan A, Landberg R, Dominguez-Bello MG, Young BK. 2022. A randomized controlled trial of the effects of whole grains versus refined grains diets on the microbiome in pregnancy. Sci Rep 12:7509.

7. Jefferson KK, Parikh HI, Garcia EM, Edwards DJ, Serrano MG, Hewison M, Shary JR, Powell AM, Hollis BW, Fettweis JM. 2019. Relationship between vitamin D status and the vaginal microbiome during pregnancy. Journal of Perinatology 39:824-836.

8. Rosen EM, Martin CL, Siega-Riz AM, Dole N, Basta PV, Serrano M, Fettweis J, Wu M, Sun S, Thorp Jr JM. 2022. Is prenatal diet associated with the composition of the vaginal microbiome? Paediatric and Perinatal Epidemiology 36:243-253.

530   9.   Peter I, Maldonado-Contreras A, Eisele C, Frisard C, Simpson S, Nair N, Rendon A, Hawkins K, Cawley C,

531   Debebe A. 2020. A dietary intervention to improve the microbiome composition of pregnant women with

532   Crohn's disease and their offspring: the MELODY (Modulating Early Life Microbiome through Dietary

533   Intervention in Pregnancy) trial design. Contemporary clinical trials communications 18:100573.

534   10.   Olendzki BC, Hsiao BS, Weinstein K, Chen R, Frisard C, Madziar C, Picker M, Pauplis C, Maldonado-

535   Contreras A, Peter I. 2023. Dietary Intake of Pregnant Women with and without Inflammatory Bowel

536   Disease in the United States. Nutrients 15.

537   11.   Harvey RF, Bradshaw JM. 1980. A simple index of Crohn's-disease activity. Lancet 1:514.

538   12.   Lewis JD, Chuai S, Nessel L, Lichtenstein GR, Aberra FN, Ellenberg JH. 2008. Use of the noninvasive

539   components of the Mayo score to assess clinical response in ulcerative colitis. Inflamm Bowel Dis 14:1660-

540   6.

541   13.   Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a dual-index

542   sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina

543   sequencing platform. Appl Environ Microbiol 79:5112-20.

544   14.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016. DADA2: High-resolution

545   sample inference from Illumina amplicon data. Nat Methods 13:581-3.

546   15.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA, Gregory Caporaso J.

547   2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-

548   classifier plugin. Microbiome 6:1-17.

549   16.   Usyk M, Zolnik CP, Castle PE, Porras C, Herrero R, Gradissimo A, Gonzalez P, Safaeian M, Schiffman M,

550   Burk RD, Costa Rica HPVVTG. 2020. Cervicovaginal microbiome and natural history of HPV in a

551   longitudinal study. PLoS Pathog 16:e1008376.

552   17.   Reitmeier S, Hitch TC, Treichel N, Fikas N, Hausmann B, Ramer-Tait AE, Neuhaus K, Berry D, Haller D,

553   Lagkouvardos I. 2021. Handling of spurious sequences affects the outcome of high-throughput 16S rRNA

554   gene amplicon profiling. ISME Communications 1:1-12.

555   18.   Linz AM, Crary BC, Shade A, Owens S, Gilbert JA, Knight R, McMahon KD. 2017. Bacterial community

556   composition and dynamics spanning five years in freshwater bog lakes. MSphere 2:e00169-17.

557   19.   Blanco-Miguez A, Beghini F, Cumbo F, McIver LJ, Thompson KN, Zolfo M, Manghi P, Dubois L, Huang KD,

558   Thomas AM, Nickols WA, Piccinno G, Piperni E, Puncochar M, Valles-Colomer M, Tett A, Giordano F,

Davies R, Wolf J, Berry SE, Spector TD, Franzosa EA, Pasolli E, Asnicar F, Huttenhower C, Segata N.

2023. Extending and improving metagenomic taxonomic profiling with uncharacterized species using

MetaPhlAn 4. Nat Biotechnol doi:10.1038/s41587-023-01688-w.

20.   McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis and graphics

of microbiome census data. PloS one 8:e61217.

21.   Anderson MJ. 2014. Permutational multivariate analysis of variance (PERMANOVA). Wiley statsref:

statistics reference online:1-15.

22.   Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL, Weingart G, Ren B, Schwager

EH, Chatterjee S, Thompson KN, Wilkinson JE, Subramanian A, Lu Y, Waldron L, Paulson JN, Franzosa

EA, Bravo HC, Huttenhower C. 2021. Multivariable association discovery in population-scale meta-omics

studies. PLoS Comput Biol 17:e1009442.

23.   France MT, Ma B, Gajer P, Brown S, Humphrys MS, Holm JB, Waetjen LE, Brotman RM, Ravel J. 2020.

VALENCIA: a nearest centroid classification method for vaginal microbial communities based on

composition. Microbiome 8:166.

24.   Hervé M, Hervé MM. 2020. Package 'RVAideMemoire'. See https://CRANR-projectorg/package=

RVAideMemoire.

25.   Krebs-Smith SM, Pannucci TE, Subar AF, Kirkpatrick SI, Lerman JL, Tooze JA, Wilson MM, Reedy J.

2018. Update of the Healthy Eating Index: HEI-2015. J Acad Nutr Diet 118:1591-1602.

26.   U.S. Department of Agriculture FaNS, Center for Nutrition Policy and Promotion 2021. Average Healthy

Eating Index-2015 Scores for Non-Pregnant Non-Lactating, Pregnant and Lactating Women 20-44 Years.

What We Eat in America, NHANES 2013-2018.

27.   Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket

CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011. Vaginal microbiome of reproductive-age

women. Proc Natl Acad Sci U S A 108 Suppl 1:4680-7.

28.   Kumar M, Murugesan S, Singh P, Saadaoui M, Elhag DA, Terranegra A, Kabeer BSA, Marr AK, Kino T,

Brummaier T, McGready R, Nosten F, Chaussabel D, Al Khodor S. 2021. Vaginal Microbiota and Cytokine

Levels Predict Preterm Delivery in Asian Women. Front Cell Infect Microbiol 11:639665.

29.   Marild K, Soderling J, Stephansson O, Axelrad J, Halfvarson J, Group SS, Broms G, Marsal J, Olen O,

Ludvigsson JF. 2022. Histological remission in inflammatory bowel disease and risk of adverse pregnancy

outcomes: A nationwide study. EClinicalMedicine 53:101722.

30.   Baird DD, Narendranathan M, Sandler RS. 1990. Increased risk of preterm birth for women with

inflammatory bowel disease. Gastroenterology 99:987-94.

31.   Masson L, Barnabas S, Deese J, Lennard K, Dabee S, Gamieldien H, Jaumdally SZ, Williamson AL, Little

F, Van Damme L, Ahmed K, Crucitti T, Abdellati S, Bekker LG, Gray G, Dietrich J, Jaspan H, Passmore JS.

2019. Inflammatory cytokine biomarkers of asymptomatic sexually transmitted infections and vaginal

dysbiosis: a multicentre validation study. Sex Transm Infect 95:5-12.

32.   Campisciano G, Zanotta N, Licastro D, De Seta F, Comar M. 2018. In vivo microbiome and associated

immune markers: New insights into the pathogenesis of vaginal dysbiosis. Sci Rep 8:2307.

33.   Chen X, Lu Y, Chen T, Li R. 2021. The female vaginal microbiome in health and bacterial vaginosis.

Frontiers in cellular and infection microbiology:271.

34.   Hillier SL, Nugent RP, Eschenbach DA, Krohn MA, Gibbs RS, Martin DH, Cotch MF, Edelman R, Pastorek

JG, Rao AV. 1995. Association between bacterial vaginosis and preterm delivery of a low-birth-weight

infant. New England Journal of Medicine 333:1737-1742.

35.   Donders G, Van Calsteren K, Bellen G, Reybrouck R, Van den Bosch T, Riphagen I, Van Lierde S. 2009.

Predictive value for preterm birth of abnormal vaginal flora, bacterial vaginosis and aerobic vaginitis during

the first trimester of pregnancy. BJOG: An International Journal of Obstetrics & Gynaecology 116:1315-

1324.

36.   Srinivasan S, Hoffman NG, Morgan MT, Matsen FA, Fiedler TL, Hall RW, Ross FJ, McCoy CO, Bumgarner

R, Marrazzo JM, Fredricks DN. 2012. Bacterial communities in women with bacterial vaginosis: high

resolution phylogenetic analyses reveal relationships of microbiota to clinical criteria. PLoS One 7:e37818.

37.   Song SD, Acharya KD, Zhu JE, Deveney CM, Walther-Antonio MRS, Tetel MJ, Chia N. 2020. Daily Vaginal

Microbiota Fluctuations Associated with Natural Hormonal Cycle, Contraceptives, Diet, and Exercise.

mSphere 5.

38.   Ling Z, Kong J, Liu F, Zhu H, Chen X, Wang Y, Li L, Nelson KE, Xia Y, Xiang C. 2010. Molecular analysis

of the diversity of vaginal microbiota associated with bacterial vaginosis. BMC genomics 11:1-16.

39. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket

CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011. Vaginal microbiome of reproductive-age

women. Proc Natl Acad Sci U S A 108 Suppl 1:4680-7.

40. Moshfegh AJ, Rhodes DG, Baer DJ, Murayi T, Clemens JC, Rumpler WV, Paul DR, Sebastian RS,

Kuczynski KJ, Ingwersen LA, Staples RC, Cleveland LE. 2008. The US Department of Agriculture

Automated Multiple-Pass Method reduces bias in the collection of energy intakes. Am J Clin Nutr 88:324-

32.

41. Casey PH, Goolsby SL, Lensing SY, Perloff BP, Bogle ML. 1999. The use of telephone interview

methodology to obtain 24-hour dietary recalls. J Am Diet Assoc 99:1406-11.

42. Bogle M, Stuff J, Davis L, Forrester I, Strickland E, Casey PH, Ryan D, Champagne C, McGee B, Mellad K,

Neal E, Zaghloul S, Yadrick K, Horton J. 2001. Validity of a telephone-administered 24-hour dietary recall in

telephone and non-telephone households in the rural Lower Mississippi Delta region. J Am Diet Assoc

101:216-22.

43. Gersovitz M, Madden JP, Smiciklas-Wright H. 1978. Validity of the 24-hr. dietary recall and seven-day

record for group comparisons. J Am Diet Assoc 73:48-55.

44. Schatzkin A, Kipnis V, Carroll RJ, Midthune D, Subar AF, Bingham S, Schoeller DA, Troiano RP, Freedman

LS. 2003. A comparison of a food frequency questionnaire with a 24-hour recall for use in an

epidemiological cohort study: results from the biomarker-based Observing Protein and Energy Nutrition

(OPEN) study. Int J Epidemiol 32:1054-62.

45. Maldonado-Contreras A, Noel SE, Ward DV, Velez M, Mangano KM. 2020. Associations between Diet, the

Gut Microbiome, and Short-Chain Fatty Acid Production among Older Caribbean Latino Adults. J Acad Nutr

Diet doi:10.1016/j.jand.2020.04.018.

46. Ockene IS, Tellez TL, Rosal MC, Reed GW, Mordes J, Merriam PA, Olendzki BC, Handelman G, Nicolosi

R, Ma Y. 2012. Outcomes of a Latino community-based intervention for the prevention of diabetes: the

Lawrence Latino Diabetes Prevention Project. Am J Public Health 102:336-42.

47. Zhao M, Chiriboga D, Olendzki B, Xie B, Li Y, McGonigal LJ, Maldonado-Contreras A, Ma Y. 2018.

Substantial Increase in Compliance with Saturated Fatty Acid Intake Recommendations after One Year

Following the American Heart Association Diet. Nutrients 10.

48.   Kirkpatrick SI, Reedy J, Krebs-Smith SM, Pannucci TE, Subar AF, Wilson MM, Lerman JL, Tooze JA.

2018. Applications of the Healthy Eating Index for Surveillance, Epidemiology, and Intervention Research:

Considerations and Caveats. J Acad Nutr Diet 118:1603-1621.

668 **<u>SUPPLEMENTARY TABLES</u>**

670 **Table S1.** List of oligos used to determine vaginal cytokine gene expression.

Serial No	Target gene (qRT primer)	Sequence (5' - 3')
1	IL-1β (F)	CAAGAGTGCTGGAGCGATAA
2	IL-1β (R)	CCGTTACAGCGGAGAGATATAAG
3	TNFα (F)	CACAGAAGACACTCAGGGAAAG
4	TNFα (R)	TCGAGGGAGTCACCCTTAAA
5	IL-6 (F)	GGAGGAGCCAAAGCTCAATAA
6	IL-6 (R)	ACTCCACACCACAAGAAGATG
7	IL-4 (F)	CCTGCTTCATGAGGGAAACT
8	IL-4 (R)	GGTGACAGAACAAGACCCTATC
9	IL-8 (F)	GAAAGGAAGTAGCTGGCAGAG
10	IL-8 (R)	GGGTGGAAAGGTTTGGAGTAT
11	IL-10 (F)	TGGAGTGAGTCCTGGAGAAATA
12	IL-10 (R)	CTCCATGTCCATCACACTTAGG
13	IFN-γ (F)	CCAGCCATCCTCAGAAATGT
14	IFN-γ (R)	CTTGCATCTCCTCACTCTAACC
15	IL-17 (F)	CCATAGTGAAGGCAGGAATCA
16	IL-17 (R)	GAGGTGGATCGGTTGTAGTAATC
17	GM-CSF (F)	CCTCCAGCAGGAATGTCTTAAT
18	GM-CSF (R)	TCTGACTCCTTGGGATGAAATG
19	GAPDH (F)	GTGCTCCCACTCCTGATTT
20	GAPDH (R)	CCTTCTCTAAGTCCCTCCTACA

680 **Table S2.** Overview number of sequences and amplicon sequence variants (ASVs) obtained from the

681 vaginal samples and stool samples.

Parameter	Vaginal samples			Stool samples	
	Raw sequences	Post-filtering	Post-filtering/rarefaction*	Raw sequences	Rarefied sequences**
N samples	48	48	48	45	44
Total number of seqs	2,751,605	2,742,919	336,000	652,433,458	236,170,044
Mean seqs/sample	57,325	57,144	7,000	14,498,521	5,367,501
Standard deviation of seqs/sample	24,010	23,945	-	7,468,723	-
Max number of seqs in a sample	123,825	123,720	7,000	30,294,951	5,367,501
Min number of seqs in a sample	12,916	12,872	7,000	715	5,367,501
Total number of different ASVs or Strains	1,425	573	563	1,420	897
Mean ASV or Strains/sample	48	36	33	115	118
Standard deviation of ASV or Strains/sample	38	23	23	46	43
Max number of ASVs or Strains in a sample	208	109	105	249	249
Min number of ASVs or Strains in a sample	6	6	5	1	57

683 * Sequences were rarefied at *7,000 sequences/sample or **5,367,501 sequences/sample, each

684 representing the highest number of sequences that included all vaginal and stool samples, respectively.

 Table S3. Demographic and clinical variables for pregnant individuals with Crohn's Disease (CD) or

 Ulcerative Colitis (UC) recruited for this study between 2019 and 2022.

Demographics and clinical variables	CD (N=18)	UC (N=5)	Overall (N=23)	P value[&]
Age				*0.822*
Mean (SD)	33.2 (4.86)	33.4 (4.22)	33.3 (4.63)	
Median [Min, Max]	33.5 [22.0, 41.0]	32.0 [28.0, 39.0]	33.0 [22.0, 41.0]	
BMI categories [&&]				*0.126*
Underweight	1 (5.6%)	0 (0%)	1 (4.3%)	
Normal	10 (55.6%)	2 (40.0%)	12 (52.2%)	
Overweight	6 (33.3%)	2 (40.0%)	8 (34.8%)	
Obese	1 (5.6%)	1 (20.0%)	2 (8.7%)	
Race				-
White	18 (100%)	5 (100%)	23 (100%)	
Asian	0 (0%)	0 (0%)	0 (0%)	
Other	0 (0%)	0 (0%)	0 (0%)	
Gestational diabetes				*0.053*
Yes	1 (5.6%)	2 (40.0%)	3 (13.0%)	
No	14 (77.8%)	2 (40.0%)	16 (69.6%)	
Information unavailable	3 (16.7%)	1 (20.0%)	4 (17.4%)	
Use of antibiotic during pregnancy				*1.000*
No	15 (83.3%)	4 (80.0%)	19 (82.6%)	
Yes	3 (16.7%)	0 (0%)	3 (13.0%)	
Information unavailable	0 (0%)	1 (20.0%)	1 (4.3%)	
Disease activity [&&&]				*0.456*
Mild disease	5 (27.8%)	1 (20.0%)	6 (26.1%)	
Remission	10 (55.6%)	3 (60.0%)	13 (56.5%)	
Information unavailable	3 (16.7%)	1 (20.0%)	4 (17.4%)	
Use of IBD medication				*0.342*
No	6 (33.3%)	3 (60.0%)	9 (39.1%)	
Yes	12 (66.7%)	2 (40.0%)	14 (60.9%)	
Preterm				*0.272*
No	12 (66.7%)	3 (60.0%)	15 (65.2%)	
Yes	2 (11.1%)	2 (40.0%)	4 (17.4%)	
Missing	4 (22.2%)	0 (0%)	4 (17.4%)	
Infant birth weight (g)				*0.411*
Mean (SD)	3100 (515)	3330 (169)	3150 (470)	
Median [Min, Max]	3230 [1810, 3710]	3230 [3180, 3540]	3230 [1810, 3710]	
Fecal calprotectin (ng/mg)				*1.000*
Mean (SD)	69.4 (65.9)	59.5 (42.7)	67.3 (60.9)	
Median [Min, Max]	45.2 [3.11, 208]	45.2 [6.89, 111]	45.2 [3.11, 208]	
Community State Types (CSTs)				*0.209*
I	7 (38.9%)	2 (40.0%)	9 (39.1%)	
II	4 (22.2%)	0 (0%)	4 (17.4%)	
III	2 (11.1%)	3 (60.0%)	5 (21.7%)	

IV-C	3 (16.7%)	0 (0%)	3 (13.0%)
V	2 (11.1%)	0 (0%)	2 (8.7%)

[&]Fisher's exact test for categorical variables and Wilcoxon test for continuous variables.

[&&]BMIs categories correspond to the WHO's classifications: Underweight (<18.5), normal weight (18.5–24.9), overweight (≥25.0), and obese (≥30).

[&&&]Disease activity was estimated using the Harvey Bradshaw Index and the Mayo score for individuals with Crohn's Disease or Ulcerative colitis, respectively.

713　**Table S4.** Vaginal microbial Community State Types (CSTs) distribution by health status, levels of fecal

714　calprotectin, and expression of vaginal cytokines

Variables	CST-I (N=21)	CST-II (N=7)	CST-III (N=14)	CST-IV-C (N=4)	CST-V (N=2)	Overall (N=48)	P value&
Health status							*0.369*
HC	12 (57.1%)	3 (42.9%)	9 (64.3%)	1 (25.0%)	0 (0%)	25 (52.1%)	
IBD	9 (42.9%)	4 (57.1%)	5 (35.7%)	3 (75.0%)	2 (100%)	23 (47.9%)	
Fecal calprotectin (ng/mg)							*0.373*
Mean (SD)	37.0 (55.8)	36.6 (47.2)	26.3 (26.5)	40.4 (47.8)	120 (124)	37.6 (51.1)	
Median	8.54	12.4	19.4	19.5	120	15.3	
IL-6							*0.114*
Mean (SD)	3.8 (4.1)	4.4 (3.4)	2.9 (2.3)	7.5 (3.9)	8.7 (0.05)	4.1 (3.7)	
Median	2.14	4.05	2.5	7.2	8.76	3.12	
IL-4							*0.474*
Mean (SD)	0.761 (0.658)	0.850 (0.762)	0.855 (0.684)	0.377 (0.301)	0.193 (0.140)	0.746 (0.651)	
Median	0.421	0.877	0.581	0.277	0.193	0.428	
IL-1							*0.086*
Mean (SD)	1.22 (0.453)	1.18 (0.403)	1.04 (0.392)	0.763 (0.519)	2.01 (0.321)	1.16 (0.469)	
Median	1.23	1.06	1.02	0.655	2.01	1.06	
TNF-alpha							*0.237*
Mean (SD)	3.46 (3.14)	4.42 (4.05)	3.00 (2.51)	2.48 (1.90)	11.2 (1.79)	3.71 (3.33)	
Median	1.32	4.74	1.69	1.93	11.2	2.1	
IFN-gamma							*0.431*
Mean (SD)	0.583 (0.671)	0.663 (0.951)	0.776 (0.751)	0.152 (0.303)	0.00340 (0.00157)	0.591 (0.715)	
Median	0.561	0.001	0.712	0.000763	0.0034	0.511	
IL-8							*0.201*
Mean (SD)	3.05 (3.05)	3.96 (4.26)	2.25 (2.12)	4.15 (4.24)	11.5 (2.62)	3.39 (3.48)	
Median	1.57	1.26	1.79	2.68	11.5	1.93	
GM-CSF							*0.076*
Mean (SD)	1.04 (0.453)	1.10 (0.561)	1.15 (0.488)	0.689 (0.565)	4.31 (0.303)	1.19 (0.815)	
Median	1.02	0.917	1.08	0.485	4.31	1.02	
IL-10							*0.176*
Mean (SD)	1.15 (0.625)	1.27 (0.745)	1.20 (0.496)	1.09 (0.500)	5.59 (0.473)	1.36 (1.06)	
Median	1	1.17	1.11	0.922	5.59	1.09	
IL-17							*0.096*
Mean (SD)	1.93 (1.20)	2.60 (1.31)	1.59 (1.08)	1.85 (1.26)	6.03 (1.09)	2.09 (1.44)	
Median	2.07	2.49	1.17	2.08	6.03	1.99	

 Table S5. HEI-2015 score and individual dietary components scoring distribution by health status.

HEI-2015 dietary components (scoring scale)	IBD (N=22)	HC (N=25)	Overall (N=47)	P value[&]
Healthy Eating Index (0-100)				*0.175*
Mean (SD)	66.6 (13.8)	61.4 (13.2)	63.8 (13.6)	
Median [Min, Max]	67.5 [45.0, 91.0]	64.0 [35.0, 84.0]	65.0 [35.0, 91.0]	
Total vegetable (0-5)				*0.361*
Mean (SD)	3.68 (1.39)	4.20 (1.12)	3.96 (1.27)	
Median [Min, Max]	4.00 [1.00, 5.00]	5.00 [1.00, 5.00]	5.00 [1.00, 5.00]	
Greens and beans (0-5)				*0.179*
Mean (SD)	4.41 (1.47)	3.48 (1.90)	3.91 (1.75)	
Median [Min, Max]	5.00 [0, 5.00]	5.00 [0, 5.00]	5.00 [0, 5.00]	
Total fruit (0-5)				*0.279*
Mean (SD)	3.23 (1.57)	2.80 (1.58)	3.00 (1.57)	
Median [Min, Max]	3.50 [0, 5.00]	3.00 [0, 5.00]	3.00 [0, 5.00]	
Whole fruit (0-5)				*0.281*
Mean (SD)	3.82 (1.76)	3.44 (1.87)	3.62 (1.81)	
Median [Min, Max]	5.00 [0, 5.00]	5.00 [0, 5.00]	5.00 [0, 5.00]	
Whole grains (0-10)				*0.728*
Mean (SD)	5.36 (3.27)	5.52 (3.27)	5.45 (3.24)	
Median [Min, Max]	5.50 [0, 10.0]	6.00 [0, 10.0]	6.00 [0, 10.0]	
Dairy (0-10)				*0.659*
Mean (SD)	6.68 (2.23)	6.76 (2.60)	6.72 (2.41)	
Median [Min, Max]	6.00 [3.00, 10.0]	7.00 [3.00, 10.0]	6.00 [3.00, 10.0]	
Total protein foods (0-5)				*0.508*
Mean (SD)	4.64 (0.727)	4.32 (1.03)	4.47 (0.905)	
Median [Min, Max]	5.00 [3.00, 5.00]	5.00 [1.00, 5.00]	5.00 [1.00, 5.00]	
Seafood and plant protein (0-5)				*0.212*
Mean (SD)	3.86 (1.83)	3.36 (1.78)	3.60 (1.80)	
Median [Min, Max]	5.00 [0, 5.00]	3.00 [0, 5.00]	5.00 [0, 5.00]	
Fatty acid ratio (0-10)				*0.162*
Mean (SD)	5.50 (3.60)	3.04 (2.94)	4.19 (3.46)	
Median [Min, Max]	5.00 [0, 10.0]	2.00 [0, 10.0]	3.00 [0, 10.0]	
Sodium (0-10)				*0.217*
Mean (SD)	5.84 (2.79)	5.82 (3.22)	5.83 (2.97)	
Median [Min, Max]	6.00 [1.00, 10.0]	6.50 [0, 10.0]	6.00 [0, 10.0]	
Refined grains (0-10)				*0.251*
Mean (SD)	6.09 (4.10)	6.76 (2.79)	6.45 (3.44)	
Median [Min, Max]	8.00 [0, 10.0]	7.00 [0, 10.0]	8.00 [0, 10.0]	
Added sugar (0-10)				*0.514*
Mean (SD)	8.18 (1.79)	8.20 (1.91)	8.19 (1.84)	
Median [Min, Max]	9.00 [5.00, 10.0]	9.00 [5.00, 10.0]	9.00 [5.00, 10.0]	
Saturated fats (0-10)				*0.847*
Mean (SD)	5.23 (3.32)	3.48 (3.00)	4.30 (3.24)	
Median [Min, Max]	5.50 [0, 10.0]	3.00 [0, 10.0]	4.00 [0, 10.0]	

& Fisher's exact test for categorical variables and Wilcoxon test for continuous variables

743 **Table S6.** HEI-2015 score and individual dietary components scoring distribution by Community State

744 Types (CSTs).

	Community State Types						
	I	II	III	IV-C	V	Overall	P
Diet variables	(N=21)	(N=6)	(N=14)	(N=4)	(N=2)	(N=47)	value&
Healthy Eating Index (total score 0-100)							*0.098*
Mean (SD)	**68.4 (11.3)**	61.0 (15.7)	**55.2 (13.7)**	69.0 (9.49)	73.5 (13.4)	63.8 (13.6)	
Median	**67**	65.5	**53**	65.5	73.5	65	
Total vegetables (0-5)							*0.007*
Mean (SD)	**4.57 (0.978)**	3.50 (1.38)	**3.21 (1.12)**	4.25 (1.50)	3.50 (2.12)	3.96 (1.27)	
Median	**5**	4	**3**	5	3.5	5	
Greens and Beans (0-5)							*0.169*
Mean (SD)	4.38 (1.36)	3.50 (2.07)	3.07 (2.20)	4.75 (0.500)	4.50 (0.707)	3.91 (1.75)	
Median	5	4.5	4	5	4.5	5	
Total fruit (0-5)							*0.016*
Mean (SD)	2.86 (1.31)	3.33 (1.86)	2.29 (1.59)	4.75 (0.500)	5.00 (0)	3.00 (1.57)	
Median	3	3.5	2	5	5	3	
Whole fruit (0-5)							*0.271*
Mean (SD)	3.81 (1.72)	3.83 (1.60)	2.79 (2.12)	4.50 (1.00)	5.00 (0)	3.62 (1.81)	
Median	5	4.5	3	5	5	5	
Whole grains (0-10)							*0.947*
Mean (SD)	5.52 (3.33)	5.33 (3.14)	5.14 (3.35)	5.50 (4.43)	7.00 (0)	5.45 (3.24)	
Median	6	5.5	4.5	6	7	6	
Dairy (0-10)							*0.919*
Mean (SD)	6.62 (2.52)	6.00 (2.45)	6.93 (2.53)	7.25 (1.71)	7.50 (3.54)	6.72 (2.41)	
Median	6	6	6.5	7.5	7.5	6	
Total protein foods (0-5)							*0.482*
Mean (SD)	4.33 (1.11)	5.00 (0)	4.50 (0.760)	4.50 (0.577)	4.00 (1.41)	4.47 (0.905)	
Median	5	5	5	4.5	4	5	
Seafood and plant protein (0-5)							*0.112*
Mean (SD)	4.24 (1.37)	3.67 (2.16)	2.79 (1.89)	2.50 (2.38)	4.50 (0.707)	3.60 (1.80)	
Median	5	5	2	2.5	4.5	5	
Fatty acid ratio (0-10)							*0.306*
Mean (SD)	4.95 (3.34)	2.67 (3.27)	3.36 (3.52)	5.25 (3.20)	4.50 (6.36)	4.19 (3.46)	
Median	4	1.5	2	4	4.5	3	
Sodium (0-10)							*0.305*
Mean (SD)	6.57 (2.60)	6.00 (3.35)	4.57 (3.27)	5.00 (2.94)	8.00 (1.41)	5.83 (2.97)	
Median	7	7	3.5	5.5	8	6	
Refined grains (0-10)							*0.627*
Mean (SD)	6.62 (3.80)	6.83 (2.86)	5.36 (3.50)	7.50 (2.52)	9.00 (1.41)	6.45 (3.44)	
Median	8	6.5	5	8	9	8	
Added sugar (0-10)							*0.02*
Mean (SD)	**9.10 (1.51)**	8.00 (1.79)	**7.07 (1.82)**	7.75 (2.06)	8.00 (1.41)	8.19 (1.84)	

Median	**10**	8.5	**6.5**	7.5	8	9	
Saturated fats (0-10)							*0.532*
Mean (SD)	5.10 (3.46)	2.67 (2.50)	3.79 (3.04)	4.75 (3.30)	3.50 (4.95)	4.30 (3.24)	
Median	5	2.5	3.5	6	3.5	4	

& Kruskal-Wallis test followed by Wilcoxon tests for significant variables; in bold are indicated the pairwise analyses yielding significant differences between groups.

SUPPLEMENTARY FIGURES

Figure S1. Comparison of vaginal microbiota diversity by health status and fecal calprotectin levels.

Shannon (A) and Simpson's (B) diversity indices to evaluate the microbial diversity. A linear regression

model assessed the comparison by health status. A Principal Coordinates Analysis (PCoA) (C) was

performed based on the Aitchison distance. The Shannon diversity index (D) by health status, with linear

773   regression modeling for analysis. Simpson's diversity index was employed separately for Healthy Controls

774   (E) and IBD patients (F), indicating no significant correlations; linear regression models were utilized for

775   these assessments. For the PCoA, PERMANOVA was applied to calculate P values and R^2, helping

776   quantify the explained variance.

796
797 **Figure S2**. Comparison of vaginal microbial diversity in IBD patients off medication vs Healthy Controls

798 (HC). Shannon's index (A) and Simpson's index (B). The significance levels were computed using a linear

799 regression model. Beta diversity patterns are depicted through Principal Coordinates Analysis (PCoA)

800 based on Aitchison distances (C), statistical significance and explained variance are reported through P

801 values and R^2, using PERMANOVA.

Figure S3. Vaginal microbial diversity and its of association with vaginal IL-4. Vaginal microbial diversity using Shannon (A) and Simpson's (B) diversity index by vaginal IL-4 relative fold change among all samples, reveals no significant associations, with linear regression models. Same analysis, although only among IBD patients using Shannon (C) and Simpson's (D) diversity index by vaginal IL-4 relative fold change, was not significant either. Principal Coordinates Analysis (PCoA), based on Aitchison distance, demonstrates no significance for vaginal microbial composition by IL-4 including only IBD patients (E). For PCoA PERMANOVA was used to obtain the P valued and to quantify the variance explained, as shown by P values and R^2.

S4. Vaginal species positively correlated with women's BMI. Number sequences by BMI for WAL 1855D

sp (A) and *Dialister* sp (B). Analysis was performed with MaAsLin2 algorithm. P values were adjusted for

multiple comparisons.

825
826

Figure S5. Comparison of fecal microbiota diversity by health status and fecal calprotectin levels. Shannon

(A) and Simpson's (B) diversity indices to evaluate the microbial diversity. A linear regression model

assessed the comparison by health status. A Principal Coordinates Analysis (PCoA) was performed based

on the Aitchison distance to compare health status (C). Taxa significantly different by health status based

on the MaAsLin2 algorithm (D). Shannon (E) and Simpson's (F) diversity indexes by fecal calprotectin

levels, with linear regression modeling for analysis. For the PCoA, PERMANOVA was applied to calculate

P values and R², helping quantify the explained variance

837   **Figure S6.** Diet quality is not associated with vaginal microbial alpha or beta diversity. HEI-2015 score for

838   alpha diversity measured with Shannon's (A) and Simpson's (B) diversity indexes. P values were

839   calculated with a linear regression model. Principal Coordinates Analysis (PCoA), based on Aitchison

840   distance, for vaginal microbial composition by HEI-2015 score (B). PERMANOVA was used to obtain the

841   P valued and to quantify the variance explained, as shown by P values and R2.

847 **Figure S7.** Microbial profile of pregnant individuals included in the study grouped by HEI-2015 total

848 vegetable score and sorted by *L. crispatus* relative abundance. Each bar represents a unique sample.