Angiotensin Converting Enzyme (ACE) expression in microglia reduces amyloid β deposition and neurodegeneration by increasing SYK signaling and endolysosomal trafficking.

Andrew R. Gomez[1], Hyae Ran Byun[1], Shaogen Wu[1], AKM Ghulam Mohammad[1], Jasmine Ikbariyeh[1], Jaelin Chen[1], Alek Muro[1], Lin Li[1], Ken E. Bernstein[1,5], Richard Ainsworth[2] and Warren G. Tourtellotte[1, 3, 4, 5,]*

Short Title: Modulation of microglial function by angiotensin converting enzyme (ACE)

Affiliations:
[1]Department of Pathology and Laboratory Medicine, Cedars-Sinai Medical Center, Los Angeles, CA 90048.
[2]Kao Autoimmunity Institute, Cedars-Sinai Medical Center, Los Angeles, CA 90048
[3]Department of Neurology, Cedars-Sinai Medical Center, Los Angeles, CA 90048
[4]Department of Neurosurgery, Cedars-Sinai Medical Center, Los Angeles, CA 90048
[5]Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, CA 90048

*Correspondence to:
Warren G. Tourtellotte, MD, PhD, FCAP
Department of Pathology and Laboratory Medicine
Cedars-Sinai Medical Center
8700 Beverly Blvd., Rm. 8719
Los Angeles, CA 90048
Tel: 310248-7736
warren.tourtellotte@cshs.org

ABSTRACT

Genome-wide association studies (GWAS) have identified many gene polymorphisms

associated with an increased risk of developing Late Onset Alzheimer's Disease (LOAD). Many

of these LOAD risk-associated alleles alter disease pathogenesis by influencing microglia innate

immune responses and lipid metabolism pathways. Angiotensin Converting Enzyme (ACE), a

GWAS LOAD risk-associated gene best known for its role in regulating systemic blood

pressure, also enhances innate immunity and lipid processing in peripheral myeloid cells, but a

role for ACE in modulating the function of myeloid-derived microglia remains unexplored. Using

novel mice engineered to express ACE in microglia and CNS associated macrophages (CAMs),

we find that ACE expression in microglia reduces Aβ plaque load, preserves vulnerable neurons

and excitatory synapses, and greatly reduces learning and memory abnormalities in the 5xFAD

amyloid mouse model of Alzheimer's Disease (AD). ACE-expressing microglia show enhanced

Aβ phagocytosis and endolysosomal trafficking, increased clustering around amyloid plaques,

and increased SYK tyrosine kinase activation downstream of the major Aβ receptors, TREM2

and CLEC7A. Single microglia sequencing and digital spatial profiling identifies downstream

SYK signaling modules that are differentially expressed by ACE expression in microglia that

mediate endolysosomal biogenesis and trafficking, mTOR and PI3K/AKT signaling, and

increased oxidative phosphorylation, while pharmacologic inhibition of SYK activity in ACE-

expressing microglia abrogates the potentiated Aβ engulfment and endolysosomal trafficking.

These findings establish a role for ACE in enhancing microglial immune function and they

identify potential utility for ACE-expressing microglia as a cell-based therapy to augment

endogenous microglial responses to Aβ in AD.

INTRODUCTION

Alzheimer's Disease (AD) is the leading cause of dementia and a progressive neurodegenerative disease which currently has no cure (1). AD is characterized by intraneuronal aggregates of paired helical filaments formed from hyper-phosphorylated microtubule associated protein tau (neurofibrillary tangles) and abnormally cleaved Amyloid Precursor Protein (APP) that aggregates into soluble amyloid-beta (Aβ) oligomers and insoluble macromers that form extracellular plaques in the brain (2). Together, these neuropathological changes lead to neurodegeneration and cognitive decline which is distinct from normal aging. Microglia, the myeloid cell-derived resident innate immune phagocytes of the central nervous system, have an important role in clearing protein aggregates and maintaining amyloid proteostasis in the brain (3, 4). While germline mutations in genes such as Presenilin and APP account for a relatively small number of early onset cases of familial AD (5), Genome-Wide Association Studies (GWAS) have provided accumulating evidence that sequence polymorphisms that influence the expression of many genes confer increased risk of the more common Late-Onset AD (LOAD) (6-8). Interestingly, many of these genes encode proteins that appear to modulate the ability of microglia to recognize, engulf, traffic and degrade protein aggregates such as Aβ (9). While a detailed understanding of the convergent signaling pathways within microglia that are altered by LOAD gene mutations is only beginning to emerge, they hold promise for discovering effective pharmacologic or cell-based therapies for AD.

Angiotensin Converting Enzyme (ACE) has been repeatedly implicated as a LOAD gene, but how it functions in AD pathogenesis is poorly understood (6, 7, 10-14). ACE is a dipeptidyl carboxypeptidase that is best known for its role in blood pressure regulation where it cleaves angiotensin I (AngI) into angiotensin II (AngII), a potent vasoconstrictor (15). Some evidence suggests that ACE may have a protective effect on the development of dementia considering that renin-angiotensin modifying anti-hypertensive drugs appear to decrease the overall rate of

dementia (16-18), but more specifically drugs that reduce blood pressure by blocking the AngII

receptor, AGTR1 rather than by blocking ACE catalytic activity, appear to be most effective at

decreasing the rate of dementia (19-21). A recent study of 6,601 patients of European ancestry

found that increased cortical ACE expression was protective against development of AD and the

effect was not associated with blood pressure regulation (22). ACE is a relatively promiscuous

peptidase which cleaves substrates not involved in blood pressure regulation (23), suggesting

that its protective role in AD may be due to non-blood pressure regulating functions such as in

myeloid cell development, myeloid immune function and cleavage of neurotoxic $A\beta_{1-42}$ protein to

less neurotoxic $A\beta_{1-40}$ protein in AD (24-26). In mice, a clinical dose of ACE inhibitor worsened

amyloid pathology in the hAPPSwInd amyloid AD model, as did induced ACE haploinsufficiency

(27). Interestingly, overexpression of ACE in myelomonocytes and granulocytes in mice leads

to a profound enhancement of their bactericidal, anti-tumor and anti-atherogenic activity (28-30).

Overexpression of ACE in myelomonocytes which included microglia in the APP/PS1 amyloid

mouse model, decreased amyloid plaque formation and improved learning and memory that

were dependent on ACE catalytic activity, but independent of blood pressure regulation (31).

Taken together, these data suggest that ACE may potentiate microglia function like it does in

peripheral myeloid-derived macrophages and neutrophils, and at least in part, provide some

insight regarding how ACE functions to modify AD pathogenesis as a LOAD-associated gene.

Here, we tested the hypothesis that ACE expression in microglia can modulate their ability to

process Aβ and to alter the course of neurodegeneration and cognitive decline in an amyloid

model of AD. We generated a novel mouse model which expresses ACE in microglia and CNS-

associated macrophages (CAMs) to interrogate whether it can alter microglia function and the

molecular, histopathologic and cognitive hallmarks of AD that have previously been well-

characterized in the 5xFAD amyloid mouse model of AD. We found that ACE-expressing

microglia have enhanced ability to migrate toward plaques and engulf neurotoxic $A\beta_{42}$, and to

reduce vulnerable excitatory synapse loss, neurodegeneration and cognitive abnormalities

associated with Aβ deposition in the brain over time. Interestingly, we found in vivo and in vitro

evidence that ACE expression in microglia increases SYK tyrosine kinase activity which is

known to mediate signaling downstream of the major microglia Aβ receptors, TREM2 and

CLEC7A, and to orchestrate the microglia immune response to Aβ (32-34). Taken together,

these results highlight the utility of modulating microglial function by enhancing ACE expression

within them to mitigate Aβ deposition and neurodegeneration in the brain.

RESULTS
In vivo microglia-specific ACE expression
ACE expression is not detectable in homeostatic microglia or in microglia activated around

amyloid plaques. However, in ACE$^{10/10}$ transgenic mice that express ACE in myelomonocytic

cells (28), there is also strong expression of ACE in microglia, suggesting that the previous

results showing decreased amyloid burden and behavioral abnormalities in APP/PS1 mice may

have also been a result of ACE function in microglia (35) (Fig. S1). To examine the function of

ACE more specifically in microglia, mice were engineered to express ACE by targeting a

construct expressing c-terminal flag-tagged human ACE (hACE-3xF) to the safe harbor Rosa

locus in the genome of mouse embryonic stem cells (36). LoxP sites flanking a transcriptional

stop sequence (LSL) downstream of the constitutively active mammalian CAG promoter was

used to regulate the expression of a self-cleaving 2A peptide for TurboGFP and hACE-3xF

proteins after loxP site recombination in the genome (37) (Fig. 1A). These novel Rosa$^{+/hACE-flox}$

mice were mated to Cx3Cr1$^{+/CreERT2}$ mice that express Tamoxifen (TMX)-dependent cre-

recombinase in myeloid cells (38) to generate RACE- (Rosa$^{+/+}$;Cx3Cr1$^{+/CreERT2}$) and RACE+

(Rosa$^{+/hACE-flox}$;Cx3Cr1$^{+/CreERT2}$) mice (Fig. 1B). To restrict hACE expression to yolk sac-derived

myeloid cells that engraft into the CNS during development (e.g., microglia and CNS-Associated

Macrophages – CAMs, that consist of a small population of dural macrophages, choroid plexus

macrophages, and perivascular macrophages) (39), RACE- and RACE+ mice received TMX

injections (125 mg/kg, IP) on four consecutive days between postnatal days 28 to 31 to activate

transgene expression in myeloid cells in RACE+ (hereafter referred to as R+), but not in RACE-

(hereafter referred to as R-) mice (Fig. 1C). After TMX treatment, peripheral myeloid cells with

loxP site recombination (transgene-ON) are replaced by turnover of non-loxP recombined

(transgene-OFF) cells from the bone marrow approximately 21 days after TMX treatment,

whereas myeloid-derived cells engrafted in the brain early during embryogenesis, such as

microglia, do not turn over in the brain and retain their transgene expression (38, 40). As

expected, 3 weeks after TMX injection, flow cytometry analysis of myeloid cells isolated from

brain, blood and spleen showed that the TurboGFP transgene was highly expressed in brain

myeloid cells from R+ mice, whereas transgene expression in peripheral blood and spleen

myeloid cells returned to undetectable levels (Fig. 1D and S2). In the brain, hACE protein

expression was restricted to >96% of microglia in R+ mice and it was undetectable in microglia

from R- mice (Fig. 1E).

ACE expression in microglia decreases Aβ$_{1-42}$ amyloid plaque burden in 5xFAD mice

RACE and 5xFAD mice were mated to examine whether ACE expression in microglia can

modulate their immune function and enhance their ability to reduce amyloid plaque burden in an

amyloid model of AD. 5xFAD transgenic mice express 5 human mutations of amyloid precursor

protein and presenilin 1, they accumulate Aβ$_{42}$ protein that aggregates into extracellular amyloid

plaques in the brain (41), and have neurodegeneration and cognitive abnormalities similar to AD

(41, 42). 5xFAD hemizygous mice from the original Tg6799 transgenic line were backcrossed

to C57BL6J (5xFAD$^{+/Tg6799}$, hereafter referred to as AD+) (41), mated to RACE mice and

injected with TMX to generate R-;AD- (Cx3Cr1$^{+/CreERT2}$;Rosa$^{+/+}$;5xFAD$^{+/+}$), R+;AD-

(Cx3Cr1$^{+/CreERT2}$;Rosa$^{+/ACE-flox}$;5xFAD$^{+/+}$), R-;AD+ (Cx3Cr1$^{+/CreERT2}$;Rosa$^{+/+}$;5xFAD$^{+/Tg6799}$), and

R+;AD+ (Cx3Cr1$^{+/CreERT2}$;Rosa$^{+/ACE-flox}$;5xFAD$^{+/Tg6799}$) mice. By 9 months of age, R-;AD+ mice

consistently accumulated a large number of Aβ$_{42}$ containing plaques throughout the brain (Fig. 2

A and A') and in R+;AD+ mice, plaque density was markedly decreased throughout the brain

(Fig. 2B and B') and where quantified, it was decreased by 3.5-fold in the hippocampal CA1

region (p < 0.0001), 2.3-fold in somatosensory cortex (p < 0.001) and 2.6-fold in the amygdala

(p< 0.001; Fig 2C).

Injury, pathogens and other insults to the brain lead to microglia activation that is

characterized in part by changes in their morphology and increased cytokine production (43).

Under homeostatic conditions, microglia have highly branched and elongated processes that

shorten and become less branched upon neuroinflammatory activation (44, 45). Appropriately

reflecting their activated state, microglia had significantly decreased branching and process

length in R-;AD+ mice compared to R-;AD- mice (p < 0.0001). Interestingly, ACE expression in

microglia significantly increased process branching and length in R+;AD- mice (p < 0.0001) and

it restored process branching and length to control (R-;AD-) levels in R+;AD+ mice (p < 0.05;

Fig. S3). Among nine pro-inflammatory cytokines examined, TNF-α (p < 0.0001), IL1-β

(p<0.0001) and CXCL1 (p < 0.05) were elevated in R-;AD+ compared to R-;AD- mice. ACE

expression in microglia did not significantly alter the expression of TNF-α or CXCL1, but it

modestly increased expression of IL1-β in R+;AD+ compared to R-;AD+ mice (p < 0.01; Fig.

S4). Similarly, ACE expression in microglia slightly decreased the ratio of neurotoxic $A\beta_{1-42}$ to

non-neurotoxic $A\beta_{1-40}$ protein production in the brain by 1.2-fold (p< 0.05; Fig. S5). Thus, ACE

expression in microglia significantly reduces plaque load in the brains of 5xFAD mice, and while

it alters microglia morphology and slightly decreases neurotoxic $A\beta_{1-42}$ production, it has very

little effect on the production of the proinflammatory cytokines that were examined.

ACE expression in microglia enhances their association with plaques and increases Aβ
phagocytosis and endolysosomal trafficking
ACE expression in microglia increased their accumulation around plaques, leading to 1.8-fold

more plaque-associated microglia (PAM) per plaque compared to R-;AD+ mice (p < 0.001; Fig.

3A). In addition, there was 2-fold more phagocytosis of $A\beta_{42}$ protein per PAM in R+;AD+

microglia compared to R-;AD- microglia (p < 0.0001; Fig. 3B), which was associated with an

increase in the volume of the CD68+ endolysosomal compartment. The endolysosomal

compartment volume was expanded by 5.4-fold in R-;AD+ compared to R-;AD- microglia (p <

0.0001), and it was enhanced another 1.7-fold with ACE expression in microglia (R+;AD+)

compared to R-;AD+ microglia (p < 0.01, Fig. 3C). These results suggest that decreased

amyloid plaque burden in the brain of R+;AD+ mice is at least in part due to increased

recruitment of microglia to Aβ-containing plaques and to their enhanced phagocytosis and

endolysosomal degradation of $A\beta_{42}$ within them.

ACE expression in microglia reduces neuron and synapse degeneration in 5xFAD mice
By 9-months of age, neuron and synapse degeneration that is correlated with abnormalities in

learning and memory behavior are present in 5xFAD mice (46). Consistent with published

reports, R-;AD+ mice had significant neuron loss in previously identified vulnerable regions of

the brain (47), with a 1.7-fold decrease in hippocampal subicular neuron density (p < 0.0001)

and a 1.3-fold decrease in layer V forebrain cortical neuron density (p < 0.01). Interestingly,

ACE expression in microglia significantly reduced neuron degeneration and restored neuron

density to control (R-;AD-) levels in the subiculum and in layer V cortical forebrain neurons (Fig

4A).

Excitatory synapse loss is correlated with neuron loss from excitatory glutaminergic neurons

in the forebrain and the loss is reduced by ACE expression in microglia. Excitatory synapses

marked by puncta labeled by the excitatory presynaptic specialization marker, vGlut1 and co-

localized with the postsynaptic specialization marker, PSD95 were reduced by 2.3-fold in R-

;AD+ compared to R-;AD- mice (p < 0.0001). ACE expression in microglia increased excitatory

synapse density by 1.9-fold (p < 0.0001) to near control (R-;AD-) levels (Fig. 4B). Synapses are

eliminated, at least in part, by mechanisms that involve labeling them with the complement

component C1q to "mark" them for microglia-dependent elimination (48). Consistent with the

loss of synapses in the forebrain of R-;AD+ mice, PSD95 labeled synapses were 3.8-fold more

frequently labeled with C1q in R-;AD+ compared to R-;AD- mice (p < 0.0001). Moreover, in

R+;AD+ mice, where excitatory synapses were increased compared to R-;AD+ mice, there was

a 2.5-fold decrease in the frequency of C1q labeling in synapses marked by PSD95 (p < 0.01;

Fig. S6A). Similarly, in R-;AD+ mice, the decreased density of PSD95 postsynaptic

specializations was accompanied by a 3.9-fold increase in phagocytosis of PSD95 by microglia

(p < 0.0001) which was decreased by 2.0-fold in microglia from R+;AD+ compared to R-;AD+

mice (p < 0.001; Fig. S6B). Taken together, these results indicate that ACE expression in

microglia reduces neuron and synapse degeneration in 5xFAD mice, at least partly through

decreasing PSD95 phagocytosis and decreasing C1q-mediated synapse turnover by microglia.

ACE expression in microglia restores learning and memory abnormalities in 5xFAD mice
5xFAD mice have learning and memory abnormalities that correlate with amyloid plaque

deposition and degeneration of neurons and synapses (49). Mice of all genotypes (R-;AD-,

R+;AD-; R-;AD+ and R+;AD+) had similar body weights (Fig. S7A), locomotor behavior (Fig.

S7B, C) and anxiety-like behavior in open field testing (Fig. S7D, E). However, in a

hippocampus-dependent Barnes maze memory test, 9 month old R-;AD+ mice showed

significant learning abnormalities with increased time to locate an object (latency) after 5 days of

training compared to R-;AD- mice (p < 0.0001). With ACE expression in microglia, R+;AD+

mice had significantly decreased learning latency compared to R-;AD+ mice over a 5 day

training period (p < 0.001; Fig. 5A). In a memory task performed two-days after training to

locate the object, R-AD+ mice had memory impairment with a 3.9-fold increase in latency

compared to R-;AD- mice (p < 0.0001). Interestingly, latency was significantly reduced by 1.6-

fold in R+;AD+ compared to R-;AD+ mice (p < 0.001; Fig. 5B). Similar results were obtained

when mice were trained to find a new object location (reversal memory). Latency to locate the

new object in R-;AD+ mice was significantly increased 3.5-fold compared to R-;AD- mice (p <

0.0001) and it was significantly decreased by 1.3-fold (p < 0.05) in R+;AD+ compared to R-;AD+

mice (Fig. 5C).

Freezing time in a fear conditioning assay is increased when an animal perceives an adverse

conditioned stimulus that is either environmentally contextual (hippocampal-dependent) or cued

(amygdala-dependent) when paired with an unconditioned shock stimulus. Time freezing was

significantly decreased during 5 training sessions in R-;AD+ mice compared to R-;AD- mice (p <

0.01), and it was similar between all other genotypes (Fig. 5D). However, in a contextual

memory test performed one day after training, R-;AD+ mice had a significant 1.9-fold reduction

in time freezing compared to R-;AD- mice (p < 0.0001). ACE expression in microglia in R+;AD+

mice significantly reduced freezing time by 1.8-fold in R+;AD+ compared to R-;AD+ mice (p <

0.001; Fig. 5E). Similarly, in a cued memory test performed 2 days after training, R-;AD+ mice

had a significant 2.4-fold decrease in time freezing (p < 0.0001) compared to R-;AD- mice,

which was significantly increased by 2.1-fold in R+;AD+ compared to R-;AD+ mice (p < 0.001).

Thus, in addition to reducing Aβ plaque load and rescuing neuron and synapse loss, ACE

expression in microglia significantly reduces learning and memory abnormalities in 5xFAD mice.

ACE expression in microglia enhances endolysosomal trafficking as well as Syk tyrosine
kinase and cell migration signaling
To characterize the molecular changes induced by ACE in microglia, single nuclei were

isolated from 6-month-old R-;AD-, R+;AD-, R-;AD+ and R+;AD+ frozen forebrains. Microglia

nuclei were enriched by fluorescence activated cell sorting (FACS) to diminish NeuN+ neuron,

Sox10+ oligodendroglia, and Sox9+ astrocyte nuclei in the samples (Fig. S8A) before single

nuclear capture and reverse transcription (10x Genomics) (Fig. S8B). snRNAseq and UMAP

analysis identified 10 transcriptionally distinct cell clusters (Fig. S9A) that were assigned to their

cellular identity by mapping well-established lineage associated gene expression to them (Fig.

S9B). Microglia were enriched approximately 3-5-fold and accounted for 10% of all nuclei

sequenced (Fig. S9A). Microglia were represented by two unique cell clusters that expressed

the microglia lineage-enriched markers, Csf1r, Cx3cr1, Tmem119, C1qa, Hexb and P2ry12 (Fig.

S9B and S10).

The two microglia clusters were pooled to perform a secondary UMAP analysis to identify 8

transcriptionally distinct microglia sub-clusters that were further demultiplexed to compare

genotype-dependent gene expression and cell distribution within them (Fig. 6A). Gene

expression analysis identified microglia sub-clusters with gene expression profiles similar to

previously reported homeostatic microglia (HM; sub-cluster 1), transitional microglia (TM; sub-

cluster 2) and disease-associated microglia (DAM) (sub-cluster 3, designated as DAM-A) (Fig.

S11) (50). In addition, a previously uncharacterized small DAM sub-cluster was identified (sub-

cluster 5) and designated as DAM-B (Fig. 6B). The frequency of microglia in DAM-A expanded

from 0.3% to 26.6% (χ^2 test, p < 10^{-34}) and in DAM-B, microglia expanded from 2.8% to 11.4%

(χ^2 test, p < 10^{-9}) in R-;AD+ compared to R-;AD- mice. With ACE expression in microglia, the

frequency of DAM-A was increased further by 4.6% and in DAM-B it was decreased by 4.9% in

R+;AD+ compared to R-;AD+ mice (Fig. 6A). Thus, ACE expression in microglia shifts

transcriptionally distinct DAMs by increasing DAM-A and decreasing DAM-B frequency.

The majority of DAM were represented by DAM-A which contained both TREM2-independent

DAM stage 1 (DAM-1) and TREM2-dependent DAM stage 2 (DAM-2) that were previously

described in 5xFAD mice (Fig. S11) (51). ACE expression in DAM-A significantly modulated the

expression of 165 genes, including the up-regulation of two late onset AD associated genes,

Apoe and Trem2 (8, 52), up-regulation of 15 additional previously described DAM genes (such

as, CD9, Cstd and Ctsl) and down-regulation of 3 previously described DAM genes (Colec12,

Dkk2 and Myo5a) (50) (Fig. 6C, Table S1). In addition, CD68 which was also upregulated at the

protein level in the expanded endolysosomal compartment of R+;AD+ microglia (Fig. 3C), was

also upregulated in the snRNAseq studies (Fig. 6C). To better understand how ACE expression

in microglia modulates their function to decrease Aβ plaque load, neuron/synapse degeneration

and learning and memory impairments, Ingenuity Pathway Analysis (IPA) was performed on the

genes differentially expressed between R+;AD+ and R-;AD+ DAM-A. Interestingly, pathways

related to endolysosomal trafficking and spleen tyrosine kinase (SYK) signaling were

significantly modulated by ACE expression in DAM-A (Fig. 6D, Table S2).

Increased endolysosomal trafficking was characterized by significantly altered gene

expression in pathways mediating CLEAR signaling, phagosome maturation and reactive

oxygen species (ROX) production (Fig. 6E, Table S3). CLEAR signaling has a major role in

lysosomal maturation and cellular clearance which recruits phagosome maturation mechanisms

and ROX production (53, 54). These molecular changes are consistent with increased

internalization of Aβ_{42} (Fig. 3B) and the expanded CD68+ endolysosomal compartment (Fig.

3C) that was observed *in vivo* in ACE-expressing PAM from R+;AD+ compared to R-;AD+ mice.

Endolysosomal trafficking mediated by the myeloid immune receptors, CLEC7A and TREM2

was also enhanced in ACE expressing microglia. CLEC7A (also known as DECTIN1) is a c-

type lectin receptor upregulated in microglia in 5xFAD mice and during neurodegeneration (50,

55, 56). Real-time fluorescence measurement in primary microglia isolated from R+;AD- and R-

;AD- mice treated with Zymosan conjugated to pHrodo, which binds and activates CLEC7A

receptors (Fig S12A), showed significantly increased phagocytosis and endolysosome loading

(Fig S12B), and degradation (Fig. S12C) in ACE-expressing microglia. Similarly, TREM2 is up-

regulated in DAM in 5xFAD mice (50, 57), up-regulated further by ACE expression in microglia

(Fig. 6C), and up-regulated in microglia from patients with AD (32). TREM2 binds Aβ_{42} to

activate microglial immune response and function, including protein clearance and cytokine

production (57). Real-time fluorescence measurement of microglia treated with oligomeric Aβ_{1-}

$_{42}$ (oAβ_{42}) conjugated to pHrodo (Fig. S12D) also showed significantly increased phagocytosis

and lysosome loading (Fig S12E), and increased degradation (Fig S12F) in microglia isolated

from R+;AD- compared to R-;AD- mice. Thus, ACE expression in microglia enhances

phagocytosis, endolysomal trafficking and degradation of extracellular Aβ through mechanisms

that involve TREM2 and CLEC7A receptor binding and downstream signaling (33).

Recent studies indicate that CLEC7A and TREM2 mediate microglia activation in response

to Aβ binding through phosphorylation of SYK tyrosine kinase which in turn activates

downstream signaling pathways such as mTOR and PI3K/AKT, and functional pathways that

regulate oxidative phosphorylation and cell migration (32, 34, 58). Interestingly, IPA identified

many downstream SYK signaling pathways that were highly upregulated in R+;AD+ compared

to R-;AD+ DAM-A, such as mTOR and PI3K/AKT signaling (Fig 6F, Table S4). Similarly, genes

involved in oxidative phosphorylation were generally upregulated, which is consistent with a

previously identified role for ACE in enhancing immune function at least in part through

increasing oxidative phosphorylation and ATP biogenesis in other myeloid cells such as

macrophages and neutrophils (59). Similarly, the IPA function for *Movement of Cells* was highly

modulated by ACE expression in microglia (Fig S13, Table S5), which likely correlates with

increased migration of microglia that accumulate around plaques in R+;AD+ relative to R-;AD+

mice *in vivo* (Fig. 3A).

To better understand the role of ACE-expression in DAM-A and their spatial relationship to

Aβ plaques in the brain, we generated tissue microarrays from the hippocampus of 9-month-old

male and female R-;AD-, R+;AD-, R-;AD+ and R+;AD+ mice, and examined them using whole

transcriptome analysis with the Nanostring GeoMx digital spatial profiling (DSP) platform.

Morphology markers for Iba1 to identify microglia, Aβ$_{42}$ to identify plaques and Syto13 to identify

cell nuclei were used to guide the selection of regions of interest (ROI) containing plaque-

associated microglia (PAM) and non-plaque-associated microglia (nPAM) for gene expression

analysis (Fig 7A). The Iba1 fluorescence channel was used to generate ultraviolet light masks

to elute probes bound to RNA within microglia in the ROI (Fig. 7B). In previous studies, DAM

appeared to correspond to PAM based on analysis of a small number of DAM proteins that were

upregulated primarily in PAM relative to nPAM (50). DSP analysis identified many genes that

were deregulated in PAM compared to nPAM (Fig. S14A, Table S6) and there was a highly

significant correlation between genes upregulated in PAM by DSP with those upregulated in

DAM-A by snRNAseq (p < 0.0001) (Fig. S14B). Thus, DAM-A appear to primarily represent

PAM, while DAM-B represent a small population of DAM that are not apparently associated with

plaques.

ACE expression in microglia led to significant differential gene expression in R+;AD+

compared to R-;AD+ PAM (Fig. 7C, Table S7), although the sensitivity to detect differentially

expressed genes was probe dependent and appeared to be decreased compared to snRNAseq

analysis performed on DAM-A (Fig. 6C). Nevertheless, Reactome Pathway Analysis (RPA)

highlighted the functional impact of ACE expression in PAM (Fig. 7D, Table S8) and it showed

similarity to pathways identified by IPA performed on gene expression identified by snRNAseq

on DAM-A. The pathways most significantly modulated by ACE in PAM were related to C-type

lectin receptor and CLEC7A receptor signaling (Fig. 7E, Table S9), pathways consistent with

SYK signaling downstream of mTOR, which included protein translation and rRNA synthesis

(Fig. 7F, Table S10), and pro-inflammatory signaling through NFκB and IL1-β (Fig. S15, Table

S11). Taken together, these results indicate that ACE expression in microglia modulates their

ability to process Aβ_{42} by driving signaling through TREM2 and CLEC7A receptors, and their

downstream tyrosine kinase effector SYK, to potentiate many recently identified functions of

SYK signaling in microglia, including increased plaque association, Aβ engulfment,

neuroprotection, and decreased Aβ plaque formation (32-34, 60).

Activated intracellular SYK signaling can be identified by its phosphorylation at Y352 (pSYK)

(61, 62). There was no detectable difference in the amount of pSYK in microglia mediated by

ACE expression between R+;AD+ and R-;AD+ nPAM (Fig. 8A). However, pSYK was increased

by 2.0-fold (p < 0.05) in PAM compared to nPAM in R-;AD+ mice (Fig. 8A), consistent with

previous evidence indicating that SYK signaling is increased in PAM in AD (32, 60).

Interestingly, in ACE expressing PAM from R+;AD+ mice, there was a further 2.4-fold increase

in the amount of pSYK within them compared to R-;AD+ PAM (p < 0.0001), consistent with the

pathway analyses showing molecular signatures related to increased SYK signaling in ACE

expressing microglia (Fig. 8A). To test whether increased Aβ_{42} uptake and endolysosomal

trafficking in ACE expressing microglia is mediated by increased SYK signaling, primary

microglia were isolated from R-;AD- and R+;AD- mice and cultured in vitro. Treatment of ACE

expressing microglia from R+;AD- mice with oligomeric Aβ_{42} (oAβ_{42}) showed a 1.7-fold increase

in uptake and trafficking into the CD68 endolysosomal compartment when compared to R-;AD-

microglia (p < 0.0001; Fig. 8B), consistent with a role for ACE in potentiating oAβ_{42} uptake and

trafficking that is thought to be mediated by SYK signaling (32-34). Accordingly, treatment with

the selective SYK inhibitor PRT062607 (P505) decreased $oA\beta_{42}$ uptake and endolysosomal

trafficking by 1.7-fold in the absence of ACE expression in R-;AD- microglia (p < 0.01), and it

decreased uptake and degradation of $oA\beta_{42}$ by 2.2-fold in ACE expressing microglia from

R+;AD- mice (p < 0.0001). Thus, inhibition of SYK signaling abrogated the potentiating effects

of ACE expression in R+;AD- microglia on $oA\beta_{42}$ uptake and endolysosomal trafficking to levels

similar to non-ACE expressing microglia (Fig. 8B). Taken together, these results indicate that

ACE expression in microglia potentiates $oA\beta_{42}$ uptake and degradation through mechanisms

that involve SYK signaling and the coordination of cellular processes mediating Aβ engulfment,

plaque association, decreased plaque formation and neuroprotection (32-34).

DISCUSSION
ACE expression in myeloid-derived macrophages and neutrophils potentiates their immunologic

function (28, 29, 63). In addition, ACE haploinsufficiency or administration of a CNS penetrant

ACE inhibitor exacerbates amyloid deposition and neurodegeneration in hAPPSw mice (27). By

contrast, $ACE^{10/10}$ mice that express ACE in peripheral myelomonocytes (and microglia - Fig.

S1) have reduced Aβ deposition, neurodegeneration and improved learning and memory

deficits when crossed to the APP/PS1 Aβ producing mice (31). Here, we found that selective

expression of ACE in microglia/CAMs potentiates their ability to process Aβ. We found that in

R+;AD+ mice there was decreased Aβ plaque burden, reduced neurodegeneration and

excitatory synapse loss, and reduced learning and memory impairments. Moreover, ACE-

expressing microglia in R+;AD+ mice were more frequently recruited to plaques, and they

engulfed more Aβ that was targeted to an expanded endolysosomal compartment. Accordingly,

pathway analysis from snRNAseq and DSP studies identified many ACE-regulated pathways in

DAM. For example, there was evidence of altered pro-inflammatory NFκB and IL-1β signaling

(Fig. S15), and significant alteration of the IPA *Movement of Cells Network* in DAM (Fig. S13),

which may drive their altered immune activation and increased aggregation around Aβ plaques.

Similarly, while in vitro and in vivo labeling of the endolysosomal compartment with CD68

showed increased endolysosomal trafficking of Aβ in ACE expressing microglia, functional

pathway analysis also showed upregulation of the *CLEAR Signaling* network, a master regulator

of lysosomal biogenesis and dynamics, and *Phagosome Maturation* and *Production of Reactive*

Oxygen Species. Specifically, expression of several pH-sensitive cathepsin lysosomal

proteases (cathepsin A, B, C, D, L, S and Z) were markedly upregulated in ACE-expressing

DAM, which may have contributed to their enhanced ability to degrade Aβ. This appeared to be

a cell-autonomous enhancement of microglial function as ACE-expressing microglia derived

from R+;AD- mice treated in vitro with Zymosan and $oA\beta_{42}$ conjugated to pHrodo showed clear

evidence of enhanced phagocytosis and Aβ degradation mediated by CLEC7A and TREM2

receptors.

The complement component C1q opsonizes damaged synaptic membranes, and in AD

models with Aβ deposition there is increased C1q-labeling of synapses that leads to widespread

synapse loss mediated by microglia (48). We found that while C1q subunits, C1qa, C1qb and

C1qc were upregulated in ACE-expressing DAM, PSD95+ post-synaptic membranes were less

frequently labeled by C1q in R+;AD+ brain (Fig. S6A). This correlated with less synapse loss

and decreased phagocytosis of PSD95-labeled synaptic membranes by microglia (Fig. S6B).

Together, these findings suggest that decreased Aβ plaque burden, as well as decreased

neuron degeneration and synapse loss in R+;AD+ mice contributed to the functional

improvement in learning and memory behavior.

GWAS studies have identified >40 LOAD risk-associated polymorphic gene loci, which

influences many genes expressed in microglia (64). While the impact of these polymorphisms

on microglia function is only partially understood, two of the most important risk-associated

variants APOE4 and TREM2[R47H] appear to impair microglial immunometabolic function. For

example, expression of APOE4 in microglia/CAMs reduces their accumulation around Aβ

plaques, reduces expression of complement and lysosomal pathways in microglia, accelerates

amyloid plaque burden and impairs synaptic plasticity (65). Similarly, TREM2 which is a high

affinity receptor for Aβ and APOE4 (57, 66), is essential for transducing DAM activation

response to Aβ (50). Microglia expressing the LOAD risk-associated variant TREM2[R47H] have

reduced affinity for their Aβ and APOE4 ligands, decreased Aβ plaque accumulation, and they

poorly acquire a DAM activation state (67, 68). These phenotypes are associated with altered

signaling from the TREM2/DAP12 complex that phosphorylates/activates the tyrosine kinase

SYK to engage downstream signaling pathways mediated in part by mTOR, PI3K/AKT and

mitochondrial oxidative phosphorylation (32, 34). Thus, emerging evidence points to SYK as an

intracellular signaling hub downstream of TREM2 that drives microglial immunometabolic fitness

and a neuroprotective phenotype in response to Aβ. Interestingly, we found clear evidence of

increased SYK phosphorylation in ACE-expressing microglia in R+;AD+ PAM in response to Aβ,

and that pharmacologic SYK inhibition abrogated the enhanced phagocytosis and

endolysosomal trafficking of Aβ within ACE-expressing (R+;AD-) microglia. ACE-enhanced

SYK signaling was also identified from snRNAseq pathway analysis that showed upregulation of

several signaling pathways downstream of SYK in DAM, including mTOR, PI3/AKT and

oxidative phosphorylation. Similarly, DSP pathway analysis in PAM from R+;AD+ mice

identified increased translation and rRNA synthesis, known to be downstream of mTOR

signaling, and increased c-lectin/CLEC7A receptor signaling, known to transduce Aβ binding

and drive SYK phosphorylation and signaling (32). Taken together, these results indicate that

ACE-expression in microglia increases SYK signaling and downstream pathway activation to

enhance their AD protective function, and accordingly, the phenotypic effects of ACE expression

in microglia are opposite to those observed in SYK-deficient microglia in 5xFAD mice (32, 34).

Microglia encircle Aβ plaques and they engage a DAM transcriptional activation state that

regulates their immune response to process Aβ (50). In these studies, we identified two DAM

populations in AD+ mice, DAM-A which appeared to represent classic DAM that consisted of

both DAM stage I and DAM stage II microglia (50), and a much smaller DAM-B population.

ACE expression in microglia eliminated nearly 50% of the DAM-B and increased the frequency

of DAM-A, and it induced many genes in DAM-A related to enhanced endolysosomal trafficking,

migration, inflammation, oxidative phosphorylation and SYK signaling. Microglia that engage

the DAM transcriptional state are thought to represent PAM (50), although single cell/nuclei

transcriptomic studies have not definitively resolved the spatial relationship of DAM with Aβ

plaques. By using snRNAseq combined with DSP, we found a significant correlation between

PAM gene signatures and those in DAM-A indicating they were associated with Aβ plaques,

while DAM-B were apparently not plaque-associated (Fig. S14). DAM-B appear to represent an

Nkain2, Pde4b, St18 high expressing population of microglia that were previously described

when APOE was ablated in neurons in a tauopathy mouse model (69). Future studies should

further characterize the functional relevance of DAM-B in 5xFAD mice.

ACE is normally expressed in human and mouse neurons (70), but not in microglia (Fig. S1),

raising an interesting question about whether ACE has any endogenous role in regulating

microglia function in the brain. ACE is a membrane bound protein with two catalytic domains

located outside of the cell membrane. Similar to cell surface APP, ACE is released from the cell

surface by cleavage-secretion to produce soluble, catalytically active ACE under normal

physiological conditions (71). While its function is very poorly understood in neurons,

expression on the surface of neurons and cleavage-secretion could provide a trans-signaling

mechanism to modulate microglia function. Hints that such a signaling mechanism may exist

come from studies identifying altered microglial function in ACE haploinsufficient mice where

neuronal ACE levels are decreased by 50% (27) and studies showing that ACE levels are

increased in the brain from AD patients and in neurons treated with Aβ$_{42}$ (70, 72). In the RACE

mice developed for these studies, where ACE is abundantly expressed on the cell surface of

microglia (Fig. 1E), it seems plausible that cleavage-secretion could enhance an ACE signaling

pathway through local cis-signaling in microglia which may have evolved to respond to ACE

cleavage products from neurons. Future studies should interrogate whether a neuron-microglia-

ACE signaling pathway exists and if so, to identify the ACE cleavage product(s) that modulate

neuron protective mechanisms in microglia. Presuming that future studies confirm a similar role

for ACE in human microglia, ACE expression in microglia may be useful to override

immunometabolic defects caused by single or multiple LOAD risk-associated genes. Given that

ACE-expressing microglia seem to complement signaling pathways that are impaired in

microglia that express the major AD risk-associated proteins, such as TREM2^{R47H} and APOE4, it

may be feasible to use ACE-expressing microglia in a cell-based therapy for AD. Promising

cell-based therapies are in development for many diseases, and our recent first-in-human

clinical trial demonstrated the safety profile of transplanted mesenchymal stem-cells expressing

glial derived neurotrophic factor (GDNF) for the treatment of ALS (73). Indeed, many pre-

clinical studies are underway to engraft human microglia into mice as proof-of-concept that they

may eventually be effective in human microglial transplantation therapy (74).

METHODS
Study Approval.

All animal procedures were reviewed and approved by the Institutional Animal Care and Use

Committee (IACUC) at Cedars Sinai Medical Center in Los Angeles California (IACUC approval:

IACUC007292).

Experimental Models.
Mice and genotyping:

Rosa-hACE-flox mice: Rosa-hACE-flox mice were generated using homologous recombination

in C57BL/6 embryonic stem cells (ESC; MUBES-01001, Cyagen, Santa Clara, CA) to generate

100% ESC-derived founder mice using a targeting construct containing a self-deleting

Neomycin phosphotransferase (sdNeo) positive-selection cassette (TurboKnockout, Cyagen,

Santa Clara, CA). To generate the gene targeting construct, homologous recombination

domains (HRDs) were generated using high fidelity Taq DNA polymerase (Q5 high-fidelity DNA

polymerase, NEB, Ipswich, MA) and polymerase chain reaction (PCR) with a BAC plasmid

containing the Gt(Rosa)26Sor gene sequence (RP23-401D9) as PCR DNA template. A 2647

bp 5' HRD and a 4107 bp 3' HRD flanking the XbaI site in intron 1 of the Rosa genomic locus

was generated and the sdNeo-CAG-LSL-TurboGFP-P2A-hACE-3xF construct was cloned in

reverse orientation into the Xba1 site flanked by the HRDs using isothermal assembly. The final

targeting construct was verified using restriction nuclease mapping and Sanger sequencing.

The targeting construct was linearized using AscI restriction endonuclease, electroporated into

MUBES ESCs and selected for 24 hours after transfection in G418 (200 mg/mL). 187 clones

were isolated and screened, and 27 candidate clones were identified using long range (LR)

PCR across the 3' HRD arm using primers PR1 and PR2. Additional PCR screening was

performed to verify the presence of the Neo selection cassette and the SV40 poly adenylation

sequence in the candidate clone ESC genome.

Southern blot analysis was performed on MfeI restricted genomic DNA from PCR confirmed

candidate clones to verify appropriate genome targeting using a probe upstream of the 5' HRD.

The double stranded DNA probe was generated by PCR using BAC plasmid DNA (RP23-

401D9) to amplify a ~500 bp amplicon using primers PR3 and PR4. The probe DNA was

labeled with Digoxigenin using DIG-High Prime DNA labeling kit and Southern blot detection

was performed using DIG Nucleic Acid Detection kit and accompanying protocols. The

restriction fragments hybridized to the probe were expected to generate a 12.6kb fragment on

the wild type allele and a 10.3kb fragment on the targeted allele in MfeI restricted genomic DNA

from heterozygous clones. Five ESC candidate clones identified by PCR (2A3, 2A4, 1B1, 1E2

and 2H3) were confirmed by Southern blotting analysis. Clone 2A3 was selected for blastocyst

injection to generate 3 male and 2 female germline Rosa[+/hACE-flox] mice. Founder mice were

mated to C57BL/6J mice and heterozygotes were maintained on an isogenic C57BL/6J genetic

background.

Rosa[+/hACE-flox] mice were genotyped using genomic DNA isolated from tail biopsy or ear punch

tissue with the following primers: OT2138 and OT2139 to generate a 247 bp Rosa wild type allele

amplicon and OT2138 and OT2141 to generate a 451 bp hACE-flox knockin allele amplicon.

Cx3Cr1-CreERT2 mice: Cx3Cr1$^{+/CreERT2}$ knockin mice been previously characterized to drive cre-

recombinase expression in myeloid cells and microglia.   Approximately 21 days after

intraperitoneal Tamoxifen injection myeloid cells are regenerated in their transgene-OFF state,

but in microglia which are permanently engrafted into the brain during embryogenesis, loxP site

recombination endures in a transgene-on state (40).   Heterozygous mice were obtained from The

Jackson Laboratory (Jax strain: #020940) and backcrossed to an isogenic C57BL/6J line.

Mice were genotyped using genomic DNA isolated from tail biopsy tissue with the following

primers: OT2011 and OT2012: to generate a 291 bp wild type allele amplicon and OT2011 and

OT2013 to generate a 455 bp CreERT2 knockin allele amplicon.

5xFAD mice: 5xFAD mice were previously generated to express five human familial amyloid

mutations and they rapidly produce extracellular amyloid-beta (Aβ) deposition in the brain.   5xFAD

mice exhibit cognitive and neurodegenerative changes, similar to Alzheimer's Disease (41).   The

original 5xFAD transgenic line, Tg6799 was obtained from The Jackson Laboratory

(5xFAD$^{+/Tg6799}$) and heterozygous mice were used in these studies.

5xFAD mice were genotyped by using the previously identified transgene integration site to

genotype heterozygous and homozygous mice by PCR (75).   Mice were genotyped using

genomic DNA isolated from tail biopsy or ear punch tissues with the following primers: OT2275

and OT2276 to generate a 402 bp wild type allele amplicon and OT2275 and OT2274 to generate

a 274 bp transgenic allele amplicon.

RACE mice: RACE mice were generated by mating Cx3Cr1-CreERT2 mice to Rosa-hACE-flox

mice.   Mice with genotype Cx3Cr1$^{+/CreERT2}$; Rosa$^{+/+}$ were designated RACE- and mice with

genotype Cx3Cr1$^{+/CreERT2}$; Rosa$^{+/hACE-flox}$ were designated RACE+.

All mouse lines were maintained on C57BL/6J isogenic background.   Specific PCR reaction

conditions for genotyping are available upon request.

SUPPLEMENTAL INFORMATION

Supplemental methods and extended data can be found online.
2
AUTHOR CONTRIBUTIONS
W.G.T. conceived the project and designed the studies. A.G., S.W., H.B., A.K.M., L.L and A.M.

performed experiments, W.G.T, A.G., S.W., H.B., A.K.M. and L.L. analyzed the data, and

W.G.T. and A.G. wrote the manuscript.

ACKNOWLEGEMENTS
Supported by NIH: RF1-AG074365. The authors thank Lin Karman for technical assistance.

CONFLICT OF INTEREST
The authors declare that no conflicts of interest exist.

REFERENCES
1. Association As. 2023: Alzheimer's Disease Facts and Figures. 2023.

2. Frisoni GB, Altomare D, Thal DR, Ribaldi F, van der Kant R, Ossenkoppele R, et al. The
probabilistic model of Alzheimer disease: the amyloid hypothesis revised. *Nature reviews*
Neuroscience. 2022;23(1):53-66.

3. Clayton KA, Van Enoo AA, and Ikezu T. Alzheimer's Disease: The Role of Microglia in Brain
Homeostasis and Proteopathy. *Front Neurosci.* 2017;11:680.

4. Xu Y, Jin M-Z, Yang Z-Y, and Jin W-L. Microglia in neurodegenerative diseases. *Neural*
Regeneration Research. 2021;16(2):270-80.

5. Chávez-Gutiérrez L, and Szaruga M. Mechanisms of neurodegeneration - Insights from
familial Alzheimer's disease. *Semin Cell Dev Biol.* 2020;105:75-85.

6. Wightman DP, Jansen IE, Savage JE, Shadrin AA, Bahrami S, Holland D, et al. A genome-
wide association study with 1,126,563 individuals identifies new risk loci for Alzheimer's
disease. *Nat Genet.* 2021;53(9):1276-82.

7. Bellenguez C, Küçükali F, Jansen IE, Kleineidam L, Moreno-Grau S, Amin N, et al. New
insights into the genetic etiology of Alzheimer's disease and related dementias. *Nature*
Genetics. 2022;54(4):412-36.

8. Kunkle BW, Grenier-Boley B, Sims R, Bis JC, Damotte V, Naj AC, et al. Genetic meta-
analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Abeta, tau,
immunity and lipid processing. *Nat Genet.* 2019;51(3):414-30.

9. Miao J, Ma H, Yang Y, Liao Y, Lin C, Zheng J, et al. Microglia in Alzheimer's disease:
pathogenesis, mechanisms, and therapeutic potentials. *Front Aging Neurosci.*
2023;15:1201982.

10.   Wingo AP, Liu Y, Gerasimov ES, Gockley J, Logsdon BA, Duong DM, et al. Integrating
human brain proteomes with genome-wide association data implicates new proteins in
Alzheimer's disease pathogenesis. *Nature Genetics.* 2021;53(2):143-6.

11.   Kehoe PG, Russ C, McIlory S, Williams H, Holmans P, Holmes C, et al. Variation in DCP1, encoding ACE, is associated with susceptibility to Alzheimer disease. *Nat Genet.* 1999;21(1):71-2.

12.   Lehmann DJ, Cortina-Borja M, Warden DR, Smith AD, Sleegers K, Prince JA, et al. Large meta-analysis establishes the ACE insertion-deletion polymorphism as a marker of Alzheimer's disease. *American journal of epidemiology.* 2005;162(4):305-17.

13.   Marioni RE, Harris SE, Zhang Q, McRae AF, Hagenaars SP, Hill WD, et al. GWAS on family history of Alzheimer's disease. *Transl Psychiatry.* 2018;8(1):99.

14.   Meng Y, Baldwin CT, Bowirrat A, Waraska K, Inzelberg R, Friedland RP, et al. Association of polymorphisms in the Angiotensin-converting enzyme gene with Alzheimer disease in an Israeli Arab community. *American journal of human genetics.* 2006;78(5):871-7.

15.   Esther CR, Marino EM, Howard TE, Machaud A, Corvol P, Capecchi MR, et al. The critical role of tissue angiotensin-converting enzyme as revealed by gene targeting in mice. *J Clin Invest.* 1997;99(10):2375-85.

16.   Murray MD, Hendrie HC, Lane KA, Zheng M, Ambuehl R, Li S, et al. Antihypertensive Medication and Dementia Risk in Older Adult African Americans with Hypertension: A Prospective Cohort Study. *Journal of General Internal Medicine.* 2018;33(4):455-62.

17.   Hughes D, Judge C, Murphy R, Loughlin E, Costello M, Whiteley W, et al. Association of Blood Pressure Lowering With Incident Dementia or Cognitive Impairment: A Systematic Review and Meta-analysis. *JAMA : the journal of the American Medical Association.* 2020;323(19):1934-44.

18.   Ding J, Davis-Plourde KL, Sedaghat S, Tully PJ, Wang W, Phillips C, et al. Antihypertensive medications and risk for incident dementia and Alzheimer's disease: a meta-analysis of individual participant data from prospective cohort studies. *The Lancet Neurology.* 2020;19(1):61-70.

19.   Li N-C, Lee A, Whitmer RA, Kivipelto M, Lawler E, Kazis LE, et al. Use of angiotensin receptor blockers and risk of dementia in a predominantly male population: prospective cohort analysis. *BMJ (Clinical research ed).* 2010;340:b5465.

20.   van Dalen JW, Marcum ZA, Gray SL, Barthold D, Moll van Charante EP, van Gool WA, et al. Association of Angiotensin II–Stimulating Antihypertensive Use and Dementia Risk. *Neurology.* 2021;96(1):e67-e80.

21.   Barthold D, Joyce G, Wharton W, Kehoe P, and Zissimopoulos J. The association of multiple anti-hypertensive medication classes with Alzheimer's disease incidence across sex, race, and ethnicity. *PLOS ONE.* 2018;13(11):e0206705.

22.   Ryan DK, Karhunen V, Su B, Traylor M, Richardson TG, Burgess S, et al. Genetic Evidence for Protective Effects of Angiotensin-Converting Enzyme Against Alzheimer Disease But Not Other Neurodegenerative Diseases in European Populations. *Neurology Genetics.* 2022;8(5):e200014.

23. Rao A, Bhat SA, Shibata T, Giani JF, Rader F, Bernstein KE, et al. Diverse biological functions of the renin-angiotensin system. *Med Res Rev.* 2024;44(2):587-605.

24. Sinka L, Biasch K, Khazaal I, Peault B, and Tavian M. Angiotensin-converting enzyme (CD143) specifies emerging lympho-hematopoietic progenitors in the human embryo. *Blood.* 2012;119(16):3712-23.

25. Bernstein KE, Khan Z, Giani JF, Cao DY, Bernstein EA, and Shen XZ. Angiotensin-converting enzyme in innate and adaptive immunity. *Nature reviews Nephrology.* 2018;14(5):325-36.

26. Zou K, Yamaguchi H, Akatsu H, Sakamoto T, Ko M, Mizoguchi K, et al. Angiotensin-Converting Enzyme Converts Amyloid-beta protein Ab 1-42 to Ab 1-40, and Its Inhibition Enhances Brain Ab Deposition. *Journal of Neuroscience.* 2007;27(32):8628-35.

27. Liu S, Ando F, Fujita Y, Liu J, Maeda T, Shen X, et al. A clinical dose of angiotensin-converting enzyme (ACE) inhibitor and heterozygous ACE deletion exacerbate Alzheimer's disease pathology in mice. *J Biol Chem.* 2019;294(25):9760-70.

28. Shen XZ, Li P, Weiss D, Fuchs S, Xiao HD, Adams JA, et al. Mice with enhanced macrophage angiotensin-converting enzyme are resistant to melanoma. *Am J Pathol.* 2007;170(6):2122-34.

29. Khan Z, Shen XZ, Bernstein EA, Giani JF, Eriguchi M, Zhao TV, et al. Angiotensin-converting enzyme enhances the oxidative response and bactericidal activity of neutrophils. *Blood.* 2017;130(3):328-39.

30. Okwan-Duodu D, Weiss D, Peng Z, Veiras LC, Cao DY, Saito S, et al. Overexpression of myeloid angiotensin-converting enzyme (ACE) reduces atherosclerosis. *Biochemical and biophysical research communications.* 2019;520(3):573-9.

31. Bernstein KE, Koronyo Y, Salumbides BC, Sheyn J, Pelissier L, Lopes DH, et al. Angiotensin-converting enzyme overexpression in myelomonocytes prevents Alzheimer's-like cognitive decline. *J Clin Invest.* 2014;124(3):1000-12.

32. Wang S, Sudan R, Peng V, Zhou Y, Du S, Yuede CM, et al. TREM2 drives microglia response to amyloid-β via SYK-dependent and -independent pathways. *Cell.* 2022;185(22):4153-69.e19.

33. Schafer DP, and Stillman JM. Microglia are SYK of Aβ and cell debris. *Cell.* 2022;185(22):4043-5.

34. Ennerfelt H, Frost EL, Shapiro DA, Holliday C, Zengeler KE, Voithofer G, et al. SYK coordinates neuroprotective microglial responses in neurodegenerative disease. *Cell.* 2022;185(22):4135-52.e22.

35. Bernstein KE, Gonzalez-Villalobos RA, Giani JF, Shah K, Bernstein E, Janjulia T, et al. Angiotensin-converting enzyme overexpression in myelocytes enhances the immune response. *Biol Chem.* 2014;395(10):1173-8.

36. Soriano P. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet.* 1999;21(1):70-1.

37. Srinivas S, Watanabe T, Lin CS, William CM, Tanabe Y, Jessell TM, et al. Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC developmental biology.* 2001;1(1):4.

38. Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, et al. Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity.* 2013;38(1):79-91.

39. McKinsey GL, Lizama CO, Keown-Lang AE, Niu A, Santander N, Larpthaveesarp A, et al. A new genetic strategy for targeting microglia in development and disease. *eLife.* 2020;9:e54590.

40. Parkhurst CN, Yang G, Ninan I, Savas JN, Yates JR, 3rd, Lafaille JJ, et al. Microglia promote learning-dependent synapse formation through brain-derived neurotrophic factor. *Cell.* 2013;155(7):1596-609.

41. Oakley H, Cole SL, Logan S, Maus E, Shao P, Craft J, et al. Intraneuronal beta-amyloid aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial Alzheimer's disease mutations: potential factors in amyloid plaque formation. *J Neurosci.* 2006;26(40):10129-40.

42. Ohno M. Failures to reconsolidate memory in a mouse model of Alzheimer's disease. *Neurobiol Learn Mem.* 2009;92(3):455-9.

43. Ferro A, Auguste YSS, and Cheadle L. Microglia, Cytokines, and Neural Activity: Unexpected Interactions in Brain Development and Function. *Front Immunol.* 2021;12:703527.

44. Stence N, Waite M, and Dailey ME. Dynamics of microglial activation: a confocal time-lapse analysis in hippocampal slices. *Glia.* 2001;33(3):256-66.

45. Navarro V, Sanchez-Mejias E, Jimenez S, Munoz-Castro C, Sanchez-Varo R, Davila JC, et al. Microglia in Alzheimer's Disease: Activated, Dysfunctional or Degenerative. *Front Aging Neurosci.* 2018;10:140.

46. Roy ER, Chiu G, Li S, Propson NE, Kanchi R, Wang B, et al. Concerted type I interferon signaling in microglia and neural cells promotes memory impairment associated with amyloid beta plaques. *Immunity.* 2022;55(5):879-94 e6.

47. Eimer WA, and Vassar R. Neuron loss in the 5XFAD mouse model of Alzheimer's disease correlates with intraneuronal Abeta42 accumulation and Caspase-3 activation. *Molecular neurodegeneration.* 2013;8:2.

48. Hong S, Beja-Glasser VF, Nfonoyim BM, Frouin A, Li S, Ramakrishnan S, et al. Complement and microglia mediate early synapse loss in Alzheimer mouse models. *Science.* 2016;352(6286):712-6.

49.   Roy ER, Wang B, Wan Y-W, Chiu GS, Cole AL, Yin Z, et al. Type I interferon response drives neuroinflammation and synapse loss in Alzheimer disease. *The Journal of Clinical Investigation.* 2020.

50.   Keren-Shaul H, Spinrad A, Weiner A, Matcovitch-Natan O, Dvir-Szternfeld R, Ulland TK, et al. A Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease. *Cell.* 2017;169(7):1276-90 e17.

51.   Huang Y, Happonen KE, Burrola PG, O'Connor C, Hah N, Huang L, et al. Microglia use TAM receptors to detect and engulf amyloid β plaques. *Nature immunology.* 2021;22(5):586-94.

52.   Chen X, and Holtzman DM. Emerging roles of innate and adaptive immunity in Alzheimer's disease. *Immunity.* 2022;55(12):2236-54.

53.   Palmieri M, Impey S, Kang H, di Ronza A, Pelz C, Sardiello M, et al. Characterization of the CLEAR network reveals an integrated control of cellular clearance pathways. *Human Molecular Genetics.* 2011;20(19):3852-66.

54.   Settembre C, and Medina DL. TFEB and the CLEAR network. *Methods Cell Biol.* 2015;126:45-62.

55.   Herre J, Marshall ASJ, Caron E, Edwards AD, Williams DL, Schweighoffer E, et al. Dectin-1 uses novel mechanisms for yeast phagocytosis in macrophages. *Blood.* 2004;104(13):4038-45.

56.   Krasemann S, Madore C, Cialic R, Baufeld C, Calcagno N, El Fatimy R, et al. The TREM2-APOE Pathway Drives the Transcriptional Phenotype of Dysfunctional Microglia in Neurodegenerative Diseases. *Immunity.* 2017;47(3):566-81.e9.

57.   Zhao Y, Wu X, Li X, Jiang LL, Gui X, Liu Y, et al. TREM2 Is a Receptor for beta-Amyloid that Mediates Microglial Function. *Neuron.* 2018;97(5):1023-31.e7.

58.   Torres-Hernandez A, Wang W, Nikiforov Y, Tejada K, Torres L, Kalabin A, et al. Targeting SYK signaling in myeloid cells protects against liver fibrosis and hepatocarcinogenesis. *Oncogene.* 2019;38(23):4512-26.

59.   Cao DY, Spivia WR, Veiras LC, Khan Z, Peng Z, Jones AE, et al. ACE overexpression in myeloid cells increases oxidative metabolism and cellular ATP. *J Biol Chem.* 2020;295(5):1369-84.

60.   Schweig JE, Yao H, Beaulieu-Abdelahad D, Ait-Ghezala G, Mouzon B, Crawford F, et al. Alzheimer's disease pathological lesions activate the spleen tyrosine kinase. *Acta Neuropathol Commun.* 2017;5(1):69.

61.   Makhoul S, Dorschel S, Gambaryan S, Walter U, and Jurk K. Feedback Regulation of Syk by Protein Kinase C in Human Platelets. *International journal of molecular sciences.* 2019;21(1).

62.   Tsang E, Giannetti AM, Shaw D, Dinh M, Tse JK, Gandhi S, et al. Molecular mechanism of the Syk activation switch. *J Biol Chem.* 2008;283(47):32650-9.

63. Okwan-Duodu D, Datta V, Shen XZ, Goodridge HS, Bernstein EA, Fuchs S, et al. Angiotensin-converting enzyme overexpression in mouse myelomonocytic cells augments resistance to Listeria and methicillin-resistant Staphylococcus aureus. *J Biol Chem.* 2010;285(50):39051-60.

64. Sudwarts A, and Thinakaran G. Alzheimer's genes in microglia: a risk worth investigating. *Molecular neurodegeneration.* 2023;18(1):90.

65. Liu CC, Wang N, Chen Y, Inoue Y, Shue F, Ren Y, et al. Cell-autonomous effects of APOE4 in restricting microglial response in brain homeostasis and Alzheimer's disease. *Nature immunology.* 2023;24(11):1854-66.

66. Atagi Y, Liu CC, Painter MM, Chen XF, Verbeeck C, Zheng H, et al. Apolipoprotein E Is a Ligand for Triggering Receptor Expressed on Myeloid Cells 2 (TREM2). *J Biol Chem.* 2015;290(43):26043-50.

67. Song WM, Joshita S, Zhou Y, Ulland TK, Gilfillan S, and Colonna M. Humanized TREM2 mice reveal microglia-intrinsic and -extrinsic effects of R47H polymorphism. *J Exp Med.* 2018;215(3):745-60.

68. Zhou Y, Song WM, Andhey PS, Swain A, Levy T, Miller KR, et al. Human and mouse single-nucleus transcriptomics reveal TREM2-dependent and TREM2-independent cellular responses in Alzheimer's disease. *Nature medicine.* 2020;26(1):131-42.

69. Koutsodendris N, Blumenfeld J, Agrawal A, Traglia M, Grone B, Zilberter M, et al. Neuronal APOE4 removal protects against tau-mediated gliosis, neurodegeneration and myelin deficits. *Nature Aging.* 2023;3(3):275-96.

70. Cuddy LK, Prokopenko D, Cunningham EP, Brimberry R, Song P, Kirchner R, et al. Abeta-accelerated neurodegeneration caused by Alzheimer's-associated ACE variant R1279Q is rescued by angiotensin system inhibition in mice. *Sci Transl Med.* 2020;12(563).

71. Ramchandran R, Sen GC, Misono K, and Sen I. Regulated cleavage-secretion of the membrane-bound angiotensin-converting enzyme. *J Biol Chem.* 1994;269(3):2125-30.

72. Miners S, Ashby E, Baig S, Harrison R, Tayler H, Speedy E, et al. Angiotensin-converting enzyme levels and activity in Alzheimer's disease: differences in brain and CSF ACE and association with ACE1 genotypes. *Am J Transl Res.* 2009;1(2):163-77.

73. Baloh RH, Johnson JP, Avalos P, Allred P, Svendsen S, Gowing G, et al. Transplantation of human neural progenitor cells secreting GDNF into the spinal cord of patients with ALS: a phase 1/2a trial. *Nature medicine.* 2022;28(9):1813-22.

74. Ifediora N, Canoll P, and Hargus G. Human stem cell transplantation models of Alzheimer's disease. *Front Aging Neurosci.* 2024;16:1354164.

75. Goodwin LO, Splinter E, Davis TL, Urban R, He H, Braun RE, et al. Large-scale discovery of mouse transgenic integration sites reveals frequent structural variation and insertional mutagenesis. *Genome research.* 2019;29(3):494-505.

1   **Figure Legends**
2
Fig. 1. Enforcing ACE expression in microglia in vivo. **(A)** Embryonic stem cells were

electroporated with a targeting construct designed to express TurboGFP and flag tagged human

ACE (hACE-3xF) in cells that express active cre-recombinase. G418 resistant clones were

screened by PCR across the 3' homology domain and by Southern blotting across the 5'

homology domain. Five properly targeted clones were identified and clone 2A3 was used to

generate 3 male and 2 female germline heterozygous ACE-flox (Rosa$^{+/hACE-flox}$) mice. **(B)** ACE-

flox mice were mated to Cx3Cr1$^{+/CreERT2}$ mice to generate RACE- (Cx3Cr1$^{+/CreERT2}$; Rosa$^{+/+}$) and

RACE+ (Cx3Cr1$^{+/CreERT2}$; Rosa$^{+/hACE-flox}$) mice. **(C)** RACE- and RACE+ mice were injected with

Tamoxifen (TMX; 125 mg/kg, IP) on four consecutive days from postnatal day 28-31 to express

the TurboGFP and hACE transgene in myeloid-derived cells. TMX injected RACE+ and RACE-

mice are referred to as R+ and R- mice, respectively. **(D)** After TMX injection, R+ mice had

detectable transgene expression in peripheral myeloid cells (spleen and blood) as well as in the

brain. By 21 days after TMX injection, transgene expression in the peripheral myeloid cells

(spleen and blood) became undetectable as the cells with somatic mutations were replaced by

cells from the bone marrow stem cell niche that retained the LSL (transgene-OFF) mutation. In

the brain, transgene expression remained on and was highly expressed in nearly all

Cx3Cr1/CD11b co-expressed myeloid cells. **(E)** R+ mice had high ACE expression in myeloid-

derived microglia that was not detectable in microglia from R- mice. (Scale bar = 10 µM).

21
Fig. 2. ACE expression in microglia reduced Aβ plaque burden in brains from 5xFAD mice. In

6-month-old mice Aβ$_{1-42}$ containing plaques were markedly decreased in **(B)** R+;AD+ compared

to **(A)** R-;AD+ brains. **(A', B')** Aβ$_{1-42}$ containing plaques (green) were surrounded by Iba1

expressing microglia (red). **(C)** Plaque density was reduced in CA1 of the hippocampus by 3.5-

fold ($p < 0.0001$), in somatosensory cortex by 2.3-fold ($p < 0.001$) and in the amygdala by 2.6-

fold ($p < 0.001$) in R+; AD+ compared to R-; AD+ brains (A, B, scale bar = 500 µM and A', B'

inset = 50 µM).

Fig. 3. Increased microglia migration, Aβ phagocytosis and endolysosomal trafficking in ACE

expressing microglia. **(A)** Plaques were identified by staining them with Amylo-Glo (blue) and

plaque associated microglia (PAM) labeled with an antibody to Iba1 were defined as Iba1+

microglia within 10 µM of a plaque edge. There were 1.8-fold more PAM surrounding plaques in

R+;AD+ compared to R-;AD+ mice ($p < 0.001$). **(B)** Immunofluorescence using antibodies to

Iba1 to identify microglia (red) and Aβ$_{1-42}$ to identify plaques (green) showed increased Aβ

protein phagocytosis by ACE expressing microglia. Individual PAM were masked in 3-

dimensions (3D) on 50 µM thick sections using iMaris image analysis software and the volume

of Aβ$_{1-42}$ within individual PAM was determined. Microglia from R+;AD+ mice had 2-fold more

internalized Aβ$_{1-42}$ compared to R-;AD+ microglia ($p < 0.0001$). **(C)** Immunofluorescence using

antibodies to Iba1 to detect microglia (red), CD68 to detect the cellular endolysomal

compartment (green), and Amylo-Glo to detect amyloid plaques (blue) showed increased

endolysosomal trafficking in R-;AD+ and R+;AD+ compared to R-;AD- PAM. Imaris filament

tracing was used to generate 3D masks representing individual PAM and then to quantify the

volume of CD68 protein within them. R-;AD+ PAM had 5.4-fold increase in CD68 labeling

compared to R-;AD- microglia, reflecting the increased endolysosomal trafficking within PAM

surrounding amyloid plaques ($p < 0.0001$). In R+;AD+ PAM, there was an additional 1.7-fold

increase in CD68 labeling compared to R-;AD+ PAM ($p < 0.01$), suggesting that ACE

expression in microglia enhances endolysosomal trafficking and protein degradation within

them. (Scale bar in A = 10 µM and B, C = 20 µM; square symbol = male and circular symbol =

female)

Fig. 4. ACE expression in microglia reduces neuron and synapse degeneration in 5xFAD mice.

(A) Neurons in the subiculum of the hippocampal formation (Sub) and layer V forebrain cortical

neurons (V) were labeled by immunofluorescence with an antibody to NeuN, and neuron density

in these regions was measured (representative images from R-;AD+ and R+;AD+ shown).

Neuron density was decreased by 1.7-fold in Sub ($p < 0.0001$) and 1.3-fold in V ($p < 0.01$) in R-

;AD+ compared to R-;AD- mice, consistent with neuron loss in these brain regions in 5xFAD

mice. Neuron density in R+;AD+ Sub and V were increased to levels found in R-;AD- mice. **(B)**

Synaptic density in the forebrain was measured using immunofluorescence to detect

presynaptic excitatory (vGlut1) and postsynaptic (PSD95) specializations. Excitatory synapses

with co-expression of vGlut1 and PSD95 (arrowheads) were decreased by 2.3-fold in R-;AD+

compared to R-;AD- mice ($p < 0.0001$) and they were increased by 1.9-fold ($p < 0.0001$) in

R+;AD+ compared to R-;AD- mice. (A, scale bar = 1 µM, B, scale bar = 200 µM; square symbol

= male and round symbol = female)

Fig. 5. ACE expression in microglia significantly ameliorates learning and memory

abnormalities in 5xFAD mice. Barnes maze and fear conditioning were used to test

hippocampal-dependent and amygdala-dependent learning and memory behavior. **(A)** Training

for five days to locate an object in a Barnes maze showed significant increase in time to locate

the object (latency) in R-;AD+ compared to R-;AD- mice ($p < 0.0001$). However, latency to learn

the location of the object was significantly reduced in R+;AD+ compared to R-;AD+ mice ($p <$

0.001). **(B)** In a memory test, latency to locate the object was significantly increased in R-;AD+

by 3.9-fold compared to R-;AD- mice two days after training ($p < 0.0001$) and latency to locate

the object was significantly decreased in R+;AD+ by 1.6-fold compared to R-;AD+ mice ($p <$

0.001). **(C)** In a reversal memory test, by moving the object to a new location and re-training, R-

;AD+ mice again had a significantly increased latency to locate the object by 3.5-fold compared

to R-;AD- mice ($p < 0.0001$) which was significantly decreased by 1.3-fold in R+;AD+ compared

to R-;AD+ mice ($p < 0.05$). **(D)** In fear conditioning behavioral tests, R-;AD+ mice had

significantly impaired learning of contextual and cued stimuli during training compared to R-;AD-

mice (p < 0.01). **(E)** In contextual memory testing one-day after training, R-;AD+ mice had

significant memory impairment with a 1.9-fold decrease in freezing time compared to R-;AD-

mice (p < 0.0001). Memory was significantly improved with freezing time increased by 1.8-fold

in R+;AD+ compared to R-;AD+ mice (p < 0.001). **(F)** In cued memory testing two-days after

training, R-;AD+ mice had significant memory impairment with a 2.4-fold decrease in freezing

time compared to R-;AD- mice (p < 0.0001). Memory was significantly improved with a 2.1-fold

increase in freezing time in R+;AD+ compared to R-;AD+ mice (p < 0.001).

8
Fig. 6. ACE expression in disease associated microglia (DAM) modulates endolysosomal

trafficking and SYK signaling. **(A)** snRNAseq and UMAP subcluster analysis of microglia from

the forebrain of 9-month-old R-;AD-, R+;AD-, R-;AD+ and R+;AD+ mice identified 8

transcriptionally distinct microglia subclusters. Several microglia subclusters had gene

expression profiles consistent with previously described homeostatic (HM; subcluster 1),

transitional (TM; subcluster 2) and disease associated (DAM-A; subcluster 3) microglia. The

DAM-A subcluster contained previously identified TREM2-independent DAM stage 1 (DAM-1)

and TREM2-dependent DAM stage 2 (DAM-2) microglia (50). A smaller and poorly defined

DAM subcluster (subcluster 5) was identified and designated as DAM-B. The frequency of

DAM-A and DAM-B were highly altered by genotype. The frequency of DAM-A was increased

from 0.3% to 26.6% and in DAM-B they were increased from 2.8% to 11.4% in R-;AD+

compared to R-;AD- mice. With ACE expression in microglia the frequency of DAM-A was

increased by 4.6% (to 31.2%) and in DAM-B they were decreased by 4.9% (to 6.5%). **(B)** DAM-

A and DAM-B microglia were only observed in the presence of amyloid in AD+ brains, defining

them as DAM, and their frequency was skewed by ACE expression in them. **(C)** Ingenuity

Pathway Analysis (IPA) identified many highly significantly altered signaling pathways in DAM-A

R+;AD+ compared to R-;AD+ microglia. Although human ACE (hACE) was elevated in R+

microglia, it was not detected in the mouse sequencing data that used the mouse reference

genome for sequence mapping. **(D)** Endolysosomal trafficking involving CLEAR signaling,

phagosome maturation and production of ROX as well as SYK signaling involving mTOR,

PI3/AKT signaling, and oxidative phosphorylation were upregulated by ACE expression in

microglia. **(E, F)** IPA analysis identified pathways associated with endolysosomal trafficking and

SYK signaling that where upregulated by ACE expression in DAM-A (R+;AD+).

6
Fig. 7. ACE modulates CLEC7A, mTOR and inflammatory signaling in microglia. **(A)** Regions

of interest (ROI) containing plaque associated microglia (PAM) and non-plaque associated

microglia (nPAM) were selected using immunofluorescence morphology markers to localize

microglia (Iba1) and plaques ($A\beta_{1-42}$), and the nuclear stain SYTO Greeen. **(B)** Nanostring digital

spatial profiling (GeoMx) and whole transcriptome analysis were used on regions of interest

(ROI) selected using the morphology markers, and the Iba1 channel was used to generate a UV

mask to photo-elute probes hybridized primarily to microglia. **(C)** ACE expression in PAM

significantly modulated gene expression within them. Although human ACE (hACE) was highly

expressed in microglia, it was not detected by the mouse-specific probes in GeoMx. **(D)**

Reactome pathway analysis showed that many of the genes regulated by ACE in PAM

modulate signaling pathways related to mTOR, CLEC7A, and inflammatory signaling. **(E)** ACE

expression in PAM modulates many genes related to the c-lectin receptor CLEC7A signaling in

microglia and **(F)** genes related to signaling pathways downstream of mTOR. (scale bar = 100

µM)

21
Fig. 8: ACE expression in microglia enhances SYK phosphorylation and SYK-dependent $A\beta$

phagocytosis and endolysosomal trafficking. **(A)** SYK phosphorylation was increased 2-fold in

PAM relative to nPAM ($p < 0.05$). With ACE expression in microglia, levels of p-SYK within

PAM increased by 2.4-fold ($p < 0.0001$), consistent with increased SYK signaling relative to

microglia not expressing ACE. **(B)** Treatment of primary microglia with $oA\beta_{1-42}$ resulted in a 1.7-

fold increase in $oA\beta_{1-42}$ trafficking to the CD68+ endolysosomal compartment ($p < 0.0001$) in

microglia expressing ACE (R+) compared to microglia without ACE expression (R-). Treatment

of microglia with the SYK inhibitor (P505 - PRT062607) decreased $oA\beta_{1-42}$ trafficking by 1.7-fold

in microglia not expressing ACE ($p < 0.0001$) and abrogated trafficking by 2.2-fold ($p < 0.0001$)

in microglia expressing ACE, and to levels indistinguishable from microglia not expressing ACE.

(A, B, scale = 10 µM).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

A Iba1/Aβ₁₋₄₂/SYTO

ROI (PAM)
ROI (PAM)
ROI (nPAM)

B MG (Iba1) UV Mask

C PAM: R+;AD+ vs. R-;AD+

-Log10 (p) vs. Average Log2(FC)

Phf21a
Pml
Rhox2a
Gjb5
Isy1
Rnf113a1
Shfl
Prag1
Rtel1
Rad54b
Tacc3
Lrrc47
Olfr1349
Taf1b
Postn

C130050O18Rik
Ttyh1
Kcnn2
Pde2a
Defb21
Ccdc84
Poglut3
Med6
Lrfn1
Thada

D Reactome Pathway Analysis
R+;AD+ vs. R-;AD+ (PAM)

-Log10(FDR) vs. Normalized enrichment Score

Downstream of mTOR signaling (rRNA synthesis, protein translation)
CLEC7A (Dectin-1) signaling
Activation of NF-kappaB in B cells
Beta-catenin independent WNT signaling
C-type lectin receptors (CLRs)
Degradation of beta-catenin by the destruction complex
Innate Immune System
Macroautophagy
Autophagy
Interleukin-1 family signaling

E CLEC7A Signaling

C-lectin receptor   CLEC7A receptor

R-; AD+
R+; AD+

F Downstream of mTOR

Translation   rRNA synthesis

R-; AD+
R+; AD+

Fig. 8