Quinolone resistance genes *qnr*, *aac(6')-Ib-cr*, *oqxAB*, and *qepA* in

environmental *Escherichia coli*: insights into their genetic

contexts from comparative genomics

Ryota Gomi[a, *] and Fumie Adachi[b]

[a]Department of Environmental Engineering, Graduate School of Engineering, Kyoto

University, Katsura, Nishikyo-ku, Kyoto 615-8540, Japan

[b]Department of Environmental Health, Osaka Institute of Public Health, 1-3-3 Nakamichi,
Higashinari-ku, Osaka 537-0025, Japan

[*]Corresponding author

E-mail address: gomi.ryota.34v@kyoto-u.jp; Tel: +81-75-383-3354

Running title: Quinolone resistance genes in environmental *E. coli*

ABSTRACT

Previous studies have reported the occurrence of transferable quinolone resistance

determinants in environmental *Escherichia coli*. However, little is known about their vectors

and genetic contexts. To gain insights into these genetic characteristics, we analyzed the

complete genomes of 53 environmental *E. coli* isolates, including 20 sequenced in this study

and 33 sourced from RefSeq. The following transferable quinolone resistance determinants

were detected: *qnrS1* (n = 33), *aac(6')-Ib-cr* (n = 12), *qnrS2* (n = 5), *oqxAB* (n = 4), *qnrB4* (n

= 3), *qnrD1* (n = 3), *qnrB7* (n = 2), *qnrB19* (n = 2), *qepA1* (n = 1), and *qnrA1* (n = 1). These

resistance genes were detected on plasmids of diverse replicon types; however, *aac(6')-Ib-cr*,

qnrS1, and *qnrS2* were also detected on the chromosome. The genetic contexts surrounding

these genes included not only those previously reported in clinical isolates but also novel

contexts, such as *qnrD1* embedded within a composite transposon-like structure bounded by

Tn*3*-derived inverted-repeat miniature elements (TIMEs). This study provides deep insights

into mobile genetic elements associated with transferable quinolone resistance determinants,

highlighting the importance of genomic surveillance of antimicrobial resistant bacteria in the

environment.

Keywords: *Escherichia coli*, Antimicrobial resistance, Transferable mechanisms of quinolone

resistance (TMQR), Plasmid-mediated quinolone resistance (PMQR), Genetic context

 INTRODUCTION

Escherichia coli is a commensal member of the gut microbiota of humans and animals but can

also cause intestinal and extraintestinal infections.[1] Another concern besides pathogenicity is

the rise of antimicrobial resistance (AMR) in *E. coli*.[2] This is due to the accumulation of AMR

determinants, including acquired AMR genes and chromosomal point mutations.[3]

Quinolones are antibiotics that are widely used for treatment of infections caused by a variety

of bacteria, including *E. coli*.[4] Quinolone resistance in *E. coli* is mainly attributed to mutations

in the quinolone resistance-determining regions (QRDRs) of the topoisomerase genes *gyrA* and

parC.[5,6] Although these chromosomal mutations are generally not horizontally transferred,

there are quinolone resistance determinants that can be transmitted between bacteria, called

transferable mechanisms of quinolone resistance (TMQR).[7] These resistance determinants

include target protection (*qnr*), antibiotic efflux (mainly *qepA* and *oqxAB*), and antibiotic

modification (*aac(6')-Ib-cr* and *crpP*). Although these genes are usually called plasmid-

mediated quinolone resistance (PMQR) genes, the term PMQR is not always correct because

these transferable genes can sometimes be integrated into the chromosome (hence we use the

term TMQR, not PMQR, throughout this paper).[8] It is known that the levels of resistance

conferred by TMQR are relatively low; however, TMQR can play an important role in the

development of quinolone resistance because (i) the effect of TMQR on quinolone minimum

inhibitory concentrations (MICs) is additive to that conferred by other transferable or

chromosome-mediated quinolone resistance determinants, and (ii) TMQR can facilitate the

selection of mutants with higher-level quinolone resistance.[7,9]

Information on the vectors and genetic contexts of AMR genes is important for multiple

reasons: for example, (i) it allows identification of frequently associated/linked resistance

genes and thus can provide information on co-resistance; (ii) it facilitates better understanding

of their evolution and how resistance genes spread.[10,11] Although some previous studies

performed in-depth analysis of mobile elements carrying TMQR determinants in clinical *E.*

coli, limited information exists with respect to the vectors and genetic contexts of TMQR

determinants for environmental *E. coli*.[7,8] Here, we analyzed the complete genomes of

environmental *E. coli* isolates, sequenced in this study or obtained from the NCBI reference

sequence (RefSeq) database, with the aim to acquire genomic insights into TMQR, including

their vectors and genetic contexts.

MATERIALS AND METHODS

E. coli **isolates**

Twenty-four *E. coli* isolates obtained from surface water and wastewater in Japan were

characterized in this study. All these isolates were confirmed to carry TMQR determinants by

molecular methods such as PCR. These 24 isolates included 19 isolates from our previous

collections: (i) three isolates from river water samples collected in 2010,[12] (ii) 10 isolates from

river water samples collected between 2011–2013,[13] and (iii) six extended-spectrum β-

lactamase (ESBL)-producing *E. coli* isolates obtained from municipal wastewater and hospital

wastewater in 2015.[14] Five isolates were obtained in the present study: three isolates were

screened from *E. coli* isolated from municipal wastewater using CHROMagar ECC (Kanto

Chemical Co., Tokyo, Japan) supplemented with cefotaxime (1 mg/L) between 2019–2020,

and two isolates were screened from *E. coli* isolated from river water using CHROMagar ECC

supplemented with ciprofloxacin (0.03 mg/L) between 2021–2022. Antimicrobial

susceptibility testing of the 24 isolates was performed by microdilution using Dry Plate Eiken

(Eiken, Tokyo, Japan) according to CLSI specifications (see **Table S1** for antimicrobials used).

The MIC for ciprofloxacin was also determined using the Etest (bioMérieux, Marcy-l'Étoile,

France). The results were evaluated according to CLSI criteria[15] and EUCAST epidemiological

cutoff (ECOFF) values (https://mic.eucast.org/search/).

All 24 isolates were subjected to short- and long-read sequencing in the present study as

described below (note that short reads were obtained for some of the isolates in our previous

studies but we obtained short reads again in this work because the short reads in our previous

studies had low/uneven sequencing depth).

Genome sequencing and assembly

DNA was extracted from each isolate using a DNeasy PowerSoil Pro Kit (Qiagen, Hilden,

Germany). Short-read sequencing libraries were prepared using an Illumina DNA PCR-Free

Prep kit (Illumina, San Diego, CA). The libraries were sequenced on a NovaSeq 6000 platform

(Illumina), generating paired-end reads of 151 bp each. Long-read sequencing libraries were

prepared using the SQK-LSK109 kit (Oxford Nanopore Technologies, Oxford, UK) and

sequenced on the MinION with FLO-MIN106 flow cells.

Short reads were trimmed using fastp (v0.23.2),[16] and long reads were filtered using Filtlong

(v0.2.0, https://github.com/rrwick/Filtlong). Hybrid assembly was performed with a long-read-

first approach, i.e., long-read assembly using Flye (v2.9.1-b1780)[17] followed by long-read

polishing with Medaka (v1.8.0, https://github.com/nanoporetech/medaka) and short-read

polishing with Polypolish (v0.5.0).[18] Hybrid assembly was also performed with a short-read-

first approach using Unicycler (v0.5.0).[19] If the long-read-first approach generated a complete

genome, the polished Flye assembly was chosen as a final assembly. We note that long-read-

first assemblies sometimes lack small plasmids, so we recovered small plasmids that were not

present in the Flye assembly but were present in the Unicycler assembly.[20] If the long-read-

first approach did not generate a complete genome but the short-read-first approach did, the

Unicycler assembly was chosen as a final assembly.

If the above approaches failed to generate a complete genome, we performed long-read

sequencing again using DNA prepared from the same bacterial pellet but with a different

extraction method, which employs enzymatic digestion followed by proteinase K digestion and

AMPure XP (Beckman Coulter, Inc. Brea, CA, USA) bead purification, aiming to reduce DNA

shearing and to obtain longer reads. The obtained long reads with less fragmentation were

subjected to hybrid assembly as described above.

Retrieving complete *E. coli* genomes with TMQR determinants from the NCBI database

RefSeq *E. coli* genomes with an assembly level of "complete genome" (n = 2815) were

downloaded using the ncbi-genome-download tool (v0.3.1, https://github.com/kblin/ncbi-

genome-download) in July 2023. Metadata information, including strain name and isolation

source, was extracted from the downloaded files, and genomes determined to be of

environmental origin (i.e., surface water, sediment, wastewater, and soil) (n = 203) were

retained. AMR genes were detected using ABRicate (v1.0.1,

https://github.com/tseemann/abricate) with the NCBI database. This identified 48 genomes

with TMQR determinants. For each of the 48 genomes, we searched PubMed and Google

Scholar for the associated literature report. This identified literature reports for 33 genomes,

which were further analyzed in this study. Detailed information on these 33 genomes is

summarized in **Table S2**.

Genomic analysis

The *E. coli* genomes sequenced in this study and those retrieved from RefSeq were subjected

to the following genomic analysis. AMR genes were detected using ResFinder 4.1[21] and

ABRicate (v1.0.1, https://github.com/tseemann/abricate) with the NCBI database. Multilocus

sequence   typing   (MLST)   was   performed   using   mlst   (v2.19.0,

https://github.com/tseemann/mlst). Plasmid replicons were detected using PlasmidFinder 2.1,

and plasmids were typed using pMLST 2.0.[22] Genomes were annotated using the RAST

server,[23] ISfinder,[24] and the BLAST algorithm (https://blast.ncbi.nlm.nih.gov/Blast.cgi). The

blastn analysis was performed using the nucleotide collection (nr/nt) database in April 2024.

Data availability

The complete genomes and sequence reads obtained in the present study have been deposited

in GenBank and the NCBI SRA under BioProject PRJNA1078256 (also see **Table 1** for the

nucleotide accession numbers of replicons with TMQR determinants). **Table S1** also

summarizes the nucleotide accession numbers of closed replicons for completed genomes and

SRA accession numbers for the remaining genomes.

RESULTS AND DISCUSSION

Basic characteristics of *E. coli* isolates sequenced in this study

Twenty-four *E. coli* isolates with TMQR determinants were sequenced in the present study.

These *E. coli* isolates were highly diverse and belonged to 20 different STs, including clinically

important lineages such as ST131 (**Table 1**). TMQR determinants detected include *qnrS1* (n =

12), *qnrD1* (n = 6), *aac(6')-Ib-cr* (n = 4), *qnrB7* (n = 3), *qnrS2* (n = 3), *qepA1* (n = 1), *qnrB19*

(n = 1), and *oqxAB* (n = 1) (**Table 1**). Five isolates carried multiple TMQR determinants: three

isolates carried *qnrB7* and *qnrD1*, one isolate carried *aac(6')-Ib-cr* and two copies of *qnrS2*,

and one isolate carried *aac(6')-Ib-cr*, *qnrS2*, and *oqxAB*.

Antimicrobial susceptibility testing revealed that the isolates were resistant to a wide variety

of antibiotics: nonsusceptibility rates ranged from 0% (piperacillin-tazobactam, ceftolozane-

tazobactam, cefmetazole, imipenem, meropenem, and amikacin) to 70.8% (ampicillin and

piperacillin) (**Table S1**). Nonsusceptibility rates for quinolones were 29.2% for nalidixic acid,

37.5% for ciprofloxacin (evaluated by Etest), and 37.5% for levofloxacin. All isolates were

non-susceptible to nalidixic acid, ciprofloxacin, and levofloxacin when they carried a QRDR

mutation(s). Isolates with TMQR determinants but without QRDR mutations were susceptible

to nalidixic acid but either susceptible or non-susceptible to ciprofloxacin and levofloxacin.

This is consistent with previous studies reporting that isolates with TMQR determinants but

without QRDR mutations sometimes show an unusual phenotype of nalidixic acid

susceptibility and ciprofloxacin resistance.[7] We also note that isolates with TMQR

determinants but without QRDR mutations were all classified as non-wild type to ciprofloxacin

and levofloxacin according to the ECOFF criteria, indicating that interpreting results with

ECOFF values can detect TMQR determinants when isolates do not carry QRDR mutations.

Of the 24 isolates, the genomes of 20 isolates could be completed, and these 20 completed

genomes were further analyzed to elucidate the genetic contexts of TMQR determinants.

Basic characteristics of RefSeq *E. coli* genomes with TMQR determinants

Thirty-three published RefSeq *E. coli* genomes with TMQR determinants were retrieved in the

present study. These were from wastewater (n = 18) or environmental water (n = 15), isolated

between 2012 and 2022, and from various geographical locations, with Switzerland (n = 14)

and Japan (n = 10) being dominant (**Table 2**). In total, 24 different STs were detected among

the genomes. The TMQR determinants detected include *qnrS1* (n = 21), *aac(6')-Ib-cr* (n = 9),

qnrB4 (n = 3), *oqxAB* (n = 3), *qnrS2* (n = 2), *qnrA1* (n = 1), and *qnrB19* (n = 1). Six genomes

carried multiple TMQR determinants: three genomes carried *qnrS1* and *qnrS2*, one genome

carried *oqxAB* and *aac(6')-Ib-cr*, one genome carried *qnrB19* and *qnrS1*, and one genome

carried *aac(6')-Ib-cr*, *qnrB4*, and *qnrS1*.

Genetic contexts of *qnr* genes

A RefSeq genome analyzed in this study carried *qnrA1* on an IncA/C plasmid, p009_A. (*qnrA1*

was the first TMQR gene to be discovered, in 1998[25]). IS*CR1* is usually present upstream of

qnrA1, whereas downstream structures of *qnrA1* are known to be divergent.[26] In p009_A, the

structure IS*CR1-qnrA1-ampR-qacEΔ1-sul1-orf5-orf6*-IRt-IS*6100* was identified (**Figure 1**).

This structure was previously reported in the class 1 integrons In36 and In37,[27] though a

structure (*intI1-bla*$_{VIM-1}$*-aac(6')-II-dfrA1-ΔaadA-smr*-IS*Pa21*) seen in another class 1 integron,

In4873, was detected in the upstream region in p009_A.[28] The association of In4873, which

carries a *bla*$_{VIM}$ metallo-β-lactamase gene, and the *qnrA*-containing region in In36/In37 is

worrisome from a drug-resistance perspective but seems to be rare: blastn analysis found only

one closely related structure in plasmid pME-1a (CP041734.1) from a clinical *Enterobacter*

hormaechei isolate, which carried *bla*$_{VIM-4}$ instead of *bla*$_{VIM-1}$.

Three RefSeq genomes carried *qnrB4* on different IncF plasmids (F1:A1:B1, F2:A-:B-,

F31:F36:A4:B37). In one IncF plasmid (p142_A-OXA181 [F1:A1:B1]), a ColKP3 replicon

was also detected, but its *repA* gene was interrupted. *qnrB4* is known to be globally distributed

and often associated with the AmpC type β-lactamase gene *bla*$_{DHA-1}$.[26] In all three plasmids,

bla$_{DHA-1}$ was found upstream and a *ΔintI1-dfrA17-ΔaadA5* structure was identified downstream

of *qnrB4* (**Figure 2a**). In p53_A, the *ΔintI1-dfrA17-ΔaadA5* structure was inverted and the

ΔintI1 sequence was 63-bp longer compared with the other two plasmids. Detection of the

(almost) identical genetic context in plasmids from Japan and Switzerland indicates a global

prevalence of this multiresistance region in environmental *E. coli*.

Two isolates sequenced to closure in the present study carried *qnrB7*. *qnrB7* was previously

reported to be associated with the ESBL gene *bla*$_{SHV-12}$.[26] However, reports of complete plasmid

(or chromosome) sequences carrying *qnrB7* are scarce; blastn analysis identified only four

complete plasmid sequences carrying *qnrB7*. One of them, the IncX3 plasmid

pLHST2018_IncX3 harbored by a clinical *Salmonella enterica* isolate from India, carried both

qnrB7 and *bla*$_{SHV-12}$.[29] In our isolates, *qnrB7* was also carried by IncX3 plasmids and associated

with *bla*$_{SHV-12}$ (**Figure 2b**). Various insertion sequences, including IS*26* and IS*3000*, and

remnants of transposons were found around *qnrB7* and *bla*$_{SHV-12}$, which seem to have

contributed to the mobilization of these resistance genes. Genetic contexts of *qnrB7* and *bla*$_{SHV-}$

$_{12}$ were almost identical in pKTa005_4 and pKFu015_3 except that a 1.9 kbp fragment of

IS*3000* was inserted next to another IS*3000* in pKTa005_4. The same genetic context was

identified in pLHST2018_IncX3 but was interrupted by IS*26* in *deoR*.

Two genomes, one sequenced by us and the other from RefSeq, carried *qnrB19*. *qnrB19* is

usually mobilized by transposition units (TPU) mediated by IS*Ecp1*, such as Tn*2012* and

Tn*5387*, or small ColE1-like plasmids containing fragments of these transposons.[30] The most

common *qnrB19*-carrying ColE1-like plasmid is a 2699 bp plasmid named pPAB19-

1/pSGI15/pECY6-7/pMK100, with others being minor variants of this plasmid.[30] The two

qnrB19-positive genomes carried this gene on small plasmids, p20A24_1 (3180 bp) and p32-

4_E (4417 bp), both with a ColE-like replicon Col(pHAD28).[31] These plasmids contained

putative ColE1-like *ori* (RNAI, RNAII, and *oriV*), *oriT*, and Xer sites (sites used for converting

plasmid multimers into monomers) (**Figure 2c**). p20A24_1 was found to be closely related to

plasmid pPAB19-3 (JN985534) from a clinical *E. coli* isolate, with some minor differences

(e.g., the Tn*5387* fragment was ~240bp longer in p20A24_1). p32-4_E was identified as a

cointegrate plasmid comprising the common *qnrB19*-carrying plasmid pPAB19-1 (GQ412195)

and a 1718 bp plasmid almost identical (single nucleotide difference) to pMB9272_7

(CP103528), with putative *oriT* regions being the cointegration points. pMB9272_7 carries

only two genes encoding hypothetical proteins, and no plasmid replicon was identified by

PlasmidFinder. Cointegrates comprising ColE1-like plasmids and large plasmids were

previously shown to play an important role in the evolution of AMR.[31] The example of p32-

4_E indicates cointegrates of ColE1-like plasmids and other small plasmids may also

contribute to the spread of AMR.

Three genomes sequenced to closure in the present study carried *qnrD1*. *qnrD1* was first

described on a 4270 bp plasmid, p2007057, in a clinical *Salmonella enterica* strain isolated in

China.[32] *qnrD1* has since been detected on multiple plasmids; however, blastn analysis

identified that *qnrD1* is rare in *Enterobacteriaceae* and mostly detected on ~2.7 kbp plasmids

in *Morganellaceae*. Two of the three genomes carried *qnrD1* on a plasmid identical to

p2007057 (**Figure 3**). This plasmid contains four ORFs, one of which is *mob*, in addition to

qnrD1. The remaining genome carried *qnrD1* on a 6657 bp plasmid, named pKFu015_4, with

no close blastn hits (all hits were below 50% query coverage). The *qnrD1* region, including

qnrD1 and ORF2, in pKFu015_4 was identical to ~2.7 kbp *Morganellaceae* plasmids such as

pRS12-11 (KF364953) and was ~95% identical to the corresponding region in p2007057 (and

thus pKMi029_6 and pKTa005_7). This plasmid also carried regions corresponding to the *ori*,

bom (basis of mobility), *rom* (RNA one inhibition modulator), *mob*, and *cer* (ColE1 resolution

sequence) regions in plasmid ColE1.[33] Analysis of sequences surrounding the *qnrD1* region

revealed that there were 209 bp directly oriented repeats upstream and downstream of the

region. This repeat carried imperfect 33 bp inverted repeats (IRs) closely related to some Tn*3*-

family transposons (e.g., Tn*Ec4*), though it did not encode a transposase. The blastn analysis

revealed that this repeat is identical to a Tn*3*-derived inverted-repeat miniature element (TIME),

named $TIME_{IS101}$.[34] Pairs of TIMEs are known to be involved in mobilization of intervening

sequences when the corresponding Tn*3* family transposases are provided *in trans*, and these

composite transposon-like structures were previously named TIME-COMP.[34,35] Thus, the

TIME-COMP structure, $TIME_{IS101}$-*qnrD1*-ORF2-$TIME_{IS101}$, seems to have been mobilized as

a single unit into a ColE1-like plasmid, generating pKFu015_4. The presence of 5 bp direct

repeats (AGCTA) surrounding this structure supports this idea. Moreover, we also found a

ColE1-like plasmid without insertion of this TIME-COMP structure among the genomes

sequenced in this study (pKMi029_5, CP147114), further supporting this. *qnrD1* was

previously suggested to be mobilized as a mobile insertion cassette (mic) element (explained

below),[36] but this observation indicates another mode of *qnrD1* mobilization, namely TIME-

COMP.

qnrS1 was the most prevalent TMQR gene both in our isolates (n = 12) and in RefSeq genomes

(n = 21), indicating the high prevalence of this gene in environmental *E. coli*. *qnrS1* was

detected on plasmids of diverse replicon types, and on the chromosome in two genomes (**Table**

1 and **Table 2**). Analysis of the flanking sequences revealed that the genetic contexts of *qnrS1*

can largely be classified into 13 types (type A to type M in **Figure 4**). Types B, E, and I are the

contexts found only in the genomes analyzed in this study and no other hits were found by

blastn analysis. Type A is the context detected in the original *qnrS1*-harboring plasmid

pAH0376 from a clinical *Shigella flexneri* isolate and was also detected in five isolates

analyzed in this study (note that the available pAH0376 sequence ends at *tnpR* and the presence

of IS*Kpn19* in the plasmid is unknown).[37] Insertion of a TPU (IS*Ecp1*-*bla*$_{\text{CTX-M-15}}$-*Δorf477*) into

tnpA of Tn*2* in type A, which is a context prevalent in genomes deposited in GenBank (e.g.,

CP052151) but not detected in this study, seems to have been followed by (i) truncation of *tnpR*

by IS*26* (type B), (ii) transposition of a longer TPU, including the original TPU (IS*Ecp1*-*bla*$_{\text{CTX-}}$

$_{\text{M-15}}$-*Δorf477*) and the *qnrS1* region, into the chromosome by IS*Ecp1* (type C), or (iii) partial

deletion of Tn*2* and IS*Ecp1* and IS*26*-mediated truncation of IS*Kpn19* (type D). Truncation of

type A by two IS*26* elements might have led to generation of type E. IS*Kpn19* elements are

present both upstream and downstream of *qnrS1* in type F. The blastn analysis identified the

type F context in only two replicons, *E. coli* plasmid pA1.S1.126.c2 (CP146587) from human

urine and *E. coli* plasmid pKT58A (JX065631) from a wild bird in Slovakia.[38] In type G, IS*Ecl2*

is truncated by IS*26*, and class 1 integrons with different cassette arrays are present upstream

of *qnrS1*. The type H context contained the *ftsI-bla*$_{LAP-2}$-IS*Ecl2* structure upstream of *qnrS1*,

which was reported previously in clinical *Klebsiella pneumoniae* and was also detected in

multiple *Enterobacteriaceae* plasmids by blastn analysis.[39] Truncation of type H by IS*26* at

bla$_{LAP-2}$ seems to have given rise to type I. Type J and type K contexts are inserted in the same

position in each IncX1 plasmid and flanked by 5-bp direct repeats of ATAAC (or the reverse

complement sequence GTTAT). Insertion of Tn*2* in the ancestral IncX1 plasmid followed by

IS*26*-mediated rearrangements, including insertion of the region containing *qnrS1*, can explain

these contexts. Type L and Type M seem to have been derived from the *qnrS1*-bearing

transposon Tn*6292*. Tn*6292* itself was not detected in the present study, though blastn analysis

identified that intact Tn*6292* is almost restricted to *Enterobacteriaceae* plasmids from China.

Two genomes sequenced to closure in this study and two RefSeq genomes carried *qnrS2*. *qnrS2*

was previously reported to be part of a mic element bracketed by 22 bp imperfect IRs in the

Aeromonas punctata plasmid p37.[40] A mic element is a nonautonomous element bracketed by

two (imperfect) IRs. A mic does not carry a transposase gene but can be mobilized *in trans* by

a transposase that can recognize the IR. In two RefSeq genomes, *qnrS2* was embedded within

the same genetic context on IncX1 plasmids, but the mic element was truncated by IS*26*

(**Figure 5**). The IS*26*-*qnrS2*-*ΔmpR*-IS*26* structure, which was also detected in *E. coli* plasmids

such as pRW7-1_235k_tetX (MT219825), was followed by the *fosA3* genetic context type K,[41]

linking *qnrS2* to *fosA3*, *Δbla*_{TEM}, and *bla*_{CTX-M-55}. Two ST2179 genomes sequenced to closure

in this study carried *qnrS2* on the chromosome. In both genomes, *qnrS2* was within the

truncated mic element and was surrounded by directly-oriented IS*26* sequences. This IS*26*-

qnrS2-*ΔmpR*-IS*26* structure (note that this structure is different from that in the IncX1 plasmids,

in that the IS*26* sequences are directly-oriented and a ~200 bp fragment is present between the

truncated mic and the left IS*26*) was also detected in some *Enterobacteriaceae* plasmids such

as pSH16G4525 (MH522424). Interestingly, this *qnrS2* region was duplicated in 19M19. In

both ST2179 genomes, the *qnrS2* region was followed by *E. coli* chromosomal genes, but the

gene contents were different, likely due to the rearrangement caused by intramolecular

replicative transposition of IS*26*.[35]

Genetic contexts of *aac(6')-Ib-cr*

Three genomes sequenced to closure in this study and nine RefSeq genomes carried *aac(6')-*

Ib-cr. Notably, *aac(6')-Ib-cr* was detected on the chromosome in six isolates (50%) belonging

to five different STs (ST10, ST410, ST648, ST1266, and ST2179). A chromosomal location

for *aac(6')-Ib-cr* was also reported previously in clinical ST131 and ST648 isolates from

Spain.[8] These examples indicate chromosomal integration of *aac(6')-Ib-cr* is frequent and has

occurred multiple times in different STs, again highlighting that the term PMQR is not

appropriate to describe transferable quinolone resistance genes. Plasmidic *aac(6')-Ib-cr* genes

were detected on different IncF plasmids (F1:A1:B16, F1:A1:B49, F10:A1:B49, F36:A4:B-,

F36:A4:B1, F47:A-:B75). One plasmid, pWP4-W18-ESBL-09_1 (F1:A1:B16), also carried a

Col156-like replicon, but the *repA* gene seems to be truncated.

aac(6′)-Ib-cr has been identified in a gene cassette as part of class 1 integrons.[42] Although

aac(6′)-Ib-cr is sometimes linked to the 5′-conserved segment (5′-CS) and the 3′-conserved

segment (3′-CS) in class 1 integrons, integrons containing *aac(6′)-Ib-cr* are often interrupted

by IS*26*.[8,43] Our analysis supports this observation, having identified deletion/truncation of

either or both of 5′-CS and 3′-CS by IS*26* in 11 of 12 cases (**Figure 6**). Apparently, the

truncated structures seen in the 11 genomes were derived from the class 1 integron *intI1-*

aac(6′)-Ib-cr-bla$_{OXA-1}$*-catB3-arr-3-qacEΔ1-sul1*. Among them, the IS*26-aac(6′)-Ib-cr-bla*$_{OXA-}$

$_{1}$*-ΔcatB3*-IS*26* structure was prevalent and detected in seven genomes (58%), while the

upstream and downstream sequences were divergent. The blastn analysis revealed that this

structure is prevalent among the public genomes (more than 1000 hits with >99% coverage and

with >99% identity). It should be noted, however, that the IS*26* sequences are inversely

oriented and thus this structure is neither a transposon nor a pseudo compound transposon.[44]

The prevalence of this structure among various plasmids/chromosomes can partially be

explained by the presence of other mobile genetic elements surrounding it, which might have

aided the movement of multiresistance regions containing this structure.

Genetic contexts of *oqxAB*

Mobile *oqxAB* genes are commonly found within the IS*26-oqxA-oqxB-oqxR*-IS*26* structure,

named Tn*6010*.[45] All four *oqxAB* genes identified in the present study were situated within

Tn*6010* (or a truncated Tn*6010*) and carried by plasmids of different Inc types (**Figure 7**).

While the downstream sequences flanking Tn*6010* were divergent in the four plasmids, three

RefSeq plasmids carried common sequences upstream of Tn*6010*, containing *tnpA* of Tn*2*,

IS*26*, *bleO*, *nimC*/*nimA*, and *ΔISEnca1*. The blastn analysis using this upstream structure +

Tn*6010* as a query sequence identified more than 100 genomes carrying this structure. On the

other hand, the upstream structure of Tn*6010* was replaced by a truncated Tn*1721* and a

truncated Tn*2* in pJKHS004_1, linking *tetA* and a truncated *bla*$_{TEM}$ to *oqxAB*. No genomes were

found by blastn analysis to carry the multiresistance region detected in pJKHS004_1.

Genetic context of *qepA1*

qepA genes are typically associated with IS*CR3* and embedded within complex integrons.[46]

One isolate sequenced in this study carried *qepA1*, and the gene was located between a

truncated *intI1* and IS*CR3* on an incF plasmid (C4:F18:A-:B1), pKMi012_1 (**Figure 8**). This

region was followed by a *ΔintI1*/*groEL-dfrB4-qacEΔ1-sul1-Δorf5* region, the chromate

resistance (*chrA*) region, and the macrolide resistance (*mph*(A)) region. The blastn analysis of

the entire resistance region bounded by IS*26* revealed almost identical sequences (100% query

coverage and >99% identity) in three *E. coli* plasmids, namely pJJ1887-5 (CP014320.1) from

a clinical isolate in the USA, pM216_mF (LC492469.1) from a clinical isolate in Myanmar,

and pMyNCGM604 (LC744503.1) from an isolate of unknown origin in Myanmar, indicating

a wide distribution of this multiresistance region.

Study limitations

This study has some limitations. The genomes of four isolates (KMi014, KTa007, KTa009,

JKHS019) could not be completed despite repeated long-read sequencing. This is probably due

to the presence of long repeats in these genomes. We also included RefSeq genomes in our

analysis to extend our dataset, but most RefSeq genomes were from Japan and Switzerland.

This seems to reflect the situation where long-read sequencing is still not widely used to study

antibiotic resistance in environmental bacteria.

CONCLUSIONS

Here, we performed in-depth analysis of the genetic contexts of TMQR determinants in

environmental *E. coli*. The genetic contexts described in this study included those closely

related to the contexts previously reported in clinical isolates. We also detected novel genetic

contexts, such as *qnrD1* embedded within TIME-COMP. Some of the TMQR genes were

frequently associated with other AMR genes in the described contexts (e.g., *qnrB7* and $bla_{SHV\text{-}}$

$_{12}$), and this information will contribute to our understanding of how multidrug resistance

emerges in bacteria. We also found that certain TMQR determinants (i.e., *qnrS1*, *qnrS2*, and

aac(6′)-Ib-cr) are located not only on plasmids but also on the chromosome, encouraging the

use of the term TMQR rather than PMQR to avoid erroneous interpretations in future studies.

Overall, this study provides valuable insights into mobile genetic elements associated with

TMQR determinants and highlights the importance of genomic surveillance of antimicrobial

resistant bacteria in the environment.

ACKNOWLEDGEMENTS

This work was supported by JSPS KAKENHI (grant number JP22K18038), the Kurita Water

and Environment Foundation (22B006), the Sumitomo Foundation (2230113), and the

Environment Research and Technology Development Fund (JPMEERF20235R01) of the

Environmental Restoration and Conservation Agency, provided by the Ministry of the

Environment of Japan. Computations were partially performed on the NIG supercomputer at

ROIS National Institute of Genetics. We acknowledge the NGS core facility of the Genome

Information Research Center at the Research Institute for Microbial Diseases of Osaka

University for the support in DNA sequencing. We also thank Isao Nakamoto and Yasufumi

Matsumura for their support in antimicrobial susceptibility testing.

AUTHOR CONTRIBUTIONS

Conceptualization: R.G.; methodology: R.G.; validation: R.G.; formal analysis: R.G.;

investigation: R.G. and F.A; resources: R.G. and F.A; data curation: R.G.; visualization: R.G.;

writing – original draft: R.G.; writing – review & editing: R.G. and F.A.

References

1   Kaper, J. B., Nataro, J. P. & Mobley, H. L. Pathogenic *Escherichia coli*. *Nat Rev*
Microbiol **2**, 123-140 (2004). https://doi.org/10.1038/nrmicro818
2   Denamur, E., Clermont, O., Bonacorsi, S. & Gordon, D. The population genetics of
pathogenic *Escherichia coli*. *Nat Rev Microbiol* **19**, 37-54 (2021).
https://doi.org/10.1038/s41579-020-0416-x
3   Poirel, L. *et al*. Antimicrobial Resistance in *Escherichia coli*. *Microbiol Spectr* **6** (2018).
https://doi.org/10.1128/microbiolspec.ARBA-0026-2017
4   Aldred, K. J., Kerns, R. J. & Osheroff, N. Mechanism of quinolone action and resistance.
Biochemistry **53**, 1565-1574 (2014). https://doi.org/10.1021/bi5000564
5   Aoike, N. *et al*. Molecular characterization of extraintestinal *Escherichia coli* isolates
in Japan: relationship between sequence types and mutation patterns of quinolone
resistance-determining regions analyzed by pyrosequencing. *J Clin Microbiol* **51**,
1692-1698 (2013). https://doi.org/10.1128/JCM.03049-12
6   van der Putten, B. C. L. *et al*. Quantifying the contribution of four resistance
mechanisms to ciprofloxacin MIC in *Escherichia coli*: a systematic review. *J*
Antimicrob Chemother **74**, 298-310 (2019). https://doi.org/10.1093/jac/dky417
7   Ruiz, J. Transferable Mechanisms of Quinolone Resistance from 1998 Onward. *Clin*
Microbiol Rev **32** (2019). https://doi.org/10.1128/CMR.00007-19
8   Ruiz, E. *et al*. qnr, aac(6')-Ib-cr and qepA genes in *Escherichia coli* and *Klebsiella* spp.:
genetic environments and plasmid and chromosomal location. *J Antimicrob Chemother*
67, 886-897 (2012). https://doi.org/10.1093/jac/dkr548
9   Hooper, D. C. & Jacoby, G. A. Mechanisms of drug resistance: quinolone resistance.
Ann N Y Acad Sci **1354**, 12-31 (2015). https://doi.org/10.1111/nyas.12830
10   Partridge, S. R. Analysis of antibiotic resistance regions in Gram-negative bacteria.
FEMS Microbiol Rev **35**, 820-855 (2011). https://doi.org/10.1111/j.1574-
6976.2011.00277.x
11   Partridge, S. R. & Tsafnat, G. Automated annotation of mobile antibiotic resistance in
Gram-negative bacteria: the Multiple Antibiotic Resistance Annotator (MARA) and

database. *J Antimicrob Chemother* **73**, 883-890 (2018).
https://doi.org/10.1093/jac/dkx513

12   Adachi, F., Yamamoto, A., Takakura, K. & Kawahara, R. Occurrence of
fluoroquinolones and fluoroquinolone-resistance genes in the aquatic environment. *Sci*
Total Environ **444**, 508-514 (2013). https://doi.org/10.1016/j.scitotenv.2012.11.077

13   Gomi, R. *et al*. Whole-Genome Analysis of Antimicrobial-Resistant and Extraintestinal
Pathogenic *Escherichia coli* in River Water. *Appl Environ Microbiol* **83** (2017).
https://doi.org/10.1128/AEM.02703-16

14   Gomi, R. *et al*. Occurrence of Clinically Important Lineages, Including the Sequence
Type 131 C1-M27 Subclone, among Extended-Spectrum-beta-Lactamase-Producing
Escherichia coli in Wastewater. *Antimicrob Agents Chemother* **61** (2017).
https://doi.org/10.1128/AAC.00564-17

15   CLSI. Performance Standards for Antimicrobial Susceptibility Testing—Thirtieth
Edition: M100.   (2020).

16   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**, i884-i890 (2018).
https://doi.org/10.1093/bioinformatics/bty560

17   Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone
reads using repeat graphs. *Nat Biotechnol* **37**, 540-546 (2019).
https://doi.org/10.1038/s41587-019-0072-8

18   Wick, R. R. & Holt, K. E. Polypolish: Short-read polishing of long-read bacterial
genome assemblies. *PLoS Comput Biol* **18**, e1009802 (2022).
https://doi.org/10.1371/journal.pcbi.1009802

19   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**,
e1005595 (2017). https://doi.org/10.1371/journal.pcbi.1005595

20   Wick, R. R., Judd, L. M. & Holt, K. E. Assembling the perfect bacterial genome using
Oxford Nanopore and Illumina sequencing. *PLoS Comput Biol* **19**, e1010905 (2023).
https://doi.org/10.1371/journal.pcbi.1010905

21   Bortolaia, V. *et al*. ResFinder 4.0 for predictions of phenotypes from genotypes. *J*
Antimicrob Chemother **75**, 3491-3500 (2020). https://doi.org/10.1093/jac/dkaa345

22   Carattoli, A. *et al*. *In silico* detection and typing of plasmids using PlasmidFinder and
plasmid multilocus sequence typing. *Antimicrob Agents Chemother* **58**, 3895-3903
(2014). https://doi.org/10.1128/AAC.02412-14

23   Aziz, R. K. *et al*. The RAST Server: rapid annotations using subsystems technology.
BMC Genomics **9**, 75 (2008). https://doi.org/10.1186/1471-2164-9-75

24 Siguier, P., Perochon, J., Lestrade, L., Mahillon, J. & Chandler, M. ISfinder: the
reference centre for bacterial insertion sequences. *Nucleic Acids Res* **34**, D32-36 (2006).
https://doi.org/10.1093/nar/gkj014

25 Martinez-Martinez, L., Pascual, A. & Jacoby, G. A. Quinolone resistance from a
transferable plasmid. *Lancet* **351**, 797-799 (1998). https://doi.org/10.1016/S0140-
6736(97)07322-4

26 Strahilevitz, J., Jacoby, G. A., Hooper, D. C. & Robicsek, A. Plasmid-mediated
quinolone resistance: a multifaceted threat. *Clin Microbiol Rev* **22**, 664-689 (2009).
https://doi.org/10.1128/CMR.00016-09

27 Wang, M. *et al*. Plasmid-mediated quinolone resistance in clinical isolates of
Escherichia coli from Shanghai, China. *Antimicrob Agents Chemother* **47**, 2242-2248
(2003). https://doi.org/10.1128/AAC.47.7.2242-2248.2003

28 Papagiannitsis, C. C. *et al*. Survey of metallo-beta-lactamase-producing
Enterobacteriaceae colonizing patients in European ICUs and rehabilitation units,
2008-11. *J Antimicrob Chemother* **70**, 1981-1988 (2015).
https://doi.org/10.1093/jac/dkv055

29 Jacob, J. J. *et al*. *Salmonella* Typhi acquires diverse plasmids from other
Enterobacteriaceae to develop cephalosporin resistance. *Genomics* **113**, 2171-2176
(2021). https://doi.org/10.1016/j.ygeno.2021.05.003

30 Tran, T. *et al*. Small plasmids harboring *qnrB19*: a model for plasmid evolution
mediated by site-specific recombination at *oriT* and Xer sites. *Antimicrob Agents*
Chemother **56**, 1821-1827 (2012). https://doi.org/10.1128/AAC.06036-11

31 Ares-Arroyo, M., Rocha, E. P. C. & Gonzalez-Zorn, B. Evolution of ColE1-like
plasmids across gamma-Proteobacteria: From bacteriocin production to antimicrobial
resistance. *PLoS Genet* **17**, e1009919 (2021).
https://doi.org/10.1371/journal.pgen.1009919

32 Cavaco, L. M., Hasman, H., Xia, S. & Aarestrup, F. M. *qnrD*, a novel gene conferring
transferable quinolone resistance in *Salmonella enterica* serovar Kentucky and
Bovismorbificans strains of human origin. *Antimicrob Agents Chemother* **53**, 603-608
(2009). https://doi.org/10.1128/AAC.00997-08

33 Chan, P. T., Ohmori, H., Tomizawa, J. & Lebowitz, J. Nucleotide sequence and gene
organization of ColE1 DNA. *J Biol Chem* **260**, 8925-8935 (1985).

34 Szuplewska, M., Ludwiczak, M., Lyzwa, K., Czarnecki, J. & Bartosik, D. Mobility and
generation of mosaic non-autonomous transposons by Tn*3*-derived inverted-repeat
miniature elements (TIMEs). *PLoS One* **9**, e105010 (2014).
https://doi.org/10.1371/journal.pone.0105010

Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. Mobile Genetic Elements Associated with Antimicrobial Resistance. *Clin Microbiol Rev* **31** (2018). https://doi.org/10.1128/CMR.00088-17

Guillard, T. *et al*. Mobile insertion cassette elements found in small non-transmissible plasmids in *Proteeae* may explain *qnrD* mobilization. *PLoS One* **9**, e87801 (2014). https://doi.org/10.1371/journal.pone.0087801

Hata, M. *et al*. Cloning of a novel gene for quinolone resistance from a transferable plasmid in *Shigella flexneri* 2b. *Antimicrob Agents Chemother* **49**, 801-803 (2005). https://doi.org/10.1128/AAC.49.2.801-803.2005

Dolejska, M., Villa, L., Hasman, H., Hansen, L. & Carattoli, A. Characterization of IncN plasmids carrying $bla_{CTX-M-1}$ and *qnr* genes in *Escherichia coli* and *Salmonella* from animals, the environment and humans. *J Antimicrob Chemother* **68**, 333-339 (2013). https://doi.org/10.1093/jac/dks387

Karah, N. *et al*. Plasmid-mediated quinolone resistance determinants *qnr* and *aac(6')-Ib-cr* in *Escherichia coli* and *Klebsiella* spp. from Norway and Sweden. *Diagn Microbiol Infect Dis* **66**, 425-431 (2010). https://doi.org/10.1016/j.diagmicrobio.2009.12.004

Cattoir, V., Poirel, L., Aubert, C., Soussy, C. J. & Nordmann, P. Unexpected occurrence of plasmid-mediated quinolone resistance determinants in environmental *Aeromonas* spp. *Emerg Infect Dis* **14**, 231-237 (2008). https://doi.org/10.3201/eid1402.070677

Lv, L. *et al*. Multiple Plasmid Vectors Mediate the Spread of *fosA3* in Extended-Spectrum-beta-Lactamase-Producing *Enterobacterales* Isolates from Retail Vegetables in China. *mSphere* **5** (2020). https://doi.org/10.1128/mSphere.00507-20

Rodriguez-Martinez, J. M. *et al*. Plasmid-mediated quinolone resistance: Two decades on. *Drug Resist Updat* **29**, 13-29 (2016). https://doi.org/10.1016/j.drup.2016.09.001

Partridge, S. R., Zong, Z. & Iredell, J. R. Recombination in IS*26* and Tn*2* in the evolution of multiresistance regions carrying $bla_{CTX-M-15}$ on conjugative IncF plasmids from *Escherichia coli*. *Antimicrob Agents Chemother* **55**, 4971-4978 (2011). https://doi.org/10.1128/AAC.00025-11

Harmer, C. J., Pong, C. H. & Hall, R. M. Structures bounded by directly-oriented members of the IS*26* family are pseudo-compound transposons. *Plasmid* **111**, 102530 (2020). https://doi.org/10.1016/j.plasmid.2020.102530

Norman, A., Hansen, L. H., She, Q. & Sorensen, S. J. Nucleotide sequence of pOLA52: a conjugative IncX1 plasmid from *Escherichia coli* which enables biofilm formation and multidrug efflux. *Plasmid* **60**, 59-74 (2008). https://doi.org/10.1016/j.plasmid.2008.03.003

46 Cattoir, V., Poirel, L. & Nordmann, P. Plasmid-mediated quinolone resistance pump
QepA2 in an *Escherichia coli* isolate from France. *Antimicrob Agents Chemother* **52**,
3801-3804 (2008). https://doi.org/10.1128/AAC.00638-08

47 Lin, Y. *et al*. Metadata Analysis of mcr-1-Bearing Plasmids Inspired by the Sequencing
Evidence for Horizontal Transfer of Antibiotic Resistance Genes Between Polluted
River and Wild Birds. *Front Microbiol* **11**, 352 (2020).
https://doi.org/10.3389/fmicb.2020.00352

48 Long, H. *et al*. The co-transfer of plasmid-borne colistin-resistant genes *mcr-1* and *mcr-*
3.5, the carbapenemase gene bla_{NDM-5} and the 16S methylase gene *rmtB* from
Escherichia coli. *Sci Rep* **9**, 696 (2019). https://doi.org/10.1038/s41598-018-37125-1

49 Che, Y. *et al*. High-resolution genomic surveillance elucidates a multilayered
hierarchical transfer of resistance between WWTP- and human/animal-associated
bacteria. *Microbiome* **10**, 16 (2022). https://doi.org/10.1186/s40168-021-01192-w

50 Ju, X., Wu, Y., Chen, G. & Zhang, R. *Escherichia coli* high-risk clone ST410 harboring
bla_{NDM-13} isolated from hospital wastewater in China. *Environ Sci Pollut Res Int* **30**,
91487-91491 (2023). https://doi.org/10.1007/s11356-023-28193-6

51 Gomi, R., Yamamoto, M., Tanaka, M. & Matsumura, Y. Chromosomal integration of
bla_{CTX-M} genes in diverse *Escherichia coli* isolates recovered from river water in Japan.
Curr Res Microb Sci **3**, 100144 (2022). https://doi.org/10.1016/j.crmicr.2022.100144

52 Sekizuka, T., Inamine, Y., Segawa, T. & Kuroda, M. Characterization of NDM-5- and
CTX-M-55-coproducing *Escherichia coli* GSH8M-2 isolated from the effluent of a
wastewater treatment plant in Tokyo Bay. *Infect Drug Resist* **12**, 2243-2249 (2019).
https://doi.org/10.2147/IDR.S215273

53 Sekizuka, T., Tanaka, R., Hashino, M., Yatsu, K. & Kuroda, M. Comprehensive
Genome and Plasmidome Analysis of Antimicrobial Resistant Bacteria in Wastewater
Treatment Plant Effluent of Tokyo. *Antibiotics (Basel)* **11** (2022).
https://doi.org/10.3390/antibiotics11101283

54 Finton, M. D., Meisal, R., Porcellato, D., Brandal, L. T. & Lindstedt, B. A. Whole
Genome Sequencing and Characterization of Multidrug-Resistant (MDR) Bacterial
Strains Isolated From a Norwegian University Campus Pond. *Front Microbiol* **11**, 1273
(2020). https://doi.org/10.3389/fmicb.2020.01273

55 Biggel, M., Zurfluh, K., Treier, A., Nuesch-Inderbinen, M. & Stephan, R.
Characteristics of *fosA*-carrying plasmids in *E. coli* and *Klebsiella* spp. isolates
originating from food and environmental samples. *J Antimicrob Chemother* **76**, 2004-
2011 (2021). https://doi.org/10.1093/jac/dkab119

56 Bleichenbacher, S. *et al*. Environmental dissemination of carbapenemase-producing
Enterobacteriaceae in rivers in Switzerland. *Environ Pollut* **265**, 115081 (2020).
https://doi.org/10.1016/j.envpol.2020.115081

57 Kurekci, C. *et al*. Emergence and Characterization of Tigecycline Resistance Gene
tet(X4) in ST609 *Escherichia coli* Isolates from Wastewater in Turkey. *Microbiol*
Spectr **10**, e0073222 (2022). https://doi.org/10.1128/spectrum.00732-22
58 Liedhegner, E. *et al*. Similarities in Virulence and Extended Spectrum Beta-Lactamase
Gene Profiles among Cefotaxime-Resistant *Escherichia coli* Wastewater and Clinical
Isolates. *Antibiotics (Basel)* **11** (2022). https://doi.org/10.3390/antibiotics11020260

TABLES

Table 1. Characteristics of 24 *E. coli* isolates sequenced in this study[a]

Isolate	Source	Year of isolation	ST	TMQR determinant(s)[b]	Replicon with TMQR determinant(s) (size, nucleotide accession number)	QRDRs[c]
1002W03	river water	2010	ST91	*qnrS1* (type F)	IncN plasmid (43766 bp, CP147054)	SDSE (0)
1008S13	river water	2010	ST1141	*qnrS1* (type I)	IncF plasmid (65333 bp, CP147052)	SDSE (0)
1008S19	river water	2010	ST216	*qnrS1* (type H)	IncF/IncN plasmid (50257 bp, CP147051)	SDSE (0)
KFu015	river water	2011	ST215	*qnrB7, qnrD1*	*qnrB7*: IncX3 plasmid (46557 bp, CP147133) *qnrD1*: Col3M/ColRNAI plasmid (6657 bp, CP147134)	SDSE (0)
KMi012	river water	2011	ST46	*qepA1*	IncF plasmid (140174 bp, CP147126)	LNIE (3)
KMi029	river water	2011	ST216	*qnrD1*	Col3M plasmid (4270 bp, CP147115)	SDSE (0)
KMi014[d]	river water	2011	ST607	*qnrB7, qnrD1*	-	SDSE (0)
KKa014	river water	2012	ST58	*qnrS1* (type A)	IncY plasmid (131602 bp, CP147106)	SDSE (0)
KFu021	river water	2012	ST58	*qnrS1* (type A)	IncY plasmid (130264 bp, CP147102)	SDSE (0)
KOr024	river water	2013	ST101	*qnrS1* (type J)	IncX1 plasmid (41491 bp, CP147097)	SDSE (0)

KTa005	river water	2013	ST542	*qnrB7, qnrD1*	*qnrB7*: IncX3 plasmid (48469 bp, CP147088) *qnrD1*: Col3M plasmid (4270 bp, CP147091)	SDSE (0)
KTa007[d]	river water	2013	ST218	*qnrD1*	-	SDSE (0)
KTa009[d]	river water	2013	ST218	*qnrD1*	-	SDSE (0)
JSWP006	municipal wastewater	2015	ST393	*qnrS1* (type E)	IncF plasmid (123573 bp, CP147082)	LNIE (3)
JSWP021	municipal wastewater	2015	ST224	*aac(6')-Ib-cr*	IncF plasmid (120365 bp, CP147076)	LNIE (3)
JKHS004	hospital wastewater	2015	ST2179	*qnrS2, aac(6')-Ib-cr, oqxA, oqxB*	*qnrS2, aac(6')-Ib-cr*: chromosome (4949596 bp, CP147069) *oqxA, oqxB*: IncF plasmid (143870 bp, CP147070)	LDIE (2)
JKHS006	hospital wastewater	2015	ST131	*qnrS1* (type L)	IncN plasmid (63327 bp, CP147064)	LDSE (1)
JKHS016	hospital wastewater	2015	ST43	*qnrS1* (type D)	IncI1-I(Alpha) plasmid (100666 bp, CP147060)	SDSE (0)
JKHS019[d]	hospital wastewater	2015	ST410	*aac(6')-Ib-cr*	-	LNIE (3)
19M19	municipal wastewater	2019	ST2179	*qnrS2* (two copies), *aac(6')-Ib-cr*	*qnrS2* (two copies), *aac(6')-Ib-cr*: chromosome (4870130 bp, CP147048)	LDIE (2)
19A14	municipal wastewater	2019	ST38	*qnrS1* (type C)	chromosome (4879959 bp, CP147046)	SDSE (0)
20A24	municipal wastewater	2020	ST10	*qnrB19*	Col(pHAD28) plasmid (3180 bp, CP147045)	SDSE (0)

| 21F61 | river water | 2021 | ST657 | *qnrS1* (type J) | IncX1 plasmid (48446 bp, CP147040) | SDSE (0) |
| 22F46 | river water | 2022 | ST540 | *qnrS1* (type A) | IncF plasmid (100207 bp, CP147033) | SDSE (0) |

[a]Detailed information on each isolate, such as other AMR genes detected, is available in **Table**

S1.

[b]The context type is shown for *qnrS1*.

[c]GyrA codons 83 and 87 and ParC codons 80 and 84 are shown in this order. SDSE is the wild-

type. The number of amino acid substitutions is shown in parentheses.

[d]The genomes of these four isolates could not be completed.

Table 2. Characteristics of 33 RefSeq *E. coli* genomes with TMQR determinants[a]

Isolate[(reference)]	Source	Location	Year of isolation	ST	TMQR determinant(s)[b]	Replicon with TMQR determinant(s) (size, nucleotide accession number)	QRDRs[c]
W5-6[47]	river water	China	2015	ST744	*oqxA, oqxB*	IncHI2/IncHI2A plasmid (241043 bp, NZ_CP032993.1)	LNIE (3)
WCHEC025943[48]	hospital wastewater	China	2017	ST410	*oqxA, oqxB, aac(6')-Ib-cr*	*oqxA*, *oqxB*: IncHI2/IncHI2A/IncN plasmid (265538 bp, NZ_CP027202.2) *aac(6')-Ib-cr*: IncF plasmid (75779 bp, NZ_CP027200.1)	LNIE (3)
SWHEFF_58[49]	municipal wastewater	China	2018	ST38	*qnrS1* (type K)	IncX1 plasmid (46569 bp, NZ_CP055050.1)	SDSE (0)
STLEFF_47[49]	municipal wastewater	China	2018	ST117	*qnrS1* (type G)	IncI1-I(Alpha) plasmid, (121650 bp, NZ_CP058876.1)	SDSE (0)
C-SRM-3[50]	hospital wastewater	China	2022	ST410	*aac(6')-Ib-cr*	chromosome (4799348 bp, NZ_CP123197.1)	LNIE (3)
KFu023[51]	river water	Japan	2013	ST10	*aac(6')-Ib-cr*	chromosome (4672002 bp, NZ_CP091662.1)	LNIE (3)

GSH8M-2[52]	municipal wastewater	Japan	2018	ST542	*qnrS1* (type G), *qnrS2*	*qnrS1*: IncF plasmid (145389 bp, NZ_AP019676.1) *qnrS2*: IncX1 plasmid (49365 bp, NZ_AP019678.1)	SDSE (0)
WP5-S17-ESBL-11[53]	municipal wastewater	Japan	2017	ST58	*qnrS1* (type C)	chromosome (4781831 bp, NZ_AP022107.1)	SDSE (0)
WP2-W18-CRE-03[53]	municipal wastewater	Japan	2018	ST602	*oqxA, oqxB*	IncF/IncQ1 plasmid (158754 bp, NZ_AP021915.1)	LNIE (3)
WP4-W18-ESBL-09[53]	municipal wastewater	Japan	2018	ST405	*aac(6')-Ib-cr*	IncF/Col156 plasmid (117617 bp, NZ_AP022070.1)	LNIE (3)
WP7-S17-ESBL-04[53]	municipal wastewater	Japan	2017	ST10	*qnrS1* (type B)	IncF/IncN plasmid (123262 bp, NZ_AP022178.1)	SDSE (0)
WP8-S17-ESBL-05[53]	municipal wastewater	Japan	2017	ST641	*qnrS1* (type G)	IncX1 plasmid (53216 bp, NZ_AP022218.1)	SDSE (0)
WP8-S18-CRE-02[53]	municipal wastewater	Japan	2018	ST542	*qnrS1* (type G), *qnrS2*	*qnrS1*: IncF plasmid (145389 bp, NZ_AP022246.1)	SDSE (0)

						qnrS2: IncX1 plasmid (49365 bp, NZ_AP022248.1)	
WP8-S18-ESBL-07[53]	municipal wastewater	Japan	2018	ST2858	*qnrS1* (type D)	IncB/O/K/Z plasmid (91233 bp, NZ_AP022263.1)	SDSE (0)
BEC1-S17-ESBL-09[53]	sea water	Japan	2017	ST130	*qnrB4*	IncF plasmid (83140 bp, NZ_AP022300.1)	LDSE (1)
NMBU_W12E19[54]	surface water	Norway	2019	ST1286	*qnrS1* (type D)	IncI1-I(Alpha) plasmid (88772 bp, NZ_CP042886.1)	SDSE (0)
OW1E2[55]	surface water	Switzerland	2012	ST11280	*qnrS1* (type H)	IncF plasmid (74070 bp, NZ_CP067247.1)	LDSE (1)
ABW_S38[55]	wastewater	Switzerland	2015	ST215	*qnrS1* (type H)	IncF plasmid (53010 bp, NZ_CP067268.1)	SDSE (0)
027_SK3[55]	wastewater	Switzerland	2016	ST215	*qnrS1* (type H)	IncF plasmid (104584 bp, NZ_CP067313.1)	SDSE (0)
ABW_A19[55]	wastewater	Switzerland	2015	ST1266	*aac(6')-Ib-cr*	chromosome (5250599 bp, NZ_CP067307.1)	LNIE (3)
CF009[56]	surface water	Switzerland	2019	ST73	*qnrA1*	IncA/C plasmid (159959 bp, NZ_CP048305.1)	SDSE (0)
CF032[56]	surface water	Switzerland	2019	ST1284	*qnrB19, qnrS1* (type M)	*qnrB19*: Col(pHAD28) plasmid (4417 bp, NZ_CP048315.1)	LNIE (3)

						qnrS1: IncX3/ColKP3 plasmid (50168 bp, NZ_CP048321.1)	
CF064[56]	surface water	Switzerland	2019	ST940	*qnrS1* (type M)	IncX3/ColKP3 plasmid (50136 bp, NZ_CP048325.1)	SDSE (0)
CF142[56]	surface water	Switzerland	2019	ST648	*qnrS1* (type M)*, qnrB4, aac(6')-Ib-cr*	*qnrS1*, *qnrB4*: IncF/ColKP3 plasmid (155486 bp, NZ_CP048338.1) *aac(6')-Ib-cr*: chromosome (5073188 bp, NZ_CP048337.1)	LNIE (3)
CF061[56]	surface water	Switzerland	2019	ST940	*qnrS1* (type M)	IncX3/ColKP3 plasmid (51471 bp, NZ_CP048327.1)	SDSE (0)
CF010[56]	surface water	Switzerland	2019	ST656	*qnrS1* (type M)	IncX3/ColKP3 plasmid (51479 bp, NZ_CP048332.1)	LDSE (1)
CF124[56]	surface water	Switzerland	2019	ST410	*qnrS1* (type M)	IncX3/ColKP3 plasmid (51466 bp, NZ_CP048346.1)	LNIE (3)
CF053[56]	surface water	Switzerland	2019	ST205	*qnrB4*	IncF plasmid (144585 bp, NZ_CP048360.1)	LNIE (3)
CF164[56]	surface water	Switzerland	2019	ST167	*aac(6')-Ib-cr*	IncF plasmid (132482 bp, NZ_CP048368.1)	LNIE (3)

CF038[56]	surface water	Switzerland	2019	ST410	*aac(6')-Ib-cr*	IncF plasmid (86996 bp, NZ_CP048377.1)	LNIE (3)
TKEC21-17[57]	municipal wastewater	Turkey	2021	ST609	*qnrS1* (type A), *qnrS1* (type L)	*qnrS1* (type A): IncF plasmid (91489 bp, NZ_CP092450.1) *qnrS1* (type L): IncX3 plasmid (46799 bp, NZ_CP092451.1)	SDSE (0)
TKEC21-59[57]	municipal wastewater	Turkey	2021	ST609	*qnrS1* (type A)	IncF plasmid (86914 bp, NZ_CP092453.1)	SDSE (0)
1000C-3[58]	hospital wastewater	USA	2018	ST167	*aac(6')-Ib-cr*	IncF plasmid (145445 bp, NZ_CP091428.1)	LNIE (3)

ᵃDetailed information on each isolate, such as other AMR genes detected, is available in **Table**

S2.

ᵇThe context type is shown for *qnrS1*.

ᶜGyrA codons 83 and 87 and ParC codons 80 and 84 are shown in this order. The number of

amino acid substitutions is shown in parentheses.

FIGURES

Figure 1. The genetic context of *qnrA1*. A red arrow indicates the TMQR determinant, pink

arrows indicate other AMR genes, gray pointed boxes indicate ISs, and black arrows indicate

other genes. Diagrams are not exactly to scale (this also applies to other figures). Inc type,

nucleotide accession number, and country of origin are shown in parentheses next to the

plasmid name. IR: inverted repeat. IRi: inverted repeat at *intI1* end. IRt: inverted repeat at *tni*

end.

Figure 2. Genetic contexts of (a) *qnrB4*, (b) *qnrB7*, and (c) *qnrB19*. Genes and elements are

shown as in **Figure 1**. Nucleotide sequences with >99% nucleotide identity are indicated by

the blue shaded regions in panel (a). Panel (c) is enlarged compared with panels (a) and (b),

and describes the whole plasmid structures. For p32-4_E, a circular structure is also shown to

describe the relationship to pPAB19-1 and the pMB9272_7-like plasmid. The putative *oriT*

sequence in the pMB9272_7-like plasmid was predicted by comparison with well-

characterized plasmids such as ColE1. hp: hypothetical protein.

Figure 3. Structures of plasmids carrying *qnrD1*. Genes and elements are shown as in earlier figures. The light blue shaded regions indicate nucleotide sequences with ~95% nucleotide identity. Annotation of pKFu015_4 was partially performed by sequence comparison with ColE1. IR_L: inverted repeat left. IR_R: inverted repeat right. IR_L and IR_R of $TIME_{IS101}$ are shown at the bottom of the figure, and different nucleotides are marked in red.

Figure 4. Genetic contexts of *qnrS1*. Genes and elements are shown as in earlier figures. The

blue shaded regions indicate nucleotide sequences with >99% nucleotide identity. Note that for

simplicity, only *qnrS1* and directly flanking regions without complex rearrangements (such as

inversions) are shaded in blue (e.g., the left IS*Kpn19* in the type F context shares >99%

nucleotide identity with IS*Kpn19* in the type G context, but this region is not shaded). Tn*6292*

and its derivatives (Tn*6360* and Tn*6361*) are shown for the purpose of comparison. See **Table**

1 for information on genomes carrying each context type.

Figure 5. Genetic contexts of *qnrS2*. Genes and elements are shown as in earlier figures.

JKHS004 and 19M19 carry *qnrS2* on the chromosome, and chromosomal STs are indicated

instead of plasmid types. The blue shaded regions indicate nucleotide sequences with >99%

nucleotide identity.

Figure 6. Genetic contexts of *aac(6′)-Ib-cr*. The contexts are shown separately for plasmids

and chromosomes. *aac(6′)-Ib-cr* is abbreviated as *aac* in the figure. Genes and elements are

shown as in earlier figures. The blue shaded regions indicate nucleotide sequences with >99%

nucleotide identity. Note that for simplicity, only *aac(6′)-Ib-cr* and flanking regions are shaded

in blue (e.g., IS*26* sequences in JKHS004 and 19M19 share >99% nucleotide identity to IS*26*

sequences in C-SRM-3, but these regions are not shaded).

641 **Figure 7**. Genetic contexts of *oqxAB*. Genes and elements are shown as in earlier figures. The

642 blue shaded regions indicate nucleotide sequences with >99% nucleotide identity. Note that for

643 simplicity, only Tn*6010* and flanking regions are shaded in blue (e.g., the left IS*26* in Tn*4352*

644 shares >99% nucleotide identity to other IS*26* sequences, but these regions are not shaded).

647 **Figure 8**. The genetic context of *qepA1*. Genes and elements are shown as in earlier figures.