Pronounced expression of extracellular matrix proteoglycans regulated by Wnt pathway underlies the parallel evolution of lip hypertrophy in East African cichlids

Nagatoshi Machii[1], Ryo Hatashima[1], Tatsuya Niwa[1,2], Hideki Taguchi[1,2], Ismael A. Kimirei[3], Hillary D. J. Mrosso[4], Mitsuto Aibara[1], Tatsuki Nagasawa[1], Masato Nikaido[1,*]

[1]School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan

[2]Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology, Yokohama, Japan

[3]Tanzania Fisheries Research Institute (TAFIRI), Dar es Salaam, Tanzania

[4]Tanzania Fisheries Research Institute (TAFIRI), Mwanza Fisheries Research Center, Mwanza, Tanzania

*Corresponding author: Masato Nikaido

Email: mnikaido@life.isct.ac.jp

Tel: +81-3-5734-2659, Fax: +81-3-5734-02946

Abstract

Cichlid fishes inhabiting the East African Great Lakes, Victoria, Malawi, and

Tanganyika, are textbook examples of parallel evolution, as they have acquired similar

traits independently in each of the three lakes during the process of adaptive radiation.

In particular, "hypertrophied lip" has been highlighted as a prominent example of

parallel evolution. However, the underlying molecular mechanisms remain poorly

understood. In this study, we conducted an integrated comparative analysis between

the hypertrophied and normal lips of cichlids across three lakes based on histology,

proteomics, and transcriptomics. Histological and proteomic analyses revealed that the

hypertrophied lips were characterized by enlargement of the proteoglycan-rich layer, in

which versican and periostin proteins were abundant. Transcriptome analysis revealed

that the expression of extracellular matrix-related genes, including collagens,

glycoproteins and proteoglycans, was higher in hypertrophied lips, regardless of their

phylogenetic relationships. In addition, the genes in Wnt signaling pathway, which is

involved in promoting proteoglycan expression, was highly expressed in both the

juvenile and adult stages of hypertrophied lips. Our comprehensive analyses showed

that hypertrophied lips of the three different phylogenetic origins can be explained by

similar proteomic and transcriptomic profiles, which may provide important clues into

the molecular mechanisms underlying phenotypic parallelisms in East African cichlids.

Introduction

Parallel evolution is evidence of natural selection that drives adaptive solutions to the environment (Losos, 2011). Acquisition of similar traits in distantly related lineages can be observed in multiple lineages, such as the loss of eyes in cave fish, patterns of wing colors in Heliconius butterflies, and body armor of sticklebacks (Krishnan and Rohner, 2017; Stern, 2013). Since it is unlikely that such a complex phenotype was acquired repeatedly by neutral evolution alone, the same direction of natural selection among species is likely to drive the parallel phenotypic change (Losos, 2011; Stern, 2013). Therefore, research on parallel evolution enables us to compare the commonalities and differences in the mechanisms behind the acquisition of novel traits and elucidate how adaptation occurs in similar environments.

East African cichlids offer one of the best opportunities to examine the molecular basis of parallel evolution (Kocher et al., 1993; Muschick et al., 2012). There are hundreds of endemic cichlid species in the East African Great Lakes, Victoria, Malawi, and Tanganyika, as a result of explosive adaptive radiation (Salzburger, 2018; Santos et al., 2023). As a result of independent adaptations in each lake, parallel evolution is often observed among lakes, such as that of the nose snout (Duenser et al., 2023), jaw morphology, and body shape (Kocher et al., 1993; Muschick et al., 2012). To reveal their evolutionary mechanisms, genetic bases of the phenotypes have been investigated. For example, the parallel evolution of body stripes in cichlids of Lake Victoria and Lake Malawi was explained by relatively simple changes in pigmental cell arrangement caused by mutations in the cis-regulatory element of agouti-related peptide 2 (agrp2) (Kratochwil et al., 2018). Although this parallel evolution of body stripe patterns can be explained by a single gene, most traits are likely to be polygenic, hindering elucidation of the parallel evolution mechanism.

62   Cichlid fishes have acquired a huge diversity in their oral morphologies, reflecting ecological and trophic specializations (Albertson and Kocher, 2006; Kocher, 2004; Lecaudey et al., 2021, 2019). Hypertrophied lips, one of the most prominent examples of oral diversity, have been independently acquired in cichlids inhabiting each of the East African Great Lakes, as well as in the Nicaraguan crater lakes and South American rivers (Henning et al., 2017; Kautt et al., 2020; Masonick et al., 2023, 2022). Several hypotheses have been proposed for the adaptive roles of hypertrophied lips, such as for effectively sucking prey out from narrow crevices (Lukas et al., 2015), reducing mechanical stress during foraging in rocky habitats (Fryer, 1972; Greenwood, 1974), and facilitating prey detection by providing an enlarged area for taste receptors (Arnegard and Snoeks, 2001; Oliver and Arnegard, 2010). Thus, hypertrophied lips are considered an adaptive trait for a specific trophic niche, making them an ideal model for understanding the mechanism of parallel evolution.

The molecular mechanisms underlying hypertrophied lips have been investigated in various lineages of cichlids. For example, transcriptome comparisons between hypertrophied and normal lips of cichlids in Lake Tanganyika (Colombo et al., 2013) and South American crater lakes (Manousaki et al., 2013) have identified some differentially expressed genes (DEGs) as candidates for lip hypertrophy; however, their primary regulators remain to be elucidated. In addition, quantitative trait locus (QTL) mapping (Henning et al., 2017) and genome-wide association study (GWAS) (Masonick et al., 2023) have been conducted to elucidate the genomic loci responsible for hypertrophied lips in the cichlids of Lake Victoria and Lake Malawi. These genome-wide studies concluded that lip hypertrophy was a polygenic trait involving several candidate genes. However, the molecular mechanism underlying lip hypertrophy has not yet been elucidated.

In this study, we comprehensively compared the hypertrophied and normal lips

of cichlids across all East African Great Lakes at various biological levels using

histology, proteomics, and transcriptomics. As a result, we showed that a novel key

pathway commonly involved in the formation of hypertrophied lips, providing insight

into a better understanding of the molecular basis of a typical example of parallel

evolution.

Materials and Methods

Animals

Three species of Victoria cichlids (*Haplochromis chilotes* with hypertrophied

lips; *H. sauvagei,* and *H. pyrrhocephalus* with normal lips), four species of Malawi

cichlids (*Placidochromis milomo* with hypertrophied lips; *Maylandia lombardoi, M. zebra,*

and *Labidochromis caeruleus* with normal lips), and two species of Taganyika cichlids

(*Lobochilotes labiatus* with hypertrophied lips; *Tropheus moorii* with normal lips) were

used in the experiments. *H. chilotes, H. sauvagei*, and *H. pyrrhocephalus* were collected

from the Mwanza gulf of Lake Victoria during an exploration that was led by the Nikaido

Laboratory in 2018. The other fish were obtained from a commercial supplier. Fish were

maintained and bred at 27°C on a 12/12 h light/dark cycle. One to 12 individuals were

kept in a plastic tank (40 cm × 25 cm × 36 cm) and fed food pellets 2-3 times a day. The

sample information is provided in Table S1. All experiments were conducted in

accordance with the Institutional Animal Experiment Committee of the Tokyo Institute of

Technology.

Histological observation

Cichlids were anesthetized with 0.02% ethyl p-aminobenzoate. The lower lips

were then dissected, along with the lower jaw, and were fixed in 4% paraformaldehyde

(PFA, Fujifilm-Wako, Chuo-ku, Osaka, Japan) in phosphate-buffered saline (PBS) at

4°C for 24 h. Tissues were demineralized in 0.34 M ethylenediaminetetraacetic acid

(EDTA, Fujifilm-Wako, Chuo-ku, Osaka, Japan) in 100 mM, pH7.0 tris (hydroxymethyl)

aminomethane (Tris, Fujifilm-Wako, Chuo-ku, Osaka, Japan) buffer at 4°C for 24 h.

After demineralization, tissues were treated with 20% sucrose/PBS at 4°C for 24 h.

Tissues were then embedded in Tissue-Tek O.C.T. compound (Sakura Finetek, Chuo-

ku, Tokyo, Japan) and frozen with liquid nitrogen. Embedded blocks were sliced in the

sagittal plane into 20 μm sections and placed on a coated glass slide (MAS-01,

Matsunami, Bunkyo-ku, Tokyo, Japan). Sections were stored at –80°C until they were

used for experiments. To observe the histological structure, hematoxylin–eosin (HE)

staining, Alcian blue (AB) staining, and Van Gieson (VG) staining were conducted.

Mayer's hematoxylin solution (Fujifilm-Wako, Chuo-ku, Osaka, Japan), Eosin Y

solution (Fujifilm-Wako, Chuo-ku, Osaka, Japan), ALCIAN BLUE 8GX (Sigma-Aldrich,

St. Louis, MO, USA), Van Gieson solution P (Fujifilm-Wako, Chuo-ku, Osaka, Japan),

and Van Gieson solution F (Fujifilm-Wako, Chuo-ku, Osaka, Japan) were used for

each staining. Experiments were performed according to the manufacturer's protocol

and previous studies (Arnegard and Snoeks, 2001; Oliver and Arnegard, 2010). For AB

staining, the pH was adjusted to 2.5 with 3% acetic acid (Fujifilm-Wako, Chuo-ku,

Osaka, Japan). For VG staining, Van Gieson solution P and Van Gieson solution F

were prepared at a ratio of 7.55:1. The stained area was calculated using ImageJ

version 1.5.1 (Schneider et al., 2012) after trimming the image to include only the lips.

The images were captured using a Zeiss Axioplan fluorescence microscope and a

Zeiss Axiocam 503 color CCD camera (Carl Zeiss, Oberkochen, Germany). Images

were level-corrected, contrast adjusted, and tiled using Adobe Photoshop CC 2022.

Chondroitinase digestion

Chondroitinase ABC from *Proteus vulgaris* (Sigma-Aldrich, St. Louis, MO, USA)

was used for chondroitinase digestion. 2 mUnits/0.1% bovine serum albumin

solutionBSA in 20 mmol/L Tris-HCl pH 7.7 were prepared as the enzyme solution. In

the experiment, the sections were first treated with 100 μL of 0.1% BSA solution for 5

min. Next, 100 μL of enzyme solution, or BSA solution for the control, was added.

During the reaction, the sections were placed in a moist chamber. The enzymatic

reaction was performed at 37°C for 1 h. After the reaction, the samples were washed

with water for 15 min, and then AB staining was examined.

Identification of extracellular matrix (ECM)-related genes in cichlids

To compare the transcriptome of cichlid lips, cichlid extracellular matrix (ECM)-related genes were identified using the data from a previous study (Hynes and Naba, 2012). The amino acid sequences of human, mouse, and zebrafish ECM-related genes registered in Human GENCODE 41, Mouse GENCODE M30, Ensembl, and UniProtKB were used as queries, and those of *M. zebra* and *O. niloticus* were identified from their protein sequences registered in Ensembl and Uniprot using OrthoFinder2 v.2.5.4. The longest amino acid sequence of each isoform was selected to represent the gene. After Orthfinder2 analysis, ortholog correspondences were corrected manually, if necessary. A list of genes is provided in Table S2.

Sample preparation for proteomic analysis

Proteomic analysis was conducted on the lips of cichlids, *H. chilotes* (hypertrophied) and *H. sauvagei* (normal), from Lake Victoria. The lips of *H. chilotes* were dissected in two half anterior (tip) and half posterior (base), which are analyzed separately. For proteomics analysis of pellet fractions, dissected samples were homogenized with a pestle in 1% sodium dodecyl sulfate in PBS supplemented with a protease inhibitor cocktail (cOmpleteTM mini, EDTA free, Roche, Basel, Switzerland). The resultant lysate was centrifuged at 20,000× g for 15 min, and the supernatant was extracted. A fourfold volume of methanol, an equal volume of chloroform, and a threefold volume of distilled water were added to the supernatants, and the suspension was vortexed vigorously for 1 min after each addition. After vortexing, the supernatants were centrifuged at 15,000× g for 2 min and the upper water layer was removed. A threefold volume of methanol was added, and the samples were

centrifuged at 15,000× g for 2 min. The supernatant was removed by aspiration and

the pellet fraction was obtained.

Subsequently, sample preparation for LC−MS/MS was performed according to

a previous study (Masuda et al. 2009), with some modifications. Proteins in the pellet

fraction were solubilized in phase transfer surfactant (PTS) solution (12 mM sodium

deoxycholate, 12 mM sodium lauryl sulfate, 100 mM Tris-HCl, pH 9.0), and the protein

concentration was determined using a Pierce BCA Protein Assay Kit (Thermo Fisher

Scientific, Waltham, MA). Then, 50 μg of protein in 50 μL of PTS was prepared for the

following procedure. First, the proteins were reduced by treatment with 5 mM

dithiothreitol (DTT) at room temperature for 30 min, and then alkylated with 50 mM

iodoacetamide in the dark at room temperature for 30 min. The reaction was stopped

by a fivefold dilution with 50 mM ammonium bicarbonate. Denatured proteins in the

solution were digested into peptide fragments in two steps: first, incubation with 0.5 μg

of Trypsin/Lys-C Mix (Promega, Madison, WI) at room temperature for 3 h, followed by

incubation with 1.0 μg of Trypsin/Lys-C Mix at 37°C overnight. After digestion, an equal

volume of ethyl acetate and 0.5% trifluoroacetic acid (final concentration) were added

and the mixture was shaken vigorously for 2 min. The mixture was then centrifuged at

15,700 × g for 2 min and the upper ethyl acetate layer was removed. The solvent was

dried using a centrifugal evaporator. The residual pellet was redissolved in 600 μL of

0.1% TFA and 2% acetonitrile and then desalted using a handmade StageTip

composed of an SDB-XC Empore Disk (3 M, Maplewood, MN, USA). The solution was

applied to a StageTip equilibrated with 0.1% TFA and 2% acetonitrile, washed with

0.1% TFA and 2% acetonitrile, and eluted with 0.1% TFA and 80% acetonitrile. After

the desalting, the solvent was dried using a centrifugal evaporator, and the residual

peptides were redissolved in 120 μL of 0.1% TFA and 2% acetonitrile. The solution

was centrifuged at 20,000 × g for 5 min, and the supernatant was collected and

subjected to LC–MS/MS measurement.

LC–MS/MS measurement and data processing

LC–MS/MS measurements and SWATH-MS acquisition (Gillet et al., 2012)

were performed using an Eksigent NanoLC Ultra and TripleTOF 4600 tandem-mass

spectrometer or an Eksigent nanoLC 415 and TripleTOF 6600 mass spectrometer (AB

Sciex, USA). The trap column used for nanoLC was a 5.0 mm × 0.3 mm ODS column

(L-column2, CERI, Bunkyo-ku, Tokyo, Japan), and the separation column was a 12.5

cm × 75 µm capillary column packed with 3 µm C18- silica particles (Nikkyo Technos,

Bunkyo-ku, Tokyo, Japan). The flow rate of nanoLC was 300 nL/min, and peptides

were separated by applying a 10-40% linear acetonitrile gradient over 70 min in the

presence of 0.1% formic acid. Data were acquired in DIA/SAWTH (Gillet et al., 2012)

mode controlled by Analyst software (version 1.7, AB Sciex, USA) with the following

settings: 15 Da fixed width window in 350–1100 m/z (50 cycles), 45 msec acquisition

time, and 100–1600 fragment ion m/z range. The measurements were performed three

times for each sample. Quantitative values were calculated by DIA-NN software

(version 1.8.1) with default settings (Demichev et al., 2020). The library for

DIA/SWATH quantification was generated using DIA-NN with the amino acid

sequences of *M. zebra* total proteins obtained from UniProtKB. The P-value and fold

change were calculated by Python libraries (scipy.stats and pandas) using the proteins

detected in all three measurements for three parts of the lips. Quantitative values were

scaled by log10 before statistical tests (Welch's t-test, two sided).

RNA-seq analysis

RNA was extracted from dissected lower lips using TRI Reagent (Molecular

Research Center, Inc.). Three species of Lake Victoria cichlids (*Haplochromis chilotes*

[n = 4]*, H. sauvagei* [n = 4]*, H. pyrrhocephalus* [n = 2]), three species of Lake Malawi

cichlids (*Placidochromis milomo* [n = 5], *Maylandia zebra* [n = 2]*, Labidochromis*

caeruleus [n = 5]), and two species of Lake Taganyika cichlids (*Lobochilotes labiatus*

[n = 4]*, Tropheus moorii* [n = 4]) were used for RNA-seq (Table S1). The extracted total

RNA was sequenced at 100□bp paired-end reads on a Illumina NovaSeq 6000 by

Macrogen Japan Corp., using a TruSeq stranded mRNA Library Prep Kit (Illumina).

The raw FASTQ data were qualified using fastp version 0.20.1 (Chen et al., 2018) in

default option. The high-quality reads were mapped to *M. zebra* genome (Conte and

Kocher, 2015) using STAR version 2.7.10a (Dobin et al., 2013) and quantified using

rsem-calculate-expression version 1.3.1 (Li and Dewey, 2011). Principal component

analysis (PCA) was conducted using Python scikit-learn package. Low-expression

genes (average transcripts per kilobase million (TPM) below 5.0) were removed, and

expression rates were normalized by Z score. The resulting 12,231 genes were used

for PCA analysis. For PCA of ECM-related genes, we used 186 genes as inputs, which

were extracted in the same manner. DEGs were detected using the TCC package

(Sun et al., 2013) with the TMM-edgeR-edgeR combination. Enrichment analysis was

conducted using DAVID version 2023q3 (Huang et al., 2009a, 2009b) after the gene

IDs of *M. zebra* were converted to human gene IDs using ensembl biomart. The

annotation terms used for functional annotation clustering were collected from

biological process (Gene Ontology), KEGG, and Reactome.

Results

Comparative histology of hypertrophied and normal lips among cichlids from the

East African Great Lakes

We compared the histological structures of hypertrophied and normal lips

among cichlids from Lake Victoria, Lake Malawi, and Lake Tanganyika. For an

accurate comparison, we first observed serial sections of the lower lips with the jaws to

determine the midline of the lips in each species, which was defined by the shape of

the lower dentary bone (Figure S1). Based on this definition, we compared the

distribution of collagens and proteoglycans, the main ECM components, at the midline

of the lower lips using VG stain (red) and AB stain (blue). Hypertrophied lips of cichlids

from all lakes generally consisted of two mutually exclusive structures in the

hypodermis: a collagen-rich layer at the base and a proteoglycan-rich layer at the tip.

Moreover, comparison of hypertrophied lips and normal lips showed that the area of

the proteoglycan-rich (AB positive) layer was notably enlarged in hypertrophied lips,

which was statistically significant in cichlids from Lake Victoria (Figure 1; Figure S2).

Proteoglycans and their related proteins were abundant in the hypertrophied

lips

To explore the proteins associated with lip hypertrophy, we used shotgun

proteomics to compare the protein profiles of hypertrophied lips and normal lips in

Lake Victoria cichlids. Using LC–MS/MS in DIA/SWATH acquisition mode, we detected

2,720 proteins in total from three types of lip samples: the tip of hypertrophied lips

(proteoglycan-rich loose connective tissue), the base of hypertrophied lips (collagen-

rich loose connective tissue), and normal lips. First, we observed few differences

between the tip of hypertrophied lips and the base of the hypertrophied lips (Figure S3).

On the other hand, we identified 138 differentially accumulated proteins between the

tip of hypertrophied and normal lips (133 proteins upregulated and 5 proteins

downregulated in the hypertrophied lip) under the criteria (P-value < 0.05; log2 fold

change > 1) (Figure 2A). Compared to normal lips, tip of hypertrophied lips is abundant

in proteoglycans and their related proteins such as versican a/b (vcan), brevican

(bcan), periostin (postn), hyaluronan and proteoglycan link protein 1 (hapln1), and

hyaluronan binding protein 2 (habp2), which were highly accumulated in hypertrophied

lips with statistical significance (Figures 2B-D). The results of the above proteomic

analyses were consistent with the histological observation that proteoglycan-rich loose

connective tissue was enlarged in hypertrophied lips compared to that in normal lips.

Chondroitin sulfate proteoglycans such as vcan are the main component of

hypertrophied lips

Next, we performed glycosaminoglycan digestion experiments to examine

whether loose connective tissue was composed of the proteoglycans, which were

detected as highly accumulated proteins in hypertrophied lips. Among the

proteoglycans, we focused on vcan, which showed a clear difference in protein amount

between hypertrophied and normal lips. Vcan is a high-molecular-weight core protein

(approximately 370 kDa) that is glycosylated with chondroitin sulfate (Iozzo and

Schaefer, 2015; Wight et al., 2020). Therefore, we stained sections of hypertrophied

lips with AB stain after the treatment with chondroitinase, which specifically digests

glycosylated chondroitin sulfate. As a result, AB staining in loose connective tissue

became drastically weak in treated sections, which was common among lakes

(Figures 2E-G). Our data suggested that the main components of enlarged loose

connective tissue in hypertrophied lips are chondroitin sulfate proteoglycans such as

vcan.

Convergent expression in ECM-related genes

Given that proteomic analysis is not well suited for detecting high-turnover proteins (Decaris et al., 2014), we additionally conducted transcriptomic comparisons between hypertrophied and normal lips across Lake Victoria, Lake Malawi and Lake Tanganyika. PCA with all expressed genes clustered according to each lake, reflecting genetic closeness (Figure 3A). Next, we compared the gene expression of ECM-related genes, for which the proteomics profiles were clearly distinct between hypertrophied and normal lips. A PCA plot was drawn using only ECM-related genes in cichlids, which were identified based on a previous study (Hynes and Naba, 2012) (Figure 3B). In contrast to the PCA plot of all expressed genes, that of ECM-related genes reflected differences of lip morphology (hypertrophied and normal), regardless of the genetic closeness. To evaluate the results of these PCAs, we calculated the average silhouette (AS) score, which assesses the cohesiveness of the cluster. In the PCA of all genes, the AS score of the cluster by lake (0.1601) was higher than that of the cluster by lip morphology (0.0849). In contrast, in the PCA of ECM-related genes, the AS score of the cluster by lip morphology (0.1798) was higher than that of the cluster by lake (0.0774). These results indicate that hypertrophied lips are characterized by the expression of ECM-related genes.

Next, we compared the expression of each ECM-related gene between hypertrophied and normal lips of cichlids from the three lakes (Figures 3C, D). As a result, several ECM-related genes showed higher expression in hypertrophied lips than in normal lips, such as type-I collagen fibrils (col1a1a, col1a1b, and col1a2), hyaluronan-binding proteoglycans (vcanb, hapln1a/b, and hapln3), periostins (postna/b), small leucine-rich proteoglycans (dcn, lum, and aspn), and microfibril-

related genes (mfap2, emilin1b, emilin2a, emilin2b, and emilin3b). Consistent with the

proteomics comparison, some proteoglycans and their related proteins (vcanb,

hapln1a, and postna) showed significantly higher expression in hypertrophied lips than

in normal lips. In conclusion, we observed convergent expression of ECM-related

genes in hypertrophied lips of cichlids among the East African Great Lakes.

High expression of ECM-related genes in the juvenile and adult stages of

hypertrophied lips

Next, we performed transcriptome analysis by comparing juvenile and adult

cichlids from Lake Victoria to examine the possible expression changes in ECM-

related genes during development (Figure 4A). In the adult stage, the expression

profiles of ECM-related genes between hypertrophied and normal lips showed obvious

differences (low correlation value 0.75; orange square in Figure 4B), which was

consistent with previous analyses (Figure 3B). In the juvenile stage, the difference was

weaker than in adult stage (0.85; blue square in Figure 4B). Interestingly, the

difference in the expression profiles between the juvenile and adult stages of

hypertrophied lips (0.90; Figure 4B black square) was lower than that of normal lips

(0.83; Figure 4B black dotted square). To examine the genes contributing to the

differences in expression profiles, we performed hierarchical clustering analysis for the

relative expression levels of each ECM-related gene (Figure 4C, D). The analyses

consistently showed the higher expression of proteoglycans and their related genes

(e.g. postna, vcanb, and hapln1b) in hypertrophied lips at both the juvenile and adult

stages. These results indicated that the accumulation of proteoglycans, which leads to

the lip hypertrophy, begins at the juvenile stage.

The Wnt pathway-related genes are highly expressed in hypertrophied lips

To explore the molecular pathways involved in lip hypertrophy, we performed

an enrichment analysis using DAVID (Huang et al., 2009a, 2009b) against DEGs

obtained from the transcriptome. As a result of enrichment analysis, DEGs were

categorized in the canonical and noncanonical Wnt signaling pathways, suggesting

that high expression of genes in the Wnt signaling pathway is likely to be involved in

the hypertrophied lips of juvenile and adult fish. Comparison of the developmental

stages revealed differences in the expression patterns between the canonical and

noncanonical Wnt pathways: the genes of the canonical Wnt pathway were highly

expressed in both stages (Figures 5C–I), whereas those of the non-canonical pathway

were highly expressed only in the adult stage (Figures 5J–N). We further compared

the expression patterns of these Wnt signaling pathways in cichlids from Lake Malawi

and Lake Tanganyika. Notably, the expression of genes for the Wnt pathway tend to be

higher in the hypertrophied-lip cichlids in Lake Malawi and Lake Tanganyika, although

the differences were not as distinct as those found in cichlids from Lake Victoria.

These results were consistent with those of previous studies that showed that the

expression of ECM-related genes, such as vcan, was promoted by the Wnt signaling

pathway (Rahmani et al., 2005). Continuous activation of the Wnt signaling pathway

throughout all developmental stages may lead to lip hypertrophy via the deposition of

ECM-related proteins.

Discussion

In this study, we investigated the differences between hypertrophied and normal lips using histology, proteomics, and transcriptomics. Histological comparisons revealed that cichlids with hypertrophied lips from all three East African Great Lakes exhibited an enlargement of the proteoglycan-rich loose connective tissue. By integrating proteomics and transcriptomics, we identified several proteins that were significantly accumulated in the hypertrophied lips. Among them, proteoglycans and their related proteins such as vcan and postn, could be a primary factor that explains lip hypertrophy. Interestingly, these proteins are known to be involved in the formation of human keloids, a skin disease characterized by an overabundance of ECM in the dermis (Deng et al., 2021; Naitoh et al., 2005). It has been discussed that excessive deposition of vcan retains large amounts of water in the sulfate or carbonyl moieties, contributing to the increase in keloid volume (Yagi et al., 2013). In addition, postn is involved in fibrosis through its ability to interact with multiple ECM proteins. For example, upregulation of postn leads to strengthening of the ECM by crosslinking with collagens and excessive deposition of ECM proteins by differentiating quiescent fibroblasts into activated myofibroblasts (Ashley et al., 2017; Yamaguchi et al., 2013). In cichlid, vcan and postn may have a similar function, leading to hypertrophied loose connective tissue.

The Wnt signaling pathway was also elevated in both the adult and juvenile stages of hypertrophied lips. The Wnt signaling pathway is known to be a key regulator of the expression of ECM-related genes; for example, vcan is directly regulated by the Wnt/b-catenin and tcf4 signaling pathways, leading to vcan upregulation (Rahmani et al., 2005). Additionally, Wnt5a, a noncanonical pathway, plays a critical role in dermal condensation during mouse and sugar glider development (Feigin et al., 2023). Cellular-level localization of gene expression for the canonical and non-canonical Wnt

pathway is required for future examination of the involvement of the Wnt signaling

pathway in lip hypertrophy.

From a genomic perspective, several studies have investigated the genetic

basis of hypertrophied lip cichlids (Masonick et al., 2023; Nakamura et al., 2021).

Importantly, some Wnt pathway-related genes (tcf4 and daam2) and ECM-related

genes (postna, col12a1a, and col12a1b) have been found to be under positive

selection in cichlids with hypertrophied lips of Lake Victoria (see Nakamura et al., 2021

Table S3). For future research, examining whether these genes are under selection in

other lakes is crucial to understand the genetic mechanisms underlying the parallel

evolution of hypertrophied lips.

Although the parallel evolution of hypertrophied lips in cichlids has received

considerable attention in evolutionary biology, the commonality and diversity

underlying their molecular basis remain poorly understood. This might be primarily due

to the fact that many genetic factors are possibly involved in the dynamic organization

of hypertrophied lips. We uncovered the apparent relationships between hypertrophied

lips and the expression profiles of ECM proteins, in particularly proteoglycans. The

trends for the overall expression of ECM-related genes were similar across

hypertrophied lip species, but we rarely observed a specific gene that was commonly

expressed at high or low levels in all three examples of hypertrophied lips across all

East African Great Lakes. Furthermore, although we focused primarily on the

relationship between the Wnt signaling pathway and lip hypertrophy, several other

pathways may be involved in the development of hypertrophied lips. For example, our

enrichment analysis in adult cichlids identified VEGF-related pathways, which could

contribute to lip hypertrophy by increasing vascularization and nutrient supply to the lip

tissue. In addition, previous quantitative trait locus (QTL) analysis by Henning et al.

(2017) concluded that lip hypertrophy is likely influenced by numerous loci with small

additive effects. These lines of data imply that although enlargement of proteoglycan-

rich loose connective tissue is common in hypertrophied lips, the developmental

pathways to accomplish this are diverse in each lake. The pathways responsible for lip

hypertrophy therefore require further investigation. In conclusion, our approach of

integrating histology, proteomics and transcriptomics revealed complex evolutionary

pathways for hypertrophied lips in cichlids of the East African Great Lakes, providing

an opportunity to further understand the molecular mechanism underlying the parallel

evolution of polygenic traits, such as jaw morphology or body shape.

425
Fig.1 Histological comparisons of hypertrophied and normal lips of cichlids among lakes, focusing on proteoglycan-rich loose connective tissue (A,B). Examples of typical cichlids with hypertrophied lips (A, *Haplochromis chilotes*) and normal lips (B, *H. sauvagei*). The lower jaw and lip are highlighted by black squares. (C) Phylogenetic relationships among cichlids from Lake Victoria, Lake Malawi, and Lake Tanganyika. (D–O) The midline of the lips, based on the lower dentary bone of the lower jaw (marked by asterisks), in cichlids from Lake Victoria (D,E, *H. chilotes* with

hypertrophied lips; F,G, *H. sauvagei* with normal lips), Lake Malawi (H,I,

Placidochromis milomo with hypertrophied lips; J,K, *Maylandia lombardoi* with normal

lips), and Lake Tanganyika (L,M, *Lobochilotes labiatus* with hypertrophied lips; N,O,

Tropheus moorii with normal lips) were sectioned in sagittal, and stained with AB:

Alcian blue (D, H, L, F, J, and N), or VG: Van Gieson (E, I, M, F, J, and N). Each

section is shown with the dorsal side up and the anterior side to the right. The area

surrounded by dotted lines shows proteoglycan-rich tissue (stained dense blue by AB)

or collagen-rich tissue (stained dense red by VG). Each species showed mutually

exclusive staining by AB and VG, and AB-positive (proteoglycan-rich) tissues were

commonly enlarged in hypertrophied lips among the cichlids from all three lakes with

statistical significance. Scale bars are 1mm.

444
Fig.2 Chondroitin sulfate proteoglycans accumulate in hypertrophied lips. (A) Volcano

plot of proteomics comparison between the tips of hypertrophied and normal lips.

Statistical significance was examined using Welch's t-test. Yellow plots highlight highly

accumulated proteins in hypertrophied lips, and blue plots highlight highly accumulated

proteins in normal lips with the criteria of P-value < 0.05 & log2 fold change > 1. (B,C)

The top 20 proteins highly accumulated in normal (B) and hypertrophied lips (C) are

shown along with log2 fold change relative to normal lips. (D) The relative amount of

each proteoglycan-related protein. MS2 intensity indicates the intensity of peptides

obtained from LC–MS/MS and DIA/SWATH analysis using DIA-NN software by

comparing three parts of the lip: the base of hypertrophied lips (HLip), the tip of HLip,

and normal lips (NLip). Welch's t-test was used to examine statistical significance. (E–

G) Digestion of chondroitin sulfate by treatment with (upper panels) and without (lower

panels) chondroitinase in hypertrophied lips of cichlids from Lake Victoria (E), Lake

Malawi (F), and Lake Tanganyika (G). Each section is shown with the dorsal side up

and the anterior side to the right. Scale bars are 1mm.

460
Fig.3 Hypertrophied lips of cichlids are characterized by higher expression of ECM-

related genes. Transcriptomic analysis using three species of Lake Victoria cichlids (*H.*

chilotes (Hch) with hypertrophied lips; *H. sauvagei* (Hsa) and *H. pyrrhocephalus* (Hpy)

with normal lips), three species of Lake Malawi cichlids (*Placidochromis milomo* (Pmi)

with hypertrophied lips; *Maylandia zebra* (Mze) and *Labidochromis caeruleus* (Lca)

with normal lips), and two species of Lake Tanganyika cichlids (*Lobochilotes labiatus*

(Lla) with hypertrophied lips; *Tropheus moorii* (Tmo) with normal lips). (A, B) PCA plots

of all expressed genes showed differences depending on the lakes (A), while those of

ECM-related genes showed differences depending on lip morphology: hypertrophied

or normal (B). (C) Heatmap of ECM-related gene expression between hypertrophied

and normal lips. The expression rate (TPM) normalized by Z score of each gene is

compared for Lake Victoria (yellow; Hch, Has and Hpy), Lake Malawi (blue; Pmi, Lca

and Mze), and Lake Tanganyika (red; Lla and Tmo). The intensity of the relative

expression rate is shown in the bottom right of the figure. (D) Cumulative fold change

of the ECM-related genes comparing hypertrophied and normal lips. The gene symbol

corresponds to Figure 3C. The color represents the lake information. The gray dotted

line represents the value at which higher expression was observed in the

hypertrophied lips.

480
Fig.4 Comparison of juvenile and adult cichlids to examine gene expression changes

during the development of hypertrophied lips. (A) The lips of the cichlids used in this

analysis are shown with the dorsal side up and the anterior side to the right: the

juvenile and adult stages of *Haplochromis chilotes* (Hch, representative of

hypertrophied lips: HLip) and *H. sauvagei* (Hsa, representative of normal lips: NLip)

from Lake Victoria. Hypertrophied lips at juvenile stage are not notably enlarged. Scale

bars are 0.5 mm. (B) Spearman correlation matrix of the ECM-related gene expression

rate (TPM) normalized by Z score. The raw correlation values are indicated on the

right side and the average correlation values within each stage and species are

indicated on the left side. The orange frame highlights the correlations between the

hypertrophied and normal lips at the adult stage, and the blue frame highlights those at

the juvenile stage. The black frame highlights the correlations between juvenile and

adult samples of each species. (C) Cluster heatmap of the ECM-related gene

expression rate (TPM) normalized by Z score in each gene. Each gene is clustered by

similarity of expression rates. The intensity of the relative expression rate is shown on

the bottom left of the figure. (D) The log10 cumulative fold change of the ECM-related

genes comparing hypertrophied and normal lips. The gene symbol corresponds to

Figure 4C. The positive values indicate high expression in hypertrophied lips, whereas

the negative values indicate low expression in hypertrophied lips. The colors of the

bars indicate the stage of the samples.

502
Fig.5 Wnt pathway-related genes are highly expressed in hypertrophied lips of juvenile

and adult cichlids. (A, B) Enrichment analysis of highly expressed genes in the

hypertrophied lips. Labels are according to the top annotation in DAVID annotation

clusters. Asterisks indicate statistically significant annotation clusters (FDR < 0.05).

The Wnt-related annotations are highlighted in yellow. (C–N) Expression of the specific

Wnt pathway-related genes. (C, J) The size of the dots represents the log fold change,

and the color of the dots represents the P-value. Blue dots represent genes that are

highly expressed in hypertrophied lips (HLip), and red dots represent genes that are

highly expressed in normal lips (NLip). (D–I, K–N) Box plots show the expression rate

(TPM) of Wnt pathway-related genes. Yellow boxes represent genes that are highly

expressed in hypertrophied lips and blue boxes represent genes that are highly

expressed in normal lips, respectively. Mann–Whitney U-test was used to examine the

statistical significance (*: P-value < 0.05). (C–I) The canonical Wnt pathway-related

genes, and (J–N) show the non-canonical Wnt pathway-related genes.

518
Fig.6 The trend of higher expression of Wnt-related genes in hypertrophied lips. (A)

Box plots of expression rate (TPM) of Wnt related genes of hypertrophied lips (HLip:

yellow) and normal lips (NLip: blue). Mann–Whitney U-test was used to examine the

statistical significance (*: P-value < 0.05, **: P-value < 0.01). (B) Illustration of the

canonical Wnt pathway. DEGs are marked in the bold.

Albertson RC, Kocher TD. 2006. Genetic and developmental basis of cichlid trophic
diversity. *Heredity* **97**:211–221.
Arnegard ME, Snoeks J. 2001. New Three-Spotted Cichlid Species with Hypertrophied
Lips (Teleostei: Cichlidae) from the Deep Waters of Lake Malaŵi/Nyasa, Africa.
The American Society of Ichthyologists and Herpetologists. doi:10.1643/0045-
8511(2001)001[0705:NTSCSW]2.0.CO;2
Ashley SL, Wilke CA, Kim KK, Moore BB. 2017. Periostin regulates fibrocyte function
to promote myofibroblast differentiation and lung fibrosis. *Mucosal Immunol*
10:341–351.
Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**:i884–i890.
Colombo M, Diepeveen ET, Muschick M, Santos ME, Indermaur A, Boileau N,
Barluenga M, Salzburger W. 2013. The ecological and genetic basis of convergent
thick-lipped phenotypes in cichlid fishes. *Mol Ecol* **22**:670–684.
Conte MA, Kocher TD. 2015. An improved genome reference for the African cichlid,
Metriaclima zebra. *BMC Genomics* **16**:724.
Decaris ML, Gatmaitan M, FlorCruz S, Luo F, Li K, Holmes WE, Hellerstein MK, Turner
SM, Emson CL. 2014. Proteomic analysis of altered extracellular matrix turnover
in bleomycin-induced pulmonary fibrosis. *Mol Cell Proteomics* **13**:1741–1752.
Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M. 2020. DIA-NN: neural
networks and interference correction enable deep proteome coverage in high
throughput. *Nat Methods* **17**:41–44.
Deng C-C, Hu Y-F, Zhu D-H, Cheng Q, Gu J-J, Feng Q-L, Zhang L-X, Xu Y-P, Wang D,
Rong Z, Yang B. 2021. Single-cell RNA-seq reveals fibroblast heterogeneity and
increased mesenchymal fibroblasts in human fibrotic skin diseases. *Nat Commun*
12:3709.
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29:15–21.
Duenser A, Singh P, Lecaudey LA, Sturmbauer C, Albertson RC, Gessl W, Ahi EP.
2023. Conserved Molecular Players Involved in Human Nose Morphogenesis
Underlie Evolution of the Exaggerated Snout Phenotype in Cichlids. *Genome Biol*
Evol **15**. doi:10.1093/gbe/evad045
Feigin CY, Moreno JA, Ramos R, Mereby SA, Alivisatos A, Wang W, van Amerongen R,
Camacho J, Rasweiler JJ 4th, Behringer RR, Ostrow B, Plikus MV, Mallarino R.
2023. Convergent deployment of ancestral functions during the evolution of
mammalian flight membranes. *Sci Adv* **9**:eade7511.
Fryer G. 1972. The cichlid fishes of the Great Lakes of Africa. *Their Biology and*
Evolution **641**.
Gillet LC, Navarro P, Tate S, Röst H, Selevsek N, Reiter L, Bonner R, Aebersold R.
2012. Targeted Data Extraction of the MS/MS Spectra Generated by Data-
independent Acquisition: A New Concept for Consistent and Accurate Proteome
Analysis*. *Mol Cell Proteomics* **11**:O111.016717.
Greenwood PH. 1974. Cichlid fishes of Lake Victoria, East Africa: the biology and
evolution of a species flock. *Bull Brit Mus (Nat Hist) Zool Suppl* **6**:1–134.
Henning F, Machado-Schiaffino G, Baumgarten L, Meyer A. 2017. Genetic dissection
of adaptive form and function in rapidly speciating cichlid fishes. *Evolution*
71:1297–1312.
Huang DW, Sherman BT, Lempicki RA. 2009a. Bioinformatics enrichment tools: paths
toward the comprehensive functional analysis of large gene lists. *Nucleic Acids*
Res **37**:1–13.

Huang DW, Sherman BT, Lempicki RA. 2009b. Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**:44–57.
Hynes RO, Naba A. 2012. Overview of the matrisome--an inventory of extracellular
matrix constituents and functions. *Cold Spring Harb Perspect Biol* **4**:a004903.
Iozzo RV, Schaefer L. 2015. Proteoglycan form and function: A comprehensive
nomenclature of proteoglycans. *Matrix Biol* **42**:11–55.
Kautt AF, Kratochwil CF, Nater A, Machado-Schiaffino G, Olave M, Henning F, Torres-
Dowdall J, Härer A, Hulsey CD, Franchini P, Pippel M, Myers EW, Meyer A. 2020.
Contrasting signatures of genomic divergence during sympatric speciation. *Nature*
588:106–111.
Kocher T, Conroy JA, McKaye KR, Richard SJ. 1993. Similar Morphologies of Cichlid
Fish in Lakes Tanganyika and Malawi Are Due to Convergence.
Kocher TD. 2004. Adaptive evolution and explosive speciation: the cichlid fish model.
Nat Rev Genet **5**:288–298.
Kratochwil CF, Liang Y, Gerwin J, Woltering JM, Urban S, Henning F, Machado-
Schiaffino G, Hulsey CD, Meyer A. 2018. Agouti-related peptide 2 facilitates
convergent evolution of stripe patterns across cichlid fish radiations. *Science*
362:457–460.
Krishnan J, Rohner N. 2017. Cavefish and the basis for eye loss. *Philos Trans R Soc*
Lond B Biol Sci **372**. doi:10.1098/rstb.2015.0487
Lecaudey LA, Singh P, Sturmbauer C, Duenser A, Gessl W, Ahi EP. 2021.
Transcriptomics unravels molecular players shaping dorsal lip hypertrophy in the
vacuum cleaner cichlid, Gnathochromis permaxillaris. *BMC Genomics* **22**:506.
Lecaudey LA, Sturmbauer C, Singh P, Ahi EP. 2019. Molecular mechanisms underlying
nuchal hump formation in dolphin cichlid, Cyrtocara moorii. *Sci Rep* **9**:20296.
Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* **12**:323.
Losos JB. 2011. Convergence, adaptation, and constraint. *Evolution* **65**:1827–1840.
Lukas B, Gonzalo M-S, Frederico H, Axel M. 2015. What big lips are good for: on the
adaptive function of repeatedly evolved hypertrophied lips of cichlid fishes. *Biol J*
Linn Soc Lond **115**:448–455.
Manousaki T, Hull PM, Kusche H, Machado-Schiaffino G, Franchini P, Harrod C, Elmer
KR, Meyer A. 2013. Parsing parallel evolution: ecological divergence and
differential gene expression in the adaptive radiations of thick-lipped Midas cichlid
fishes from Nicaragua. *Mol Ecol* **22**:650–669.
Masonick P, Meyer A, Darrin Hulsey C. 2023. A Kiss of Deep Homology: Partial
Convergence in the Genomic Basis of Hypertrophied Lips in Cichlid Fish and
Human Cleft Lip. *Genome Biol Evol* **15**. doi:10.1093/gbe/evad072
Masonick P, Meyer A, Hulsey CD. 2022. Phylogenomic Analyses Show Repeated
Evolution of Hypertrophied Lips Among Lake Malawi Cichlid Fishes. *Genome Biol*
Evol **14**. doi:10.1093/gbe/evac051
Muschick M, Indermaur A, Salzburger W. 2012. Convergent evolution within an
adaptive radiation of cichlid fishes. *Curr Biol* **22**:2362–2368.
Naitoh M, Kubota H, Ikeda M, Tanaka T, Shirane H, Suzuki S, Nagata K. 2005. Gene
expression in human keloids is altered from dermal to chondrocytic and
osteogenic lineage. *Genes Cells* **10**:1081–1091.
Nakamura H, Aibara M, Kajitani R, Mrosso HDJ, Mzighani SI, Toyoda A, Itoh T, Okada
N, Nikaido M. 2021. Genomic Signatures for Species-Specific Adaptation in Lake
Victoria Cichlids Derived from Large-Scale Standing Genetic Variation. *Mol Biol*
Evol **38**:3111–3125.

Okuda S, Watanabe Y, Moriya Y, Kawano S, Yamamoto T, Matsumoto M, Takami T,
Kobayashi D, Araki N, Yoshizawa AC, Tabata T, Sugiyama N, Goto S, Ishihama Y.
2017. jPOSTrepo: an international standard data repository for proteomes. *Nucleic*
Acids Res **45**:D1107–D1111.
Oliver MK, Arnegard ME. 2010. A new genus for Melanochromis labrosus, a
problematic Lake Malawi cichlid with hypertrophied lips (Teleostei: Cichlidae).
Ichthyol Explor Freshwaters **21**:209–232.
Rahmani M, Read JT, Carthy JM, McDonald PC, Wong BW, Esfandiarei M, Si X, Luo Z,
Luo H, Rennie PS, McManus BM. 2005. Regulation of the Versican Promoter by
the β-Catenin-T-cell Factor Complex in Vascular Smooth Muscle Cells*. *J Biol*
Chem **280**:13019–13028.
Salzburger W. 2018. Understanding explosive diversification through cichlid fish
genomics. *Nat Rev Genet* **19**:705–717.
Santos ME, Lopes JF, Kratochwil CF. 2023. East African cichlid fishes. *Evodevo* **14**:1.
Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* **9**:671–675.
Stern DL. 2013. The genetic causes of convergent evolution. *Nat Rev Genet* **14**:751–
764.
Sun J, Nishiyama T, Shimizu K, Kadota K. 2013. TCC: an R package for comparing tag
count data with robust normalization strategies. *BMC Bioinformatics* **14**:219.
Wight TN, Kang I, Evanko SP, Harten IA, Chang MY, Pearce OMT, Allen CE, Frevert
CW. 2020. Versican-A Critical Extracellular Matrix Regulator of Immunity and
Inflammation. *Front Immunol* **11**:512.
Yagi Y, Muroga E, Naitoh M, Isogai Z, Matsui S, Ikehara S, Suzuki S, Miyachi Y, Utani
A. 2013. An ex vivo model employing keloid-derived cell-seeded collagen sponges
for therapy development. *J Invest Dermatol* **133**:386–393.
Yamaguchi Y, Ono J, Masuoka M, Ohta S, Izuhara K, Ikezawa Z, Aihara M, Takahashi
K. 2013. Serum periostin levels are correlated with progressive skin sclerosis in
patients with systemic sclerosis. *Br J Dermatol* **168**:717–725.
655

Fig S1 The midlines of the lips were defined by the shape of the lower dentary bone.

Hematoxylin–eosin stain (HE stain) was used to examine serial sections of the lower

lips and lower jaws of cichlids from Lake Victoria (A, *H. chilotes* with hypertrophied lips;

B, *H. sauvagei* with normal lips), Lake Malawi (C, *Placidochromis milomo* with

hypertrophied lips; D, *Maylandia lombardoi* with normal lips), and Lake Tanganyika (E,

Lobochilotes labiatus with hypertrophied lips; F, *Tropheus moorii* with normal lips).

Each section is shown with the dorsal side up and the anterior side to the left. The

lower dentary bone became smaller as it approached the midline, and just at the

midline, the bone was composed of cartilage. Scale bars are 1mm.

Fig. S2 The relative area of the proteoglycan-rich layer was significantly enlarged in

the hypertrophied lips of Victoria cichlids. The stained area percentage of alcian blue

stain (AB stain; A) and Van Gieson stain (VG stain; B) relative to the whole-lip area

was calculated using ImageJ software. *H. chilotes* with hypertrophied lips (yellow box)

and *H. sauvagei* with normal lips (blue box) were used in the experiment. Each plot is

an average value calculated by three sections near the midline of the lips. Mann–

Whitney U-test was used to examine the statistical significance (*: P-value < 0.05, **:

P-value < 0.01).

677
Fig. S3 Volcano plot of proteomics comparison between the tip of hypertrophied lips

and the base of hypertrophied lips (A) and the base of hypertrophied lips and normal

lips (B). Statistical significance was examined by Welch's t-test. Differentially

accumulated proteins in each comparison are highlighted in the black plot with the

following criteria: P-value < 0.05, log2 fold change > 1.

Table. S1 Basic information of the cichlids used in this study.

Table. S2 List of ECM-related genes of cichlids identified in this study. The

classification of ECM-related genes (the column of matrisome_annotation) follows a

previous study (Hynes and Naba, 2012).

Table. S3 All differentially accumulated proteins detected by LC–MS/MS under the

criteria of P-value < 0.05, log2 fold change > 1).

Table. S4 Results of enrichment analysis of highly expressed genes in hypertrophied

lips of juvenile and adult cichlids. The annotation terms were clustered by DAVID

annotation clustering.

Table. S5 All differentially expressed genes between hypertrophied lips and normal lips.

We compared adult Victoria cichlids (*H. chilotes* with hypertrophied lips and *H.*

sauvagei with normal lips), juvenile Victoria cichlids (*H. chilotes* and *H. sauvagei*),

Malawi cichlids (*Placidochromis milomo* with hypertrophied lips and *Labidochromis*

caeruleus with normal lips), and Taganyika cichlids (*Lobochilotes labiatus* with

hypertrophied lips and *Tropheus moorii* with normal lips).

Acknowledgement

This study was supported by JSPS KAKENHI (23KJ0951 to N. M., 17H04606,

20KK0167 to M.N.) and JST SPRING (JPMJSP2106). Computations were partially

performed on the NIG supercomputer at ROIS National Institute of Genetics.   The

authors thank the Open Research Facilities for Life Science and Technology, Tokyo

Institute of Technology for LC–MS/MS analysis (for technical assistance). The authors

thank the Hongoh laboratory members. Nagatoshi Machii thank my family, Nagashi

Machii, Mitsue M. and Sawa M.

Data availability

RNA-seq short reads and raw proteome data are stored at NCBI and jPOST (Okuda et

al., 2017) (Accession: PRJDB3401, PRJDB17757, and JPST003042/PXD051571).