Sex-peptide targets distinct higher order processing neurons in the brain to induce the female post-mating response

MOHANAKARTHIK P. NALLASIVAN[1], DEEPANSHU N.D. SINGH[1,2], MOHAMMED SYAHIR R. S. SALEH[1] AND MATTHIAS SOLLER[1,2,3]

[1]School of Biosciences, College of Life and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

[2]Birmingham Centre for Genome Biology, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

Running title: SP targets CNS neurons

Key Words: sex peptide, post-mating behaviors, neuronal wiring, sex peptide receptor (SPR), *fruitless* (*fru*), *doublesex* (*dsx*) and *pickpocket* (*ppk*) neurons

[3]Corresponding author: m.soller@bham.ac.uk

Abstract

Sex-peptide (SP) transferred during mating induces female post-mating responses including refractoriness to re-mate and increased oviposition in *Drosophila*. Yet, where SP target neurons reside, remained uncertain. Here we show that expression of membrane-tethered SP (mSP) in the head or trunk either reduces receptivity or increases oviposition, respectively. Using fragments from large regulatory regions of *Sex Peptide Receptor*, *fruitless* and *doublesex* genes together with intersectional expression of mSP, we identified distinct interneurons in the brain and abdominal ganglion controlling receptivity and oviposition. These interneurons can induce post-mating responses through SP received by mating. Trans-synaptic mapping of neuronal connections reveals input from sensory processing neurons and two post-synaptic trajectories as output. Hence, SP target neurons operate as key integrators of sensory information for decision of behavioural outputs. Multi-modularity of SP targets further allows females to adjust SP-mediated male manipulation to physiological state and environmental conditions for maximizing reproductive success.

Introduction

Reproductive behaviors are to a large degree hard-wired in the brain to guarantee reproductive success making the underlying neuronal circuits amenable to genetic analysis *(Anderson, 2016; Dulac and Kimchi, 2007; Rings and Goodwin, 2019; Yamamoto and Koganezawa, 2013)*.

During development, sex-specific circuits are built into the brain under the control of the sex determination genes *doublesex* (*dsx*) and *fruitless* (*fru*) in *Drosophila* *(Billeter et al., 2006; Schutt and Nothiger, 2000)*. They encode transcription factors that are alternatively spliced in a male or female specific mode *(Schutt and Nothiger, 2000)*. By default, the *dsx* gene generates the male-specific isoform Dsx^M, while a female-specific isoform Dsx^F is generated by alternative splicing and expressed in about ~700 distinct neurons in the brain important for female reproductive behaviors directing readiness to mate and egg laying *(Rezaval et al., 2012; Rideout et al., 2010)*. Fru^M is expressed in about ~1000 neurons in males and implements development of neuronal circuitry key to display male courtship behavior, but is switched off in females through alternative splicing by incorporation of a premature stop codon *(Demir and Dickson, 2005; Manoli et al., 2005; Stockinger et al., 2005)*.

The circuitry of female specific behaviors including receptivity to courting males for mating and egg laying have been mapped using intersectional gene expression via the *split-GAL4* system to restrict expression of activators or inhibitors of neuronal activity to very few neurons *(Aranha and Vasconcelos, 2018; Cury and Axel, 2023; Wang et al., 2020a; Wang et al., 2020b; Wang et al., 2021)*. Through this approach, sensory neurons in the genital tract have been identified as key signal transducers for the readiness to mate and the inhibition of egg laying connecting to central parts of the brain via projection to abdominal ganglion neurons *(Feng et al., 2014; Hasemeyer et al., 2009; Rezaval et al., 2012; Yang et al., 2009)*. This circuit then projects onto centrally localized pattern generators in the brain to direct a

behavioral response via efferent neurons *(Wang et al., 2020a; Wang et al., 2020b; Wang et al., 2021)*.

Once females have mated, they will reject courting males and lay eggs *(Manning, 1967)*. These post-mating responses (PMRs) are induced by male derived sex-peptide (SP) transferred during mating *(Avila et al., 2011; Chen et al., 1988; Hopkins and Perry, 2022)*. In addition, SP will induce a number of other behavioral and physiological changes including increased egg production, feeding, a change in food choice, sleep, memory, constipation, midgut morphology, stimulation of the immune system, and sperm storage and release *(Carvalho et al., 2006; Domanitskaya et al., 2007; Kim et al., 2010; Peng et al., 2005; Ribeiro and Dickson, 2010; Scheunemann et al., 2019; Soller et al., 1999) (Cognigni et al., 2011) (Avila et al., 2010; Isaac et al., 2010; Wainwright et al., 2021; White et al., 2021)*. SP binds to broadly expressed Sex Peptide Receptor (SPR), an ancestral receptor for myoinhibitory peptides (MIPs) *(Jang et al., 2017; Kim et al., 2010; Yapici et al., 2008)*. Although MIPs seem not to induce PMRs, excitatory activity of MIP expressing neurons underlies re-mating *(Jang et al., 2017; Kim et al., 2010; Yapici et al., 2008)*. Expression of membrane-tethered SP (mSP) induces PMRs in an autocrine fashion when expressed in neurons, but not glia *(Haussmann et al., 2013; Nakayama et al., 1997)*.

First attempts to identify SP target neurons by enhancer *GAL4* induced expression of *UASmSP* only identified lines with broad expression in the nervous system *(Nakayama et al., 1997)*. Later, drivers with more restricted expression including *dsx*, *fru* and *pickpocket* (*ppk*) genes were identified, but they are expressed in all parts of the nervous system throughout the body eluding to reveal the location of SP target sites unambiguously *(Hasemeyer et al., 2009; Haussmann et al., 2013; Rezaval et al., 2012; Yang et al., 2009; Yapici et al., 2008)*.

To delineate where in the *Drosophila* SP target neurons are located which induce the main PMRs, refusal to mate and egg laying, we expressed mSP specifically only in the head or

trunk. These experiments separate reduction of receptivity induced in the head from trunk induction of egg laying. To further restrict our search for SP target neurons, we focused on three genes, *SPR*, *dsx* and *fru,* because SPR is broadly expressed but anticipated to induce PMRs only from few neurons, and because *GAL4* inserted in the endogenous *dsx* and *fru* loci induces PMRs from mSP expression. Using *GAL4* tiling lines with fragments encompassing the regulatory regions of complex *SPR*, *fru* and *dsx* genes *(Jenett et al., 2012; Kvon et al., 2014; Pfeiffer et al., 2008)*, we identified one regulatory region in each gene reducing receptivity and inducing egg laying upon mSP expression, and one additional region in *SPR* only inducing egg laying. To further refine this analysis, we used intersectional gene expression using *split-GAL4* and *flipase* (*flp*) mediated excision of stop cassettes in *UAS* reporters *(Luan et al., 2006; Struhl and Basler, 1993)*. Consistent with previous results that the SP response can be induced via multiple pathways *(Haussmann et al., 2013)*, we found distinct sets of neurons in the central brain and the abdominal ganglion that can induce PMRs via expression of mSP. Mapping the pre- and post-synaptic connections of the distinct SP target neurons by *retro-* and *trans*-Tango *(Sorkaç et al., 2023; Talay et al., 2017)* revealed that SP target neurons direct higher order sensory processing in the central brain. These neurons feed into two common post-synaptic neuronal subtypes indicating that SP interferes with the integration of diverse sensory inputs to build a stereotyped output either reducing receptivity and/or increasing egg laying.

Results

Reduction of receptivity and induction of egg laying are separable by head and trunk expression of membrane-tethered SP

Due to the complex behavioral and physiological changes induced by SP, neurons in the central nervous system have been suspected as main targets for SP *(Kubli, 1992)*. To express

mSP only in the head we used an *elav FRTstopFRT GAL4* in combination with *otdflp*, that

expresses in the head to drive recombination and head-specific expression of mSP from *UAS*

(Figure 1A) *(Asahina et al., 2014; Haussmann et al., 2008; Nallasivan et al., 2021;*

Zaharieva et al., 2015). To express mSP only in the trunk we used *tshGAL4* (Figure 1B)

(Soller et al., 2006).

When we expressed mSP in the head, females reduced receptivity indistinguishable from

mated females, but did not lay eggs, thereby again demonstrating that the two main PMRs

can be separated (Figures 1C and 1D) *(Haussmann et al., 2013)*. In contrast, when we

expressed mSP in the trunk, females remained receptive, but laid eggs in numbers

indistinguishable from mated females (Figures 1C and 1D).

Moreover, *tshGAL4* is expressed in *fru*, *dsx*, *ppk* genital tract sensory neurons (Figures 1E-

1H). Since mSP expression with *tshGAL4* does not affect receptivity, these genital tract

neurons unlikely are direct targets for SP *(Haussmann et al., 2013)*. Taken together, these

results indicate presence of SP target neurons in the brain and ventral nerve cord (VNC) for

the reduction of receptivity and induction of egg laying, respectively.

Few restricted regulatory regions in large *SPR*, *fru* and *dsx* genes can induce the SP
response

Expression of mSP from *UAS* via *GAL4* inserts in *fru* and *dsx* genes induces a robust

reduction in receptivity and increase in egg laying *(Haussmann et al., 2013; Rezaval et al.,*

2012). To identify SP target neurons, we thought to dissect the broad expression pattern of

complex *SPR*, *fru* and *dsx* genes spanning 50-80 kb by identifying regulatory DNA fragments

in the enhancer regions that drive *UAS mSP* in a subset of neurons. For these experiments, we

analysed 22, 27 and 25 *GAL4* lines from the VDRC and Janelia tiling *GAL4* projects *(Jenett*

et al., 2012; Kvon et al., 2014; Pfeiffer et al., 2008) (Figures 2A-2C).

Strikingly, in *SPR*, *fru* and *dsx* genes we identified only one regulatory region in each gene

(*SPR8*, *fru11/12* and *dsx24*) that reduced receptivity and induced egg laying through *GAL4*

UAS expression of mSP (Figures 2D and 2E). In addition, we identified one line (*SPR12*) in

the *SPR* gene, that induced egg laying, but did not reduce receptivity consistent with previous

results that SP regulation of receptivity and egg laying can be split *(Haussmann et al., 2013)*.

All of these lines expressed in subsets of neurons in the central brain and the ventral nerve

cord in distinct, but reduced patterns compared to the expression of the *SPR*, *fru* and *dsx*

genes *(Rideout et al., 2010; Yapici et al., 2008; Zhou et al.)* (Figures 2F-2O). Moreover,

these lines showed prominent labelling of abdominal ganglion neurons in the VNC (Figures

2K-2O). In addition, all of these lines except *SPR12* are also expressed in genital tract

sensory neurons (Figure S1).

From all the 74 lines that we have analyzed for PMRs from *SPR*, *fru* and *dsx* genes, we also

analysed expression in genital tract sensory neurons as they had been postulated to be the

primary targets of SP *(Hasemeyer et al., 2009; Rezaval et al., 2012; Yang et al., 2009;*

Yapici et al., 2008). Apart from PMR inducing lines *SPR8*, *fru11*, *fru12* and *dsx24*, that

showed expression in genital tract sensory neurons, we identified three lines (*SPR3*, *SPR 21*

and *fru9*), which also robustly expressed in genital tract sensory neurons but did not induce

PMRs from expression of mSP (Figure S2).

Secondary ascending abdominal ganglion neurons can induce the PMRs from mSP

expression

A set of neurons has been identified in the abdominal ganglion important for egg laying by

silencing neuronal activity from a screen expressing the rectifying potassium channel Kir2.1

that identified six driver lines (*FD1-6*) *(Feng et al., 2014)*. These neurons have been termed

SAG (secondary ascending abdominal ganglion neurons) neurons, that are also

interconnected with myoinhibitory peptide sensing neurons *(Jang et al., 2017)*. Since

enhancer lines identified in *SPR*, *fru* and *dsx* genes are prominently expressed in the

abdominal ganglion, we tested whether mSP expression from these SAG neuron expressing

lines induced PMRs.

From these six lines, one robustly suppressed receptivity and induced egg laying

(*FD6/VT3280*), while two lines only induced egg laying (*FD3/VT4515* and *FD4/V000454*)

similar to controls from mSP expression (Figure 3). Again, all three lines also expressed in

subsets of neurons in the central brain and VNC, particularly in the abdominal ganglion

(Zhou et al.). In addition, *FD3* and *FD4* did not express in genital tract sensory neurons, in

contrast to *FD6 (Feng et al., 2014)*.

Intersectional expression reveals distinct mSP responsive neurons in the central brain

and abdominal ganglion

To further restrict the expression to fewer neurons, we intersected the expression patterns of

those lines that induced robust reduction of receptivity and increase of egg laying using *split-*

GAL4 (*SPR8*, *fru11/12*, *dsx* and *FD6*, for further experiments we used *dsxGAL4-DBD*,

because *dsx24* is less robust and *fru11* and *fru12* were made into one fragment) that activates

the *UAS* reporter when *GAL4* is reconstituted via dimerization of activation (AD-GAL4) and

DNA binding (GAL4-DBD) domains *(Luan et al., 2006)* (Figure 4A).

Again, intersection of *SPR8* with *fru11/12*, *dsx* or *FD6*, and *fru11/12* with *dsx* or *FD6*

expression robustly reduced receptivity and increased egg laying upon expression of mSP

(Figures 4B and 4C).

When we further analyzed the expression of these *split-GAL4* intersections in the brain, we

found that each combination first showed very restricted expression, but second, that none of

these combinations labeled the same neurons (Figures 4D-4H). For *dsx* neurons, *split-GAL4*

intersections correspond to a subset of dPC2l (*SPR8 ∩ dsx*) and dPCd-2 (*fru11/12 ∩ dsx*) neurons *(Deutsch et al., 2020; Nojima et al., 2021; Schretter et al., 2020)*. These results suggest the SP targets interneurons in the brain that feed into higher processing centers from different entry points likely representing different sensory input.

In the ventral nerve cord, we found expression in the abdominal ganglion with all *split-GAL4* combinations (Figures 4I-4M). In particular, intersection of *dsx* with *SPR8* or *fru11/12* showed exclusive expression in the abdominal ganglion, while the other combinations also expressed in other cells of the VNC. All together, these data suggest that the abdominal ganglion harbors several distinct type of neurons involved in directing PMRs *(Oliveira-Ferreira et al., 2023)*.

In the female genital tract, these split-Gal4 combinations show expression in genital tract neurons with innervations running along oviduct and uterine walls (Figures S3A-S3E). In addition, *SPR8 ∩ fru11/12* and *SPR8 ∩ dsx* were also expressed in the spermathecae (Figures S3A-S3B).

Both inhibitory and activating neurons have been attributed to impact on PMRs *(Kvitsiani and Dickson, 2006; Rezaval et al., 2012; Yapici et al., 2008)*. These neurons seem to be part of intersecting circuitry as general inhibition of *ppk* neurons by tetanus toxin (TNT) only partially blocks the SP response in contrast to inhibition of *ppk* neurons in the brain alone *(Nallasivan et al., 2021)*.

When we inhibited neuronal activity by expression of TNT *(Sweeney et al., 1995)*, we observed a significant reduction of receptivity for all *split-Gal4* combinations, though only partially for inhibition in *fru11/12 ∩ FD6* neurons. Likewise, all *split-Gal4* combinations induced a significant increase in egg laying (Figures S4A and S4B). Ablation of these neurons by expression of apoptosis inducing *reaper* and *hid* genes essentially replicated the results from neuronal inhibition indicating that SPR target neurons are modulatory and are

not part of motor circuits because females laid eggs and performed normally in receptivity assays (Figures S4C and S4D).

To evaluate the composition of the intersected expression patters into inhibitory and activating neurons we also expressed the *Bacillus halodurans* sodium channel (NaChBac) *(Feng et al., 2014)* to activate all of the intersected neurons. Here, we found a significant reduction of receptivity for four of the five *split-GAL4* combinations, though only partially for activation of *SPR8 ∩ dsx* neurons (Figure S4E). Activating *fru11/12 ∩ FD6* neurons did not reduce receptivity (Figure S4E). Likewise, we found the same pattern for the induction of egg laying (Figure S4F). Four of the five *split-GAL4* combinations induced a significant increase which was only partial in *SPR8 ∩ dsx* neurons and no egg laying was induced by activating *fru11/12 ∩ FD6* neurons.

Essentially, these results are consistent with previous findings that inhibitory neurons prevail *(Nallasivan et al., 2021)*, possibly as input from trunk neurons as found for *ppk* expressing neurons (see below).

***ppk* neurons do not intersect with SPR, fru, dsx and FD6 neurons in inducing PMRs by mSP**

Expression of mSP in *ppk* neurons can also induce PMRs (Figures S5A and S5B) *(Hasemeyer et al., 2009; Yang et al., 2009)*. The complement of *ppk* neurons labeled with *ppkGAL4* consists of at least two populations including sensory neurons and eight interneurons in the central brain *(Nallasivan et al., 2021)*. These brain neurons show severe developmental defects in SP-insensitive *Nup54* mutant alleles, but they receive inhibitory input from sensory neurons *(Nallasivan et al., 2021)*.

To evaluate whether *ppk* neurons are part of the previously identified expression patterns, we intersected them by crossing *GAL4-AD* lines *SPR8*, *SPR12* and *fru11/12* with a *ppk GAL4-*

DBD line containing the previously used 3 kb promoter fragment *(Grueber et al., 2003; Seidner et al., 2015)*. Surprisingly, none of these *split-GAL4* combinations reduced female receptivity or increased egg laying (Figures S5A and S5B). Likewise, no GFP expressing neurons were detected in the brain, abdominal ganglion or the genital tract (Figures S5C-S5E, S5F-S5H and S5I-S5K). Also, inhibiting or activating neurons with these split-Gal4 combinations did not reduce receptivity or induce egg laying (Figures S5L and S5O). How exactly *ppk* neurons impact on PMRs, however, needs to be further evaluated in follow-up studies.

mSP responsive neurons rely on SPR and are required for PMRs induced by SP delivered through mating

Next, we tested whether PMRs induced by mSP expression in the *SPR8 ∩ dsx, fru11/12 ∩ dsx* or *SPR8 ∩ fru11/12* rely on *SPR*. Expression of mSP in *dsx ∩ SPR8* and *dsx ∩ fru11/12* neurons in *SPR* mutant females did not reduce receptivity or induce egg laying (Figures 5A and B, see also Figures 4A and B), while a partial response was observed for *SPR8 ∩ fru 11/12* induced mSP expression in *SPR* mutant females, which is consistent with presence of additional receptors for SP *(Haussmann et al., 2013)*.

Since SP is transferred during mating to females and enters the hemolymph *(Haussmann et al., 2013)*, we wanted to test whether SPR is required in these neurons for inducing PMRs after mating. For *SPR RNAi* in *dsx ∩ fru11/12* and *SPR8 ∩ fru 11/12* neurons, no reduction, or a partial reduction, of receptivity was observed, respectively, while *SPR RNAi* in *dsx ∩ SPR8* neurons turned virgin females unreceptive (Figure 5A). Expression of mSP in *dsx ∩ fru11/12* neurons in the context of *SPR RNAi* partially reduced receptivity again suggesting additional receptors for SP *(Haussmann et al., 2013)*.

Strikingly, however, *SPR RNAi* in these neurons prevented egg laying independent of

whether SP was delivered by mating or when tethered to the membrane of these neurons

(Figure 5B).

These results demonstrate that neurons identified by *split-GAL4* intersected expression of

SPR8 with *dsx* or *fru11/12*, or *fru11/12* with *dsx* are genuine SP targets as they rely on *SPR*

and PMRs are induced by SP delivered through mating.

Expression of mSP in distinct neurons in the brain induces PMRs

The analysis of *ppk* neurons in SP-insensitive *Nup54* alleles revealed a hierarchy of trunk

neurons that dominate over central brain neurons *(Nallasivan et al., 2021)*. To focus on the

role of central brain neurons, we generated a *UAS mSP* line with a stop cassette (*UAS*

FRTstopFRT mSP) that allows to restrict expression of mSP to the head in the presence of

otdflp, which only expresses in the head (Figure 6A), but not in the trunk *(Asahina et al.,*

2014; Nallasivan et al., 2021).

In combination with the intersectional approach, we now can restrict mSP expression to few

central brain neurons, or alternatively activate or silence these neurons (Figure 6B).

Expression of mSP in *SPR8 ∩ dsx, fru11/12 ∩ dsx* or *SPR8 ∩ fru11/12* neurons mSP in the

central brain significantly reduced receptivity, but oviposition was only substantially induced

in *SPR8 ∩ dsx* brain neurons (Figures 6C and 6D). These results clearly demonstrate a role

for brain neurons in the SP response. However, we noticed that the flipase approach can

result in false negatives as *fruflp* inserted in the same position in the endogenous locus as

fruGAL4 does not induce a response with *UAS FRTstopFRT* mSP in contrast to *fruGAL4*

induced expression of mSP. In contrast, the same experiment with *dsxGAL4* and *dsxflp*

results in a positive SP response indistinguishable from mated females *(Haussmann et al.,*

2013).

Next, we tested whether neuronal activation or inhibition would induce a post-mating response. Strikingly, conditional activation of *SPR8 ∩ dsx, fru11/12 ∩ dsx* or *SPR8 ∩ fru11/12* brain neurons with TrpA1in adult females completely inhibited receptivity and induced egg laying comparable to mated females (Figures 6C and 6D). In contrast, inhibition of these neurons with tetanus toxin (TNT) did not alter the virgin state, e.g. receptivity was not reduced and egg laying was not induced (Figures 6C and 6D).

mSP responsive neurons operate in higher order sensory processing in the brain

With the split-GAL4 approach we identified five distinct neuronal sub-types that can induce PMRs. To find out whether these neurons receive input from distinct entry points in the brain and to identify the target neurons of these mSP responsive neurons, we used the *retro-* and *trans*-Tango technique to specifically activate reporter gene expression in up- and down-stream neurons *(Sorkaç et al., 2023; Talay et al., 2017)*(Figures 7A-O).

In the brain, the *retro*-Tango analysis did not identify primary sensory neurons, but higher order neurons in the central brain in all five *split-GAL4* combinations (Fig 7A-E). In addition, neurons in the suboesophagal ganglion were marked from *SPR8* intersections with *dsx* and *FD6*, and in *dsx ∩ fru11/12*. In *dsx ∩ fru11/12*, neurons in the optic lobe (medulla) were marked. In addition, a strong signal was observed in all five *split-GAL4* combinations in the mushroom bodies (Figs 7A-E). Although mushroom bodies are dispensable for PMRs *(Fleischmann et al., 2001)* their connection to SP target neurons indicates an experience dependent component of PMRs .

The *trans*-Tango analysis identified a subset of neurons with cell bodies in the suboesophageal ganglion with projections to the *pars intercerebralis* for *SPR8 ∩ dsx* and *fru11/12 ∩ dsx* neurons (Figures 7K and 7L). For *SPR8 ∩ fru11/12* and *SPR8 ∩ FD6* neurons common target neurons were found in the antennal mechanosensory and motor centre

(AMMC) region with a single neuron identified near the mushroom body region (Figures 7M and 7N) *(Ishimoto and Kamikouchi, 2021)*. For *fru11/12 ∩ FD6* no obvious targets were identified in the central brain (Figure 7O).

In the VNC, the *trans*-Tango analysis showed post-synaptic targets within the abdominal ganglion with all five *split-GAL4* combinations indicating an interconnected neuronal network (Figure S6A-S6O), which needs to be elaborated in detail. In the genital tract, no post-synaptic targets were detected indicating that these are afferent neurons integrating sensory input (Figure S6P-S6AD).

Taken together, circuitries identified via *retro-* and *trans-*Tango place SP target neurons at the interface of sensory processing interneurons connecting to two commonly shared post-synaptic processing neuronal populations in the brain. Hence, our data indicate that SP interferes with sensory input processing from multiple modalities that are canalized to higher order processing centres to generate a behavioural output.

Discussion

Much has been learned about the neuronal circuitry governing reproductive behaviors in *Drosophila* from interfering with neuronal activity in few neurons selected by intersectional expression using split-GAL4 *(Wang et al., 2020a; Wang et al., 2020b; Wang et al., 2021)*. However, how sex-peptide signaling as main inducer of the post-mating response, prominently consisting of refractoriness to re-mate and induction of egg laying, is integrated in this circuitry is not completely understood *(Haussmann et al., 2013)*. Here, we addressed this gap by identifying regulatory regions in *SPR*, *fru* and *dsx* genes driving membrane-tethered expression of SP in subsets of neurons to delineate SP targets to very few neurons in the central brain and the ventral nerve cord by intersectional expression. Consistent with previous analysis describing multiple pathways for the SP response *(Haussmann et al.,*

2013), we find five distinct populations of interneurons in the central brain directing PMRs.

In SP target neurons in the central brain, SPR is essential to induce PMRs when receiving SP

from males through mating. From mapping post-synaptic targets by *trans*-Tango, we

identified two populations of interneurons. The architecture of this circuitry is reminiscent

for processing of sensory input transmitted to central brain pattern generators for behavioral

output. Hence, SP interferes at several levels for coordinating PMRs, but also leaves the

female the opportunity to interfere under unfavorable conditions with specific elements of

PMRs, e.g., if there is no egg laying substrate, females will still not remate *(Haussmann et*

al., 2013). Likewise, mated females will not lay eggs despite suitable egg laying substrates if

parasitoid wasp are present *(Kacsoh et al., 2015)*. Thus, the architecture of female PMRs

contrasts with male-courtship behavior consisting of a sequel of behavioral elements that

once initiated will always follow stereotypically to the end culminating in mating, or start

from the beginning when interrupted *(Greenspan and Ferveur, 2000; Hall, 1994)*.

SP induces PMRs via entering the hemolymph to target neurons in the central brain

and ventral nerve cord

Early characterization of the SP signaling cascade demonstrated induction of PMRs from

various other sources than mating including transgenic secretion from the fat body,

expression as membrane-tethered form on neurons or injection of synthetic peptide into the

hemolymph *(Aigaki et al., 1991; Chen et al., 1988; Nakayama et al., 1997; Schmidt et al.,*

1993). Likewise, SP is detected in the hemolymph after mating at a PMR inducing

concentration *(Haussmann et al., 2013)*. Moreover, PMRs are induced faster, when SP is

injected compared to induction by mating *(Haussmann et al., 2013)*. This delay, however, is

not attributed to sperm binding of SP as it is unchanged after mating with spermless males.

These results suggest that SP reaches its targets through entering the circulatory system to

target neurons and contrasts a previously proposed model favoring genital tract neurons as SP

sensors from the lumen of the genital tract *(Hasemeyer et al., 2009; Rezaval et al., 2012;*

Yang et al., 2009).

In further support of the internalization model, we identified *GAL4* drivers that express mSP

in genital tract neurons, but do not induce PMRs. Also, *SPR12* does not express in genital

tract neurons, but induces egg laying by expression of mSP. Moreover, expression of mSP in

the trunk (including all genital tract sensory neurons), only induces egg laying, but does not

change receptivity. Likewise, expression of mSP specifically in the brain (*SPR8 ∩ dsx*) can

reduce receptivity and induce egg laying indistinguishable from mated females.

These results are in strong favor for SP entering the hemolymph to target neurons in the

ventral nerve cord for inducing egg laying, and in the central brain for reducing receptivity

and inducing egg laying.

Integration of SP signaling into the circuitry directing reproductive behaviours

Reduction of receptivity and induction of egg laying are both induced by the same critical

concentration of injected SP *(Haussmann et al., 2013; Schmidt et al., 1993)* initially

suggesting a simple on/off system for PMRs likely initiated from a small population of

neurons. However, such model would not allow to split the SP response into individual PMR

components by expression of mSP.

Here, we identified several *GAL4* drivers, that can induce only egg laying (*SPR12*, *FD3*, *FD4*

and *tsh GAL4*), but do not reduce receptivity. Most striking is *tshGAL4*, that expresses only in

the trunk. All of these lines express in the abdominal ganglion and *dsx* neurons in the

abdominal ganglion have been identified to induce egg laying *(Rezaval et al., 2012; Zhou et*

al.). Hence, this neuronal structure has a key role in regulating egg laying. Since more than a

single neuronal population seems to direct egg laying, further high-resolution mapping is

required to identify individual neuronal population within the abdominal ganglion *(Jang et al., 2017; Oliveira-Ferreira et al., 2023)*.

Since *tshGAL4* only induces egg laying, neurons in the brain must direct reduction of receptivity. Through intersectional expression in combination with head-specific expression of *otdflp*, we could express mSP only in the brain by FLP mediated brain-specific excision of a stop cassette. We observed a significant reduction in receptivity for all five intersections tested, but for four the response is only partial likely due to the inefficiency of FLP mediated recombination.

Moreover, brain neurons can also induce egg laying when *SPR8* is intersected with *dsx*, and to some extent also from *SPR8* intersection with *fru11/12*. Due to the inefficiency of FLP mediated recombination, however, this is likely an underestimate and solving this issue requires development of more robust tools.

In any case, however, our results show that PMRs can be induced from mSP expression from several sites suggesting interference with processing of sensory information at the level of interneurons. In particular, *SPR8* ∩ *fru11/1*2 neurons resemble auditory AMMC-B2 neurons involved in processing of information of the male love song *(Yamada et al., 2018)*. Likewise, *SPR8* ∩ *dsx* neurons seem to overlap with dimorphic *dsx* pCL2 interneurons that are part of the 26 neurons constituting the pC2 neuronal population involved in courtship song sensing, mating acceptance and ovipositor extrusion for rejection of courting males *(Deutsch et al., 2019; Kimura et al., 2015; Wang et al., 2020a)*. The *SPR8* ∩ *FD6* neurons resemble dopaminergic *fru* P1 neurons involved in courtship and the *fru11/12* ∩ *dsx* neurons seem to overlap with *dsx* pCd and neuropeptide F neurons involved in courtship *(Zhang et al., 2021)*. In females, pC1d neurons have been linked to aggression *(Deutsch et al., 2020; Schretter et al., 2020)*. The *fru11/12* ∩ *FD6* neurons resemble a class of gustatory pheromone sensing neurons *(Sakurai et al., 2013)*. Although we likely have not identified all SP sensing

neurons, our resources will provide a handle to future exploration of the details of this

neuronal circuitry incorporating SP signaling for inducing PMRs.

Conclusions

We have identified distinct SP sensing neurons in the central brain and the ventral nerve cord.

Since these five different SP sensing neuronal populations in the central brain converge into

two target sites, our data suggest a model (Figure 7P), whereby SP signaling interferes with

integration of sensory input. Independent interference with different sensory modalities opts

for the female to counteract male manipulation at the level of perception of individual

sensory cues to adapt to varying physiological and environmental conditions to maximize

reproductive success.

Acknowledgments: We thank T. Aigaki, G. Barnea, P. Soba, W.J. Joiner, B. Dickson, S.

Goodwin, C. Rezaval, D. Anderson, J.J. Hodge, A. Hidalgo, S. Collier, the Bloomington

stock center, the Vienna Drosophila RNAi Center for flies, T. Aigaki and W.J. Joiner for

plasmids, the University of Cambridge Department of Genetics Fly Facility and FlyORF for

injections, D. Scocchia for help with PCR and, I.U. Haussmann, Y.J. Kim, J.C. Billeter and

J-R Martin for comments on the manuscript. We acknowledge funding by the Biotechnology

and Biological Science Research Council to MS.

Author contributions: MS conceived and directed the project. MPN performed genetic

experiments and imaging. MS and SS performed genetic and DS imaging experiments. MPN

and MS analyzed data. MS wrote the manuscript with support from MPN. All authors read

and approved the final manuscript.

Data availability: Data is included in the paper and supplementary files. Source data is

provided.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Competing interests: The authors declare no competing interests.

Materials and methods

Key resources table

Reagent type or resource	Source	Identifier
Antibodies		
Anti-HA (rat, 3F10)	Roche	
Anti-GFP (rabbit)	Molecular Probes	
Goat anti-rabbit Alexa Fluor 488	Molecular Probes	
Goat anti-rabbit Alexa Fluor 546	Molecular Probes	
Goat anti-rabbit Alexa Fluor 647	Molecular Probes	
Goat anti-rat Alexa Fluor 647	Molecular Probes	
Bacterial and Virus Strains		
For recombinant DNA cloning: *Escherichia coli*: DH5α	New England Biolabs	
Experimental Models: Organisms/Strains		
Drosophila: wild-type: Canton S	Soller lab	
Drosophila ST stock: w; Sco/CyO actGFP; PrDr/TM6 Sb DfdGFP	Soller lab	
Drosophila CT stock: w; Sco/SM6 Roi; PrDr/TM3 Sb	Soller lab	
Drosophila: w*; UASmSP (3rd, 61C)	T. Aigaki	
Drosophila: dsx-GAL4 inserted into the endogenous dsx gene (84E5-84E6)	S. Goodwin	
Drosophila: fru-GAL4 inserted into the endogenous fru gene (91A6-91B3)	B. Dickson	
Drosophila: nSyb GAL4 (3rd)	S. Goodwin	
Drosophila: ppk-GAL4/CyO	Bloomington Stock Centre	BDSC 49021
Drosophila: tshGAL4-1/CyO	Bloomington Stock Centre	BDSC 3040
elav FRTstopFRT GAL4	Soller lab	
Otdflp (2nd)	D. Anderson	

UASmCD8GFP (X)	Bloomington Stock Centre	BDSC 5136
UASmCD8GFP (2nd)	Bloomington Stock Centre	BDSC 5137
UAS-H2B::YFP (2nd)	A. Hidalgo	
elavLexA (2nd)	Bloomington Stock Centre	BDSC 52676
LexAop NLStomato (2nd)	Bloomington Stock Centre	BDSC 66690
UAS TNT (2nd)	J.J. Hodge	
UAS TrpA1 (3rd)	Bloomington Stock Centre	BDSC 26264
UASFlybow 1.1 (myrGFP, 2nd)	Bloomington Stock Centre	BDSC 35537
UAS-NaCh::BacGFP (3rd)	Bloomington Stock Centre	BDSC9467
UAS reaper/FM7;UAS hid/CyO	Bloomington Stock Centre	BDSC 5823 BDSC65403
UAS FRTstopFRT GFP/CyO	B. Dickson	
UAS FRTstopFRT TNT/CyO	B. Dickson	
UAS FRTstopFRT TrpA1/CyO	B. Dickson	
UAS FRTstopFRT mSP (3rd)	This study	
UAS dicer2; UAS SPR RNAi (X, 3rd)	B. Dickson	
SPR	Bloomington Stock Centre	BDSC 7708
Df(1)JC70/FM7c	Bloomington Stock Centre	BDSC 944
Drosophila SPR8 AD: VT033616-p65.AD (attP40)	Bloomington Stock Centre	BDSC71392
Drosophila Fru11/12 AD: VT043695-p65.AD (attP40)	Bloomington Stock Centre	BDSC72065
Drosophila dsx DBD	S. Goodwin	
Drosophila dsx24 DBD: R42G02-GAL4.DBD (attP2)	Bloomington Stock Centre	
Drosophila SPR8 DBD: VT057286-Gal4.DBD (attP2)	Bloomington Stock Centre	BDSC71425
Drosophila fru11/12 DBD: VT043695-GAL4.DBD (attP2)	Bloomington Stock Centre	BDSC72788
Drosophila FD6 DBD: VT003280-GAL4.DBD (attP2)	Bloomington Stock Centre	BDSC75877
Drosophila ppk DBD: ppk-GAL4.DBD (VK00027, 89E11)	W. J. Joiner	
Drosophila SPR12 AD: VT057292-p65.AD (attP40)	Bloomington Stock Centre	BDSC72924
UAS-myrGFP QUAS-mtdTomato-3xHA; trans-Tango	Bloomington Stock Centre	BDSC 95317
QUAS-mtdTomato-3xHA; retro-Tango	G. Barnea	
Drosophila: fru1 GAL4: R23C03-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49021
Drosophila: fru2 GAL4, R22H11-GAL4 (attP2)	Bloomington Stock Centre	BDSC 48043

Drosophila: fru3 GAL4, R21H09-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49867
Drosophila: fru4 GAL4, R23C12-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49026
Drosophila: fru5 GAL4, R22F06-GAL4 (attP2)		KDRC 11848
Drosophila: fru6 GAL4, R23D03GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: fru7 GAL4, R22B09-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: fru8 GAL4, R23B12-GAL4 (attP2)	Korea Drosophila Resource Centre	KDRC 11849
Drosophila: fru9 GAL4, R22A02-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49868
Drosophila: fru10 GAL4, R22C05-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49301
Drosophila: fru11 GAL4, R22C11-lexA (attP40)	Bloomington Stock Centre	BDSC 52604
Drosophila: fru12 GAL4, R22A11-GAL4 (attP2)	Bloomington Stock Centre	BDSC 48966
Drosophila: fru13 GAL4, R23A06-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49009
Drosophila: fru14 GAL4, R22C03-GAL4 (attP2)	Korea Drosophila Resource Centre	KDRC 11868
Drosophila: fru15 GAL4, R23B04-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49016
Drosophila: fru16 GAL4, R22C07-GAL4 (attP2)	Bloomington Stock Centre	BDSC 48975
Drosophila: fru17 GAL4, R23C08-GAL4 (attP2)	Korea Drosophila Resource Centre	KDRC 11835
Drosophila: fru18 GAL4, R23C07GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: fru19 GAL4, R22B10-GAL4 (attP2)	Bloomington Stock Centre	BDSC 48969
Drosophila: fru20 GAL4, R22E10-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49302
Drosophila: fru21 GAL4, R22D11-GAL4 (attP2)	Bloomington Stock Centre	BDSC 48982
Drosophila: fru22 GAL4, R22H07-GAL4 (attP2)	Bloomington Stock Centre	BDSC 490003
Drosophila: fru23 GAL4, R21H02-GAL4 (attP2)	Korea Drosophila Resource Centre	KDRC 11847
Drosophila: fru24 GAL4, R23B11-GAL4 (attP2)	Bloomington Stock Centre	BDSC 49019
Drosophila: fru25 GAL4: VT043674-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: fru26 GAL4, VT043675-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: fru27 GAL4, VT043676-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx1 GAL4, R39E06-GAL4 (attP2)	Bloomington Stock Centre	BDSC 50051
Drosophila: dsx2 GAL4, R40A05-GAL4 (attP2)	Bloomington Stock Centre	BDSC 48138
Drosophila: dsx3 GAL4, R40F03-GAL4 (attP2)	Bloomington Stock Centre	BDSC 47355
Drosophila: dsx4 GAL4, R40F04-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx5 GAL4, R41A01-GAL4 (attP2)	Bloomington Stock Centre	N/A

Drosophila: dsx6 GAL4, R41D01GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx7 GAL4, R41F06-GAL4 (attP2)	Bloomington Stock Centre	BDSC 47584
Drosophila: dsx8 GAL4, R42C06-GAL4 (attP2)	Bloomington Stock Centre	BDSC 50150
Drosophila: dsx9 GAL4, R42D02-GAL4 (attP2)	Bloomington Stock Centre	BDSC 41250
Drosophila: dsx10 GAL4, R42D04-GAL4 (attP2)	Bloomington Stock Centre	BDSC 47588
Drosophila: dsx11 GAL4, VT038171-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx12 GAL4, VT038169-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx13 GAL4, VT038167-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx14 GAL4, VT038166-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx15 GAL4, VT038161-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx16 GAL4, VT038159-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx17 GAL4, VT038157-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx18 GAL4, VT038155-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx19 GAL4, VT038151-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx20 GAL4, VT038149-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx21 GAL4, P{VT038148-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx22 GAL4, P{VT038147-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx23 GAL4, R22H07-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx24 GAL4, R21H02-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: dsx25 GAL4, R21B01-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR1 GAL4, R78F09-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR2 GAL4, R78F11-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR3 GAL4, R78E11-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR4 GAL4, R78E12-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40002
Drosophila: SPR5 GAL4, R78G09-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40015
Drosophila: SPR6 GAL4, R78G08-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR7 GAL4, R78F07-GAL4 (attP2)	Bloomington Stock Centre	BDSC 47409
Drosophila: SPR8 GAL4, R78F10-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40007
Drosophila: SPR9 GAL4, R78G02-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40010

Drosophila: SPR10 GAL4, R78G07-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR11 GAL4, R78G04-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40012
Drosophila: SPR12 GAL4, R78F05-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR13 GAL4, R78G05-GAL4 (attP2)	Bloomington Stock Centre	BDSC 41308
Drosophila: SPR14 GAL4, R78G06-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR15 GAL4, R78G03-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40011
Drosophila: SPR16 GAL4, R78F06-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR17 GAL4, R78F12-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR18 GAL4, R78F03-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR19 GAL4, R78F01-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40003
Drosophila: SPR20 GAL4, R78G01-GAL4 (attP2)	Bloomington Stock Centre	BDSC 40009
Drosophila: SPR21 GAL4, R78F02-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: SPR22 GAL4, R78F08-GAL4 (attP2)	Bloomington Stock Centre	N/A
Drosophila: FD1 GAL4, VT050405-GAL4 (attP2)	Vienna Drosophila Stock Centre	VDSC
Drosophila: FD2 GAL4, VT007068-GAL4 (attP2)	Vienna Drosophila Stock Centre	VDSC
Drosophila: FD3 GAL4, VT045154-GAL4 (attP2),	Vienna Drosophila Stock Centre	VDSC
Drosophila: FD4 GAL4, VT000454-GAL4 (attP2)	Vienna Drosophila Stock Centre	VDSC
Drosophila: FD5 GAL4, VT050247-GAL4 (attP2)	Vienna Drosophila Stock Centre	VDSC
Drosophila: FD6 GAL4, VT003280-GAL4 (attP2)	Vienna Drosophila Stock Centre	VDSC

pUAST-GGTmSP FRTGFPstopFRT gBlock (FRT underlined) GAATTGGGAATTCGTTAACAGATCTGCGATCGC<u>GGCCCGGGGAT</u> <u>CTTGAAGTTCCTATTCCGAAGTTCCTATTCTCT</u>AGAAAGTATAGG AACTTCAGAGCGCTTTTGAAGCTAGCTAAAGAGCCTGCTAAAGC AAAAAAGAAGTCACCATGGTGTCGAGCGCAAGCAAGGGCGAGG AGCTGTTCACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGC GACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGG GCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCA CCGGCAAGCTGCCCGTGCCCTGGCCCACCCTCGTGACCACCCTGA CCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAG CAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAG GAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCG CGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATC GAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGG GGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTATATC ATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAAGA TCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCAC TACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCC CGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACC CCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACC GCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAGTACTC AGATCTTTGCAAGCTTGTAGAGTTTCCCATTTAATAATTCATATTA TCTCGAATCTAGTCAATTACGGCTTTCCTCAAATAGAAAAATAAA AAAAATGAAAAAATGCACTTGCCATTTAAACTTAGACGCGATAAC GAATTC<u>CGGGGATCTTGAAGTTCCTATTCCGAAGTTCCTATTCTCT</u> <u>AGAAAGTATAGGAACTTCAGAGCGCTTTTGAAGCT</u>GCGGCCGCG GCTCGACGGTATCGATAAGCTTG	IDT	

Fly strains and husbandry

Flies were kept on standard cornmeal-agar food (1%industrial-grade agar, 2.1% dried yeast, 8.6% dextrose, 9.7% cornmeal and 0.25% Nipagin, all in (w/v)) in a 12 h light : 12 h dark cycle. Propionic acid was omitted from fly food as acidity affects egg laying *(Gou et al., 2014)*. Genetic crosses were done in vials and kept at low density to ensure larvae were not competing for food and if necessary, additional live yeast was added. For all behavioral assays, virgin and mated Canton-S were used as controls. Virgin females, e.g. from crosses of *GAL4* with *UASmSP*, were collected after emergence within a 5 h window and well-fed with live yeast sprinkled on food for maximum egg production and allowed to sexually mature (3–5 days).

To recombine 2nd chromosome inserts for *splitGAL4AD* (*attP40*) and 3rd chromosome *splitGAL4DBD* (*attP2*), standard genetic crossing schemes were used and final stocks were

balanced with CyO and TM3 Sb (combined from from ST and CT stock, see key resource

list). *SplitGal4AD* and *DBD* combination lines were then crossed to *UASmSP*. For meiotic

recombination, final stocks were validated by behavioral analysis for *UAS mSP*, for *flp* with

eFeG UASCD8GFP to monitor GFP expression and for *otdflp UASstopTrpA* and *otdflp*

UASstopTNT by crossing to *elavGAL4* and monitored by lethality.

For enhanced recombination with *flp*, virgin females were transferred to 30º C after eclosion

and kept for 5 d at this temperature before performing the behavioral assays. For induction of

neuronal activity by temperature sensitive TrpA1, females were kept at 30º C.

To make *UAS FRTstopFRT mSP*, a gBlock (IDT) stop cassette with the FRT sequences used

in the eFeG plasmid *(Haussmann et al., 2008)* was inserted into NotI cut *pUAST-GGTmSP*

(gift from T. Aigaki) by Gibson assembly. In the stop-cassette, the *FRT* sequence is followed

by a *GFP* with a 3'UTR from *ewg* containing polyA site 1 from intron 6 *(Haussmann et al.,*

2011). Flies were transformed by *P*-element mediated transgenesis and a inserts on each

chromosome were established that show a robust post-mating response with *dsxflp*

indistinguishable from mated females.

Behavioral analysis

Females were examined for the main post-mating behaviors receptivity and oviposition as

described previously and as follows *(Soller et al., 1999; Soller et al., 2006)*. To generate

mated females, one female and three males were added to fly vials and observed until mating

and males were removed after mating. For receptivity tests, mature virgin or mated females

were added to fly vials (95 mm length and 24 mm diameter) containing Canton S males with

an aspirator and observed for 1 h, generally 3 females and 7 males. For these experiments,

males were separated from females at least one day before the experiment. Receptivity tests

were done in the afternoon with virgins, or 5-24 h after mating for controls. For oviposition,

females were placed individually in fly vials in the afternoon and the number of eggs laid was

counted the next day.

Statistical analysis

Sample size was based on previous studies, non-blinded and not pre-determined by statistical

methods (*Haussmann et al., 2013; Nallasivan et al., 2021; Soller et al., 1997*). Behavioral

data are representatives of at least three replicates that were performed on three different

days. Statistical analysis of behavioral experiments were performed using GraphPadPrism 9

(GraphPad by Dotmatics) using one way Anova followed by pairwise comparisons with

Tukey's test.

Immunohistochemistry and imaging

For the analysis of adult neuronal projection from *UAS CD8GFP, UAS*

H2BYFP,UASmyrGFP, lexAopNLStomato o*r QUAS mtdtomato3xHA* expressing brains,

ventral nerve cords or genital tracts, tissues were dissected in PBS (137 mM NaCl, 10 mM

phosphate, 2.7 mM KCl, pH7.4), fixed in 4% (w/v in PBS) paraformaldehyde for 15 minutes,

washed three times in PBST (PBS with 1% BSA and 0.3% Triton-X100), then once in PBS

for 10 mins, mounted in Vectashield (Vector Labs) and visualized with confocal microscopy

using a Leica TCS SP8. If signals were weak, antibody *in-situ* stainings were done as

described previously (*Haussmann et al., 2008)* for validation using rat anti-HA (MAb 3F10,

1:20; Roche), rabbit anti-GFP (Molecular Probes, 1:100) and visualized with Alexa Fluor 488

(1:250; Molecular Probes or Invitrogen), Alexa Fluor 546 (1:250; Molecular Probes or

Invitrogen) or Alexa Fluor 647 (1:250; Molecular Probes or Invitrogen). For imaging, tissues

were mounted in Vectashield (Vector Labs).

Confocal microscopy and image processing

Adult tissues were scanned using a Leica SP8 confocal microscope equipped with a set of fluorescent filters and hybrid detector (HyD). Adult brains were scanned using a 40x HC PL APO 40x/1.30 lens with oil, 1024 x 1024 resolution and 0.96 μm Z-step. VNC and genital tracts were scanned using a HC PL APO CS2 20X/0.75 with oil, 1024 x 1024 resolution and 0.96 μm Z-step. Images were obtained using Leica Application Suite X (LAS X) imaging acquisition software. Raw data files were in LIF format and were processed using FIJI. For high resolution mapping neurons were identified in the virtual fly brain based on registered *GAL4* expression and traces retrieved for modelling *(Galili et al., 2022; Phelps et al., 2021; Scheffer et al., 2020)*.

References

Aigaki, T., Fleischmann, I., Chen, P.S., Kubli, E., 1991. Ectopic expression of sex peptide alters reproductive behavior of female *D. melanogaster*. *Neuron* **7**, 557-563.

Anderson, D.J., 2016. Circuit modules linking internal states and social behaviour in flies and mice. *Nat Rev Neurosci* **17**, 692-704.

Aranha, M.M., Vasconcelos, M.L., 2018. Deciphering Drosophila female innate behaviors. *Curr Opin Neurobiol* **52**, 139-148.

Asahina, K., Watanabe, K., Duistermars, B.J., Hoopfer, E., Gonzalez, C.R., Eyjolfsdottir, E.A., Perona, P., Anderson, D.J., 2014. Tachykinin-expressing neurons control male-specific aggressive arousal in Drosophila. *Cell* **156**, 221-235.

Avila, F.W., Ravi Ram, K., Bloch Qazi, M.C., Wolfner, M.F., 2010. Sex peptide is required for the efficient release of stored sperm in mated Drosophila females. *Genetics* **186**, 595-600.

Avila, F.W., Sirot, L.K., LaFlamme, B.A., Rubinstein, C.D., Wolfner, M.F., 2011. Insect seminal fluid proteins: identification and function. *Annu Rev Entomol* **56**, 21-40.

Billeter, J.C., Rideout, E.J., Dornan, A.J., Goodwin, S.F., 2006. Control of male sexual

behavior in Drosophila by the sex determination pathway. *Curr Biol* **16**, R766-776.

Carvalho, G.B., Kapahi, P., Anderson, D.J., Benzer, S., 2006. Allocrine modulation of

feeding behavior by the Sex Peptide of Drosophila. *Curr Biol* **16**, 692-696.

Chen, P.S., Stumm-Zollinger, E., Aigaki, T., Balmer, J., Bienz, M., Bohlen, P., 1988. A male

accessory gland peptide that regulates reproductive behavior of female D. melanogaster. *Cell*

54, 291-298.

Cognigni, P., Bailey, A.P., Miguel-Aliaga, I., 2011. Enteric neurons and systemic signals

couple nutritional and reproductive status with intestinal homeostasis. *Cell Metab* **13**, 92-104.

Cury, K.M., Axel, R., 2023. Flexible neural control of transition points within the egg-laying

behavioral sequence in Drosophila. *Nat Neurosci* **26**, 1054-1067.

Demir, E., Dickson, B.J., 2005. fruitless splicing specifies male courtship behavior in

Drosophila. *Cell* **121**, 785-794.

Deutsch, D., Clemens, J., Thiberge, S.Y., Guan, G., Murthy, M., 2019. Shared Song Detector

Neurons in Drosophila Male and Female Brains Drive Sex-Specific Behaviors. *Curr Biol* **29**,

3200-3215.e3205.

Deutsch, D., Pacheco, D., Encarnacion-Rivera, L., Pereira, T., Fathy, R., Clemens, J.,

Girardin, C., Calhoun, A., Ireland, E., Burke, A., Dorkenwald, S., McKellar, C., Macrina, T.,

Lu, R., Lee, K., Kemnitz, N., Ih, D., Castro, M., Halageri, A., Jordan, C., Silversmith, W.,

Wu, J., Seung, H.S., Murthy, M., 2020. The neural basis for a persistent internal state in

Drosophila females. *Elife* **9**.

Domanitskaya, E.V., Liu, H., Chen, S., Kubli, E., 2007. The hydroxyproline motif of male

sex peptide elicits the innate immune response in Drosophila females. *FEBS J* **274**, 5659-

5668.

Dulac, C., Kimchi, T., 2007. Neural mechanisms underlying sex-specific behaviors in

vertebrates. *Curr Opin Neurobiol* **17**, 675-683.

Feng, K., Palfreyman, M.T., Hasemeyer, M., Talsma, A., Dickson, B.J., 2014. Ascending

SAG neurons control sexual receptivity of Drosophila females. *Neuron* **83**, 135-148.

Fleischmann, I., Cotton, B., Choffat, Y., Spengler, M., Kubli, E., 2001. Mushroom bodies

and post-mating behaviors of Drosophila melanogaster females. *J Neurogenet* **15**, 117-144.

Galili, D.S., Jefferis, G.S., Costa, M., 2022. Connectomics and the neural basis of behaviour.

Current opinion in insect science **54**, 100968.

Gou, B., Liu, Y., Guntur, A.R., Stern, U., Yang, C.H., 2014. Mechanosensitive neurons on

the internal reproductive tract contribute to egg-laying-induced acetic acid attraction in

Drosophila. *Cell reports* **9**, 522-530.

Greenspan, R.J., Ferveur, J.F., 2000. Courtship in Drosophila. *Annual review of genetics* **34**,

205-232.

Grueber, W.B., Ye, B., Moore, A.W., Jan, L.Y., Jan, Y.N., 2003. Dendrites of distinct classes

of Drosophila sensory neurons show different capacities for homotypic repulsion. *Curr Biol*

13, 618-626.

Hall, J.C., 1994. The mating of a fly. *Science* **264**, 1702-1714.

Hasemeyer, M., Yapici, N., Heberlein, U., Dickson, B.J., 2009. Sensory neurons in the

Drosophila genital tract regulate female reproductive behavior. *Neuron* **61**, 511-518.

Haussmann, I.U., Hemani, Y., Wijesekera, T., Dauwalder, B., Soller, M., 2013. Multiple

pathways mediate the sex-peptide-regulated switch in female Drosophila reproductive

behaviours. *Proc Biol Sci* **280**, 20131938.

Haussmann, I.U., Li, M., Soller, M., 2011. ELAV-mediated 3'-end processing of ewg

transcripts is evolutionarily conserved despite sequence degeneration of the ELAV-binding

site. *Genetics* **189**, 97-107.

Haussmann, I.U., White, K., Soller, M., 2008. Erect wing regulates synaptic growth in

Drosophila by integration of multiple signaling pathways. *Genome Biol* **9**, R73.

Hopkins, B.R., Perry, J.C., 2022. The evolution of sex peptide: sexual conflict, cooperation,

and coevolution. *Biological reviews of the Cambridge Philosophical Society* **97**, 1426-1448.

Isaac, R.E., Li, C., Leedale, A.E., Shirras, A.D., 2010. Drosophila male sex peptide inhibits

siesta sleep and promotes locomotor activity in the post-mated female. *Proc Biol Sci* **277**, 65-

70.

Ishimoto, H., Kamikouchi, A., 2021. Molecular and neural mechanisms regulating sexual

motivation of virgin female Drosophila. *Cell Mol Life Sci* **78**, 4805-4819.

Jang, Y.H., Chae, H.S., Kim, Y.J., 2017. Female-specific myoinhibitory peptide neurons

regulate mating receptivity in Drosophila melanogaster. *Nature communications* **8**, 1630.

Jenett, A., Rubin, G.M., Ngo, T.T., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D.,

Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, J.H., Peng, H., Trautman,

E.T., Svirskas, R.R., Myers, E.W., Iwinski, Z.R., Aso, Y., et al., 2012. A GAL4-driver line

resource for Drosophila neurobiology. *Cell reports* **2**, 991-1001.

Kacsoh, B.Z., Bozler, J., Ramaswami, M., Bosco, G., 2015. Social communication of

predator-induced changes in Drosophila behavior and germ line physiology. *Elife* **4**.

Kim, Y.J., Bartalska, K., Audsley, N., Yamanaka, N., Yapici, N., Lee, J.Y., Kim, Y.C.,

Markovic, M., Isaac, E., Tanaka, Y., Dickson, B.J., 2010. MIPs are ancestral ligands for the

sex peptide receptor. *Proc Natl Acad Sci U S A* **107**, 6520-6525.

Kimura, K., Sato, C., Koganezawa, M., Yamamoto, D., 2015. Drosophila ovipositor

extension in mating behavior and egg deposition involves distinct sets of brain interneurons.

PLoS One **10**, e0126445.

Kubli, E., 1992. The sex-peptide. *Bioessays* **14**, 779-784.

Kvitsiani, D., Dickson, B.J., 2006. Shared neural circuitry for female and male sexual

behaviours in Drosophila. *Curr Biol* **16**, R355-356.

Kvon, E.Z., Kazmar, T., Stampfel, G., Yáñez-Cuna, J.O., Pagani, M., Schernhuber, K.,

Dickson, B.J., Stark, A., 2014. Genome-scale functional characterization of Drosophila

developmental enhancers in vivo. *Nature* **512**, 91-95.

Luan, H., Peabody, N.C., Vinson, C.R., White, B.H., 2006. Refined spatial manipulation of

neuronal function by combinatorial restriction of transgene expression. *Neuron* **52**, 425-436.

Manning, A., 1967. The control of sexual receptivity in female Drosophila. *Animal*

Behaviour **15**, 239-250.

Manoli, D.S., Foss, M., Villella, A., Taylor, B.J., Hall, J.C., Baker, B.S., 2005. Male-specific

fruitless specifies the neural substrates of Drosophila courtship behaviour. *Nature*.

Nakayama, S., Kaiser, K., Aigaki, T., 1997. Ectopic expression of sex-peptide in a variety of

tissues in Drosophila females using the P[GAL4] enhancer-trap system. *Mol Gen Genet* **254**,

449-455.

Nallasivan, M.P., Haussmann, I.U., Civetta, A., Soller, M., 2021. Channel nuclear pore

protein 54 directs sexual differentiation and neuronal wiring of female reproductive behaviors

in Drosophila. *BMC biology* **19**, 226.

Nojima, T., Rings, A., Allen, A.M., Otto, N., Verschut, T.A., Billeter, J.C., Neville, M.C.,

Goodwin, S.F., 2021. A sex-specific switch between visual and olfactory inputs underlies

adaptive sex differences in behavior. *Curr Biol* **31**, 1175-1191.e1176.

Oliveira-Ferreira, C., Gaspar, M., Vasconcelos, M.L., 2023. Neuronal substrates of egg-

laying behaviour at the abdominal ganglion of Drosophila melanogaster. *Sci Rep* **13**, 21941.

Peng, J., Chen, S., Busser, S., Liu, H., Honegger, T., Kubli, E., 2005. Gradual Release of

Sperm Bound Sex-Peptide Controls Female Postmating Behavior in Drosophila. *Current*

Biology **15**, 207-213.

Pfeiffer, B.D., Jenett, A., Hammonds, A.S., Ngo, T.T., Misra, S., Murphy, C., Scully, A.,

Carlson, J.W., Wan, K.H., Laverty, T.R., Mungall, C., Svirskas, R., Kadonaga, J.T., Doe,

C.Q., Eisen, M.B., Celniker, S.E., Rubin, G.M., 2008. Tools for neuroanatomy and

neurogenetics in Drosophila. *Proc Natl Acad Sci U S A* **105**, 9715-9720.

Phelps, J.S., Hildebrand, D.G.C., Graham, B.J., Kuan, A.T., Thomas, L.A., Nguyen, T.M.,

Buhmann, J., Azevedo, A.W., Sustar, A., Agrawal, S., Liu, M., Shanny, B.L., Funke, J.,

Tuthill, J.C., Lee, W.A., 2021. Reconstruction of motor control circuits in adult Drosophila

using automated transmission electron microscopy. *Cell* **184**, 759-774.e718.

Rezaval, C., Pavlou, H.J., Dornan, A.J., Chan, Y.B., Kravitz, E.A., Goodwin, S.F., 2012.

Neural circuitry underlying Drosophila female postmating behavioral responses. *Curr Biol*

22, 1155-1165.

Ribeiro, C., Dickson, B.J., 2010. Sex peptide receptor and neuronal TOR/S6K signaling

modulate nutrient balancing in Drosophila. *Curr Biol* **20**, 1000-1005.

Rideout, E.J., Dornan, A.J., Neville, M.C., Eadie, S., Goodwin, S.F., 2010. Control of sexual

differentiation and behavior by the doublesex gene in Drosophila melanogaster. *Nat Neurosci*

13, 458-466.

Rings, A., Goodwin, S.F., 2019. To court or not to court - a multimodal sensory decision in

Drosophila males. *Current opinion in insect science* **35**, 48-53.

Sakurai, A., Koganezawa, M., Yasunaga, K., Emoto, K., Yamamoto, D., 2013. Select

interneuron clusters determine female sexual receptivity in Drosophila. *Nature*

communications **4**, 1825.

Scheffer, L.K., Xu, C.S., Januszewski, M., Lu, Z., Takemura, S.Y., Hayworth, K.J., Huang,

G.B., Shinomiya, K., Maitlin-Shepard, J., Berg, S., Clements, J., Hubbard, P.M., Katz, W.T.,

Umayam, L., Zhao, T., Ackerman, D., Blakely, T., Bogovic, J., Dolafi, T., Kainmueller, D.,

et al., 2020. A connectome and analysis of the adult Drosophila central brain. *Elife* **9**.

Scheunemann, L., Lampin-Saint-Amaux, A., Schor, J., Preat, T., 2019. A sperm peptide

enhances long-term memory in female Drosophila. *Sci Adv* **5**, eaax3432.

Schmidt, T., Choffat, Y., Klauser, S., Kubli, E., 1993. The *Drosophila melanogaster* SP: A

molecular analysis of structure-function relationships. *J Insect Physiol* **39**, 361-368.

Schretter, C.E., Aso, Y., Robie, A.A., Dreher, M., Dolan, M.J., Chen, N., Ito, M., Yang, T.,

Parekh, R., Branson, K.M., Rubin, G.M., 2020. Cell types and neuronal circuitry underlying

female aggression in Drosophila. *Elife* **9**.

Schutt, C., Nothiger, R., 2000. Structure, function and evolution of sex-determining systems

in Dipteran insects. *Development* **127**, 667-677.

Seidner, G., Robinson, J.E., Wu, M., Worden, K., Masek, P., Roberts, S.W., Keene, A.C.,

Joiner, W.J., 2015. Identification of Neurons with a Privileged Role in Sleep Homeostasis in

Drosophila melanogaster. *Curr Biol* **25**, 2928-2938.

Soller, M., Bownes, M., Kubli, E., 1997. Mating and Sex Peptide Stimulate the Accumulation

Of Yolk In Oocytes Of Drosophila Melanogaster. *European Journal of Biochemistry* **243**,

732-738.

Soller, M., Bownes, M., Kubli, E., 1999. Control of oocyte maturation in sexually mature

Drosophila females. *Dev Biol* **208**, 337-351.

Soller, M., Haussmann, I.U., Hollmann, M., Choffat, Y., White, K., Kubli, E., Schafer, M.A.,

2006. Sex-peptide-regulated female sexual behavior requires a subset of ascending ventral

nerve cord neurons. *Curr Biol* **16**, 1771-1782.

Sorkaç, A., Moșneanu, R.A., Crown, A.M., Savaș, D., Okoro, A.M., Memiș, E., Talay, M.,

Barnea, G., 2023. retro-Tango enables versatile retrograde circuit tracing in Drosophila. *Elife*

12.

Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirian, L., Dickson, B.J., 2005. Neural circuitry

that governs Drosophila male courtship behavior. *Cell* **121**, 795-807.

Struhl, G., Basler, K., 1993. Organizing activity of wingless protein in Drosophila. *Cell* **72**,

527-540.

Sweeney, S.T., Broadie, K., Keane, J., Niemann, H., O'Kane, C.J., 1995. Targeted expression

of tetanus toxin light chain in Drosophila specifically eliminates synaptic transmission and

causes behavioral defects. *Neuron* **14**, 341-351.

Talay, M., Richman, E.B., Snell, N.J., Hartmann, G.G., Fisher, J.D., Sorkaç, A., Santoyo,

J.F., Chou-Freed, C., Nair, N., Johnson, M., Szymanski, J.R., Barnea, G., 2017.

Transsynaptic Mapping of Second-Order Taste Neurons in Flies by trans-Tango. *Neuron* **96**,

783-795.e784.

Wainwright, S.M., Hopkins, B.R., Mendes, C.C., Sekar, A., Kroeger, B., Hellberg, J., Fan,

S.J., Pavey, A., Marie, P.P., Leiblich, A., Sepil, I., Charles, P.D., Thézénas, M.L., Fischer, R.,

Kessler, B.M., Gandy, C., Corrigan, L., Patel, R., Wigby, S., Morris, J.F., Goberdhan, D.C.I.,

Wilson, C., 2021. Drosophila Sex Peptide controls the assembly of lipid microcarriers in

seminal fluid. *Proc Natl Acad Sci U S A* **118**.

Wang, F., Wang, K., Forknall, N., Parekh, R., Dickson, B.J., 2020a. Circuit and Behavioral

Mechanisms of Sexual Rejection by Drosophila Females. *Curr Biol* **30**, 3749-3760.e3743.

Wang, F., Wang, K., Forknall, N., Patrick, C., Yang, T., Parekh, R., Bock, D., Dickson, B.J.,

2020b. Neural circuitry linking mating and egg laying in Drosophila females. *Nature* **579**,

101-105.

Wang, K., Wang, F., Forknall, N., Yang, T., Patrick, C., Parekh, R., Dickson, B.J., 2021.

Neural circuit mechanisms of sexual receptivity in Drosophila females. *Nature* **589**, 577-581.

White, M.A., Bonfini, A., Wolfner, M.F., Buchon, N., 2021. Drosophila melanogaster sex

peptide regulates mated female midgut morphology and physiology. *Proc Natl Acad Sci U S*

A **118**.

Yamada, D., Ishimoto, H., Li, X., Kohashi, T., Ishikawa, Y., Kamikouchi, A., 2018.

GABAergic Local Interneurons Shape Female Fruit Fly Response to Mating Songs. *J*

Neurosci **38**, 4329-4347.

Yamamoto, D., Koganezawa, M., 2013. Genes and circuits of courtship behaviour in

Drosophila males. *Nat Rev Neurosci* **14**, 681-692.

Yang, C.H., Rumpf, S., Xiang, Y., Gordon, M.D., Song, W., Jan, L.Y., Jan, Y.N., 2009.

Control of the postmating behavioral switch in Drosophila females by internal sensory

neurons. *Neuron* **61**, 519-526.

Yapici, N., Kim, Y.J., Ribeiro, C., Dickson, B.J., 2008. A receptor that mediates the post-

mating switch in Drosophila reproductive behaviour. *Nature* **451**, 33-37.

Zaharieva, E., Haussmann, I.U., Brauer, U., Soller, M., 2015. Concentration and localization

of co-expressed ELAV/Hu proteins control specificity of mRNA processing. *Mol Cell Biol*

35, 3104-3115.

Zhang, S.X., Glantz, E.H., Miner, L.E., Rogulja, D., Crickmore, M.A., 2021. Hormonal

control of motivational circuitry orchestrates the transition to sexuality in Drosophila. *Sci Adv*

7, eabg6926.

Zhou, C., Pan, Y., Robinett, C.C., Meissner, G.W., Baker, B.S., Central brain neurons

expressing doublesex regulate female receptivity in Drosophila. *Neuron* **83**, 149-163.

Figure legends

Figure 1: The main PMRs in females can be separated.

A, B) Schematic depiction of head and trunk expression in *Drosophila elav FRTstopFRT*

GAL4; *otdflp* (A) and in *tshGAL4* (B) visualized by *UAS GFP* (green).

C, D) Receptivity (C) and oviposition (D) of wild type control virgin (red) and mated (orange) females, and virgin females expressing *UAS mSP* (green) pan-neuronally with *nsybGAL4* or in head and trunk patterns shown as means with standard error from three repeats for receptivity (21 females per repeat) by counting the number of females mating within a 1 h period or for oviposition by counting the eggs laid within 18 hours from 30 females. Statistically significant differences from ANOVA post-hoc comparison are indicated by different letters ($p < 0.0001$).

E-H) Representative adult female genital tract showing *tshGAL4 UAS H2BYFP* (green) and *elavLexA LexAop NLStomato* (red) nuclear expression. The magnification (F-H) shows sensory genital tract neurons. Scale bar shown in E and H are 100 μm and 20 μm, respectively.

Figure 2: Distinct regulatory regions in *SPR*, *fru* and *dsx* genes induce PMRs from mSP expression.

A-C) Schematic representation of *SPR*, *fru*, and *dsx* chromosomal regions depicting coding and non-coding exons as black or white boxes, respectively, and splicing patterns in solid lines. Vertical lines below the gene model depict enhancer *GAL4* lines with names and those in red showed PMRs by expression of mSP.

D, E) Receptivity (D) and oviposition (E) of wild type control virgin (red) and mated (orange) females, and virgin females expressing *UAS mSP* (green) under the control of GAL4 pan-neuronally in *nsyb* or in *SPR8*, *SPR12*, *fru11*, *fru12*, and *dsx24* patterns shown as means with standard error from three repeats for receptivity (21 females per repeat) by counting the number of females mating within a 1 h period or for oviposition by counting the eggs laid within 18 hours from 30 females. Statistically significant differences from ANOVA post-hoc comparison are indicated by different letters ($p \leq 0.0001$).

F-O) Representative adult female brains (F-J) and ventral nerve cords (VNC, K-O) expressing *UAS CD8GFP* under the control of *SPR8*, *SPR12*, *fru11*, *fru12* and *dsx24 GAL4*. Scale bars shown in J and O are 50 μm and 100 μm, respectively.

Figure 3: Expression of mSP in secondary ascending abdominal ganglion neurons induces PMRs.

A, B) Receptivity (A) and oviposition (B) of wild type control virgin (red) and mated (orange) females, and virgin females expressing *UAS mSP* (green) under the control of GAL4 pan-neuronally in *nsyb* or in *FD1*, *FD2*, *FD3*, *FD4*, *FD5*, and *FD6* patterns shown as means with standard error from three repeats for receptivity (21 females per repeat) by counting the number of females mating within a 1 h period or for oviposition by counting the eggs laid within 18 hours from 30 females. Statistically significant differences from ANOVA post-hoc comparison are indicated by different letters (p<0.0001).

Figure 4: Distinct circuits from intersection of *SPR*, *fru*, *dsx* and *FD6* patterns in the brain and VNC induce PMRs from mSP expression.

A) Schematic showing the intersectional gene expression approach: GAL4 activation (AD, orange) and DNA binding domains (DBD, blue) are expressed in different, but overlapping patterns. Leucine zipper dimerization reconstitutes a functional split-GAL4 in the intersection (pink) to express *UAS* reporters.

B, C) Receptivity (B) and oviposition (C) of wild type control virgin (red) and mated (orange) females, and virgin females expressing *UAS mSP* (green) under the control of *split-GAL4* intersecting *SPR8 ∩ fru11/12, SPR8 ∩ dsx, SPR8 ∩ FD6, fru11/12 ∩ dsx* and *fru11/12 ∩ FD6* patterns shown as means with standard error from three repeats for receptivity (21 females per repeat) by counting the number of females mating within a 1 h period or for

oviposition by counting the eggs laid within 18 hours from 30 females. Statistically

significant differences from ANOVA post-hoc comparison are indicated by different letters

(p<0.0001).

D-M) Representative adult female brains and ventral nerve cords (VNC) expressing *UAS*

CD8GFP under the control of *SPR8 ∩ fru11/12, SPR8 ∩ dsx, SPR8 ∩ FD6, fru11/12 ∩ dsx*

and *fru11/12 ∩ FD6*. Scale bars shown in H and M are 50 μm and 100 μm, respectively.

Figure 5: Distinct neuronal circuitries from intersection of *SPR*, *fru* and *dsx* sense SP

after mating to induce PMRs.

A, B) Receptivity (A) and oviposition (B) of wild type control virgin (red) and mated

(orange) females, and virgin females expressing *UAS mSP* (green) under the control of *split-*

Gal4 intersecting *SPR8 ∩ dsx, fru11/12 ∩ dsx,* and *SPR8 ∩ fru11/12* patterns in *SPR/Df*

mutant females or *SPR* RNAi knock-down shown as means with standard error from three

repeats for receptivity (21 females per repeat) by counting the number of females mating

within a 1 h period or for oviposition by counting the eggs laid within 18 hours from 30

females. Statistically significant differences from ANOVA post-hoc comparison are indicated

by different letters (p<0.0001 except p=0.002 and p=0.006 for c and d in A, and p=0.004 for

c in B).

Figure 6: Distinct neuronal circuitries in the brain senses SP to induce PMRs.

A, B) Schematic depiction of *UAS GFP* (green) expression in the head of *Drosophila* (A)

combining *split-GAL4* intersectional expression (*AD-GAL4* and *GAL4-DBD*) with brain-

expressed *otdflp* mediated recombination of *UAS FRTGFPstopFRTmSP* (B).

C, D) Receptivity (C) and oviposition (D) of wild type control virgin (red) and mated

(orange) females, and virgin females expressing *UAS FRTGFPstopFRTmSP* (grey), *UAS*

FRTGFPstopFRTTrpA1 (purple) and *UAS FRTGFPstopFRTTNT* (pink) under the control of

split-GAL4 intersecting *SPR8 ∩ dsx, fru11/12 ∩ dsx* and *SPR8 ∩ fru11/12* patterns with

brain-specific FRT-mediated recombination by *otdflp* shown as means with standard error

from three repeats for receptivity (21 females per repeat) by counting the number of females

mating within a 1 h period or for oviposition by counting the eggs laid within 18 hours from

30 females. Statistically significant differences from ANOVA post-hoc comparison are

indicated by different letters ($p < 0.0001$ except $p < 0.0004$ for c in C, $p < 0.007$ for c in D).

Figure 7: *retro*- and *trans*-Tango identification of pre- and post-synaptic neurons of SP

target neurons reveals higher order neuronal input canalized into shared output

circuitries.

A-O) Representative adult female brains expressing *QUAST tomato3xHA retro*-Tango (left,

A-E), *UAS myrGFP* (middle, F-J) and *QUAST tomato3xHA trans*-Tango (right, K-O) in

SPR8 ∩ dsx, fru11/12 ∩ dsx , SPR8 ∩ fru11/12, SPR8 ∩ FD6 and *fru11/12 ∩ FD6* split-

GAL4s. The presynaptic (A-E, left), *split-GAL4* (F-J, middle) and postsynaptic (K-O, right)

neuronal circuitries are shown in an inverted grey background. Arrows (magenta) indicate

neurons and their corresponding projections in different regions in female brain. The scale

bar shown in O is 50 μm.

P) Model for the SP induced post-mating response. SP interferes with interpretation of

sensory cues, e.g. vison, hearing, smell, taste, and touch at distinct sites in the brain indicated

by higher order projections revealed by intersectional expression in the following patterns:

SPR8 ∩ dsx (blue)*, fru11/12 ∩ dsx* (black)*, SPR8 ∩ fru11/12* (yellow)*, SPR8 ∩ FD6* (pink)

and *fru11/12 ∩ FD6* (olive). and VNC (*fru11/12 ∩ dsx*) during higher order neuronal

processing.

Supplementary information

Supplementary Figure S1: Expression analysis of PMR-inducing *GAL4* in the genital tract.

A-E) Representative adult female genital tracts expressing *UAS CD8GFP* under the control of *SPR8*, *SPR12*, *fru11*, *fru12* and *dsx24 GAL4*, and *LexAop NLStomato* under the control of *elavLexA*. Arrows indicate genital tract sensory neurons. The insert shows expression of GFP in the genital tract sensory neurons. Scale bars shown in A and insets are 100 μm and 20 μm, respectively.

Supplementary Figure S2: Expression analysis of non-PMR-inducing *GAL4* in the genital tract.

A-C) Representative adult female genital tracts expressing *UAS CD8GFP* under the control of *SPR3*, *SPR21* and *fru9 GAL4*. Arrows indicate genital tract sensory neurons. The insert shows expression of GFP in the genital tract sensory neurons. Scale bars shown in C and insets are 100 μm and 20 μm, respectively.

Supplementary Figure S3: Expression analysis of split-GAL4 in the genital tract.

A-E) Representative adult female genital tracts expressing *UAS CD8GFP* under the control of *SPR8 ∩ fru11/12, SPR8 ∩ dsx, SPR8 ∩ FD6, fru11/12 ∩ dsx* and *fru11/12 ∩ FD6 split-GAL4* intersectional patterns. The scale bar shown in E is 100 μm.

Supplementary Figure S4: PMRs after neuronal inhibition, ablation or activation of distinct circuits from intersection of *SPR*, *fru*, *dsx* and *FD6* patterns in the brain and VNC.

A-F) Receptivity (A, C and E) and oviposition (B, D and F) of wild type control virgin (red) and mated (orange) females, and virgin females expressing either *UAS TNT* (azure, A and B) or *UAS reaper hid* to inhibit or ablate neurons (yellow, C and D), respectively, or *UAS NaChBac* (brown, E and F) to activate neurons in *SPR8 ∩ fru11/12, SPR8 ∩ dsx, SPR8 ∩ FD6, fru11/12 ∩ dsx* and *fru11/12 ∩ FD6 split-Gal4* patterns shown as means with standard error from three repeats for receptivity (21 females per repeat) by counting the number of females mating within a 1 h period or for oviposition by counting the eggs laid within 18 hours from 30 females. Statistically significant differences from ANOVA post-hoc comparison are indicated by letters (p≤0.0095 in A and B, p<0.0001 in C and D except p=0.016 for c in D, p<0.0001 in E and p<0.0002 in F).

Supplementary Figure S5: *ppk* is not part of the *SPR8, SPR12 and fru11/12* PMR-inducing neuronal circuitry

A, B) Receptivity (A) and oviposition (B) of wild type control virgin (red) and mated (orange) females, and virgin females expressing *UAS mSP* (green) under the control of *GAL4* in *ppk* or in *SPR8 ∩ ppk, SPR12 ∩ ppk,* and *fru11/12 ∩ ppk* patterns shown as means with standard error from three repeats for receptivity (21 females per repeat) by counting the number of females mating within a 1 h period or for oviposition by counting the eggs laid within 18 hours from 30 females. Statistically significant differences from ANOVA post-hoc comparison are indicated by different letters (p<0.0001).

C-K) Representative adult female brains, ventral nerve cords (VNC) and genital tracts expressing *UAS CD8GFP* under the control of *UAS* by *SPR8 ∩ ppk, SPR12 ∩ ppk,* and *fru11/12 ∩ ppk*. Scale bars shown in E are 50 μm and in H and K are 100 μm, respectively.

L-O) Receptivity (L, N) and oviposition (M, O) of wild type control virgin (red) and mated (orange) females, and virgin females expressing either *UAS TNT* (azure) or *UAS NaChBac*

(brown) to inhibit or activate neurons in *SPR8 ∩ ppk, SPR12 ∩ ppk,* and *fru11/12 ∩ ppk*

patterns shown as means with standard error from three repeats for receptivity (21 females

per repeat) by counting the number of females mating within a 1 h period or for oviposition

by counting the eggs laid within 18 hours from 30 females. Statistically significant

differences from ANOVA post-hoc comparison are indicated by different letters (p<0.001 for

b, and p<0.01 for c in L and N).

Supplementary Figure S6: *trans*-Tango identifies post-synaptic proceeding neurons of

SP targets in the VNC, but not the genital tract

A-O) Representative adult female ventral nerve cords (VNC, A-O) and genital tracts (P-AD)

expressing *UAS myrGFP; QUAST tomato3xHA trans*-Tango in *SPR8 ∩ fru11/12, SPR8 ∩*

dsx, SPR8 ∩ FD6, fru11/12 ∩ dsx and *fru11/12 ∩ FD6* split-*GAL4s*. The presynaptic (A-E

and P-T) and postsynaptic (F-J and U-Y) neuronal circuitries are shown in an inverted grey

background and the merge is shown in colour. In the merged picture (K-O and Z-AD), the

pre-synaptic and post synaptic neuronal circuitry is shown in green and magenta,

respectively. Scale bars shown in O and AD are 100 μm.

Figure 1

A Head expression

B Trunk expression

C Virgin ▮ Mated ▮
UAS mSP (in virgins) ▮

Receptivity (%)

Wild type | Neuron | Head | Trunk

D

Oviposition (eggs/18h)

Wild type | Neuron | Head | Trunk

E tshGal4 UAS H2BYFP
elavLexA LexAop NLStomato

GFP
RFP

F **G** **H**

Figure 3

UAS CD8GFP

Figure 5

Figure 6

Figure 7