Ecological divergence is marked by species specific plastic gene expression and distinct rhizosphere among sibling allopolyploid marsh orchids (*Dactylorhiza*)

Katie Emelianova[1*], Anna-Sophie Hawranek[1], Mimmi C. Eriksson[1,2,4], Thomas M. Wolfe[1,2,3], Ovidiu Paun[1*]

[1]*Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, A-1030 Vienna, Austria.*

[2]*Vienna Graduate School of Population Genetics, Vienna, Austria.*

[3]*Institute of Forest Entomology, Forest Pathology and Forest Protection (IFFF), BOKU, Peter-Jordan-Straße 82/I, A−1190 Vienna, Austria.*

[4]*Department of Plant Biology, Swedish University of Agricultural Sciences, Uppsala, Sweden*

*corresponding authors: katherine.emelianova@univie.ac.at; ovidiu.paun@univie.ac.at

Abstract:

- Phenotypic plasticity, the dynamic adjustment of traits to environmental variations, is pivotal to enabling species to exploit broader niches and withstand suboptimal conditions. This adaptability is expected to be particularly relevant for newly formed allopolyploids, which have redundant gene copies and must establish themselves in diverse environments, distinct from their parents and other relatives.
- Here we investigate ecologically divergent sibling allopolyploid marsh orchids (*Dactylorhiza*) through reciprocal transplantation experiments at two localities to understand the drivers of ecological divergence and species persistence in the face of interspecific gene flow.
- Although consistent abiotic differences characterise the alternative environments, the great majority of gene expression differences between the sibling allopolyploids is plastic. Few genes exhibit fixed expression differences, suggesting that despite ecological divergence, plasticity may mediate gene flow between sibling allopolyploids, thereby preventing differentiation. Extending our investigations to the rhizosphere, we uncover a much more diverse root mycobiome for either species when growing in the environment with nutrient-poor soils.
- Our finding suggests that both biotic and abiotic factors contribute to the distribution and delimitation of *Dactylorhiza* sibling allopolyploids. Altogether, our results indicate that plasticity can simultaneously promote diversification and homogenization of lineages, influencing the establishment and persistence of recurrently formed allopolyploid species.

Keywords: constitutive gene expression divergence, *Dactylorhiza*, fungal community, gene expression plasticity, marsh orchids, polyploidy.

Running title: Expression plasticity in marsh orchids

Plain language summary

A duplicated genome can enhance plasticity in polyploids, widening their environmental tolerance, but the long term effects of this plasticity remain unclear. Studying ecologically distinct marsh orchids, we show that sibling allopolyploids exhibit high plasticity with only very few fixed transcriptomic differences. Instead, we observe divergent mycobiomes associated with each orchid's environment, revealing a complex interplay in polyploid speciation.

Introduction

Polyploidy, or the duplication of the whole genome, introduces substantial genetic redundancy, which has been recognised as a diversifying force that may enhance the capacity to express different gene copies under varying environmental conditions (de Jong & Adams, 2023). This elevated genetic redundancy is expected to increase phenotypic plasticity (Mattingly & Hovick, 2023), the ability of an organism to adjust to environmental fluctuations and thus track a dynamically changing trait optimum beyond the limits of a fixed genotype (Schlichting & Pigliucci, 1998). However, increased phenotypic plasticity is often regarded as inherently costly as soon as the population has reached an adaptive optimum (DeWitt et al., 1998), in particular in extreme environments - habitats frequently occupied by polyploids and hybrids, which thereby avoid direct competition with diploid and other relatives. Moreover, some authors have argued that plasticity might slow down the evolutionary process, by reducing the efficiency of natural selection (Charlesworth et al., 1982) or even resulting in maladaptation (Arnold et al., 2019; Van Kleunen & Fischer, 2005). In stark contrast, others have documented adaptive benefits of plasticity, for example related to responses to abiotic or biotic stress (Auld & Relyea, 2011; Nicotra et al., 2015; Solé-Medina et al., 2022), whereas others have involved plasticity in initial stages of phenotypic evolution as a key mechanism fostering local adaptation (Mallard et al., 2020; Szukala et al., 2023).

Polyploid species typically originate as small founding populations that face additional challenges associated with minority cytotype disadvantage when sympatric with a majority of diploid cytotypes (Husband, 2000; Levin, 2002). Polyploids often form recurrently, increasing the genetic diversity of the new lineages (Soltis & Soltis, 1999). If they form polytopically or at different evolutionary times, they may encounter different environments and therefore selective pressures, which may lead to formation of distinct sibling polyploid species (Paun et al., 2007). Phenotypic plasticity can allow polyploids to overcome an initially depauperate gene pool by extending their habitable niche beyond their fixed genotypic capacity (Castillo et al., 2018), at the same time potentially enabling colonisation of environments that are free from homo- or heteroploid relatives (Hahn et al., 2012). However, increased plasticity may also facilitate gene flow between establishing polyploids and with their relatives, alleviating in the long term some of the founder effects of low genetic diversity. Studies addressing the role of plasticity in polyploid evolution often focus on the immediate implications of polyploid plasticity, such as whether plasticity is heightened in polyploids compared to diploids (Drunen & Johnson, 2022). However, the role of plasticity in the demographic dynamics between young sibling polyploid species remains poorly understood, despite its likely significant implications for how whole genome duplication contributes to biodiversity (Mattingly & Hovick, 2023).

To understand the interplay between plasticity, gene flow and ecological delimitation in polyploids, we use the marsh orchids *Dactylorhiza majalis* (Rchb.) P.F.Hunt & Summerh. and *D. traunsteineri* (Saut. ex Rchb.) Soó (Fig. 1a). These are sibling allopolyploids ($2n = 80$) resulting from two independent and unidirectional polyploidisation events between diploids *D. incarnata* (the paternal parent) and *D. fuchsii* (the maternal parent) (Brandrud et al., 2020; Pillon et al., 2007). Typically, *D. majalis* includes larger plants, with more flowers per inflorescence and broader leaves than *D. traunsteineri* (Fig. 1a). The two orchids have generalist pollination

syndromes, corresponding to a food-deception strategy (i.e., no reward is offered to pollinators) (Ostrowiecka et al., 2019). It is generally accepted that *D. traunsteineri* is the younger of the two sibling allopolyploids, and estimates of formation place both origins at different times in the recent half of the Quaternary (Brandrud et al., 2020; Hawranek, 2021; Nordström & Hedrén, 2009). Both sibling allopolyploids have a wide European distribution, with *D. majalis* having a more continuous distribution towards southern locations from the Pyrenees to southern Scandinavia, while *D. traunsteineri* occupies disjunct northerly locations in Scandinavia and the British Isles, with their distributions overlapping in the Alps. Gene flow between the two allopolyploids has been documented in areas of sympatry, with a tendency for more intense gene flow from *D. majalis* to *D. traunsteneri* (Balao et al., 2016; Brandrud et al., 2020). However, the sibling allopolyploids maintain distinct ecologies differentiated by soil chemistry, including nutrient content and water availability (Paun et al., 2011; Wolfe et al., 2023), and both species are found in their respective ecologies even at local scales in regions of sympatry. Despite the documented gene flow, previous studies uncovered ecologically-relevant transcriptomic differences between the allopolyploids (Wolfe et al. 2023), accompanied by a stable epigenetic divergence correlated with eco-environmental variables (Paun et al., 2010).

We investigate leaf and root gene expression of *D. majalis* and *D. traunsteineri* at two localities in the Austrian Alps, where the sibling allopolyploids coexist in sympatry and experience varying levels of gene flow (Balao et al., 2016; Brandrud et al., 2020). Using a reciprocal transplantation design, we grow individuals from each species in the environment of the other to understand how much their ecological preference is genetically encoded versus how much is underpinned by plasticity. This approach enables us to identify genes with fixed expression differences between the two species, representing genes which have become canalised (have adapted to the environment of each respective species and are no longer plastic), and avoid the homogenising effects of gene flow in sympatric populations. Comparing the proportion and function of plastic and fixed genes in admixed populations can enable us to understand the basis that delimits the ecological ranges of *D. majalis* and *D. traunsteineri*, and the proportion of the gene space (Dodsworth et al., 2015) involved in this delimitation.

We also investigate the potential for fungal communities to reinforce species boundaries, utilising the populations of *D. majalis* and *D. traunsteineri* at St. Ulrich and Kitzbuhel to test the consistency of these profiles across independent localities. Mycorrhizal fungi provide vital support, especially during early stages of the developmental cycle of orchids, and can ameliorate effects of nutrient deficiencies in host plants (Bhantana et al., 2021; Saboor et al., 2021). However the broader role of mycorrhizal fungi in determining host plant success and, in general, species boundaries remains poorly understood. Leveraging the capture of contaminants during RNA-seq experiments, we perform fungal taxonomic profiling of root RNA-seq data from native and transplanted samples to test whether the environments of *D. traunsteineri* and *D. majalis* differed in fungal community profiles. Differences in soil fungal communities may represent an ecologically relevant distinction between the distinct allopolyploid habitats (Wolfe et al., 2023), suggesting that symbiotic organisms could play a role in the delimitation of species boundaries.

This work uses *in situ* measurements of gene expression, functional response to different environments, and fungal community profiling to investigate the consequences of

transgressing ecological optima between two sibling allopolyploid species. Our final aim is to provide a deeper understanding of how resilience to environmental heterogeneity interplays with ecological adaptation to maintain species boundaries between sibling allopolyploid marsh orchids in the face of gene flow.

Methods

Study sites

Two localities where the two allopolyploids exist within a few hundred metres from each other have been selected for transplantation experiments. The first locality is near Lake Schwarzsee, north-west of Kitzbühel (Tyrol, Austria), which is the type locality of *D. traunsteineri* (Foley, 1990). In this area, the populations of *D. majalis* and *D. traunsteineri* are separated by a region lacking *Dactylorhiza* orchids, with the shortest distance between them being approximately 100m. Here, *D. majalis* grows in shady places under trees. *Dactylorhiza traunsteineri* plants are found in the open marshland closer to the lake, primarily on elevated and thus dryer locations (hummocks), nevertheless soil humidity is higher than in the *D. majalis* habitat. In general, the soil specific for *D. majalis* populations is significantly richer in most macro- and micronutrients as compared to *D. traunsteineri* (Wolfe et al., 2023). However, at the specific site in Kitzbuhel, the soil shows the opposite trend for phosphorus content (Fig. 1b).

The second locality is situated south of Pillersee, northwest of St. Ulrich am Pillersee (Tyrol, Austria). Here the two species are also separated by an area (shortest distance ca 200m) that lacks either species, but it hosts rare individuals of the diploid parent *D. incarnata*. Within this area, the two allopolyploids are also separated by a ca 50m wide tree stand. Both species at St Ulrich am Pillersee occupy very humid soils, but the soil chemistry at this site agrees with the characteristic difference observed across their range (Fig. 1b). Previous genetic analyses have shown that gene flow between the two species is more prominent at Kitzbuhel, whereas admixture is also present to some extent at St. Ulrich am Pillersee, at least partly via bacrosses with diploid *D. incarnata* (Balao et al., 2016; Brandrud et al., 2020).

Reciprocal transplantation and RNA sequencing

Ten wild individuals each of *D. majalis* and *D. traunsteineri* were randomly selected at each of the two localities. For each locality and species, five individuals were transplanted and grown in the environment of the other species, and five individuals were transplanted and grown in their own environment to control for transplantation effects. *Dactylorhiza* plants are perennials and grow a new tuber each growing season to support growth in the following season (Wolfe et al., 2023), therefore plants were sampled after two growing seasons to eliminate as much as possible the effects of the original environment. Leaf and root tissues were collected at each locality in the morning and stored in RNAlater (Sigma-Aldrich) at 4°C overnight, before transferring to -80°C. At collection, the transplanted plants have been entirely removed from the sites.

Figure 1. The study system and experiment design. (a) The study species: t, *Dactylorhiza traunsteineri* and m, *D. majalis* growing at one of the transplantation sites at St. Ulrich am Pillersee, Austria. T and M show impressions of the two environments at the same locality. (b) Divergent preference for selected soil characteristics between *D. traunsteineri* (blue) and *D. majalis* (yellow) across multiple European populations (data from Wolfe et al., 2023). With filled circles two measurements per species are indicated at the experimental site at Kitzbuhel, and with triangles at the site at St. Ulrich am Pillersee, both Austria. (c) Scheme of main comparisons made in differential expression (DE) testing. As above, lower case letters indicate sample species, superscript capital letters indicate the environment in which the sample was experimentally grown. ↔ indicates test of differential expression between sample groups, ∩ indicates intersection of genes found to be differentially expressed between sample groups.

Total RNA was extracted using the mirVana Isolation Kit (Thermo Fisher Scientific) according to manufacturer's instructions, and quantitated using a NanoDrop ND-1000 Spectrophotometer (Thermo Scientific), using the wavelength ratio of A260/280 to estimate

purity. A Plant RNA Pico kit was used on a 2100 Bioanalyzer (Agilent Technologies) to confirm the quality of the RNA isolates. DNA digestion was performed with DNase I (Promega), to remove any DNA contamination. Ribosomal RNA was depleted using the RiboZero Plant Kit for RNA-Seq (Illumina) according to manufacturer's instructions. RNA was fragmented by hydrolysis for 2 minutes at 94° and cDNA synthesis was performed with random hexamers following the TruSeq Stranded RNA kit protocol (Illumina). To achieve strand specificity, the second strand was synthesised with dUTPs. The libraries were sequenced using directional pair-end sequencing on an Illumina NovaSeq instrument at the Next Generation Sequencing Facility of the Vienna BioCenter Core facilities (VBCF, www.viennabiocenter.org/vbcf). The RNA-seq data are available from GenBank BioProject XXXX.

Few accessions did not survive during the experiment, whereas some accessions used have been later genetically identified as heavily introgressed between the two species or backcrossed towards the diploid *D. incarnata* (the latter at the locality of St. Ulrich). Few additional accessions have been excluded either because of insufficient quality of the RNA extraction (i.e., RNA integrity number lower than six across multiple RNA extractions) or of the data obtained (i.e., showing significantly more genes with zero counts than the rest of the accessions). Across both allopolyploids, 16 leaf and 17 root samples from Kitzbuhel have been retained for the bioinformatic analyses, and 13 leaf and 14 root samples from St Ulrich.

Genetic clustering

Quality filtered and adapter trimmed reads were mapped to the *D. incarnata* v.1.0 reference genome (Wolfe et al., 2023) using its .gff annotations and STAR v2.7.11a (Dobin et al., 2013) in two-pass mode, excluding any reads which mapped to more than ten positions. To evaluate the genetic structure of the randomly selected accessions, we used ANGSD v0.941 (Korneliussen et al., 2014) for estimating genotype likelihoods, as this approach is suitable for RNA-seq data (Szukala et al., 2023), which is expected to include highly variable sequencing depth among genes and exons. For this purpose the mapped bam files of the two tissues for each accession were merged with SAMtools v1.7 (Li et al., 2009). ANGSD was further run based on a SAMtools model (-GL 2), inferring the major/minor alleles (-doMajorMinor 1) and estimating allele frequencies (-doMaf 1). In the first round we retained SNPs that were identified outside annotated repeat regions, and which had a p-value lower than 2e-6 (-SNP_pval 2e-6), a minimum mapping quality of 20 (-minMapQ 20), a minimum base quality of 20 (-minQ 20), with data for at least 80% of individuals (-minInd 26) and a minor allele frequency corresponding to two accessions (-minMaf 0.06). In a second run we retained from the resulting variants only one per 2-kb window, aiming to reduce linkage disequilibrium between the markers. Covariance matrices computed from the obtained genotype likelihoods were used for principal components analyses (PCA) using pcANGSD v0.99 (Meisner & Albrechtsen 2018) and were plotted in R.

Differential expression analysis

Uniquely mapped reads were summarised per tissue type across the coding sequence, excluding introns, for each sample using featureCounts from Rsubread v2.0.4 (Liao et al.,

2019). Prior to differential expression analysis, the counts matrix was filtered to exclude low expression genes, retaining only those expressed at greater than one count per million in at least three samples, corresponding to the number of retained biological replicates in the smallest group in the analyses. DESeq2 v1.36.0 (Love et al., 2014) was used to identify differentially expressed genes (DEGs), normalising counts using median of ratios. To avoid incorporating signals of population structure and history, DEGs were detected for each locality (St Ulrich or Kitzbuhel) separately. Each test was performed to compare either the same species in different environments, or the two species in the same environment.

First, we identified constitutive gene expression differences, defined as DEGs between the sibling allopolyploids consistently identified in either environment, by finding genes which are significantly ($p < 0.05$, LogFC > 1.5) up- or downregulated between species in both environments i.e. the intersection of DEGs found in comparisons m^M vs t^M and m^T vs t^T (where m stands for *D. majalis*, t for *D. traunsteineri*, superscript M for the *D. majalis* environment and superscript T for that of *D. traunsteineri*; see also "constitutive DE" comparison in Fig. 1c). To determine whether there is a significant overlap of DEGs between each comparison (i.e. whether we find more constitutive DEG than we expect by chance) we use the super exact test implemented in the SuperExactTest package (Wang et al., 2015) in R , using the number of genes included in the DESeq2 analyses as the background set from which constitutively DEGs are sampled.

Then we identified genes which showed plastic expression patterns (significantly differentially expressed, $p < 0.05$, LogFC > 1.5) in each species and locality when transplanted to the environment of the opposite species, by quantifying DEGs between t^M and t^T for *D. traunsteineri*, and between m^T and m^M for *D. majalis* (see "*D. traunsteineri* plastic" and "*D. majalis* plastic" comparisons respectively in Fig. 1c). Functional enrichment analyses were performed on the resultant groups of genes using the Bioconductor package topGO v2.48.0 (Alexa & Rahnenfuhrer, 2023) using the weight01 algorithm and all genes in the reference genome as background set.

To focus on species-specific patterns of constitutive and plastic gene expression, Z-score was calculated for significantly enriched functional terms to indicate the difference in number of genes mapped to GO terms or GO terms mapped to GOSlim terms between *D. majalis* and *D. traunsteineri*, a higher absolute number reflecting greater directionality towards one species or another. Z-score was calculated for number of genes mapping to GO terms as (#genes^{majalis} - #genes^{traunsteineri}) / (genes^{majalis} + genes^{traunsteineri}) , and for proportion of GO terms mapping to GO slim terms as (%GOterms^{majalis} - %GOterms^{traunsteineri}) / (%GOterms^{majalis} + %GOterms^{traunsteineri}).

Fungal taxonomic abundance analysis

We performed community profiling of fungal taxa in the RNA-seq root samples of *D. majalis* and *D. traunsteineri* plants grown in the two environments using MiCoP (LaPierre et al., 2019), a mapping based approach for classification and abundance of viral and fungal sequences. MiCoP uses a rapid read mapping approach to achieve more granular classification estimates which are more sensitive to low-abundance species than comparable methods such

as kmers or megablast (Altschul et al., 1990). Paired end reads from each root sample were mapped with bwa mem v0.7.11 (Li et al., 2009) to the fungal RefSeq database accessed 7th November 2017 (Robbertse & Tatusova, 2011) included with the MiCoP release, followed by computation of fungal abundances from the alignment file. Quantification of diversity was performed with Shannon Index in the R package vegan v2.6.4 (Oksanen et al., 2009), calculating the Shannon index on relative abundance of fungal orders identified in each sample. Statistically significant differences in the Shannon diversity indices of the *D. majalis* and *D. traunsteineri* environments was tested per locality using a Welch Two Sample T-Test. Differential association of specific fungal taxa with the environment of either *D. majalis* or *D. traunsteineri* was analysed using LDA effect size (LEfSe, Segata et al., 2011). Features were scaled to sum to 10,000, using a p-value of 0.05 for both Kruskal-Wallis and Wilcoxon tests, and an effect size of discriminative features threshold of LDA > 2.

Results

Global effects of transplantation

Leaf tissue libraries consisted on average of 65 ± 30 million pairs of reads, whereas root libraries, of 107 ± 18 million. Of these, 43.9% and, respectively, 44.2% mapped uniquely to the *D. incarnata* reference genome (see Supplementary Table S1 for further details).

After excluding low confidence variants, those associated with repetitive regions, and keeping only putatively unlinked loci, we retained 315,846 polymorphic positions for investigating the genetic structure of accessions with a PCA. The first axis of the PCA explaining 4.2% of the variation (Supplementary Fig. 1) separates *D. traunsteineri* accessions from St. Ulrich from the rest, whereas the second axis (representing 1.4% of the variation) separates *D. traunsteineri* accessions from Kitzbühel from the rest. In line with previous results (Balao et al. 2016, Brandrud et al. 2020) *D. majalis* accessions from either locality form a more cohesive genetic cluster, whereas *D. traunsteineri* individuals separate very clearly by locality. Finally, whereas *D. traunsteineri* shows no genetic separation by environment, the samples of *D. majalis* show a minor tendency of separation by environment on the second PCA axis. As the individuals for the experiment were selected by chance, this slight separation may track back to a somewhat different expression rate of this species between the two environments, resulting in slightly different genotype likelihoods.

We further used expression-based PCAs to visualise the global transcriptional profiles of the accessions, separated by tissue type and locality (Fig. 2 A and C). The expression profiles tended to group samples first by species, and then by environment. This effect was particularly evident in root tissue, which showed a clear separation of species. Furthermore, the PCAs separate *D. traunsteineri* individuals grown in native versus non-native environments to a greater extent than for *D. majalis*, a trend that is especially pronounced in leaves.

Figure 2. Variability of expression profiles of *Dactylorhiza majalis* and *D. traunsteineri* across locality, tissue and environment. **A, C** PCA plots for leaf and root data, respectively, separately per locality (K, Kitzbühel; S, St. Ulrich). Points represent biological replicates of *D. majalis* and *D. traunsteineri* (rectangles and circles respectively) grown in *D. majalis* or *D. traunsteineri* environments (symbols with blue or yellow lines, respectively). **B, D** Numbers of genes up- (green bars) or down-regulated (orange bars) between *D. majalis* (yellow symbols) and *D. traunsteineri* (blue symbols) and environments (yellow line, *D. majalis* environment; blue line, *D. traunsteineri* environment). Left side bars show the number of DE genes at Kitzbühel (K), and the right side bars at St. Ulrich am Pillersee (S). The number of symbols in the experimental set up in the centre does not necessarily correspond to the actual analysed accessions per group.

Differential expression tests indicated that the two allopolyploids have more distinct profiles in the *D. majalis* environment (hereafter M environment), compared to the nutrient-poor habitat of *D. traunsteneri* (hereafter T environment), and this pattern was highly consistent between the localities and tissue types (Fig. 2B and D). In leaves and roots, respectively, 978 and 2,066 genes were differentially regulated between the allopolyploids in Kitzbuhel in the M environment, and 857 and 923 in St. Ulrich (Supplementary Table 2). However, only 143 and 501, and, respectively 163 and 512 genes were differentially expressed for the same comparisons in the T environment.

On the other hand, *D. traunsteineri* exhibited a greater number of DEGs in leaf tissue between native and non-native environments at both localities compared to *D. majalis* under the same conditions (Fig. 2B), however this trend was visible in root tissue only at St. Ulrich (Fig. 2D).

Constitutive and plastic expression

Root tissue had more genes expressed (i.e., with at least one count per million in at least three individuals) overall compared to leaf tissue: 35,251 and 34,844 genes were expressed in root tissue in Kitzbuhel and St. Ulrich respectively, compared to 30,808 and 28,872 genes in leaf tissue for the same localities. Of these, only 30 and 29 genes in leaf tissue were constitutively differentially expressed between the two allopolyploids in Kitzbuhel and St. Ulrich respectively, and 51 and 63 genes in roots for the same localities (Fig. 3). All four comparisons showed a significantly greater number of constitutively expressed genes than expected by chance ($p < 0.05$, super exact test, Supplementary Tables 2-4). Electron transport during photosynthesis, pH regulation and transmembrane transport appear to be enriched among constitutively differentially expressed genes between the allopolyploids in leaves, whereas in roots detoxification activity, carbohydrate transmembrane transport, and nucleotide and nucleoside metabolism (Fig. 4; Supplementary Table 5). Functional terms differentially regulated between the allopolyploids in the M environment only included cell wall biogenesis and defense response genes that are downregulated in the leaves of *D. traunsteineri*, and photosynthesis and signalling pathway that are upregulated in the same conditions. In turn, in the T environment, the leaves of *D. majalis* upregulate some photosynthesis genes. In roots, in the M environment several components of transmembrane transport differ between the two allopolyploids, whereas in the T environment apart from differences in transmembrane and water transport between the allopolyploids, some genes involved in response to heat and response to nitrogen compound are downregulated, and growth is upregulated in *D. majalis* (Fig. 4; Supplementary Table 5).

Figure 3. A Number of genes differentially expressed between *Dactylorhiza majalis* and *D. traunsteineri* in both environments (pink), in the *D. majalis* environment only (yellow), and in the *D. traunsteineri* environment only (blue). **B** Log fold change in gene expression between *Dactylorhiza majalis* and *D. traunsteineri* in the *D. traunsteineri* environment (Y-axis) and between *D. majalis* and *D. traunsteineri* in the *D. majalis* environment (X-axis). *D. majalis* was arbitrarily set as the base condition, therefore positive values indicate upregulation in *D. traunsteineri* relative to *D. majalis* and negative values indicate upregulation of *D. majalis* relative to *D. traunsteineri*. Genes shown in colour are designated differentially expressed (FDR < 0.05).

The environmental differences between the native habitats of *D. majalis* and *D. traunsteineri* elicit species specific plastic responses in either species when arriving in the alternative environment (see *D. traunsteineri* plastic and *D. majalis* plastic, respectively, in Figure 1). In general, root tissues showed more transcriptional plasticity than leaves after transplantation of each species to the alternative environment (except *D. traunsteineri* in Kitzbühel), and both tissues tended to upregulate more genes than downregulate (except the roots of *D. traunsteineri*; Fig. 2B and D; Supplementary Table 2). The great majority of differentially regulated genes between environments were unique to a species and locality (Fig. 5). *Dactylorhiza traunsteineri* leaf tissue showed the most overlap in plastic genes between Kitzbuhel and St. Ulrich of 17.6%, however most other comparisons showed fewer than 10% of plastic genes were replicated across both localities.

Figure 4. Top five GO terms significantly enriched (Fisher's exact test p < 0.05) in genes that are differentially expressed between *Dactylorhiza majalis* and *D. traunsteineri* in both species' environments (constitutive DEGs, pink panel), genes which are differentially expressed only in only the *D. majalis* environment (yellow panel) and genes which are differentially expressed only in the *D. traunsteineri* environment (blue panel). GO term categories are coloured by Z-score, indicating the directionality of gene expression underlying the enrichments, GO terms coloured in red contain more genes upregulated in *D. majalis*, GO terms coloured in blue contain more genes upregulated in *D. traunsteineri*. Results for leaf tissue samples are shown in the left panel, results for root tissue samples in the right panel. Differentially expressed genes for comparisons in Kitzbuhel and St. Ulrich are combined prior to functional enrichment analysis. Functional enrichment was performed on differentially expressed genes in each category (FDR < 0.05) using TopGO using the weight01 algorithm.

Figure 5. Venn diagram of plastic DE genes in each species (m, *D. majalis*; t, *D. traunsteineri*) when transplanted to the alternative environment (M, *D. majalis* environment; T, *D. traunsteineri* environment) (FDR < 0.05, log fold change > 1.5) within the same locality (K, Kitzbuhl; S, St. Ulrich). Labels follow Fig 1b and 1d.

To better understand the functional context of plasticity in *D. majalis* and *D. traunsteineri*, we performed GO term enrichment analysis to identify significantly overrepresented functional categories (Fisher's exact test p < 0.05) in genes exhibiting plasticity in their expression pattern (log fold change > 1.5, FDR < 0.05 in a single species grown in non-native versus native environment). To emphasise species-specific plastic responses we focussed on functional categories which tended to be found disproportionately in either *D. majalis* or *D. traunsteineri,* calculating Z-score on proportion of GO terms from each species mapping to GOSlim terms and excluding functional categories with a Z-score of less than 0.4, and thus deemed to be found in roughly equal proportions in both species (see methods, for a full list of enriched functional terms see Supplementary Table 6). GOSlim terms disproportionately found to be enriched in *D. majalis* plastic gene expression were signal transduction, anatomical structure morphogenesis, cellular homeostasis and growth, and those in *D. traunsteineri* were DNA metabolic process, the response to abiotic stimulus (including light, ionizing radiation, heat and water deprivation), and photosynthesis. GO terms classified under these GOSlim terms are shown in Fig. 6.

Figure 6. GO terms significantly enriched (Fisher's exact test p < 0.05) in genes differentially expressed in each species upon transplantation to the environment of the opposite species, by tissue and locality. GO terms enriched in upregulated genes are shown in upper panels, and downregulated in lower panels. Only GO terms mapped in different proportions to respective GOSlim terms between *D. majalis* and *D. traunsteineri* are visualised, showing GO terms with a Z-score of > 0.4 (see methods, for a full list of enriched GO terms see Supplementary Table 6).

Photosynthetic genes, including light response, and genes related to sensing are upregulated in *D. traunsteineri* leaves grown in the M environment, as were genes responding to hyperosmotic salinity stress and double strand break repair (Fig. 6). In turn, at both localities it consistently downregulated in the same conditions genes involved in response to heat, response to phosphate starvation and innate immune response. The leaves in *D. majalis* showed no enrichments, except for downregulated genes when grown in the T environment at Kitzbuhel, for which growth and regulation of growth through auxin and cytokinin activated pathways were overrepresented.

In root tissues, enrichments for *D. traunsteineri* upregulated genes were obtained only at Kitzbuhel, and they were related to mitosis and DNA replication, and systemic acquired resistance genes when grown in the M environment. *D. traunsteineri* roots further

downregulated genes related to phosphate starvation, water deprivation, and salt and heat stress, as well as genes related to innate immune response. When grown in the T environment at Kitzbühel, *D. majalis* root tissue upregulated homeostasis of glucose and metal ions cadmium and zinc, the formation of secondary shoots, and genes related to the serine/threonine kinase transmembrane receptor pathway. *D. majalis* downregulated genes related to multidimensional cell growth in root tissue when grown in the T environment at St. Ulrich.

Fungal taxonomic profiles of native and transplanted root samples

We sought to characterise the fungal species associated with the roots of *Dactylorhiza majalis* and *D. traunsteineri* because mycorrhizal and other fungi play an important role as determinants of fitness through orchid life cycles (Perotto & Balestrini, 2024; H. N. Rasmussen, 1990; H. N. Rasmussen & Rasmussen, 2009). Mapping RNA-seq reads to the Fungal RefSeq database with MiCoP identified fungal community profiles which differed the most between the environments of *D. majalis* or *D. traunsteineri*, rather than by species (Fig. 7).

Figure 7. Percent abundance of fungal orders identified in root RNA-seq per sample. Fungal orders which were unclassified or present at a proportion of less than 1% are labelled "Other". Transplantation of *D. majalis* and *D. traunsteineri* to the environment of the opposite species had the effect of shifting the fungal community profile of root tissue to that of the surrounding environment, rather than root tissue reflecting native fungal community profiles.

The M environment was characterised by the dominance of the order Wallemiales, a small group of xerophilic fungi in Basidiomycota (Zalar et al., 2005). While this taxon was also often also well represented in root tissues from the T environment, it's lower incidence compared to the alternative environment was accompanied by a higher mean relative abundance of non-Wallemiales fungal taxa (46.8% and 19.7% mean relative abundance of non-Wallemiales fungal taxa in the M environment in *D. majalis* and *D. traunsteineri* respectively,

compared to 74.4% and 55.7% in the T environment for the same species). The remaining taxonomic composition of the fungal community profiles included taxa which are known to be mycorrhizal partners to some orchids, such as Russulales (Paduano et al., 2011), Mortierellales (Herrera-Rus et al., 2020), Polyporales (Downing et al., 2020), and Hypocreales (Jiang et al., 2019), as well as orders containing potentially pathogenic fungal species such as Glomerellales (Baroncelli et al., 2018).

Alpha diversity as measured by Shannon index (Fig. 8) showed a clear distinction in the diversity and evenness of fungal taxa in the M environment compared to the T environment, the latter showing near-consistently higher diversity for both species at both localities. Higher fungal diversity in the T environment compared to M was replicated across both Kitzbühel (M mean = 1.02, T mean=1.55) and St. Ulrich (M mean = 0.55, T mean=1.28). However a statistically significant difference in Shannon Index was found only between individuals at M and T environments in St. Ulrich (Welch t-test, p=0.049), with no significant difference identified in Kitzbühel (Welch t-test, p=0.075).

LEfSe analysis (Supplementary Fig. 2) confirmed that Wallemiales was the only order significantly discriminative of the M environment (LEfSe LDA=3.14, p=0.0092). Wallemiales showed a higher mean relative abundance in the M environment, with between 49.4% and 90.3% mean relative abundance of Wallemiales regardless of species, compared to between 18.4% and 43.4% in the T environment. In contrast, three orders, Mortierellales (LEfSe LDA=2.44, p=0.0011), Spizellomycetales (LEfSe LDA=2.48, p=0.0010) and Tremellales (LEfSe LDA=2.21, p=0.022), were significantly discriminative of the T environment. The two most significantly discriminative orders of the T environment, Mortierellales and Spizellomycetales, had a mean relative abundance of between ca. 5 and 10% in *D. majalis* transplants and *D. traunsteineri* native individuals in both St. Ulrich and Kitzbühel. In contrast, these orders were found only in Kitzbühel in the M environment at a mean relative abundance of over 2%, with a maximum mean relative abundance of ca. 4%. Only one order (Tremellales, LEfSe LDA=2.32, p=0.0022) was found to be discriminative of the St. Ulrich and Kitzbuhel localities when testing for differential abundance using locality as the main class and environment as a subclass, suggesting that the M and T environments were the predominant drivers of fungal community differences and that this effect is relatively stable across the two localities.

Figure 8. Alpha diversity of each environment (that of *Dactylorhiza majalis* or *D. traunsteineri*, denoted M and T respectively) and each locality (Kitzbuhel or St. Ulrich, left and right panels respectively) measured using Shannon Index. Shannon index was calculated using relative abundance of fungal order, and is shown across samples from both species per locality and environment as circled points (*D. majalis* in yellow and *D. traunsteineri* in blue).

Discussion

Dominant plastic responses are punctuated by fixed transcriptional differences between species

Ecological plasticity is an important phenomenon that allows species to extend their genetically encoded trait optima, enabling them to withstand environmental fluctuations (Fox et al., 2019) or colonise new ranges (Richards et al., 2006). Polyploid plasticity is expected to be extensive, but it is often examined in the context of neopolyploid establishment, as it plays a critical role in overcoming obstacles such as reduced initial genetic diversity, minority cytotype exclusion and competition with diploids (Anneberg et al., 2023; Husband, 2000). Polyploids form recurrently, which enriches their genetic pool, but it can also lead to the development of distinct lineages, starting from different diploid populations, and each following independent evolutionary trajectories. The formation of stable polyploid species relies, among others, on the reinforcement of reproductive barriers to prevent genetic assimilation by more established

cytotypes. However, adaptive introgression of genes can also increase the fitness of establishing polyploid taxa (Chapman & Abbott, 2010; Han et al., 2015). In this context, ecological plasticity can allow polyploids to colonise and establish themselves in novel environments, but also tolerate environments inhabited by related taxa where genetic exchange can occur, thus potentially exposing establishing polypoids to opposing evolutionary forces.

To explore the further implications of ecological plasticity for repeated polyploid formation, we conducted reciprocal transplantation with *Dactylorhiza majalis* and *D. traunsteineri.* These sibling allopolyploids occupy distinct habitats (meadows with mesic soils versus very low nutrient fens), an ecological divergence that has been previously shown to be associated with distinct features of leaf chemistry, light harvesting, photoprotection, nutrient transport and stomata activity between the two allopolyploids (Wolfe et al., 2023). Our goal here was to quantify the plastic range of each species relative to the amount of fixed expression differences accumulated between them. We found very few constitutive DEGs, with more such genes identified in roots compared to leaves, and this consistently among the two localities investigated. The higher number of constitutive DEGs in roots aligns with the widespread differences in soil nutrient profiles between the specific environments of the two allopolyploids (Fig. 1; see also Paun et al., 2011; Wolfe et al., 2023), suggesting that the below-ground environment exerts greater divergent selective pressure on *D. majalis* and *D. traunsteineri*.

Rather than finding many constitutive DEGs, most DEGs between *D. majalis* and *D. traunsteineri* were environment-specific (Fig. 3). Notably, many more DEGs between the allotetraploids were found in the M environment than in the T. Our results further show that *D. traunsteineri* undergoes in general greater transcriptional remodelling in the alternative environment, than the vice versa situation. More widespread and competitive species like *D. majalis* (Paun et al., 2011) are thought to have higher levels of plasticity (Hiatt & Flory, 2020) than those with narrow ecological preferences such as *D. traunsteineri*. The transplantation of *D. majalis* to the T environment may fall more within its tolerated range, and hence require fewer changes. Alternatively or in addition, as the oldest of the two allopolyploids, *D. majalis* may be further along the transcriptional diploidization process.

Adaptation to the nutritionally depauperate environment of *D. traunsteineri* requires significant metabolic modifications (Wolfe et al., 2023). If the basis for these modifications is rather plastic, the transplantation of *D. traunsteineri* to the *D. majalis* environment may allow it to return to a transcriptional baseline to make use of the newly available resources, a possibility that is reflected in the functions of plastic genes (Fig. 6 and discussed later). Plasticity can mitigate unfavourable interspecific interactions (Otte et al., 2016), providing a buffer that prevents competitive exclusion or genetic swamping, thereby facilitating species coexistence (Hess et al., 2022). Polyploid plasticity, especially along environmental clines, is important for achieving the ecological speciation required for successful polyploid establishment (Shimizu-Inatsugi et al., 2017). The heterogeneous soil types occupied by *D. majalis* and *D. traunsteineri* likely served as the foundation for the ecological divergence between them.

Transcriptional plasticity allowing *D. traunsteineri* to survive in nutritionally poor environments, characterised by low competition, may be rewarded by allowing it to successfully establish as a young allopolyploid, developing genetic isolation away from the swamping effects of hybridisation when grown in sympatry with its siblings and diploid parents. While global

transcriptional profiles do not always clearly discriminate *D. majalis* and *D. traunsteineri* (Fig. 2A and 2C), the small number of constitutive DEGs may represent evidence of nascent differentiation in response to habitat-specific conditions.

Sibling allopolyploids plastically alter nutrient acquisition, photosynthetic capacity and osmotic regulation in the alternative habitat

Growing each sibling allopolyploid in the habitat of the other stimulated remodelling of functions reflecting their previously described habitat differences such as soil chemistry and nutrient availability (Wolfe et al., 2023); *D. traunsteineri* downregulated genes concerned with cellular response to phosphate starvation in leaves and roots, suggesting an alleviation of pressure in nutrient availability when grown in the *D. majalis* environment.

While upregulation of nutrient starvation functional terms is not seen in *D. majalis*, terms involving root system architecture remodelling and zinc homeostasis are upregulated in *D. majalis* root tissue (Supplementary Table 6, Fig. 6), reflecting a suite of processes responsive to nutrient starvation and abiotic stress (Bechtaoui et al., 2021; Shao et al., 2020). Root architecture remodelling increases root surface area and thus access to nutrients (Kiba & Krapp, 2016; Sánchez-Calderón et al., 2005), while phosphate limitation has been shown to be a key modulator of zinc bioavailability, forming a complex homeostatic interaction in plant responses to nutrient deficiency (Bouain et al., 2014; Lay-Pruitt et al., 2022).

Consistent with differences in water availability between the environments of *D. majalis* and *D. traunsteineri,* the latter shows a broad response to osmotic pressure, downregulating genes involved with water deprivation and salt stress, and upregulating genes annotated to hyperosmotic salinity response. Osmotic regulation is required to maintain efficient stomatal conductance (Bandurska, 2022) and is an important modulator of root system depth and breadth (Deak & Malamy, 2005), and the preponderance of *D. traunsteineri* plastic genes annotated to such terms may reflect the extent of the osmotic remodelling usually required in its native environment. To a lesser extent, *D. majalis* root tissue showed signs of plastically adjusting to an osmotically altered environment, upregulating important abiotic stress sensitive functions such as calcium-mediated signalling (Wilkins et al., 2016), as well as functions modulating glucose homeostasis, a key osmoprotectant (Ahmad et al., 2020). Wolfe et al. (2023) showed a lower energetic efficiency of photosynthesis in *D. traunsteineri* compared to *D. majalis*, and a higher rate of photooxidative regulation through quenching. Leaf tissue of *D. traunsteineri* grown in the *D. majalis* environment shows plastically reduced levels of gene expression in pathways known to alleviate oxidative stress and maintain efficient photosynthesis such as glutathione (Müller-Schüssele et al., 2020) and L-ascorbate (Chen et al., 2021) (Supplementary Table 6). Moreover, plastically upregulated genes are enriched for photsynthetic and photomorphongenic functions, indicating a plastic increase in *D. traunsteineri* photosynthetic efficiency upon release from pressures in its native environment.

Interestingly, up- and downregulated genes in *D. traunsteineri* root tissues were enriched in some photosynthesis related terms. Photosynthesis tends to be inhibited under low nutrient availability (Farhat et al., 2016; Mu & Chen, 2021), however due to roots being non-photosynthetic tissue, study of photosynthesis regulation in root tissue is sparse. Kang et al.

(2014) identified photosynthetic genes downregulated in the root tissue of *Arabidopsis thaliana* during phosphorous starvation, speculating that phosphorous depletion may stimulate aberrant photosynthesis to produce elevated levels of reactive oxygen species (ROS). Both nutrient deficiency (Ahanger et al., 2017; Kim et al., 2010) and photosystem I of the photosynthetic apparatus (Furutani et al., 2020) have been linked to elevated levels of ROS. The response to ROS appears to be diminished in *D. traunsteineri* when grown in the non-native *D. majalis* environment, as indicated by the downregulation of genes enriched in L-ascorbic acid and glutathione pathways, known for their role in ROS scavenging (Akram et al., 2017).

Interspecific fungal community differences

Whereas fungal reads were extracted from the *Dactylorhiza* root RNA-seq data, it is not clear what proportion of the sequence reads originate from genuine orchid mycorrhizal partners rather than bystander fungi present in the surrounding rhizosphere. Nevertheless, we find a consistent difference in the mycobiome associated to the roots of the sibling allopolyploids; diversity measures show higher alpha diversity in the *D. traunsteineri* environment (Fig. 8) regardless of species or locality compared with the *D. majalis* environment, which was dominated by Wallemiales taxa and exhibited smaller proportions of taxa well represented in the *D. traunsteineri* environment such as Spizellomycetales and Mortierellales. Three orders were discriminant of the *D. traunsteineri* environment (Mortierellales, Spizellomycetales and Tremellales) while Wallemiales was the only fungal order which was significantly discriminant of the *D. majalis* environment. This result is interesting given the limitation of resources such as nitrogen and phosphorus in the *D. traunsteineri* environment, which may prompt adapted plants to cultivate relationships with a more diverse set of specialist mycorrhizal partners to alleviate such nutrient deficiencies (Kohler et al., 2015; Wyatt et al., 2014).

Numerous environmental factors are important for successful symbiotic germination of orchid seeds with their mycorrhizal partners, and studies in *D. majalis* have found this process to be highly temperature dependent (H. Rasmussen et al., 1990). Interestingly, climatic niches for *D. traunsteineri* and *D. majalis* were found to differ in particular according to temperature seasonality and mean temperature of the driest quarter (Wolfe et al., 2023), suggesting key factors involved in symbiosis-dependent germination success differ between *D. majalis* and *D. traunsteineri*. The study presented here used adult plants to perform reciprocal transplantations, therefore individuals tested had germinated in native soil, and it is not known whether germination success of each sibling allopolyploid is modulated by the fungal community profile of their respective environments. Further studies are needed to conclusively determine whether fungal partners play a role in the delimitation of *D. majalis* and *D. traunsteineri* native ranges, and more broadly whether such interactions help drive ecological divergence in nascent species.

Conclusions

Hybridization and polyploidy are rich sources of biodiversity in the genus *Dactylorhiza*, but consequences of these processes can have contradictory effects. We show that divergent

ecological preferences in nascent species *D. majalis* and *D. traunsteineri* are circumvented by transcriptomic plasticity, impeding the build-up of fixed transcriptomic differences. Rather, we show that the nutritionally depauperate soil of *D. traunsteineri* harbours a more diverse community of fungal taxa than the nutritionally abundant soil of *D. majalis.* Taken with the broader role the rhizosphere plays in orchid germination and nutrient acquisition, we argue that divergent symbiotic interactions at key life stages may play a delimiting role in the distribution of this pair of sibling *Dactylorhiza* species, and merits further investigation.

Acknowledgements

We thank Juliane Baar, Richard Bateman, Marie Brandrud, Mark Chase, Mikael Hedren, Marcel Hirsch, Marie Huber and Daniela Paun for their contribution to the ideas and results presented here. Sequencing was performed at the Vienna BioCenter Core Facilities (VBCF; https://www.viennabiocenter.org/). Computational resources were provided by the Life Science Compute Cluster (LiSC) of the University of Vienna. Regional county administrations in Austria are acknowledged for issuing necessary permits.

Funding information

This research was funded by the Austrian Science Fund (FWF) through the START grant (https://doi.org/10.55776/Y661) to O.P. Katie Emelianova was funded through the Marie Skłodowska-Curie grant "StrucRadiation" (https://doi.org/10.3030/101066772).

Data availability statement

The raw Illumina sequencing data will be deposited on NCBI SRA (GenBank BioProject PRJNAXXXX, accessions YYYY) before the paper will be accepted for publication. The table of counts and the bioinformatics scripts are available from GitHub (https://github.com/katieemelianova/dactylorhiza/blob/main/dactylorhiza_DE.R).

References

Ahanger, M. A., Tomar, N. S., Tittal, M., Argal, S., & Agarwal, R. M. (2017). Plant growth under water/salt stress: ROS production; antioxidants and significance of added potassium under such conditions. *Physiology and Molecular Biology of Plants*, *23*(4), 731–744. https://doi.org/10.1007/s12298-017-0462-7

Ahmad, F., Singh, A., & Kamal, A. (2020). Osmoprotective Role of Sugar in Mitigating Abiotic

Stress in Plants. In *Protective Chemical Agents in the Amelioration of Plant Abiotic Stress* (pp. 53–70). John Wiley & Sons, Ltd. https://doi.org/10.1002/9781119552154.ch3

Akram, N. A., Shafiq, F., & Ashraf, M. (2017). Ascorbic Acid-A Potential Oxidant Scavenger and Its Role in Plant Development and Abiotic Stress Tolerance. *Frontiers in Plant Science*, *8*. https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2017.00613

Alexa, A., & Rahnenfuhrer, J. (2023). *Gene set enrichment analysis with topGO*. https://doi.org/doi:10.18129/B9.bioc.topGO,

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Anneberg, T. J., O'Neill, E. M., Ashman, T.-L., & Turcotte, M. M. (2023). Polyploidy impacts population growth and competition with diploids: Multigenerational experiments reveal key life-history trade-offs. *New Phytologist*, *238*(3), 1294–1304. https://doi.org/10.1111/nph.18794

Arnold, P. A., Kruuk, L. E. B., & Nicotra, A. B. (2019). How to analyse plant phenotypic plasticity in response to a changing climate. *The New Phytologist*, *222*(3), 1235–1241. https://doi.org/10.1111/nph.15656

Auld, J. R., & Relyea, R. A. (2011). Adaptive plasticity in predator-induced defenses in a common freshwater snail: Altered selection and mode of predation due to prey phenotype. *Evolutionary Ecology*, *25*(1), 189–202. https://doi.org/10.1007/s10682-010-9394-1

Balao, F., Tannhäuser, M., Lorenzo, M. T., Hedrén, M., & Paun, O. (2016). Genetic differentiation and admixture between sibling allopolyploids in the Dactylorhiza majalis complex. *Heredity*, *116*(4), Article 4. https://doi.org/10.1038/hdy.2015.98

Bandurska, H. (2022). Drought Stress Responses: Coping Strategy and Resistance. *Plants*, *11*(7), 922. https://doi.org/10.3390/plants11070922

Baroncelli, R., Sukno, S. A., Sarrocco, S., Cafà, G., Le Floch, G., & Thon, M. R. (2018). Whole-Genome Sequence of the Orchid Anthracnose Pathogen Colletotrichum orchidophilum. *Molecular Plant-Microbe Interactions®*, *31*(10), 979–981. https://doi.org/10.1094/MPMI-03-18-0055-A

Bechtaoui, N., Rabiu, M. K., Raklami, A., Oufdou, K., Hafidi, M., & Jemo, M. (2021). Phosphate-Dependent Regulation of Growth and Stresses Management in Plants. *Frontiers in Plant Science*, *12*. https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2021.679916

Bhantana, P., Rana, M. S., Sun, X., Moussa, M. G., Saleem, M. H., Syaifudin, M., Shah, A., Poudel, A., Pun, A. B., Bhat, M. A., Mandal, D. L., Shah, S., Zhihao, D., Tan, Q., & Hu, C.-X. (2021). Arbuscular mycorrhizal fungi and its major role in plant growth, zinc nutrition, phosphorous regulation and phytoremediation. *Symbiosis*, *84*(1), 19–37. https://doi.org/10.1007/s13199-021-00756-6

Bouain, N., Shahzad, Z., Rouached, A., Khan, G. A., Berthomieu, P., Abdelly, C., Poirier, Y., & Rouached, H. (2014). Phosphate and zinc transport and signalling in plants: Toward a better understanding of their homeostasis interaction. *Journal of Experimental Botany*, *65*(20), 5725–5741. https://doi.org/10.1093/jxb/eru314

Brandrud, M. K., Baar, J., Lorenzo, M. T., Athanasiadis, A., Bateman, R. M., Chase, M. W., Hedrén, M., & Paun, O. (2020). Phylogenomic Relationships of Diploids and the Origins of Allotetraploids in Dactylorhiza (Orchidaceae). *Systematic Biology*, *69*(1), 91–109. https://doi.org/10.1093/sysbio/syz035

Castillo, J. M., Gallego-Tévar, B., Figueroa, E., Grewell, B. J., Vallet, D., Rousseau, H., Keller, J., Lima, O., Dréano, S., Salmon, A., & Aïnouche, M. (2018). Low genetic diversity contrasts with high phenotypic variability in heptaploid Spartina densiflora populations invading the Pacific coast of North America. *Ecology and Evolution*, *8*(10), 4992–5007. https://doi.org/10.1002/ece3.4063

Chapman, M. A., & Abbott, R. J. (2010). Introgression of fitness genes across a ploidy barrier. *New Phytologist*, *186*(1), 63–71. https://doi.org/10.1111/j.1469-8137.2009.03091.x

Charlesworth, B., Lande, R., & Slatkin, M. (1982). A Neo-Darwinian Commentary on Macroevolution. *Evolution*, *36*(3), 474–498. https://doi.org/10.2307/2408095

Chen, X., Zhou, Y., Cong, Y., Zhu, P., Xing, J., Cui, J., Xu, W., Shi, Q., Diao, M., & Liu, H. (2021). Ascorbic Acid-Induced Photosynthetic Adaptability of Processing Tomatoes to Salt Stress Probed by Fast OJIP Fluorescence Rise. *Frontiers in Plant Science*, *12*. https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2021.594400

de Jong, G. W., & Adams, K. L. (2023). Subgenome-dominant expression and alternative splicing in response to Sclerotinia infection in polyploid Brassica napus and progenitors. *The Plant Journal*, *114*(1), 142–158. https://doi.org/10.1111/tpj.16127

Deak, K. I., & Malamy, J. (2005). Osmotic regulation of root system architecture. *The Plant Journal*, *43*(1), 17–28. https://doi.org/10.1111/j.1365-313X.2005.02425.x

DeWitt, T. J., Sih, A., & Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends in Ecology & Evolution*, *13*(2), 77–81. https://doi.org/10.1016/S0169-5347(97)01274-3

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Dodsworth, S., Leitch, A. R., & Leitch, I. J. (2015). Genome size diversity in angiosperms and its influence on gene space. *Current Opinion in Genetics & Development*, *35*, 73–78. https://doi.org/10.1016/j.gde.2015.10.006

Downing, J. L., Liu, H., McCormick, M. K., Arce, J., Alonso, D., & Lopez-Perez, J. (2020). Generalized mycorrhizal interactions and fungal enemy release drive range expansion of orchids in southern Florida. *Ecosphere*, *11*(8), e03228. https://doi.org/10.1002/ecs2.3228

Drunen, W. E. V., & Johnson, M. T. J. (2022). Polyploidy in urban environments. *Trends in*

Ecology & Evolution, *37*(6), 507–516. https://doi.org/10.1016/j.tree.2022.02.005

Farhat, N., Elkhouni, A., Zorrig, W., Smaoui, A., Abdelly, C., & Rabhi, M. (2016). Effects of

magnesium deficiency on photosynthesis and carbohydrate partitioning. *Acta*

Physiologiae Plantarum, *38*(6), 145. https://doi.org/10.1007/s11738-016-2165-z

Foley, M. J. Y. (1990). An assessment of populations of Dactylorhiza traunsteineri (Sauter) S06

in the British Isles and a comparison with others from Continental Europe. *Watsonia*,

18(2), 153–172.

Fox, R. J., Donelson, J. M., Schunter, C., Ravasi, T., & Gaitán-Espitia, J. D. (2019). Beyond

buying time: The role of plasticity in phenotypic adaptation to rapid environmental

change. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

374(1768), 20180174. https://doi.org/10.1098/rstb.2018.0174

Furutani, R., Ifuku, K., Suzuki, Y., Noguchi, K., Shimakawa, G., Wada, S., Makino, A., Sohtome,

T., & Miyake, C. (2020). Chapter Five—P700 oxidation suppresses the production of

reactive oxygen species in photosystem I. In T. Hisabori (Ed.), *Advances in Botanical*

Research (Vol. 96, pp. 151–176). Academic Press.

https://doi.org/10.1016/bs.abr.2020.08.001

Hahn, M. A., Kleunen, M. van, & Müller-Schärer, H. (2012). Increased Phenotypic Plasticity to

Climate May Have Boosted the Invasion Success of Polyploid Centaurea stoebe. *PLOS*

ONE, *7*(11), e50284. https://doi.org/10.1371/journal.pone.0050284

Han, T.-S., Wu, Q., Hou, X.-H., Li, Z.-W., Zou, Y.-P., Ge, S., & Guo, Y.-L. (2015). Frequent

introgressions from diploid species contribute to the adaptation of the tetraploid

Shepherd's purse (Capsella bursa-pastoris). *Molecular Plant*, *8*(3), 427–438.

https://doi.org/10.1016/j.molp.2014.11.016

Hawranek, A.-S. (2021). *The impact of recurrent origins and gene flow on the genetic structure*

of allopolyploid marsh orchids (Dactylorhiza, Orchidaceae).

https://doi.org/10.25365/thesis.70360

Herrera-Rus, I., Pastor, J. E., & Juan, R. (2020). Fungal colonization associated with phenological stages of a photosynthetic terrestrial temperate orchid from the Southern Iberian Peninsula. *Journal of Plant Research*, *133*(6), 807–825. https://doi.org/10.1007/s10265-020-01225-9

Hess, C., Levine, J. M., Turcotte, M. M., & Hart, S. P. (2022). Phenotypic plasticity promotes species coexistence. *Nature Ecology & Evolution*, *6*(9), Article 9. https://doi.org/10.1038/s41559-022-01826-8

Hiatt, D., & Flory, S. L. (2020). Populations of a widespread invader and co-occurring native species vary in phenotypic plasticity. *New Phytologist*, *225*(1), 584–594. https://doi.org/10.1111/nph.16225

Husband, B. C. (2000). Constraints on polyploid evolution: A test of the minority cytotype exclusion principle. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *267*(1440), 217–223. https://doi.org/10.1098/rspb.2000.0990

Jiang, J., Zhang, K., Cheng, S., Nie, Q., Zhou, S.-X., Chen, Q., Zhou, J., Zhen, X., Li, X. T., Zhen, T. W., Xu, M., Hsiang, T., Sun, Z., & Zhou, Y. (2019). Fusarium oxysporum KB-3 from Bletilla striata: An orchid mycorrhizal fungus. *Mycorrhiza*, *29*(5), 531–540. https://doi.org/10.1007/s00572-019-00904-3

Kang, J., Yu, H., Tian, C., Zhou, W., Li, C., Jiao, Y., & Liu, D. (2014). Suppression of Photosynthetic Gene Expression in Roots Is Required for Sustained Root Growth under Phosphate Deficiency. *Plant Physiology*, *165*(3), 1156–1170. https://doi.org/10.1104/pp.114.238725

Kiba, T., & Krapp, A. (2016). Plant Nitrogen Acquisition Under Low Availability: Regulation of Uptake and Root Architecture. *Plant and Cell Physiology*, *57*(4), 707–714. https://doi.org/10.1093/pcp/pcw052

Kim, M. J., Ciani, S., & Schachtman, D. P. (2010). A Peroxidase Contributes to ROS Production during Arabidopsis Root Response to Potassium Deficiency. *Molecular Plant*, *3*(2), 420–

427. https://doi.org/10.1093/mp/ssp121

Kohler, A., Kuo, A., Nagy, L. G., Morin, E., Barry, K. W., Buscot, F., Canbäck, B., Choi, C.,
Cichocki, N., Clum, A., Colpaert, J., Copeland, A., Costa, M. D., Doré, J., Floudas, D.,
Gay, G., Girlanda, M., Henrissat, B., Herrmann, S., … Martin, F. (2015). Convergent
losses of decay mechanisms and rapid turnover of symbiosis genes in mycorrhizal
mutualists. *Nature Genetics*, *47*(4), Article 4. https://doi.org/10.1038/ng.3223

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics*, *15*(1), 356. https://doi.org/10.1186/s12859-014-
0356-4

LaPierre, N., Mangul, S., Alser, M., Mandric, I., Wu, N. C., Koslicki, D., & Eskin, E. (2019).
MiCoP: Microbial community profiling method for detecting viral and fungal organisms in
metagenomic samples. *BMC Genomics*, *20*(5), 423. https://doi.org/10.1186/s12864-019-
5699-9

Lay-Pruitt, K. S., Wang, W., Prom-u-thai, C., Pandey, A., Zheng, L., & Rouached, H. (2022). A
tale of two players: The role of phosphate in iron and zinc homeostatic interactions.
Planta, *256*(2), 23. https://doi.org/10.1007/s00425-022-03922-2

Levin, D. A. (2002). *The Role of Chromosomal Change in Plant Evolution*. Oxford University
Press. https://doi.org/10.1093/oso/9780195138597.001.0001

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence
Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352

Liao, Y., Smyth, G. K., & Shi, W. (2019). The R package Rsubread is easier, faster, cheaper
and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids
Research*, *47*(8), e47. https://doi.org/10.1093/nar/gkz114

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

https://doi.org/10.1186/s13059-014-0550-8

Mallard, F., Nolte, V., & Schlötterer, C. (2020). The Evolution of Phenotypic Plasticity in

Response to Temperature Stress. *Genome Biology and Evolution*, *12*(12), 2429–2440.

https://doi.org/10.1093/gbe/evaa206

Mattingly, K. Z., & Hovick, S. M. (2023). Autopolyploids of Arabidopsis thaliana are more

phenotypically plastic than their diploid progenitors. *Annals of Botany*, *131*(1), 45–58.

https://doi.org/10.1093/aob/mcab081

Mu, X., & Chen, Y. (2021). The physiological response of photosynthesis to nitrogen deficiency.

Plant Physiology and Biochemistry, *158*, 76–82.

https://doi.org/10.1016/j.plaphy.2020.11.019

Müller-Schüssele, S. J., Wang, R., Gütle, D. D., Romer, J., Rodriguez-Franco, M., Scholz, M.,

Buchert, F., Lüth, V. M., Kopriva, S., Dörmann, P., Schwarzländer, M., Reski, R.,

Hippler, M., & Meyer, A. J. (2020). Chloroplasts require glutathione reductase to balance

reactive oxygen species and maintain efficient photosynthesis. *The Plant Journal: For

Cell and Molecular Biology*, *103*(3), 1140–1154. https://doi.org/10.1111/tpj.14791

Nicotra, A. B., Segal, D. L., Hoyle, G. L., Schrey, A. W., Verhoeven, K. J. F., & Richards, C. L.

(2015). Adaptive plasticity and epigenetic variation in response to warming in an Alpine

plant. *Ecology and Evolution*, *5*(3), 634–647. https://doi.org/10.1002/ece3.1329

Nordström, S., & Hedrén, M. (2009). Evolution, phylogeography and taxonomy of allopolyploid

Dactylorhiza (Orchidaceae) and its implications for conservation. *Nordic Journal of

Botany*, *27*(6), 548–556. https://doi.org/10.1111/j.1756-1051.2009.00548.x

Oksanen, J., Kindt, R., Legendre, P., Hara, B., Simpson, G., Solymos, P., Henry, M., Stevens,

H., Maintainer, H., & Oksanen@oulu, jari. (2009). *The vegan Package*.

Ostrowiecka, B., Tałałaj, I., Brzosko, E., Jermakowicz, E., Mirski, P., Kostro-Ambroziak, A.,

Mielczarek, Ł., Lasoń, A., Kupryjanowicz, J., Kotowicz, J., & Wróblewska, A. (2019).

Pollinators and visitors of the generalized food-deceptive orchid Dactylorhiza majalis in North-Eastern Poland. *Biologia*, *74*(10), 1247–1257. https://doi.org/10.2478/s11756-019-00285-0

Otte, T., Hilker, M., & Geiselhardt, S. (2016). Phenotypic plasticity of mate recognition systems prevents sexual interference between two sympatric leaf beetle species. *Evolution*, *70*(8), 1819–1828. https://doi.org/10.1111/evo.12976

Paduano, C., Rodda, M., Ercole, E., Girlanda, M., & Perotto, S. (2011). Pectin localization in the Mediterranean orchid Limodorum abortivum reveals modulation of the plant interface in response to different mycorrhizal fungi. *Mycorrhiza*, *21*(2), 97–104. https://doi.org/10.1007/s00572-010-0315-5

Paun, O., Bateman, R. M., Fay, M. F., Hedrén, M., Civeyrel, L., & Chase, M. W. (2010). Stable Epigenetic Effects Impact Adaptation in Allopolyploid Orchids (Dactylorhiza: Orchidaceae). *Molecular Biology and Evolution*, *27*(11), 2465–2473. https://doi.org/10.1093/molbev/msq150

Paun, O., Bateman, R. M., Fay, M. F., Luna, J. A., Moat, J., Hedrén, M., & Chase, M. W. (2011). Altered gene expression and ecological divergence in sibling allopolyploids of Dactylorhiza (Orchidaceae). *BMC Evolutionary Biology*, *11*(1), 113. https://doi.org/10.1186/1471-2148-11-113

Paun, O., Fay, M. F., Soltis, D. E., & Chase, M. W. (2007). Genetic and epigenetic alterations after hybridization and genome doubling. *TAXON*, *56*(3), 649–656. https://doi.org/10.2307/25065849

Perotto, S., & Balestrini, R. (2024). At the core of the endomycorrhizal symbioses: Intracellular fungal structures in orchid and arbuscular mycorrhiza. *New Phytologist*, *242*(4), 1408–1416. https://doi.org/10.1111/nph.19338

Pillon, Y., Fay, M. F., Hedrén, M., Bateman, R. M., Devey, D. S., Shipunov, A. B., Van Der Bank, M., & Chase, M. W. (2007). Evolution and temporal diversification of western

European polyploid species complexes in *Dactylorhiza* (Orchidaceae). *TAXON*, *56*(4), 1185–1208. https://doi.org/10.2307/25065911

Rasmussen, H., Andersen, T. F., & Johansen, B. (1990). Temperature sensitivity of in vitro germination and seedling development of Dactylorhiza majalis (Orchidaceae) with and without a mycorrhizal fungus. *Plant, Cell & Environment*, *13*(2), 171–177. https://doi.org/10.1111/j.1365-3040.1990.tb01289.x

Rasmussen, H. N. (1990). Cell differentiation and mycorrhizal infection in Dactylorhiza majalis (Rchb. F.) Hunt & Summerh. (Orchidaceae) during germination in vitro. *New Phytologist*, *116*(1), 137–147. https://doi.org/10.1111/j.1469-8137.1990.tb00519.x

Rasmussen, H. N., & Rasmussen, F. N. (2009). Orchid mycorrhiza: Implications of a mycophagous life style. *Oikos*, *118*(3), 334–345. https://doi.org/10.1111/j.1600-0706.2008.17116.x

Richards, C. L., Bossdorf, O., Muth, N. Z., Gurevitch, J., & Pigliucci, M. (2006). Jack of all trades, master of some? On the role of phenotypic plasticity in plant invasions. *Ecology Letters*, *9*(8), 981–993. https://doi.org/10.1111/j.1461-0248.2006.00950.x

Robbertse, B., & Tatusova, T. (2011). Fungal genome resources at NCBI. *Mycology*, *2*(3), 142–160. https://doi.org/10.1080/21501203.2011.584576

Saboor, A., Ali, M. A., Danish, S., Ahmed, N., Fahad, S., Datta, R., Ansari, M. J., Nasif, O., Rahman, M. H. ur, & Glick, B. R. (2021). Effect of arbuscular mycorrhizal fungi on the physiological functioning of maize under zinc-deficient soils. *Scientific Reports*, *11*(1), Article 1. https://doi.org/10.1038/s41598-021-97742-1

Sánchez-Calderón, L., López-Bucio, J., Chacón-López, A., Cruz-Ramírez, A., Nieto-Jacobo, F., Dubrovsky, J. G., & Herrera-Estrella, L. (2005). Phosphate starvation induces a determinate developmental program in the roots of Arabidopsis thaliana. *Plant & Cell Physiology*, *46*(1), 174–184. https://doi.org/10.1093/pcp/pci011

Schlichting, C., & Pigliucci, M. (1998). *Phenotypic Evolution: A Reaction Norm Perspective*.

Sinauer.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., & Huttenhower, C.

(2011). Metagenomic biomarker discovery and explanation. *Genome Biology*, *12*(6),

R60. https://doi.org/10.1186/gb-2011-12-6-r60

Shao, C.-H., Qiu, C.-F., Qian, Y.-F., & Liu, G.-R. (2020). Nitrate deficiency decreased

photosynthesis and oxidation-reduction processes, but increased cellular transport,

lignin biosynthesis and flavonoid metabolism revealed by RNA-Seq in Oryza sativa

leaves. *PLoS ONE*, *15*(7), e0235975. https://doi.org/10.1371/journal.pone.0235975

Shimizu-Inatsugi, R., Terada, A., Hirose, K., Kudoh, H., Sese, J., & Shimizu, K. K. (2017). Plant

adaptive radiation mediated by polyploid plasticity in transcriptomes. *Molecular Ecology*,

26(1), 193–207. https://doi.org/10.1111/mec.13738

Solé-Medina, A., Robledo-Arnuncio, J. J., & Ramírez-Valiente, J. A. (2022). Multi-trait genetic

variation in resource-use strategies and phenotypic plasticity correlates with local climate

across the range of a Mediterranean oak (Quercus faginea). *New Phytologist*, *234*(2),

462–478. https://doi.org/10.1111/nph.17968

Soltis, D. E., & Soltis, P. S. (1999). Polyploidy: Recurrent formation and genome evolution.

Trends in Ecology & Evolution, *14*(9), 348–352. https://doi.org/10.1016/s0169-

5347(99)01638-9

Szukala, A., Bertel, C., Frajman, B., Schönswetter, P., & Paun, O. (2023). Parallel adaptation to

lower altitudes is associated with enhanced plasticity in Heliosperma pusillum

(Caryophyllaceae). *The Plant Journal*, *115*(6), 1619–1632.

https://doi.org/10.1111/tpj.16342

Van Kleunen, M., & Fischer, M. (2005). Constraints on the evolution of adaptive phenotypic

plasticity in plants. *New Phytologist*, *166*(1), 49–60. https://doi.org/10.1111/j.1469-

8137.2004.01296.x

Wang, M., Zhao, Y., & Zhang, B. (2015). Efficient Test and Visualization of Multi-Set

Intersections. *Scientific Reports*, *5*(1), Article 1. https://doi.org/10.1038/srep16923

Wilkins, K. A., Matthus, E., Swarbreck, S. M., & Davies, J. M. (2016). Calcium-Mediated Abiotic Stress Signaling in Roots. *Frontiers in Plant Science*, *7*. https://doi.org/10.3389/fpls.2016.01296

Wolfe, T. M., Balao, F., Trucchi, E., Bachmann, G., Gu, W., Baar, J., Hedrén, M., Weckwerth, W., Leitch, A. R., & Paun, O. (2023). Recurrent allopolyploidizations diversify ecophysiological traits in marsh orchids (Dactylorhiza majalis s.l.). *Molecular Ecology*, *32*(17), 4777–4790. https://doi.org/10.1111/mec.17070

Wyatt, G. A. K., Kiers, E. T., Gardner, A., & West, S. A. (2014). A Biological Market Analysis of the Plant-Mycorrhizal Symbiosis. *Evolution*, *68*(9), 2603–2618.

Zalar, P., Sybren de Hoog, G., Schroers, H.-J., Frank, J. M., & Gunde-Cimerman, N. (2005). Taxonomy and phylogeny of the xerophilic genus Wallemia (Wallemiomycetes and Wallemiales, cl. Et ord. Nov.). *Antonie van Leeuwenhoek*, *87*(4), 311–328. https://doi.org/10.1007/s10482-004-6783-x

Supplementary figures

Supplementary Figure 1. PCA plot based on genotypic variability expressed as genotype likelihoods in *Dactylorhiza majalis* and *D. traunsteineri* in the two localities tested (Kitzbühel and St. Ulrich am Pillersee). Symbols represent replicates of *D. majalis* (squares) and *D. traunsteineri* (circles) in native and transplanted habitats (according to the legend).

Supplementary Figure 2. Clades identified to be explanatory of the difference between fungal community profiles in *D. majalis* and *D. traunsteineri* environments using linear discriminant analysis effect size (LEFSe). Only clades with a minimum log(LDA) score of 3 and a Kruskal-Wallis p-value < 0.05 are shown.

(c) Expression Category   **Comparison**

Constitutive DE	$m^T \longleftrightarrow t^T \;\cap\; t^M \longleftrightarrow m^M$
D. traunsteineri plastic	$t^M \longleftrightarrow t^T$
D. majalis plastic	$m^M \longleftrightarrow m^T$

	Constitutive	Constitutive	
anion transmembrane transport	○	○	carbohydrate transmembrane transport
photosynthetic electron transport in photosystem II	●	●	detoxification
potassium ion transmembrane transport	○	●	seed dormancy process
regulation of intracellular pH	○	●	deoxyribonucleoside triphosphate catabolic process
sodium ion import across plasma membrane	○	●	pyrimidine deoxyribonucleotide metabolic process

	D. majalis only	D. majalis only	
cell wall organization	●	○	transmembrane transport
negative regulation of catalytic activity	●	●	cellular oxidant detoxification
cell wall biogenesis	●	●	oligopeptide transmembrane transport
xyloglucan metabolic process	●	○	response to wounding
carotene catabolic process	●	●	sesquiterpene biosynthetic process

	D. traunsteineri only	D. traunsteineri only	
microtubule–based process	●	○	transmembrane transport
photosynthetic electron transport in photosystem II	●	○	cell wall organization
negative regulation of phosphoprotein phosphatase activity	●	●	response to water
response to herbicide	●	●	response to hydrogen peroxide
regulation of protein serine/threonine phosphatase activity	●	●	water transport

Plastic Upregulated (left)

GO term	Kitzbuhel	St. Ulrich
photosynthesis, light harvesting in photosystem I	● D. traunsteineri	
hyperosmotic salinity response	● D. traunsteineri	
photosynthetic electron transport in photosystem II	● D. traunsteineri	
nuclear DNA replication		▲ D. traunsteineri
response to light stimulus		▲ D. traunsteineri
response to ionizing radiation		▲ D. traunsteineri
double−strand break repair via break−induced replication		▲ D. traunsteineri
mitotic recombination		▲ D. traunsteineri
hyperosmotic salinity response		▲ D. traunsteineri
reductive pentose−phosphate cycle		▲ D. traunsteineri
response to absence of light		▲ D. traunsteineri
regulation of photomorphogenesis		▲ D. traunsteineri
dbl−strand break repair via synth−dependent strand annealing		▲ D. traunsteineri
response to blue light		▲ D. traunsteineri
photosynthesis, light harvesting in photosystem I		▲ D. traunsteineri

Plastic Downregulated (left)

GO term	Kitzbuhel	St. Ulrich
cytokinin−activated signaling pathway	● D. majalis	● D. traunsteineri
auxin−activated signaling pathway	● D. majalis	● D. traunsteineri
unidimensional cell growth	● D. majalis	● D. traunsteineri
phosphorelay signal transduction system	● D. majalis	● D. traunsteineri
regulation of innate immune response	● D. traunsteineri	▲ D. traunsteineri
response to heat	● D. traunsteineri	▲ D. traunsteineri
cellular response to phosphate starvation	● D. traunsteineri	▲ D. traunsteineri
response to salt stress	▲ D. traunsteineri	▲ D. traunsteineri
innate immune response	▲ D. traunsteineri	▲ D. traunsteineri
response to heat	▲ D. traunsteineri	▲ D. traunsteineri
cellular response to phosphate starvation	▲ D. traunsteineri	▲ D. traunsteineri

Plastic Upregulated (right)

GO term	D. majalis	D. traunsteineri
transmembrane receptor protein serine/threonine kinase signaling pathway	● D. majalis	
cellular zinc ion homeostasis	● D. majalis	
cellular cadmium ion homeostasis	● D. majalis	
calcium−mediated signaling	● D. majalis	
cellular glucose homeostasis	● D. majalis	
secondary shoot formation	● D. majalis	
negative regulation of DNA−templated DNA replication		● D. traunsteineri
nuclear DNA replication		● D. traunsteineri
photosynthesis, light harvesting in photosystem I		● D. traunsteineri
mitotic DNA replication		● D. traunsteineri
systemic acquired resistance		● D. traunsteineri
DNA replication initiation		● D. traunsteineri
DSB repair via break−induced replication		● D. traunsteineri

Plastic Downregulated (right)

GO term	D. majalis	D. traunsteineri
multidimensional cell growth	▲ D. majalis	
response to salt stress	● D. traunsteineri	
response to osmotic stress	● D. traunsteineri	
cellular response to heat	● D. traunsteineri	
response to heat	● D. traunsteineri	
+ve reg. transcription from RNApolII prmtr in response to heat stress	● D. traunsteineri	
cellular response to phosphate starvation	▲ D. traunsteineri	
photosynthesis, light reaction	▲ D. traunsteineri	
response to heat	▲ D. traunsteineri	
response to water deprivation	▲ D. traunsteineri	
innate immune response	▲ D. traunsteineri	
+ve reg. transcription from RNApolI prmtr in response to heat stress	▲ D. traunsteineri	
cellular response to heat	▲ D. traunsteineri	

Legend:
● Kitzbuhel
▲ St. Ulrich
● D. majalis (yellow)
● D. traunsteineri (blue)