Revival of traditional agricultural systems – A multidisciplinary on-farm survey of maize-bean

intercropping reveals unexpected competition effects on bean.

Noa Vazeux-Blumental[1], Laura Mathieu[1], Théo Trabac[2], Carine Palaffre[3], Bernard Lagardère[3], Maryse

Carraretto[4], Cyril Bauland[1], Martine Le Guilloux[1], Christine Paysant Le Roux[5,6], José Caïus[5,6], Anne

Marmagne[7], Jérôme Enjalbert[1], Timothée Flutre[1], Edith Le Cadre[8], Virginie Parnaudeau[8], Daniel

Muller[2], Yvan Moënne-Loccoz[2], Domenica Manicacci[1], Maud I. Tenaillon[1]

[1] Université Paris-Saclay, INRAE, CNRS, AgroParisTech, GQE-Le Moulon, 91190, Gif-sur-Yvette,

France

[2] Université Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup, UMR5557 Ecologie Microbienne,

69622, Villeurbanne, France

[3] INRAE, Unité expérimentale du maïs, 40390, Saint Martin de Hinx, France

[4] Université de Toulouse, Université Toulouse, CNRS, EHESS, ENSFEA, INU Champollion, LISST,

31058, Toulouse, France

[5] Université Paris-Saclay, CNRS, INRAE, Université Evry, IPS2, 91190, Gif-sur-Yvette, France

[6] Université Paris Cité, CNRS, INRAE, IPS2, 91190, Gif-sur-Yvette, France

[7] INRAE, AgroParisTech, Université Paris-Saclay, IJPB, 78026, Versailles, France

[8] I NRAE, Institut Agro Rennes-Angers, SAS, 35042, Rennes, France

ORCID IDs

N.V.-B.: 0009-0001-1971-1019; L.M.: 0009-0003-4083-1298; C.P.: 0000-0002-8427-1726; C.B.: 0000-

0001-9916-2744; C.P.-L.: 0000-0003-2046-6492; J.C.: 0000-0001-8692-042X; A.M.: 0000-0003-3466-

9477; J.E.: 0000-0003-0408-2253; T.F.: 0000-0003-4489-4782; E.L.C.: 0000-0001-5859-5682; V.P.:

0000-0002-4442-6703; D.M.: 0000-0002-6619-4691; Y.M.-L.: 0000-0002-9817-1953; Do.M.: 0000-

0002-6779-113X; M.I.T.:0000-0002-0867-3678;

Correspondance

Domenica Manicacci, Université Paris-Saclay, INRAE, CNRS, AgroParisTech, GQE-Le Moulon,

91190, Gif-sur-Yvette, France

Email: domenica.manicacci@inrae.fr

Maud Tenaillon, Université Paris-Saclay, INRAE, CNRS, AgroParisTech, GQE-Le Moulon, 91190, Gif-

sur-Yvette, France

Email: maud.tenaillon@inrae.fr

Word count = 7,656

Number of tables = 1

Number of figures = 7

Supporting information = 14 supplementary figures; 13 supplementary datasets (supplied separately);

Societal Impact Statement

Cereal-legume intercropping is a promising strategy for sustainable agroecosystems that leverages the biological complementarities between species, reducing the need for chemical inputs while enhancing field biodiversity. Here, we focused on maize-bean intercropping, which is experiencing a revival in modern conventional agricultural settings. We found a trend towards an increased soil bacterial diversity in intercropping. But competition between the two crops dominated and primarily impacted yield and gene expression in beans. Despite lower yield, bean seeds exhibited greater size. Our study indicates that implementing variety selection of both species and adapting farming practices is essential to fully harness the potential of intercropping.

Summary

- Cereal-legume intercropping is a promising strategy for sustainable agroecosystems. The traditional intercropping of maize and bean is experiencing a revival in some modern agricultural settings, such as in southwestern France, where maize hybrids are intercropped with the commercialized Tarbais bean. We conducted on-farm surveys and a field assay to address the following questions: How do intercropping and sole cropping impact yield, nutrient uptake and rhizosphere bacterial assemblages? Do positive or negative interactions between maize and bean dominate in intercropping? What is the effect of intercropping on plant transcriptomics?

- We recorded farming practices, conducted yield and nutrient measurements, and characterized soil bacterial assemblages to compare sole-cropped maize and beans with intercropped plants. We also set up a controlled field assay to extend this comparison to plant gene expression differences.

- Intercropping was associated with a trend towards increased bacterial diversity. The cropping system significantly influenced agronomic traits, with frequent farm-by-cropping system interactions underscoring the critical role of farming practices. Competition dominated maize-bean intercropping, with 34 negative correlations among the 47 significant ones between maize and bean traits. This competition had effects on both yield and nutrition but primarily impacted beans, which produced fewer but bigger/heavier seeds. Transcriptomic results concurred with these findings revealing no differentially expressed genes in maize but 5,070 in beans under competition.

- Overall, our findings suggest that beneficial interactions between the two crops are hindered, underscoring the importance of carefully considering partner varieties and farming practices in the revival of traditional agricultural systems.

Keywords

agroecology, bacterial metabarcoding, cereal-legume mixture, competition, plant nutrition, RNAseq

1. INTRODUCTION

Conventional agriculture mainly relies on simplified cropping systems, with a single crop species grown per field (sole cropping) and environmental uniformization through the use of chemical inputs to ensure high yields. Given the current environmental challenges, there is a growing concern for more sustainable agroecosystems based on ecological regulations (Beillouin 2021; Fréville et al. 2022; Gaba et al. 2015). In this context, mixed intercropping (thereafter intercropping), where several crops are grown simultaneously with no distinct row arrangements, relies on biological complementarities and beneficial species interactions that diminish inputs requirement (Bedoussac et al. 2015). This is the case for cereal-legume intercrops whose success relies on their complementarity, which results primarily from the capacity of legumes (*Fabaceae*) to assimilate nitrogen fixed by symbiotic rhizobia. By mixing cereals (*Poaceae*) with legumes, the need for nitrogen-based fertilizers is reduced as compared to sole-cropped cereals. Cereals promote symbiotic nitrogen fixation as they deplete soil nitrogen accessible for legumes but also through their root exudation, stimulating nodulation and rhizobia symbiotic activity (Rodriguez 2020; Li et al. 2016a). Cereal-legume interactions can yield additional benefits. First, improved resource utilization may occur through complementarity of their aerial and root architecture. This has been demonstrated at the root level in maize-bean-squash intercropping (Zhang et al. 2014), where complementarity increases biomass production in N-deficient soils. Second, facilitative processes involving soil acidification by legume root exudates, which locally increases the availability of inorganic phosphorus in the soil, may benefit the cereal crop as exemplified in maize-faba bean (Li et al. 2007; Zhang et al. 2016), maize-chickpea (Li et al. 2004), wheat-faba bean (Li et al. 2016b), and wheat-common bean (Li et al. 2008) intercropping. Third, weed reduction can result from enhanced competitive abilities of the crop mixture, which decreases the need for herbicides (Poggio 2005). Fourth, transfer of N and P may be accrued between plant species interconnected through arbuscular mycorrhizal fungi (Johansen and Jensen 1996; Zhang et al. 2019). Fifth, the recruitment of plant growth-promoting rhizobacteria, such as phloroglucinol-producing *Pseudomonas* on cereal roots, can promote plant growth and provide control over various root pathogens (Mazzola 2004).

Although beneficial interactions between cereals and legumes are well-documented from an agronomic perspective, the genetic and molecular determinants of their interactions, as well as their impact on soil ecology, remain poorly explored (Becker et al. 2023). This is particularly true for the intercropping of maize with climbing common bean, where the maize stalk supports the climbing bean, thereby promoting its exposure to light. Yet this association is of major interest, for both economic and historical reasons. Economically, maize is the second most widely grown crop in the world and common bean is the most widely used legume for direct human consumption. Historically, both crops were domesticated in Mexico (Bitocchi et al. 2013; Matsuoka et al. 2002) and their association within the same cropping system dates back to 7,000 to 4,400 years BP (Zizumbo-Villarreal and Colunga-GarcíaMarín 2010). This cropping system, central to the iconic "milpa", featured maize, squash, and

beans—collectively known as the "three sisters"—and was the predominant multi-cropping subsistence system in Mesoamerica. Notably, the practice of intercropping maize and bean has spread beyond its region of origin, such as on small farms in sub-Saharan Africa (Nassary et al. 2020) and in various traditional European farming systems (Vazeux-Blumental et al. 2024). Thus, the intercropping of maize and bean appears to have been introduced and maintained as these crops spread beyond their centers of domestication.

While being abandoned in favor of sole cropping during the 20th century, this practice has persisted on small-scale subsistence farms in many parts of the world and has recently experienced a revival in some modern agricultural settings. An example is the Tarbais bean in southwestern France, where maize and bean were introduced in the 18th century and widely intercropped, becoming an essential part of the culinary heritage (Bonnain-Dulon and Brochot 2004). However, during the 1950s, the cultivation of beans considerably declined due to the intensification of agriculture and the introduction of maize hybrids, i.e. maize varieties mostly used for animal feed and developed through crosses of inbred lines, significantly increasing yield and supporting industrial farming. As beans were not suitable for mechanization, their cultivation around Tarbes (Hautes-Pyrénées area) decreased from approximately 12,000 hectares at the beginning of the 20th century to around 55 hectares by 1970 (Bonnain-Dulon and Brochot 2004). It was not until 1986 that the Tarbais bean was developed to diversify local agriculture and enhance farming profitability, thereby promoting a local crop used in traditional dishes, such as the cassoulet stew. Tarbais bean obtained a quality label and a protected geographic indication in 2000, which makes it an emblematic crop in the region, with high added value. Along with the increase in acreages of Tarbais bean, several farmers reintroduced intercropping with maize as an alternative to the use of plastic nets in sole-cropped beans (Figure 1).

The objective of our study was to gain a deeper understanding of the functioning and potential benefits of intercropping maize and common bean. We focused on the Tarbais-maize system, which represents a revival of a traditional farming system using modern varieties and, most often, conventional farming practices (notably relatively high levels of nitrogen fertilization). While these practices may not be ideal for leveraging beneficial interactions between crops (Fréville et al. 2022), we hypothesized that maize-bean intercropping could still offer benefits. Due to the lack of research on this particular intercropping, we adopted a descriptive multidisciplinary approach combining agronomy, microbial ecology, and transcriptomics to answer three main questions: How do intercropping and sole cropping impact yield, nutrient uptake and rhizosphere bacterial assemblages? Do positive or negative interactions between maize and bean dominate in intercropping at the phenotypic level? What is the effect of intercropping at plant transcriptomic level? We recorded farming practices, performed agronomic and nutrient measurements on plants collected from both intercropped and sole-cropped plots in farmers' fields, and we examined bacterial communities and corresponding soils in the rhizosphere. Because the detection of molecular signatures of intercropping requires a controlled environment with

150 replicates, we established a controlled field assay in which we compared both agronomic traits and gene
151 expression between sole crop and intercropping of Tarbais bean and two maize hybrids.
152

153 **2. MATERIALS AND METHODS**
154

155 **2.1 On-farm analyses in the Tarbes region**
156

2.1.1 Farms and farming practices
We focused on the Tarbes region (Figure S1) located in southwestern France. In order to compare sole
and intercropping, we identified multiple farms that practiced at least one of the two sole cropping
(maize or bean) in addition to intercropping, seven of which agreed to participate to this study. In this
region, most farmers use the recent "Label Rouge" bean variety Alaric, an inbred line derived from
crosses among local landraces registered in the European Community's varietal catalog in 1999. The
Lapujole variety, registered in 2014, was derived from Alaric by introgression of an anthracnose
resistance allele from scarlet runner bean (*Phaseolus coccineus*). Both varieties are therefore genetically
similar (hereafter we used Tarbais bean to designate both). Among the seven farmers included in our
study, five used the Tarbais bean and two farmers used local bean landraces of their own. Several maize
varieties were grown, most of which were modern hybrids (Dataset S1). A single farmer (F4) used a
local maize landrace. Only two farms (F6 and F7) practiced the three cropping systems (sole-cropped
bean, sole-cropped maize and maize-bean intercropping).

Farming practices in each farm were recorded (Dataset S1). The sowing dates for the different
farms ranged from April 15th to June 10th. Since manual harvest of beans requires wide space between
rows, sole-cropped maize was cultivated at a much higher density (8 plants per m^2) compared to both,
intercropped maize and bean (2.5 plants per m^2 for each of the species) and sole-cropped bean (2.5 plants
per m^2). Except for the organic farm F4, with low N-fertilization (50 kg N ha^{-1}) for sole-cropped maize
and no fertilization for intercropped maize and bean), N-fertilizer input was on average 224 N units ha^{-1}
for sole-cropped maize, 130 N units ha^{-1} for sole-cropped bean, and 135 N units ha^{-1} for intercropped
maize and bean. Farm F4 did not practice irrigation. For the other farms, irrigation along the growing
season was, on average, 181 mm for sole-cropped maize, 150 mm for sole-cropped bean, and 117 mm
for intercropped maize and bean.
180

2.1.2 Plant and soil sampling for agronomic analyses
All plant and soil samples were collected before harvest in October 2017, from 16 fields (one to three
fields per farm, Dataset S1). In each field, we randomly sampled three plots (located > 5 m apart) that
served as replicates in the analyses. Each plot consisted of 4 to 5 (sole cropping) or 7 to 10
(intercropping) plants each. Among the 16 fields, seven fields corresponded to intercropping (for a total
of 21 plots), five to maize sole cropping (for a total of 15 plots) and four to bean sole cropping (for a

total of 12 plots). Given that we collected maize and bean separately from the intercropping plots, these
48 plots led to a total of 69 plant samples of 4 to 5 pooled plants each (with the exception of one plot of
3 plants). In addition, we retrieved about 500 g of soil from each plot to determine soil properties.
190

2.1.3 Soil properties determination

We measured 13 variables on soil samples taken at 5-15 cm depth (Dataset S2). Analyses were
performed at the Laboratoire des sols d'Arras (INRAE) accredited by the French Accreditation
Committee (Cofrac). Samples were first gently dried at 40°C, then sieved and ground at 2 mm. Soil
texture was determined according to LAS-SOL-0302 procedure with 5-fraction granulometry without
decarbonation (g kg^{-1}). From this first sample, a second representative sample was obtained and further
ground at 250 µm (ISO 11464) for analyses requiring finer grinding. Finally, ammoniacal (Am, NH_4^+)
and nitric (Ni, NO_3^-) nitrogen were quantified on all fresh samples (i.e., without pooling samples;
Dataset S3), and samples from the same field, albeit different plots, were pooled for the subsequent
analyses (Dataset S3).

Sampling and sieving were carried out after destruction of the organic matter by hydrogen
peroxide (H_2O_2). The soil water pH was determined (soil:water ratio of 1:5) according to NF EN ISO
10390. The Metson Cationic Exchange Capacity (Metson 1957) was determined following the
procedure NF X 31-130. Total organic carbon and nitrogen were determined by heat combustion
(1000°C) on 250 µm particles obtained from grinding the 2-mm dried soil, with decarbonation
performed for the former. Calcium carbonate content was measured from CO_2 release after acidification
of the sample by HCl according to NF EN ISO 10693.
208

2.1.4 Plant phenotyping

All phenotypic and nutrient measures on plants were performed by pooling all plants from a same plot.
In addition to the number of plants per sample, we measured 22 (for maize) or 24 traits (for bean, Table
1). Note that we separated vegetative (including maize cobs and bean pod shells) from reproductive
(kernels for maize or seeds for beans, referred to as "propagules" in Table 1) parts. Propagules were
counted per plant, but also per ear for maize and per pod for bean – referred to as "sets" in Table 1). To
measure plant nutrient parameters related to nitrogen and carbon content, we ground the samples into a
fine and homogeneous powder. A subsample (0.8–1.2 mg) of each powder sample was assayed for total
N and C contents and ^{15}N abundance using the FLASH 2000 Organic Elemental Analyzer coupled to an
isotope ratio mass spectrometer (ThermoFisher Scientific, Waltham, MA). The N and C concentrations
(N% and C%) of each sample were expressed as mg (of N or C) per 100 mg of dry weight for both
vegetative tissues (NV, CV) and propagules (NP, CP). We also computed N and C content per plant both
for vegetative parts (NV_p, CV_p) and propagules (NP_p, CP_p) by multiplying NV, CV, NP, CP with

the dry weight per plant, which generated four additional variables. Collected plant phenotypic data are
given in Dataset S4.
224
2.1.5 Plant nutrient simulations
To assess the effect of intercropping on the nitrogen cycle, we used the N mineralization as a proxy
during presence of the intercropping in fields. Accordingly, we simulated soil nitrogen mineralization
(kg N.ha^{-1}) in the agricultural plots using Nsim (Parnaudeau et al. 2012). Nsim includes an agronomic
model that estimates the potential soil N mineralization according to the characteristics of the soil (Clivot
2019), and uses soil water content and daily temperatures recorded in the Tarbais region in 2017 to
calculate the effective N mineralization from organic matter. Soil water content is itself a function of
soil characteristics, weather and canopy conditions. We used as input parameters: calendar dates of
sowing and harvest as well as the irrigation and nitrogen supply (Dataset S1); because mineral N in the
soil was not measured at the beginning of the experiment, the previous harvest values were used as
default values at the beginning of the simulations. We also considered soil texture as determined by soil
analyses. Since bean parameters were not available for Nsim, we parametrized the model using default
parameters for pea (*Pisum sativum*).
238
2.1.6 Isotope analysis
The ^{15}N abundance was calculated as atom% (percentage of total N) and defined as
%^{15}N=100 × (^{15}N)/(^{15}N + ^{14}N) (Marmagne et al. 2022). We determined ^{15}N /^{14}N ratio (*d15N*) for the
vegetative parts (*d15NV*) and propagules (*d15NP*) of maize and bean using the ^{15}N abundance method
as discussed in Shearer and Kohl (1988), with the following equation:
$d15N = 1000[(\%^{15}N - \%^{15}N_{atm})/(\%^{15}N_{atm})]$
where %^{15}N$_{atm}$ is the percentage of ^{15}N in the atmosphere, a constant value of 0.3663.
The ^{13}C /^{12}C ratio (*d13C*) was calculated for the vegetative parts (*d13CV*) and propagules (*d13CP*) of
maize and bean using the following equation:
$d13C = 1000[(^{13}C /^{12}C)/(^{13}C /^{12}C_{atm})-1]$
where ^{13}C /^{12}C$_{atm}$ is the actual ratio of ^{13}C to ^{12}C of the atmosphere, a constant equal to 0.011142.
We estimated symbiotic nitrogen fixation of the bean from the *d15N* by calculating the percent
nitrogen derived from atmosphere (*d15Ndfa*) for the vegetative parts (*d15NVdfa*) and the propagules
(*d15NPdfa*) separately, using the following equation (Shearer et al. 1980):
$d15Ndfa = 100[(d15N_{non.fix} - d15N_{fix})/(d15N_{non.fix} - B)]$
where *d15N$_{non.fix}$* is the *d15N* of a non-fixing plant such as maize. We therefore estimated it from the
average value of maize *d15N* between the replicates of each farm. B = *d15Nfix* when plant N only
comes from the atmospheric N (when plants are grown in N-free soil). Here, we took the B value of
d15N common bean equal to -1.988. This value was proposed by Farid and Navabi (2015) as the

average of *d15N* measurements of a total of 20 randomly-selected bean genotypes (Peoples et al.
2002).
260
2.1.7 Statistical analysis on soil characteristics and plant phenotypic traits

We employed ANOVA (R package "stats") to examine whether the 13 soil property variables, simulated nitrogen mineralization (Nsim), irrigation and N fertilization (Nin) varied according to farm and cropping systems (sole-cropped bean, sole-cropped maize or maize-bean intercropping). Since cropping conditions were not replicated within farms, we were unable to test the interaction between those factors. Therefore, we utilized a linear model including farm and cropping system as fixed effects. Note that the farm effect incorporates differences in the plant varieties, in farming practices and in soil characteristics.

We assessed the plant phenotypes of the maize and bean separately, initially employing Principal Component Analysis (PCA) and ellipse plots for graphical analysis using the factoextra R package where ellipses represent the 95% confidence interval (https://github.com/kassambara/factoextra). Subsequently, we examined the farm and cropping system effects on plant phenotypes through ANOVA. To assess the interaction between farm and cropping system for each crop, we restricted our analysis to farms practicing both sole cropping and intercropping (F3, F5, F6, F7 for beans, and F1, F2, F4, F6, F7 for maize), and used the three plots per field as replicates to test the farm-by-cropping system interaction.

To address multiple testing, we used PCA plots to determine effective number of independent traits in the data sets as the number of principal components required to explain more than 90% of the variation in soil, bean and maize, respectively. Each *P* value was adjusted in accordance, i.e. multiplied by the corresponding number of components.

To investigate the type of interaction between maize and bean plants, we examined correlations between phenotypic traits of both species using intercropping data, correcting for multiple testing through False Discovery Rate (FDR; Benjamini and Hochberg 1995) with a threshold at 20%. Negative correlations are expected in case of competition between species, while positive correlations suggest beneficial interactions.

We did not compute the Land Equivalent Ratio, often used to compare production in intercropping versus sole cropping (Mead and Willey 1980), because the farming practices differed between cropping conditions.

2.1.8 Soil processing for metabarcoding

Metabarcoding analysis of the soil bacterial community was carried out using 39 composite samples from the three plots in each of the 16 fields. Non-rhizospheric samples were collected from each of the 16 fields, while rhizospheric samples were taken separately for each crop in the seven intercropped maize and bean fields (14 samples), in addition to those from the sole-cropped fields (9 samples). Composite samples of non-rhizosphere soil and rhizosphere soil (for each crop) were taken at 5-15 cm depth. For rhizosphere soil, 2 or 3 plants of each species were studied per plot (6-9 plants of each species

per field) and soil in contact with roots was taken from 2 to 3 different roots per plant. Therefore, soil from roughly 18 roots of each species was sampled per field (~1 g per composite sample) with a sterile scalpel. Likewise, non-rhizosphere soil was taken from 2 to 3 root-free soil samples in each of the 3 plots (~1 g non-rhizosphere soil per composite sample).

Each composite sample was placed in a 2 ml ZR BashingBead Lysis tube containing beads and ZR stabilization reagent (Zymo Research, Irvine, CA), and was mixed immediately in the field using a bead-beating TerraLyzer lysis system (Zymo Research) for 30 s. DNA extraction was carried out in the laboratory using the FastDNA kit (MP Biomedicals, Santa Ana, CA), following the manufacturer's instructions. We sequenced 16S rRNA genes using Illumina MiSeq technology (2 × 300 bp).

2.1.9 Operational Taxonomic Units assignation and analysis

Sequence trimming and quality control (removal of sequences < 150 bp or with ambiguous base calls) were performed by MR DNA (Shallowater, TX). The remaining sequences were denoised, Operational Taxonomic Units (OTUs; arbitrarily defined with a threshold of 3% divergence) identified, and chimeras removed. Final OTUs were taxonomically classified using BLASTn against RDPII (http://rdp.cme.msu.edu). OTU quantification and taxonomic assignation are given in Tables S5 and S6. We used datasets without singletons (i.e., sequences found a single time among the 39 sequenced samples) to generate rarefaction curves.

Data analysis was primarily conducted using the phyloseq R package (McMurdie and Holmes 2013). At each taxonomic level, we visualized the 10 most abundant taxa in the entire dataset using percentage normalization and assessed the proportion of these OTU sequences for each condition. For other analyses, we utilized the rarefy_even_depth function to normalize sample sizes based on the smallest sample size. We computed alpha diversity indices (Observed diversity, Chao1) that represent measures of biodiversity within individual samples. The Observed diversity index simply counts the number of unique taxa observed in a sample, while the Chao1 index estimates the true richness of the sample, taking into account the presence of rare taxa. These indices were calculated using the plot_richness function on the normalized data. Additionally, Unifrac distances, a measure of dissimilarity between microbial communities, were computed between samples and then visualized using Non-metric MultiDimensional Scaling (NMDS) plots. Permutational Multivariate Analysis of Variance (PERMANOVA) was employed to test differences among conditions using the adonis2 function from the vegan R package (https://github.com/vegandevs/vegan), with sample distances as input. The unique and common OTUs between conditions were determined on non-normalized data using Venn diagrams created with the limma package (Ritchie et al. 2015). Furthermore, variations among conditions were assessed using Between Class Analysis (BCA).

We relied on differential analysis based on OTU count using negative binomial (a.k.a. Gamma-Poisson) distribution and Wald significance tests to identify discriminant OTUs (with FDR Q value)

between sole cropped and intercropped conditions for maize and bean taken separately. This analysis was performed using the DESeq2 R package (https://github.com/thelovelab/DESeq2).

Subsequently, we compared OTU composition among soil types and farming practices using co-inertia analysis. The RV coefficient between the two datasets was computed, and its significance was tested using Monte-Carlo tests with 10,000 replicates. Both BCA and co-inertia analyses were conducted using the ade4 R package (https://adeverse.github.io/ade4/).

2.2 Experimental assays of maize-bean intercropping in Saclay

2.2.1 Experimental design and field management

To assess the effect of maize-bean intercropping at the transcriptomic level, we set up a controlled field experiment in Saclay in 2022 (France). A single bean genotype (B, the Tarbais variety Alaric) was associated with two maize hybrids, DG0801 (M1, INRAE hybrid used as a control variety for low-nitrogen trials in southwestern France) and LG30.275 (M2, commercial hybrid used by INRAE as a control variety in northern France). Each of the three genotypes (B, M1, M2) was grown in sole cropping along with the two possible intercropping (M1B and M2B) in a three-blocks randomized design. Each block therefore contained five randomized plots with 160 cm between plots and two stand types: sole and intercropping. Each plot encompassed two rows 5.2m length of 12 plants for sole cropping with 24 cm intra-row spacing and 80 cm inter-row spacing, and similarly for intercropping albeit with 12 maize and 12 beans per row. Bean and maize plants were sown at a similar density as practiced on-farm for intercropping, that is 2.36 plants m^{-2} for both sole cropping and intercropping conditions (meaning that plant density is doubled for intercropping plots).

The soil was prepared before sowing by adding nitrogen fertilization in line with the farming practices recommended by the Tarbais bean cooperative (residual nitrogen of 26 kg N ha^{-1} after winter and application after sowing of 75 kg N ha^{-1} of nitrogen in the form of 27% ammonium nitrate). Such nitrogen supply corresponded to a grain yield target of 30% of their potential for commercial maize hybrids. No other chemical treatment was applied. Sowing occurred on April 28th 2022 for maize. Beans were sown after maize emergence on May 24th 2022, and the seeds were inoculated with a mix of symbiotic rhizobia (LEGUMEFIX Phaseolus Inoculant, Legume Technology LTD, Nottingham, UK). Stakes were used for support in bean sole cropping. Note that we encountered a drought with only 94.5 mm of cumulated rainfall for the three first months of the growing season (188 mm 20 years average). Manual irrigation was applied once during the course of the experiment.

2.2.2 Sampling and phenotyping

Leaf sampling was performed on the morning of July 11[th], 2022, with a collection minimized to ensure physiological consistency across plants. This date coincided to the flowering stage for M2, an advanced vegetative stage for M1, and an early flowering stage in beans where nodules are likely most active

(Peña-Cabriales and Castellanos 1993). We pooled 1 leaf per plant for 10 randomly-selected plants per

plot, for a total of 21 samples (9 sole-cropping samples i.e. 3 plots × 3 blocks, and 12 intercropping

samples i.e. 2 plots × 3 blocks × 2 species; Dataset S7). For maize, we sampled 11th leaf, and for bean,

we chose the terminal leaflet of the last fully developed leaf on each plant. We retained 2 cm^2 of the

central portion of the blade without the midrib from the 10 leaves. Samples were flash-frozen and stored

at -80°C. Traits measured on maize and bean plants are given in Dataset S8.

2.2.3 Quantification of gene expression with QuantSeq 3'RNA-seq

Total RNAs were extracted from 100 mg of tissue with the Nucleospin RNA plant kit (Macherey-Nagel,

Düren, Germany) following the manufacturer instructions. Extracted RNAs were checked for purity and

quantity with a bioanalyzer (Agilent, Santa Clara, CA). Barcoded libraries were constructed using

QuantSeq 3'RNA-Seq Library Prep Kit FWD for Illumina (Lexogen, Vienna, Austria) with Unique

Molecular Identifiers (UMIs) to reduce amplification biases, and sequencing was conducted by the

POPS transcriptomics platform (IPS2, Université Paris-Saclay). QuantSeq 3'RNA-seq produce a single

read per transcript close to the 3′ end of polyadenylated RNAs. From 3.4 to 17.4 million 75-bp reads

were obtained per sample (Dataset S7).

Pre-processing steps and read counts included deduplication of reads based on UMIs with UMI-

tools v1.0.1 (Smith et al. 2017). Reads with UMI base quality score falling below 10 were removed. We

performed removal of adapter sequences and quality trimming with BBduk from the BBmap suite

(v38.84, https://sourceforge.net/projects/bbmap/) with the options k=13 ktrim=r useshortkmers=t

mink=5 qtrim=r trimq=10 minlength=30. The nf-core/rnaseq pipeline version 3.10.1 (Ewels et al. 2020;

Patel et al. 2024, https://doi.org/10.5281/zenodo.7505987) was then used, as implemented with

"nextflow" version 22.10.6 (Di Tommaso et al. 2017). Fastq files were sub-sampled with fq version

2022-02-15 (https://github.com/stjude-rust-labs/fq) and strandedness was automatically inferred using

Salmon version 1.4.0 (Patro et al. 2017). Trimmed reads were then mapped using STAR (v2.7.9a, Dobin

et al. 2013) with default parameters, on the Zm-B73-REFERENCE-NAM-5.0 maize reference genome

(https://phytozome-next.jgi.doe.gov/info/Zmays_Zm_B73_REFERENCE_NAM_5_0_55) and on the

Phaseolus   vulgaris   v2.1   bean   reference   genome   (https://phytozome-

next.jgi.doe.gov/info/Pvulgaris_v2_1). Quantification was then made with Salmon version 1.4.0, and

the file "salmon.merged.gene_counts.tsv" was used for subsequent analysis.

2.2.4 Differential gene expression in sole cropping and intercropping

From the raw read counts, we verified reproducibility among replicates (blocks) by computing R^2. We

removed the library B1_M2B1_1, which was identified as R^2 outlier in the two comparisons involving

blocks 2 and 3 (Figure S2). We used the standard procedure proposed by EdgeR (Lun et al. 2016) for

filtering and normalization. In order to compute scaling factors for each library, we used the Trimmed

Mean of M-values (TMM) normalization method (Robinson and Oshlack 2010) implemented in EdgeR
(function calcNormFactors) with the following parameters: min.count=10, min.total.count=15,
large.n=3, min.prop=0.6. We subsequently performed MultiDimensional Scaling (MDS) on the resulting
normalized counts to compute pairwise distances between libraries.

In order to identify differentially expressed (DE) genes, we further used EdgeR to estimate the
trended dispersion parameters (function estimateDisp), fit a quasi-likelihood negative binomial
generalized log-linear model (glmQLFit), and employed a quasi-likelihood F-test as described in Lun et
al. (2016) (function glmQLFTest) to compute contrasts and associated P values for each transcript. The
P values were then adjusted to control the FDR at 5% (Benjamini and Hochberg 1995). First, using only
the intercropped beans, we tested the maize genotype effect by contrasting M1 vs M2. Second, we
decomposed the bean expression variance into a cropping system (sole vs intercropping) effect and a
block effect. As for maize, we decomposed the expression variance into a cropping system effect, a
genotype effect, a block effect and a (cropping system-by-genotype) interaction. Just like for beans, DE
genes and adjusted P values were identified from the sole vs intercropping contrast.

3. RESULTS

3.1 Soil properties and farming practices
We examined soil properties among farms. Overall, the soils of the 16 fields were all loamy but their
contents varied from 17.1% to 30.0% for clay, from 36.1% to 62.0% for silt and from 10.3% to 44.8%
for sand (Figure S3). We found significant differences among farms for 6 of the 13 soil variables (Dataset
S9) and variability between fields was mainly due to soil fertility variables such as total N content (N),
organic matter content (OM), C:N ratio (CN) and Cation Exchange Capacity (CEC), as exhibited on the
first principal component of the PCA (Dim1, Figure 2a).

Among the three variables related to farming practices (Nsim, Nin and irrigation), the simulated
soil nitrogen mineralization from organic matter (Nsim) and the N fertilization (Nin) differed among
farms (Dataset S9). Notably, the only organic farm in our panel (F4) stood out as an outlier along the
first principal component of the PCA, suggesting superior soil fertility as expected from the longer crop
rotation and high pre-sowing manure inputs in this farming system (Figure 2b). Conversely, farm F6
(which used the highest quantity of nitrogen fertilizers; Dataset S1) displayed low values for fertility
soil variables (Figure 2). Note that Nsim was positively correlated with N, OM and CEC (r > 0.63) while
it was negatively correlated with irrigation (r = -0.79, P = 0.0003, Dataset S10). Additionally, Nsim
exhibited a negative correlation with N fertilization (r = -0.27, P = 0.037), as expected, indicating that
soils with greater soil mineralization require less fertilization.

3.2 Soil bacterial communities associated with plants are shaped by agronomic practices and cropping system.

We described rhizospheric and non-rhizospheric bacterial communities. A total of 4,632,917 16S rRNA sequences were obtained. Their abundance varied from 80,394 to 160,393 sequences (Dataset S5). Sequence assignation resulted in 12,171 bacterial OTUs, classified at seven taxonomic levels (Dataset S6). When combining data from all farms, the most prevalent phyla were *Proteobacteria*, *Actinobacteria*, *Bacteroidetes*, *Chloroflexi* and *Firmicutes* (Figure S4).

Interestingly, we observed no significant difference in alpha diversity between maize and bean. However, intercropping conditions displayed a trend toward higher alpha diversity (Observed OTUs and Chao1 indexes) irrespective of the rhizosphere/non-rhizosphere status for both maize and bean soils (Figure S5). This increase in diversity under intercropping conditions was significant for soils associated with maize only (FDR Q = 0.014). The observed diversity of non-rhizosphere soil in intercropping conditions appeared to be intermediate between the non-rhizosphere soils of intercropped maize and intercropped bean.

We examined distances between samples considering all OTUs (Figure S6) and revealed a closer proximity among samples originating from the same farm (farm effect at P < 0.001 in PERMANOVA, Dataset S11), regardless of soil conditions (rhizosphere vs non-rhizosphere), plant species (maize vs bean), or cropping system (intercropping vs sole cropping). OTUs whose counts differed among samples, however, separated rhizospheric from non-rhizospheric samples with a clearer pattern for beans than for maize (Figure 3). The second axis of the BCA differentiated samples by cropping system for both crops (Figure 3).

Co-inertia analysis unveiled a significant association between OTU composition and the 13 soil variables, irrigation, N input, and Nsim (correlation coefficient RV = 0.58; P = 0.025, Figure S7). The first dimension of soil PCA (Figure 2) exhibited a negative contribution to the first dimension of co-inertia (-0.93) while the second dimension of soil PCA contributed positively to the second dimension of co-inertia (0.93). The first two axes of the co-inertia accounted for 61.2% of the total inertia (Figure S7). Notably, Irrigation and Nsim presented the highest values of canonical weights (respectively 0.45 and -0.43, Dataset S12) that defined the second axis of co-inertia, indicating that these two variables highly impacted OTU composition. F4 and F6 emerged as the most distinct farms on the co-inertia plot along the first axis (Figure S7). Interestingly, F4 was the only organic farm in our sample, applying the lowest level of mineral inputs (Dataset S1) while displaying superior soil fertility (Figure 2).

Finally, Wald multiple comparison tests carried out with individual OTUs indicated that *Vibrio*, *Acinetobacter*, *Pseudonocardia alaniniphila*, *Hydrogenophaga palleronii* and *Pseudoxanthomona koreensis* were more abundant in sole-cropped than intercropped bean rhizosphere, while *Janthinobacterium lividum* was more abundant in the intercropped bean rhizosphere (Figure S8). In the maize rhizosphere, *Actinospica*, *Sphingomonas mali, Ktedonobacter robiniae* (previously bacterium SOSP1-30) and *Alcaligenes* were more abundant in sole-cropped samples while *Thermoactinomyces*

and *Bdellovibrio* were more abundant in intercropped samples (Figure S9). These last results need however to be taken with caution as the data are scarce and significance is, in some instances, driven by a single data point.

3.3 Cropping system impacts maize agronomic traits

We examined the effect of the cropping system on maize agronomic traits using a multivariate analysis. We found that the first principal component explained 40.9% of the variation and was strongly associated with yield-related variables, including propagule dimensions and number (Dim 1, Figure 4a). This principal component correlated positively with nitrogen fertilization (Nin, r = 0.77, $P < 0.01$) and irrigation (r = 0.68, $P < 0.01$), and negatively with simulated nitrogen mineralization (Nsim, r = -0.78, $P < 0.01$). In addition, different fields of a given farm tended to cluster together along the first principal component, which suggests that farming practices and local soil properties primarily determined maize yield (Figure 4b). The ellipse plot further revealed a clear distinction between sole (M) and intercropped maize (IC) along this axis (Figure 4b).

The difference between cropping systems was stable across farms for three traits, intercropping being associated to higher values of two N-nutrition related traits (NP and *d15NV*) and lower values of kernel weight per ear (Sw), regardless of the farm (Figure 5, Dataset S13). The percentage of carbon in vegetative parts (CV), although showing significant farm-by-cropping system interaction, exhibited consistently smaller values in intercropping than in sole cropping (Figure S10). For eight other traits, the difference between cropping systems varied among farms (i.e., significant farm-by-cropping system interactions, Dataset S13).

3.4 Cropping system impacts bean agronomic traits

We examined the effect of the cropping system on bean agronomic traits. The first two principal components explained over 59% of the variation (Figure 4c). The first principal component was associated with yield and nutrition traits (Pw, NbS, NbP, PS, CP_p, CV_p, NP_p, Vw), while the second one was linked to propagule size (Pwi, Ple and Psu). Sole cropped (B) and intercropped (IC) beans were clearly separated along the first principal component (Figure 4d). Intercropping negatively affected yield components across all farms, although the intensity varied, as indicated by significant farm-by-cropping system interactions (Figure S11). This included vegetative weight (Vw), seed weight (Pw), the number of seeds per plant (NbP), and the number of pods per plant (NbS). A similar pattern was observed for traits related to nitrogen and carbon content per plant, such as CV_p, CP_p, and NP_p (Figure 5, Dataset S13). Interestingly, as compared to sole cropping, intercropped beans displayed higher values for seed size traits (Ple, Pwi, Psu) as well as for nutrition-related traits (NV, CP) across all farms (Figure 5, Dataset S13).

3.5 Competition prevails in maize-bean intercropping

Correlations between maize and bean traits were computed to explore the interplay between the two crops, positive correlations being expected under facilitation and negative correlations under competition. Of the 528 correlations (22 traits for maize and 24 traits for bean), 47 were significant among which 34 were negative and 13 were positive (Figure S12), pointing towards competition rather than beneficial interactions. Considering the subset of traits impacted either in the maize or the bean by intercropping (Figure 5), we found that maize traits related to nitrogen nutrition (*d15NV*; *d15NP*; CV_p; Table 1) and vegetative dry weight (Vw) were negatively correlated both with yield-related traits (NbS, Pw, Sw, Tpw, NbP) and two nutrition-related traits (CP_p, NP_p) measured in bean (Table 1). These results indicate inter-species competition for resources such as light, water and mineral N needed for nitrogen assimilation and CO_2 fixation through photosynthesis, resulting in reduced bean yield. Conversely, the nitrogen percentage in vegetative parts (NV) of bean was positively correlated with four maize traits (Pw, NbP, CP_p, *d13CP* ; Table 1). This most likely resulted from an indirect effect linked to the lower seed production—less nitrogen mobilized into the reproductive organs—in intercropped beans.

3.6 Controlled field assay shows that species competition affects bean transcriptome

We conducted a controlled field assay in INRAE plant breeding station of Saclay (northern France) to further assess the impact of intercropping on plant gene expression. Agronomic measures revealed that 6 of 13 yield and development traits exhibited lower values in intercropped bean as compared to sole-cropped bean, i.e. total seed mass per plant (reduced by 87%; Figure 6a), number of seeds per plant (reduced by 89%), number of pods per plant (reduced by 89%), width of the main leaflets (reduced by 19%), foliage density (reduced by 56%), and plant height (reduced by 59%; Figure 6b) (Figure S13). Although the two maize varieties strongly differed for all yield and development traits (except for the number of ears per plant and the number of lodged plants), they were both unaffected by the cropping condition (Figure 6c,d and Figure S14).

We investigated transcriptomic changes in sole cropping versus intercropping stands. First, we tested differences in maize gene expression between genotypes M1 and M2. We found clear distinctions between the two genotypes on the first axis of the MDS plot, that accounted for 66% of the total variation; Figure 7a). We further investigated differentially expressed (DE) genes, with the prediction that their number would be higher in beans than in maize, since bean phenotypes (but not maize phenotypes) were severely affected by the cropping condition. Following our prediction, we detected no DE genes between sole-cropped and intercropped maize plants. In bean, we identified only 11 DE genes between M1B and M2B and therefore concluded on the near absence of impact of maize variety on bean transcriptome. We thus considered M1B and M2B as biological replicates for the detection of DE genes. Out of 14,993 bean genes, nearly a third (5,070 genes) were DE between cropping conditions (first MDS axis, Figure 7b).

4. DISCUSSION

Intercropping systems are gaining in popularity, appreciated for their low-input requirements as well as their yield in low-input settings. Such positive effects result from beneficial plant-plant interactions (Fréville et al. 2022). In this study, we undertook a multidisciplinary approach to get insights into maize-bean intercropping, a practice that was recently revived in southwestern France for the cultivation of the Tarbais bean. The primary strength, yet limitation, of this study lies in the direct sampling of plants from on-farm fields. This approach introduces numerous confounding factors such as maize variety, inputs, abiotic and biotic environmental factors, that are challenging to control. Here, these factors were collectively captured as the "farm" effect.

Soil analyses revealed notable differences in fertility and pH across farms (Dataset S9), while soil texture remained relatively consistent (Figure S3). Globally, we observed a positive correlation between nitrogen mineralization by microorganisms and soil fertility, i.e., with total N content (r = 0.64; $P < 0.01$) and organic matter content (r = 0.64; $P < 0.01$), alongside a negative trend with nitrogen fertilizer application. These correlations driven by the organic farm suggest that farmers compensate for poor soil fertility with nitrogen inputs, and they also highlight the intricate connections between farming practices, soil fertility, and the capacity of microorganisms to release nutrients into the soil.

While it is well known that bulk bacterial communities are shaped by various soil variables— including texture (Girvan et al. 2003), nitrogen content (Zeng 2016), pH (Lauber et al. 2008), and water (Li 2021), the rhizosphere communities are also shape by crop genotypes (Bouffaud et al. 2016) and physiology, which collectively determine the compounds released in the immediate vicinity of roots as exudates (Ulbrich et al. 2022). Due to the complexity of interactions, studying the impact of intercropping on bacterial composition remains challenging (Duan et al. 2024). Here, we found a trend towards higher diversity in soil samples obtained from intercropping than from sole cropping (significant in maize; Figure S5), a pattern observed in sugarcane–soybean intercropping (Malviya et al. 2021). Interestingly, alpha diversity levels of maize and bean soils were not different, meaning that the greater diversity observed in intercropped maize more likely results from a complementarity effect rather than a selective effect (Loreau and Hector 2001). The combination of root exudates released by both crops may indeed enhance the diversity of the associated bacterial communities. In addition, we found in bean that two of the OTUs that discriminated the cropping systems include plant growth-promoting rhizobacteria (Figure S8). The *Acinetobacter* spp., which were predominant in sole-cropped bean rhizosphere, comprise strains that can produce indole-3-acetic acid (IAA), siderophores, gibberellin, antibiotics, biosurfactants/bioemulsifiers and solubilize phosphate, potassium, and zinc (Rokhbakhsh-Zamin et al. 2011). Also, *Janthinobacterium lividum* was more abundant in the bean rhizosphere when intercropped with maize. Interestingly, strains of this species can inhibit a number of soil-borne pathogens, including *Pythium ultimum*, *Rhizoctonia solani* and *Rhizoctonia oryzae*, and display biocontrol activity in the wheat rhizosphere (Yin et al. 2021).

In cereal-legume intercropping, cereal roots develop more quickly, utilizing nitrogen before
legumes thereby stimulating nodule formation and nitrogen fixation in beans (Hauggaard-Nielsen et al.
2008). Although we did not conduct a detailed examination of root nodules, initial observations revealed
no nodules on the bean in topsoil. This is likely caused by the sampling timing (close to harvest), which
is not ideal to make such observations, but nevertheless suggests limited symbiotic nitrogen fixation,
despite the nodulation capability of Tarbais beans (Laguerre et al. 2006). This could be due to two
factors: the non-restrictive nitrogen fertilization conditions, which can inhibit nodulation (Voisin et al.
2002), or the absence of the specific rhizobial strains needed for nodulation in the soil (Yan et al. 2014).
Because we evidenced presence of *Rhizobium leguminosarum* (Dataset S6), involved in bean
nodulation, our findings suggest that years of conventional practices of maize monoculture without a
Fabaceae cover crop may have diminished the potential benefits of plant-microbial interactions in these
fields.

In order to assess the potential beneficial interactions between maize and bean in intercropping,
we examined the effect of intercropping on plant nutrition and development. Pairwise correlations
between traits of the two crops revealed that most significant correlations were negative, indicating that
competition outweighed beneficial interactions. Notably, yield components in both crops were
negatively affected by intercropping, with beans experiencing greater reductions compared to sole
cropping situations (Figures S10, S11 and S12). Interestingly, however, bean seed size traits (length,
width and surface) were larger in intercropped than sole cropping, indicating that competition with maize
translates into beans producing fewer pods and seeds but larger seeds. This pattern illustrates the trade-
off in resource allocation between fecundity and seed size, the latter providing offspring with more
resources (Aarssenand and Keog 2002) and therefore a potential competitive advantage during early
development, as shown in other species (Houssard and Escarré 1991). In intercropping, maize also
displayed increased nitrogen acquisition in kernels and higher ^{15}N abundance in vegetative tissues
(Figure 5), suggesting an enhanced capacity of maize to mobilize soil nitrogen – the elevated ^{15}N levels
indicate that maize primarily relies on soil nitrogen. This contrasts with bean, which showed a higher
nitrogen percentage in vegetative tissues (Figure S12). The reduced capacity of certain beans to compete
with maize for soil nitrogen (Dawo et al. 2009) likely explains their lower yield. In this setting, nitrogen
fixation was limited, as evidenced by no significant increase in nitrogen fixation in intercropping.
Enhanced nitrogen in vegetative parts of beans may hence simply reflect a reduced demand for nitrogen
in seeds due to fewer seeds being produced, and a lower dilution of nitrogen assimilates. These findings
suggest that competition between species dominates the observed resource allocation patterns, with
maize benefiting more from shared soil nitrogen resources while beans face limitations.

Finally, our field assay in Saclay provided a first glimpse on the characterization of the
molecular signatures of maize-bean intercropping (Becker et al. 2023). Our results show that
competition in intercropping was much stronger than what we observed on-farm. It is interesting to note
that under such increased competition, the burden falls on the bean and resulted into a strong decline in

yield (~ 87%) and vegetative development as compared to sole cropping beans. Note that early growth
of bean remained unaffected. In contrast, maize showed no difference between cropping conditions
across all measured traits. Interestingly, gene expression analysis concurred with phenotypic data,
revealing no differentially expressed genes in maize but 5070 DE genes in beans, which indicated a
complete rewiring of bean gene expression when suffering drastic competition.
630
631 **5. CONCLUSION**
632
Despite the intrinsic limitations of field surveys in farmers' settings, including confounding effects and
scarce sampling, our results demonstrate that intercropping leads to plant competition that significantly
impacts the agronomic traits of maize and bean in the Tarbes region. However, this impact is highly
dependent on farming practices. Key factors such as crop variety, plant row arrangement and density,
soil properties and farm inputs that affect soil fertility and microbial composition play critical roles in
shaping these outcomes. Differences observed between on-farm surveys and the field experiment, for
instance, illustrate the potential importance of maize genotype and environmental conditions, such as
soil composition and drought, in driving plant competition dynamics. These findings underscore the
pivotal role of farming practices and soil fertility in mediating species interactions, extending even to
the diversity of bacterial soil communities. Interestingly, intercropping tends to foster more diverse
microbial communities compared to sole cropping, a potentially beneficial effect for soil health. Despite
this advantage, intercropping remains characterized by strong interspecific competition, particularly to
the detriment of bean. This competition negatively affects both gene expression and yield-related traits
in bean. However, a notable outcome is the production of larger seeds by intercropped beans compared
to those grown in sole cropping—an attribute that could be valued by farmers as larger seed size may
be a criterion for premium labels, guaranteeing higher market prices and stable demand.
This interdisciplinary study paves the way for recommendations on maize-bean intercropping
and outlines several research perspectives. First, the genetic basis of beans in the Tarbes region is
particularly narrow and the two bean varieties have been selected without explicit consideration of their
symbiotic N-fixation potential in soils of this area, where rhizobial inoculation is not customary. Second,
beans are the primary focus in this system, and selecting an appropriate maize variety is crucial. The
ideal maize partner should be robust enough to support the beans but not overly competitive. Hybrids
optimized for sole cropping and mineral nitrogen absorption are unsuitable in this context. Molecular
studies, such as those conducted here but expanded to other growth stages, plant organs, and varieties,
could help identify the metabolic pathways involved and guide partner selection using available genetic
resources. Third, the study encourages a reevaluation of the criteria for the success of this intercropping
system. Beyond productivity, traits such as the nutritional quality of seeds and/or their specific
characteristics could be shaped by consumer preferences, while acknowledging the positive impacts of
intercropping on the agroecosystem.

ACKNOWLEDGEMENTS

We are extremely grateful to Hélène Corti, a retired technician who helped us collecting the data. In addition, we would like to thank Marie-Hélène Jeuffroy and Etienne Delannoy for insightful discussion at the start of the project. This work would not have been possible without the contribution of farmers and the Coopérative du Haricot Tarbais. Three anonymous reviewers greatly contributed to improve the manuscript.

AUTHOR CONTRIBUTIONS

Conceptualization, M.C., T.F., J.E., Y.M-L., Do.M., M.I.T.; Methodology, N.V.-B., M.C., C.B., T.F., J.E., V.P., E.L.C., Da.M., Y.M.-L., Do.M., M.I.T.; Software, N.V-B, T.T., C.P.-L., T.F., Da.M., Y.M.-L., Do.M., M.I.T.; Validation, N.V.-B., L.M., T.T.; Formal analysis, N.V.-B., L.M., T.T., V.P.; Investigation, N.V.-B., B.L., C.P., M.C., M.L.G., J.C., A.M., Y.M.-L., Do.M., M.I.T.; Resources, C.P., M.C., C.B., J.C., C.P.-L., T.F., M.L.G, V.P., A.M., Y.M.-L.; Writing – Original Draft, N.V.-B., L.M., Do.M., M.I.T.; Writing –Review & Editing, N.V.-B., L.M., T.F., J.E., E.L.C., V.P., Da.M., Y.M.-L., Do.M. M.I.T.; Vizualization, N.V.-B., L.M., T.T.; Supervision, Y.M.-L., Do.M. M.I.T.; Project administration, Y.M.-L., M.I.T.; Funding Acquisition, N.V.-B., C.P.-L., T.F., Y.M.-L., Do.M., M.I.T.

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest to declare that are relevant to the content of this article.

DATA AVAILABILITY STATEMENT

Raw reads from metabarcoding data have been deposited in the DDBJ/EMBL/GeneBank under accession number PRJNA 1112577 and similarly for raw reads from QuantSeq 3'RNA-seq under accession number PRJNA 1115135. Raw phenotypic data as well as input files to run the code were deposited on recherche.data.gouv.fr with the DOI https://doi.org/10.57745/BZMPT9. Codes for RNAseq and phenotypic data analyses are available at recherche.data.gouv.fr with the DOI https://doi.org/10.57745/BZMPT9.

SUPPORTING INFORMATION

This paper contains a supporting information file with 14 supplementary figures 13 supplementary datasets.

REFERENCES

Aarssen, L. W., & Keog, T. (2002). Conundrums of competitive ability in plants: what to measure? *Oikos*, *96*(3), 531-542. https://doi.org/10.1034/j.1600-0706.2002.960314.x

Becker, C., Berthomé, R., Delavault, P., Flutre, T., Fréville, H., Gibot-Leclerc, S., Le Corre, V., Morel, J-B., Moutier, N., Munos, S., Richard-Molard., C., Westwood, J., Courty, P-E., de Saint Germain, A., Louarn, G. & Roux, F. (2023). The ecologically relevant genetics of plant-plant interactions. *Trends in Plant Science*, 28(1): 31-42. https://doi.org/10.1016/j.tplants.2022.08.014

Bedoussac, L., Journet, E.-P., Hauggaard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen, E. S., Prieur, L., & Justes, E. (2015). Ecological principles underlying the increase of productivity achieved by cereal-grain legume intercrops in organic farming. A review. *Agronomy for Sustainable Development*, *35*(3), 911-935. https://doi.org/10.1007/s13593-014-0277-7

Beillouin, D., Ben-Ari, T., Malézieux, E., Seufert, V., & Makowski, D. (2021). Positive but variable effects of crop diversification on biodiversity and ecosystem services. *Global Change Biology*, *27*(19), 4697-4710. https://doi.org/10.1111/gcb.15747

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate : A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289-300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bitocchi, E., Bellucci, E., Giardini, A., Rau, D., Rodriguez, M., Biagetti, E., Santilocchi, R., Zeuli, P. S., Gioia, T., Logozzo, G., Attene, G., Nanni, L., & Papa, R. (2013). Molecular analysis of the parallel domestication of the common bean (*Phaseolus vulgaris*) in Mesoamerica and the Andes. *New Phytologist*, *197*(1), 300-313. https://doi.org/10.1111/j.1469-8137.2012.04377.x

Bonnain-Dulon, R., & Brochot, A. (2004). De l'authenticité des produits alimentaires. *Ruralia, 14*, 969. http:// journals.openedition.org/ruralia/969

Bouffaud, M.-L., Renoud, S., Moënne-Loccoz, Y., & Muller, D. (2016). Is plant evolutionary history impacting recruitment of diazotrophs and *nifH* expression in the rhizosphere? *Scientific Reports, 6*(1), 21690. https://doi.org/10.1038/srep21690

Brooker, R. W., Bennett, A. E., Cong, W. F., Daniell, T. J., George, T. S., Hallett, P. D., Hawes, C., Iannetta, P. P. M., Jones, H. G., Karley, A. J., Li, L., McKenzie, B. M., Pakeman, R. J., Paterson, E., Schob, C., Shen, J. B., Squire, G., Watson, C. A., Zhang, C. C., … White, P. J. (2015). Improving intercropping : A synthesis of research in agronomy, plant physiology and ecology. *New Phytologist, 206*(1), 107-117. https://doi.org/10.1111/nph.13132

Clivot, H., Mouny, J.-C., Duparque, A., Dinh, J.-L., Denoroy, P., Houot, S., Vertès, F., Trochard, R., Bouthier, A., Sagot, S., & Mary, B. (2019). Modeling soil organic carbon evolution in long-term arable experiments with AMG model. *Environmental Modelling & Software, 118*, 99-113. https://doi.org/10.1016/j.envsoft.2019.04.004

Dawo, M. I., Wilkinson, J. M., & Pilbeam, D. J. (2009). Interactions between plants in intercropped
maize and common bean. *Journal of the Science of Food and Agriculture, 89*(1), 41-48.
https://doi.org/10.1002/jsfa.3408

Di Tommaso, P., Chatzou, M., Floden, E. W., Barja, P. P., Palumbo, E., & Notredame, C. (2017).
Nextflow enables reproducible computational workflows. *Nature Biotechnology, 35*(4),
316-319. https://doi.org/10.1038/nbt.3820

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., &
Gingeras, T. R. (2013). STAR : Ultrafast universal RNA-seq aligner. *Bioinformatics* (Oxford,
England), *29*(1), 15-21. https://doi.org/10.1093/bioinformatics/bts635

Duan, Y., Lei, X., Cao, Y., Liu, L., Zou, Z., Ma, Y., Zhu, X., & Fang W. (2024). Leguminous green
manure intercropping changes the soil microbial community and increases soil nutrients and
key quality components of tea leaves. *Horticultural research*, *11*(3): uhae0188.
https://doi.org/10.1093/hr/uhae018

Ewels, P. A., Pelzer, A., Fillinger, S., Patel, H., Alneberg, J., Wilm, A., Garcia, M. U., Di Tommaso, P.,
& Nahsen, S. (2020). The nf-core framework for community-curated bioinformatics pipelines.
Nature Biotechnology, 38(3), 271-271. https://doi.org/10.1038/s41587-020-0435-1

Farid, M., & Navabi, A. (2015). N$_2$ fixation ability of different dry bean genotypes. *Canadian Journal
of Plant Science, 95*(6), 1243-1257. https://doi.org/10.4141/cjps-2015-084

Fréville, H., Montazeaud, G., Forst, E., David, J., Papa, R., & Tenaillon, M. I. (2022). Shift in
beneficial interactions during crop evolution. *Evolutionary Applications, 15*(6), 905-918.
https://doi.org/10.1111/eva.13390

Gaba, S., Lescourret, F., Boudsocq, S., Enjalbert, J., Hinsinger, P., Journet, E. P., Navas, M. L., Wery,
J., Louarn, G., Malezieux, E., Pelzer, E., Prudent, M., & Ozier-Lafontaine, H. (2015). Multiple
cropping systems as drivers for providing multiple ecosystem services : From concepts to
design. *Agronomy for Sustainable Development, 35*(2), 607-623.
https://doi.org/10.1007/s13593-014-0272-z

Girvan, M. S., Bullimore, J., Pretty, J. N., Osborn, A. M., & Ball, A. S. (2003). Soil type is the primary
determinant of the composition of the total and active bacterial communities in arable soils.
Applied and Environmental Microbiology, 69(3), 1800-1809.
https://doi.org/10.1128/AEM.69.3.1800-1809.2003

Hauggaard-Nielsen, H., Jørnsgaard, B., Kinane, J., & Jensen, E. S. (2008). Grain legume–cereal
intercropping : The practical application of diversity, competition and facilitation in arable and
organic cropping systems. *Renewable Agriculture and Food Systems, 23*(1), 3-12.
https://doi.org/10.1017/S1742170507002025

Houssard C. & Escarré J. (1991). The effects of seed weight on growth and competitive ability of
Rumex acetosella from two successional old-fields. *Oecologia, 86*(2), 236-242.
https://doi.org/10.1007/BF00317536.

Hupe, A., Naether, F., Haase, T., Bruns, C., Heß, J., Dyckmans, J., Joergensen, R. G., & Wichern, F. (2021). Evidence of considerable C and N transfer from peas to cereals via direct root contact but not via mycorrhiza. *Scientific Reports, 11*(1), 11424. https://doi.org/10.1038/s41598-021-90436-8

Johansen, A., & Jensen, E. S. (1996). Transfer of N and P from intact or decomposing roots of pea to barley interconnected by an arbuscular mycorrhizal fungus. *Soil Biology & Biochemistry, 28*(1), 73-81. https://doi.org/10.1016/0038-0717(95)00117-4

Laguerre, G., Courde, L., Nouaım, R., Lamy, I., Revellin, C., Breuil, M. C., & Chaussod, R. (2006). Response of rhizobial populations to moderate copper stress applied to an agricultural soil. *Microbial Ecology, 52*, 426-435. https://doi.org/10.1007/s00248-006-9081-5

Lauber, C. L., Strickland, M. S., Bradford, M. A., & Fierer, N. (2008). The influence of soil properties on the structure of bacterial and fungal communities across land-use types. *Soil Biology and Biochemistry, 40*(9), 2407-2415. https://doi.org/10.1016/j.soilbio.2008.05.021

Li, B., Li, Y.-Y., Wu, H.-M., Zhang, F.-F., Li, C.-J., Li, X.-X., Lambers, H., & Li, L. (2016a). Root exudates drive interspecific facilitation by enhancing nodulation and N2 fixation. *Proceedings of the National Academy of Sciences of the United States of America, 113*(23), 6496-6501. https://doi.org/10.1073/pnas.1523580113

Li, C., Dong, Y., Li, H., Shen, J., & Zhang, F. (2016b). Shift from complementarity to facilitation on P uptake by intercropped wheat neighboring with faba bean when available soil P is depleted. *Scientific Reports, 66*(1), 18663. https://doi.org/10.1038/srep18663

Li, C., Stomph, T.-J., Makowski, D., Li, H., Zhang, C., & Zhang, F. (2023). The productive performance of intercropping. *Proceedings of the National Academy of Sciences of the United States of America, 120*(2), e2201886120. https://doi.org/10.1073/pnas.2201886120

Li, H. (2021). Irrigation has a higher impact on soil bacterial abundance, diversity and composition than nitrogen fertilization. *Scientific Reports, 11*(1), 16901. https://doi.org/10.1038/s41598-021-96234-6

Li, H., Shen, J., Zhang, F., Clairotte, M., Drevon, J.-J., Le Cadre, E., & Hinsinger, P. (2008). Dynamics of phosphorus fractions in the rhizosphere of common bean (*Phaseolus vulgaris* L.) and durum wheat (*Triticum turgidum durum* L.) grown in monocropping and intercropping systems. *Plant and Soil, 312*(1), 139-150. https://doi.org/10.1007/s11104-007-9512-1

Li, L., Li, S. M., Sun, J. H., Zhou, L. L., Bao, X. G., Zhang, H. G., & Zhang, F. S. (2007). Diversity enhances agricultural productivity via rhizosphere phosphorus facilitation on phosphorus-deficient soils. *Proceedings of the National Academy of Sciences of the United States of America, 104*(27), 11192-11196. https://doi.org/10.1073/pnas.0704591104

Li, S. M., Li, L., Zhang, F. S., & Tang, C. (2004). Acid phosphatase role in chickpea/maize intercropping. *Annals of Botany, 94*(2), 297-303. https://doi.org/10.1093/aob/mch140

Loreau, M. & Hector A. (2001). Partitioning selection and complementarity in biodiversity experiments. *Nature, 412*, 72-76. https://doi.org /10.1038/35083573

Lun, A. T. L., Chen, Y., & Smyth, G. K. (2016). It's DE-licious : a recipe for differential expression analyses of RNA-seq experiments using quasi-likelihood methods in edgeR. In E. Mathé & S. Davis (Eds.), *Statistical Genomics* (Vol. 1418, p. 391-416). Springer New York. https://doi.org/10.1007/978-1-4939-3578-9_19

Ma, H. (2023). Maize/alfalfa intercropping enhances yield and phosphorus acquisition. *Field Crops Research, 303*, 109136. https://doi.org/10.1016/j.fcr.2023.109136

Maitra, S., Hossain, A., Brestic, M., Skalicky, M., Ondrisik, P., Gitari, H., Brahmachari, K., Shankar, T., Bhadra, P., Palai, J. B., Jena, J., Bhattacharya, U., Duvvada, S. K., Lalichetti, S., & Sairam, M. (2021). Intercropping—A low input agricultural strategy for food and environmental security. *Agronomy, 11*(2), 343. https://doi.org/ 10.3390/agronomy11020343

Malviya, M. K., Solanki, M. K., Li, C.-N., Wang, Z., Zeng, Y., Verma, K. K., Singh, R. K., Singh, P., Huang, H.-R., Yang, L.-T., Song, X.-P., & Li, Y.-R. (2021). Sugarcane-legume intercropping can enrich the soil microbiome and plant growth. *Frontiers in Sustainable Food Systems, 5*, 606595. https://doi.org/10.3389/fsufs.2021.606595

Marmagne, A., Masclaux-Daubresse, C., & Chardon, F. (2022). Modulation of plant nitrogen remobilization and postflowering nitrogen uptake under environmental stresses. *Journal of Plant Physiology, 277*, 153781. https://doi.org/10.1016/j.jplph.2022.153781

Matsuoka, Y., Mitchell, S. E., Kresovich, S., Goodman, M., & Doebley, J. (2002). Microsatellites in *Zea* – variability, patterns of mutations, and use for evolutionary studies. *Theoretical and Applied Genetics, 104*(2), 436-450. https://doi.org/10.1007/s001220100694

Mazzola, M. (2004). Assessment and management of soil microbial community structure for disease suppression. *Annual Review of Phytopathology, 42*, 35-59. https://doi.org/10.1146/annurev.phyto.42.040803.140408

McMurdie, P. J., & Holmes, S. (2013). phyloseq : an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One, 8*(4), e61217. https://doi.org/10.1371/journal.pone.0061217

Mead, R., & Willey, R. W. (1980). The concept of a 'land equivalent ratio' and advantages in yields from intercropping. *Experimental Agriculture, 16*(3), 217-228. https://doi.org/10.1017/S0014479700010978

Metson, A. J. (1957). Methods of chemical analysis for soil survey samples. *Soil Science, 83*(3), 245. https://doi.org/10.7931.dl1-sbb-012

Moyer-Henry, K. A., Burton, J. W., Israel, D. W., & Rufty, T. W. (2006). Nitrogen transfer between plants : A ^{15}N natural abundance study with crop and weed species. *Plant and Soil, 282*(1), 7-20. https://doi.org/10.1007/s11104-005-3081-y

Nassary, E. K., Baijukya, F., & Ndakidemi, P. A. (2020). Productivity of intercropping with maize and
common bean over five cropping seasons on smallholder farms of Tanzania. *European*
Journal of Agronomy, 113, 125964. https://doi.org/10.1016/j.eja.2019.125964

Parnaudeau, V. V., Reau, R. R., & Dubrulle, P. P. (2012). Un outil d'évaluation des fuites d'azote vers
l'environnement à l'échelle du système de culture : Le logiciel Syst'N. *Innovations*
Agronomiques, 21, 59-70. http://www7.inra.fr/ciag/revue/volume_21_septembre_2012

Patel, Y., Zhu, C., Yamaguchi, T. N., Bugh, Y. Z., Tian, M., Holmes, A., Fitz-Gibbon, S. T., & Boutros,
P. C. (2024). NFTest : Automated testing of Nextflow pipelines. *Bioinformatics, 40*(2),
btae081. https://doi.org/10.1093/bioinformatics/btae081

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and
bias-aware quantification of transcript expression. *Nature Methods, 14*(4), 417-419.
https://doi.org/10.1038/nmeth.4197

Peña-Cabriales, J. J., & Castellanos, J. Z. (1993). Effects of water stress on N2 fixation and grain yield
of *Phaseolus vulgaris* L. *Plant and Soil, 152*(1), 151-155. https://doi.org/10.1007/BF00016345

Peoples, M. B., Boddey, R. M., & Herridge, D. F. (2002). Quantification of nitrogen fixation (Chapter
13). In G. J. Leigh (Ed.), Nitrogen Fixation at the Millennium (p. 357-389). Elsevier Science.
https://doi.org/10.1016/B978-044450965-9/50013-6

Poggio, S. L. (2005). Structure of weed communities occurring in monoculture and intercropping of
field pea and barley. *Agriculture, Ecosystems & Environment, 109*(1), 48-58.
https://doi.org/10.1016/j.agee.2005.02.019

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). Limma
powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic*
Acids Research, 43(7), e47. https://doi.org/10.1093/nar/gkv007

Robinson, M. D., & Oshlack, A. (2010). A scaling normalization method for differential expression
analysis of RNA-seq data. *Genome Biology, 11*(3), R25. https://doi.org/10.1186/gb-2010-11-3-
r25

Rodriguez, C., Carlsson, G., Englund, J.-E., Flöhr, A., Pelzer, E., Jeuffroy, M.-H., Makowski, D., &
Jensen, E. S. (2020). Grain legume-cereal intercropping enhances the use of soil-derived and
biologically fixed nitrogen in temperate agroecosystems. A meta-analysis. *European Journal*
of Agronomy, 118, 126077. https://doi.org/10.1016/j.eja.2020.126077

Rokhbakhsh-Zamin, F., Sachdev, D., Kazemi-Pour, N., Engineer, A., Pardesi, K. R., Zinjarde, S. S.,
Dhakephalkar, P. K., & Chopade, B. A. (2011). Characterization of plant-growth-promoting
traits of *Acinetobacter* species isolated from rhizosphere of *Pennisetum glaucum. Journal of*
Microbiology and Biotechnology, 21(6), 556-566. https://doi.org/10.4014/jmb.1012.12006

Schlatter, D. C., Bakker, M. G., Bradeen, J. M., & Kinkel, L. L. (2015). Plant community richness and
microbial interactions structure bacterial communities in soil. *Ecology, 96*(1), 134-142.
https://doi.org/10.1890/13-1648.1

Shearer, G., & Kohl, D. H. (1988). Natural ^{15}N abundance as a method of estimating the contribution
of biologically fixed nitrogen to N$_2$-fixing systems : Potential for non-legumes. *Plant and*
Soil, 110(2), 317-327. https://doi.org/10.1007/BF02226812

Shearer, G., Kohl, D. H., & Harper, J. E. (1980). Distribution of ^{15}N among plant parts of nodulating
and nonnodulating isolines of soybeans. *Plant Physiology, 66*(1), 57-60.
https://doi.org/10.1104/pp.66.1.57

Smith, T., Heger, A., & Sudbery, I. (2017). UMI-tools : Modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. *Genome Research, 27*(3), 491-499.
https://doi.org/10.1101/gr.209601.116

Ulbrich, T. C., Rivas-Ubach, A., Tiemann, L. K., Friesen, M. L., & Evans, S. E. (2022). Plant root
exudates and rhizosphere bacterial communities shift with neighbor context. *Soil Biology &*
Biochemistry, 172, 108753. https://doi.org/10.1016/j.soilbio.2022.108753

Vazeux-Blumental, N., Manicacci, D., & Tenaillon, M. I. (2024). The milpa, from Mesoamerica to
present days, a traditional agricultural system serving agroecology. *Comptes rendus de*
l'Académie des Sciences Biologies, 6(347), 159-173. https://doi.org/10.5802/crbiol.164

Voisin, A.-S., Salon, C., Munier-Jolain, N. G., & Ney, B. (2002). Effect of mineral nitrogen on
nitrogen nutrition and biomass partitioning between the shoot and roots of pea (*Pisum sativum*
L.). *Plant and Soil, 242*(2), 251-262. https://doi.org/10.1023/A:1016214223900

Xing, Y., Yu, R.-P., An, R., Yang, N., Wu, J.-P., Ma, H.-Y., Zhang, J.-D., Bao, X.-G., Lambers, H., &
Li, L. (2023). Two pathways drive enhanced nitrogen acquisition via a complementarity effect
in long-term intercropping. *Field Crops Research, 293*, 108854.
https://doi.org/10.1016/j.fcr.2023.108854

Yan, J., Han, X. Z., Ji, Z. J., Li, Y., Wang, E. T., Xie, Z. H., & Chen, W. F. (2014). Abundance and
diversity of soybean-nodulating rhizobia in black soil are impacted by land use and crop
management. *Applied and Environmental Microbiology, 80*(17), 5394-5402.
https://doi.org/10.1128/AEM.01135-14

Yin, C., Casa Vargas, J. M., Schlatter, D. C., Hagerty, C. H., Hulbert, S. H., & Paulitz, T. C. (2021).
Rhizosphere community selection reveals bacteria associated with reduced root disease.
Microbiome, 9(1), 86. https://doi.org/10.1186/s40168-020-00997-5

Zeng, J., Liu, X., Song, L., Lin, X., Zhang, H., Shen, C., & Chu, H. (2016). Nitrogen fertilization
directly affects soil bacterial diversity and indirectly affects bacterial community composition.
Soil Biology & Biochemistry, 92, 41-49. https://doi.org/10.1016/j.soilbio.2015.09.018

Zhang, C., Postma, J. A., York, L. M., & Lynch, J. P. (2014). Root foraging elicits niche
complementarity-dependent yield advantage in the ancient 'three sisters' (maize/bean/squash)
polyculture. *Annals of Botany, 114*(8), 1719-1733. https://doi.org/10.1093/aob/mcu191

Zhang, D., Zhang, C., Tang, X., Li, H., Zhang, F., Rengel, Z., Whalley, W. R., Davies, W. J., & Shen,
J. (2016). Increased soil phosphorus availability induced by faba bean root exudation
stimulates root growth and phosphorus uptake in neighbouring maize. *New Phytologist,*
209(2), 823-831. https://doi.org/10.1111/nph.13613
Zhang, S., Lehmann, A., Zheng, W., You, Z., & Rillig, M. C. (2019). Arbuscular mycorrhizal fungi
increase grain yields : A meta-analysis. *New Phytologist, 222*(1), 543-555.
https://doi.org/10.1111/nph.15570
Zizumbo-Villarreal, D., & Colunga-GarcíaMarín, P. (2010). Origin of agriculture and plant
domestication in West Mesoamerica. *Genetic Resources and Crop Evolution, 57*(6), 813-825.
https://doi.org/10.1007/s10722-009-9521-4

 FIGURE LEGENDS

FIGURE 1. The local Tarbais bean intercropped with maize (left) or as a sole crop (right).

FIGURE 2. Principal Component Analysis (PCA) computed on 13 soil variables per field. The two first

axes are shown. (a) Correlation circle of the 13 soil variables with three supplementary variables added

to the PCA (in blue), i.e. Nsim (which estimates soil nitrogen mineralization), Irrigation, and Nin

(nitrogen units supplied during the growing season). (b) Projection of fields with colors indicating farms

and symbols the cropping system, i.e. maize sole cropping (M), bean sole cropping (B) or maize-bean

intercropping (IC). Variable names are indicated in Dataset S2.

FIGURE 3. Between Class Analysis (BCA) computed on 11,586 OTUs. Samples are named according

to rhizosphere (R) or non-rhizosphere (NR) soil, followed by maize (M) or bean (B), and then sole

cropping (SC) or intercropping (IC). Projection on the first two dimensions are presented, with the

corresponding percentages of total inertia explained by these axes.

FIGURE 4. Principal Component Analysis (PCA) on maize and bean phenotypic and nutrition

variables. The first two PCs computed from 22 maize variables (Table 1) are shown with corresponding

correlation circle (a) and data projection (b). Similarly, for beans, the first two PCs computed on 24

variables (Table 1) are shown with corresponding correlation circle (c) and data projection (d). The three

variables Irrigation, Nin (N input) and Nsim (simulated soil mineralization) shown in blue (a and c)

were added to the PCA. In b and d, colours indicate farms, symbols the cropping system with

corresponding ellipses, i.e. sole-cropped maize in b (M, dashed lines), sole-cropped bean in d (B, dashed

line), intercropped maize in b and intercropped bean in d (IC, full lines).

FIGURE 5. Effect of the cropping system on maize and bean phenotypic and nutrition variables. A two-

way ANOVA model with interaction was used to test the effects of the Farm and the Cropping system

(sole cropping versus intercropping) on phenotypic and nutrition variables (Table 1). Significant P

values for the cropping system effect are indicated by *, ** and *** for $P < 0.05$, $P < 0.01$ and $P <$

0.001, respectively. Pink asterisks indicate significance of tests performed on maize traits and green

asterisks those for bean traits. Asterisks are underlined when the value is higher in intercropping than in

sole cropping system. Variables (see Table 1 for full details) impacted by intercropping are *d15NV* (delta

15N among total N of the Vegetative parts), NV (Percent Nitrogen in Vegetative parts), CV (Percent

Carbon in Vegetative parts), CV_p (Per plant Carbon content in Vegetative parts), Vw (Vegetative parts

dry weight per plant), Pw (Propagule dry weight per plant), TPw (Thousand Propagules dry weight),

NbP (Number of Propagules produced per plant), NbS (Number of Sets of propagules per plant), Sw

(Average propagule weight per Set), Ple (Average Propagule length), Pwi (Average Propagule width),

Psu (Average Propagule surface), NP (Percent Nitrogen in Propagules), CP (Percent Carbon in
Propagules), CP_p (Per plant Carbon content in propagules) and, NP_p (Per plant Nitrogen content in
propagules).

FIGURE 6. Development and yield of maize and bean plants under sole cropping and intercropping in
the field assay at Saclay. Stripcharts display the total seed weight per bean plant (a) and the height of
bean plants (b), the total kernel weight per maize plant (c) and the height of maize plants (d). Blue dots
represent the average value in each of the three blocks and horizontal lines indicate the means. Data are
shown in a and b for sole-cropped maize genotype 1 (M1), sole-cropped maize genotype 2 (M2), maize-
bean intercropping (M1B and M2B), and in c and d for sole-cropped bean (B), and maize-bean
intercropping (M1B and M2B). Pictures illustrate sole-cropped beans (e) and intercropped beans and
maize (f).

FIGURE 7. MDS plots of distances between gene expression profiles in sole-cropped and intercropped
maize and bean. The distances observed between samples in the two MDS plots approximate the typical
$\log_2$ fold changes between the samples. MDS plots were computed from 16,021 genes for maize samples
(a) and 14,993 genes for bean samples (b). Colors and labels indicate the cropping system and the
sample, for sole-cropped maize of the two different genotypes (M1 and M2), sole-cropped bean (B), and
maize-bean intercropping (M1B and M2B). The last digit in the labels indicate the corresponding block
(1, 2, or 3). The percentage of variation explained by each dimension is shown.

TABLE 1. Description of phenotypes and nutrient variables measured on plants. [a]: only measured for the bean.

Variable names	Variable description
Plants	Number of Plants per sample
Vw	Vegetative parts dry weight per plant (g)
Pw	Propagule dry weight per plant (g)
NbP	Number of Propagules produced per plant
Ple	Average Propagule length – measured on 50 propagules (mm)
Pwi	Average Propagule width – measured on 50 propagules (mm)
Psu	Average Propagule surface – measured on 50 propagules (mm^2)
d15NP	Delta ^{15}N among total N of the Propagules
d13CP	Delta ^{13}C among total C of the Propagules
d15NV	Delta ^{15}N among total N of the Vegetative parts
d13CV	Delta ^{13}C among total C of the Vegetative parts
NP[a]	Percent Nitrogen (measured in dry matter) in Propagules
CP[a]	Percent Carbon (measured in dry matter) in Propagules
NV[a]	Percent Nitrogen (measured in dry matter) in Vegetative parts
CV[a]	Percent Carbon (measured in dry matter) in Vegetative parts
NP_p	Per plant Nitrogen content (g) in propagules
CP_p	Per plant Carbon content (g) in propagules
NV_p	Per plant Nitrogen content (g) in Vegetative parts
CV_p	Per plant Carbon content (g) in Vegetative parts
NbS	Number of Sets of propagules (ears or pods) per plant
PS (=NbP/NbS)	Average number of Propagules per Set – ears or pods
Sw (=Pw/NbS)	Average propagule weight per Set – ears or pods (g)
TPw (=1000xPw/NbP)	Thousand Propagules dry weight (g)
d15NVdfa[a]	N in the vegetative parts derived from symbiotic fixation (for bean only)
d15NPdfA[a]	N in the propagules derived from symbiotic fixation (for bean only)

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Maize

Bean

Figure 5.

Figure 6.

Figure 7.