1   **Genetic architecture underlying response to the fungal pathogen *Dothistroma septosporum***

2   **in lodgepole pine, jack pine, and their hybrids**

3
4   **Abstract**
5

In recent decades, *Dothistroma* needle blight (DNB), a pine tree disease caused by the fungal
pathogen *Dothistroma septosporum,* has severely damaged lodgepole pine (*Pinus contorta*
Dougl. ex. Loud.) in British Columbia, Canada, and raised health concerns for jack pine (*Pinus*
banksiana Lamb.). The pathogen has already shown signs of host shift eastward to the hybrid
populations between lodgepole pine and jack pine (*Pinus contorta* × *P. banksiana*), and possibly
into pure jack pine. However, we have little knowledge about mechanisms of resistance to *D.*
septosporum, especially the underlying genetic basis of variation in pines. In this study, we
conducted controlled inoculations to induce infection by *D. septosporum* and performed a
genome-wide case-control association study with pooled sequencing (pool-seq) data to dissect
the genetic architecture underlying response in lodgepole pine, jack pine, and their hybrids. We
identified candidate genes associated with *D. septosporum* response in lodgepole pine and in
hybrid samples. We also assessed genetic structure in hybrid populations and inferred how
introgression may affect the distribution of genetic variation involved in *D. septosporum*
response in the studied samples. These results can be used to develop genomic tools to evaluate
DNB risk, guide forest management strategies, and potentially select for resistant genotypes.

21
22
23   **KEY WORDS**
24   Conifer, fungal pathogen resistance, F_{ST}, hybrid, introgression, pool-GWAS
25
26
27   **1 | INTRODUCTION**
28
Pests and pathogens constantly threaten forest trees, and their impacts are changing as climates
are altered (Seidl et al., 2017; Simler-Williamson et al., 2019). Knowledge about the genetic
architecture of pathogen resistance and the identified candidate resistance (R) genes provides the
basis for genomic breeding to improve tree resilience to forest pathogens. Genes and alleles that
confer disease resistance or tolerance can eventually be used to develop genomic tools for
rapidly selecting resistant genotypes (Isabel et al., 2019). Current breeding practices for forest
trees involve recurrent selection, which includes repeated cycles of selection, breeding, and
testing, to search for genotypes with better disease resistance, and then deployment of these
selected tree genotypes (White, 2004). This takes years of effort to design and implement.
Increasingly, genome editing is used to develop crops resistant to disease (van Esse et al., 2020).
For fruit trees, CRISPR/Cas9 system has been applied to enhance the disease resistance in citrus
(Jia et al., 2017; Wang et al., 2019), apple (Zhou et al., 2020), and others (Min et al., 2022).
Given the long generation time in conifers and the frequent threats from diseases and insects,
identifying R genes that can be targeted for selection or editing is a useful step in developing
resistant genotypes for reforestation.
44

Dothistroma needle blight (DNB), which is caused by the fungal pathogen *Dothistroma septosporum,* is a foliar disease of a wide range of pine trees (Gibson, 1972), with infected trees showing reddish-brown bands in needles, defoliation, and growth reduction. DNB has caused increased damage to lodgepole pine (*Pinus contorta* Dougl. ex. Loud., LP) in recent decades in Western Canada (Dale et al., 2011; Woods, 2003), and is now expanding its range eastwards, threatening jack pine (*Pinus banksiana* Lamb., JP). In severe cases, DNB causes extensive mortality, even amongst mature trees in LP plantations (Woods, 2003). Historically, DNB had only minor impacts on native forest trees in North America, but emerged as a severe forest disease during 1950s to 1960s in *Pinus radiata* plantations in Africa, New Zealand, and South America (Gibson, 1972). Recently, DNB has caused increased outbreak incidence and host range expansion in North America and Europe, which is likely correlated with climate change (Boroń et al., 2021; Welsh et al., 2009; Woods et al., 2005). In North America, while *D. septosporum* tends to be more common in moist western forests closer to the Pacific Ocean, *Dothistroma*-like symptoms were observed in drier LP and LP × JP in northern Alberta in 2012 and 2013 with further tests confirming *D. septosporum* was the cause of the disease observed (Ramsfield et al., 2021). Feau et al. (2021) performed a controlled inoculation experiment and demonstrated that LP, JP, and LP × JP are susceptible to *D. septosporum*, with JP showing a higher disease severity than LP or LP × JP, though there is no report showing DNB infection in natural JP forests. Considering *D. septosporum*'s eastward shift reaching towards the natural range of JP and JP's high susceptibility to *D. septosporum*, it is important to understand how these species respond to DNB infection.

The evolutionary history and biogeography of JP and LP have likely played an important role in shaping the genetic basis of resistance to DNB. JP is widely distributed across boreal forests in eastern North America, extending in Canada from Northwest Territories to Nova Scotia (Rudolph & Laidly, 1990), areas largely outside of the historical range of *Dothistroma*. LP and JP evolved in allopatry, with an estimated divergence time of ~8 million years before present (Hao et al., 2015), but hybridized in areas of contact, primarily in central and northwestern Alberta (Burns et al., 2019; Rudolph & Yeatman, 1982). Following secondary contact of these two species, additional patchy regions of introgression have been developed in central Alberta north through to the Northwest Territories and northeastern British Columbia to the Alberta-Saskatchewan border (Burns et al., 2019). Natural hybrids between LP and JP (*Pinus contorta × P. banksiana,* LP × JP) display a wide range of phenotypes that are intermediate or more typical of one parent, depending on the level of introgression (Wood et al., 2009; Yeatman & Teich, 1969).

In light of the geographic distribution of LP and JP, the longer history of host-pathogen coevolution in LP might explain its stronger resistance to DNB infection, compared to JP. A former study has evidenced coevolution between *Cronartium harknessii* lineages with LP and JP, with genetic basis underlying both pathogen virulence and host resistance (McAllister et al., 2022). For DNB infection caused by *D. septosporum*, though we lack direct evidence to prove a coevolutionary history between *D. septosporum* and different pine hosts, we have already identified candidate resistance genes to *D. septosporum* by RNA-seq analysis on experimentally infected LP seedlings (Lu et al., 2021). These candidate genes include 43 genes that are present in the plant-pathogen interaction pathway, as well as nine R genes that contained sites under positive selection. It seems that both constitutive (baseline) and induced (defenses activated after

being attacked) defenses have been developed in LP. However, little is known about genetic
basis of resistance to *D. septosporum* in JP. When subjected to other biotic agents, LP and JP
showed unique monoterpene profiles in response to mountain pine beetle (Hall et al., 2013) and
significant differences in chitinase gene expression in response to the fungal pathogens
Grosmannia clavigera and *Cronartium harknessii* (Peery et al., 2021). These previous studies
revealed the species-specific pathogen responses in a focused set of gene families. Nevertheless,
since many complex traits are determined by many genes of small effect, to identify the causal
genes underlying disease response traits, a genome-wide gene survey might give us a high-
resolution perspective (Alonso-Blanco & Méndez-Vigo, 2014; Tam et al., 2019).
100
To dissect the genetic architecture of pathogen resistance and to identify candidate genes,
researchers have employed genetic markers, mainly SNPs, and methods like quantitative trait
locus (QTL) mapping and genome-wide association analysis (GWAS). Polygenic disease
responses, comprising numerous loci of small effect, appear to be the rule (Stocks et al., 2019).
Large numbers of SNPs were associated with pitch canker resistance in *Pinus taeda* L. (De La
Torre et al., 2019; Lu et al., 2017; Quesada et al., 2010), with white pine blister rust resistance in
Pinus lambertiana Dougl. (Weiss et al., 2020), and with *Heterobasidion* root rot resistance in
Picea abies (L.) Karst. (Capador-Barreto et al., 2021). To identify the candidate R genes, most
GWAS studies on forest trees employ individual-based methods, with both phenotypic and
genotypic data collected on each sampled individual. Case-control methods have been developed
to study disease where individuals are pooled into healthy vs. infected groups, but such
approaches typically still use individual-based sequencing to call genotypes before pooling (Wu
et al., 2010). When multiple replicates are available, it is possible to combine pool-seq and the
case-control design: Endler et al. (2016) analyzed the genetic differences underlying abdominal
pigmentation variation among *Drosophila* populations, while Stocks et al. (2019) identified
SNPs associated with low versus high ash dieback damage in *Fraxinus excelsior* L. using this
approach. Recently, Singh et al. (2024) used a case-control pool-seq approach similar to that
deployed here to search for signatures of resistance to Swiss needle cast in *Pseudotsuga*
menziesii (Mirb.) Franco. Pool-seq is a cost-effective alternative approach to sequencing of
individuals, but the weakness and limitations are also evident, such as unequal representation of
individuals and suboptimal allele frequency estimate when the pool size is small, as well as
misaligned mapping and sequencing errors (Schlötterer et al., 2014). Nonetheless, in a study on
conifer genotyping, Lind et al. (2022) found that allele frequencies estimated from pooled DNA
sequencing samples were highly correlated with frequencies estimated from individual
sequencing.
126
Another factor in understanding the genetics of resistance is whether there is interplay between
any alleles conferring resistance and introgression between the species. Bechsgaard et al. (2017)
showed that plant R genes can adaptively introgress between closely related species. The LP × JP
hybrid zone has been very well studied by Cullingham et al. (2012), who found that this hybrid
zone presents a mosaic zone with variable introgression and patchy distributions of hybrids
(Burns et al., 2019). While it is clear that these species share a broad hybrid zone, it is unclear
whether any alleles for disease resistance will be more or less introgressed than the average
region in the genome.
135

In this study, our aim is to identify the genetic basis of DNB resistance in LP, JP, and LP × JP
samples and to explore spatial patterning in any identified loci. The specific objectives of our
study were to: 1) develop a GWAS case-control approach using pool-seq samples to identify loci
with consistent associations with *D. septosporum* response across replicates; and 2) compare the
genetic architecture underlying DNB resistance within the studied pine trees and infer how
introgression may affect the genetic basis of *D. septosporum* responses. This study provides an
important step towards identifying candidate genes to develop genomic tools for screening trees
resistant to DNB infection.
144
145 **2 | METHODS**
146
2.1 | Plant materials
Seeds were obtained from 40 natural seedlots across Western Canada (Figure 1, seedlot numbers
and locations can be found in Table S1, seed contributors http://adaptree.forestry.ubc.ca/seed-
contributors/), including 25 LP seedlots from British Columbia (BC_LP) and three from Alberta
(AB_LP), seven LP × JP seedlots from Alberta (AB_LPxJP), and five JP seedlots from Alberta
(AB_JP). The range maps were downloaded from
https://sites.ualberta.ca/~ahamann/data/rangemaps.html (Hamann et al., 2005). When the seeds
were collected in the wild, they were assigned to pure LP, pure JP, or LP × JP, based on their
location and morphological traits such as cone and branch characteristics, microfibril angle, and
cell area (Wheeler & Guries, 1987; Wood et al., 2009; Yeatman & Teich, 1969). The proportion
of LP and JP ancestry of the collected seeds were genotyped using 11 microsatellite loci by
Cullingham et al. (2012). Briefly, seeds were germinated to obtain seedlings, then DNA was
isolated from the seedlings. DNA was used to amplify 11 microsatellite loci and allele sizes were
determined for genotyping as described by Cullingham et al. (2011).
161
For the present study, seeds from the 40 seedlots were used to grow seedlings in a greenhouse at
University of British Columbia, Vancouver, BC, for their first year (a flow chart of experimental
procedure is shown in Figure S1). Resistance was phenotyped by inoculating 100 individuals per
seedlot with one of two *D. septosporum* isolates (D1 & D2, 50 individuals of each isolate). D1
was isolated from needles of an infected LP × JP seedlot in Alberta, Canada, while D2 was
isolated from an infected LP seedlot in the Kispiox Valley region close to Smithers in
Northwestern British Columbia, Canada (Feau et al., 2021; Ramsfield et al., 2021). The seedlots
were identified as LP or LP × JP as aforementioned (Cullingham et al., 2012). The controlled
inoculation experiment was performed as described in Kabir et al. (2013) and Feau et al. (2021).
Briefly, the one-year-old seedlings were placed in a completely randomized experimental design
in a growth chamber with the condition of 16 hours daylight at 20 ℃, 8 hours a night at 12 ℃,
and minimum relative humidity of 80 %. *D. septosporum* conidia were harvested from colonies
grown for 10~16 days on *Dothistroma* sporulation medium plates and then suspended in sterile
distilled water. Each seedling was sprayed twice at 10-day intervals with a standard inoculum of
approximately 3 mL of 1.6×10^6 conidia mL^{-1}, using a household trigger action atomizer. Once
the needles were dry (~45 minutes), the seedlings were wrapped in transparent plastic bags and
kept in the growth chamber. After 48h the plastic bags were removed and the seedlings were
kept in the growth chamber for 15 weeks till phenotyping. A mister spraying tap water was
activated hourly for 3 minutes during the trial to maintain needle wetness. Control seedlings, at
least one for each seedlot, were inoculated with sterile water.

Fifteen weeks after inoculation, seedlings were rated for proportion of necrotic needles with red
bands and/or fruiting bodies according to a disease severity scale of 1 to 5 (1 = least, 5 = most
disease severity). Following disease rating, the ten most- and ten least-infected individuals per
seedlot (five individuals for each *D. septosporum* isolate) were identified and retained for DNA
extraction. Genomic DNA of each of these individuals was extracted as described in Lind et al.
(2022), using the Nucleospin 96 Plant II Core kit (Macherey–Nagel GmbH & Co. KG, Germany)
on an Eppendorf epMotion 5075 liquid-handling platform. The DNA of five most-infected and
five least-infected individuals by each of the two *D. septosporum* isolates was combined in
equimolar amounts to compose four pooled libraries per population, two susceptible (D1-S, D2-S)
and two resistant libraries (D1-R, D2-R). Hence,160 pooled DNA libraries were generated.
192

193
194
FIGURE 1 Sampling locations (Table S1) of the 40 pine seedlots analyzed in this study, plotted
within the natural range of *Pinus contorta* (lodgepole pine) and *Pinus banksiana* (jack pine) in
Canada and USA. The topright map shows the context of the sampling locations. The range
maps were downloaded from https://sites.ualberta.ca/~ahamann/data/rangemaps.html. BC_LP
(green diamonds) — *Pinus contorta* seedlots from British Columbia (N=25); AB_LP (orange
diamonds) — *Pinus contorta* seedlots from Alberta (N=3); AB_ LPxJP (purple diamonds) —
Pinus contorta × P. banksiana seedlots from Alberta (N=7); AB_JP (brown diamonds) — *Pinus*
banksiana seedlots from Alberta (N=5).
203

2.2 | Sequence capture and pool-seq genotyping

The capture probes were comprised of two sets of probes. The first set of probes was designed based on an existing LP sequence capture array (Suren et al., 2016) by removing the probes that did not yield successful genotyping in Yeaman et al. (2016). Probe sequences of the existing capture array were aligned to the reference genome using GMAP v2019-03-15 (Wu & Watanabe, 2005). Probes that covered the genomic regions with called SNPs in the dataset from Yeaman et al. (2016) were retained, otherwise the probes were discarded. Since there is no available LP reference genome, a masked *Pinus taeda* reference genome Pita.2_01.masked3k2.fa (https://treegenesdb.org/FTP/Genomes/Pita/v2.01/genome/) (Neale et al., 2014) was used instead. *Pinus taeda* is a closely related species to LP (Jin et al., 2021). The second set of probes was newly designed probes derived from the *D. septosporum*-induced genes, which were based on a LP reference transcriptome assembled using the RNA-seq data of *D. septosporum* infected LP samples (Lu et al., 2021). To avoid duplicates, only those *D. septosporum*-induced genes with low homology to the retained probe sequences were used to design the new probes. To do so, the retained probe sequences were aligned to this transcriptome using blastn v2.9.0 with an E-value of 1e-10. A total of 8,778 *D. septosporum*-induced genes did not have any aligned probe sequences. These non-duplicate *D. septosporum*-induced genes were subsequently aligned to the reference genome to predict the exon-intron boundaries using GMAP v2019-03-15. Exon sequences from these induced genes with a length of at least 100 bp were combined with the previously designed working probe sequences, and this combined set of sequences was submitted to Roche NimbleGen (Roche Sequencing Solutions, Inc., CA USA) for Custom SeqCap EZ probe design (design name: 180321_lodgepole_v2_EZ). Combining the two sets of probes, this updated LP sequence capture array has a capture space of 44 Mbp, containing roughly 35,467 assembled genes. Most LP genes responsive to environment stress and fungal pathogen attack were included in the current capture probe design. Though genes expressed in different development periods might be missed, genes that have evidence of substantial expression have been covered in this capture probe design.

The capture libraries for each of the 160 pools were constructed following NimbleGen SeqCap EZ Library SR User's Guide and as described in Lind et al. (2022). Then the R (resistant) and S (susceptible) libraries (two libraries per capture, R1+S1 or R2+S2, indexed with different barcodes) per population and per isolate were combined for sequence capture and enrichment. Sequencing was performed using the Illumina NovaSeq 6000 S4 PE 150 platform in Centre d'expertise et de services Génome Québec. Our in-house pool-seq pipeline (Lind, 2021) was employed to align the reads to the reference genome and call SNPs. For raw SNPs, only bi-allelic loci in regions without annotated repetitive elements or potentially paralogous genes were retained. The annotated repetitive elements were acquired from the LP genome annotation (Wegrzyn et al., 2014). The potentially paralogous genes were identified as described in Lind et al. (2022) using haploid megagametophyte sequences, for the heterozygous SNP calls for haploid sequences are likely to represent misalignments of paralogs. Afterwards, the SNP loci with depth (DP) < 10, DP > 400, global minor allele frequency < 0.05, or > 25% missing data were also removed.

2.3 | Genetic structure analyses

The genetic structure among the 160 pooled samples, which represented four pooled libraries
(D1-R, D1-S, D2-R, D2-S) for each of the 40 seedlots, was detected by using principal
component analysis (PCA), reconstructing a phylogenetic tree, and by assessing correlation of
allele frequencies and population differentiation (F_{ST}) between samples using unlinked SNPs. To
reduce the impact of linkage disequilibrium on estimation, the SNP set was thinned using
vcftools (Danecek et al., 2011), so that no two sites were within 100,000 bp. PCA and genetic
distance were calculated using the major allele depth and the R package "adegenet" (Jombart,
2008). An unrooted phylogenetic tree was constructed using the distance output and the
Neighbour-Joining algorithm, which was implemented by the R package "ape" (Paradis &
Schliep, 2019). A bootstrap analysis was performed using 5000 bootstrap replicates. Correlation
coefficients of major allele frequencies between all pairs within the 160 samples were calculated
and plotted using the R package "corrplot" (Wei & Simko, 2017). F_{ST} was estimated between all
pairs within the 160 samples using the R package "poolfstat" (Hivert et al., 2018). The clustering
patterns output from these genetic structure analyses can be used to judge the subtle population
structure leading to false positive problems in GWAS.
264
265 **2.4 | Dissection of genetic architecture underlying *D. septosporum* response**
266
The alleles and candidate genes associated with *D. septosporum* response were detected using
the pool-GWAS method. A GWAS case-control approach was developed using pool-seq samples
to identify allele frequency differences between susceptible and resistant pines inoculated with *D.*
septosporum. This method was based on Cochran-Mantel-Haenszel (CMH) method and
implemented in our in-house pipeline (Lind, 2021). While sequencing read depth was commonly
used as an estimation of allele count in such applications of test (Futschik & Schlötterer, 2010;
Schlötterer et al., 2014), reliable results would be only achieved when the haploid size of the
sample is much greater than the depth of coverage (so that most reads arise from uniquely
sampled haplotypes). As the pool size used in the present study (diploid size of 2N=10) is often
smaller than the sequencing depth of coverage, read counts were converted into allele
frequencies and then multiplied by the ploidy to yield an approximate count for each pool that
represents the real replication level. A simulation was conducted to compare the false positive
rate from the CMH test using uncorrected or corrected allele counts across a range of number of
demes, individuals, and depths. Results showed that the false positive rate of the CMH method is
high, especially when the sequencing depth is much higher than the pool size (Figure S2a).
Using corrected allele counts, the allele frequencies multiplied by the ploidy (N=10) tends to
lower the false positive rate (Figure S2b). Thus, the corrected allele counts were used for CMH
test.
285
As the genetic structure analysis showed that LP × JP and JP samples tend to cluster, JP samples
and LP × JP samples were combined as JP + LP × JP samples for downstream analyses. Pool-
GWAS was performed separately within LP and JP + LP × JP samples. Analysis of R and S
samples was conducted by combining individuals inoculated with D1 or D2 isolates to maximize
statistical power. To identify candidate genes, SNPs in the top 1% of -$\log_{10}(p)$ values from a
pool-GWAS test were classified as outliers. As linkage disequilibrium can amplify signatures of
selection, candidate resistance genes were identified as those with a large number of outlier
SNPs relative to the average genome-wide expectation, as represented by an index based on the
binomial distribution as per Yeaman et al. (2016). The identified loci on the enriched genes were

realigned to the 12 LP linkage groups (MacLachlan et al., 2021) and Manhattan plots were
drawn. The LP transcriptome (Lu et al., 2021) was used to annotate the reference genome. The
aligned transcripts to the reference genome were identified as those with a minimum alignment
identity of 90 % and a minimum alignment coverage of 85 % using GMAP v2019-03-15.
299
300 **2.5 | Linkage disequilibrium (LD) between loci**
301
A non-linear model (Hill & Weir, 1988) was used to estimate the decay of LD with physical
distance. Pairwise correlation (r^2) of allele frequencies between loci on the same scaffold were
calculated. The r^2 values and the physical distances between loci on all scaffolds (genome-wide)
or on scaffolds containing the identified top candidate genes (significant) were used to fit the
non-linear model as described by Marroni et al. (2011). The LD decay graph was plotted using R
(R Core Team, 2018).
308
309 **2.6 | Introgression**
310
To study patterns of introgression affecting different regions of the genome and among different
samples, the mean pairwise F_{ST} was calculated for each locus using the R package "poolfstat".
Loci with high F_{ST} values have high genetic differentiation, which may represent species barriers
or genomic regions under divergent selection, while loci with low F_{ST} values imply introgression.
To estimate genetic similarity of the studied JP and LP seedlots, the five JP samples (p36 to p40)
and six pure LP samples (the six westmost LP samples, p15, p16, p18, p20, p23, p25) were used
to calculate F_{ST} values, which were then averaged. To evaluate whether regions associated with
D. septosporum response have atypical patterns of introgression, the F_{ST} values of the top 1 % *D.*
septosporum response outliers identified from LP (N = 3,358) and those from JP + LP × JP (N =
3,041) were compared, with F_{ST} values of randomly drawn SNPs (N = 3,400).
321
Similarly, F_{ST} between the JP + LP × JP samples (p29 to p40) and six pure LP samples (p15, p16,
p18, p20, p23, p25) were averaged to estimate genetic similarity of the studied JP + LP × JP and
LP. Different genomic regions were represented by different SNP sets, including *D. septosporum*
response outliers identified from LP (N = 3,358) and outliers identified from JP + LP × JP (N =
3,041), as well as the unlinked SNP set (N = 31,716), which was used in 2.3 for detecting genetic
structure and representing random genomic region. The genetic similarity (F_{ST}) of JP + LP × JP
and LP was regressed to the longitude of these JP + LP × JP samples. The regression slopes for
the three SNPs sets were compared using analysis of covariance method implemented by the R
package "lsmeans" (Lenth, 2016; R Core Team, 2018).
331
332 **3 | RESULTS**
333
3.1 | Patterns of genetic structure
After filtering, a total of 364,691 SNP loci were retained for downstream analyses, and most loci
had a minor allele frequency between 0.05 and 0.1 (Figure S3). A thinned set of 31,716 unlinked
SNPs was used for studying genetic structure among the 160 pooled samples (Figure 2). The
most prominent patterns of structure were associated with species (Figure 2a & b, Figure S4),
with LP samples readily distinguishable from JP and LP × JP samples. LP × JP samples clustered
more closely with JP than LP. Compared to those sampled from the eastern hybrid zone, western

LP × JP tend to be more similar to LP (samples were arranged according to longitude from left to
right in Figure 2 c & d). Given this clear pattern of genetic structure, we conducted GWAS in LP
and JP + LP × JP pine samples separately.
344
345

346
347
FIGURE 2 Genetic structure among the 160 pooled samples. a) Principal component analysis. b)
Unrooted phylogenetic tree. The bootstrap values for each branch were shown in Figure S4. c)
Correlation plot of the major allele frequencies between samples. d) Pairwise F_{ST} values between
pools. BC_LP — *Pinus contorta* (lodgepole pine) samples from British Columbia; AB_LP —
Pinus contorta (lodgepole pine) samples from Alberta; AB_LPxJP — *Pinus contorta × P.*
banksiana samples from Alberta; AB_JP — *Pinus banksiana* (jack pine) samples from Alberta.
D1 — *D. septosporum* isolate 1; D2 — *D. septosporum* isolate 2. R — resistant tree; S —
susceptible tree. In c) and d), each column represents a pooled sample. The samples were
arranged from left to right according to the longitude of sampling location, and four pooled
libraries per seedlot (D1-R, D1-S, D2-R, D2-S) were arranged together.
358
359
360
361
362 **3.2 | Signatures of association to infection response**

363
Out of the 100 top-ranked candidate genes in each test (Table S2), three were identified within
both the LP and JP + LP × JP samples (Figure 3). This is significantly more than the
hypergeometric expectation, based on the probability of overlap for two draws of 100 genes from
20,026 genes. A previous study anchored 10,093 scaffolds on the LP linkage map (MacLachlan
et al., 2021), so we plotted SNPs and labeled the enriched plant-pathogen interaction candidate
genes on the 12 linkage groups (Figure 3). The three genes identified in both species encode
ABC transporter, F-box Kelch-repeat protein, and serine/threonine protein kinase. Genes
encoding F-box protein and serine protein kinase were highlighted in Figure 3. Genes encoding
ABC transporter were not included in the linkage groups, so they were not included in Figure 3.
We also checked the enrichment of differentially expressed genes (DEGs) in the top 100 ranked
genes. Out of 631 DEGs identified in the previous study (Lu et al., 2021), four were also
candidate genes in the present study. Fisher's Exact test showed there is no significant
enrichment of DEGs in the candidate genes.
377

378
379
FIGURE 3 Manhattan plot showing SNPs associated with *D. septosporum* response within *Pinus*
contorta (lodgepole pine, LP) and combined *Pinus banksiana* and *Pinus contorta* × *P. banksiana*
(JP + LPxJP) samples. The x axis represents 12 linkage groups. The y axis represents
significance of associations identified from pool-GWAS. Candidate genes were identified using

a binomial test for enrichment of outlier SNPs per gene. Candidate genes and their surrounding
50 kbp regions are highlighted in red. Of the three top-ranked candidate genes identified within
both the LP and JP + LPxJP samples, one was not present in the linkage map, while the other
two are highlighted in blue ("Serine" on linkage group 4 and "F-box" on linkage group 10).
Labels in the graphs represent candidate genes encoding plant-pathogen interaction proteins:
bHLH — bHLH transcription factor; Serine — serine/threonine-protein kinase; Auxin — auxin
response factor; LRR — proteins with leucine-rich repeat; F-box — F-box Kelch-repeat protein;
MADS — MADS transcription factor; MYB — MYB transcription factor; WRKY — WRKY
transcription factor; TIF — translation initiation factor. The candidate gene IDs (in Table S2) are:
In LP, bHLH — Scaffold_2015-465162-569336; Serine_1 — Scaffold_3728-494595-736720;
Serine_2 — Scaffold_3141-1-105566; Serine_3 — Scaffold_1072-1299306-1374553; Auxin —
Scaffold_3823-1571412-1689102; LRR — Scaffold_1361-487806-570406; F-box —
Scaffold_3698-733755-839782; In JP + LPxJP, Serine — Scaffold_3728-494595-736720;
MADS — Scaffold_427-2324701-2717580; F-box_1 — Scaffold_3698-733755-839782; MYB
— Scaffold_4478-441537-627389; F-box_2 — Scaffold_342-1965787-2253515; WRKY —
Scaffold_830-1136302-1239031; TIF — Scaffold_207-7583953-7695067.
400
Loci on scaffolds with outliers (the SNPs with top 1 % of $-\log_{10}(p)$ values from pool-GWAS test)
tend to exhibit a more gradual decay in LD compared with the genome-wide loci on all scaffolds
(Figure 4), with JP + LP × JP pine outliers having a much slower decay rate than LP. In LP, the
distance at which LD decays to half of its maximum value is 96 bp for genome-wide loci, and
174 bp for outlier loci. In JP + LP × JP, the half LD decay distance is 1,357 bp for genome-wide
loci, and 68,215 bp for outlier loci. The slower LD decay pattern for the outliers may indicate
selection on these identified loci.
408

409
410 FIGURE 4 Linkage disequilibrium (LD) decay for genome-wide loci and outlier loci identified
411 from *Pinus contorta* (lodgepole pine, LP) and combined *Pinus banksiana* and *Pinus contorta* × *P.*
412 *banksiana* (JP + LPxJP) samples. The r^2 values (pairwise correlation of allele frequencies
413 between loci) and the physical distances between loci were used to fit the non-linear model as
414 described by Marroni et al. (2011).

3.3 | Introgression

By calculating average F_{ST} between pairs of LP and JP samples, we found that the *D. septosporum* response outliers, which were identified separately in LP or JP + LP × JP samples, have higher F_{ST} values than randomly drawn SNPs (Wilcoxon test *p*-value < 0.01, Figure 5a & 5b), with outliers identified in JP + LP × JP samples showing higher F_{ST} values than those identified in LP samples (Wilcoxon test *p*-value < 0.01). These results suggest that the outliers associated with disease response constitute or are linked to genomic regions for divergent selection or species barriers, rather than facilitating introgression. F_{ST} estimates between JP + LP × JP samples and a set of pure LP samples showed a pattern of increased differentiation with distance eastwards of the JP + LP × JP samples (Figure 5c), which were higher for the outliers than randomly chosen SNPs. Given the small number of samples, significance of linear model fitting is borderline, with *p*-value = 0.06 for outliers identified in LP, *p*-value = 0.05 for outliers identified in JP + LP × JP, and *p*-value = 0.06 for randomly chosen SNPs (df = 10). The differences between regression slopes on different gene sets (LP outliers, JP + LP × JP outliers, and unlinked SNPs) were not significant (*p*-values of pairwise comparisons > 0.3). Though it is not possible to test whether a linear vs. discontinuous model would fit better to these data, the results suggest that isolation-by-distance is occurring here.

FIGURE 5 F_{ST} values of *D. septosporum* response outliers (a & b) and regression patterns of F_{ST} values with longitude. For a & b, the F_{ST} values were calculated between the *Pinus banksiana* (jack pine, JP) samples (p36 to p40) and the six pure *Pinus contorta* (lodgepole pine, LP) samples (p15, p16, p18, p20, p23, p25).JP + LPxJP represents the combined JP and *Pinus contorta* × *P. banksiana* samples. F_{ST} values of *D. septosporum* response outliers identified from LP (N=3,358) and JP + LPxJP (N=3,041) samples were compared, with F_{ST} values of randomly drawn SNPs (N=3,400). Outliers identified from both species have higher F_{ST} values than randomly drawn SNPs (*p*-value < 0.01). For c, the F_{ST} values were calculated between the JP + LPxJP samples (p29 to p40) and the six pure LP samples (p15, p16, p18, p20, p23, p25) and then

446 averaged for each JP + LPxJP sample. Different genomic regions were represented by different
447 SNP sets, including LP outliers (N=3,358), JP + LPxJP outliers (N=3,041), and unlinked SNPs
448 (N=31,716). The F_{ST} values represented by different SNP sets were regressed to the longitude of
449 these JP + LPxJP samples. Differences between slopes are not significant (ANCOVA, *p*-value >
450 0.3).
451
452 **4 | DISCUSSION**
453
In recent decades, the fungal pathogen *D. septosporum* has become an increasing threat to LP in
British Columbia (Feau et al., 2021; Welsh et al., 2009). Although *Dothistroma* needle blight
(DNB) has not been observed in natural stands of JP in Alberta, Ramsfield et al. (2021)
confirmed JP's susceptibility to infection by *D. septosporum* using a trial under natural
conditions. The potential of climate change to allow this pathogen to spread across the wide
natural distribution of JP raises serious concerns. The hybrid zone of contact between the sister
species LP and JP may act as a bridge, allowing this fungal pathogen host shift (Taylor et al.,
2015), as evidenced by the intermediate phenotype of gall development in LP × JP seedlings
compared to that in LP and JP (McAllister et al., 2022). The genetic basis of pathogen resistance
in each species is therefore an important factor affecting the potential for host shift. To study the
genetics of resistance to DNB, we analyzed the genetic architecture underlying *D. septosporum*
response using the study samples of LP, JP, and LP × JP. We identified candidate genes
associated with *D. septosporum* response and inferred the impact of introgression on the genetic
structure underlying *D. septosporum* response in the studied samples. We detected largely non-
overlapping sets of candidate genes in LP and JP + LP × JP in response to *D. septosporum*,
which might be shaped by different evolutionary forces of selection and introgression.
Alternatively, the non-overlapping sets of candidate genes might reflect that many non-causal
genes exist in the top 1 % GWAS outliers of each species, given that we did not test for
significance at the gene-level. In either case, it seems clear that resistance to DNB is a highly
polygenic trait with alleles of small effect.
474
This study is the first report of genome-wide scanning of *D. septosporum* response genes in LP,
JP, and LP × JP. We identified a few top candidate genes encoding leucine-rich repeats (LRR)-
containing protein domain and serine threonine kinases in LP samples (Figure 3, Table S2). Plant
breeders have long used nucleotide-binding LRR (NB-LRR) receptor R genes for preventing or
reducing diseases in crops. When attacked by infectious pathogens, plant NB-LRR receptors can
recognize pathogen effectors by either direct or indirect mechanisms (Dodds & Rathjen, 2010).
Such strategy is called effector-triggered immunity (ETI), which occurs after a basal resistance,
pathogen-associated-molecular-patterns-triggered immunity (PTI). In a previous study using
RNA-seq data, Lu et al. (2021) identified genes involved in PTI and ETI, as well as R genes with
positive selection signals in LP. These results indicate that the long coevolutionary history
between LP and *D. septosporum* (Capron et al., 2021; Welsh et al., 2009) has given rise to some
tolerance or resistance mechanism against DNB. Since most of the identified associated markers
are likely not causal but rather are in LD with the causal loci, the observed fast LD decay rate in
lodgepole pine implies promising fine mapping resolutions as the identified markers are likely to
be proximal to the causative genes.
490

On the contrary, JP, as a potential new host, may not have coevolved such a specific defense, and the long LD blocks surrounding candidate genes identified from JP + LP × JP samples, which are typical of hybrid zones, limit the resolution of association mapping, so it may be difficult to locate the causative genes for follow-up studies (Goulet et al., 2017; Wilson & Goldstein, 2000). Additionally, we caution that the capture probes were designed using LP not JP genes, ascertainment bias might give rise to the underrepresentation of JP alleles (Lachance & Tishkoff, 2013). However, we can use these JP + LP × JP candidate genes to infer how introgression affects the frequencies of disease-resistant alleles. Since we analyzed pool-seq data and lacked the pure JP samples out of the hybrid zone, it is not possible to analyze parental origin information, thus we inferred the ancestry using clustering. The clustering patterns in Figure 2 show that most hybrid pine trees in our study had greater JP than LP ancestry, likely in part due to our limited sampling of JP from further Eastwards. Western LP × JP tend to be more genetically similar to LP, while eastern LP × JP tend to be more genetically dissimilar to LP. This is a typical characteristic for ongoing introgression and hybridization in the contact zone with a possible trend of LP expanding its range eastward (Harrison & Larson, 2014; Moran et al., 2021).

In hybrid zones, alleles in some genomic regions are able to introgress across species boundaries, whereas alleles that constitute species barriers or are under divergent selection will remain differentiated (Harrison & Larson, 2014). We found significantly high F_{ST} values between JP and LP samples for the candidate genes associated with *D. septosporum* infection response. This can occur if these candidate genes tend to reside in genomic regions with restricted gene flow, driven by linkage with genes for divergent selection or species barriers. McAllister et al. (2022) observed a gradient of resistance to *C. harknessii* in LP, which could be associated with introgression of resistance genes from JP into LP. Since LP might have a long coevolutionary history with *D. septosporum* (Capron et al., 2021; Welsh et al., 2009), the introgression of LP genes into hybrids may increase resistance to *D. septosporum*. However, variation in F_{ST} along longitude (Figure 5c) did not differ significantly between background and candidate loci, so we cannot draw strong conclusions about adaptive introgression here.

5 | CONCLUSION

The recent outbreaks of DNB in LP and its ability to infect JP pose a pressing need to understanding the genetic architecture of resistance to *D. septosporum*. The present study provides an array of candidate genes associated with *D. septosporum* response using pool-GWAS. The identified candidate genes within LP and JP + LP × JP samples are largely non-overlapping, and the adaptive introgression is yet to be found. This knowledge can be used to develop genomic tools to minimize DNB risk and guide forest management strategies. Further testing and validation of these associations using a SNP array is currently underway.

DATA AVAILABILITY STATEMENT

The raw pool-seq data were deposited in NCBI SRA (accession number: PRJNA602898; http://www.ncbi.nlm.nih.gov/sra). The assembled transcriptomes and their annotation were deposited in Dryad https://doi.org/10.5061/dryad.prr4xgxtk .The scripts used for analyses

537   were deposited in https://github.com/Mengmeng-Lu/Genetic-architecture-underlying-response-
538   to-the-fungal-pathogen-Dothistroma-septosporum-in-Pinus-con

539   REFERENCES

Alonso-Blanco, C. & Méndez-Vigo, B. (2014). Genetic architecture of naturally occurring
quantitative traits in plants: an updated synthesis. *Current opinion in plant biology, 18*,
37-43. https://doi.org/10.1016/j.pbi.2014.01.002

Bechsgaard, J., Jorgensen, T. H., & Schierup, M. H. (2017). Evidence for adaptive introgression of
disease resistance genes among closely related *Arabidopsis* species. *G3 (Bethesda), 7*(8),
2677-2683. https://doi.org/10.1534/g3.117.043984

Boroń, P., Lenart-Boroń, A., Mullett, M., Grad, B., & Nawrot-Chorabik, K. (2021). Population
structure of Dothistroma septosporum in Poland: revealing the genetic signature of a
recently established pathogen. *Plant Pathology, 70*(6), 1310-1325.
https://doi.org/10.1111/ppa.13383

Burns, I., James, P. M. A., Coltman, D. W., & Cullingham, C. I. (2019). Spatial and genetic
structure of the lodgepole × jack pine hybrid zone. *Canadian Journal of Forest Research,*
49(7), 844-853. https://doi.org/10.1139/cjfr-2018-0428

Capador-Barreto, H. D., Bernhardsson, C., Milesi, P., Vos, I., Lundén, K., Wu, H. X., . . . Elfstrand,
M. (2021). Killing two enemies with one stone? Genomics of resistance to two sympatric
pathogens in Norway spruce. *Molecular Ecology, 30*(18), 4433-4447.
https://doi.org/10.1111/mec.16058

Capron, A., Feau, N., Heinzelmann, R., Barnes, I., Benowicz, A., Bradshaw, R. E., . . . Hamelin, R.
C. (2021). Signatures of post-glacial genetic isolation and human-driven migration in the
Dothistroma needle blight pathogen in Western Canada. *Phytopathology®, 111*(1), 116-
127. https://doi.org/10.1094/phyto-08-20-0350-fi

Cullingham, C. I., Cooke, J. E., Dang, S., Davis, C. S., Cooke, B. J., & Coltman, D. W. (2011).
Mountain pine beetle host-range expansion threatens the boreal forest. *Molecular*
Ecology, 20(10), 2157-2171. https://doi.org/10.1111/j.1365-294X.2011.05086.x

Cullingham, C. I., James, P. M., Cooke, J. E., & Coltman, D. W. (2012). Characterizing the physical
and genetic structure of the lodgepole pine× jack pine hybrid zone: mosaic structure and
differential introgression. *Evolutionary Applications, 5*(8), 879-891.
https://doi.org/10.1111/j.1752-4571.2012.00266.x

Dale, A. L., Lewis, K. J., & Murray, B. W. (2011). Sexual reproduction and gene flow in the pine
pathogen *Dothistroma septosporum* in British Columbia. *Phytopathology, 101*(1), 68-76.
https://doi.org/10.1094/phyto-04-10-0121

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., . . . 1000 Genomes
Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics,*
27(15), 2156-2158. https://doi.org/10.1093/bioinformatics/btr330

De La Torre, A. R., Puiu, D., Crepeau, M. W., Stevens, K., Salzberg, S. L., Langley, C. H., & Neale,
D. B. (2019). Genomic architecture of complex traits in loblolly pine. *New Phytologist,*
221(4), 1789-1801. https://doi.org/110.1111/nph.15535

Dodds, P. N. & Rathjen, J. P. (2010). Plant immunity: towards an integrated view of plant–
pathogen interactions. *Nature review genetics, 11*(8), 539-548.
https://doi.org/10.1038/nrg2812

Drenkhan, R., Tomešová-Haataja, V., Fraser, S., Bradshaw, R. E., Vahalík, P., Mullett, M. S., . . .
Barnes, I. (2016). Global geographic distribution and host range of Dothistroma species:
a comprehensive review. *Forest Pathology, 46*(5), 408-442.
https://doi.org/10.1111/efp.12290
Endler, L., Betancourt, A. J., Nolte, V., & Schlötterer, C. (2016). Reconciling differences in pool-
GWAS between populations: A case study of female abdominal pigmentation in
Drosophila melanogaster. Genetics, 202(2), 843-855.
https://doi.org/10.1534/genetics.115.183376
Feau, N., Ramsfield, T. D., Myrholm, C. L., Tomm, B., Cerezke, H. F., Benowicz, A., . . . Hamelin, R.
C. (2021). DNA-barcoding identification of *Dothistroma septosporum* on *Pinus contorta*
var. *latifolia*, *P. banksiana* and their hybrid in northern Alberta, Canada. *Canadian*
Journal of Plant Pathology, 43(3), 472-479.
https://doi.org/10.1080/07060661.2020.1829065
Futschik, A. & Schlötterer, C. (2010). The next generation of molecular markers from massively
parallel sequencing of pooled DNA Samples. *Genetics, 186*(1), 207-218.
https://doi.org/10.1534/genetics.110.114397
Gibson, I. A. S. (1972). Dothistroma blight of *Pinus Radiata*. *Annual Review of Phytopathology,*
10(1), 51-72. https://doi.org/10.1146/annurev.py.10.090172.000411
Goulet, B. E., Roda, F., & Hopkins, R. (2017). Hybridization in plants: Old ideas, new techniques.
Plant physiology, 173(1), 65-78. https://doi.org/10.1104/pp.16.01340
Hall, D. E., Yuen, M. M. S., Jancsik, S., Quesada, A. L., Dullat, H. K., Li, M., . . . Bohlmann, J. (2013).
Transcriptome resources and functional characterization of monoterpene synthases for
two host species of the mountain pine beetle, lodgepole pine (*Pinus contorta*) and jack
pine (*Pinus banksiana*). *BMC plant biology, 13*(1), 80. https://doi.org/10.1186/1471-
2229-13-80
Hamann, A., Smets, P., Yanchuk, A. D., & Aitken, S. N. (2005). An ecogeographic framework for
in situ conservation of forest trees in British Columbia. *Canadian Journal of Forest*
Research, 35(11), 2553-2561. https://doi.org/10.1139/x05-181
Hao, Z.-Z., Liu, Y.-Y., Nazaire, M., Wei, X.-X., & Wang, X.-Q. (2015). Molecular phylogenetics and
evolutionary history of sect. *Quinquefoliae* (*Pinus*): Implications for Northern
Hemisphere biogeography. *Molecular Phylogenetics and Evolution, 87*, 65-79.
https://doi.org/10.1016/j.ympev.2015.03.013
Harrison, R. G. & Larson, E. L. (2014). Hybridization, introgression, and the nature of species
boundaries. *Journal of Heredity, 105*(S1), 795-809.
https://doi.org/10.1093/jhered/esu033
Hill, W. G. & Weir, B. S. (1988). Variances and covariances of squared linkage disequilibria in
finite populations. *Theoretical population biology, 33*(1), 54-78.
https://doi.org/10.1016/0040-5809(88)90004-4
Hivert, V., Leblois, R., Petit, E. J., Gautier, M., & Vitalis, R. (2018). Measuring genetic
differentiation from pool-seq data. *Genetics, 210*(1), 315-330.
https://doi.org/10.1534/genetics.118.300900
Hulse, S. V., Antonovics, J., Hood, M. E., & Bruns, E. L. (2023). Host–pathogen coevolution
promotes the evolution of general, broad-spectrum resistance and reduces foreign

pathogen spillover risk. *Evolution Letters, 7*(6), 467-477.
https://doi.org/10.1093/evlett/qrad051
Jin, W.-T., Gernandt, D. S., Wehenkel, C., Xia, X.-M., Wei, X.-X., & Wang, X.-Q. (2021).
Phylogenomic and ecological analyses reveal the spatiotemporal evolution of global
pines. *Proceedings of the National Academy of Sciences, 118*(20), e2022302118.
https://doi.org/10.1073/pnas.2022302118
Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics, 24(11), 1403-1405. https://doi.org/10.1093/bioinformatics/btn129
Kabir, M. S., Ganley, R. J., & Bradshaw, R. E. (2013). An improved artificial pathogenicity assay
for Dothistroma needle blight on *Pinus radiata. Australasian Plant Pathology, 42*(4),
503-510. https://doi.org/10.1007/s13313-013-0217-z
Lachance, J. & Tishkoff, S. A. (2013). SNP ascertainment bias in population genetic analyses:
why it is important, and how to correct it. *Bioessays, 35*(9), 780-786.
https://doi.org/10.1002/bies.201300014
Lenth, R. V. (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical*
Software, 69(1), 1 - 33. https://doi.org/10.18637/jss.v069.i01
Lind, B. M. (2021). GitHub.com/brandonlind/cmh_test: preprint release (Version 1.0.0). Zenodo.
http://doi.org/10.5281/zenodo.5083798
Lind, B. M. (2021). GitHub.com/CoAdapTree/varscan_pipeline: Publication release (Version
1.0.0). Zenodo. https://doi.org/10.5281/zenodo.5083302
Lind, B. M., Lu, M., Obreht Vidakovic, D., Singh, P., Booker, T. R., Aitken, S. N., & Yeaman, S.
(2022). Haploid, diploid, and pooled exome capture recapitulate features of biology and
paralogy in two non-model tree species. *Molecular Ecology Resources, 22*(1), 225-238.
https://doi.org/10.1111/1755-0998.13474
Lotan, J. E. & Critchfield, W. B. (1990). Lodgepole pine. In R. M. Burns & B. H. Honkala (Eds.),
Silvics of North America (Vol. Volume 1: Conifers, pp. 302-315). Washington, D.C.: USDA
Forest Service.
Lu, M., Feau, N., Obreht Vidakovic, D., Ukrainetz, N., Wong, B., Aitken, S. N., . . . Yeaman, S.
(2021). Comparative gene expression analysis reveals mechanism of *Pinus contorta*
response to the fungal pathogen *Dothistroma septosporum. Molecular plant-microbe*
interactions: MPMI, 34(4), 397-409. https://doi.org/10.1094/mpmi-10-20-0282-r
Lu, M., Krutovsky, K. V., Nelson, C. D., West, J. B., Reilly, N. A., & Loopstra, C. A. (2017).
Association genetics of growth and adaptive traits in loblolly pine (*Pinus taeda* L.) using
whole-exome-discovered polymorphisms. *Tree Genetics & Genomes, 13*(3), 57.
https://doi.org/10.1007/s11295-017-1140-1
MacLachlan, I. R., McDonald, T. K., Lind, B. M., Rieseberg, L. H., Yeaman, S., & Aitken, S. N.
(2021). Genome-wide shifts in climate-related variation underpin responses to selective
breeding in a widespread conifer. *Proceedings of the National Academy of Sciences,*
118(10), e2016900118. https://doi.org/10.1073/pnas.2016900118
Marroni, F., Pinosio, S., Zaina, G., Fogolari, F., Felice, N., Cattonaro, F., & Morgante, M. (2011).
Nucleotide diversity and linkage disequilibrium in *Populus nigra* cinnamyl alcohol
dehydrogenase (CAD4) gene. *Tree Genetics & Genomes, 7*(5), 1011-1023.
https://doi.org/10.1007/s11295-011-0391-5

McAllister, C. H., Cullingham, C. I., Peery, R. M., Mbenoun, M., McPeak, E., Feau, N., . . . Cooke, J.
E. K. (2022). Evidence of coevolution between *Cronartium harknessii* lineages and their
corresponding hosts, lodgepole pine and jack pine. *Phytopathology, 112*(8), 1795-1807.
https://doi.org/10.1094/phyto-09-21-0370-r
Moran, B. M., Payne, C., Langdon, Q., Powell, D. L., Brandvain, Y., & Schumer, M. (2021). The
genomic consequences of hybridization. *eLife, 10*, e69016.
https://doi.org/10.7554/eLife.69016
Neale, D. B., Wegrzyn, J. L., Stevens, K. A., Zimin, A. V., Puiu, D., Crepeau, M. W., . . . Langley, C.
H. (2014). Decoding the massive genome of loblolly pine using haploid DNA and novel
assembly strategies. *Genome Biology, 15*(3), R59. https://doi.org/10.1186/gb-2014-15-
3-r59
Paradis, E. & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics, 35*(3), 526-528.
https://doi.org/10.1093/bioinformatics/bty633
Peery, R. M., McAllister, C. H., Cullingham, C. I., Mahon, E. L., Arango-Velez, A., & Cooke, J. E. K.
(2021). Comparative genomics of the chitinase gene family in lodgepole and jack pines:
contrasting responses to biotic threats and landscape level investigation of genetic
differentiation. *Botany, 99*(6), 355-378. https://doi.org/10.1139/cjb-2020-0125
Quesada, T., Gopal, V., Cumbie, W. P., Eckert, A. J., Wegrzyn, J. L., Neale, D. B., . . . Davis, J. M.
(2010). Association mapping of quantitative disease resistance in a natural population of
loblolly pine (*Pinus taeda* L.). *Genetics, 186*(2), 677-686.
https://doi.org/10.1534/genetics.110.117549
R Core Team. (2018). R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing. https://www.R-project.org/. In.
Ramsfield, T., Myrholm, C., & Tomm, B. (2021). Natural infection of *Pinus contorta* var. *latifolia*,
Pinus banksiana and their hybrid by *Dothistroma septosporum* in Alberta, Canada.
Forest Pathology, 51(5), e12717. https://doi.org/10.1111/efp.12717
Rudolph, T. D. & Laidly, P. R. (1990). *Pinus banksiana* Lamb. Jack Pine. In R. M. Burns & B. H.
Honkala (Eds.), *Silvics of North America* (Vol. Volume 1: Conifers. Agriculture Handbook
654, pp. 280-293). Washington, D.C.: USDA Forest Service.
Rudolph, T. D. & Yeatman, C. W. (1982). Genetics of jack pine. In *Research Paper WO-38* (pp.
35-37). Washington, D.C.: USDA Forest Service.
Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals —
mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics,*
15(11), 749-763. https://doi.org/10.1038/nrg3803
Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., . . . Reyer, C. P.
O. (2017). Forest disturbances under climate change. *Nature Climate Change, 7*(6), 395-
402. https://doi.org/10.1038/nclimate3303
Simler-Williamson, A. B., Rizzo, D. M., & Cobb, R. C. (2019). Interacting effects of global change
on forest pest and pathogen dynamics. *Annual Review of Ecology, Evolution, and*
Systematics, 50(Volume 50, 2019), 381-403. https://doi.org/10.1146/annurev-ecolsys-
110218-024934

Singh, P., StClair, J. B., Lind, B., Cronn, R., Wilhelmi, N. P., Feau, N., . . . Yeaman, S. (2024).
Genetic architecture of disease resistance and tolerance in Douglas-fir trees. *New*
Phytologist. https://doi.org/10.32942/osf.io/s5rd7
Stocks, J. J., Metheringham, C. L., Plumb, W. J., Lee, S. J., Kelly, L. J., Nichols, R. A., & Buggs, R. J.
A. (2019). Genomic basis of European ash tree resistance to ash dieback fungus. *Nature*
Ecology & Evolution, 3(12), 1686-1696. https://doi.org/10.1038/s41559-019-1036-6
Suren, H., Hodgins, K. A., Yeaman, S., Nurkowski, K. A., Smets, P., Rieseberg, L. H., . . . Holliday, J.
A. (2016). Exome capture from the spruce and pine giga-genomes. *Molecular Ecology*
Resources, 16(5), 1136-1146. https://doi.org/10.1111/1755-0998.12570
Tam, V., Patel, N., Turcotte, M., Bossé, Y., Paré, G., & Meyre, D. (2019). Benefits and limitations
of genome-wide association studies. *Nature Reviews Genetics, 20*(8), 467-484.
https://doi.org/10.1038/s41576-019-0127-1
Taylor, S. A., Larson, E. L., & Harrison, R. G. (2015). Hybrid zones: windows on climate change.
Trends in Ecology & Evolution, 30(7), 398-406.
https://doi.org/10.1016/j.tree.2015.04.010
Wegrzyn, J. L., Liechty, J. D., Stevens, K. A., Wu, L. S., Loopstra, C. A., Vasquez-Gross, H. A., . . .
Neale, D. B. (2014). Unique features of the loblolly pine (*Pinus taeda* L.) megagenome
revealed through sequence annotation. *Genetics, 196*(3), 891-909.
https://doi.org/10.1534/genetics.113.159996
Wei, T. & Simko, V. (2017). R package "corrplot": Visulaization of a correlation matrix.
https://github.com/taiyun/corrplot.
Weiss, M., Sniezko, R. A., Puiu, D., Crepeau, M. W., Stevens, K., Salzberg, S. L., . . . De La Torre, A.
R. (2020). Genomic basis of white pine blister rust quantitative disease resistance and its
relationship with qualitative resistance. *The Plant Journal, 104*(2), 365-376.
https://doi.org/10.1111/tpj.14928
Welsh, C., Lewis, K., & Woods, A. (2009). The outbreak history of Dothistroma needle blight: an
emerging forest disease in northwestern British Columbia, Canada. *Canadian Journal of*
Forest Research, 39(12), 2505-2519. https://doi.org/10.1139/X09-159
Wheeler, N. C. & Guries, R. P. (1987). A quantitative measure of introgression between
lodgepole and jack pines. *Canadian Journal of Botany, 65*, 1876-1885.
https://doi.org/10.1139/b87-257
Wilson, J. F. & Goldstein, D. B. (2000). Consistent long-range linkage disequilibrium generated
by admixture in a Bantu-Semitic hybrid population. *The American Journal of Human*
Genetics, 67(4), 926-935. https://doi.org/10.1086/303083
Wood, L. J., Hartley, I. D., & Watson, P. (2009). Determining hybridization in jack pine and
lodgepole pine from British Columbia. *Wood and Fiber Science, 41*, 386-395.
Woods, A. J. (2003). Species diversity and forest health in northwest British Columbia. *The*
Forestry Chronicle, 79(5), 892-897. https://doi.org/10.5558/tfc79892-5
Woods, A. J., Coates, K. D., & Hamann, A. (2005). Is an unprecedented Dothistroma needle
blight epidemic related to climate change? *BioScience, 55*(9), 761-769.
https://doi.org/10.1641/0006-3568(2005)055[0761:IAUDNB]2.0.CO;2
Wu, M. C., Kraft, P., Epstein, M. P., Taylor, D. M., Chanock, S. J., Hunter, D. J., & Lin, X. (2010).
Powerful SNP-set analysis for case-control genome-wide association studies. *American*
Journal of Human Genetics, 86(6), 929-942. https://doi.org/10.1016/j.ajhg.2010.05.002

Wu, T. D. & Watanabe, C. K. (2005). GMAP: a genomic mapping and alignment program for
mRNA and EST sequences. *Bioinformatics, 21*(9), 1859-1875.
https://doi.org/10.1093/bioinformatics/bti310
Yeaman, S., Hodgins, K. A., Lotterhos, K. E., Suren, H., Nadeau, S., Degner, J. C., . . . Aitken, S. N.
(2016). Convergent local adaptation to climate in distantly related conifers. *Science,*
353(6306), 1431-1433. https://doi.org/10.1126/science.aaf7812
Yeatman, C. W. & Teich, A. H. (1969). Genetics and breeding of jack and lodgepole pines in
Canada. *The Forestry Chronicle, 45*(6), 428-433. https://doi.org/10.5558/tfc45428-6
761