Genetic architecture underlying response to the fungal pathogen *Dothistroma septosporum*

in lodgepole pine, jack pine, and their hybrids

Abstract

In recent decades, *Dothistroma* needle blight (DNB), a pine tree disease caused by the fungal
pathogen *Dothistroma septosporum,* has severely damaged lodgepole pine (*Pinus contorta*
Dougl. ex. Loud.) in British Columbia, Canada. The pathogen has already shown signs of host
shift eastward to lodgepole × jack hybrid pine populations, and possibly into pure jack pine
(*Pinus banksiana* Lamb.). However, we have little knowledge about mechanisms of resistance to
D. septosporum, especially the underlying genetic basis in pines. In this study, we conducted
controlled inoculations to induce infection by *D. septosporum* and performed a genome-wide
case-control association study with pooled sequencing (pool-seq) data to dissect the genetic
architecture underlying response in lodgepole pine, jack pine, and their hybrids. We identified
candidate genes associated with *D. septosporum* response in lodgepole pine and in lodgepole ×
jack hybrid pine samples. We also assessed genetic structure in hybrid populations and inferred
how introgression may affect the distribution of genetic variation involved in *D. septosporum*
response in the studied samples. These results can be used to develop genomic tools to evaluate
DNB risk, guide forest management strategies, and potentially select for resistant genotypes.
20
21
22 **KEY WORDS**
23 Conifer, fungal pathogen resistance, F_{ST}, hybrid, introgression, pool-GWAS
24

1 | INTRODUCTION

Pests and pathogens constantly threaten forest trees, and their impacts are changing as climates
are altered (Seidl et al., 2017; Simler-Williamson et al., 2019). Knowledge about the genetic
architecture of pathogen resistance and the identified candidate resistance (R) genes provides the
basis for breeding practices to improve tree resilience to forest pathogens. Genes and alleles that
confer disease resistance or tolerance can eventually be used to develop genomic tools for
rapidly selecting resistant genotypes (Isabel et al., 2019). Current breeding practices for forest
trees involve recurrent selection, which includes repeated cycles of selection, breeding, and
testing, to search for genotypes with better disease resistance, and then deployment of these
selected tree genotypes (White, 2004). This takes years of effort to design and implement.
Increasingly, genome editing is used to develop crops resistant to disease (van Esse et al., 2020).
CRISPR/Cas9 system has been applied to enhance the disease resistance in citrus (Jia et al.,
2017; Wang et al., 2019), apple (Zhou et al., 2020), and timberland species (Min et al., 2022).
Given the long generation time in conifers and the frequent threats from diseases and insects,
identifying R genes that can be targeted for selection or editing is a useful step in developing
resistant genotypes for reforestation.
43

Dothistroma needle blight (DNB), which is caused by the fungal pathogen *Dothistroma*
septosporum, has caused increased damage to lodgepole pine (*Pinus contorta* Dougl. ex. Loud.)
in recent decades in Western Canada (Dale et al., 2011; Woods, 2003). DNB is a foliar disease of
a wide range of pine trees (Gibson, 1972), with infected trees showing reddish-brown bands in
needles, defoliation, and growth reduction. Historically, DNB has had only minor impacts on
native forest trees in western North America (Woods et al., 2005). Recently, however, DNB is
causing increased outbreak incidence and extent, which is correlated with climate change (Welsh
et al., 2009). While *D. septosporum* tends to be more common in moist western forests closer to
the Pacific Ocean, *Dothistroma*-like symptoms were observed in drier lodgepole and lodgepole ×
jack hybrid pines in northern Alberta in 2012 and 2013 with further tests confirming *D.*
septosporum was the cause of the disease observed (Ramsfield et al., 2021). Feau et al. (2021)
performed a controlled inoculation experiment and demonstrated that lodgepole pine, jack pine
(*Pinus banksiana* Lamb.), and their hybrids are susceptible to *D. septosporum*, with jack pine
showing a higher disease severity than lodgepole or hybrid pine. Although there is no report
showing DNB infection in natural jack pine forests, their higher susceptibility to *D. septosporum*
raises serious concern.
60
The history of host-pathogen coevolution might explain the stronger resistance to DNB infection
in lodgepole than in jack pine. A previous study has identified candidate resistance genes by
RNA-seq analysis on experimentally infected lodgepole pine seedlings (Lu et al., 2021). These
candidate genes include 43 genes that are present in the plant-pathogen interaction pathway, as
well as nine R genes that contained sites under positive selection. It seems that both constitutive
(baseline) and induced (defenses activated after being attacked) defenses have been developed in
lodgepole pine. Moreover, the host-pathogen coevolution can promote a broad-spectrum
resistance (Hulse et al., 2023). It is likely that the induced and increased general defense have
contributed to the lodgepole pine's resistance to DNB infection, however, little is known about
resistance in jack pine. Previous studies have compared the molecular responses between
lodgepole and jack pine when subjected to other biotic agents. Lodgepole and jack pine showed
unique monoterpene profiles in response to mountain pine beetle (Hall et al., 2013) and
significant differences in chitinase gene expression in response to the fungal pathogens
Grosmannia clavigera and *Cronartium harknessii* (Peery et al., 2021). These previous studies
used particular gene families to examine the species-specific pathogen responses in a focused set
of candidates, whereas a genome-wide gene survey might give us a broad perspective.
77
To dissect the genetic architecture of pathogen resistance and to identify candidate genes,
researchers have employed genetic markers, mainly SNPs, and methods like quantitative trait
locus (QTL) mapping and genome-wide association analysis (GWAS). Polygenic disease
responses, comprising numerous loci of small effect, appear to be the rule (Stocks et al., 2019).
Large numbers of SNPs were associated with the pitch canker resistance in *Pinus taeda* L. (De
La Torre et al., 2019; Lu et al., 2017; Quesada et al., 2010), with the white pine blister rust
resistance in *Pinus lambertiana* Dougl. (Weiss et al., 2020), and with the *Heterobasidion* root rot
resistance in *Picea abies* (L.) Karst. (Capador-Barreto et al., 2021). To identify the candidate R
genes, most GWAS studies on forest trees employ individual-based methods, with both
phenotypic and genotypic data collected on each sampled individual. Case-control methods have
been developed to study disease where individuals are pooled into healthy vs. infected groups,
but such approaches typically still use individual-based sequencing to call genotypes before

pooling (Wu et al., 2010). When multiple replicates are available, it is possible to combine pool-seq and the case-control design: Endler et al. (2016) analyzed the genetic differences underlying abdominal pigmentation variation among *Drosophila* populations, while Stocks et al. (2019) identified SNPs associated with low versus high ash dieback damage in *Fraxinus excelsior* L. using this approach. Recently, Singh et al. (2024) used a case-control pool-seq approach similar to that deployed here to search for signatures of resistance to Swiss needle cast in *Pseudotsuga menziesii* (Mirb.) Franco. Pool-seq is a cost-effective alternative approach to sequencing of individuals, but the weakness and limitations are also evident, such as unequal representation of individuals and suboptimal allele frequency estimate when the pool size is small, as well as misaligned mapping and sequencing errors (Schlötterer et al., 2014). Nonetheless, in a study on conifer genotyping, Lind et al. (2022) found that allele frequencies estimated from pooled DNA sequencing samples were highly correlated with frequencies estimated from individual sequencing.

An important factor in understanding the genetics of resistance in these species is whether there is any interplay between any alleles conferring resistance and introgression between the species. Lodgepole pine mainly inhabits montane regions of western North America (Figure 1), throughout the interior of British Columbia and the Rocky Mountains of Alberta (Lotan & Critchfield, 1990), whereas jack pine is widely distributed across boreal forests in eastern North America, extending in Canada from Northwest Territories to Nova Scotia (Rudolph & Laidly, 1990). Lodgepole and jack pine evolved in allopatry, with an estimated divergence time of ~8 million years before present (Hao et al., 2015), but hybridized in areas of contact, primarily in central and northwestern Alberta (Burns et al., 2019; Rudolph & Yeatman, 1982). Following secondary contact of these two species, additional patchy regions of introgression have been developed in central Alberta north through to the Northwest Territories and northeastern British Columbia to the Alberta-Saskatchewan border (Burns et al., 2019). Natural hybrids display a wide range of phenotypes that are intermediate or more typical of one parent, depending on the level of introgression (Wood et al., 2009; Yeatman & Teich, 1969). The hybrid zone between these species has been very well studied by Cullingham et al. (2012), who found that the lodgepole × jack pine hybrid zone presents a mosaic zone with variable introgression and patchy distributions of hybrids (Burns et al., 2019). While it is clear that these species share a broad hybrid zone, it is unclear whether any alleles for disease resistance will be more or less introgressed than the average region in the genome.

In this study, our aim is to identify the genetic basis of DNB resistance in the studied pine species and to explore spatial patterning in any identified loci. The specific objectives of our study were to: 1) develop a GWAS case-control approach using pool-seq samples to identify loci with consistent associations with *D. septosporum* response across replicates; and 2) compare the genetic architecture underlying DNB resistance within the studied pine trees and infer how introgression may affect the genetic basis of *D. septosporum* responses. This study provides an important step towards identifying candidate genes to develop tools for screening trees resistant to DNB infection.

2 | METHODS

2.1 | Plant materials

Seeds were obtained from 40 natural seedlots across Western Canada (Figure 1, seedlot numbers
and locations can be found in Table S1, seed contributors http://adaptree.forestry.ubc.ca/seed-
contributors/), including 25 lodgepole pine seedlots from British Columbia (BC_pl) and three
from Alberta (AB_pl), seven lodgepole × jack hybrid pine seedlots from Alberta (AB_px), and
five jack pine seedlots from Alberta (AB_pj). The range maps were downloaded from
https://sites.ualberta.ca/~ahamann/data/rangemaps.html (Hamann et al., 2005). When the seeds
were collected in the wild, they were assigned to pure lodgepole pine, pure jack pine, or
lodgepole × jack hybrid pine, based on their location and morphological traits such as cone and
branch characteristics, microfibril angle, and cell area (Wheeler & Guries, 1987; Wood et al.,
2009; Yeatman & Teich, 1969). The ancestry of the collected seeds were genotyped using 11
microsatellite loci by Cullingham et al. (2012). Briefly, seeds were germinated to obtain
seedlings, then DNA was isolated from the seedlings. DNA was used to amplify 11
microsatellite loci and allele sizes were determined for genotyping as described by Cullingham et
al. (2011).
150
For the present study, seeds from the 40 seedlots were used to grow seedlings in a greenhouse at
University of British Columbia, Vancouver, BC, for their first year (a flow chart of experimental
procedure is shown in Figure S1). Resistance was phenotyped by inoculating 100 individuals per
seedlot with one of two *D. septosporum* isolates (D1 & D2, 50 individuals of each isolate). D1
was isolated from needles of an infected lodgepole × jack hybrid pine seedlot in Alberta,
Canada, while D2 was isolated from an infected lodgepole pine seedlot in the Kispiox Valley
region close to Smithers in Northwestern British Columbia, Canada (Feau et al., 2021; Ramsfield
et al., 2021). The seedlots were identified as lodgepole or lodgepole × jack hybrid pine as
aforementioned (Cullingham et al., 2012). The controlled inoculation experiment was performed
as described in Feau et al. (2021). Briefly, the one-year-old seedlings were placed in a
completely randomized experimental design in a growth chamber with the condition of 16 hours
daylight at 20 ℃, 8 hours a night at 12 ℃, and minimum relative humidity of 80 %. *D.*
septosporum conidia were harvested from colonies grown for 10~16 days on *Dothistroma*
sporulation medium plates and then suspended in sterile distilled water. Each seedling was
sprayed twice at 10-day interval with a standard inoculum of approximately 3 mL of 1.6×10^6
conidia mL^{-1}, using a household trigger action atomizer. Once the needles were dry (~45
minutes), the seedlings were wrapped in transparent plastic bags and kept in the growth chamber.
After 48h the plastic bags were removed and the seedlings were kept in the growth chamber for
15 weeks till phenotyping. A mister spraying tap water was activated hourly for 3 minutes during
the trial to maintain needle wetness. Control seedlings, at least one for each seedlot, were
inoculated with sterile water.
172
After 15 weeks since the beginning of the experiment, seedlings were rated for proportion of
necrotic needles with red bands and/or fruiting bodies according to a disease severity scale of 1
to 5 (1 = least, 5 = most disease severity). Following disease rating, the ten most- and ten least-
infected individuals per seedlot (five individuals for each *D. septosporum* isolate) were identified
and retained for DNA extraction. Genomic DNA of each of these individuals was extracted as
described in Lind et al. (2022), using the Nucleosin 96 Plant II Core kit (Macherey–Nagel
GmbH & Co. KG, Germany) on an Eppendorf epMotion 5075 liquid-handling platform. The
DNA of five most-infected and five least-infected individuals by each of the two *D. septosporum*
isolates was combined in equimolar amounts to compose four pooled libraries per population,

two susceptible (D1-S, D2-S) and two resistant libraries (D1-R, D2-R). Hence,160 pooled DNA libraries were generated.

FIGURE 1 Sampling locations (Table S1) of the 40 pine seedlots analyzed in this study, plotted within the natural range of lodgepole and jack pine in Canada and USA. The topright map shows the context of the sampling locations. The range maps were downloaded from https://sites.ualberta.ca/~ahamann/data/rangemaps.html. BC_pl (green diamonds) — lodgepole pine seedlots from British Columbia (N=25); AB_pl (orange diamonds) — lodgepole pine seedlots from Alberta (N=3); AB_px (purple diamonds) — lodgepole × jack hybrid pine seedlots from Alberta (N=7); AB_pj (brown diamonds) — jack pine seedlots from Alberta (N=5).

2.2 | Sequence capture and pool-seq genotyping

The capture probes were designed based on an existing lodgepole pine sequence capture array (Suren et al., 2016) by removing the probes that did not yield successful genotyping in Yeaman et al. (2016) and adding new probes derived from newly assembled *D. septosporum*-induced genes. Probe sequences of the existing capture array were aligned to the reference genome using

GMAP v2019-03-15 (Wu & Watanabe, 2005). Probes that covered the genomic regions with called SNPs in the dataset from Yeaman et al. (2016) were retained, otherwise the probes were discarded. Since there is no available lodgepole pine reference genome, a masked loblolly pine reference genome Pita.2_01.masked3k2.fa (https://treegenesdb.org/FTP/Genomes/Pita/v2.01/genome/) (Neale et al., 2014) was used instead. Loblolly pine is a closely related species to lodgepole pine (Jin et al., 2021). New probes targeting *D. septosporum*-induced genes were designed based on a lodgepole pine reference transcriptome assembled using the RNA-seq data of *D. septosporum* infected lodgepole pine samples (Lu et al., 2021). To avoid duplicates, only those *D. septosporum*-induced genes with low homology to the retained probe sequences were used to design the new probes. To do so, the retained probe sequences were aligned to this transcriptome using blastn v2.9.0 with an E-value of 1e-10. A total of 8,778 *D. septosporum*-induced genes did not have any aligned probe sequences. These non-duplicate *D. septosporum*-induced genes were subsequently aligned to the reference genome to predict the exon-intron boundaries using GMAP v2019-03-15. Exon sequences from these induced genes with a length of at least 100 bp were combined with the previously designed working probe sequences, and this combined set of sequences was submitted to Roche NimbleGen (Roche Sequencing Solutions, Inc., CA USA) for Custom SeqCap EZ probe design (design name: 180321_lodgepole_v2_EZ). This updated lodgepole pine sequence capture array has a capture space of 44 Mbp, containing roughly 35,467 assembled genes. Most lodgepole pine genes responsive to environment stress and fungal pathogen attack were included in the current capture probe design, though genes expressed in different development periods might be missed, genes that have evidence of substantial expression have been covered in this capture probe design.

The capture libraries for each of the 160 pools were constructed following NimbleGen SeqCap EZ Library SR User's Guide and as described in Lind et al. (2022). Then the R (resistant) and S (susceptible) libraries (two libraries per capture, R1+S1 or R2+S2, indexed with different barcodes) per population and per isolate were combined for sequence capture and enrichment. Sequencing was performed using the Illumina NovaSeq 6000 S4 PE 150 platform in Centre d'expertise et de services Génome Québec. Our in-house pool-seq pipeline (Lind, 2021) was employed to align the reads to the reference genome and call SNPs. For raw SNPs, only bi-allelic loci in regions without annotated repetitive elements or potentially paralogous genes were retained. The annotated repetitive elements were acquired from the loblolly pine genome annotation (Wegrzyn et al., 2014). The potentially paralogous genes were identified as described in Lind et al. (2022) using haploid megagametophyte sequences, for the heterozygous SNP calls for haploid sequences are likely to represent misalignments of paralogs. Afterwards, the SNP loci with depth (DP) < 10, DP > 400, global minor allele frequency < 0.05, or > 25% missing data were also removed.

2.3 | Genetic structure analyses

The genetic structure among the 160 pooled samples, which represented four pooled libraries (D1-R, D1-S, D2-R, D2-S) for each of the 40 populations, was detected by using principal component analysis (PCA), reconstructing a phylogenetic tree, and by assessing correlation of allele frequencies and population differentiation (F_{ST}) between samples using unlinked SNPs. To reduce the impact of linkage disequilibrium on estimation, the SNP set was thinned using

vcftools (Danecek et al., 2011), so that no two sites were within 100,000 bp. PCA and genetic
distance were calculated using the major allele depth and the R package "adegenet" (Jombart,
2008). An unrooted phylogenetic tree was constructed using the distance output and the
Neighbour-Joining algorithm, which was implemented by the R package "ape" (Paradis &
Schliep, 2019). A bootstrap analysis was performed using 5000 bootstrap replicates. Correlation
coefficients of major allele frequencies between all pairs within the 160 samples were calculated
and plotted using the R package "corrplot" (Wei & Simko, 2017). F_{ST} was estimated between all
pairs within the 160 samples using the R package "poolfstat" (Hivert et al., 2018).
258
259 **2.4 | Dissection of genetic architecture underlying *D. septosporum* response**
260
The alleles and candidate genes associated with *D. septosporum* response were detected using
the pool-GWAS method. A GWAS case-control approach was developed using pool-seq samples
to identify allele frequency differences between susceptible and resistant pines inoculated with
D. septosporum. This method was based on Cochran-Mantel-Haenszel (CMH) method and
implemented in our in-house pipeline (Lind, 2021). While sequencing read depth was commonly
used as an estimation of allele count in such applications of test (Futschik & Schlötterer, 2010;
Schlötterer et al., 2014), reliable results would be only achieved when the haploid size of the
sample is much greater than the depth of coverage (so that most reads arise from uniquely
sampled haplotypes). As the pool size used in the present study (diploid size of 2N=10) is often
smaller than the sequencing depth of coverage, read counts were converted into allele
frequencies and then multiplied by the ploidy to yield an approximate count for each pool that
represents the real replication level. A simulation was conducted to compare the false positive
rate from the CMH test using uncorrected or corrected allele counts across a range of number of
demes, individuals, and depths. Results showed that the false positive rate of the CMH method is
high, especially when the sequencing depth is much higher than the pool size (Figure S2a).
Using corrected allele counts, the allele frequencies multiplied by the ploidy (N=10) tends to
lower the false positive rate (Figure S2b). Thus, the corrected allele counts were used for CMH
test.
279
As the genetic structure analysis showed that lodgepole × jack hybrid pine and jack pine samples
tend to cluster, jack pine samples and lodgepole × jack hybrid pine samples were combined as
jack-hybrid samples for downstream analyses. Pool-GWAS was performed separately within
lodgepole and jack-hybrid pine samples. Analysis of R and S samples was conducted by
combining individuals inoculated with D1 or D2 isolates to maximize statistical power. To
identify candidate genes, SNPs in the top 1% of $-\log_{10}(p)$ values from a pool-GWAS test were
classified as outliers. As linkage disequilibrium can amplify signatures of selection, candidate
resistance genes were identified as those with a large number of outlier SNPs relative to the
average genome-wide expectation, as represented by an index based on the binomial distribution
as per Yeaman et al. (2016). The identified loci on the enriched genes were realigned to the 12
lodgepole pine linkage groups (MacLachlan et al., 2021) and Manhattan plots were drawn. The
lodgepole pine transcriptome (Lu et al., 2021) was used to annotate the reference genome. The
aligned transcripts to the reference genome were identified as those with a minimum alignment
identity of 90 % and a minimum alignment coverage of 85 % using GMAP v2019-03-15.
294
295

2.5 | Linkage disequilibrium (LD) between loci

A non-linear model (Hill & Weir, 1988) was used to estimate the decay of LD with physical distance. Pairwise correlation (r^2) of allele frequencies between loci on the same scaffold were calculated. The r^2 values and the physical distances between loci on all scaffolds (genome-wide) or on scaffolds containing the identified top candidate genes (significant) were used to fit the non-linear model as described by Marroni et al. (2011). The LD decay graph was plotted using R (R Core Team, 2024).

2.6 | Introgression

To study patterns of introgression affecting different regions of the genome and among different samples, the mean pairwise F_{ST} was calculated for each locus using the R package "poolfstat". Loci with high F_{ST} values have high genetic differentiation, which may represent species barriers or genomic regions under divergent selection, while loci with low F_{ST} values imply introgression. To estimate the genetic similarity of studied jack pine and lodgepole pine populations, the five jack pine samples (p36 to p40) and six pure lodgepole samples (the six westmost lodgepole pine samples, p15, p16, p18, p20, p23, p25) were used to calculate F_{ST} values, which were then averaged. To evaluate whether regions associated with *D. septosporum* response had atypical patterns of introgression, the F_{ST} values of the top 1 % *D. septosporum* response outliers identified from lodgepole pine (N = 3,358) and those from jack-hybrid pine (N = 3,041) were compared, with F_{ST} values of randomly drawn SNPs (N = 3,400).

Similarly, F_{ST} between the jack-hybrid samples (p29 to p40) and six pure lodgepole samples (p15, p16, p18, p20, p23, p25) were averaged to estimate the genetic similarity of studied jack-hybrid and lodgepole pine. Different genomic regions were represented by different SNP sets, including *D. septosporum* response outliers identified from lodgepole pine (N = 3,358) and outliers identified from jack-hybrid pine (N = 3,041), as well as the unlinked SNP set (N = 31,716), which was used for detecting genetic structure and representing random genomic region. The genetic similarity (F_{ST}) of jack-hybrid and lodgepole pine was regressed to the longitude of these jack-hybrid samples. The regression slopes for the three SNPs sets were compared using analysis of covariance method implemented by the R package "lsmeans" (Lenth, 2016; R Core Team, 2024).

3 | RESULTS

3.1 | Patterns of genetic structure
After filtering, a total of 364,691 SNP loci were retained for downstream analyses, and most loci had a minor allele frequency between 0.05 and 0.1 (Figure S3). A thinned set of 31,716 unlinked SNPs was used for studying genetic structure among the 160 pooled samples (Figure 2). The most prominent patterns of structure were associated with species (Figure 2a & b, Figure S4), with lodgepole pine samples readily distinguishable from jack and hybrid pine samples. Hybrid samples clustered more closely with jack than lodgepole pine. Compared to those sampled from the eastern hybrid zone, western hybrid pines tend to be more similar to lodgepole pine (samples were arranged according to longitude from left to right in Figure 2 c & d). Given this clear

342 pattern of genetic structure, we conducted GWAS in lodgepole pine and jack-hybrid pine
343 samples separately.
344

345
346
FIGURE 2 Genetic structure among the 160 pooled samples. a) Principal component analysis. b)
Unrooted phylogenetic tree. The bootstrap values for each branch were shown in Figure S4. c)
Correlation plot of the major allele frequencies between samples. d) Pairwise F_{ST} values between
pools. BC_pl — lodgepole pine samples from British Columbia; AB_pl — lodgepole pine
samples from Alberta; AB_px — lodgepole × jack hybrid pine samples from Alberta; AB_pj —
jack pine samples from Alberta. D1 — *D. septosporum* isolate 1; D2 — *D. septosporum* isolate
2. R — resistant tree; S — susceptible tree. In c) and d), each column represents a pooled
sample. The samples were arranged from left to right according to the longitude of sampling
location, and four pooled libraries per seedlot (D1-R, D1-S, D2-R, D2-S) were arranged together.
356
357
358 **3.2 | Signatures of association to infection response**
359
Out of the 100 top-ranked candidate genes in each test (Table S2), three were identified within
both the lodgepole and jack-hybrid pine samples (Figure 3). This is significantly more than the
hypergeometric expectation, based on the probability of overlap for two draws of 100 genes from
20,026 genes. A previous study anchored 10,093 scaffolds on the lodgepole pine linkage map
(MacLachlan et al., 2021), so we plotted SNPs and labeled the enriched plant-pathogen
interaction candidate genes on the 12 linkage groups (Figure 3). The three genes identified in
both species encode ABC transporter, F-box Kelch-repeat protein, and serine/threonine protein

kinase (F-box protein and serine protein kinase were highlighted in Figure 3, ABC transporter
genes were not included in the linkage groups, so they are not included in the figure). We also
checked the enrichment of differentially expressed genes (DEGs) in the top 100 ranked genes.
Out of 631 DEGs identified in the previous study (Lu et al., 2021), four were also candidate
genes in the present study. Fisher's Exact test showed there is no significant enrichment of DEGs
in the candidate genes.
373

374
375
FIGURE 3 Manhattan plot showing SNPs associated with *D. septosporum* response within
lodgepole and jack-hybrid pine samples. The x axis represents 12 linkage groups. The y axis

represents significance of associations identified from pool-GWAS. Candidate genes were
identified using a binomial test for enrichment of outlier SNPs per gene. Candidate genes and
their surrounding 50 kbp regions are highlighted in red. Of the three top-ranked candidate genes
identified within both the lodgepole and jack-hybrid pine samples, one was not present in the
linkage map, while the other two are highlighted in blue ("Serine" on linkage group 4 and "F-
box" on linkage group 10). Labels in the graphs represent candidate genes encoding plant-
pathogen interaction proteins: bHLH — bHLH transcription factor; Serine — serine/threonine-
protein kinase; Auxin — auxin response factor; LRR — proteins with leucine-rich repeat; F-box
— F-box Kelch-repeat protein; MADS — MADS transcription factor; MYB — MYB
transcription factor; WRKY — WRKY transcription factor; TIF — translation initiation factor.
The candidate gene IDs (in Table S2) are: In lodgepole pine, bHLH — Scaffold_2015-465162-
569336; Serine_1 — Scaffold_3728-494595-736720; Serine_2 — Scaffold_3141-1-105566;
Serine_3 — Scaffold_1072-1299306-1374553; Auxin — Scaffold_3823-1571412-1689102;
LRR — Scaffold_1361-487806-570406; F-box — Scaffold_3698-733755-839782; in jack-
hybrid pine, Serine — Scaffold_3728-494595-736720; MADS — Scaffold_427-2324701-
2717580; F-box_1 — Scaffold_3698-733755-839782; MYB — Scaffold_4478-441537-627389;
F-box_2 — Scaffold_342-1965787-2253515; WRKY — Scaffold_830-1136302-1239031; TIF
— Scaffold_207-7583953-7695067.
396
397
Loci on scaffolds with outliers (the SNPs with top 1 % of -$\log_{10}(p)$ values from pool-GWAS test)
tend to exhibit a more gradual decay in LD compared with the genome-wide loci on all scaffolds
(Figure 4), with jack-hybrid pine outliers having a much slower decay rate than lodgepole pine.
In lodgepole pine, the distance at which LD decays to half of its maximum value is 96 bp for
genome-wide loci, and 174 bp for outlier loci. In jack-hybrid pine, the half LD decay distance is
1,357 bp for genome-wide loci, and 68,215 bp for outlier loci. The slower LD decay pattern for
the outliers may indicate selection on these identified outlier loci.
405

406
407
408 FIGURE 4 Linkage disequilibrium (LD) decay for genome-wide loci and outlier loci identified
409 from lodgepole and jack-hybrid pine samples. The r^2 values (pairwise correlation of allele
410 frequencies between loci) and the physical distances between loci were used to fit the non-linear
411 model as described by Marroni et al. (2011).
412

3.3 | Introgression

By calculating average F_{ST} between pairs of jack pine and lodgepole pine samples, we found that
the *D. septosporum* response outliers, which were identified separately in lodgepole or jack-
hybrid samples, have higher F_{ST} values than randomly drawn SNPs (Wilcoxon test *p*-value <
0.01, Figure 5a & 5b), with outliers identified in jack-hybrid samples showing higher F_{ST} values
than those identified in lodgepole samples (Wilcoxon test *p*-value < 0.01). These results suggest
that the outliers associated with disease response constitute or are linked to genomic regions for
divergent selection or species barriers, rather than facilitating introgression. F_{ST} estimates
between jack-hybrid samples and a set of pure lodgepole pine samples showed a pattern of
increased differentiation with distance eastwards of the jack-hybrid samples (Figure 5c), which
was higher for the outliers than randomly chosen SNPs. Given the small number of samples,
significance of linear model fitting is borderline, with *p*-value = 0.06 for outliers identified in
lodgepole, *p*-value = 0.05 for outliers identified in jack-hybrid, and *p*-value = 0.06 for randomly
chosen SNPs (df = 10). The differences between regression slopes on different gene sets
(lodgepole outliers, jack-hybrid outliers, and unlinked SNPs) were not significant (*p*-values of
pairwise comparisons > 0.3). Given the limited sample size, it is not possible to test whether a
linear vs. discontinuous model would fit better to these data, but these results suggest that
isolation-by-distance is occurring here.

433
434
435

FIGURE 5 F_{ST} values of *D. septosporum* response outliers (a & b) and regression patterns of F_{ST}
values with longitude. For a & b, the F_{ST} values were calculated between the jack pine samples
(p36 to p40) and six pure lodgepole samples (p15, p16, p18, p20, p23, p25). F_{ST} values of *D.*
septosporum response outliers were identified from lodgepole (N=3,358) and jack-hybrid
(N=3,041) samples, and randomly drawn SNPs (N=3,400). Outliers identified from both species
have higher F_{ST} values than randomly drawn SNPs (*p*-value < 0.01). For c, the F_{ST} values were
calculated between jack-hybrid samples (p29 to p40) and six pure lodgepole samples (p15, p16,
p18, p20, p23, p25) and then averaged for each jack-hybrid sample. Different genomic regions
were represented by different SNP sets, including lodgepole outliers (N=3,358), jack-hybrid

outliers (N=3,041), and unlinked SNPs (N=31,716). The F_{ST} values represented by different SNP
sets were regressed to the longitude of these jack-hybrid samples. Differences between slopes are
not significant (ANCOVA, *p*-value > 0.3).
448
449   **4 | DISCUSSION**
450
In recent decades, the fungal pathogen *D. septosporum* has become an increasing threat to
lodgepole pine in British Columbia (Feau et al., 2021; Welsh et al., 2009). Although
Dothistroma needle blight (DNB) has not been observed in natural stands of jack pine in Alberta,
Ramsfield et al. (2021) confirmed jack pine's susceptibility to infection by *D. septosporum* using
a trial under natural conditions. The potential of climate change to allow this pathogen to spread
across the wide natural distribution of jack pine raises serious concerns. The hybrid zone of
contact between the sister species lodgepole and jack pine may act as a bridge, allowing this
fungal pathogen host shift (Taylor et al., 2015). The genetic basis of pathogen resistance in each
species is therefore an important factor affecting the potential for host shift. To study the genetics
of resistance to DNB, we analyzed the genetic architecture underlying *D. septosporum* response
using the study samples of lodgepole pine, jack pine, and their hybrids. We identified candidate
genes associated with *D. septosporum* response and inferred the impact of introgression on the
genetic structure underlying *D. septosporum* response in the studied samples. We detected
largely non-overlapping sets of candidate genes in lodgepole and jack-hybrid pine in response to
D. septosporum, which might be shaped by different evolutionary forces of selection and
introgression. Alternatively, the non-overlapping sets of candidate genes might reflect that many
non-causal genes exist in the top 1 % GWAS outliers of each species, given that we did not test
for significance at the gene-level. In either case, it seems clear that resistance to DNB is a highly
polygenic trait with alleles of small effect.
470
This study is the first report of genome-wide scanning of *D. septosporum* response genes in
lodgepole pine, jack pine, and their hybrids. We identified a few top candidate genes encoding
leucine-rich repeats (LRR)-containing protein domain and serine threonine kinases in lodgepole
pine samples (Figure 3, Table S2). Plant breeders have long used nucleotide-binding leucine-rich
repeat (NB-LRR) receptor resistance genes (R genes) for preventing or reducing diseases in
crops. When attacked by infectious pathogens, plant NB-LRR receptors can recognize pathogen
effectors by either direct or indirect mechanisms (Dodds & Rathjen, 2010). Such strategy is
called effector-triggered immunity (ETI), which occurs after a basal resistance, pathogen-
associated-molecular-patterns-triggered immunity (PTI). In a previous study using RNA-seq
data, Lu et al. (2021) identified genes involved in PTI and ETI, as well as R genes with positive
selection signals in lodgepole pine. These results indicate that the long coevolutionary history
between lodgepole pine and *D. septosporum* (Capron et al., 2021; Welsh et al., 2009) has given
rise to some tolerance or resistance mechanism against DNB. Since most of the identified
associated markers are likely not causal but rather are in LD with the causal loci, the observed
fast LD decay rate in lodgepole pine implies promising fine mapping resolutions as the identified
markers are likely to be proximal to the causative genes.
487
On the contrary, jack pine, as a potential new host, may not have coevolved such a specific
defense, and the long LD blocks surrounding jack-hybrid pine candidate genes, which are typical
of hybrid zones, limit the resolution of association mapping, so it may be difficult to locate the

causative genes for follow-up studies. Additionally, we caution that the capture probes were
designed using lodgepole pine not jack pine genes, ascertainment bias might give rise to the
underrepresentation of jack pine alleles. However, we can use these jack-hybrid pine candidate
genes to infer how introgression affects the frequencies of disease-resistant alleles. Since we
analyzed pool-seq data and lacked the pure jack pine samples out of the hybrid zone, it is not
possible to analyze parental origin information, thus we inferred the ancestry using clustering.
The clustering patterns in Figure 2 show that most hybrid pine trees in our study had greater jack
pine than lodgepole pine ancestry, likely in part due to our limited sampling of jack pines from
further Eastwards. Western hybrid pines tend to be more genetically similar to lodgepole pine,
while eastern hybrid pines tend to be more genetically dissimilar to lodgepole pine. This is a
typical characteristic for ongoing introgression and hybridization in the contact zone with a
possible trend of lodgepole pine expanding its range eastward.
503
In hybrid zones, alleles in some genomic regions are able to introgress across species boundaries,
whereas alleles that constitute species barriers or are under divergent selection will remain
differentiated (Harrison & Larson, 2014). We found significantly high F_{ST} values between jack
and lodgepole pine samples for the candidate genes associated with *D. septosporum* infection
response. This can occur if these candidate genes tend to reside in genomic regions with
restricted gene flow, driven by linkage with genes for divergent selection or species barriers.
Bechsgaard et al. (2017)showed that R genes can adaptively introgress between closely related
species. Since lodgepole pine has a long coevolutionary history with *D. septosporum* (Capron et
al., 2021; Welsh et al., 2009), the introgression of lodgepole pine genes into hybrids may
increase resistance to *D. septosporum*. However, variation in F_{ST} along longitude (Figure 5c) did
not differ significantly between background and candidate loci, so we cannot draw strong
conclusions about adaptive introgression here.
516
517 **5 | CONCLUSION**
518
The recent outbreaks of DNB in lodgepole pine and its ability to infect jack pine pose a pressing
need for understanding the genetic architecture of resistance to *D. septosporum*. The present
study provides an array of candidate genes associated with *D. septosporum* response using pool-
GWAS. The identified candidate genes within lodgepole and jack-hybrid pines are largely non-
overlapping, and the adaptive introgression is yet to be found. This knowledge can be used to
develop genomic tools to minimize DNB risk and guide forest management strategies. Further
testing and validation of these associations using a SNP array is currently underway.
526
527
528 **DATA AVAILABILITY STATEMENT**
529
530 The raw pool-seq data were deposited in NCBI SRA (accession number: PRJNA602898;
531 http://www.ncbi.nlm.nih.gov/sra). The assembled transcriptomes and their annotation were
532 deposited in Dryad https://doi.org/10.5061/dryad.prr4xgxtk .The scripts used for analyses
533 were deposited in https://github.com/Mengmeng-Lu/Genetic-architecture-underlying-response-
534 to-the-fungal-pathogen-Dothistroma-septosporum-in-Pinus-con
535
536

REFERENCES

Bechsgaard, J., Jorgensen, T. H., & Schierup, M. H. (2017). Evidence for adaptive introgression of disease resistance genes among closely related *Arabidopsis* species. *G3 (Bethesda), 7*(8), 2677-2683. https://doi.org/10.1534/g3.117.043984

Burns, I., James, P. M. A., Coltman, D. W., & Cullingham, C. I. (2019). Spatial and genetic structure of the lodgepole × jack pine hybrid zone. *Canadian Journal of Forest Research, 49*(7), 844-853. https://doi.org/10.1139/cjfr-2018-0428

Capador-Barreto, H. D., Bernhardsson, C., Milesi, P., Vos, I., Lundén, K., Wu, H. X., . . . Elfstrand, M. (2021). Killing two enemies with one stone? Genomics of resistance to two sympatric pathogens in Norway spruce. *Molecular Ecology, 30*(18), 4433-4447. https://doi.org/10.1111/mec.16058

Capron, A., Feau, N., Heinzelmann, R., Barnes, I., Benowicz, A., Bradshaw, R. E., . . . Hamelin, R. C. (2021). Signatures of post-glacial genetic isolation and human-driven migration in the *Dothistroma* needle blight pathogen in Western Canada. *Phytopathology®, 111*(1), 116-127. https://doi.org/10.1094/phyto-08-20-0350-fi

Cullingham, C. I., Cooke, J. E., Dang, S., Davis, C. S., Cooke, B. J., & Coltman, D. W. (2011). Mountain pine beetle host-range expansion threatens the boreal forest. *Molecular Ecology, 20*(10), 2157-2171. https://doi.org/10.1111/j.1365-294X.2011.05086.x

Cullingham, C. I., James, P. M., Cooke, J. E., & Coltman, D. W. (2012). Characterizing the physical and genetic structure of the lodgepole pine× jack pine hybrid zone: mosaic structure and differential introgression. *Evolutionary Applications, 5*(8), 879-891. https://10.1111/j.1752-4571.2012.00266.x

Dale, A. L., Lewis, K. J., & Murray, B. W. (2011). Sexual reproduction and gene flow in the pine pathogen *Dothistroma septosporum* in British Columbia. *Phytopathology, 101*(1), 68-76. https://doi.org/10.1094/phyto-04-10-0121

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., . . . 1000 Genomes Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics, 27*(15), 2156-2158. https://doi.org/10.1093/bioinformatics/btr330

De La Torre, A. R., Puiu, D., Crepeau, M. W., Stevens, K., Salzberg, S. L., Langley, C. H., & Neale, D. B. (2019). Genomic architecture of complex traits in loblolly pine. *New Phytologist, 221*(4), 1789-1801. https://doi.org/110.1111/nph.15535

Dodds, P. N. & Rathjen, J. P. (2010). Plant immunity: towards an integrated view of plant–pathogen interactions. *Nature review genetics, 11*(8), 539-548. https://doi.org/10.1038/nrg2812

Endler, L., Betancourt, A. J., Nolte, V., & Schlötterer, C. (2016). Reconciling differences in pool-GWAS between populations: A case study of female abdominal pigmentation in *Drosophila melanogaster*. *Genetics, 202*(2), 843-855. https://doi.org/10.1534/genetics.115.183376

Feau, N., Ramsfield, T. D., Myrholm, C. L., Tomm, B., Cerezke, H. F., Benowicz, A., . . . Hamelin, R. C. (2021). DNA-barcoding identification of *Dothistroma septosporum* on *Pinus contorta* var. *latifolia*, *P. banksiana* and their hybrid in northern Alberta, Canada. *Canadian Journal of Plant Pathology, 43*(3), 472-479. https://doi.org/10.1080/07060661.2020.1829065

Futschik, A. & Schlötterer, C. (2010). The next generation of molecular markers from massively
parallel sequencing of pooled DNA Samples. *Genetics, 186*(1), 207-218.
https://doi.org/10.1534/genetics.110.114397

Gibson, I. A. S. (1972). *Dothistroma* blight of *Pinus radiata*. *Annual Review of Phytopathology,*
10(1), 51-72. https://10.1146/annurev.py.10.090172.000411

Hall, D. E., Yuen, M. M. S., Jancsik, S., Quesada, A. L., Dullat, H. K., Li, M., . . . Bohlmann, J.
(2013). Transcriptome resources and functional characterization of monoterpene
synthases for two host species of the mountain pine beetle, lodgepole pine (*Pinus*
contorta) and jack pine (*Pinus banksiana*). *BMC plant biology, 13*(1), 80.
https://doi.org/10.1186/1471-2229-13-80

Hamann, A., Smets, P., Yanchuk, A. D., & Aitken, S. N. (2005). An ecogeographic framework for
in situ conservation of forest trees in British Columbia. *Canadian Journal of Forest*
Research, 35(11), 2553-2561. https://doi.org/10.1139/x05-181

Hao, Z.-Z., Liu, Y.-Y., Nazaire, M., Wei, X.-X., & Wang, X.-Q. (2015). Molecular phylogenetics and
evolutionary history of sect. *Quinquefoliae* (*Pinus*): Implications for Northern
Hemisphere biogeography. *Molecular Phylogenetics and Evolution, 87*, 65-79.
https://doi.org/10.1016/j.ympev.2015.03.013

Harrison, R. G. & Larson, E. L. (2014). Hybridization, introgression, and the nature of species
boundaries. *Journal of Heredity, 105*(S1), 795-809.
https://doi.org/10.1093/jhered/esu033

Hill, W. G. & Weir, B. S. (1988). Variances and covariances of squared linkage disequilibria in
finite populations. *Theoretical population biology, 33*(1), 54-78.
https://doi.org/10.1016/0040-5809(88)90004-4

Hivert, V., Leblois, R., Petit, E. J., Gautier, M., & Vitalis, R. (2018). Measuring genetic
differentiation from pool-seq data. *Genetics, 210*(1), 315-330.
https://doi.org/10.1534/genetics.118.300900

Hulse, S. V., Antonovics, J., Hood, M. E., & Bruns, E. L. (2023). Host–pathogen coevolution
promotes the evolution of general, broad-spectrum resistance and reduces foreign
pathogen spillover risk. *Evolution Letters, 7*(6), 467-477.
https://doi.org/10.1093/evlett/qrad051

Isabel, N., Holliday, J. A., & Aitken, S. N. (2019). Forest genomics: Advancing climate adaptation,
forest health, productivity, and conservation. *Evolutionary Applications, 13*(1), 3-10.
https://doi.org/10.1111/eva.12902

Jia, H., Zhang, Y., Orbović, V., Xu, J., White, F. F., Jones, J. B., & Wang, N. (2017). Genome editing
of the disease susceptibility gene CsLOB1 in citrus confers resistance to citrus canker.
Plant biotechnology journal, 15(7), 817-823. https://doi.org/10.1111/pbi.12677

Jin, W.-T., Gernandt, D. S., Wehenkel, C., Xia, X.-M., Wei, X.-X., & Wang, X.-Q. (2021).
Phylogenomic and ecological analyses reveal the spatiotemporal evolution of global
pines. *Proceedings of the National Academy of Sciences, 118*(20), e2022302118.
https://doi.org/10.1073/pnas.2022302118

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics, 24(11), 1403-1405. https://doi.org/10.1093/bioinformatics/btn129

Lenth, R. V. (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical*
Software, 69(1), 1 - 33. https://doi.org/10.18637/jss.v069.i01

Lind, B. M. (2021). GitHub.com/brandonlind/cmh_test: preprint release (Version 1.0.0).
Zenodo. http://doi.org/10.5281/zenodo.5083798

Lind, B. M. (2021). GitHub.com/CoAdapTree/varscan_pipeline: Publication release (Version
1.0.0). Zenodo. https://doi.org/10.5281/zenodo.5083302

Lind, B. M., Lu, M., Obreht Vidakovic, D., Singh, P., Booker, T. R., Aitken, S. N., & Yeaman, S.
(2022). Haploid, diploid, and pooled exome capture recapitulate features of biology and
paralogy in two non-model tree species. *Molecular Ecology Resources, 22*(1), 225-238.
https://doi.org/10.1111/1755-0998.13474

Lotan, J. E. & Critchfield, W. B. (1990). Lodgepole pine. In R. M. Burns & B. H. Honkala (Eds.),
Silvics of North America (Vol. Volume 1: Conifers, pp. 302-315). Washington, D.C.: USDA
Forest Service.

Lu, M., Feau, N., Obreht Vidakovic, D., Ukrainetz, N., Wong, B., Aitken, S. N., . . . Yeaman, S.
(2021). Comparative gene expression analysis reveals mechanism of *Pinus contorta*
response to the fungal pathogen *Dothistroma septosporum*. *Molecular plant-microbe*
interactions: MPMI, 34(4), 397-409. https://doi.org/10.1094/mpmi-10-20-0282-r

Lu, M., Krutovsky, K. V., Nelson, C. D., West, J. B., Reilly, N. A., & Loopstra, C. A. (2017).
Association genetics of growth and adaptive traits in loblolly pine (*Pinus taeda* L.) using
whole-exome-discovered polymorphisms. *Tree Genetics & Genomes, 13*(3), 57.
https://doi.org/10.1007/s11295-017-1140-1

MacLachlan, I. R., McDonald, T. K., Lind, B. M., Rieseberg, L. H., Yeaman, S., & Aitken, S. N.
(2021). Genome-wide shifts in climate-related variation underpin responses to selective
breeding in a widespread conifer. *Proceedings of the National Academy of Sciences,*
118(10), e2016900118. https://doi.org/10.1073/pnas.2016900118

Marroni, F., Pinosio, S., Zaina, G., Fogolari, F., Felice, N., Cattonaro, F., & Morgante, M. (2011).
Nucleotide diversity and linkage disequilibrium in *Populus nigra* cinnamyl alcohol
dehydrogenase (CAD4) gene. *Tree Genetics & Genomes, 7*(5), 1011-1023.
https://doi.org/10.1007/s11295-011-0391-5

Min, T., Hwarari, D., Li, D., Movahedi, A., & Yang, L. (2022). CRISPR-based genome editing and
its applications in woody plants. *International journal of molecular sciences, 23*(17).
https://doi.org/10.3390/ijms231710175

Neale, D. B., Wegrzyn, J. L., Stevens, K. A., Zimin, A. V., Puiu, D., Crepeau, M. W., . . . Langley, C.
H. (2014). Decoding the massive genome of loblolly pine using haploid DNA and novel
assembly strategies. *Genome Biology, 15*(3), R59. https://doi.org/10.1186/gb-2014-15-
3-r59

Paradis, E. & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics, 35*(3), 526-528.
https://doi.org/10.1093/bioinformatics/bty633

Peery, R. M., McAllister, C. H., Cullingham, C. I., Mahon, E. L., Arango-Velez, A., & Cooke, J. E. K.
(2021). Comparative genomics of the chitinase gene family in lodgepole and jack pines:
contrasting responses to biotic threats and landscape level investigation of genetic
differentiation. *Botany, 99*(6), 355-378. https://doi.org/10.1139/cjb-2020-0125

Quesada, T., Gopal, V., Cumbie, W. P., Eckert, A. J., Wegrzyn, J. L., Neale, D. B., . . . Davis, J. M.
(2010). Association mapping of quantitative disease resistance in a natural population of

loblolly pine (*Pinus taeda* L.). *Genetics, 186*(2), 677-686.
https://doi.org/10.1534/genetics.110.117549

R Core Team. (2024). R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing. https://www.R-project.org/.

Ramsfield, T., Myrholm, C., & Tomm, B. (2021). Natural infection of *Pinus contorta* var. *latifolia*,
Pinus banksiana and their hybrid by *Dothistroma septosporum* in Alberta, Canada.
Forest Pathology, 51(5), e12717. https://doi.org/10.1111/efp.12717

Rudolph, T. D. & Laidly, P. R. (1990). *Pinus banksiana* Lamb. Jack Pine. In R. M. Burns & B. H.
Honkala (Eds.), *Silvics of North America* (Vol. Volume 1: Conifers. Agriculture Handbook
654, pp. 280-293). Washington, D.C.: USDA Forest Service.

Rudolph, T. D. & Yeatman, C. W. (1982). Genetics of jack pine. In *Research Paper WO-38* (pp.
35-37). Washington, D.C.: USDA Forest Service.

Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals —
mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics,*
15(11), 749-763. https://doi.org/10.1038/nrg3803

Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., . . . Reyer, C. P.
O. (2017). Forest disturbances under climate change. *Nature Climate Change, 7*(6), 395-
402. https://doi.org/10.1038/nclimate3303

Simler-Williamson, A. B., Rizzo, D. M., & Cobb, R. C. (2019). Interacting effects of global change
on forest pest and pathogen dynamics. *Annual Review of Ecology, Evolution, and*
Systematics, 50(Volume 50, 2019), 381-403. https://doi.org/10.1146/annurev-ecolsys-
110218-024934

Singh, P., StClair, J. B., Lind, B., Cronn, R., Wilhelmi, N. P., Feau, N., . . . Yeaman, S. (2024).
Genetic architecture of disease resistance and tolerance in Douglas-fir trees. *New*
Phytologist. https://doi.org/10.32942/osf.io/s5rd7

Stocks, J. J., Metheringham, C. L., Plumb, W. J., Lee, S. J., Kelly, L. J., Nichols, R. A., & Buggs, R. J.
A. (2019). Genomic basis of European ash tree resistance to ash dieback fungus. *Nature*
Ecology & Evolution, 3(12), 1686-1696. https://doi.org/10.1038/s41559-019-1036-6

Suren, H., Hodgins, K. A., Yeaman, S., Nurkowski, K. A., Smets, P., Rieseberg, L. H., . . . Holliday, J.
A. (2016). Exome capture from the spruce and pine giga-genomes. *Molecular Ecology*
Resources, 16(5), 1136-1146. https://doi.org/10.1111/1755-0998.12570

Taylor, S. A., Larson, E. L., & Harrison, R. G. (2015). Hybrid zones: windows on climate change.
Trends in Ecology & Evolution, 30(7), 398-406.
https://doi.org/10.1016/j.tree.2015.04.010

van Esse, H. P., Reuber, T. L., & van der Does, D. (2020). Genetic modification to improve
disease resistance in crops. *New Phytologist, 225*(1), 70-86.
https://doi.org/10.1111/nph.15967

Wang, L., Chen, S., Peng, A., Xie, Z., He, Y., & Zou, X. (2019). CRISPR/Cas9-mediated editing of
CsWRKY22 reduces susceptibility to *Xanthomonas citri subsp. citri* in Wanjincheng
orange (*Citrus sinensis* (L.) Osbeck). *Plant Biotechnology Reports, 13*(5), 501-510.
https://doi.org/10.1007/s11816-019-00556-x

Wegrzyn, J. L., Liechty, J. D., Stevens, K. A., Wu, L. S., Loopstra, C. A., Vasquez-Gross, H. A., . . .
Neale, D. B. (2014). Unique features of the loblolly pine (*Pinus taeda* L.) megagenome

revealed through sequence annotation. *Genetics, 196*(3), 891-909.
712   https://doi.org/10.1534/genetics.113.159996

713 Wei, T. & Simko, V. (2017). R package "corrplot": Visulaization of a correlation matrix.
714   *https://github.com/taiyun/corrplot*.

715 Weiss, M., Sniezko, R. A., Puiu, D., Crepeau, M. W., Stevens, K., Salzberg, S. L., . . . De La Torre,
716   A. R. (2020). Genomic basis of white pine blister rust quantitative disease resistance and
717   its relationship with qualitative resistance. *The Plant Journal, 104*(2), 365-376.
718   https://doi.org/10.1111/tpj.14928

719 Welsh, C., Lewis, K., & Woods, A. (2009). The outbreak history of *Dothistroma* needle blight: an
720   emerging forest disease in northwestern British Columbia, Canada. *Canadian Journal of*
721   *Forest Research, 39*(12), 2505-2519. https://doi.org/10.1139/X09-159

722 Wheeler, N. C. & Guries, R. P. (1987). A quantitative measure of introgression between
723   lodgepole and jack pines. *Canadian Journal of Botany, 65*, 1876-1885.
724   https://doi.org/10.1139/b87-257

725 White, T. L. (2004). TREE BREEDING, PRINCIPLES | Breeding Theory and Genetic Testing. In J.
726   Burley (Ed.), *Encyclopedia of Forest Sciences* (pp. 1551-1561). University of Oxford,
727   Oxford, UK: Elsevier.

728 Wood, L. J., Hartley, I. D., & Watson, P. (2009). Determining hybridization in jack pine and
729   lodgepole pine from British Columbia. *Wood and Fiber Science, 41*, 386-395.

730 Woods, A. J. (2003). Species diversity and forest health in northwest British Columbia. *The*
731   *Forestry Chronicle, 79*(5), 892-897. https://doi.org/10.5558/tfc79892-5

732 Woods, A. J., Coates, K. D., & Hamann, A. (2005). Is an unprecedented *Dothistroma* needle
733   blight epidemic related to climate change? *BioScience, 55*(9), 761-769.
734   https://doi.org/10.1641/0006-3568(2005)055[0761:IAUDNB]2.0.CO;2

735 Wu, M. C., Kraft, P., Epstein, M. P., Taylor, D. M., Chanock, S. J., Hunter, D. J., & Lin, X. (2010).
736   Powerful SNP-set analysis for case-control genome-wide association studies. *American*
737   *Journal of Human Genetics, 86*(6), 929-942. https://doi.org/10.1016/j.ajhg.2010.05.002

738 Wu, T. D. & Watanabe, C. K. (2005). GMAP: a genomic mapping and alignment program for
739   mRNA and EST sequences. *Bioinformatics, 21*(9), 1859-1875.
740   https://doi.org/10.1093/bioinformatics/bti310

741 Yeaman, S., Hodgins, K. A., Lotterhos, K. E., Suren, H., Nadeau, S., Degner, J. C., . . . Aitken, S. N.
742   (2016). Convergent local adaptation to climate in distantly related conifers. *Science,*
743   *353*(6306), 1431-1433. https://doi.org/10.1126/science.aaf7812

744 Yeatman, C. W. & Teich, A. H. (1969). Genetics and breeding of jack and lodgepole pines in
745   Canada. *The Forestry Chronicle, 45*(6), 428-433. https://doi.org/10.5558/tfc45428-6

746 Zhou, H., Bai, S., Wang, N., Sun, X., Zhang, Y., Zhu, J., & Dong, C. (2020). CRISPR/Cas9-mediated
747   mutagenesis of MdCNGC2 in apple callus and VIGS-mediated silencing of MdCNGC2 in
748   fruits improve resistance to *Botryosphaeria dothidea*. *Frontiers in Plant Science, 11*.
749   https://doi.org/10.3389/fpls.2020.575477