Bayesian parameter inference for epithelial mechanics

Xin Yan[1,†], Goshi Ogita[2,†,‡], Shuji Ishihara[3,4], and Kaoru Sugimura[1,4,5,‡]

[1]Department of Computational Biology and Medical Sciences, Graduate School of
Frontier Sciences, The University of Tokyo, Chiba 277-8561, Japan
[2]Laboratory for Physical Biology, RIKEN Center for Biosystems Dynamics
Research, Kobe 650-0047, Japan
[3]Department of Integrated Sciences, Graduate School of Arts and Sciences, The
University of Tokyo, Tokyo 153-8902, Japan
[4]Universal Biology Institute, The University of Tokyo, Tokyo 113-0033, Japan
[5]Department of Biological Sciences, Graduate School of Science, The University of
Tokyo, Tokyo 113-0032, Japan
[†]Joint first authors.
[‡]Correspondence: goshi.ogita@riken.jp (GO), sugimura@bs.s.u-tokyo.ac.jp (KS)

May 2, 2024

Abstract

Cell-based mechanical models, such as the Cell Vertex Model (CVM), have
proven useful for studying the mechanical control of epithelial tissue dynam-
ics. We recently developed a statistical method called image-based parameter
inference for formulating CVM model functions and estimating their parame-
ters from image data of epithelial tissues. In this study, we employed Bayesian
statistics to improve the utility and flexibility of image-based parameter in-
ference. Tests on synthetic data confirmed that both our non-hierarchical and
hierarchical Bayesian models provide accurate estimates of model parameters.
By applying this method to *Drosophila* wings, we demonstrated that the reli-
ability of parameter estimation is closely linked to the mechanical anisotropies
present in the tissue. Moreover, we revealed that the cortical elasticity term
is dispensable for explaining force-shape correlations *in vivo*. We anticipate
that the flexibility of the Bayesian statistical framework will facilitate the in-
tegration of various types of information, thereby aiding in the quantitative
dissection of the mechanical control of tissue dynamics.

1 Introduction

Mathematical modeling provides means for testing hypotheses that are experimen-
tally inaccessible, deciphering underlying mechanisms, and guiding new research di-
rections through testable predictions [1–3]. These advantages make it indispensable
for understanding complex biological phenomena, such as tissue development and
repair. Mathematical modeling typically involves three key steps: model construc-
tion, parameter estimation, and model selection [4, 5]. First, model equations are
formulated to represent the biological process of interest, often resulting in multiple
candidate models. Subsequently, parameter estimation is carried out to determine

the parameter values that best recapitulate experimental data. Assessing the uncertainty associated with parameter estimation is important, especially when making decisions based on theoretical predictions [6–8]. Finally, the most appropriate model is selected from the candidate models based on statistical criteria that balance the goodness-of-fit and model complexity. Despite its importance, a statistical framework to evaluate parameter estimation uncertainty and model goodness is yet to be fully established in developmental biology [8, 9].

Various modeling frameworks have been employed to investigate the physical mechanisms by which mechanics control tissue dynamics [10–13]. Among them, the cell vertex model (CVM) is one of the most frequently used models for studying epithelial mechanics [14–16]. The CVM treats epithelial tissue as a tile of polygons, and computes the displacement and reconnection of cell vertices via the minimization of virtual work, by which the balance between junction tensions and cell pressure is considered (Fig. 1A-C). In the CVM, cellular forces, such as junction tension and cell pressure, are dependent on the morphological characteristics of cells (*e.g.,* junction length and cell area), for which specific forms of model functions and parameter values are assumed. These model functions and parameters must be determined based on experimental data to ensure the validity of the modeling; however, most existing approaches do not fully leverage the quantitative information embedded in experimental data.

With some recent exceptions [17–20], many previous studies using the CVM have relied on standard model functions that qualitatively represent epithelial mechanics [15, 16]. Parameter values are determined by iteratively running numerical simulations with different parameter sets and fitting the simulation results to experimental observations [7, 21–25]. Summary statistics, such as the polygonal distribution of cells, are used for this fitting process, making the comparison between simulation and experimental data indirect. Moreover, information criteria are often underutilized for comparing multiple candidate models.

To address these technical challenges concerning the CVM, we recently developed a data-driven, statistical method to deduce model functions and estimate their parameters directly from experimental data [26] (Fig. 1D; explained in detail in Methods). First, we formulate model functions (tension and pressure functions) based on correlations between force and cell shape obtained from image data of epithelial tissue. Substitution of the model functions into force-balance equations at the cell vertex leads to an equation with respect to the parameters of the model. Solving this equation using a least-squares method allows for the estimation of parameter values, without performing numerical simulations. Finally, Akaike Information Criterion (AIC) [27] is employed for selecting the most suitable model among candidates. Tests using synthetic and *in vivo* data have validated the accuracy and robustness of parameter estimation and model selection [26]. Overall, our method offers a simple, efficient, and data-driven approach for cell-based modeling in epithelial mechanics.

Nevertheless, the current method has some limitations. It provides only point estimates of parameters, leaving the uncertainty in estimation unaddressed. In addition, the point estimation makes it difficult to identify representative models and parameters for groups categorized by tissue type, developmental stage, genotype, and so on. Furthermore, certain model functions result in physically unrealistic estimates, likely due to multicollinearity. Bayesian statistics have been known to offer solutions for overcoming related limitations [28, 29]. The uncertainty in parameter estimation can be directly quantified from the posterior distribution. Hierarchi-

cal Bayesian modeling enables the integration of multi-layered information, such as groups and their individual samples. Using informative priors to restrict the parameter space may extend the range of model functions that can be analyzed.

In this study, we aimed to improve the image-based parameter inference using Bayesian statistics. We developed both non-hierarchical and hierarchical Bayesian models and evaluated their accuracy in parameter estimation. With these methods, we explored how uncertainty in parameter estimation correlates with developmental changes in tissue mechanics, and examined whether mathematical representation of the cortical elasticity is appropriate for modeling epithelial tissues. We foresee that the flexibility in Bayesian statistical framework will aid in integrating various types of information such as time and space, thereby facilitating the quantitative dissection of mechanical control of tissue dynamics.

2 Methods

In this study, we reformulated image-based parameter inference [26] using Bayesian statistics. The original method and the new Bayesian method follow the same procedures until constructing candidate models (section 2.1) and writing force-balance equations at cell vertices (section 2.2). After that, the former uses the least-squares method for parameter estimation (section 2.3), while the latter formulates the problem using Bayes' theorem (section 2.4).

2.1 Mechanical model for epithelial tissue

In the monolayer epithelium, actomyosin cables running along the plane of adherens junction (AJ) generate junction tension, while cell pressure counteracts junction tension to maintain the cell area (Fig. 1A–C). In the CVM, the partial differentiation of virtual work $U_0(\{\vec{x}\})$ by the length of the junction between the i-th and j-the cell, l_{ij}, leads to the junction tension T_{ij} as $T_{ij} \equiv \partial U_0/\partial l_{ij}$, while the partial differentiation by the area of the i-th cell, A_i, leads to the cell pressure P_i as $P_i \equiv -\partial U_0/\partial A_i$. The junction tension and the cell pressure are determined by model equations that include parameters representing cell mechanical properties, such as the elastic coefficient of cell.

We have established data-driven formulation of tension and pressure equations in the CVM (Fig. 1D) [26]. Briefly, we first employed Bayesian force inference [30] to quantify the relative values of junction tension and pressure difference among cells from image data of epithelial tissues. Subsequently, we plotted the inferred forces against the morphological characteristics of cells, such as the junction length and the cell area. Through this analysis of force-shape correlation, we formulated candidate model functions for tension and pressure (for detailed information, see [26]).

2.1.1 Model equations for junction tension

The function for junction tension is of the following form:

$$T(l_{ij}, \theta_{ij}) = \Lambda_0(\theta_{ij}) - \Lambda_1(\theta_{ij})l_{ij} \tag{1}$$

where θ_{ij} and l_{ij} are the direction and length of junction between i-th and j-th cells. The anisotropic line tension $\Lambda_0(\theta)$ (Fig. 2A) and the anisotropic spring constant

$\Lambda_1(\theta)$ (Fig. 2A) are defined as bellow:

$$\Lambda_0(\theta_{ij}) = \lambda_0\{1 + \mu_0 \cos 2(\theta_{ij} - \varphi_0)\} \tag{2}$$

$$\Lambda_1(\theta_{ij}) = \lambda_1\{1 + \mu_1 \cos 2(\theta_{ij} - \varphi_1)\}. \tag{3}$$

λ_0 and λ_1 represent the magnitude of $\Lambda_0(\theta_{ij})$ and $\Lambda_1(\theta_{ij})$, respectively. μ_0 and μ_1 denote the magnitudes of their orientation-dependent (anisotropic) parts, which peak at φ_0 and φ_1. λ_0, λ_1, μ_0 and μ_1 are the linear parameters and φ_0 and φ_1 are the axial parameters with interval $[0, \pi]$.

The anisotropic spring constant, $\Lambda_1(\theta_{ij})$, represents feedback between myosin-II (myo-II)-generated junction tension and junction shrinkage (Fig. 2B). The sign of $\Lambda_1(\theta_{ij})$ indicates whether the feedback is positive (*i.e.,* accelerating junction shortening/lengthening) or negative (*i.e.,* stabilizing the junction length) . This feedback between tension/myo-II and junction shrinkage is supported by experimental observations that myo-II is getting concentrated on shrinking junctions during cell rearrangement [17, 31, 32] and has been studied in mathematical models for *Drosophila* embryogenesis [17, 33, 34].

Our previous study identified distinctive biological functions of $\Lambda_0(\theta_{ij})$ and $\Lambda_1(\theta_{ij})$ during the development of the *Drosophila* pupal wing (Fig. 2C–E) [26]. Specifically, $\Lambda_0(\theta_{ij})$ contributes to resistance against the extrinsic pulling force from the hinge. This is supported by observations that μ_0 increases following the onset of hinge constriction at 15 hours after puparium formation (h APF), which exerts a pulling force along the proximal-distal (PD) axis, and that φ_0 aligns with the PD axis (Fig. 2C, D) [35–39]. On the other hand, φ_1 is aligned with the AP axis and $\Lambda_1(\theta_{ij})$ promotes the shrinkage of AP junctions, running perpendicular to the global tension anisotropy (Fig. 2E) [36, 39, 40].

In section 3.4, we consider the following alternative model incorporating cortical elasticity:

$$T(l_{ij}, \theta_{ij}, L_i, L_j) = \Lambda_0(\theta_{ij}) - \Lambda_1(\theta_{ij})l_{ij} + \Gamma(L_i + L_j), \tag{4}$$

where Γ and L_i represent the elastic coefficient of cell cortex and the perimeter of i-th cell, respectively.

In addition to the models described above, simpler models were considered as summarized in Fig. S1. Model A is hereafter referred to as the anisotropic spring model. Model E' corresponds to the standard equation of junction tension commonly used in the CVM [15, 16, 21]:

$$T(L_i, L_j) = \lambda_0 + \Gamma(L_i + L_j). \tag{5}$$

2.1.2 Model equations for cell pressure

The pressure function used in this study is as follows:

$$P(A_i) = -kA_i + P_0, \tag{6}$$

where A_i is the area of i-th cell. k and P_0 are the elastic coefficient of a cell and the hydro-static pressure, respectively [15, 16, 21]. We have tested alternative pressure functions and demonstrated that the choice of pressure function had a negligible impact on parameter estimation [26].

2.2 Force balance equation for epithelial tissue

The parameters involved in tension/pressure functions are estimated by considering the balance of forces acting on the cell vertices [26]. Suppose that the pressure of the i-th cell is P_i and the tension of the cell junction between the i-th and j-th cell is T_{ij} (Fig. 1B). Under the approximation that epithelial sheet is regarded as a tessellation of polygons, the following equations describe the net forces acting at the 0-th vertex, $\vec{f_0} = (f_0^x, f_0^y)$ in Fig. 1C [30, 41–43].

$$f_0^x = a_1^x T_{12} + a_2^x T_{23} + a_3^x T_{31} + b_1^x P_1 + b_2^x P_2 + b_3^x P_3 \tag{7}$$

$$f_0^y = a_1^y T_{12} + a_2^y T_{23} + a_3^y T_{31} + b_1^y P_1 + b_2^y P_2 + b_3^y P_3. \tag{8}$$

The coefficients are calculated only from the vertex coordinates, such as $a_1^x = (x_1 - x_0)/l_{12} = \cos(\theta_{12})$, $a_1^y = (y_1 - y_0)/l_{12} = \sin(\theta_{12})$, $b_1^x = (y_2 - y_1)/2$ and $b_1^y = -(x_2 - x_1)/2$, and are measurable from image data.

By repeating this procedure for all vertices in tissue, the following equation is obtained:

$$\vec{F} = (f_0^x, f_0^y, f_1^x, f_1^y, \cdots)^T = C\vec{X}, \tag{9}$$

where C is an $n \times m$ matrix composed of coefficients a_i^x, a_i^y, b_j^x and b_j^y and $\vec{X} = (\vec{T}, \vec{P})^T$ is an m-dimensional vector composed of T_{ij} and P_i [30]. n is the number of force balance equations and equals twice the number of vertices. m is the sum of the number of junctions and the number of cells, which correspond to the dimension of the vector $\vec{X}$. Given that epithelial morphogenesis is assumed to be a quasi-static process, wherein forces acting on the vertices are almost balanced, the following force-balance equation holds:

$$\vec{F} = C\vec{X} \approx \vec{0} . \tag{10}$$

Substitution of the model equations (Eqs. 1 and 6) into Eq. 10 gives an equation of the parameter vector $\vec{\beta} = (\lambda_0, \mu_0, \varphi_0, \lambda_1, \mu_1, \varphi_1, \Gamma, k)^T$:

$$\vec{F} = C\vec{X}(\vec{\beta}; D) \approx \vec{0} , \tag{11}$$

where D is the list of cell shape characteristics, namely the junction length l_{ij}, the junction angle θ_{ij}, the cell perimeter length L_i, and the cell area A_i quantified from the image data. Since the solution to this equation has arbitrariness in the scaling, we chose λ_0 as a scaling factor and fixed it to 1.0. This enables us to decompose $\vec{X}(\vec{\beta})$ into the constant term derived from λ_0 and the data-dependent terms involving the other parameters:

$$\vec{X}(\vec{\beta}) = \vec{X}_{\lambda_0} + \vec{X}'(\vec{\beta}'; D). \tag{12}$$

$\vec{X}_{\lambda_0}$ is an m-dimensional vector, whose component $X_{\lambda_0,i}$ is 1 ($i = 1, \ldots, m_j$) or 0 ($i = m_j + 1, \ldots, m$), where m_j is the number of junctions. $\vec{\beta}'$ is the vector composed of parameters other than λ_0. $\vec{X}'(\vec{\beta}')$ represents the cellular force with the constant term subtracted. Substitution of Eq. 12 into Eq. 11 leads to

$$\vec{F}(\vec{\beta}'; D) = C\vec{X}_{\lambda_0} + C\vec{X}'(\vec{\beta}'; D) \approx \vec{0} , \tag{13}$$

which is the equation of the parameter to be solved.

2.3 Parameter estimation by the least-squares method

In this section, we outline the procedure used in [26] to estimate parameters by transforming Eq. 13 into the multiple-linear regression problem and solving it by the least-squares method.

The estimated values of the mechanical parameters $\hat{\vec{\beta}}'$ are obtained by minimizing the L^2 norm of $\vec{F}(\vec{\beta}'; D)$ in Eq. 13:

$$\hat{\vec{\beta}}' = \underset{\vec{\beta}'}{\operatorname{argmin}} ||\vec{F}(\vec{\beta}'; D)||^2. \tag{14}$$

To transform this minimization problem in Eq. 14 into the multivariate regression problem, we rewrite the anisotropic line tension $\Lambda_0(\theta_{ij})$ and the anisotropic spring constant of junction $\Lambda_1(\theta_{ij})$ as:

$$\Lambda_0(\theta_{ij}) = \lambda_0 + \lambda_0' \sin(2\theta_{ij}) + \lambda_0'' \cos(2\theta_{ij}) \tag{15}$$

$$\Lambda_1(\theta_{ij}) = \lambda_1 + \lambda_1' \sin(2\theta_{ij}) + \lambda_1'' \cos(2\theta_{ij}) . \tag{16}$$

The transformed parameters $\lambda_0', \lambda_0'', \lambda_1', \lambda_1''$ are defined as $\lambda_0' = \lambda_0 \mu_0 \sin(2\varphi_0), \lambda_0'' = \lambda_0 \mu_0 \cos(2\varphi_0), \lambda_1' = \lambda_1 \mu_1 \sin(2\varphi_1), \lambda_1'' = \lambda_1 \mu_1 \cos(2\varphi_1)$. Next, we define the matrices S_T and S_P as follows:

$$S_T = \begin{pmatrix} \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & \sin 2\theta_{ij} & \cos 2\theta_{ij} & -l_{ij} & -l_{ij} \sin 2\theta_{ij} & -l_{ij} \cos 2\theta_{ij} & L_i + L_j \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} , \tag{17}$$

$$S_P = \begin{pmatrix} \vdots \\ -A_i \\ \vdots \end{pmatrix}. \tag{18}$$

Note that S_T and S_P are calculated based on D, the list of cell shape characteristics quantified from image data. Consequently, the vectors $\vec{T}$ and $\vec{P}$ representing junction tension and cellular pressure are given by

$$\vec{T} = S_T \vec{\beta}_T , \quad \vec{P} = S_P \vec{\beta}_P, \tag{19}$$

respectively, where $\vec{\beta}_T$ and $\vec{\beta}_P$ are composed of the transformed tension and pressure parameters.

$$\vec{\beta}_T = \left(\lambda_0, \lambda_0', \lambda_0'', \lambda_1, \lambda_1', \lambda_1'', \Gamma\right)^T , \quad \vec{\beta}_P = \left(k\right)^T. \tag{20}$$

By using Eq. 19, the cellular force vector X is denoted as:

$$\vec{X} = S\vec{\beta}, \tag{21}$$

with the matrix S and the parameter vector $\vec{\beta}$ defined as

$$S = \begin{pmatrix} S_T & 0 \\ 0 & S_P \end{pmatrix} , \quad \vec{\beta} = \begin{pmatrix} \vec{\beta}_T \\ \vec{\beta}_P \end{pmatrix}. \tag{22}$$

Substituting Eq. 21 into Eq. 12 yields:

$$\begin{aligned} \vec{F} &= C\vec{X}_{\lambda_0} + C\vec{X}'(\vec{\beta}'; D) \\ &= C\vec{X}_{\lambda_0} + CS'\vec{\beta}', \end{aligned} \tag{23}$$

where S' is obtained by eliminating the first column from S. The minimization problem of L^2 norm of $\vec{F}$ given in Eq. 14 is equivalent to the linear multiple regression problem with $C\vec{X}_{\lambda_0}$ as the observation vector, $-CS'$ as the design matrix, and $\vec{\beta}'$ as the parameter vector. This means that the estimation of the parameter vector $\hat{\vec{\beta}}'$ can be obtained analytically as follows:

$$\hat{\vec{\beta}}' = -((CS')^T(CS'))^{-1}(CS')^T C\vec{X}_{\lambda_0}. \tag{24}$$

We solved the lest-squares problem using the function **OLS.fit()** in the Python package statsmodels.

When analyzing individual samples separately, we derived C and S from the image data of each sample and then performed parameter estimation. For pooled estimation, where individual samples are aggregated into a single dataset, we combined C (or S) for the image data of all samples in the group before proceeding with parameter estimation.

2.4 Parameter estimation by Bayesian statistics

In this study, we constructed both a non-hierarchical Bayesian model and a hierarchical Bayesian model (Fig. 1E–H). The former is designed for estimating parameters of individual samples, while the latter is designed for estimating parameters at both the group-level and individual-level (Fig. 1G, H) [28].

2.4.1 Non-hierarchical Bayesian model

The posterior distribution of the non-hierarchical Bayesian model, $p(\vec{\beta}', \sigma | C, D)$, is defined as follows:

$$p(\vec{\beta}', \sigma | C, D) \propto p(C\vec{X}_{\lambda_0} | C\vec{X}'(\vec{\beta}'; D), \sigma) p(\vec{\beta}') p(\sigma). \tag{25}$$

$p(C\vec{X}_{\lambda_0} | C\vec{X}'(\vec{\beta}'; D), \sigma)$ is the likelihood function with a scale parameter σ and represents the probability of observing the data at given parameter values $\vec{\beta}'$. $p(\vec{\beta}')$ and $p(\sigma)$ are the prior distributions for $\vec{\beta}'$ and σ, respectively.

The likelihood function is derived from the force-balance equation at cell vertices (section 2.2). We assume that the net forces acting on cell vertices are normally distributed with zero mean and $\sigma^2 I$ covariance [30]. This assumption is equivalent to $C\vec{X}_{\lambda_0} \sim \mathcal{N}(C\vec{X}'(\vec{\beta}'; D), \sigma^2 I)$ based on Eq. 12. The likelihood function thus reads:

$$p(C\vec{X}_{\lambda_0} | C\vec{X}'(\vec{\beta}'; D), \sigma) = \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{\|C\vec{X}_{\lambda_0} - C\vec{X}'(\vec{\beta}'; D)\|^2}{2\sigma^2}\right). \tag{26}$$

We employed the following non-informative priors. For the prior distribution of λ_1, μ_0, μ_1 and k, we applied a uniform distribution on the interval $[0, 1]$. For the prior distribution of the elastic coefficient of the cell cortex, Γ, we applied a half-Cauchy distribution with a scale of 5. For the prior distribution of φ_0 and φ_1, we applied an axial uniform distribution, which is a uniform distribution on the semi-circle $[0, \pi)$ [44, 45]. We set a half-Cauchy prior with the scale 5 on σ. We tested other prior distributions, such as half normal distribution, instead of half Cauchy distribution, and confirmed that the posterior distributions were not significantly affected by the choice of the prior distributions: the difference between estimated parameter values was at most 1.26×10^{-5} for three test data of *Drosophila* epithelial tissues.

2.4.2 Hierarchical Bayesian model

The posterior distribution of a hierarchical Bayesian model, $p(\vec{\beta}'_0, \sigma | \{C_i\}, \{D_i\})$, is defined as follows:

$$p(\vec{\beta}'_0, \sigma | \{C_i\}, \{D_i\}) \propto \prod_{i:\text{individual}} p(C_i \vec{X}_{\lambda_0,i} | C\vec{X}'(\vec{\beta}'_i; D_i), \sigma) p(\vec{\beta}'_i | \vec{\beta}'_0, \vec{\sigma}_0) \times p(\sigma) p(\vec{\beta}'_0) p(\vec{\sigma}_0), \tag{27}$$

where subscript i indicates the index of samples in a group, and $\{C_i\}$ and $\{D_i\}$ are the list of coefficient matrices and cell shape quantities in a group-data. The likelihood function $p(C_i \vec{X}_{\lambda_0,i} | C\vec{X}'(\vec{\beta}'_i; D_i), \sigma)$ is the same as that in the non-hierarchical Bayesian model. $p(\vec{\beta}' | \vec{\beta}'_0, \vec{\sigma}_0)$ is the group-level distribution of mechanical parameters, which the mechanical parameters of each sample follows. $\vec{\beta}'_0$ is representative mechanical parameters of a group. $p(\vec{\beta}'_0)$ is the prior distribution of the group-level meta parameters $\vec{\beta}'_0$.

We employed unimodal distributions for the group level distribution of the mechanical parameters $p(\vec{\beta}'_0)$. For the linear parameters $(\lambda_1, \mu_0, \mu_1, k)$, we tested the beta distributions. For instance, the group-level distribution of λ_1 is given by:

$$p(\lambda_1 | \alpha_{\lambda_1}, \beta_{\lambda_1}) = \frac{\lambda_1^{\alpha_{\lambda_1}-1}(1-\lambda_1)^{\beta_{\lambda_1}-1}}{B(\alpha_{\lambda_1}, \beta_{\lambda_1})}, \tag{28}$$

where α_{λ_1} and β_{λ_1} are parameters of the beta distribution and $B(\alpha_{\lambda_1}, \beta_{\lambda_1})$ is the beta function. The group level distribution of other linear parameters were set in the same way. For the axial parameters (φ_0, φ_1), we utilized the axial von-Mises distribution [45]. For instance, the group level distribution of φ_0 reads

$$p(\varphi_0 | \varphi_0^g, \kappa_0) = \frac{\exp(\kappa_0 \cos 2(\varphi_0 - \varphi_0^g))}{\pi I_0(\kappa)}, \tag{29}$$

where $I_0(\kappa)$ is the modified Bessel function of order zero. φ_0^g and κ_0 represent the location and concentration parameters of a group, respectively. We applied half-Cauchy priors with the scale 5 on $1/\kappa_0$. The group level distribution of φ_1 was set following the same procedure.

2.5 MCMC Sampling and MAP estimation

We conducted Markov chain Monte Carlo (MCMC) sampling using PyMC3, a python package for Bayesian inference (version 3.11.4; [46]). We applied the ADVI and NUTS algorithms to initialize and sample the posterior distribution [47, 48]. In the NUTS sampler, we set $target\_accept = 0.99$. For the non-hierarchical Bayesian method, we ran two MCMC chains, each with 15,000 steps, of which the first 5,000 steps were discarded as burn-in. We assessed the quality of sampling using the Gelman-Rubin statistic $\hat{R}$ [28, 49, 50] and divergences. If $\hat{R}$ exceeded 1.05 and/or divergences occurred frequently, the entire run was discarded and repeated. The MAP estimate was calculated using the find_MAP function in PyMC3, in which the numerical optimization is performed using the CG algorithm. For the hierarchical Bayesian method, we ran six MCMC chains. If most chains converged but one or two chains clearly deviated, we discarded the deviating chains and used the remaining chains for the sampling results. We calculated the MAP estimate from the mode of the posterior distribution sampled by MCMC. The approximate time required for MCMC sampling was 1–2 hours per sample, regardless of using non-hierarchical or hierarchical Bayesian models.

2.6 Model selection

For model selection in the Bayesian methods, we employed Widely applicable information criterion (WAIC) and leave-one-out cross-validation (LOOCV) [50–52]. WAIC and LOOCV are respectively calculated by the functions **waic** and **loo** in a python package Arviz [53]. We found that WAIC and LOOCV selected the same model both in the non-hierarchical and hierarchical Bayesian methods in our data sets of *Drosophila* epithelial tissues.

AIC was used for model selection in the original least-squares method [26]. AIC was calculated using **OLS.fit()** in the Python package statsmodels.

2.7 Uncertainty quantification

We quantified the uncertainty associated with parameter estimation using the 95% highest posterior density (HDI) interval, within which a parameter value falls with a 95% probability in the posterior distribution (Fig. 1F) [28]. The means of the 95% HDIs for each selected model at every developmental stage were calculated as follows: For linear parameters, we normalized the 95% HDIs by the MAP value in each sample and then computed the average of these normalized values. For axial parameters, we first calculated the double lengths of the 95% HDIs using circular statistics [44, 45], and then divided these values by 2.

2.8 Generation of synthetic data set

In this study, we used synthetic data to assess the accuracy of parameter estimation. The synthetic data on the cell configuration was generated through numerical simulation of the CVM as previously described [26]. The positions and connectivity of vertices $\{\vec{x}\}$ were changed according to the minimization of virtual work $U_0(\{\vec{x}\})$. The virtual work of the anisotropic spring model (Fig. 2; model A in Fig. S1) is given as

$$U_0(\{\vec{x}\}) = \sum_{[ij]:junction} \left\{ \Lambda_0(\theta_{ij})l_{ij} - \frac{1}{2}\Lambda_1(\theta_{ij})l_{ij}^2 \right\} + \sum_{i:cell} \frac{1}{2}k(A_i - A_0)^2, \quad (30)$$

where A_0 is the target area of a cell and set at $A_0 = 1.0$. We solved the following equation in simulations.

$$\frac{d\vec{x}_k}{dt} = -\frac{\partial U_0(\{\vec{x}\})}{\partial \vec{x}_k} = -T_{ij}\frac{\partial l_{ij}}{\partial \vec{x}_k} + P_i\frac{\partial A_i}{\partial \vec{x}_k}, \quad (31)$$

where derivatives of anisotropy coefficients such as $\partial \Lambda_0(\theta_{ij})/\partial \vec{x}_k$ are ignored [54]. The numerical simulations were implemented by C++. We set an initial cell configuration as a 20×20 cell tile, which was generated from randomly distributed centroids with a mean cell area of A_0. We adopted a free boundary condition. The discretized time-step was set to $\Delta t = 0.1$, and the simulation was run until a configuration of cells became stable (t = 5,000). To implement a T1 transition, junctions shorter than $0.05 \times \sqrt{A_0}$ were allowed to perform a T1 transition if this transition reduces the virtual work. We generated 15 synthetic data. λ_0 was fixed at 1.0, in line with the parameter estimation formulation, and the other parameters were randomly generated from a logit-normal distribution. For instance, we set the distribution function of λ_1 as follows:

$$p(\lambda_1|\lambda_1^g, \sigma_{\lambda_1}) = \frac{1}{\sqrt{2\pi}\sigma_{\lambda_1}} \frac{1}{\lambda_1(1 - \lambda_1)} \exp\left(-\frac{(logit(\lambda_1) - \lambda_1^g)^2}{2\sigma_{\lambda_1}^2} \right), \quad (32)$$

where λ_1^g is the location parameter and σ_{λ_1} is the scale parameter for the group. The distribution function of the other parameters were set in the same way. The location parameters of the logit-normal distribution were set at $(\mu_0^g, \lambda_1^g, \mu_1^g, k'^g) = (0.15, 0.15, 0.5, 0.5)$, where $k' = 0.1k$. The scale parameter was set at 0.3 for all parameters.

2.9 *In vivo* image data

In vivo image data used in this study have been described in [26]. *Drosophila* lines used were *sqhp-sqh-GFP* [55], *UAS-Dα-cat-TagRFP* [30], *apterous (ap)-Gal4*, and *DE-cad-GFP* [56]. The fly genotypes were *sqhp-sqh-GFP*, *ap-Gal4/sqhp-sqh-GFP*, *UAS-Dα-cat-TagRFP* for experiments in pupal wing and notum, and *DE-cad-GFP* for embryo. The total number of samples is 165. For the pupal wing, the number of samples at each developmental stage is as follows: 20 at 13-14 h APF, 23 at 16.5-18.5 h APF, 16 at 21-23 h APF, 21 at 25.5-27.5 h APF, and 25 at 30-32 h APF. For the pupal notum, there are 10 samples at each of five developmental stages (13-14 h, 16.5-18.5 h, 21-23 h, 25.5-27.5 h, and 30-32 h APF). For the embryo, there are 5 samples at each of two developmental stages (before and during the germband elongation).

2.10 Image processing and analysis

Images of the pupal wing and notum were segmented by custom-made macro and plugins in ImageJ/Fiji [30]. Images of the embryonic germband were segmented by EPySEG [57]. Vertex position and connectivity were extracted from skeletonized images by using custom-made code in OpenCV [30].

We preprocessed input image data as follows. First, we excluded junctions shorter than 3 pixels (*in vivo* data) or $0.05 \times \sqrt{A_0}$ (synthetic data; A_0 is a target cell area) and cells larger than twice the median area of cells from the analysis. We then normalized the spatial scale using the median cell area. Finally, we rotated the vertex coordinates so that the body axes of each image taken from the same tissue were aligned. We aligned the pupal wing images along the proximal-distal axis, while the pupal notum and embryonic germband images were aligned along the anterior-posterior axis. This step is necessary because pooled and hierarchical Bayesian estimation require the alignment of the body axis prior to conducting parameter estimation. Note that in our previous study [26], the axial parameters were corrected post-estimation rather than during the pre-processing of image data.

2.11 Bayesian force inference

Here, we briefly describe the principle of Bayesian force/stress inference [30]. Bayesian force/stress inference solves an ill-conditioned, inverse problem between forces (junction tension and cellular pressure; denoted as $\vec{X}$ in section 2.2) and cell shape. It derives the posterior distribution of cellular forces by combining the likelihood from the force-balance equation at cell vertices (Eq. 10) with the prior informed by experimental observations. The MAP estimate of $\vec{X}$ is obtained by maximizing the marginal likelihood function. Among various priors tested, the one assuming the positivity of junction tension has been shown to yield most accurate and robust estimates of the cellular forces [30, 42, 43, 58]. Details of the formalism, along with results of *in silico* and *in vivo* validation, have been reported in [30, 36, 42, 59–61].

3 Results

3.1 Validation of non-hierarchical Bayesian parameter estimation using synthetic data

The primary aim of this study is to extend and improve the image-based parameter inference [26] by reformulating it in a framework of Bayesian statistics. Image-based parameter inference involves the construction of models and the subsequent inference of parameters (Fig. 1D). For model construction, we followed the approach outlined in our previous study (section 2.1) [26]. The model equations for junction tension (referred to as the anisotropic spring model) and cell pressure were constructed based on image data of *Drosophila* epithelial tissues . The anisotropic spring model comprises two terms, $\Lambda_0(\theta_{ij})$ and $\Lambda_1(\theta_{ij})l_{ij}$ (Fig. 2 and Fig. S1). $\Lambda_0(\theta_{ij})$ represents the line tension, whose magnitude is determined by the baseline λ_0 with the anisotropic component (μ_0, φ_0). Similarly, $\Lambda_1(\theta_{ij})$ consists of the baseline λ_1 and an anisotropic component (μ_1, φ_1), which governs the dependency of junction tension on the junction length. In this and previous studies, λ_0 was fixed at 1.0 as the scaling factor (section 2.2). We employed the standard model with area elasticity coefficient k for cell pressure (section 2.1). For parameter inference, by integrating non-informative prior distributions of these parameters with the likelihood function derived from the force balance at cell vertices, we established the non-hierarchical Bayesian parameter inference framework (section 2.4). The posterior distribution of parameters was computed through MCMC sampling (section 2.5).

To validate this newly developed non-hierarchical Bayesian method, we evaluated the accuracy of parameter estimation using synthetic data with known true parameter values. Synthetic data of the two-dimensional cell configuration was generated through numerical simulation of the CVM, as outlined in section 2.8 (Fig. 3A). Using the vertex position and connectivity in synthetic data as input, we obtained the posterior distribution and MAP estimates of the parameters. The estimation error was negligible, with the maximum error in absolute value being 1.50% (Fig. 3B-E). These data clearly indicate that our non-hierarchical Bayesian method gives accurate estimates of the mechanical parameters *in silico*.

3.2 Non-hierarchical Bayesian method yields parameter estimates consistent with those obtained by the least-squares method *in vivo*

Next, we evaluated the non-hierarchical Bayesian method on *in vivo* data. For this, we assessed its consistency with the least-squares method [26], which is generally expected when employing non-informative priors in a non-hierarchical Bayesian model [4]. Specifically, we applied the non-hierarchical Bayesian method to 165 image datasets of *Drosophila* epithelial tissues (section 2.9), conducted parameter estimation using the best-fitting model selected by the least-squares method, and then compared the estimated values between the two methods. Using the model selected by the least-squares method, the posterior distributions converged for all samples, confirming the reliability of MCMC sampling. The MAP estimates were found to closely match the values inferred by the least-squares method, with a maximum difference in absolute value of 1.84×10^{-6} (Fig. 4A–F).

We further assessed the consistency of model selection between the least-squares method and the non-hierarchical Bayesian method. The former selects models based

on AIC [26], while the latter employs the widely applicable information criterion (WAIC) and leave-one-out cross-validation (LOOCV), which are commonly used model selection criteria in Bayesian statistics [50–52,62] (section 2.6). We found that among candidate models A–E listed in Fig. S1, both WAIC and LOOCV consistently selected the same model for each sample. Using the model selected by WAIC and LOOCV, we could not obtain a converged posterior distribution for one sample, which was subsequently excluded from further analysis. Our data indicated that the selected models were identical in the original and new methods for 158 out of the 164 samples (Fig. 4G). For the remaining six samples, the values of the model selection criteria were nearly indistinguishable between the rank-1 and rank-2 models, suggesting that the inconsistency in model selection arose from subtle variations in the model selection criteria. Overall, the tests on synthetic and *in vivo* data sets indicates that the image-based parameter inference has been adequately reformulated using Bayesian statistics.

3.3 Quantification of uncertainty in parameter estimation revealed its link to mechanical anisotropies in tissue

Assessing the uncertainty in parameter estimation offers insights into the reliability of obtained results [28]. In this study, we used 95% Highest Posterior Density Interval (HDI), the range within which a parameter value fall with a probability of 0.95, as a quantitative measure for the uncertainty associated with parameter estimation (Fig. 1F; section 2.7). Fig. 5A presents the 95% HDI values quantified for each parameter at each developmental stage in *Drosophila* pupal wing, in which different colors represent different models selected by WAIC. The relatively larger 95% HDI of μ_1 may be attributed to the higher order of its explanatory variable, *i.e.*, the product of l_{ij} and $\cos 2(\theta_{ij} - \varphi_1)$. This could also explain why the 95% HDI values were smaller for simpler models.

We then closely examined the 95% HDI values for individual parameters, focusing on their dependence on developmental stages. Our previous study has uncovered the distinctive functions of the tension parameters in the *Drosophila* pupal wing [26]: μ_0 and φ_0, representing the anisotropy in line tension, contribute to resistance against the proximal-distal (PD) tissue stretching induced by extrinsic forces (Fig. 2C, D). On the other hand, μ_1 and φ_1, representing the anisotropy in feedback between junction shrinkage and tension, facilitate the junction shrinkage along the anterior-posterior (AP) direction, perpendicular to tissue stretching (Fig. 2B, E). Here, we demonstrated that the 95% HDI of φ_0 became narrower after 16.5 h APF, consistent with the timing of the onset of tissue stretching (Fig. 5A–D). The 95% HDI of μ_0 exhibited a similar trend. These results suggest that as the wing begins to stretch, the estimation of μ_0 and φ_0 becomes more robust, reflecting the stabilization of tissue-scale mechanical structure. The 95% HDI of μ_1 and φ_1 took smaller values around the AP direction at 21–23 hr APF and 25.5–27.5 h APF, consistent with a stronger need to elevate the junction tension at remodeling AP junctions in these developmental stages (Fig. 5A, E-G). Other two parameters, k and λ_1, did not exhibit significant changes in the 95% HDI values during the wing development (Fig. 5A, Fig. S2, Fig. S3). In summary, the results of uncertainty quantification revealed that the reliability of parameter estimation is tightly linked to the mechanical anisotropies present in the tissue.

3.4 The cortical elasticity term is dispensable for explaining force-shape correlations *in vivo*

Building on the non-hierarchical Bayesian method with non-informative priors, we further extended the image-based parameter inference by adding an informative prior on the cortical elasticity (section 3.4) and by constructing a hierarchical model (sections 3.5 and 3.6).

In the standard model function of the CVM, $T(L_i, L_j) = \lambda_0 + \Gamma(L_i + L_j)$ (Eq. 5), the term $\Gamma(L_i + L_j)$ mathematically represents the elasticity of the cell cortex, where Γ is the elastic coefficient of the cell cortex, and L_i denotes the perimeter length of the i-th cell. In this study, we statistically evaluated the impact of the cortical elasticity term on the mechanical properties of epithelial cells, which are characterized by force-shape correlations. We first analyzed the correlation between junction tension and cell perimeter length to see if they were positively correlated as the cortical elasticity term suggests (Fig. 6A). Despite considerable variation in cell perimeter length among wing cells, no significant correlation with junction tension was observed, suggesting a negligible influence of the cortical elasticity term on the variation in tension magnitude among junctions.

To support this finding, we sought to examine a model described by Eq. 4, which integrates the cortical elasticity term into the anisotropic spring model (model A' in Fig. S1), and compared models with and without this term for its ability to fit to experimental data. However, with the original least-squares method, parameter estimation in model A' frequently led to physically unrealistic, negative estimates for Γ. We therefore decided to adopt the non-hierarchical Bayesian method, imposing a prior that assumes the positivity of Γ (see section 2.4.1 for details). This extension enabled us to perform parameter estimation and model selection using model A' and its simpler counterparts. We found that the estimated values of Γ were close to zero (Fig. 6B), consistent with the minimal correlation between junction tension and cell perimeter length. Moreover, both the WAIC and LOOCV selected models in the category of the anisotropic spring model over those including the cortical elasticity term (Fig. 6C). In summary, all of our data suggest that the cortical elasticity term is not essential for explaining force-shape correlations *in vivo*.

3.5 Validation of hierarchical Bayesian parameter estimation using synthetic data

Classifying samples into groups, rather than treating them individually, and quantifying the characteristics of each group, provides valuable information for understanding the system of interest. Hierarchical Bayesian modeling offers a statistical framework that integrates individual variance with overall group properties, thereby yielding parameters at both the individual and group levels and identifying a most appropriate model of the group [63, 64]. To this end, we extended our Bayesian model into a hierarchical Bayesian model to estimate cell mechanical properties at both the group and individual levels (section 2.4.2; Fig. 1H). Specifically, we adopted the beta distribution as a prior for linear parameters to represent the variance among individual samples within the group. The half-Cauchy distribution is used for a prior for two parameters of the beta distribution, serving as the hyper prior. For axial parameters, we utilized the axial von Mises distribution, setting location parameters as the axial uniform distribution and concentration parameters as the reciprocal of a half-Cauchy distribution with a scale of 5.

We assessed the precision of parameter estimation of the hierarchical Bayesian method using synthetic data. Synthetic data was generated in the following procedures (Fig. 7A; section 2.8). We set true distributions to each parameter of model A, and generated random numbers from these distributions to create 15 sets of true parameters. Numerical simulation of the CVM was then carried out to obtain the polygonal cell configuration. Using the vertex positions and connectivity from the synthetic data as input, we conducted parameter estimation and obtained the posterior distributions at both the group- (Fig. 7B) and the individual level (Fig. 7C). At the individual level, the true and estimated parameter values closely matched (black dots in Fig. 7D–G). Estimation accuracy was high, evidenced by a maximum relative error of only 1.52%. Furthermore, the MAP estimates of the group-level parameters were highly close to the mode of the true distribution (orange squares in Fig. 7D-G). For comparison, we aggregated the input data into a single dataset. The results from the pooled estimation deviated substantially from the true distributions (blue diamonds in Fig. 7D-G), presumably because the distinct characteristics of each input data were averaged out during the aggregation process. Collectively, our data highlight the effectiveness of hierarchical Bayesian modeling.

3.6 Application of hierarchical Bayesian method to *in vivo* data

Having established the effectiveness of our hierarchical Bayesian modeling, we applied the method to *in vivo* data from the *Drosophila* pupal wing. We conducted MCMC sampling across five developmental stages and for five candidate models A–E to obtain representative parameter values and models for each developmental stage. Our inference results indicated that the posterior distributions of mechanical parameters from all candidate models were effectively estimated at developmental stages starting from 16.5 h APF. The MCMC sampling did not converge for models A–C at 13–14 h APF, when cell and tissue mechanics do not exhibit strong anisotropies (see the Discussion for possible solutions). In the four developmental stages where MCMC sampling converged for all models, both WAIC and LOOCV selected the full model, model A, over its simpler counterparts. We then compared the parameter values estimated for model A by the hierarchical Bayesian method with those obtained by the least-squares method. We observed that compared to the results from the least-squares method, parameter values estimated by the hierarchical Bayesian method at the individual level showed a trend of shrinkage toward the group-level estimate (gray arrows in Fig. 8). This shrinkage could be attributed to the incorporation of a group-level distribution of parameters [28]. The group-level estimates aligned closely, regardless of the method used (green and orange dashed lines, and blue solid lines in Fig. 8), presumably because individual samples at the same developmental stage possessed similar parameter values.

4 Discussion

To ensure the validity of mathematical modeling, selecting the most appropriate model among the candidates and determining its parameters based on experimental data is essential. By utilizing Bayesian statistics, this study extended the image-based parameter inference [26], a statistical framework designed for cell-based models in epithelial mechanics that computes model selection and parameter estimation.

We demonstrated that Bayesian formulation enables the quantification of estimation uncertainty, broadens the range of models that can be analyzed, and facilitates the integration of information across different layers. These advantages significantly expand the utility of image-based parameter inference.

Our quantification of estimation uncertainty revealed significant changes in the 95% HDI values during wing development: the 95% HDI of anisotropy parameters in tension functions increased concomitantly with the magnitude of anisotropy in cell/tissue mechanics. These findings strongly suggest that the estimation of mechanical parameters becomes more robust when cell/tissue mechanics are aligned across the tissue, leading to the stabilization of the tissue-scale mechanical structure. Such information helps researchers gauge the reliability of predictions derived from cell-based mechanical models.

Our previous study revealed that the anisotropic spring model is favored over the standard CVM for fitting experimental data of *Drosophila* epithelial tissues [26]. However, the original least-squares method could not accommodate a model that integrates the cortical elasticity term with the anisotropic spring model. Bayesian modeling addressed this limitation by incorporating prior knowledge about the elasticity of cell cortex. This reformulation enabled a comparison between the anisotropic spring model with and without the cortical elasticity term based on statistical criterion. The model selection analysis revealed that for all *in vivo* tested in this study, models excluding the cortical elasticity term were selected. These findings suggest that the cortical elasticity term may not be essential for explaining the force-shape correlation that represent the cell mechanical properties. Alternatively, because there needs to be significant variation in the geometry among cells in the input image data for image-based parameter inference to work properly, the strong physical constraint on the cell perimeter length can hinder the detection of the cortical elasticity term. However, our measurement of cell geometry indicated significant variance in the perimeter length among cells, which contradicts the existence of such a strong physical constraint. Recent studies also highlight the need to revisit the standard equations in cell-based models in epithelial mechanics [17–20]. A quantitative comparison between modeling and experimental data will be crucial for rewriting and testing model functions to more accurately represent the mechanical nature of epithelial junction.

In constructing the hierarchical Bayesian inference, we employed prior distributions based on the assumption that each parameter for a given developmental stage is centered around a specific value. While this hierarchical Bayesian model successfully estimated model parameters for most developmental stages, MCMC sampling became unstable at certain developmental stages where the parameters controlling the strength of tension anisotropy, specifically μ_0 or μ_1, were small. During these stages, the parameters governing anisotropy orientation, namely φ_0 or φ_1, showed significant variation among individual samples that the aforementioned prior distribution could not adequately capture, leading to the instability of MCMC sampling. Therefore, the convergence of MCMC sampling in the hierarchical Bayesian model exhibited a similar trend to estimation uncertainty in the non-hierarchical Bayesian model. We observed that MCMC sampling was stabilized upon adopting a uniform distribution as the prior for the concentration parameter of the axial Von-Mises distribution (κ), which controls the variation of φ_0 and φ_1. These findings suggest the possibility of modifying the hyper prior distribution to incorporate the characteristics of the system of interest.

Adapting Bayesian parameter inference to handle various types of models and

data will pave the way for future research. For instance, by segmenting a whole-tissue image into smaller regions and applying tissue-level hyper-priors in hierarchical Bayesian modeling, one can infer a spatially smooth distribution of parameters while accounting for spatial variations within the tissue. Similarly, using time-lapse data as input, hierarchical Bayesian modeling can integrate both global trend in time-dependent changes and the variance among time-series of mechanical parameters. Furthermore, given recent advancements in cell-based modeling, especially those regarding feedback regulations in epithelial junction [18–20, 65, 66], analyzing recently proposed model functions by using the Bayesian parameter inference will be of great interest. These extensions will contribute to deepening our understanding of cell mechanics during tissue development and repair.

Data availability

The authors declare that the data supporting the findings of this study are available within the paper and its Supplementary files. The data are available from the lead contact upon reasonable request. The code used for Bayesian parameter inference can be downloaded from
https://github.com/Sugimuralab/BayesianParameterInferenceForEpithelialMechanics.

CRediT authorship contribution statement

Xin Yan: formal analysis, investigation, methodology, software, writing - original draft; **Goshi Ogita**: conceptualization, project administration, formal analysis, investigation, methodology, writing - original draft; **Shuji Ishihara**: conceptualization, supervision; writing - review and editing, funding acquisition; **Kaoru Sugimura**: conceptualization, project administration, supervision, writing - original draft, writing - review and editing, funding acquisition. All authors gave their final approval for publication and agreed to be held accountable for the work performed therein.

Funding

c

Acknowledgments

The authors would like to thank Yang Hong, Roger Kares, the Bloomington Stock Center, and the Kyoto Stock Center for fly strains; and Mayu Miyakawa for technical assistance. X.Y. and G.O. are respectively supported by World-leading Innovative Graduate Study for Frontiers of Mathematical Sciences and Physics (WINGS-FMSP) and RIKEN Pioneering Project "Prediction for Science".

References

[1] P. K. Maini, Using mathematical models to help understand biological pattern formation, Comptes Rendus Biologies 327 (3) (2004) 225–234. `doi:10.1016/j.crvi.2003.05.006`.

[2] C. J. Tomlin, J. D. Axelrod, Biology by numbers: Mathematical modelling in developmental biology, Nature Reviews Genetics 8 (5) (2007) 331–340. `doi:10.1038/nrg2098`.

[3] B. P. Ingalls, Mathematical Modeling in Systems Biology: An Introduction, MIT Press, Cambridge, Massachusetts, 2022.

[4] C. M. Bishop, Pattern Recognition and Machine Learning., Springer, New York, 2006.

[5] S. Konishi, Introduction to Multivariate Analysis: Linear and Nonlinear Modeling, Chapman and Hall/CRC, New York, 2014. `doi:10.1201/b17077`.

[6] T. Sullivan, Introduction to Uncertainty Quantification, Vol. 63 of Texts in Applied Mathematics, Springer International Publishing, Cham, 2015. `doi:10.1007/978-3-319-23395-6`.

[7] J. Kursawe, R. E. Baker, A. G. Fletcher, Approximate Bayesian computation reveals the importance of repeated measurements for parameterising cell-based models of growing tissues, Journal of Theoretical Biology 443 (2018) 66–81. `doi:10.1016/j.jtbi.2018.01.020`.

[8] A. G. Fletcher, J. M. Osborne, Seven challenges in the multiscale modeling of multicellular tissues, WIREs Mechanisms of Disease 14 (1) (2022) e1527. `doi:10.1002/wsbm.1527`.

[9] E. D. Mitra, W. S. Hlavacek, Parameter estimation and uncertainty quantification for systems biology models, Current Opinion in Systems Biology 18 (2019) 9–18. `doi:10.1016/j.coisb.2019.10.006`.

[10] A. G. Fletcher, F. Cooper, R. E. Baker, Mechanocellular models of epithelial morphogenesis, Philosophical Transactions of the Royal Society B: Biological Sciences 372 (1720) (2017) 20150519. `doi:10.1098/rstb.2015.0519`.

[11] T. Hirashima, E. G. Rens, R. M. H. Merks, Cellular Potts modeling of complex multicellular behaviors in tissue morphogenesis, Development, Growth & Differentiation 59 (5) (2017) 329–339. `doi:10.1111/dgd.12358`.

[12] L. A. Taber, Continuum Modeling in Mechanobiology, Springer International Publishing, 2020. `doi:10.1007/978-3-030-43209-6`.

[13] H. Honda, T. Nagai, Mathematical Models of Cell-Based Morphogenesis: Passive and Active Remodeling | SpringerLink, Springer-Nature., 2022.

[14] H. Honda, Geometrical Models for Cells in Tissues, in: G. H. Bourne, J. F. Danielli, K. W. Jeon (Eds.), International Review of Cytology, Vol. 81, Academic Press, 1983, pp. 191–248. `doi:10.1016/S0074-7696(08)62339-6`.

[15] A. G. Fletcher, M. Osterfield, R. E. Baker, S. Y. Shvartsman, Vertex Models of Epithelial Morphogenesis, Biophysical Journal 106 (11) (2014) 2291–2304. `doi:10.1016/j.bpj.2013.11.4498`.

[16] S. Alt, P. Ganguly, G. Salbreux, Vertex models: From cell mechanics to tissue morphogenesis, Philosophical Transactions of the Royal Society B: Biological Sciences 372 (1720) (2017) 20150520. `doi:10.1098/rstb.2015.0520`.

[17] R. J. Tetley, G. B. Blanchard, A. G. Fletcher, R. J. Adams, B. Sanson, Unipolar distributions of junctional Myosin II identify cell stripe boundaries that drive cell intercalation throughout Drosophila axis extension, eLife 5 (2016) e12094. doi:10.7554/eLife.12094.

[18] M. F. Staddon, K. E. Cavanaugh, E. M. Munro, M. L. Gardel, S. Banerjee, Mechanosensitive Junction Remodeling Promotes Robust Epithelial Morphogenesis, Biophysical Journal 117 (9) (2019) 1739–1750. doi:10.1016/j.bpj.2019.09.027.

[19] R. Sknepnek, I. Djafer-Cherif, M. Chuai, C. Weijer, S. Henkes, Generating active T1 transitions through mechanochemical feedback, eLife 12 (2023) e79862. doi:10.7554/eLife.79862.

[20] K. Nishizawa, S.-Z. Lin, C. Chardès, J.-F. Rupprecht, P.-F. Lenne, Two-point optical manipulation reveals mechanosensitive remodeling of cell–cell contacts in vivo, Proceedings of the National Academy of Sciences 120 (13) (2023) e2212389120. doi:10.1073/pnas.2212389120.

[21] R. Farhadifar, J.-C. Röper, B. Aigouy, S. Eaton, F. Jülicher, The Influence of Cell Mechanics, Cell-Cell Interactions, and Proliferation on Epithelial Packing, Current Biology 17 (24) (2007) 2095–2104. doi:10.1016/j.cub.2007.11.049.

[22] O. Canela-Xandri, F. Sagués, J. Casademunt, J. Buceta, Dynamics and Mechanical Stability of the Developing Dorsoventral Organizer of the Wing Imaginal Disc, PLOS Computational Biology 7 (9) (2011) e1002153. doi:10.1371/journal.pcbi.1002153.

[23] Y. Mao, A. L. Tournier, P. A. Bates, J. E. Gale, N. Tapon, B. J. Thompson, Planar polarization of the atypical myosin Dachs orients cell divisions in Drosophila, Genes & Development 25 (2) (2011) 131–136. doi:10.1101/gad.610511.

[24] A. Nestor-Bergmann, G. Goddard, S. Woolner, O. E. Jensen, Relating cell shape and mechanical stress in a spatially disordered epithelium using a vertex-based model, Mathematical Medicine and Biology: A Journal of the IMA 35 (Supplement_1) (2018) i1–i27. doi:10.1093/imammb/dqx008.

[25] A. Tsuboi, D. Umetsu, E. Kuranaga, K. Fujimoto, Inference of Cell Mechanics in Heterogeneous Epithelial Tissue Based on Multivariate Clone Shape Quantification, Frontiers in Cell and Developmental Biology 5 (2017). doi:10.3389/fcell.2017.00068.

[26] G. Ogita, T. Kondo, K. Ikawa, T. Uemura, S. Ishihara, K. Sugimura, Image-based parameter inference for epithelial mechanics, PLOS Computational Biology 18 (6) (2022) e1010209. doi:10.1371/journal.pcbi.1010209.

[27] H. Akaike, Information theory and an extension of the maximum likelihood principle (1973).

[28] J. K. Kruschke, Doing Bayesian Data Analysis, Academic Press, Boston, 2015.

[29] I. G. N. M. Jaya, B. Tantular, Y. Andriyana, A Bayesian approach on multicollinearity problem with an Informative Prior, Journal of Physics: Conference Series 1265 (1) (2019) 012021. doi:10.1088/1742-6596/1265/1/012021.

[30] S. Ishihara, K. Sugimura, Bayesian inference of force dynamics during morphogenesis, Journal of Theoretical Biology 313 (2012) 201–211. `doi:10.1016/j.jtbi.2012.08.017`.

[31] M. Rauzi, P.-F. Lenne, T. Lecuit, Planar polarized actomyosin contractile flows control epithelial junction remodelling, Nature 468 (7327) (2010) 1110–1114. `doi:10.1038/nature09566`.

[32] P.-L. Bardet, B. Guirao, C. Paoletti, F. Serman, V. Léopold, F. Bosveld, Y. Goya, V. Mirouse, F. Graner, Y. Bellaïche, PTEN Controls Junction Lengthening and Stability during Cell Rearrangement in Epithelial Tissue, Developmental Cell 25 (5) (2013) 534–546. `doi:10.1016/j.devcel.2013.04.020`.

[33] H. Lan, Q. Wang, R. Fernandez-Gonzalez, J. J. Feng, A biomechanical model for cell polarization and intercalation during *Drosophila* germband extension, Physical Biology 12 (5) (2015) 056011. `doi:10.1088/1478-3975/12/5/056011`.

[34] L. C. Siang, R. Fernandez-Gonzalez, J. J. Feng, Modeling cell intercalation during *Drosophila* germband extension, Physical Biology 15 (6) (2018) 066008. `doi:10.1088/1478-3975/aad865`.

[35] B. Aigouy, R. Farhadifar, D. B. Staple, A. Sagner, J.-C. Röper, F. Jülicher, S. Eaton, Cell Flow Reorients the Axis of Planar Polarity in the Wing Epithelium of Drosophila, Cell 142 (5) (2010) 773–786. `doi:10.1016/j.cell.2010.07.042`.

[36] K. Sugimura, S. Ishihara, The mechanical anisotropy in a tissue promotes ordering in hexagonal cell packing, Development 140 (19) (2013) 4091–4101. `doi:10.1242/dev.094060`.

[37] R. Etournay, M. Popović, M. Merkel, A. Nandi, C. Blasse, B. Aigouy, H. Brandl, G. Myers, G. Salbreux, F. Jülicher, S. Eaton, Interplay of cell dynamics and epithelial tension during morphogenesis of the Drosophila pupal wing, eLife 4 (2015) e07090. `doi:10.7554/eLife.07090`.

[38] B. Guirao, S. U. Rigaud, F. Bosveld, A. Bailles, J. López-Gay, S. Ishihara, K. Sugimura, F. Graner, Y. Bellaïche, Unified quantitative characterization of epithelial tissue development, eLife 4 (2015) e08519. `doi:10.7554/eLife.08519`.

[39] A. Matamoro-Vidal, I. Salazar-Ciudad, D. Houle, Making quantitative morphological variation from basic developmental processes: Where are we? The case of the Drosophila wing, Developmental Dynamics 244 (9) (2015) 1058–1073. `doi:10.1002/dvdy.24255`.

[40] K. Ikawa, K. Sugimura, AIP1 and cofilin ensure a resistance to tissue tension and promote directional cell rearrangement, Nature Communications 9 (1) (2018). `doi:10.1038/s41467-018-05605-7`.

[41] K. K. Chiou, L. Hufnagel, B. I. Shraiman, Mechanical Stress Inference for Two Dimensional Cell Arrays, PLOS Computational Biology 8 (5) (2012) e1002512. `doi:10.1371/journal.pcbi.1002512`.

[42] S. Ishihara, K. Sugimura, S. J. Cox, I. Bonnet, Y. Bellaïche, F. Graner, Comparative study of non-invasive force and stress inference methods in tissue, The European Physical Journal E 36 (4) (Apr. 2013). doi:10.1140/epje/i2013-13045-8.

[43] C. Roffay, C. J. Chan, B. Guirao, T. Hiiragi, F. Graner, Inferring cell junction tension and pressure from cell geometry, Development 148 (18) (2021) dev192773. doi:10.1242/dev.192773.

[44] N. I. Fisher, Statistical Analysis of Circular Data, Cambridge University Press, Cambridge, Mass., 1995.

[45] S. R. Jammalamadaka, A. SenGupta, Topics in Circular Statistics, Vol. 5 of Series on Multivariate Analysis, WORLD SCIENTIFIC, 2001. doi:10.1142/4031.

[46] J. Salvatier, T. V. Wiecki, C. Fonnesbeck, Probabilistic programming in Python using PyMC3, PeerJ Computer Science 2 (2016) e55. doi:10.7717/peerj-cs.55.

[47] A. Kucukelbir, Automatic Differentiation Variational Inference, Journal of machine learning research 18 (14) (2017) 1–45.

[48] M. D. Hoffman, A. Gelman, The No-U-Turn Sampler: Adaptively Setting Path Lengths in Hamiltonian Monte Carlo, J. Mach. Learn. Res. 15 (1) (2014) 1593–1623.

[49] A. Gelman, D. B. Rubin, Inference from Iterative Simulation Using Multiple Sequences, Statistical Science 7 (4) (1992) 457–472. doi:10.1214/ss/1177011136.

[50] A. Gelman, J. B. Carlin, H. S. Stern, D. B. Dunson, A. Vehtari, D. B. Rubin, Bayesian Data Analysis Third Edition, Chapman and Hall/CRC, New York, 2013.

[51] S. Watanabe, Asymptotic Equivalence of Bayes Cross Validation and Widely Applicable Information Criterion in Singular Learning Theory, The Journal of Machine Learning Research 11 (2010) 3571–3594.

[52] A. Vehtari, A. Gelman, J. Gabry, Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC, Statistics and Computing 27 (5) (2017) 1413–1432. doi:10.1007/s11222-016-9696-4.

[53] R. Kumar, C. Carroll, A. Hartikainen, O. Martin, ArviZ a unified library for exploratory analysis of Bayesian models in Python, Journal of Open Source Software 4 (33) (2019) 1143. doi:10.21105/joss.01143.

[54] T. Hiraiwa, F.-L. Wen, T. Shibata, E. Kuranaga, Mathematical Modeling of Tissue Folding and Asymmetric Tissue Flow during Epithelial Morphogenesis, Symmetry 11 (1) (2019) 113. doi:10.3390/sym11010113.

[55] A. Royou, C. Field, J. C. Sisson, W. Sullivan, R. Karess, Reassessing the Role and Dynamics of Nonmuscle Myosin II during Furrow Formation in Early Drosophila Embryos, Molecular Biology of the Cell 15 (2) (2004) 838–850. doi:10.1091/mbc.e03-06-0440.

[56] J. Huang, W. Zhou, W. Dong, A. M. Watson, Y. Hong, Directed, efficient, and versatile modifications of the Drosophila genome by genomic engineering, Proceedings of the National Academy of Sciences 106 (20) (2009) 8284–8289. doi:10.1073/pnas.0900641106.

[57] B. Aigouy, C. Cortes, S. Liu, B. Prud'Homme, EPySeg: A coding-free solution for automated segmentation of epithelia using deep learning, Development (Cambridge, England) 147 (24) (2020) dev194589. arXiv:https://journals.biologists.com/dev/article-pdf/147/24/dev194589/1969911/dev194589.pdf, doi:10.1242/dev.194589.

[58] K. Sugimura, P.-F. Lenne, F. Graner, Measuring forces and stresses in situ in living tissues, Development 143 (2) (2016) 186–196. doi:10.1242/dev.119776.

[59] K. Sugimura, Y. Bellaïche, F. Graner, P. Marcq, S. Ishihara, Robustness of force and stress inference in an epithelial tissue, in: 2013 35th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), 2013, pp. 2712–2715. doi:10.1109/EMBC.2013.6610100.

[60] W. Kong, O. Loison, P. Chavadimane Shivakumar, E. H. Chan, M. Saadaoui, C. Collinet, P.-F. Lenne, R. Clément, Experimental validation of force inference in epithelia from cell to tissue scale, Scientific Reports 9 (1) (2019) 14647. doi:10.1038/s41598-019-50690-3.

[61] S. Miyasaka, K. Izumi, S. Okuda, Y. Miki, Numerical assessment of the applicability of geometry-based force inference on homogeneous and heterogeneous cells, PLOS ONE 19 (4) (2024) e0299016. doi:10.1371/journal.pone.0299016.

[62] O. Martin, Bayesian Analysis with Python: Introduction to Statistical Modeling and Probabilistic Programming Using PyMC3 and ArviZ, 2nd Edition, Packt Publishing, Birmingham, 2018.

[63] K. Katahira, How hierarchical models improve point estimates of model parameters at the individual level, Journal of Mathematical Psychology 73 (2016) 37–58. doi:10.1016/j.jmp.2016.03.007.

[64] R. van de Schoot, S. Depaoli, R. King, B. Kramer, K. Märtens, M. G. Tadesse, M. Vannucci, A. Gelman, D. Veen, J. Willemsen, C. Yau, Bayesian statistics and modelling, Nature Reviews Methods Primers 1 (1) (2021) 1–26. doi:10.1038/s43586-020-00001-2.

[65] J. Rozman, J. M. Yeomans, R. Sknepnek, Shape-Tension Coupling Produces Nematic Order in an Epithelium Vertex Model, Physical Review Letters 131 (22) (2023) 228301. doi:10.1103/PhysRevLett.131.228301.

[66] S.-Z. Lin, M. Merkel, J.-F. Rupprecht, Structure and Rheology in Vertex Models under Cell-Shape-Dependent Active Stresses, Physical Review Letters 130 (5) (2023) 058202. doi:10.1103/PhysRevLett.130.058202.

[67] G. Ogita, Image-based model construction and parameter estimation for epithelial mechanics, doctoral thesis, Kyoto University (Sep. 2022). doi:10.14989/doctor.k24269.

Figure 1: Bayesian parameter inference for epithelial mechanics. (A) The structure of the monolayer epithelium. Left: The top view of the *Drosophila* pupal wing epithelium, in which the adherens junction (AJ) is labeled with DE-cadherin-GFP. Right: Epithelial cells are connected to each other via cell adhesion molecules such as cadherin (green). Actomyosin cables (red) run along the cell cortex in the AJ plane. Scale bar: 10 μm. (B) Forces acting along the AJ plane. Left: Junction tension. Right: Cell pressure. (C) Balance of junction tensions (T) and cell pressures (P) acting on the cell vertex. Black dots indicate cell vertices. (D) Flowchart of the method for estimating mechanical parameters from image data, in which the cell contour is visualized [26]. (E) According to Bayes' theorem, the posterior distribution P(parameter|data) is calculated from the likelihood P(data|parameter) and the prior distribution P(parameter). It provides an estimate of a parameter and uncertainty associated with estimation. In the image-based parameter inference, the likelihood function is derived from the force balance equation at cell vertices. (F) An example of the posterior distribution obtained by MCMC sampling. The red arrow indicates the maximum a posteriori (MAP) estimate, while the orange double-headed arrow represents the 95% highest density interval (HDI), which quantifies the estimation uncertainty. (G, H) The Kruschke style diagram [28] of the non-hierarchical and hierarchical Bayesian models used in this study (G: non-hierarchical, H: hierarchical). The characters represent the parameters or variables whose dependencies are indicated by arrows. The symbols $\sim$ and $=$ next to the arrows respectively indicate probabilistic and deterministic dependency. See section 2.4 for details about the formulation of Bayesian models. (A)–(D) are adapted from [26].

Figure 2: The anisotropic spring model for junction tension. (A) The anisotropic spring model consists of two terms, $\Lambda_0(\theta_{ij})$ and $\Lambda_1(\theta_{ij})l_{ij}$ (section 2.1) [26]. $\Lambda_0(\theta_{ij})$ represents line tension, with the magnitude and orientation of the anisotropy determined by μ_0 and φ_0, respectively. $\Lambda_1(\theta_{ij})$ represents junction length-dependent junction tension, with its magnitude and orientation of anisotropy determined by μ_1 and φ_1, respectively. Schematics show how the values of μ_0 and μ_1 alter the dependence of tension on junction length and orientation, in which line colors denote the orientation of junctions. (B) Schematics illustrating the feedback between junction shrinkage and myosin-II-generated junction tension. When the junction tension is inversely related to the junction length, shrinkage of a junction (indicated by the left-pointing arrow) leads to an increase in junction tension (indicated by the upward arrow), which in turn shortens the junction. This feedback is associated with the accumulation of myosin-II at shrinking junctions. Red arrows represent junction tension, while line colors indicate myosin-II concentration along the junctions. (C) Schematic of the adult fly. Orange rectangle indicates the region studied in this study. Purple line indicates the orientation of tissue tension. Schematics of junction are shown in the right. The hinge is shaded gray. (D, E) Schematics illustrating the biological functions of $\Lambda_0(\theta_{ij})$ and $\Lambda_1(\theta_{ij})$ during the development of the *Drosophila* pupal wing [26]. Red arrows represent the junction tension, and purple double-sided arrows indicate the extrinsic pulling force from the hinge. Line colors denote the myosin-II (myo-II) concentration along the junctions. (D) $\Lambda_0(\theta_{ij})$ contributes to resistance against the extrinsic pulling force from the hinge. (E) $\Lambda_1(\theta_{ij})$ facilitates the shrinkage of AP junctions by upregulating the junction tension through the feedback depicted in (B). The myo-II enrichment along the remodeling junctions is indicated by the change in line colors from pale pink to red along the vertical shrinking junction. (B) and (C)–(E) are adapted from [67] and [26], respectively.

Figure 3: Validation of of the non-hierarchical Bayesian method using synthetic data. (A) Schematic of a perfect model experiment to asses the accuracy of parameter estimation. (B–E) Estimated values of parameters are plotted against their true values for μ_0 (B; the magnitude of anisotropy in the line tension), λ_1 (C; the spring constant of junction), μ_1 (D; the magnitude of anisotropy in the spring constant of junction), and k (E; the area elasticity of cell). A blue dashed line indicates $y = x$. (A) is modified from [26].

Figure 4: Comparison between the non-hierarchical Bayesian method and the least-squares method using *in vivo* data. (A-F) The evaluation of parameter estimation of μ_0 (A), λ_1 (B), μ_1 (C), k (D), φ_0 (E), and φ_1 (F). Each dot represents the difference in parameter estimations between the non-hierarchical Bayesian Model (this study) and the original least-squares method [26] for each sample from the different developmental stages of the pupal wing. Dot colors represent the models selected by AIC in the least-squares method, which were subsequently used for parameter estimation in the non-hierarchical Bayesian method. (G) Comparison of model selection between the least-squares and non-hierarchical Bayesian methods. Each row and column represents the selected model by the least-squares (LS) and the non-hierarchical-Bayesian (Bayes) methods, respectively. MCMC sampling converged 823 out of the 835 total instances (165 samples across 5 candidate models). The remaining two instances did not affect the outcome of model selection.

A

Stage [h APF] \ Parameter	μ_0 (magnitude of anisotropy in line tension)	λ_1 (spring constant of junction)	μ_1 (magnitude of anisotropy in spring constant of junction)	k (area elasticity of cell)	φ_0 (orientation of anisotropy in line tension)	φ_1 (orientation of anisotropy in spring constant of junction)
13-14	173.0% 94.4%	13.3% 13.7% 9.9%	132.7%	50.8% 56.9% 39.4%	2.706 0.545	1.106
16.5-18.5	115.3% 29.2%	14.9% 13.0%	129.7%	44.0% 43.9%	0.523 0.144	0.816
21-23	72.6% 25.5%	12.2% 11.0%	55.8%	44.6% 49.6%	0.355 0.129	0.307
25.5-27.5	48.8% 9.7%	10.1% 8.8%	72.6%	42.2% 38.0%	0.234 0.047	0.390
30-32	49.6% 8.8%	9.8% 8.4%	112.2%	34.5% 33.4%	0.236 0.043	0.664

Figure 5: Quantification of uncertainty in parameter estimation using the non-hierarchical Bayesian method with *in vivo* data. (A) The average width of 95% HDI relative to the MAP estimate (section 2.7) is listed for each parameter at each developmental stage of the pupal wing. The colors represent the models selected by WAIC and LOOCV. Each row corresponds to a developmental stage, indicated by hours after puparium formation (h APF), and each column corresponds to a model parameter. (B-G) The 95% HDI for φ_0 (B-D) and φ_1 (E-G) at 13-14 (B, E), 16.5-18.5 (C, F) and 21-23 (D, G) h APF in the pupal wing. Lines and points indicate the 95% HDI and the MAP estimate, respectively. The index represents different individuals at different stages of development. The colors represent the models selected by WAIC and LOOCV.

Figure 6: Parameter inference in the non-hierarchical Bayesian method that incorporates the cortical elasticity term. (A) The correlation between the inferred junction tension and the cell perimeter in *Drosophila* pupal wing at 18 h 5 min APF. Junction tensions are quantified using Bayesian force inference [30]. Each junction is classified by its orientation relative to the horizontal axis (semicircle; PD: proximal-distal and AP: anterior-posterior). The horizontal axis represents the average of the perimeters of cells adjoined by the same junction, normalized by the square root of the average cell area in the sample. (B) The posterior distribution of Γ, the cortical elasticity of cell perimeter, obtained from the sample analyzed in (A). (C) Results of model selection between the anisotropic spring model with (A–E) and without (A'–E') the cortical elasticity term (see Fig. S1 for the list of models). MCMC sampling converged 1644 out of the 1650 total instances (165 samples across 10 candidate models). The remaining six instances did not affect the outcome of model selection.

Figure 7: Validation of the hierarchical Bayesian model using synthetic data. (A) Schematic diagram illustrating the generation of synthetic data for testing the hierarchical Bayesian model. Individual-level parameters $\vec{\beta}_0, \ldots, \vec{\beta}_{14}$ are sampled from the predefined group-level distribution. Synthetic data are then generated by numerically simulating the CVM using these parameters. (B, C) Posterior distribution of the group-level (B) and individual-level (C) parameters for μ_1. In (C), different colors indicate different samples from the same developmental stage. (D–G) Estimated values of parameters are plotted against their true values for μ_0 (D), λ_1 (E), μ_1 (F), and k (G). The black dots and orange squares represent the individual-level and group-level parameter estimates, respectively, obtained by the hierarchical Bayesian method. The blue diamonds show results from pooled estimation, where all individual data within a group are aggregated and then a single parameter set is estimated using the least squares method. A dashed line indicates $y = x$.

Figure 8: Application of the hierarchical Bayesian model to *in vivo* data. (A–E) Shrinkage plots for μ_0 (A), λ_1 (B), μ_1 (C) and k (D), φ_0 (E), φ_1 (F) for data at the stage of 16.5–18.5 h APF in *Drosophila* pupal wing. Estimated values using the hierarchical Bayesian method are displayed at both the individual (orange triangles) and group levels (orange dashed line), alongside values estimated by the original least-squares method (green crosses), the average of these unpooled values (green dashed line), and the estimate from the pooled sample (blue solid line). Gray arrows highlight the shrinkage of estimated values from the least-squares method to those obtained by the hierarchical Bayesian method.

A

Line colors: $\theta_{ij} = \varphi_0, \varphi_0 + \pi$; $\theta_{ij} = \varphi_0 \pm \pi/2$; No θ_{ij} dependency

B

Model	λ_0	μ_0	φ_0	λ_1	μ_1	φ_1	Γ
A	+	+	+	+	+	+	-
B	+	+	+	+	-	-	-
C	+	-	-	+	-	-	-
D	+	+	+	-	-	-	-
E	+	-	-	-	-	-	-
A'	+	+	+	+	+	+	+
B'	+	+	+	+	-	-	+
C'	+	-	-	+	-	-	+
D'	+	+	+	-	-	-	+
E'	+	-	-	-	-	-	+

Figure S1: Candidate tension models. (A) Anisotropic spring model (model A) and its simpler counterparts (model B to E). Equations for junction tension are provided for each candidate model, with parameters highlighted in orange. Schematics show how the values of μ_0 and μ_1 influence the dependence of tension on junction length and orientation. Line colors indicate the orientation of the junctions. (B)Table listing all candidate tension models used in this study. Each row corresponds to a candidate model, and each column to a parameter. The + and − signs respectively represent indicate whether a parameter was included or excluded from model equations. (A) is adapted from [26].

Figure S2: 95% HDI of parameters estimated using non-hierarchical Bayesian model with *in vivo* data. (A-L) The 95% HDI for parameters μ_0 (A, B), λ_1 (C, D), μ_1 (E, F), k (G, H), φ_0 (I, J), and φ_1 (K, L) at 13-14 (A, C, E, G, I, K) and 16.5-18.5 (B, D, F, H, J, L) h APF in pupal wing. Lines and points indicate the 95% HDI and the MAP estimate, respectively. The index represents different individuals at different stages of development. The colors represent the models selected by WAIC and LOOCV.

Figure S3: 95% HDI of parameters estimated using non-hierarchical Bayesian model with *in vivo* data. (A-R) The 95% HDI of the posterior distributions for parameters μ_0 (A-C), λ_1 (D-F), μ_1 (G-I), k (J-L), φ_0 (M-O), and φ_1 (P-R) at 21-23 (A, D, G, J, M, P), 25.5-27.5 (B, E, H, K, N, Q) and 30-32 (C, F, I, L, O, R) h APF in pupal wing. Lines and points indicate the 95% HDI and the MAP estimate, respectively. The index represents different individuals at different stages of development. The colors represent the models selected by WAIC and LOOCV.