Identification of Suitable Target/E3 Ligase Pairs for PROTAC Development using a Rapamycin-induced Proximity Assay (RiPA)

Bikash Adhikari[1], Katharina Schneider[1], Mathias Diebold[1, 2], Christoph Sotriffer[2], and Elmar Wolf[1]

[1] Institute of Biochemistry, University of Kiel, Kiel, Germany

[2] Institut für Pharmazie und Lebensmittelchemie, University of Würzburg, Würzburg, Germany.

Correspondence: elmar.wolf@biochem.uni-kiel.de

Abstract

The development of proteolysis targeting chimeras (PROTACs), which induce the degradation

of target proteins by bringing them into proximity with cellular E3 ubiquitin ligases, has

revolutionized drug development. While the human genome encodes more than 600 different

E3 ligases, current PROTACs use only a handful of them, drastically limiting their full potential.

Furthermore, many PROTAC development campaigns fail because the selected E3 ligase

candidates are unable to induce degradation of the particular target of interest. As more and

more ligands for novel E3 ligases are discovered, the chemical effort to identify the best E3

ligase for a given target is exploding. Therefore, a genetic system to identify degradation-

causing E3 ligases and suitable target/E3 ligase pairs is urgently needed. Here we used the

well-established dimerization of the FKBP12 protein and FRB domain by rapamycin to bring

the target protein WDR5 into proximity with candidate E3 ligases. Strikingly, this rapamycin-

induced proximity assay (RiPA) revealed that VHL, but not Cereblon, is able to induce WDR5

degradation - a finding previously made by PROTACs, demonstrating its predictive power. By

optimizing the steric arrangement of all components and fusing the target protein with a

minimal luciferase, RiPA can identify the ideal E3 for any target protein of interest in living

cells, significantly reducing and focusing the chemical effort in the early stages of PROTAC

development.

Introduction

New technologies such as genome sequencing and genetic screens have led to an drastic

increase of knowledge about disease-causing proteins and potential therapeutic targets.

However, the therapeutic exploitation of this knowledge was often limited by the fact that only

a minority of the newly identified targets could be blocked by conventional, i.e. monovalent

inhibitors. Thus, most of the existing therapies are so far directed against proteins with

enzymatic activity, leaving about 80% of the intracellular proteome undruggable [1]. This

limitation is largely eliminated by the concept of proteolysis targeting chimeras (PROTACs).

PROTACs are bifunctional small molecules which bind to their target and a cellular E3 ligase

and hence induce target ubiquitylation and degradation [2-4]. In addition to the ability of

PROTACs to block the primary function of many intracellular targets, proteolytic degradation

of disease-causing proteins offers further advantages over conventional inhibition. First,

complete degradation results in the inactivation of any protein function, such as all enzymatic

activities in multi-enzyme proteins or non-enzymatic activities in enzymes with additional

scaffolding functions. Second, because PROTACs remain active after once having induced

the degradation of their target (= catalytic mode of action), they act more persistently and at

lower drug concentrations, making them more specific than classical inhibitors (=occupancy-

based mode of action) [5]. Third, since the transient formation of the ternary complex of target,

PROTAC, and E3 ligase is sufficient to ubiquitylate the target, PROTACs are less susceptible

to the formation of resistance-mediating mutations that reduce the affinity of the target for the

drug [6]. Therefore, despite its novelty, the PROTAC concept has transformed the development

of targeted therapies, and numerous PROTACs are already in clinical development with the

first cases of successful clinical trials recently being reported [7,8].

Despite the revolutionary potential of PROTACs, their development is also fraught with

difficulties. Probably the most serious limitation is the current restricted coverage of E3 ligases.

While the human genome encodes for more than 600 E3 ligases, only a few are used to

develop PROTACs [3], and most PROTACs are based on only two of them, Cereblon (CRBN)

and the von Hippel-Lindau tumor suppressor (VHL). VHL- and CRBN-based PROTACs are

potent in inducing degradation of their targets but also have drawbacks. For example, both E3

ligases are commonly expressed in human tissues preventing their use for tissue-specific

degradation. In addition, both are not essential for many cells in our body, which could lead to

the rapid development of resistance if the respective E3 ligase machinery is silenced, e.g. in

the context of an oncology application. Intensive efforts are therefore being made to identify

suitable ligands for additional E3 ligases which overcome these shortcomings.

A second major limitation is that the development of PROTACs is largely empirical, with few

generally applicable design rules [9]. As a result, many PROTAC development campaigns fail

due to the inability to identify PROTACs that efficiently induce target degradation, and in most

cases, these results are not published. A possible reason for this frequent failure could be the

incompatibility of a particular E3 ligase with a given target. In fact, it has been observed that

promiscuous kinase inhibitors exhibit very distinct degradation profiles when used as ligands

for PROTACs, demonstrating that the efficacy of PROTAC-mediated degradation is not

determined solely by the sheer affinity of the target and E3 ligand but rather by efficient ternary

complex formation [10]. We made similar findings on PROTACs we developed for Aurora A

kinase. While the kinase inhibitor we used for PROTAC development also binds and inhibits

the homologous enzyme Aurora B, our Cereblon-based PROTACs show remarkable

specificity towards Aurora A [11]. We observed that certain amino acids on the putative Cereblon

interaction surface of Aurora A are not conserved in Aurora B, and that mutation of these

residues decreases PROTAC-mediated degradation of Aurora A, indicating that ternary

complex formation is supported by direct interactions between Aurora A and Cereblon. In line

with our finding, the Ciulli lab demonstrated that cooperativity in ternary complex formation is

relevant for PROTAC efficacy [12,13]. Taken together, these results suggest that certain E3

ligases are better suited for specific targets than others.

The problem that only certain E3 ligases can efficiently degrade a particular target protein is

exacerbated by the rapid discovery of new ligands for additional E3 ligases (such as IAPs [14,15],

DCAF11 [16], DCAF15 [17,18], DCAF16 [19], RNF4 [20], RNF114 [21,22], AhR [23], FEM1B [24], and KEAP1

[25-27]). In fact, the availability of new E3 ligase ligands will increase the failure rate of PROTAC

development campaigns for reasons more trivial than the inability to efficiently form ternary

complexes (**Figure 1A**). Since these E3 ligases will be much less characterized than Cereblon

and VHL, it may be unknown whether the target exposes a lysine residue within reach of the

E3 ligase, or whether these E3 ligases are even able to attach degradative ubiquitin chains

(i.e. K48, K11) or are located in the same cellular compartment as the target.

We therefore see an urgent need for genetic assays that allow us to identify suitable target/E3

ligase pairs. This assay should not only be target-specific but also applicable to different

cellular systems. In addition, it should be scalable to ideally cover the entire E3 ligase

proteome while remaining easy to use. We decided to exploit the widely used rapamycin-

dimerization system to test whether induced cellular proximity of a specific target/E3 ligase

pair induces target degradation. Our system correctly predicted PROTAC-mediated

degradation of candidate target proteins and allows accurate, time-resolved estimation of

target degradation by measuring luciferase activity in living cells, and will likely streamline

medicinal chemistry efforts during PROTAC development.

Results

Rapamycin-induced proximity assay (RiPA) induces quantifiable degradation of target
proteins

To induce artificial proximity between target proteins and cellular E3 ligases, we exploited the
rapamycin-induced dimerization of the proteins, FK506 binding protein (FKBP12) and the
FKBP12-rapamycin binding (FRB) domain of FKBP12-rapamycin associated protein (**Figure**
1 - figure supplement 1A) [28]. We cloned FRB and FKBP12 into lentiviral vector systems that
allow both transient transfection and stable cell line generation by viral transduction. We also
included a multiple cloning site for easy target protein cloning and enabled robust expression
using the spleen focus forming virus (SFFV) promoter (**Figure 1B**).

To test whether this rapamycin-induced dimerization assay (RiPA) could induce target
degradation, we inserted the WD repeat-containing protein 5 (WDR5) and the von Hippel-
Lindau tumour suppressor (VHL) into the FKBP12 and FRB-containing plasmid, respectively.
We chose this target/E3 ligase pair because we [29] and others [30,31] have previously
demonstrated robust degradation of WDR5 by PROTACs harnessing VHL. We started to
optimize the RiPA system by transfecting different amounts of the WDR5 and VHL-encoding
plasmids into HEK293 cells and incubating them with 0.1 µM rapamycin for 6 hours (**Figure**
1C). Immunoblotting showed that rapamycin-induced a marked reduction in WDR5-FKBP12
when a 10 or 100-fold excess of VHL-FRB was expressed, but not when the target protein
and E3 ligase were transfected at equimolar ratios (**Figure 1D**). FRB alone did not decrease
WDR5-FKBP12 levels, demonstrating that VHL causes degradation of WDR5-FKBP12 in
response to rapamycin. For a more quantitative and convenient readout of target protein
levels, we fused a *Oplophorus gracilirostris*-based minimal luciferase [32] to either the N- (Luc-
WDR5-FKBP12) or C-terminus (WDR5-Luc-FKBP12) of WDR5 (**Figure 1 - figure**
supplement 1B) and expressed them in HEK293 cells. Both conditions allowed robust
measurement of luciferase activity, demonstrating that terminal and internal tagging are
compatible with luciferase activity. Strikingly, rapamycin reduced luciferase activity by 50.2 %
(± 5.8 %) when expressed together with a 10- or 100-fold excess of VHL-FRB (**Figure 1E**).
Immunoblotting confirmed the robust degradation of luciferase and FKBP12-tagged WDR5 by
VHL (**Figure 1F**).

RiPA correctly predicts suitability of E3 ligases for WDR5 PROTACs

We next wanted to evaluate whether the RiPA assay has predictive power, i.e. whether it can
discriminate between suitable and unsuitable target/E3 ligase pairs, or whether any E3 ligase
can lead to target degradation after induced proximity. To this end, we expressed the E3 ligase
Cereblon in fusion with FRB (CRBN-FRB) under the same conditions and compared its ability

to degrade WDR5 with VHL-FRB. Both immunoblotting and luciferase assays again revealed

a robust and time-dependent degradation of WDR5 by VHL-FRB but not by CRBN-FRB

(**Figure 2A,B**). This is relevant because several studies with PROTACs indicate efficient

degradation of WDR5 by VHL but not by CRBN [29,31]. In contrast, CRBN-based PROTACs

have been shown to efficiently degrade the mitotic kinase Aurora A [10,11,33-35]. We therefore

tested whether VHL and CRBN are able to degrade Aurora A in the RiPA system. In contrast

to WDR5, both E3 ligases induced a robust degradation of Aurora A as shown by

immunoblotting (**Figure 2C**) and luciferase activity (**Figure 2D**).

While these results suggest that the respective E3 ligase ubiquitylates the target protein

(WDR5 or Aurora A), it cannot be excluded that luciferase or FKBP12 could also serve as a

substrate for ubiquitylation. In fact, luciferase and FKBP12 proteins contain 7 and 8 lysine

residues respectively. Therefore, we analyzed the available crystal structures of both proteins

and found that all 15 lysine residues are located on the protein surfaces accessible for post-

translational modifications (**Figure 2E**). To exclude that ubiquitylation of luciferase or FKBP12

interfered with the RiPA results, we converted all lysine residues to arginine and expressed

WDR5 together with this lysine-less version of luciferase and FKBP12. While lysine

substitution reduced luciferase activity by 35.4 %, the measurement of luciferase was still

reliable (**Figure 2F**). Strikingly, rapamycin-induced proximity to VHL under these conditions

induced WDR5 degradation comparable to lysine-containing protein tags (**Figure 2G,H**),

demonstrating that VHL induces direct ubiquitylation of WDR5. We conclude that the RiPA

system is capable of inducing target ubiquitylation and predicting suitable target/E3 ligase

pairs.

RiPA can identify degradative E3 ligases not previously used for PROTACs

The RiPA system is well suited to quantify target protein degradation by VHL and Cereblon.

However, these two E3 ligases are by far the most commonly used for PROTACs, so it is

possible that they are exceptional in terms of potency or substrate promiscuity. We therefore

wondered whether the RiPA system could detect and quantify target degradation by E3 ligases

not currently used for PROTACs. We chose the E3 ligase FBXL12 as a candidate because its

degradative function was recently suggested by pooled genome-wide screens [36], but had not

been exploited by PROTACs. We cloned FBXL12 into the FRB-containing entry vector and

co-expressed it with WDR5 containing the lysine-free FKBP12 tag in HEK293 cells. Incubation

with rapamycin-induced robust degradation of WDR5 as assessed by immunoblotting and

luciferase activity assays (**Figure 3A,B**). Similar to our observation with VHL, the position of

the luciferase in the WDR5-FKBP fusion protein is irrelevant to its activity and degradation by

FBXL12 (**Figure 3 - figure supplement 1A**). A direct comparison of WDR5 degradation

induced by FBXL12 and VHL showed a strikingly superior activity of FBXL12 (**Figure 3C**).

Next, we wondered how the steric arrangement between a target protein and the FRB tag

affects its degradation. So far, we have used a linker that is 8 amino acids long (2xGSSG).

Together with the size of FRB and FKBP12, this linker exceeds the length of most PROTACs

(**Figure 3D**). We varied the linker length between WDR5 and FKBP12 from 0 to 8x GSSG and

compared its effect on expression and degradation by FBXL12. Immunoblots showed that the

cellular expression level of the WDR5 fusion protein declined with increasing linker length

(**Figure 3E**). Strikingly, the construct without any linker between WDR5 and FKBP12 is not

only the best expressed but also degraded most drastically by FBXL12 upon the addition of

rapamycin (**Figure 3E,F**). We conclude that both tight and more flexible arrangements

between the target protein and the dimerization tag allow quantification of E3 ligase activity,

but that a seamless fusion shows the most efficient degradation, possibly mimicking the

cooperativity of PROTAC-induced ternary complex formation.

Until now we have analyzed target degradation in the RiPA system by immunoblotting or by

measuring luciferase activity after cell lysis. Since it is not only relevant how completely

PROTACs induce target degradation, but also how fast, we tried to adapt the workflow for

kinetic analyses. For this purpose, we replaced the luciferase substrate furimazine with

endurazine, which is continuously taken up and activated by the cells. We incubated WDR5

and FBXL12-expressing living cells with endurazine before adding rapamycin and measured

luciferase activity every 15 minutes for 6 hours followed by every 30 minutes (**Figure 3G**).

Rapamycin-mediated dimerization with FBXL12 induced a robust and durable degradation of

WDR5, reaching a maximum after 5 hours. We concluded that RiPA is able to estimate the

degradation of "novel" E3 ligases and allows kinetic measurements in living cells.

Identification of degradation-inducing E3 ligases by designing a universal substrate

So far, our goal has been to optimize the RiPA system to identify suitable target/E3 ligase

pairs by ensuring exclusive ubiquitylation of the specific protein target. A slightly different

question is whether an E3 ligase can degrade a substrate at all, i.e., whether it can catalyze

the addition of degradation-inducing ubiquitin chains. In fact, based on the Reactome Pathway

and the UniProt database, only about 40% of 600 or more E3 ligases annotated in the human

genome are believed to be associated with ubiquitin-proteasome system [37]. Therefore, we

aimed to develop a universal substrate that is highly susceptible to ubiquitylation by increasing

the available lysine receptor residues using two different strategies. First, we analyzed

arginine residues on the surface of luciferase that can be mutated without a large impact on

protein structure or substrate binding. Based on this consideration, we designed a mutant

containing 5 (Luc[V1]) lysine residues in addition to 7 surface residues of the wild-type protein

(**Figure 2E, 4A**). As there were still surface patches without exposed lysines on Luc[V1], we

carefully designed another mutant containing 6 additional lysine residues (Luc[V2]), resulting in

a total of 18 surface lysine residues on the protein (**Figure 4A**). Second, we fused luciferase

to three different peptides containing a different number and arrangement of additional lysine

residues (Luc^{K3}, Luc^{K6}, or Luc^{K12}; **Figure 4B**).

All luciferase mutants except Luc^{K12}, which contains 12 consecutive lysine residues in the

peptide tail, are expressed to a level that allows robust measurements (**Figure 4C**). We then

co-expressed these mutants with FBXL12 and observed that the luciferase versions with

additional lysine residues on the cell surface (Luc^{V1}, Luc^{V2}) were degraded at the same rate

and to a similar extent as wild-type luciferase. In contrast, the additional lysines in Luc^{K3} and

Luc^{K6} dramatically increased their degradation by FBXL12 (**Figure 4D**). Kinetic measurements

in living cells revealed that more than 60% of Luc^{K6} is already degraded 2 hours after the

addition of rapamycin (**Figure 4E**). Taken together, these results demonstrate that mutant

versions of luciferase can be used as universal reporters to stratify E3 ligases in terms of their

degradative potential.

Discussion

While PROTACs have revolutionized drug development, the design of such molecules,

including the choice of appropriate E3 ligases, is still largely empirical [38]. A growing body of

literature indicates that often a specific E3 ligase cannot be used for PROTAC-induced

degradation of a given target protein and that certain target/E3 ligase pairs are more suitable

than others [10,39-41]. Therefore, we developed a rapamycin-based proximity assay (RiPA) to

quantitatively measure the degradation of specific targets by E3 ligases when brought

together. We have carefully designed, tested, and modified the RiPA system to demonstrate

that it has features that allow it to be used for E3 ligase selection in PROTAC development

campaigns.

First, we were able to show that RiPA is in principle able to induce target degradation by

analyzing the degradation of WDR5 when brought into proximity with VHL. Second, we

constructed a set of lentiviral vectors that can be used for straightforward cloning of the E3

ligase and target candidates, and can be delivered to eucaryotic cells by both viral transduction

and transient transfection. While viral transduction allows the use of difficult-to-transfect cell

lines and the careful characterization of a few specific target/E3 ligase pairs, transient

transfection allows broad screening, potentially with all of more than 600 annotated E3 ligases

under S1 conditions. We observed that the plasmid encoding the candidate E3 ligase must be

transfected in at least 10-fold excess over the target protein-containing plasmid for efficient

degradation. Third, we observed that both, FRB and the minimal luciferase used for

dimerization and detection of the target protein respectively, still function when all lysines are

mutated to arginine. This ensures that RiPA only reports degradation of the target protein

when it is directly ubiquitylated since ubiquitylation of the reporter can be excluded. We
envision that this setting will be useful for identifying the most suitable E3 ligase candidates
for PROTACs targeting a particular protein. Conversely, we have also constructed reporters
that contain lysines in addition to their endogenous lysines and have observed drastically
increased degradation in some cases. These tools can be used to analyze whether an
uncharacterized E3 ligase can in principle add degradation-inducing ubiquitin chains, thus
complementing data from pooled genetic screens with fluorescent reporters [36]. Comparing
degradation with different types of these universal reporters allows estimation of the degree
of E3 ligase promiscuity that is desired for PROTACs and thus identification of "PROTACable"
E3 ligases. Finally, by combining luminescent reporters with different substrates, RiPA can be
used to precisely quantify target protein degradation in living cells and in extracts of lysed
cells. Taken together, RiPA is a versatile system that is easy to use with standard molecular
biology laboratory equipment.

While this technical study focused on the RiPA system as a resource, we made some notable
observations. The most striking finding was the lack of degradation of WDR5 by Cereblon.
While VHL efficiently reduced WDR5 levels by more than 50% within 6 hours of rapamycin
addition, Cereblon did not induce WDR5 degradation under any condition or time frame. The
lack of WDR5 degradation is not due to a general inability of Cereblon to induce protein
degradation RiPA conditions, since Cereblon efficiently decreased Aurora A levels. This is
noteworthy because it recapitulates our experience and that of others with the WDR5 and
Aurora A PROTACs: while Cereblon-based PROTACs were able to efficiently degrade Aurora
A [10,11,33-35], they were either nonfunctional [29] or much less potent [30,31] at degrading WDR5
compared to VHL-based PROTACs. A possible reason for the selective potency of both E3
ligases to degrade WDR5 by PROTACs could be their different ability to bind to WDR5 through
protein-protein interactions and thus support ternary complex formation. Indeed, WDR5 could
be co-crystalized with a variety of PROTACs and the respective E3 ligase, resulting in ternary
complexes with a range of interface sizes between the two proteins. The most potent
PRTOACs however, induced the largest interaction area, underlining the importance of
achieving extensive protein-protein interactions for the most efficient target degradation [42].
Strikingly, WDR5 degradation also decreased in the RiPA setting as the linkers between
WDR5 and FKBP12 became longer, with the WDR5-FKBP12 construct without any linker
being the most efficient. We conclude that RiPA closely recapitulates the E3 ligase selectivity
of WDR5 PROTACs and hypothesize that it can, in principle, predict E3 ligases capable of
forming ternary complexes with specific target proteins.

Another finding of our study is that the SCF (SKP1-CUL1 F-box protein)-type FBXL12 can
degrade artificial substrates in the context of the RiPA system. We consider this to be relevant

since the vast majority of PROTACs today use one of the two E3 ligases Cereblon or VHL.
Therefore, it was possible that these two E3 ligases are particularly or even exclusively suited
for the development of PRTOACs, e.g., due to their high potency or promiscuity. Since
FBXL12-mediated degradation of WDR5 is even more efficient than degradation by Cereblon,
and since FBXL12 can induce degradation of every target protein tested in our study, our data
suggest that FBXL12 may be a promising new E3 ligase for PROTAC development.

Limitations of the study

While we consider RiPA to be very helpful in guiding PROTAC development campaigns, its
results will not be fully generalizable to the characteristics of PROTACs, as it is conceivable
for this system to make false-negative and false-positive predictions. Thus, suitable target/E3
ligase pairs can fail in the RiPA setting because protein tags interfere with their interaction or
reduce the activity of the E3 ligases. However, our system allows easy testing of different
tagging approaches, and due to its simple workflow, it is easy to rapidly characterize the
activity of a new E3 ligase using multiple target proteins.

Conversely, it is also conceivable that an E3 ligase that can efficiently decrease the levels of
a particular target in the RiPA setting may be less suitable for PROTACs, since PROTACs
that mimic the steric interaction of the target/E3 pair may not be easily identified in the chemical
space. However, the RiPA system can certainly identify inappropriate E3 ligases that are not
in the same cellular compartment as the target or that cannot add a degradation-inducing
ubiquitin chain. Since RiPA also correctly predicted the differential suitability of VHL and
Cereblon to degrade WDR5, we anticipate that it will be very helpful in streamlining chemical
efforts in PROTAC development campaigns, both for PRTOACs based on established and
novel E3 ligases.

Materials and Methods

Key resource table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
WDR5 antibody (G-9)	Santa Cruz Biotechnology	Cat# sc-393080
Aurora A/AIK antibody	Cell Signaling Technology	Cat# 3092S
VHL antibody (VHL40)	Santa Cruz Biotechnology	Cat# sc-135657
Mouse monoclonal anti- CRBN (D8H3S)	Cell Signaling Technology	Cat# 71810S
Mouse monoclonal anti-vinculin (clone*hv)	Sigma-Aldrich	Cat# V9131
ECL-Anti-rabbit IgG Horseradish Peroxidase	GE Healthcare	Cat# NA934V
ECL-Anti-mouse IgG Horseradish Peroxidase	GE Healthcare	Cat# NA931V
Chemicals, peptides, and recombinant proteins		
DMEM, high glucose, pyruvate	Thermo Fisher Scientific	Cat# 41966052
Fetal Bovine Serum Advanced	Capricorn Scientific GmbH	Cat# FBS-11A
Penicillin-Streptomycin	Sigma-Aldrich	Cat# P4333
Opti-MEM I Reduced Serum Medium, no phenol red	Thermo Fisher Scientific	Cat# 11058021
HEPES	Sigma-Aldrich	Cat# H0887
Dimethylsulfoxide (DMSO)	Carl Roth	Cat# 4720
Polyethylenimine	Sigma-Aldrich	
Rapamycin	Selleckchem	Cat# S1039
Protease Inhibitor Cocktail	Sigma-Aldrich	Cat# P8340
Phosphatase Inhibitor Cocktail 2	Sigma-Aldrich	Cat# P5726
Phosphatase Inhibitor Cocktail 3	Sigma-Aldrich	Cat# P0044

Phusion High-Fidelity DNA Polymerase	Thermo Fisher Scientific	Cat# F530L
Phusion Plus DNA Polymerase	Thermo Fisher Scientific	Cat# F630L
Immobilon Western Chemiluminescent HRP Substrate	Merck Millipore	Cat#WBKLS0500
NanoGlo Endurazine Live Cell Substrate	Promega	Cat# N2571
AgeI-HF	New England BioLabs	Cat# R3552L
AscI	New England BioLabs	Cat# R0558L
BamHI-HF	New England BioLabs	Cat# R3136L
EcoRI-HF	New England BioLabs	Cat# R3101L
MluI-HF	New England BioLabs	Cat# R3198L
SpeI-HF	New England BioLabs	Cat# R3133L
XhoI	New England BioLabs	Cat# R0146L
Critical commercial assays		
Nano-Glo Luciferase Assay	Promega	Cat# N1120
GeneJET Gel Extraction Kit	Thermo Fisher Scientific	Cat# K0692
PureLink HiPure Plasmid Maxiprep Kit	Invitrogen	Cat# K210007
Experimental models: Cell lines		
HEK293T	ATCC	
Oligonucleotides		
Oligonucleotides used for PCR and cloning, see Table S2	Sigma-Aldrich	N/A
IDT G-blocks (double stranded linear DNA), see Table S2	Integrated DNA technologies	N/A
Recombinant DNA		
pRRL_puro	PMID: 25043018	N/A

pRRL_hygro	PMID: 25043018	N/A
pRRL_puro_WDR5-Luc-FKBP12	This Study	N/A
pRRL_puro_Luc-WDR5-FKBP12	This Study	N/A
pRRL_hygro_FRB-VHL	This Study	N/A
pRRL_hygro_VHL-FRB	This Study	N/A
pRRL_hygro_FRB	This Study	N/A
pRRL_hygro_CRBN-FRB	This Study	N/A
pRRL_puro_AURKA-Luc-FKBP12	This Study	N/A
pRRL_puro_WDR5-Luc-FKBP12(Kless)	This Study	N/A
pRRL_hygro_FBXL12-FRB	This Study	N/A
pRRL_puro_C-term_Luc-FKBP12_EV	This Study	N/A
pRRL_puro_C-term_Luc-FKBP12(Kless)_EV	This Study	N/A
pRRL_hygro_E3-FRB_EV	This Study	N/A
pRRL_hygro_FRB-E3_EV	This Study	N/A
pRRL_puro_Luc-WDR5-FKBP12_0xGSSG	This Study	N/A
pRRL_puro_Luc-WDR5-FKBP12_1xGSSG	This Study	N/A
pRRL_puro_Luc-WDR5-FKBP12_2xGSSG	This Study	N/A
pRRL_puro_Luc-WDR5-FKBP12_4xGSSG	This Study	N/A
pRRL_puro_Luc-WDR5-FKBP12_8xGSSG	This Study	N/A

pRRL_puro_Luc-FKBP12	This Study	N/A
pRRL_puro_Luc_V1-FKBP12	This Study	N/A
pRRL_puro_Luc_V2-FKBP12	This Study	N/A
pRRL_puro_Luc_K3-FKBP12	This Study	N/A
pRRL_puro_Luc_K6-FKBP12	This Study	N/A
pRRL_puro_Luc_K12-FKBP12	This Study	N/A
Software and algorithms		
GraphPad PRISM 10.1.1	GraphPad	https://www.graphpad.com
ImageJ 1.52	PMID: 22930834	https://imagej.nih.gov/ij/
PyMOL Molecular Graphics System	Schrödinger, LLC	https://pymol.org

Methods

Cell line and cell culture

315 Human HEK293T (female, embryo kidney) cells were cultured in DMEM medium (Thermo
316 Fischer Scientific) supplemented with 10% FBS ((Capricorn Scientific)) and 1%
317 penicillin/streptomycin (Sigma). Cells were grown at 37℃ under 5% CO_2. The cells were
318 routinely checked for mycoplasma contamination in a PCR-based assay and found negative.

Cloning of Target-FKBP12 constructs

WDR5-Luc-FKBP12, AURKA-Luc-FKBP12 and Luc-WDR5-FKBP12 vectors were cloned by
PCR amplification of the template vector for each fragment, WDR5 or AURKA, Luc and
FKBP12. A flexible linker (2xGSSG) between target-Luc and Luc-FKBP12 or Luc-target and
target-FKBP12 was incorporated in the primers (Sigma) used. The corresponding fragments
were digested with appropriate restriction enzymes and ligated into pRRL_puro. For the
WDR5-Luc-FKBP12(Kless) construct, first, a gBlock (IDT) containing multiple cloning site
(MCS), the mutated Luc (K55R, K77R, K80R, K91R, K125R, K126R and K138R), a linker
(2xGSSG) and the mutated FKBP12 (K18R, K35R, K36R, K45R, K48R, K53R, K74R and
K106R) was ordered. AscI/BamHI was then used to insert the gBlock into pRRL_puro,
resulting in the C-term_Luc-FKBP12-Kless entry vector (EV). Subsequently, the WDR5
fragment was PCR amplified and inserted into the EV using AgeI/SpeI restriction sites to get
WDR5-Luc-FKBP12(Kless). Similarly, the C-term_Luc-FKBP12 EV was produced by PCR
amplification of Luc and FKBP12 fragments, respectively, followed by an overlapping PCR to

generate a Luc-FKBP12 insert. The C-term_Luc-FKBP12 EV was then obtained by replacing

Luc-FKBP12(Kless) of the C-term_Luc-FKBP12-Kless EV.

For cloning of the Luc-WDR5-FKBP12 constructs with different GSSG linker lengths, the Luc-

WDR5 fragment was PCR amplified using pRRL_puro_Luc-WDR5-FKBP12 as a template.

The reverse primer binding to WDR5 contained the required linkers (0x or 1x or 2x or 4x or 8x

GSSG). The PCR product containing various linkers was then inserted into pRRL_puro_Luc-

WDR5-FKBP12 plasmid backbone after removing Luc-WDR5 using AgeI/MluI restriction sites.

Cloning of Luc-FKBP12 universal substrate constructs

The Luc-FKBP12 vector was cloned by PCR amplification of Luc and FKBP12 fragments and

subsequent ligation into the pRRL_puro backbone. To clone the lysine-rich Luc mutants, Luc^{V1}

and Luc^{V2}, Luc_ver1 (R13K, R45K, R114K, R154K, R168K) and Luc_ver2 (R13K, A16K,

Q22K, G28K, N35K, R45K, H88K, V104K, R114K, R154K, R168K) were ordered as gBlocks.

Luc was replaced in pRRL_puro_Luc-FKBP12 with Luc_ver1 and Luc_ver2 using AgeI/MluI

restriction sites. For Luc^{K3}, Luc^{K6} and Luc^{K12} cloning, two oligos (top and bottom) were

designed that contain the additional lysine residues and overhangs for AgeI/SpeI restriction

site. The oligos were hybridized and ligated into the pRRL_puro_C-term_Luc-FKBP12 EV

using AgeI/SpeI sites.

Cloning of E3-FRB constructs

VHL-FRB, FRB-VHL, CRBN-FRB, FBXL12-FRB and FRB vectors were cloned by PCR

amplification of VHL, CRBN, FBXL12 or FRB fragments from template plasmids or cDNA

(FBXL12). A flexible linker (2xGSSG) between the respective E3-ligase and FRB was

incorporated with the primers used. The corresponding fragments were digested with

appropriate restriction enzymes and ligated into pRRL_hygro backbone using AgeI/SpeI sites.

For E3-FRB EV and FRB-E3 EV, gBlocks containing FRB, a linker (2x GSSG) and a MCS

were ordered and inserted into pRRL_hygro using AscI/BamHI sites.

Rapamycin-induced proximity assays (RiPA)

HEK293T cells were plated at a density of 6.5×10^5 in 2 ml media per well in a 6-well plate. The

cells were allowed to attach and recover for at least 6 hours before transfection. For

transfection, two Eppendorf tubes with 140 µl OptiMEM (Thermo Fischer Scientific) each were

prepared with either 6 µl polyethylenimine (PEI; Sigma) or the appropriate amounts of

FKBP12/FRB plasmid pairs. A total of about 1.6 µg of plasmids in the required ratio (FKBP12:

FRB) was used per transfection. The contents were mixed well, centrifuged and incubated for

5 minutes at room temperature (RT) before the plasmid mixture was added to the PEI mixture.

After 20 minutes of incubation at RT, the plasmid-PEI mixture was added to the attached cells.

The cells were left in the incubator for at least 20 hours for protein expression. Next, the cells

were trypsinized, collected in media, centrifuged and resuspended in 3 ml of media (DMEM

for endpoint; assay medium, OptiMEM + 4 % FBS + 1 % P/S + 15 mM HEPES for kinetic).

For endpoint and kinetic luciferase measurement, 45 µl of the cell suspension was seeded

into two sets of replicates (for control and rapamycin) per transfection condition in a black 96-

well plate. For endpoint western blot (WB), the rest of the cell suspension was equally divided

and seeded into two wells of a 6-well plate per transfection condition while maintaining a final

volume of 2 ml with the media. The plates were left in incubator overnight. The next day,

rapamycin or control (DMSO) treatment for endpoint measurements was performed by adding

either rapamycin or a corresponding amount of DMSO to a final concentration of 10 or 100

nM. For the luciferase measurement rapamycin was pre-diluted in assay media. After the

treatment time, the cells were lysed with the Nano-Glo Luciferase Assay Reagent (Promega)

according to the manufacturer's protocol, and luminescence was measured using the Tecan

Spark Multiplate reader (Tecan) with an integration time of 1 second. Similarly, for WB the

cells were lysed in RIPA lysis buffer (50 mM HEPES pH 7.9, 140 mM NaCl, 1 mM EDTA, 1%

Triton X-100, 0.1% SDS, 0.1% sodium deoxycholate) containing protease and phosphatase

inhibitor (Sigma).

For the kinetic measurement, 2-3 hours prior to rapamycin addition, 15 µl of 5x Endurazine

(prediluted in assay medium) was added. After 2-3 hours of incubation, 15 µl of 50 nM

rapamycin (prediluted in assay medium) was added to the corresponding wells. Finally, kinetic

measurement with 15–45-minute intervals was initiated on the Tecan Spark Multiplate reader

at 37 °C/ 5 % CO_2 in the presence of a humidity cassette for 20 hours.

Western Blot

Cells treated with DMSO or rapamycin were lysed in RIPA lysis buffer for 30 minutes at 4°C

in rotary mixter. The cell debris was cleared from lysate by centrifugation and BCA assay was

done to measure protein concentration. Equal amounts of protein per sample were separated

by Bis-TRIS PAGE and transferred to PVDF membranes (Millipore). The membranes were

blocked with blocking solution (5% milk in TBS-T; 20 mM Tris HCl, pH 7.5, 150 mM NaCl, and

0.1% Tween 20), cut into pieces for different proteins and incubated with corresponding

primary antibodies at 4°C overnight. Next, the membranes were washed with TBS-T and

incubated with HRP-labelled secondary antibodies at RT for 1 hour. Visualization was done

with chemiluminescent HRP substrate (Millipore) in LAS4000 Mini (Fuji).

Protein structures

For modelling and visualization all crystal structures indicated below were prepared using the

Molecular Operating Environment (MOE) [43].

For the minimal luciferase, PDB structure 7SNS (resolution of 1.55 Å) was used. Only chain
A was prepared by modelling missing side chains, adding hydrogen atoms at pH 7 with
Protonate 3D [44] and renumbering the residues starting at 1 with the first resolved residue
(methionine). This residue was then removed as it was not present in the cloned constructs.
Buffer and water molecules were also removed from the structure.

The structure of FKBP12, FRB and rapamycin was generated by combining protein residues
from structure 3FAP [45] (1.85 Å) and rapamycin from structure 1NSG [45] (2.20 Å). The higher
occupancy conformation of R175 in chain B was retained, and the system was protonated at
pH 7. To match the cloned constructs, the two N-terminal residues (V and A) were removed,
and a C-terminal lysine was added. Water molecules were deleted from both structures.

Since no full-length structure of WDR5 was available, residues 32-334 from a crystal structure
(PDB: 2H14 [46], resolution 1.48 Å) were combined with residues 1-31 of the prediction AF-
P61964-F1 from the Alphafold2 structure database [47,48]. The predicted residues were
modelled with low to very low confidence but were considered suitable to indicate the
approximate distance between WDR5 and minimal luciferase when linked together. All water
molecules were removed from the system.

The proteins were locally minimized to a gradient of 0.1 kcal/mol*Å using the AMBER14:EHT
force field and tether restraints on all atoms (σ: 0.5 Å). In order to estimate the distances
between the individual proteins in the cloned constructs, GSSG linkers were modelled and
connected to the respective systems. The linkers were generated in a linear fashion to indicate
the maximum possible distance between the proteins and the relative positions of attachment
points. However, they do not make assumptions about possible or stable conformations. To
introduce more lysine residues to the minimal luciferase, potential mutation sites were selected
by visually inspecting the prepared structure.

All figures of protein structures were rendered using PyMOL 2.5.7 [49].

Acknowledgements

We thank André Kutschke for excellent technical support. This work was supported by grants

from the German Research Foundation (DFG, WO 2108/2-1 to E.W. and GRK 2243 to B. A.),

the European Research Council (ERC, PROTAC-PDAC to EW) and the Federal Ministry of

Education and Research (BMBF, DEGRON to E.W.).

Author Contribution

B.A. conducted and interpreted most experiments with help from all authors; K.S. helped in

kinetic experiments, M.D. and C.S. performed and interpreted computational modeling, E.W.

conceptualized the work, and E.W. and B.A. wrote the manuscript with input from all authors.

Declaration of interests

The University of Würzburg has filed a patent for the RiPA system described in the study

and E.W. and B. A. are listed as inventors.

Resource availability

Plasmids generated in this study will be made available on Addgene. Further information and

requests for resources and reagents should be directed to the lead contact, Elmar Wolf

(elmar.wolf@biochem.uni-kiel.de).

References

1. Oprea, T.I., Bologa, C.G., Brunak, S., Campbell, A., Gan, G.N., Gaulton, A., Gomez, S.M., Guha, R., Hersey, A., Holmes, J., et al. (2018). Unexplored therapeutic opportunities in the human genome. Nat Rev Drug Discov *17*, 317-332. 10.1038/nrd.2018.14.

2. Bondeson, D.P., Mares, A., Smith, I.E., Ko, E., Campos, S., Miah, A.H., Mulholland, K.E., Routly, N., Buckley, D.L., Gustafson, J.L., et al. (2015). Catalytic in vivo protein knockdown by small-molecule PROTACs. Nat Chem Biol *11*, 611-617. 10.1038/nchembio.1858.

3. Ishida, T., and Ciulli, A. (2021). E3 Ligase Ligands for PROTACs: How They Were Found and How to Discover New Ones. SLAS Discov *26*, 484-502. 10.1177/2472555220965528.

4. Winter, G.E., Buckley, D.L., Paulk, J., Roberts, J.M., Souza, A., Dhe-Paganon, S., and Bradner, J.E. (2015). DRUG DEVELOPMENT. Phthalimide conjugation as a strategy for in vivo target protein degradation. Science *348*, 1376-1381.

5. Farnaby, W., Koegl, M., McConnell, D.B., and Ciulli, A. (2021). Transforming targeted cancer therapy with PROTACs: A forward-looking perspective. Curr Opin Pharmacol *57*, 175-183. 10.1016/j.coph.2021.02.009.

6. Posternak, G., Tang, X., Maisonneuve, P., Jin, T., Lavoie, H., Daou, S., Orlicky, S., Goullet de Rugy, T., Caldwell, L., Chan, K., et al. (2020). Functional characterization of a PROTAC directed against BRAF mutant V600E. Nat Chem Biol *16*, 1170-1178. 10.1038/s41589-020-0609-7.

7. Chirnomas, D., Hornberger, K.R., and Crews, C.M. (2023). Protein degraders enter the clinic - a new approach to cancer therapy. Nat Rev Clin Oncol *20*, 265-278. 10.1038/s41571-023-00736-3.

8. Montoya, S., Bourcier, J., Noviski, M., Lu, H., Thompson, M.C., Chirino, A., Jahn, J., Sondhi, A.K., Gajewski, S., Tan, Y.S.M., et al. (2024). Kinase-impaired BTK mutations are susceptible to clinical-stage BTK and IKZF1/3 degrader NX-2127. Science *383*, eadi5798. 10.1126/science.adi5798.

9. Ward, J.A., Perez-Lopez, C., and Mayor-Ruiz, C. (2023). Biophysical and Computational Approaches to Study Ternary Complexes: A 'Cooperative Relationship' to Rationalize Targeted Protein Degradation. Chembiochem *24*, e202300163. 10.1002/cbic.202300163.

10. Donovan, K.A., Ferguson, F.M., Bushman, J.W., Eleuteri, N.A., Bhunia, D., Ryu, S., Tan, L., Shi, K., Yue, H., Liu, X., et al. (2020). Mapping the Degradable Kinome Provides a Resource for Expedited Degrader Development. Cell *183*, 1714-1731 e1710. 10.1016/j.cell.2020.10.038.

11. Adhikari, B., Bozilovic, J., Diebold, M., Schwarz, J.D., Hofstetter, J., Schroder, M., Wanior, M., Narain, A., Vogt, M., Dudvarski Stankovic, N., et al. (2020). PROTAC-mediated degradation reveals a non-catalytic function of AURORA-A kinase. Nat Chem Biol *16*, 1179-1188. 10.1038/s41589-020-00652-y.

12. Roy, M.J., Winkler, S., Hughes, S.J., Whitworth, C., Galant, M., Farnaby, W., Rumpel, K., and Ciulli, A. (2019). SPR-Measured Dissociation Kinetics of PROTAC Ternary Complexes Influence Target Degradation Rate. ACS Chem Biol *14*, 361-368. 10.1021/acschembio.9b00092.

489   13.   Gadd, M.S., Testa, A., Lucas, X., Chan, K.H., Chen, W., Lamont, D.J., Zengerle, M.,
490   and Ciulli, A. (2017). Structural basis of PROTAC cooperative recognition for selective
491   protein degradation. Nat Chem Biol *13*, 514-521. 10.1038/nchembio.2329.
492   14.   Itoh, Y., Ishikawa, M., Naito, M., and Hashimoto, Y. (2010). Protein knockdown using
493   methyl bestatin-ligand hybrid molecules: design and synthesis of inducers of
494   ubiquitination-mediated degradation of cellular retinoic acid-binding proteins. J Am
495   Chem Soc *132*, 5820-5826. 10.1021/ja100691p.
496   15.   Demizu, Y., Shibata, N., Hattori, T., Ohoka, N., Motoi, H., Misawa, T., Shoda, T., Naito,
497   M., and Kurihara, M. (2016). Development of BCR-ABL degradation inducers via the
498   conjugation of an imatinib derivative and a cIAP1 ligand. Bioorg Med Chem Lett *26*,
499   4865-4869. 10.1016/j.bmcl.2016.09.041.
500   16.   Zhang, X., Luukkonen, L.M., Eissler, C.L., Crowley, V.M., Yamashita, Y., Schafroth,
501   M.A., Kikuchi, S., Weinstein, D.S., Symons, K.T., Nordin, B.E., et al. (2021). DCAF11
502   Supports Targeted Protein Degradation by Electrophilic Proteolysis-Targeting
503   Chimeras. J Am Chem Soc *143*, 5141-5149. 10.1021/jacs.1c00990.
504   17.   Han, T., Goralski, M., Gaskill, N., Capota, E., Kim, J., Ting, T.C., Xie, Y., Williams,
505   N.S., and Nijhawan, D. (2017). Anticancer sulfonamides target splicing by inducing
506   RBM39   degradation   via   recruitment   to   DCAF15.   Science   *356*.
507   10.1126/science.aal3755.
508   18.   Du, X., Volkov, O.A., Czerwinski, R.M., Tan, H., Huerta, C., Morton, E.R., Rizzi, J.P.,
509   Wehn, P.M., Xu, R., Nijhawan, D., and Wallace, E.M. (2019). Structural Basis and
510   Kinetic Pathway of RBM39 Recruitment to DCAF15 by a Sulfonamide Molecular Glue
511   E7820. Structure *27*, 1625-1633 e1623. 10.1016/j.str.2019.10.005.
512   19.   Zhang, X., Crowley, V.M., Wucherpfennig, T.G., Dix, M.M., and Cravatt, B.F. (2019).
513   Electrophilic PROTACs that degrade nuclear proteins by engaging DCAF16. Nat
514   Chem Biol *15*, 737-746. 10.1038/s41589-019-0279-5.
515   20.   Ward, C.C., Kleinman, J.I., Brittain, S.M., Lee, P.S., Chung, C.Y.S., Kim, K., Petri, Y.,
516   Thomas, J.R., Tallarico, J.A., McKenna, J.M., et al. (2019). Covalent Ligand Screening
517   Uncovers a RNF4 E3 Ligase Recruiter for Targeted Protein Degradation Applications.
518   ACS Chem Biol *14*, 2430-2440. 10.1021/acschembio.8b01083.
519   21.   Spradlin, J.N., Hu, X., Ward, C.C., Brittain, S.M., Jones, M.D., Ou, L., To, M.,
520   Proudfoot, A., Ornelas, E., Woldegiorgis, M., et al. (2019). Harnessing the anti-cancer
521   natural product nimbolide for targeted protein degradation. Nat Chem Biol *15*, 747-
522   755. 10.1038/s41589-019-0304-8.
523   22.   Luo, M., Spradlin, J.N., Boike, L., Tong, B., Brittain, S.M., McKenna, J.M., Tallarico,
524   J.A., Schirle, M., Maimone, T.J., and Nomura, D.K. (2021). Chemoproteomics-enabled
525   discovery of covalent RNF114-based degraders that mimic natural product function.
526   Cell Chem Biol *28*, 559-566 e515. 10.1016/j.chembiol.2021.01.005.
527   23.   Ohoka, N., Tsuji, G., Shoda, T., Fujisato, T., Kurihara, M., Demizu, Y., and Naito, M.
528   (2019). Development of Small Molecule Chimeras That Recruit AhR E3 Ligase to
529   Target Proteins. ACS Chem Biol *14*, 2822-2832. 10.1021/acschembio.9b00704.
530   24.   Henning, N.J., Manford, A.G., Spradlin, J.N., Brittain, S.M., Zhang, E., McKenna, J.M.,
531   Tallarico, J.A., Schirle, M., Rape, M., and Nomura, D.K. (2022). Discovery of a
532   Covalent FEM1B Recruiter for Targeted Protein Degradation Applications. J Am Chem
533   Soc *144*, 701-708. 10.1021/jacs.1c03980.
534   25.   Konopleva, M., Tsao, T., Ruvolo, P., Stiouf, I., Estrov, Z., Leysath, C.E., Zhao, S.,
535   Harris, D., Chang, S., Jackson, C.E., et al. (2002). Novel triterpenoid CDDO-Me is a

potent inducer of apoptosis and differentiation in acute myelogenous leukemia. Blood
99, 326-335. 10.1182/blood.v99.1.326.

26. Wei, J., Meng, F., Park, K.S., Yim, H., Velez, J., Kumar, P., Wang, L., Xie, L., Chen,
H., Shen, Y., et al. (2021). Harnessing the E3 Ligase KEAP1 for Targeted Protein
Degradation. J Am Chem Soc *143*, 15073-15083. 10.1021/jacs.1c04841.

27. Pei, J., Xiao, Y., Liu, X., Hu, W., Sobh, A., Yuan, Y., Zhou, S., Hua, N., Mackintosh,
S.G., Zhang, X., et al. (2023). Piperlongumine conjugates induce targeted protein
degradation. Cell Chem Biol *30*, 203-213 e217. 10.1016/j.chembiol.2023.01.004.

28. Choi, J., Chen, J., Schreiber, S.L., and Clardy, J. (1996). Structure of the FKBP12-
rapamycin complex interacting with the binding domain of human FRAP. Science *273*,
239-242. 10.1126/science.273.5272.239.

29. Dolle, A., Adhikari, B., Kramer, A., Weckesser, J., Berner, N., Berger, L.M., Diebold,
M., Szewczyk, M.M., Barsyte-Lovejoy, D., Arrowsmith, C.H., et al. (2021). Design,
Synthesis, and Evaluation of WD-Repeat-Containing Protein 5 (WDR5) Degraders. J
Med Chem *64*, 10682-10710. 10.1021/acs.jmedchem.1c00146.

30. Yu, X., Li, D., Kottur, J., Kim, H.S., Herring, L.E., Yu, Y., Xie, L., Hu, X., Chen, X., Cai,
L., et al. (2023). Discovery of Potent and Selective WDR5 Proteolysis Targeting
Chimeras as Potential Therapeutics for Pancreatic Cancer. J Med Chem *66*, 16168-
16186. 10.1021/acs.jmedchem.3c01521.

31. Yu, X., Li, D., Kottur, J., Shen, Y., Kim, H.S., Park, K.S., Tsai, Y.H., Gong, W., Wang,
J., Suzuki, K., et al. (2021). A selective WDR5 degrader inhibits acute myeloid
leukemia in patient-derived mouse models. Sci Transl Med *13*, eabj1578.
10.1126/scitranslmed.abj1578.

32. Hall, M.P., Unch, J., Binkowski, B.F., Valley, M.P., Butler, B.L., Wood, M.G., Otto, P.,
Zimmerman, K., Vidugiris, G., Machleidt, T., et al. (2012). Engineered luciferase
reporter from a deep sea shrimp utilizing a novel imidazopyrazinone substrate. ACS
Chem Biol *7*, 1848-1857. 10.1021/cb3002478.

33. Wang, R., Ascanelli, C., Abdelbaki, A., Fung, A., Rasmusson, T., Michaelides, I.,
Roberts, K., and Lindon, C. (2021). Selective targeting of non-centrosomal AURKA
functions through use of a targeted protein degradation tool. Commun Biol *4*, 640.
10.1038/s42003-021-02158-2.

34. Rishfi, M., Krols, S., Martens, F., Bekaert, S.L., Sanders, E., Eggermont, A., De Vloed,
F., Goulding, J.R., Risseeuw, M., Molenaar, J., et al. (2023). Targeted AURKA
degradation: Towards new therapeutic agents for neuroblastoma. Eur J Med Chem
247, 115033. 10.1016/j.ejmech.2022.115033.

35. Liu, F., Wang, X., Duan, J., Hou, Z., Wu, Z., Liu, L., Lei, H., Huang, D., Ren, Y., Wang,
Y., et al. (2022). A Temporal PROTAC Cocktail-Mediated Sequential Degradation of
AURKA Abrogates Acute Myeloid Leukemia Stem Cells. Adv Sci (Weinh) *9*, e2104823.
10.1002/advs.202104823.

36. Poirson, J., Dhillon, A., Cho, H., Lam, M.H.Y., Alerasool, N., Lacoste, J., Mizan, L., and
Taipale, M. (2022). Proteome-Scale Induced Proximity Screens Reveal Highly Potent
Protein Degraders and Stabilizers. . bioRxiv. 10.1101/2022.08.15.503206.

37. Schapira, M., Calabrese, M.F., Bullock, A.N., and Crews, C.M. (2019). Targeted
protein degradation: expanding the toolbox. Nat Rev Drug Discov *18*, 949-963.
10.1038/s41573-019-0047-y.

38. Bond, M.J., and Crews, C.M. (2021). Proteolysis targeting chimeras (PROTACs) come
of age: entering the third decade of targeted protein degradation. RSC Chem Biol *2*,
725-742. 10.1039/d1cb00011j.

39.   Zhang, X., Zhao, T., Sun, M., Li, P., Lai, M., Xie, L., Chen, J., Ding, J., Xie, H., Zhou,
J., and Zhang, H. (2023). Design, synthesis and biological evaluation of KRAS(G12C)-
PROTACs. Bioorg Med Chem *78*, 117153. 10.1016/j.bmc.2023.117153.

40.   Yang, N., Fan, Z., Sun, S., Hu, X., Mao, Y., Jia, C., Cai, X., Xu, T., Li, B., Li, Y., et al.
(2023). Discovery of highly potent and selective KRAS(G12C) degraders by VHL-
recruiting PROTACs for the treatment of tumors with KRAS(G12C)-Mutation. Eur J
Med Chem *261*, 115857. 10.1016/j.ejmech.2023.115857.

41.   Steinebach, C., Ng, Y.L.D., Sosic, I., Lee, C.S., Chen, S., Lindner, S., Vu, L.P., Bricelj,
A., Haschemi, R., Monschke, M., et al. (2020). Systematic exploration of different E3
ubiquitin ligases: an approach towards potent and selective CDK6 degraders. Chem
Sci *11*, 3474-3486. 10.1039/d0sc00167h.

42.   Schwalm, M.P., Kramer, A., Dolle, A., Weckesser, J., Yu, X., Jin, J., Saxena, K., and
Knapp, S. (2023). Tracking the PROTAC degradation pathway in living cells highlights
the importance of ternary complex measurement for PROTAC optimization. Cell Chem
Biol *30*, 753-765 e758. 10.1016/j.chembiol.2023.06.002.

43.   ChemicalComputingGroup ULC, 1010 Sherbrooke St. West, Suite #910, Montreal,
QC, Canada, H3A 2R7 (2022). Molecular Operating Environment (MOE) (Chemical
Computing Group).

44.   Labute, P. (2009). Protonate3D: assignment of ionization states and hydrogen
coordinates to macromolecular structures. Proteins *75*, 187-205. 10.1002/prot.22234.

45.   Liang, J., Choi, J., and Clardy, J. (1999). Refined structure of the FKBP12-rapamycin-
FRB ternary complex at 2.2 A resolution. Acta Crystallogr D Biol Crystallogr *55*, 736-
744. 10.1107/s0907444998014747.

46.   Couture, J.F., Collazo, E., and Trievel, R.C. (2006). Molecular recognition of histone
H3 by the WD40 protein WDR5. Nat Struct Mol Biol *13*, 698-703. 10.1038/nsmb1116.

47.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.
10.1038/s41586-021-03819-2.

48.   Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan,
D., Stroe, O., Wood, G., Laydon, A., et al. (2022). AlphaFold Protein Structure
Database: massively expanding the structural coverage of protein-sequence space
with high-accuracy models. Nucleic Acids Res *50*, D439-D444. 10.1093/nar/gkab1061.

49.   The PyMOL Molecular Graphics System, Version 2.5. Schrödinger, LLC.

Figures

Figure 1. Rapamycin-induced proximity assay (RiPA) induces quantifiable degradation

of target proteins

(A) Schematic illustration of scenarios where PROTAC could not induce degradation of a

target protein.

(B) Scheme of target protein (I) and E3-ligase or control (II) constructs used in RiPA. The

linker indicated is 2xGSSG in all constructs unless stated otherwise.

(C) Schematic describing the RiPA experimental protocol.

(D) Immunoblot of WDR5 and VHL. HEK293 cells were co-transfected with WDR5-Luc-

FKBP12 and VHL-FRB or FRB in the indicated ratio and treated with 100 nM rapamycin

or vehicle for 6 hr after ~24 hr of expression. Vinculin was used as a loading control (as in

all other immunoblotting experiments).

(E) WDR5 levels based on luciferase measurements. HEK293 cells were co-transfected with

WDR5-Luc-FKBP12 or Luc-WDR5-FKBP12 and VHL-FRB or FRB constructs in the

indicated ratio, expressed for ~24 hr, and treated with rapamycin overnight. Bars represent

mean ± s.d. of n=3 replicates.

(F) Immunoblot of WDR5 and VHL. HEK293 cells were co-transfected with a combination of

WDR5-Luc-FKBP12 or Luc-WDR5-FKBP12 and VHL-FRB or FRB-VHL or FRB in the ratio

of 1:10, expressed for ~24 hr and treated with rapamycin overnight.

Figure 1 - figure supplement 1. Rapamycin-induced proximity assay (RiPA) induces

quantifiable degradation of target proteins

(A) Schematic illustration of rapamycin-induced dimerization of FRB and FKBP12 and

structure of rapamycin.

(B) Structure of WDR5-Luc-FKBP12 and Luc-WDR5-FKBP12 fusion proteins. A flexible linker,

2xGSSG is present between each component.

Figure 2. RiPA correctly predicts suitability of E3 ligases for WDR5 PROTACs

(A) Immunoblot of WDR5, VHL, and CRBN. HEK293 cells were transfected with WDR5-Luc-FKBP12 and VHL-FRB or CRBN-FRB or FRB in a 1:10 ratio, expressed for 24 hr, and treated with 10 nM rapamycin for 6 hr.

(B) WDR5 levels based on luciferase measurement. Luminescence was measured in HEK293 cells as described in (**A**) after 10 nM rapamycin treatment at specified time points. Data represent mean ± s.d. of n=3 replicates.

(C) Immunoblot of AURKA, VHL and CRBN. HEK293 cells were transfected with AURKA-Luc-FKBP12 and VHL-FRB or CRBN-FRB or FRB in a 1:10 ratio and treated with 10 nM rapamycin for 6 hr.

(D) AURKA levels based on luciferase measurement. Luminescence was measured in HEK293 cells as described in (**C**) after 10 nM rapamycin treatment at indicated time points. Data represent mean ± s.d. of n=3 replicates.

(E) Structure of FKBP12 and luciferase. Molecular surface representation of FKBP12 (top) and luciferase (bottom) showing lysine residues on their surface. The lysine residues are labelled and two sides for each protein are shown.

(F) WDR5 levels based on luciferase measurement. HEK293 cells were co-transfected with either WDR5-Luc-FKBP12 (WT) or WDR5-Luc-FKBP12 construct where all lysine

residues on Luc and FKBP12 were mutated to arginine (K-less) and FRB, expressed for
~24 hr and luminescence measured. Bars represent mean ± s.d. of n=3 replicates.
(G) Immunoblot of WDR5, VHL, and CRBN. HEK293 cells expressing WDR5-Luc-FKBP12 (K-
less) and VHL-FRB or CRBN-FRB or FRB were treated with rapamycin as in (**A**).
(H) WDR5 levels based on luciferase measurement. Luminescence was measured in HEK293
cells as described in (**G**) after 10 nM rapamycin treatment at specified time points. Data
represent mean ± s.d. of n=3 replicates.

Figure 3. RiPA can identify degradative E3 ligases not previously used for PROTACs

(A) Immunoblot of WDR5. HEK293 cells were transfected with WDR5-Luc-FKBP12 (K-less) and FBXL12-FRB or FRB in a ratio of 1:10 and treated with 10 nM rapamycin for 8 hr.

(B) WDR5 levels based on luciferase measurement. Luminescence of WDR5-Luc-FKBP12
(K-less) in the same cells as in (**A**). Bars represent mean ± s.d. of n=3 replicates.

(C) WDR5 levels based on luciferase measurement. HEK293 cells were transfected with
WDR5-Luc-FKBP12 (K-less) and FBXL12-FRB or VHL-FRB or FRB in a ratio of 1:1000
and treated with 10 nM rapamycin for the indicated time point. Data represent mean ± s.d.
of n=3 replicates.

(D) Model of Luc-WDR5-FKBP12 constructs. Structure of Luc-WDR5-FKBP12 with indicated
linkers between WDR5 and FKBP12 bound to rapamycin and FRB. The linker between
Luc-WDR5 is always 2xGSSG.

(E) Immunoblot of WDR5. HEK293 cells were transfected with Luc-WDR5-FKBP12 containing
indicated linker length and FBXL12-FRB or FRB in the ratio of 1:100, expressed for ~24
hr, and treated with 10 nM rapamycin for 6 hr.

(F) WDR5 levels based on luciferase measurement. Luminescence of Luc-WDR5-FKBP12
constructs as in cells from (**E**) and treated with 10 nM rapamycin for 8 hr. Bars represent
mean ± s.d. of n=2 replicates.

(G) WDR5 levels based on kinetic luciferase measurement. HEK293 cells were transfected
with WDR5-Luc-FKBP12 (K-less) and FBXL12-FRB or FRB in a ratio of 1:100, expressed
for ~24 hr, treated with 10 nM rapamycin, and luminescence measured for 19 hr. Data
represent mean ± s.d. of n=3 replicates.

Figure 3 - figure supplement 1. RiPA can identify degradative E3 ligases not previously
used for PROTACs

(A) WDR5 levels based on luciferase measurement. HEK293 cells were transfected with Luc-
WDR5-FKBP12 or WDR5-Luc-FKBP12 and FBXL12-FRB or FRB in a ratio of 1:10,
expressed for ~24 hr, and treated with 10 nM rapamycin for 9 hr. Bars represent mean ±
s.d. from n=3 replicates.

Figure 4. Identification of degradation-inducing E3 ligases by designing a universal substrate

(A) Model of lysine-rich luciferase. Structure of mutated luciferase with 5 additional (Luc^{V1}) and 12 additional (Luc^{V2}) lysines as compared to 7 lysine residues of wild-type luciferase. The lysine residues from WT (black), Luc^{V1} (red), and Luc^{V2} (orange; red as in V1) are labeled.

(B) Scheme of wild-type luciferase (WT) or lysine-rich luciferase (V1, V2, K3, K6, and K12) containing constructs.

(C) Luciferase measurement. HEK293 cells were co-transfected with Luc-FKBP12 constructs as shown in (**B**) and FRB, expressed for ~24 hr, and luminescence was compared. Bars represent mean ± s.d. of n=2 replicates.

(D) Luciferase measurement. HEK293 cells were transfected with the indicated versions of Luc-FKBP12 and FBXL12-FRB or FRB in a ratio of 1:100, expressed for ~24 hr, and treated with 10 nM rapamycin for 8 hr. Bars represent mean ± s.d. of n=2 replicates.

(E) Kinetic luminescence measurement. HEK293 cells expressing constructs as described in

(**D**) were treated with 10 nM rapamycin or vehicle and luminescence was monitored for 22

hr. Bars represent mean ± s.d. of n=2 replicates.

Additional Files

Supplementary Table S1. Amino acid sequences. The table contains all the amino acid
sequences of the constructs used in this study.

Supplementary Table S2. Oligonucleotide sequences. The table contains all
oligonucleotide sequences, primers and gBlocks used in this study.