Disruption of WSB2-mediated NOXA Degradation Induces Synthetic Lethality to Anti-apoptotic BCL-2 Family Protein Inhibitors

Dongyue Jiao[1,8], Kun Chang[2,3,8], Yingji Chen[1], Jiamin Jin[1], Ren Mo[4], Yucong Zhang[5], Kun Gao[6,7], Yaoting Xu[8,*], Lixin Wang[5,*], and Chenji Wang[1,*]

[1]*State Key Lab of Genetic Engineering, Shanghai Stomatological Hospital & School of Stomatology, MOE Engineering Research Center of Gene Technology, Shanghai Engineering Research Center of Industrial Microorganisms, School of Life Sciences, Fudan University, Shanghai 200438, P.R. China*

[2]*Department of Urology, Fudan University Shanghai Cancer Center, Shanghai, 200032, China;*

[3]*Department of Oncology, Shanghai Medical College, Fudan University, Shanghai 200032, China;*

[4]*Department of Urology, Inner Mongolia Urological Institute, Inner Mongolia Autonomous Region people's Hospital, Hohhot 010017, Inner Mongolia, China*

[5]*Department of Clinical Laboratory, Shanghai First Maternity and Infant Hospital, School of Medicine, Tongji University, Shanghai 200092, China*

[6]*Department of Vascular Surgery, Zhongshan Hospital, Fudan University, Shanghai, China.*

[7]*Shanghai Key Laboratory of Maternal and Fetal Medicine, Shanghai First Maternity and Infant Hospital, Shanghai 200092, China*

[8]*Department of Urology, Shanghai Fourth People's Hospital Affiliated to Tongji University School of Medicine, Shanghai 200434, China.*

*Correspondence: chenjiwang@fudan.edu.cn (C.W); 2000019@tongji.edu.cn (Y.X).

Abstract

Anti-apoptotic BCL-2 family proteins are frequently overexpressed in various cancers, contributing to the initiation and development of cancer, as well as intrinsic or acquired resistance to therapy. Although BCL-2 family protein inhibitors, such as Venetoclax, have demonstrated efficacy in hematological neoplasms, their effectiveness as single agents in solid tumors is limited. Identifying alternative molecular targets that can overcome intrinsic resistance to BCL-2 family protein inhibitors is of great clinical importance. Here, we present evidence of strong synthetic lethal interactions between WSB2, a relatively unexplored substrate-binding receptor of the Cullin 5-RBX2-Elongin B/C (CRL5) E3 ubiquitin ligase complex, and multiple anti-apoptotic BCL-2 family proteins. Mechanistically, an assembled CRL5^{WSB2} E3 ubiquitin ligase complex targets NOXA, a pro-apoptotic BCL-2 family protein, for degradation via the ubiquitin-proteasomal pathway. Ablation of WSB2 leads to a remarkable accumulation of NOXA proteins in cultured cell lines and knockout mouse organs. While WSB2 deficiency alone has a minimal effect on spontaneous apoptosis, it renders cancer cells more susceptible to apoptosis when anti-apoptotic BCL-2 family proteins are genetically depleted or pharmacologically inhibited. These findings establish WSB2 as a critical regulator of mitochondrial apoptosis and highlight the dysregulation of the WSB2-NOXA regulatory axis as a contributing factor to apoptosis resistance in cancer cells. Synergistically targeting WSB2 and anti-apoptotic BCL-2 family proteins holds promising clinical potential in the treatment of human cancers.

Keywords:

Apoptosis, BCL-2, WSB2, Cullin, Ubiquitination, Venetoclax

Introduction

One of the major hallmarks of human cancers is their inherent or developed resistance to

apoptosis, a type of programmed cell death[1]. This evasive mechanism contributes significantly

to both tumor initiation and progression, as well as the resistance observed in many cancer

treatments. Importantly, most anticancer therapies available—including chemotherapy,

radiotherapy, molecularly targeted therapy, and immunotherapy—work primarily by initiating

cell death pathways like apoptosis in cancer cells[2].

Apoptosis is primarily regulated at the mitochondrial level by the B-cell lymphoma protein-2

(BCL-2) family of proteins. This family is bifurcated into two primary categories: anti-apoptotic

proteins and pro-apoptotic proteins. Anti-apoptotic proteins, such as BCL-2, BCL-XL, BCL-W,

and MCL-1, harbor between one to four BH domains and play essential roles in maintaining

outer mitochondrial membrane integrity by inhibiting their pro-apoptotic counterparts.

Pro-apoptotic proteins further segregate into multidomain effector proteins (e.g., BAK, BAX,

and BOK) that contain multiple BH domains, and an array of "BH3-only" proteins, named for

their single conserved BH3 domain (e.g., BIM, PUMA, NOXA, BAD, and BID). Various cell

death stimuli can activate or induce these proapoptotic entities, driving mitochondrial outer

membrane permeabilization followed by apoptosis. The anti-apoptotic members of the BCL-2

family have restrainer BH3-binding grooves that help to regulate the activity of "BH3-only" and

multidomain effector proteins[3, 4]. Notably, tumors often acquire resistance to apoptosis through

overexpression of anti-apoptotic proteins or downregulation/mutation of pro-apoptotic ones,

underscoring the need for therapeutic strategies targeted at this critical cellular balance[5].

A unique class of small molecule BH3-mimetics has been developed, which specifically

target and block the anti-apoptotic BCL-2 family proteins. Among these inhibitors, ABT-199

(Venetoclax) has been approved for clinical use against BCL-2 dependent tumors, especially

in patients with small lymphocytic lymphoma (SLL) or chronic lymphocytic leukemia (CLL) who

have undergone at least one prior therapy[5]. Furthermore, Venetoclax can be combined with

other drugs to treat acute myeloid leukemia (AML)[6, 7]. Despite major breakthroughs in

hematological neoplasms, the effectiveness of these drugs in solid tumors is limited, possibly

due to their reduced dependence on BCL-2 compared to hematological neoplasms[8].

Searching for new actionable targets to sensitize solid tumor cells to BCL-2 inhibitor-mediated

cytotoxicity is a promising avenue for investigation.

Cullin 5 (CUL5), a member of the cullin-RING ubiquitin ligase family, is a core component of

multisubunit E3 ubiquitin-protein ligase complexes which catalyze the conjugation of ubiquitin

moieties onto specific protein substrates. CUL5 interacts with RBX2, Elongin B/C, and a

SOCS Box-containing substrate-binding receptor to assembly a CRL5 E3 ubiquitin ligase

protein complex[9]. WD Repeat and SOCS Box Containing 2 (WSB2), has been classified as a

substate receptor for the CRL5 E3 ubiquitin ligase complex[10]. WSB2 has been shown to be

overexpressed in several cancers (lung cancer, breast cancer, and melanoma) and promote

malignant biological behavior, such as proliferation, cycle progression, and migration[11-13].

However, a lack of reported physiological substrates for the CRL5[WSB2] complex impedes

recognizing WSB2 as a potentially targetable oncoprotein for therapeutic interventions.

By analyzing the gene co-dependency dataset deposited in the DepMap Portal

(https://depmap.org/), we have unveiled a potential functional interplay between WSB2 and

multiple BCL-2 family proteins. Biochemical analyses demonstrate that WSB2 acts as a

substrate-binding receptor of CRL5 E3 to promote NOXA turnover. A series of functional

analyses were performed in cell lines, xenograft tumor models, and knockout mice models to

explore the biological significance and clinical relevance of WSB2-mediated regulation of

mitochondrial apoptosis. These findings highlight the clinical potential of targeting WSB2 for

synthetic lethality with BCL-2 family inhibitor treatment in human cancers.

Results

Multiple BCL-2 family proteins were identified as interactors of WSB2

To elucidate the unidentified molecular functions of WSB2, we analyzed the genetic

co-dependency between WSB2 and other proteins using Broad's 21Q2 DepMap dataset[14].

This dataset, derived from large-scale loss-of-function sgRNA screens for vulnerabilities in 990

cancer cell lines, allows the identification of genes with similar functions or pathways[15, 16].

Gene ontology analysis of the top 100 co-dependent genes of WSB2 revealed a significant

enrichment in multiple apoptosis-related processes (**Supplementary Figure 1A**,

Supplementary Table. 1, 2). Among the top four correlated genes, WSB2 showed a positive

correlation with anti-apoptotic BCL2L2 (BCL-W) and MCL-1, but a negative correlation with

pro-apoptotic BAX and PMAIP1 (NOXA) (**Figure 1A**). Notably, WSB2, along with BCL-2,

BCL-W, BAX, MCL-1, NOXA, and BAK1, formed a co-essential module[17] that also includes

UBE2J2/MARCH5, an E2-E3 ligase complex responsible for MCL1/NOXA turnover (**Figure**

1B)[18-20]. Through utilizing a web server, DepLink[21], to identify genetic and pharmacological

perturbations that produce similar impacts on cell viability, as determined by DepMap and drug

sensitivity datasets obtained from two high-throughput pharmacological screenings[22, 23], we

observed that WSB2 knockout exhibited the highest correlations with two BH3-mimetics

(ABT-737, ABT-263), out of hundreds of drugs tested (**Supplementary Figure 1B, C,**

Supplementary Table 3). These molecular links prompted us to investigate whether WSB2

has any impact on mitochondrial apoptosis through the regulation of BCL-2 family proteins. To

test this hypothesis, we initially examined the potential interaction between WSB2 and BCL-2

family proteins. Exogenous co-IP assay results indicated that among the eight tested BCL-2

family proteins, WSB2 interacted with MCL-1, NOXA, BAD, BCL-W, and BCL-2, but not with

BAX, BCL-W, and BAK (**Figure 1C**). These interactions were further confirmed by

semi-endogenous or endogenous co-IP assays (**Figure 1D, F-J**). In comparison, WSB1, a

closely related paralog of WSB2, did not interact with these BCL-2 family proteins (**Figure 1E**).

Considering that BCL-2 family proteins are located on the outer mitochondrial membrane,

we investigated whether WSB2 was also located on the mitochondria. Immunofluorescent

analysis revealed that WSB2 is localized throughout the cytoplasm, with a fraction of it

colocalizing with the mitochondrial marker HSP60 (**Figure 1K**). We further separated the

nuclear, mitochondrial, and cytoplasmic fractions of HeLa cells using density-gradient

centrifugation methods. WSB2 predominantly localizes in the cytoplasm, with a moderate

fraction in the mitochondria, while its presence in the nucleus is minor (**Figure. 1L**). To

determine the submitochondrial localization of WSB2, we purified mitochondria from HeLa

cells and performed Protease K digestion experiments with different mitochondrial

preparations. Cleavage by Protease K only occurred in intact mitochondria and targeted outer

membrane proteins exposed to the cytosol, such as TOM70. Swelling the mitochondria with a

hypotonic buffer disrupted the outer mitochondrial membrane but left the inner membrane

intact, resulting in the cleavage of intermembrane space proteins like SMAC. Lysis with Triton

X-100 caused the cleavage of all mitochondrial proteins including matrix proteins like HSP60.

Similar to TOM70, WSB2 was cleaved by Protease K digestion in intact mitochondria

preparation (**Figure 1M**), supporting its classification as a mitochondrial outer membrane

protein.

Collectively, these data indicate that WSB2 specifically interacts with a subset of BCL-2

family proteins on the outer membrane of the mitochondria.

CRL5^{WSB2} E3 ubiquitin ligase complex mediates the ubiquitin-proteasomal degradation

of NOXA

WSB2 was co-purified with CRL5 complex components (RBX2, CUL5, ELOB, and ELOC),

confirming that WSB2 is a potential CRL5 adaptor (**Figure 1D**). We next investigated whether

WSB2 controls the protein stability of its interacting BCL-2 family proteins. Overexpression of

WSB2 markedly reduced the levels of co-expressed NOXA in a dose-dependent manner,

while other examined BCL-2 family proteins showed minimal or no impact (**Supplementary**

Figure 1D). Furthermore, depletion of WSB2 through shRNA-mediated knockdown (KD) or

CRISPR/Cas9-mediated knockout (KO) in prostate cancer C4-2B cells or liver cancer Huh-7

cells led to a marked increase in the steady-state levels of endogenous NOXA, without

affecting other BCL-2 family proteins examined (**Figure 2A-C, Supplementary Figure 2A, B**).

Therefore, our main focus in this study was to investigate the impact of WSB2 on the stability

of NOXA. We demonstrated that the proteasome inhibitor MG132 completely reversed the

effect of WSB2 on NOXA protein levels, while WSB1 had no effect on NOXA expression levels

(**Figure 2D**). WSB2 contains a C-terminal SOCS box consisting of the BC box and the Cullin 5

(CUL5) box, which interact with Elongin B/C and CUL5, respectively[10]. To investigate the roles

of these domains, we generated two WSB2 mutants in which the BC box or CUL5 box was

deleted, respectively (**Supplementary Figure 1E**). Strikingly, WSB2-BCM or CULM mutant

failed to decrease NOXA protein levels (**Figure 2E**). Furthermore, reintroduction of the WSB2

mutant (BCM, CULM) into WSB2-KO cells was unable to reverse the accumulation of NOXA

protein caused by WSB2 deficiency (**Figure 2F**). Depletion of CRL5 complex components

(RBX2, CUL5, ELOB, or ELOC) by siRNAs in C4-2B cells also resulted in a marked increase

in NOXA protein levels (**Figure 2G**), indicating that the assembly of the CRL5^{WSB2} complex is

necessary for NOXA degradation. The half-life of NOXA was notably prolonged in WSB2-KO

cells (**Figure 2H**, **I**). Additionally, the mRNA levels of NOXA were even reduced in WSB2-KO

cells compared to parental cells, likely to counteract the accumulation of NOXA protein

(**Figure 2J**). We also found that WSB2-WT, but not the BCM or CULM mutant, could promote

polyubiquitination of NOXA (**Figure 2K**, **L**). By employing linkage-specific K48/K63-Ub

mutants, we found that ubiquitinated NOXA primarily consisted of K48-Ub linkages, which is

consistent with the expectation that K48-Ub linkage serves as the canonical signal for

proteasomal degradation (**Figure 2M**). Conversely, the levels of endogenous ubiquitinated

NOXA were diminished in WSB2-KO cells compared to parental cells (**Figure 2N**).

Collectively, these data indicate that the CRL5^{WSB2} complex mediates the

ubiquitin-proteasomal degradation of NOXA.

The C-terminal region of NOXA is crucial for WSB2-mediated NOXA degradation

Along with the BC-box and CUL5 box, WSB2 contains five WD repeats that typically serve as

protein-protein interaction modules. We generated deletion mutants to determine the region

responsible for binding to NOXA, and surprisingly found that WSB's SOCS box, rather than the

WD repeats, is required for its interaction with NOXA (**Figure 3A, B**). Deletion of this region

abolished WSB2-mediated degradation and ubiquitination of NOXA (**Figure 3C, D**).

Reciprocally, we investigated the region in NOXA that is necessary for WSB2 binding. By

generating a series of NOXA deletion mutants and conducting co-IP assays, we found that the

C-terminal region (40-54 aa) of NOXA mediates its binding to WSB2 (**Figure 3E-G**). This

region contains a mitochondrial-targeting domain (MTD)[24]. Mutation of key residues in this

domain (5A mutant) eliminated the interaction between NOXA and WSB2. In contrast,

mutation of key residues in the BH3 domain (3E mutant) did not alter the interaction between

NOXA and WSB2 (**Figure 3H**). The NOXA-5A mutant was resistant to WSB2-mediated

degradation and ubiquitination (**Figure 3I, J**), and had a longer half-life compared to

NOXA-WT (**Figure 3K, L**). NOXA contains three lysine residues that can be attached by

ubiquitin[25]. By mutating these lysine residues to arginine, we found that WSB2-mediated

NOXA ubiquitination was completely abolished, although this mutant (KR) exhibits a

comparable WSB2-binding capacity to NOXA-WT (**Figure 3H**, **J**), indicating that these

residues function as ubiquitin attachment sites targeted by the CRL5^{WSB2} complex.

Furthermore, incubating cell lysates with a C-terminal NOXA peptide (40-54 aa) before

conducting co-IP assays resulted in a dose-dependent reduction in the WSB2-NOXA

interaction (**Figure 3M**). Based on this, we hypothesized that transduction of the C-terminal

NOXA peptide into cells could competitively inhibit WSB2-mediated NOXA degradation. To

efficiently deliver this peptide into cells, we synthesized a fusion peptide in which the

C-terminal peptide was connected to the cell-penetrating poly-arginine (R8) sequence.

Treatment of cells with this fusion peptide dose-dependently increased endogenous NOXA

protein levels in C4-2B and Huh-7 cells (**Figure 3N**).

Collectively, these data indicate that the C-terminal region of NOXA is crucial for its

interaction with WSB2 and indispensable for WSB2-mediated NOXA degradation.

Co-inhibition of WSB2 and anti-apoptotic BCL-2 family proteins causes synthetic

lethality via apoptotic cell death

Despite the strong accumulation of NOXA protein in WSB2-deficient cells, we did not observe

obvious spontaneous apoptosis during daily cell culture. This indicated that NOXA

upregulation alone may not be sufficient to trigger spontaneous apoptosis. Consequently, we

investigated whether co-depletion of an anti-apoptotic BCL-2 family protein would

synergistically induce apoptosis in WSB2-deficient cells. Stable cell lines were generated in

which BCL-XL or MCL-1 was depleted by shRNAs, followed by further depletion of WSB2 with

siRNAs. As shown in **Figure 4A-D**, co-depletion of BCL-XL/WSB2 or MCL-1/WSB2 pairs

resulted in substantial apoptosis, as evidenced by Western blot (WB) analysis detecting

caspase cleavage and flow cytometry detecting early apoptotic markers. The E3 ubiquitin

ligase MARCH5 co-exists with WSB2 in a functional module (**Figure 1B**), and previous studies

have shown that depletion of MARCH5 sensitizes cells to MCL-1 inhibitors or BCL-2/BCL-XL

inhibitors[18, 26, 27]. Consistently, we observed that co-depletion of MARCH5/WSB2 also induced

substantial apoptosis (**Figure 4E, F**).

Alternatively, we employed a pharmacological approach to inhibit anti-apoptotic BCL-2

family proteins. ABT-737 is a small molecule drug that inhibits BCL-2 and BCL-XL, while

AZD5991 is a small molecule drug that inhibits MCL-1. Either ABT-737 or AZD5991 caused

only moderate apoptosis in parental C4-2B or Huh-7 cells. However, these drugs induced

substantial apoptosis in WSB2-deficient cells (Figure **4G-J**, **Supplementary Figure 3A, B**). In

addition to direct inhibitors of the BCL-2 family proteins, inhibitors of cyclin-dependent kinase 9

(CDK9) can indirectly target MCL-1 by suppressing the transcriptional activation of MCL-1

mRNAs[28]. Indeed, we found that the CDK9 inhibitor BAY-1143572 induced moderate

apoptosis in parental cells but induced substantial apoptosis in WSB2-deficient cells (**Figure**

4K, L).

Collectively, these data indicate that WSB2 shows synthetic lethality with anti-apoptotic

BCL-2 family proteins in cancer cells.

The anti-apoptotic function of WSB2 is primarily reliant on NOXA downregulation

We investigated to determine whether WSB2 plays a role in restricting apoptosis by

destabilizing the NOXA protein. As shown in **Figure 5A-D**, reducing NOXA expression

through shRNA knockdown in WSB2-deficient C4-2B or Huh-7 cells largely, though not

completely, reversed the substantial apoptosis induced by ABT-737 treatment. Likewise,

knockdown of NOXA in WSB2-deficient C4-2B cells largely reversed the apoptosis triggered

by MCL-1 inhibitor AZD5591 (**Figure 5E, F**). Moreover, we demonstrated that treatment with

the C-terminal NOXA peptide sensitized Huh-7 cells to ABT-737-induced apoptosis (**Figure**

5G, H). In xenograft tumor assays, we observed that the C-terminal NOXA peptide or

ABT-737 treatment exhibited moderate reductions in tumor growth. Strikingly,

co-administration of the C-terminal NOXA peptide and ABT-737 produced significantly

synergetic inhibitory effects on Huh-7 tumors (**Figure 5I, J**).

Collectively, these data indicate that WSB2 deficiency-induced hypersensitivity to BCL-2

family protein inhibitors was at least in part, caused by NOXA accumulation.

Wsb2 knockout mice are more susceptible to apoptosis triggered by BCL-2 family

protein inhibitors

To gain deeper insights into the physiological roles of WSB2 in apoptosis *in vivo*, we

established a Wsb2 knockout mouse model (**Supplementary Figure 4A**). Homozygous

Wsb2$^{-/-}$ knockout mice were observed to be born at Mendelian ratios and exhibited a normal

lifespan with no apparent morphological or behavioral abnormalities (**Supplementary Figure**

4B, C). Although some Wsb2$^{-/-}$ mice displayed reduced body size after birth, their adult size

generally matched that of wild-type mice. At week 4, we collected multiple mouse tissues,

including heart, liver, lung, kidney, and brain. WB analyses demonstrated varying degrees of

upregulation in NOXA proteins in the tissues of Wsb2$^{-/-}$ mice compared to wild-type mice. A

strong upregulation of NOXA proteins were observed in the liver and heart tissues of Wsb2$^{-/-}$

mice, but not in lung, kidney, and brain tissues, indicating WSB2 modulate NOXA protein

levels in a tissue-specific manner. However, the protein levels of caspase 3, 7, and 9 in these

tissues were similar between Wsb2$^{-/-}$ mice and wild-type mice (**Supplementary Figure 4D**).

Immunofluorescence and WB analyses of heart and liver tissues revealed that cleaved

caspases were nearly undetectable in both wild-type and Wsb2$^{-/-}$ mice (**Figure 6A-E**).

Consistent with the results from *in vitro* cell culture, Wsb2 deletion alone was insufficient to

induce significant apoptosis in mouse organs. We then examined whether pharmacological

inhibition of anti-apoptotic BCL-2 family proteins in Wsb2$^{-/-}$ mice would induce substantial

apoptosis at the organ level. To test this hypothesis, we utilized ABT-199 (Venetoclax), a

highly specific BCL-2 inhibitor. Immunofluorescence and WB analyses of heart and liver

tissues showed the presence of obvious cleaved caspases, cleaved PARP1, and TUNEL

positive cells in ABT-199-treated Wsb2$^{-/-}$ mice, but not in wild-type mice (**Figure 6A-E,**

Supplementary Figure. 5A-I). Furthermore, we isolated primary mouse hepatocytes and

exposed them to ABT-199. Consistent with the *in vivo* treatment, *in vitro* cultured Wsb2$^{-/-}$

mouse hepatocytes exhibited increased susceptibility to ABT-199-triggered apoptosis (**Figure**

6F). To assess whether the cardiac injury was caused by ABT-199 treatment, we measured

the levels of several cardiac enzyme markers, including CK (creatine kinase), CK-MB (creatine

kinase isoenzyme MB), α-HBDH (α-hydroxybutyrate dehydrogenase), and LDH (lactate

dehydrogenase) in serum. As shown in **Figure 6G**, ABT-199 administration led to a significant

elevation in the levels of these cardiac enzymes in Wsb2$^{-/-}$ mice, whereas no such effect was

observed in wild-type mice. This suggested that simultaneous inhibition of Wsb2 and Bcl-2

resulted in heart injury, potentially due to uncontrolled apoptosis in cardiomyocytes.

To investigate whether the anti-apoptotic function of Wsb2 is primarily reliant on mouse

NOXA, we isolated mouse embryonic fibroblasts (MEFs). NOXA protein levels were markedly

upregulated in Wsb2$^{-/-}$ MEFs compared to wild-type MEFs (**Figure 6H**). Moreover, reducing

the expression of NOXA through shRNA-mediated knockdown in Wsb2$^{-/-}$ MEFs largely

reversed the substantial apoptosis induced by ABT-199 treatment (**Figure 6I**).

Collectively, these data indicate that WSB2-mediated NOXA destabilization is evolutionally

conserved and this regulatory axis is critical for maintaining tissue homeostasis.

WSB2 is overexpressed in several human cancer types

Given the intriguing anti-apoptotic role of WSB2, it is worthwhile to investigate whether WSB2

expression is altered in human cancers. To address this, we analyzed RNA-sequencing

(RNA-seq) data from the TCGA cancer cohorts. Remarkably, WSB2 mRNA expression was

found to be significantly elevated in the majority of cancer types, including prostate

adenocarcinoma (PRAD)and liver hepatocellular carcinoma (LIHC) (**Figure 7A**). Further

analysis of the TCGA PRAD cohort revealed a positive correlation between WSB2 mRNA

expression and several clinical parameters, including Gleason score (**Figure 7B**),

pathological stage (**Figure 7C**), and clinical stage (**Figure 7D**), and nodal metastasis status

(**Figure 7E**). Similarly, in the TCGA LIHC cohort, WSB2 mRNA expression showed positive

correlations with clinical stage (**Figure 7F**), pathological grade (**Figure 7G**), and nodal

metastasis status (**Figure 7H**). Furthermore, survival analysis indicated that high WSB2

expression was significantly correlated with shorter overall survival (OS) in the TCGA LIHC

cohort, but not in the PRAD cohort (**Figure 7I, J**). To validate these findings, we performed

immunohistochemistry (IHC) analysis using WSB2-specific antibodies after confirming their

specificity (**Supplementary Figure 4C**). Notably, IHC analysis of a PRAD tissue microarray

revealed a positive correlation between WSB2 protein expression and Gleason score (**Figure**

7K, L). IHC analysis of a LIHC tissue microarray revealed a positive correlation between

WSB2 protein expression and clinical grade (**Figure 7K, L**). At last, we also noticed a higher

expression of WSB2 in sorafenib-resistant LIHC patients as compared to sorafenib-sensitive

patients by analyzing two publicly available RNA-seq datasets (**Figure 7M, N**).

Collectively, these data suggest that WSB2 expression is abnormally elevated in certain

cancer types and may be associated with increased cancer aggressiveness, therapeutic

resistance, and decreased overall survival in those types.

Discussion

The WSB2 protein has been identified in several large-scale proteome mapping analyses as

being co-purified from the CUL5 scaffold complex[29, 30]. With the presence of a SOCS box in its

protein sequence, WSB2 is believed to serve as a receptor for substrates, assisting in their

recognition by the CRL5 E3 ubiquitin ligase complex. Its close paralog, WSB1, is induced by

hypoxia and can form a CRL5^{WSB1} complex that promotes cancer metastasis by inducing VHL

degradation[31]. Additionally, the CRL5^{WSB1} complex overcomes oncogene-induced senescence

by targeting ATM for degradation[32]. However, the physiological substrates of the CRL5^{WSB2}

complex have not yet been reported. Nevertheless, prior to our current study, there are several

pieces of evidence suggesting that this complex may play a role in regulating cell death,

possibly involving BCL-2 family proteins. In a large-scale RNAi screening aimed at

understanding cancer dependencies and synthetic lethal relationships, the top correlates of

WSB2 co-essentiality were found to be MCL-1, BCL-2, and MARCH5, while the most strongly

anti-correlated gene with WSB2 was BAX[33]. Two CRISPR/Cas9 knockout screens also

showed strong synthetic relationships between WSB2 and MCL-1, BCL-2, or MARCH5[34, 35]. A

comprehensive phenotypic CRISPR-Cas9 screen of the ubiquitin pathway revealed that

knockout of CUL5, RBX2, or WSB2 resulted in cells becoming hypersensitive to the CRM1

inhibitor leptomycin[36]. Furthermore, a genome-wide CRISPR inhibition (CRISPRi) screen

conducted in lung cancer cells demonstrated that knockout of CUL5, RBX2, or UBE2F (a

specific E2 for CRL5 E3s) caused cells to become hypersensitive to a CDK9 inhibitor or

MCL-1 inhibitor[28]. By connecting the dots from these studies, our findings provide a

comprehensive scenario. We have shown that WSB2 can assemble an active $CRL5^{WSB2}$

complex, which mediates the ubiquitination and proteasomal turnover of NOXA, maintaining

its low-level expression under basal conditions. WSB2 deficiency leads to a remarkable

accumulation of NOXA, but it alone is not sufficient to trigger spontaneous apoptosis. This is

consistent with previous studies showing that enforced expression of NOXA alone is

ineffective at triggering apoptosis in various cell types[37, 38]. Striking when combined with

genetic or pharmacological inhibition of anti-apoptotic BCL-2 family proteins, massive

apoptosis occurs in WSB2-deficient cells (**Figure 8**). However, it is important to note that

knockdown of NOXA expression in WSB2-deficient cells largely, but not completely, reverses

the massive apoptosis induced by BCL-2 family protein inhibitors, implying that WSB2 may

also modulate apoptosis through other unidentified targets. Indeed, WSB2 interacts with

multiple BCL-2 family proteins (MCL-1, BCL-2, BCL-XL, and BAD) (**Figure 1F-J**). Although

WSB2 does not alter their turnover, it is still possible that WSB2 modulates the apoptotic

function of these proteins through direct binding. Further investigation is warranted to fully

elucidate the molecular mechanisms underlying WSB2-mediated anti-apoptotic function.

Lastly , it would also be interesting to explore whether there are any upstream signals capable

of overriding WSB2-mediated NOXA destabilization under specific stress conditions.

In the current study, our focus was primarily on investigating the *in vivo* anti-apoptotic

function of WSB2. Thus, we did not extensively characterize the potential morphological and

behavioral abnormalities in $Wsb2^{-/-}$ mice. However, recent large-scale mouse phenotype

analyses conducted by the International Mouse Phenotyping Consortium (IMPC) have

reported abnormalities in tooth morphology, locomotor activity, retina, heart, osmotic and

electrolyte balance, as well as male infertility in $Wsb2^{-/-}$ mice[39]. It remains unclear whether

these abnormalities are a result of NOXA accumulation and subsequent dysregulated

apoptosis. Generating Wsb2/NOXA double knockout mice would be beneficial in determining

whether these abnormalities can be reversed by further ablating NOXA expression. Despite

the fact that WSB2 is upregulated in various types of cancer and exhibits a strong

anti-apoptotic function, which makes it a promising target for cancer therapy, it is crucial to

comprehensively understand the downstream substrates regulated by the CRL5^{WSB2} complex.

This understanding will help us evaluate the potential effects of pharmacological inhibition of

WSB2, as it may lead to the dysregulation of other substrates and undesirable side effects.

Since its initial discovery as a novel phorbol-12-myristate13-acetate (PMA) responsive gene

in T cells, and subsequently as a transcriptional target of the genotoxic response regulator p53,

NOXA has emerged as a critical player in regulating cell death pathways in various cell types

under stressed conditions[40]. Notably, NOXA is implicated in fine-tuning apoptosis induction in

cancer cells treated with genotoxic anticancer drugs, including paclitaxel (a microtubule

targeting agent), bortezomib (a proteasome inhibitor), and MLN4924 (a CRL E3 ligase

inhibitor)[40]. These different agents engage distinct mechanisms, such as transcriptional

activation or protein stabilization, to upregulate NOXA protein levels and initiate apoptotic cell

death[40]. The half-life of NOXA protein was very short, as it undergoes ubiquitin-proteasomal

degradation mediated by the addition of ubiquitin to specific lysine residues[10]. In mantle cell

lymphoma (MCL) cell lines, despite high NOXA transcript levels, low NOXA protein expression

is observed due to rapid protein degradation[41]. Similarly, paradoxical downregulation of NOXA

protein is observed in Cushing's disease (CD) adenomas, despite transcriptional upregulation

caused by recurrent promoter hypomethylation[42]. These observations suggest that certain

tumor cells may exploit pathways to accelerate NOXA degradation, thus suppressing

apoptosis. The elevated NOXA mRNA levels observed in tumor cells may potentially serve as

a compensatory mechanism to counteract the reduction in NOXA protein levels. Previous

studies have only partially characterized the ubiquitin-proteasomal degradation of NOXA,

showing that UBE2F, in conjunction with RBX2, induces CUL5 neddylation, leading to CRL5

E3 activation and subsequent NOXA degradation[28]. In lung cancer tissues, high levels of

UBE2F and CUL5 correlate with reduced NOXA levels and poorer survival in patients.

However, the specific CRL5 substrate receptor responsible for NOXA destabilization has yet to

be identified[43]. In our study, we identified WSB2 as the substrate receptor for NOXA, thereby

shedding light on its role in regulating NOXA turnover. Further investigation is needed to

determine whether WSB2 dysregulation is responsible for the accelerated protein turnover

observed in various cancer types, such as MCL and CD adenomas. It should also be noted

that WSB2 only facilitates NOXA destabilization in certain tissues/organs, such as heart, and

liver, in mouse models. It is not surprising that a specific substrate can be targeted by multiple

E3 ubiquitin ligases. A previous study has indicated that treatment with a proteasome inhibitor

could further increase NOXA protein levels in CUL5 knockout cells, suggesting that the

turnover of NOXA can be regulated by additional ubiquitin ligases apart from the CRL5 E3 [28].

In fact, the RING domain containing E3 ubiquitin ligases MARCH5 have been reported to

mediate NOXA degradation [18-20, 26, 27]. Further investigation is needed to determine which E3

ubiquitin ligase(s) play the predominant roles in specific tissues/organs or types of cancer.

Although bortezomib and MLN4924 have proven effective in stabilizing the NOXA protein

and promoting NOXA-dependent apoptosis, their broad inhibition of the proteasome or all CRL

E3 ligases, respectively, inevitably leads to side effects. In this study, we have conducted

preliminary investigations on the use of a competitive peptide to effectively inhibit the binding

of WSB2 and NOXA, resulting in the accumulation of NOXA proteins and increased sensitivity

to ABT-737. In the future, other potential therapeutic strategies can be explored, such as

designing PROTAC molecules specifically for degrading WSB2 or developing small molecules

to disrupt the interaction between WSB2 and NOXA. Extensive research is needed to

determine the safety and efficacy of these approaches in preclinical cancer models.

Materials and methods

Acquisition and analysis of DepMap and drug sensitivity datasets

Gene co-dependencies were determined using the Achilles datasets

(https://depmap.org/portal/). The Achilles dataset contains dependency scores from

genome-scale essentiality screens scores of 789 cell lines. As a measure of co-dependency,

the Pearson's correlation coefficient of essentiality scores was computed for all gene pairs. GO

analysis for the top 500 genes co-dependent-with WSB2 was performed using PANTHER to

search enriched biological processes and pathways. Co-essential module assigns of cellular

component: Bcl-2 family protein complex were obtained from a previously published

dataset(Gene co-dependency (https://mitra.stanford.edu/bassik/michael/cluster_heatmaps/)[17].

To identify genetic and pharmacologic perturbations that induce similar effects on cell viability,

a Web tool, DepLink （ https://shiny.crc.pitt.edu/deplink/) was used[21].

Cell line, cell culture, transfection, and lentiviral infection

293T, HeLa, C4-2B, and Huh-7 cells were obtained from the American Type Culture

Collection (ATCC). 293T, HeLa and Huh-7 cells were maintained in Dulbecco's modified

Eagle's medium (DMEM) supplemented with 10% FBS. C4-2B cells were maintained in

RPMI1640 medium supplemented with 10% FBS. We routinely perform DNA fingerprinting and

PCR to verify the authenticity of the cell lines and to ensure they are free of mycoplasma

infection. We conducted transient transfection using EZ Trans (Shanghai Life-iLab Biotech).

For lentiviral transfection, we transfected pLKO shRNA KD and virus-packing constructs into

293 T cells. The viral supernatant was collected after 48 h. The cells were then infected with

the viral supernatant in the presence of polybrene (8 µg/ml) and selected in growth media

containing puromycin (1.5 µg/ml). The gene-specific shRNA or siRNA sequences can be found

in **Supplementary Table 4**.

Antibodies, chemicals, and kits

The information of antibodies, chemicals, and kits used in this study is listed in

Supplementary Table 5, 6.

Plasmid construction

The plasmids used for transient overexpression were constructed using the

pCMV-FLAG/Myc vector (Clontech). Point and deletion mutants were engineered utilizing the

KOD-Plus-Mutagenesis Kit (TOYOBO) following the manufacturer's instructions. Single-guide

RNAs (sgRNAs) targeting WSB2 (http://crispr.mit.edu) were subcloned into the

pSpCas9(BB)-2A-Puro (PX459) vector for gene knockout (KO). Short hairpin RNAs (shRNAs)

targeting WSB2 or BCL-2 family proteins were subcloned into the pLKO.1 puro vector

(Addgene) for gene KD. The sequences of gene-specific sgRNAs and shRNAs are listed in

supplementary table 4.

Isolation of nucleic, cytoplasmic, and mitochondrial fractions

HeLa cells were prepared for nuclear, cytoplasmic, and mitochondrial extraction by

density-gradient centrifugation. Briefly, HeLa cells were washed three times with PBS. Then

the cells are suspended by using hypotonic solution (140 mM KCl, 10 mM EDTA, 5 mM MgCl2,

20 mM HEPES (pH 7.4), and the protease inhibitor). Then 5×10^6 HeLa cells were ground with

a glass homogenizer in an ice bath for 25 strokes. Nuclear, cytoplasmic, and mitochondrial

fractions were separated through differential centrifugation (800×g, 10 min, 4°C and 12,000×g,

35 min, 4°C). The supernatant (cytoplasmic fraction) and pellet (mitochondrial fraction) were

collected, and the pellet was further washed with wash buffer (800 mM KCl, 10 mM EDTA,

5 mM MgCl2, and 20 mM HEPES (pH 7.4), and the protease inhibitor) for three times and yield

the final mitochondrial fraction. To confirm that pure extracts were obtained, the mitochondrial,

nuclear, and cytoplasmic fractions were separated by SDS-PAGE, and the presence of

mitochondrial VDAC1, BCL2, nuclear Histone H3, and cytoplasmic GAPDH was detected by

Immunoblot.

Isolation of submitochondrial fractions

Six mitochondrial fraction samples were divided into three groups, with two samples in each

group. The first group was resuspended in 300 µl of homogenization buffer, the second group

in 300 µl of hypotonic swelling buffer (10 mM HEPES/KOH, pH 7.4, 1 mM EDTA), and the third

group in 300 µl of homogenization buffer supplemented with 0.5% (V/V) Triton X-100, followed

by a 10-min incubation on ice. Subsequently, one sample from each group was exposed to

proteinase K (70 µg/ml) for 20 min on ice, while the other sample was kept untreated as a

control. Following the treatments, mitochondrial proteins were precipitated using 300 µl of 30%

TCA (W/V) and incubated on ice for 10 min. The proteins were collected by centrifugation at

18,000 g for 10 min at 4°C, washed with 1 ml of 100% ethanol, and centrifuged again. The

resulting pellets were dissolved in 100 µl of SDS Lysis Buffer, boiled at 105°C for 8 min, and

subjected to WB analyses.

CRISPR-Cas9 mediated gene KO cell lines

C4-2B or Huh-7 cells were plated and transfected with PX459 plasmids overnight. 24 h after

transfection, 1 µg/ml puromycin was used to screen cells for three days. Living cells were

seeded in 96-well plate by limited dilution to isolate monoclonal cell line. The knock out cell

clones are screened by WB and validated by Sanger sequencing. Sequences of gene-specific

sgRNAs are listed in **Supplementary Table 4**.

RT-qPCR assays

Total RNA from cells were extracted by using TRIzol reagent (TIANGEN), followed by

reverse transcription into cDNA using the HiScript III First Strand cDNA Synthesis Kit (Vazyme).

The synthesized cDNAs were then subjected to PCR amplification using ChamQ SYBR qPCR

Master Mix (Vazyme) in CFX Real-Time PCR system (Bio-Rad). The relative mRNA levels of

DDHD2 were quantified using the $2^{-\Delta\Delta CT}$ method with normalization to GAPDH. The primer

sequences are listed in the **Supplementary Table 4**.

***In vivo* ubiquitination assays**

293T cells were transfected with HA-ubiquitin and indicated constructs. After 36 h, cells were

treated with MG132 (30 µM) for 6 h and then lysed in 1% SDS buffer (Tris [pH 7.5], 0.5 mM

EDTA, 1 mM DTT) and boiled for 10 min. For immunoprecipitation, the cell lysates were diluted

10-fold in Tris-HCL buffer and incubated with anti-NOXA or IgG-conjugated beads (Sigma) for

4 h at 4 °C. The bound beads are then washed four times with BC100 buffer (20 mM Tris-Hcl,

pH 7.9,100 mM NaCl,0.2 mM EDTA, 20% glycerol) containing 0.2% Triton X-100. The proteins

were eluted with FLAG peptide for 2 h at 4°C. The ubiquitinated form of NOXA was detected by

WB using anti-HA antibody.

IF and confocal microscopy

HeLa cells were seeded on glass coverslips in 12-well plates and harvested at 70%

confluence. The cells were washed with PBS and fixed with 4% paraformaldehyde in PBS.

After permeabilization with 0.3% Triton X-100 for 5 min and then in the blocking solution (PBS

plus 5% donkey serum), for 1h at room temperature (RT). The cells were then incubated with

primary antibodies at 4 °C for overnight. After washing with PBST buffer, fluorescence-labelled

secondary antibodies were applied. DAPI was utilized to stain nuclei. The glass coverslips

were mounted on slides and imaged using a confocal microscope (LSM880, Zeiss) with a

63*/1.4NA Oil PSF Objective. Quantitative analyses were performed using ImageJ software.

For mouse tissues staining, the mouse tissues were isolated from mice after perfusion with

0.1 M PBS (pH7.4) and fixed for 3 days with 4% PFA at 4 °C. The tumor tissues were then

placed in 30% sucrose solution for 5 days for dehydration. The tumors were embedded into

the OCT block and frozen for cryostat sectioning. Cryostat sections (45-μm thick) were

washed with PBS, and then incubated in blocking solution (PBS containing 10% goat serum,

0.3% Triton X-100, pH7.4) for 2h at RT. The samples were stained with primary antibodies

overnight at 4 °C, after washing with PBST buffer, fluorescence-labelled secondary antibodies

were applied at RT for 2h. DAPI was utilized to stain nuclei. The sections were then sealed

with an anti-fluorescence quencher. The samples were visualized and imaged using a confocal

microscope (LSM880, Zeiss) with a 63*/1.4NA Oil PSF Objective. Quantitative analyses were

performed using ImageJ software.

Apoptosis assays

Annexin V-FITC (Fluorescein isothiocyanate) and propidium iodide (PI) double staining

(Dojindo) were used to detect the apoptosis rates. The cells were cultured in six-well plates at

a density of 1.2×10^5/well and allowed to adhere to culture plate overnight. Then the medium

was replaced with fresh medium containing indicated drugs for a certain time. The cells were

then trypsinized by EDTA-free trypsin and washed twice with cold PBS. Aliquots of the cells

were resuspended in 100 μl of binding buffer and stained with 5 μl of annexin V-FITC and 5 μl

of PI working solution for 15 min at RT in the dark. All flow cytometry analyses were carried out

using a Fortessa flow cytometer (BD Bioscience). The subsequent data analysis was

conducted using FlowJo software.

Generation and breeding of Wsb2 KO mice

Mice with murine Wsb2 KO were designed and generated from Shanghai Model Organisms

Center (Shanghai, China). In brief, CRISPR/Cas9 system were microinjected into the fertilized

eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which

were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained

by mating positive F0 generation mice with C57BL/6JGpt mice. The genotype of F1 mice was

identified by PCR and confirmed by sequencing. The sequences used for CRISPR-Cas9

editing and the primers used for genotyping are listed in **Supplementary Table 4**.

Mice were maintained under a 12 h/12 h light/dark cycle at 22–25 °C and 40–50% humidity

with standard food and water available ad libitum. All procedures for animal care and animal

experiments were carried out in accordance with the guidelines of the Care and Use of

Laboratory Animals proposed by Institute of Development Biology and Molecular Medicine and

Shanghai Municipality, PR China. The male C57BL/6JGpt mice (8 months old) were divided

into 2 groups (Wsb2$^{+/+}$ and Wsb2$^{-/-}$; n=5/group), each group was given by gavage of ABT-199

(100 mg/kg per day) or vehicle (10% β-cyclodextrin) for one week. Then, we collect blood

samples from the tail vein of mice before and after oral administration to measure indicators of

myocardial zymogram using an Olympus AU640 automatic biochemical analyzer (Olympus).

MEFs generation and immortalization

Timed pregnant female mice at embryonic day 12.5 to 14.5 were sacrificed, and the

embryos were carefully dissected to remove the cerebrum, internal organs, and limbs. The

remaining tissues were cut into small pieces and treated with trypsin-EDTA (0.25%) for 10 min

at 37 °C. The trypsin was neutralized with DMEM, a complete medium supplemented with 10 %

fetal bovine serum and 1% penicillin/streptomycin. The culture media were changed every 2-3

days until the cells reached confluence. To immortalize MEFs, they were passaged up to

approximately 10 times before infection with lentiviral vectors expressing the SV40 large

T-antigen. Stable transduction was achieved with puromycin selection. The successful

integration of the immortalizing gene was confirmed through Sanger sequencing and WB

analysis.

Mouse tumor implantation

All experimental protocols were approved in advance by the Ethics Review Committee for

Animal Experimentation of Fudan University. 4–6-week-old male BALB/c nu/nu mice obtained

from SLAC Laboratory Animal Co., Ltd. were bred and maintained in our institutional

pathogen-free mouse facilities. Mice were randomly divided into 4 groups (n=6/group): vehicle

(distilled water); ABT-737 (20 mg/kg); R8-C-ter (20 mg/kg); and ABT-737+R8-C-ter. Huh-7

tumors were established by subcutaneously injecting 5×106 Huh-7 cells in 100 µl of PBS

buffer into the right flank of 6-week-old nude mice. After 1 week, vehicle and indicated drug

treatments were administered once daily by intraperitoneal injection (i.p). At the end of 3

weeks, mice were killed and *in vivo* solid tumors were dissected and weighed.

Pan-cancer dataset acquisition and analysis

Pan-cancer gene expression analysis based on tumor and normal samples was derived

from the TCGA (transcriptome datasets, http://gepia2.cancer-pku.cn/). Public databases

(TCGA) were used to analyze the correlations of WSB2 expression with clinical risk factors.

Additional publicly available RNA-seq datasets (GSE104580 and GSE109211) provided for

sorafenib-response and sorafenib-non-response HCC patients.

IHC analysis

A total of 84 patients with localized PRAD, who underwent radical prostatectomy between

January 2007 and July 2014 at Fudan University Shanghai Cancer Center (FUSCC), were

included in this study. All the patients underwent regular postoperative reviews and had

long-term follow-up data. This study was in accordance with the recommendations of the

Research Ethics Committee of FUSCC according to the provisions of the Declaration of

Helsinki (as revised in Fortaleza, Brazil, October 2013). The protocol was approved by the

Research Ethics Committee of FUSCC. Informed consent for the use of clinical data was

obtained from all the patients recruited in this study. The TMA consist of 29 LIHC patient

specimens was obtained from Shanghai Biochip Co., Ltd (Shanghai). To confirm the specificity

of the anti-WSB2 antibody, we conducted genetic control for the IHC analysis using an

anti-WSB2 antibody in both parental and WSB2 KO C4-2B cells.

Statistical analysis

Statistical analysis was performed using GraphPad Prism (GraphPad Software), and the

differences between the two groups were analyzed using one-way analysis of variance

(ANOVA) or two-way analysis of variance ANOVA. All data were displayed as means ± S.D.

values for experiments conducted with at least three replicates. * represents $p < 0.05$; **

represents $p < 0.01$; *** represents $p < 0.001$, **** represents $p < 0.0001$.

Acknowledgements

Authors' contributions

C.W. conceived the study. D.J., K.C., and J.J. performed the experiments and data analyses.

C.W., D.J., K.G., and Y.X. analyzed and interpreted the data. C.W. wrote the manuscript.

Funding

This work was in part supported by the National Natural Science Foundation of China (No.

92357301, 32370726, 91957125 to C.W., 82272992, 91954106, and 81872109 to K.G.;

82270415 to L.W), the State Key Development Programs of China (No. 2022YFA1104200 to

C.W), the Natural Science Foundation of Shanghai (No. 22ZR1406600 to C.W.; 22ZR1449200

to K.G, 22ZR1448600 to Y.X), Science and Technology Research Program of Shanghai (No.

9DZ2282100). Open Research Fund of the State Key Laboratory of Genetic Engineering,

Fudan University (No. SKLGE-2111 to K.G.), Science and Technology Research Program of

Shanghai (No. 9DZ2282100), and Central Guidance on Local Science and Technology

Development Foundation (No. 2021ZY0037 to R.M.).

Competing interests

The authors declare no competing interests.

Reference

1.   Hanahan, D. & Weinberg, R.A. Hallmarks of cancer: the next generation. *Cell* **144**, 646-674 (2011).

2.   Carneiro, B.A. & El-Deiry, W.S. Targeting apoptosis in cancer therapy. *Nat Rev Clin Oncol* **17**, 395-417 (2020).

3.   Kale, J., Osterlund, E.J. & Andrews, D.W. BCL-2 family proteins: changing partners in the dance towards death. *Cell Death Differ* **25**, 65-80 (2018).

4.   Youle, R.J. & Strasser, A. The BCL-2 protein family: opposing activities that mediate cell death. *Nat Rev Mol Cell Biol* **9**, 47-59 (2008).

5.   Cory, S., Roberts, A.W., Colman, P.M. & Adams, J.M. Targeting BCL-2-like Proteins to Kill Cancer Cells. *Trends Cancer* **2**, 443-460 (2016).

6.   Wei, A.H. *et al.* Venetoclax plus LDAC for newly diagnosed AML ineligible for intensive chemotherapy: a phase 3 randomized placebo-controlled trial. *Blood* **135**, 2137-2145 (2020).

7.   DiNardo, C.D. *et al.* Azacitidine and Venetoclax in Previously Untreated Acute Myeloid Leukemia. *N Engl J Med* **383**, 617-629 (2020).

8.   Ploumaki, I. *et al.* Bcl-2 pathway inhibition in solid tumors: a review of clinical trials. *Clin Transl Oncol* **25**, 1554-1578 (2023).

9.   Zhao, Y., Xiong, X. & Sun, Y. Cullin-RING Ligase 5: Functional characterization and

its role in human cancers. *Semin Cancer Biol* **67**, 61-79 (2020).

10.   Mahrour, N. *et al.* Characterization of Cullin-box sequences that direct recruitment of

Cul2-Rbx1 and Cul5-Rbx2 modules to Elongin BC-based ubiquitin ligases. *J Biol*

Chem **283**, 8005-8013 (2008).

11.   Zhang, Y. *et al.* WD repeat and SOCS box containing protein 2 in the proliferation,

cycle progression, and migration of melanoma cells. *Biomed Pharmacother* **116**,

108974 (2019).

12.   Ma, L., Zhang, Y. & Hu, F. miR‑28‑5p inhibits the migration of breast cancer by

regulating WSB2. *Int J Mol Med* **46**, 1562-1570 (2020).

13.   Wei, X. *et al.* WSB2 as a target of Hedgehog signaling promoted the malignant

biological behavior of Xuanwei lung cancer through regulating Wnt/beta-catenin

signaling. *Transl Cancer Res* **9**, 7394-7404 (2020).

14.   Tsherniak, A. *et al.* Defining a Cancer Dependency Map. *Cell* **170**, 564-576 e516

(2017).

15.   Bayraktar, E.C. *et al.* Metabolic coessentiality mapping identifies C12orf49 as a

regulator of SREBP processing and cholesterol metabolism. *Nat Metab* **2**, 487-498

(2020).

16.   Price, C. *et al.* Genome-Wide Interrogation of Human Cancers Identifies EGLN1

Dependency in Clear Cell Ovarian Cancers. *Cancer Res* **79**, 2564-2579 (2019).

17.   Wainberg, M. *et al.* A genome-wide atlas of co-essential modules assigns function to

uncharacterized genes. *Nat Genet* **53**, 638-649 (2021).

18.   Nakao, F. *et al.* Targeting a mitochondrial E3 ubiquitin ligase complex to overcome

AML cell-intrinsic Venetoclax resistance. *Leukemia* **37**, 1028-1038 (2023).

19.   Djajawi, T.M. *et al.* MARCH5 requires MTCH2 to coordinate proteasomal turnover of

the MCL1:NOXA complex. *Cell Death Differ* **27**, 2484-2499 (2020).

20.   Haschka, M.D. *et al.* MARCH5-dependent degradation of MCL1/NOXA complexes

defines susceptibility to antimitotic drug treatment. *Cell Death Differ* **27**, 2297-2312

(2020).

21.   Nayak, T. *et al.* DepLink: an R Shiny app to systematically link genetic and

pharmacologic dependencies of cancer. *Bioinform Adv* **3**, vbad076 (2023).

22.   Corsello, S.M. *et al.* Discovering the anti-cancer potential of non-oncology drugs by

systematic viability profiling. *Nat Cancer* **1**, 235-248 (2020).

23.   Iorio, F. *et al.* A Landscape of Pharmacogenomic Interactions in Cancer. *Cell* **166**,

740-754 (2016).

24.   Seo, Y.W. *et al.* The molecular mechanism of Noxa-induced mitochondrial dysfunction

in p53-mediated cell death. *J Biol Chem* **278**, 48292-48299 (2003).

25.   Pang, X. *et al.* The carboxyl-terminal tail of Noxa protein regulates the stability of Noxa

and Mcl-1. *J Biol Chem* **289**, 17802-17811 (2014).

26.   Subramanian, A., Andronache, A., Li, Y.C. & Wade, M. Inhibition of MARCH5 ubiquitin

ligase abrogates MCL1-dependent resistance to BH3 mimetics via NOXA. *Oncotarget*

7, 15986-16002 (2016).

27.   Arai, S. *et al.* MARCH5 mediates NOXA-dependent MCL1 degradation driven by

kinase inhibitors and integrated stress response activation. *Elife* **9** (2020).

28.   Kabir, S. *et al.* The CUL5 ubiquitin ligase complex mediates resistance to CDK9 and

MCL1 inhibitors in lung cancer cells. *Elife* 8 (2019).

29.   Huttlin, E.L. *et al.* Architecture of the human interactome defines protein communities

and disease networks. *Nature* **545**, 505-509 (2017).

30.   Bennett, E.J., Rush, J., Gygi, S.P. & Harper, J.W. Dynamics of cullin-RING ubiquitin

ligase network revealed by systematic quantitative proteomics. *Cell* **143**, 951-965

(2010).

31.   Kim, J.J. *et al.* WSB1 promotes tumor metastasis by inducing pVHL degradation.

Genes Dev **29**, 2244-2257 (2015).

32.   Kim, J.J. *et al.* WSB1 overcomes oncogene-induced senescence by targeting ATM for

degradation. *Cell Res* **27**, 274-293 (2017).

33.   McDonald, E.R., 3rd *et al.* Project DRIVE: A Compendium of Cancer Dependencies

and Synthetic Lethal Relationships Uncovered by Large-Scale, Deep RNAi Screening.

Cell **170**, 577-592 e510 (2017).

34.   DeWeirdt, P.C. *et al.* Genetic screens in isogenic mammalian cell lines without single

cell cloning. *Nat Commun* **11**, 752 (2020).

35.   DeWeirdt, P.C. *et al.* Optimization of AsCas12a for combinatorial genetic screens in

human cells. *Nat Biotechnol* **39**, 94-104 (2021).

36.   Hundley, F.V. *et al.* A comprehensive phenotypic CRISPR-Cas9 screen of the

ubiquitin pathway uncovers roles of ubiquitin ligases in mitosis. *Mol Cell* **81**,

1319-1336 e1319 (2021).

37.   Akhtar, R.S. *et al.* BH3-only proapoptotic Bcl-2 family members Noxa and Puma

mediate neural precursor cell death. *J Neurosci* **26**, 7257-7264 (2006).

38.   Shibue, T. *et al.* Differential contribution of Puma and Noxa in dual regulation of

p53-mediated apoptotic pathways. *EMBO J* **25**, 4952-4962 (2006).

39.   da Silva-Buttkus, P. *et al.* Knockout mouse models as a resource for the study of rare

diseases. *Mamm Genome* **34**, 244-261 (2023).

40.   Ploner, C., Kofler, R. & Villunger, A. Noxa: at the tip of the balance between life and

death. *Oncogene* **27 Suppl 1**, S84-92 (2008).

41.   Dengler, M.A. *et al.* Discrepant NOXA (PMAIP1) transcript and NOXA protein levels: a

potential Achilles' heel in mantle cell lymphoma. *Cell Death Dis* **5**, e1013 (2014).

42.   Asuzu, D.T. *et al.* Pituitary adenomas evade apoptosis via noxa deregulation in

Cushing's disease. *Cell Rep* **40**, 111223 (2022).

43.   Zhou, W. *et al.* Neddylation E2 UBE2F Promotes the Survival of Lung Cancer Cells by

Activating CRL5 to Degrade NOXA via the K11 Linkage. *Clin Cancer Res* **23**,

1104-1116 (2017).

Figure Legends

Figure 1. WSB2 interacts with multiple members of BCL-2 family proteins in cells.

(**A**) The top 20 co-dependent genes of WSB2 in Broad's 21Q2 DepMap dataset. Red,

anti-apoptotic BCL-2 family proteins. Blue, pro-apoptotic BCL-2 family proteins.

(**B**) A co-essential module containing WSB2 and other BCL-2 family proteins using a dataset

of CRISPR screens in 485 cell lines from the Achilles DepMap project. Cellular component:

Bcl-2 family protein complex. Molecular function: BH domain binding.

(**C-E**) WB analyses of the indicated proteins in the whole cell lysates (WCL) and co-IP samples

of anti-FLAG antibody obtained from 293 T cells transfected with the indicated plasmids.

(**F-J**) WB analyses of the indicated proteins in WCL and co-IP samples of IgG or anti-NOXA

(**F**), anti-MCL-1 (**G**), anti-BCL-2 (**H**), anti-BCL-XL (**I**), or anti-BAD (**J**) antibodies obtained from

293T cells.

(**K**) Representative IF images from HeLa cells transfected with FLAG-WSB2, stained with

FLAG (WSB2), HSP60, and DAPI. Scale bar, 20 μm.

(**L**) The cytoplasmic (Cyto), mitochondrial (Mito), and nuclear fractions (Nuc) from HeLa cells

were prepared as described in Methods section. Histone H3 (nucleus), GAPDH (cytoplasm),

and HSP60 (mitochondria) were used as subcellular fraction markers. WB analyses were

performed to detect the indicated proteins in three fractions from HeLa cells.

(**M**) Mitochondria from HeLa cells were purified as intact mitochondria or treated with

hypotonic swelling buffer or lysed with Triton X−100 buffer. Different mitochondrial

preparations were then digested with or without Proteinase K. WB analyses of the indicated

proteins in three fractions were then performed. OM: outer membrane; IMS: intermembrane

space.

Figure 2. CRL5^WSB2 E3 ubiquitin ligase complex control NOXA protein turnover.

(**A**) WB analyses of the indicated proteins in the WCL from C4-2B cells infected with lentivirus

expressing WSB2-specific shRNA or negative control (NC).

(**B, C**) WB analyses of the indicated proteins in the WCL from parental or WSB KO C4-2B or

Huh-7 cells.

(**D**) WB analyses of the indicated proteins in the WCL from 293 T cells transfected with the

indicated plasmids for 24 h and treated with DMSO or MG132 (20 μM) for 8 h.

(**E**) WB analyses of the indicated proteins in the WCLs from 293 T cells transfected with the

indicated plasmids.

(**F**) WB analyses of the indicated proteins in the WCL from parental or WSB2 KO C4-2B cells

stably overexpressing empty vector (EV), WSB2-WT or its mutants.

(**G**) WB analyses of the indicated proteins in the WCL from C4-2B cells transfected with the

indicated siRNAs.

(**H, I**) WB analyses of indicated proteins in WCLs of parental and WSB2 KO C4-2B cells

treated with cycloheximide (CHX, 50 µg/ml) and harvested at different time points. (**G**) At each

time point, the intensity of NOXA was normalized to the intensity of GAPDH and then to the

value at 0h. Data are shown as means ± SD (n=3).

(**K-M**) WB analyses of the products of *in vivo* ubiquitination assays performed using WCL from

293T cells transfected with the indicated plasmids and treated with MG132(20 µM).

(**N**) WB analyses of the products of *in vivo* ubiquitination assays. Co-IP using anti-IgG or

anti-NOXA antibody in the WCL prepared from parental and WSB2 KO C4-2B cells

transfected with HA-Ub for 24 h and treated with MG132(20 µM) for 8 h.

P values are calculated by the Two-way ANOVA test in (**G**) and One-way ANOVA test in (**H**).

Figure 3. Identification of the mutual-binding regions of WSB2 and NOXA.

(**A**) Schematic representation of WSB2 deletion mutants.

(**B**) WB analyses of the indicated proteins in WCLs and co-IP samples of anti-FLAG antibody

obtained from 293T cells transfected with the indicated plasmids.

(**C**) WB analyses of the indicated proteins in the WCL from 293 T cells transfected with the

indicated plasmids.

(**D**) WB analyses of the products of *in vivo* ubiquitination assays performed using WCL from

293T cells transfected with the indicated plasmids for 24 h and treated with MG132(20 µM).

(**E**) Schematic representation of NOXA deletion or point mutants.

(**F-H**) WB analyses of the indicated proteins in the WCLs and co-IP samples of anti-FLAG

antibody obtained from 293T cells transfected with indicated plasmids.

(**I**) WB analyses of the indicated proteins in the WCL from 293 T cells transfected with the

indicated plasmids.

(**J**) WB analyses of the products of *in vivo* ubiquitination assays performed using WCL from

293T cells transfected with the indicated plasmids and treated with MG132(20 μM).

(**K, L**) WB analyses of the indicated proteins in the WCL from 293 T cells transfected with the

indicated plasmids treated with cycloheximide (CHX, 50 μg/ml) and harvested at different time

points. (**L**) At each time point, the intensity of NOXA was normalized to the intensity of GAPDH

and then to the value at 0 h. Data are shown as means ± SD (n=3). *P* values are calculated by

the Two-way ANOVA test.

(**M**) 293T cells were transfected with the indicated plasmids. WSB2-NOXA complex was

co-immunoprecipitated by anti-FLAG antibody, and then the bound complex was incubated

with the increasing amounts of C-terminal NOXA peptide (200, 400 μg/ml) or the

corresponding scramble peptide for 12 h. Bound material was subjected to WB analyses. C-ter:

C-terminal NOXA peptide.

(**N**) WB analyses of the indicated proteins in WCLs from C4-2B and Huh-7 cells treated with

the increasing concentration of C-terminal cell-penetrating peptide of NOXA or the

corresponding R8 peptide for 12 h.

Figure 4. Combined inhibition of anti-apoptotic BCL-2 family members and WSB2 induces synthetic lethality.

(**A, B**) WB analyses of the indicated proteins in the WCL from C4-2B cells infected with lentivirus expressing BCL-XL-specific shRNA or NC and then transfected with the indicated siRNAs for 36 h. (**B**) Annexin-V-FITC/PI assays were used to stain the harvested cells in (A), of which later flow cytometry analysis was performed. Data are shown as means ± SD (n=3).

(**C, D**) Similar to (A, B), but MCL-1 was knocked down in C4-2B cells.

(**E, F**) Similar to (A, B), but MARCH5 was knocked down in C4-2B cells.

(**G, H**) WB analyses of the indicated proteins in the WCL from parental and WSB2 KO C4-2B cells treated with ABT-737 (20 μM) for 6 h. (**H**) Annexin-V-FITC/PI assays were used to stain

the harvested cells in (H), of which later flow cytometry analysis was performed. Data are

shown as means ± SD (n = 3).

(**I, J**) Similar to (A, B), but AZD5991 (10 μM) was used for treatment in C4-2B cells.

(**K, L**) Similar to (A, B), but BAY-1143572 was used for treatment in C4-2B cells.

P values are calculated by the One-way ANOVA test in (B, D, F) and Two-way ANOVA test in

(H, J, L), ns no significant.

Figure 5. WSB2 primarily exerts its anti-apoptotic function largely via destabilizing

NOXA.

(**A, B**) WB analyses of indicated proteins in WCLs from parental or WSB2 KO C4-2B cells

infected with lentivirus expressing NOXA-specific shRNA or NC, treated with DMSO or

ABT-737 (20 μM) for 6 h. (**B**) Annexin-V-FITC/PI assays were used to stain the harvested cells,

of which later flow cytometry analysis was performed. Data are shown as means ± SD (n = 3).

(**C, D**) Similar to (A, B), but AZD5991 (10 μM) was used for treatment in C4-2B cells.

(**E, F**) Similar to (A, B), but Huh-7 cells were treated.

(**G, H**) WB analyses of the indicated proteins in the WCL from Huh-7 cells treated with the

cell-penetrating C-terminal NOXA peptide (5 μM) or the corresponding scramble peptide (5 μM)

for 12 h and then the cells were treated with increasing doses of ABT-737(10, 20 μM).

Annexin-V-FITC/PI assays were used to stain the harvested cells, of which later flow

cytometry analysis was performed. Data are shown as means ± SD (n = 3).

(**I, J**) Huh-7 cells were injected subcutaneously (s.c.) into the right flank of BALB/c mice and

treated with ABT-737(30 mg/kg), R8-C-ter (20 mg/kg) or R8 (20 mg/kg) as control at different

day points. 6 mice per experimental group. Tumors in each group at day 20 were harvested

and photographed (I) and tumor weight (J) was documented. Data are shown as means ± SD

(n = 6).

P values are calculated by the Two-way ANOVA test in (B, D, F, H).

Figure 6. Validation of the anti-apoptotic function of WSB2 using Wsb2 KO mouse models.

(**A, B**) WB analyses of the indicated proteins in the WCL from heart (A) or liver (B) tissues obtained from Wsb2$^{+/+}$ and Wsb2$^{-/-}$ mice after gavage administration of vehicle or ABT-199 (100 mg/kg/day) for 7 days.

(**C**) Representative IF images from the heart or liver tissues of Wsb2$^{+/+}$ and Wsb2$^{-/-}$ mice after

oral administration of vehicle or ABT-199 (100 mg/kg/day) for 7 days, and stained with

cl-CASP3 and DAPI. Scale bar, 20 μm.

(**D, E**) The mean fluorescence intensity of cl-CASP3 from the heart (D) or liver tissues (E)

obtained from Wsb2$^{+/+}$ and Wsb2$^{-/-}$ mice. Data were shown as means ± SD (n = 5).

(**F**) WB analyses of the indicated proteins in the WCL from the primary hepatocytes of Wsb2$^{+/+}$

and Wsb2$^{-/-}$ mice treated with ABT-199 (10, 20 μM) for 6 h.

(**G**) The levels of myocardial zymogram in serum from Wsb2$^{+/+}$ and Wsb2$^{-/-}$ mice after gavage

administration of vehicle or ABT-199 (100 mg/kg/day) for 7 days.

(**H, I**) WB analyses of indicated proteins in WCLs from Wsb2$^{+/+}$ and Wsb2$^{-/-}$ MEFs infected with

lentivirus expressing NOXA-specific shRNA or NC, treated with DMSO or ABT-199 (20 μM) for

6 h. (**B**) Annexin-V-FITC/PI assays were used to stain the harvested cells, of which later flow

cytometry analysis was performed. Data are shown as means ± SD (n = 3).

P values are calculated by the One-way ANOVA test in (G) and Two-way ANOVA test in (D,

E).

Figure 7. WSB2 expression is upregulated in multiple human cancers.

(**A**) WSB2 mRNA expression in normal and tumor tissues from the TCGA cohort.

(**B–E**) Relationship between WSB2 mRNA expression and Gleason score (**B**), nodal

metastasis status (**C**), clinical T stage (**D**), and pathological T stage (**E**) in PRAD patients from

the TCGA cohort.

(**F–H**) Relationship between WSB2 mRNA expression and clinical stage (**F**), pathological

grade (**G**), and nodal metastasis status (**H**) in LIHC patients from the TCGA cohort.

(**I, J**) Kaplan–Meier survival plots of OS according to WSB2 mRNA expression in PRAD (**I**) and

LIHC (**J**) patients from the TCGA cohorts.

(**K**) Representative IHC staining results for WSB2 in PRAD TMA, scale bar, 50 μm.

(**L**) Quantification analysis of WSB2 IHC staining in PRAD patients by Gleason score

categories. n=84.

(**M**) Representative IHC staining results for WSB2 in LIHC TMA, scale bar, 50 μm.

(**N**) Quantification analysis of WSB2 IHC staining in LIHC patients by tumor grade categories.

n=29.

(**O**) The WSB2 mRNA levels in 81 sorafenib-response and 66 sorafenib-non-response LIHC

patients from GSE104580 dataset.

(**P**) The WSB2 mRNA levels in 42 sorafenib-response and 98 sorafenib-non-response LIHC

patients from GSE109211 dataset.

P values are calculated by the unpaired *t* test in (A, B-H, L, N, O, P) and log-rank test in (I, J).

*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Figure 8. The schematic diagram illustrates that the disruption of the CRL5^{WSB2} E3 Ubiquitin ligase complex results in NOXA stabilization and increased sensitivity to BCL-2 family protein inhibitors.

978   Supplementary Materials for

979   Disruption of WSB2-mediated NOXA Degradation Induces Synthetic Lethality to

980   Anti-apoptotic BCL-2 Family Protein Inhibitors

982   Figures. S1 to S5

983   Other Supplementary Materials for this manuscript include the following:

Table S1 to S6

Supplementary Table 1. Top 100 co-dependent genes of WSB2.

Supplementary Table 2. The GO analysis of top 500 co-dependent genes of

WSB2.

Supplementary Table 3. The DepLink analysis of top correlated drugs with

WSB2.

Supplementary Table 4. Sequence information.

Supplementary Table 5. Antibody information.

Supplementary Table 6 Cell cultures, chemicals, and Kits.

1000

1001

1002

1003

1004

1005

1006

1008

Supplementary Figure 1. WSB2 interacts with NOXA and promotes its degradation (related to Figure. 1, 2).

(A) GO enrichment analysis of biological processes was performed on the top 100 co-dependent genes of WSB2 in the Broad's 21Q2 DepMap dataset. The biological processes were ranked based on fold enrichment values, with the most significant processes highlighted in red and the less significant processes highlighted in blue, according to log10(FDR) values. In the graph, larger dots represent a higher number of genes involved in a particular process.

(B) Gene effect scores of query gene and the drug selected from Panel 1H. Each dot is a cell line of the selected cancer type(s). Drug sensitivity is represented by log-fold change (PRISM) or IC50 (GDSC), where a more negative value denotes stronger treatment response.

(C) WB analyses of the indicated proteins in WCLs from 293 T cells transfected with the

1020   indicated plasmids.

1021   **(D)** The canonical BC box and CUL5 box sequences of WSB2, as well as the amino acid

1022   deletions corresponding to each mutant, are indicated.

1023   **(E)** WB analyses of the indicated proteins in the WCLs from 293 T cells transfected with the

1024   indicated plasmids.

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

Supplementary Figure 2. Validation of WSB2 knockout in two cancer cell lines.

(A) Schematic of CRISPR/Cas9-mediated knockout of WSB2 by sgRNA#1 or sgRNA#2 in

C4-2B cells. Sanger sequencing confirming that the *WSB2* gene was edited by sgRNA#1 or

sgRNA#2 in C4-2B cells.

(B) Schematic of CRISPR/Cas9-mediated knockout of WSB2 by sgRNA#1 or sgRNA#2 in

Huh-7 cells. Sanger sequencing confirming that the *WSB2* gene was edited by sgRNA#1 or

sgRNA#2 in Huh-7 cells.

1057

1058

1059

1060

Supplementary Figure 3. The combination of WSB2 KO and ABT-737 treatment induces

robust apoptosis in Huh-7 cells (related to Figure 7).

(A) WB analyses of the indicated proteins in WCLs from parental and WSB2 KO Huh-7 cells

treated with DMSO or ABT-737 (20 μM) for 6 h.

(B) Annexin-V-FITC/PI assays were used to stain the harvested cells in (A), of which later flow

cytometry analysis was performed. Data are shown as means ± SD (n = 3). *P* values are

calculated by the Two-way ANOVA test.

(C) IHC analysis of parental and WSB2 KO C4-2B cells from cultures on glass slides using an

anti-DPP9 antibody. Representative staining results are shown. Scale bars, 100 μm.

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

Supplementary Figure 4. Generation and Validation of Wsb2 KO mouse models (related

to Figure 6).

(A) Strategy to generate Wsb2 KO mice using CRISPR/Cas9 methods.

(B) Representative images of male Wsb2$^{-/-}$ and matched Wsb2$^{+/+}$ mice at 3 months of age.

(C) Genotyping analysis of offspring from mating between Wsb2 heterozygous mice at P10. *P*

values are calculated by the Chi-Squared Test.

(D) WB analyses of the indicated proteins in WCL from the indicated tissues of Wsb2$^{+/+}$ and

Wsb2$^{-/-}$ mice.

1090

1091

Supplementary Figure 5. The deficiency of WSB2 enhances the susceptibility of

ABT199-induced apoptosis in the heart and liver tissues of Wsb2 KO mice (related to

Figure 6).

(**A**) Representative IF images from the heart or liver tissues of Wsb2$^{+/+}$ and Wsb2$^{-/-}$ mice after

oral administration of vehicle or ABT-199 (100 mg/kg/day) for 7 days, and stained with

cl-CASP7 and DAPI. Scale bar, 20 µm.

(**B, C**) The mean fluorescence intensity of cl-CASP7 from the heart (B) or liver tissues (C)

obtained from $Wsb2^{+/+}$ and $Wsb2^{-/-}$ mice. Data were shown as means ± SD (n = 5).

(**D**) Representative IF images from the heart or liver tissues of $Wsb2^{+/+}$ and $Wsb2^{-/-}$ mice after

oral administration of vehicle or ABT-199 (100 mg/kg/day) for 7 days, and stained with

cl-PARP1 and DAPI. Scale bar, 20 µm.

(**E, F**) The mean fluorescence intensity of cl-PARP1 from the heart (E) or liver tissues (F)

obtained from $Wsb2^{+/+}$ and $Wsb2^{-/-}$ mice. Data were shown as means ± SD (n = 5).

(**G**) Representative IF images from the heart or liver tissues of $Wsb2^{+/+}$ and $Wsb2^{-/-}$ mice after

oral administration of vehicle or ABT-199 (100 mg/kg/day) for 7 days, and stained with TUNEL

and DAPI. Scale bar, 20 µm.

(**H, I**) The $TUNEL^{+}$ cell number in the heart (H) or liver tissues (I) obtained from $Wsb2^{+/+}$ and

$Wsb2^{-/-}$ mice. Data were shown as means ± SD (n = 5). Data were shown as means ± SD (n =

5).

P values are calculated by the Two-way ANOVA test in (B, C, E, F, H, I).

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127