Navigating the Night: Effects of Artificial Light on the Behaviour of Atlantic Puffin Fledglings

Brown, T.M.[a], S.I. Wilhelm[b], A.D. Slepkov[c], K. Baker[d], G.F. Mastromonaco[e], and G. Burness[d]

Author affiliations

[a]Environmental and Life Sciences Graduate Program, Trent University

[b]Environment and Climate Change Canada

[c]Department of Physics & Astronomy, Trent University

[d]Department of Biology, Trent University

[e]Wildlife Science Division, Toronto Zoo

Name, email, and complete address of the person to whom correspondence should be sent

Taylor M. Brown
taylor.m.brown1@gmail.com
Environmental and Life Sciences Graduate Program
Trent University
Peterborough, ON, Canada
K9L 0G2

Abstract

Every year in Newfoundland, young Atlantic Puffins departing their nests at night for the first
time become stranded in towns near their breeding colonies, a phenomenon thought to be caused
by attraction toward artificial light. To test this hypothesis, we conducted three behavioural
experiments. First, we illuminated beaches near a breeding colony to determine whether more
fledglings would become stranded in illuminated versus dark conditions. Next, we conducted a
Y-maze experiment to test stranded fledglings for phototactic behaviour in general and for
preferences among several different light types. Lastly, we quantified activity levels of stranded
fledglings in an open field test during exposure to several different light types. We found
significantly more fledglings stranded when beaches were illuminated, and fledglings
significantly preferred light over darkness in the Y-maze, supporting our hypothesis that Atlantic
Puffin fledglings become stranded due to light attraction. Fledglings displayed no preferences for
certain light types over others in the Y-maze, indicating that strandings in this species may not be
mitigable by changing the streetlight type in stranding-prone towns. Interestingly, fledglings
exhibited higher activity levels in darkness and High Pressure Sodium light than in LED light,
potentially holding implications for rescue, rehabilitation, and husbandry programs. Overall, our
findings demonstrate that the only evidence-based strategy for the reduction of Atlantic Puffin
strandings remains to be the reduction of coastal artificial lighting; however, further research is
needed to determine if aspects of artificial light besides bulb type may be altered to effectively
reduce light attraction in this species.

Keywords: alcid, ALAN, light attraction, seabird, phototaxis, fledging, behaviour

Introduction

As global human development continues to increase, artificial light at night (ALAN) is of

growing concern as a pollutant of dark nightscapes (Jägerbrand and Spoelstra, 2023). In addition,

the widespread conversion to energy-efficient Light-Emitting Diode lighting is expected to

worsen the ecological impacts of ALAN through a variety of mechanisms (Pawson and Bader,

2014; Davies and Smyth, 2018). Across taxa, disruption of natural diel light cycles by the

presence of ALAN can dysregulate circadian rhythms and lead to endocrinological dysfunction,

behavioural changes, and decreased fitness, among other effects (Sanders et al., 2021; Yang et

al., 2024). One way in which ALAN can alter behaviour, especially that of nocturnal animals, is

by acting as a supernormal visual stimulus that induces phototactic behaviour (either positive or

negative; Rich and Longcore, 2006). Positive phototaxis (or photopositivity), which is often

ascribed to light attraction, has been documented in various animal taxa including insects,

amphibians, fish, sea turtles, migratory songbirds, and seabirds, among others (Rich and

Longcore, 2006; Rodríguez et al., 2017a; Jägerbrand and Spoelstra, 2023; Fabian et al., 2024).

Positive phototaxis toward sources of ALAN can result in mortality due to collisions with glass

windows, lighted structures, and vehicles; stranding and dehydration or desiccation in unfamiliar

locations; and predation of stranded individuals following disorientation and/or collision

(Witherington, 1997; Rich and Longcore, 2006; Rodríguez et al., 2017a; Van Doren et al., 2021).

Seabirds of the orders Procellariiformes and Charadriiformes (family Alcidae) are prone to

becoming stranded in human settlements following phototaxis toward and disorientation by point

sources of ALAN (Rodríguez et al., 2017a; but there are other non-mutually exclusive

hypotheses for stranding; Brown et al., 2023). Fledglings of burrow-nesting and night-fledging

seabird species appear to be the most susceptible to light-induced stranding, although adults of

some species are also affected (Rodríguez et al., 2017a; Wilhelm et al., 2021; Collins et al.,

2023). There is mounting evidence that suppression of ALAN near colonies of burrow-nesting,

night-fledging procellariiform seabirds can reduce the number of stranded fledglings (Reed et al.,

1985; Miles et al., 2010; Rodríguez et al., 2014; Burt et al., 2023). However, it is not always

possible to eliminate or suppress the use of ALAN, especially at industrial and commercial sites

where lighting is necessary for regular operation and human safety. There is thus a growing

interest from industry, non-governmental conservation organizations, and various levels of

government in investigating and implementing alternate "less attractive" light types where

suppression or elimination of ALAN is not possible (e.g., Mercer, 2018). Limited evidence

suggests that lights with high correlated colour temperature (CCT; a relatively crude but simple

and widely-used metric of the perceived colour of a nominal white light source; Durmus, 2022)

and predominantly blue-violet wavelengths (such as metal halide types) may be more attractive

to fledgling procellariiforms and other wildlife than those with low CCT and red-orange

dominant spectra (such as sodium vapour types; Salamolard et al., 2001 as cited in Minatchy,

2004; Rodríguez et al., 2017b; Longcore et al., 2018), but other findings suggest the opposite

(Atchoi et al., 2024) and some indicate no difference (Urmston et al., 2022).

While research interest in procellariiform light attraction has increased in recent years,

comparatively little attention has been given to other seabirds that are similarly affected by

ALAN. Most notably, the Atlantic Puffin (*Fratercula arctica*) represents the only non-

procellariiform seabird species that gets stranded routinely in coastal communities, with insular

Newfoundland (Newfoundland and Labrador, Canada) and Iceland being the best-known

affected locations (Rodríguez et al., 2017a; Wilhelm et al., 2021; Katz, 2023). Adults are diurnal,

but fledglings leave their burrows on maiden flights to the sea at night, unattended by parents,

leaving them susceptible to the influence of ALAN. Although populations in the western Atlantic

remain robust and are even increasing (Wilhelm et al., 2015), those in the eastern Atlantic are

declining, likely due to low recruitment (Miles et al., 2015; Fayet et al., 2021). Thus, with

increasing coastal development, understanding the extent and spectral attributes through which

ALAN impacts post-fledging behaviour in this species is urgently needed.

In Newfoundland, certain well-lit commercial and industrial locations near Atlantic Puffin

breeding colonies are known to consistently yield stranded pufflings and have thus been deemed

stranding "hotspots" (Canadian Parks and Wilderness Society, unpubl. data). The existence of

such "hotspots" has been taken as anecdotal evidence in support of the light attraction hypothesis

of stranding in puffins, although it is unclear to what degree this pattern may be driven by a

spatial bias in non-systematic search effort (e.g., toward well-lit areas, where people are also

presumably more comfortable searching at night). Furthermore, while hundreds of puffins are

stranded annually, this represents only a very small proportion of the total number of individuals

fledging from the colonies (Wilhelm et al. 2021), providing support to the natural random

dispersal hypothesis which suggest that young birds become stranded because of their

inexperience in navigating a new environment (Brown et al., 2023). To date, no experimental

study has been conducted to directly test the light attraction hypothesis and confirm the role of

ALAN in causing Atlantic Puffin strandings. To do so would require a coarse-scale experimental

approach wherein, at a minimum, light levels (including specifically a "dark" condition),

geographic location, and search effort are all tightly controlled (Brown et al., 2023).

At small geographic scales, the behavior of pufflings and other stranding-prone seabird

fledglings exposed to ALAN remains largely unknown but can be studied ex- or in-situ (e.g., in a

Y- or T-maze choice experiment). In the only alcid whose fine-scale phototactic behaviour has

been studied to date, Ancient Murrelet (*Synthliboramphus antiquus*) chicks oriented more often

toward a reflected light source than toward darkness (Gaston et al., 1988). Stranded Cory's

shearwater (*Calonectris borealis*) fledglings, however, consistently chose darkness over light in a

similar experiment (Atchoi et al., 2024). In addition to testing for the presence of photopositivity

in general (i.e., comparing taxic responses to light versus dark), it is also of interest whether

stranding-prone seabird fledglings exhibit fine-scale preferential phototactic responses or

differential activity levels in response to certain spectra of artificial light over others. For

example, Atchoi et al. (2024) found that both fledgling and adult Cory's shearwaters significantly

preferred red over blue light when both were provided as options simultaneously. Collecting this

type of information for more stranding-affected species would supplement our understanding of

seabird phototaxis as well as aid rescue efforts in predicting where stranded seabirds are likely to

go and how they will behave following stranding. Such knowledge should allow for higher

rescue success.

In this study, we tested the effect of ALAN on puffin behaviour at both a coarse and fine

scale. At a coarse scale, we experimentally tested the light attraction hypothesis of stranding in

the field by directing a 30,000 lumen LED light toward the Gull Island (Witless Bay) Atlantic

Puffin breeding colony, on alternate nights from two different dark beaches during the fledging

season. Given the hypothesis that pufflings become stranded by light attraction, we expected that

more pufflings would be observed at the beaches when they were illuminated than when the

beaches were dark. To test for effects of ALAN on a fine scale, we used a Y-maze to evaluate the

propensity of already-stranded pufflings to exhibit phototactic behaviour and preferences for

various artificial light spectra. We expected that fledglings would exhibit positive phototaxis in

the Y-maze by preferentially choosing light over darkness when both were provided as options.

We further expected that when pufflings were exposed to two different artificial light types, they

would preferentially choose those with "cooler" hues (e.g., Cool White LED; Blue LED) over

those with "warmer" hues (e.g., High Pressure Sodium; Warm White LED; Orange LED). To

determine if different spectra of artificial light induce different activity levels in pufflings, we

conducted an open field test. Although we expected to observe differences in activity levels

among the light treatments, particularly when compared with darkness, we could not predict

directionality *a priori*.

Methods

Ethical Note

All procedures involving the use of animals for research were performed in accordance

with Trent University's Animal Care Protocol guidelines under the Canadian Council on Animal

Care (CCAC), protocols #26223 and #28033, and Canadian Wildlife Service scientific permit

#SC4051.

Study Sites and Species

In Newfoundland, most stranded puffin fledglings (hereafter, "pufflings") are found along

a ~15 km section of coastline between the communities of Bay Bulls and Bauline East. This

section forms the focal search area for the "Puffin Patrol", a volunteer-based puffling rescue

program that operates each year during the fledging (and stranding) season (Wilhelm et al., 2013,

2021; Figure 1). Stranded pufflings are thought to largely originate from the breeding colonies on

Gull Island (47.262297°N, 52.773389°W) and Great Island (47.187182°N, 52.813495°W), both

part of the Witless Bay Ecological Reserve and located less than 3 km from shore at their closest

points (Wilhelm et al., 2013). Collectively, the reserve hosts some 304,000 breeding pairs

(Wilhelm et al., 2021), forming the largest concentration in North America (Lowther et al. 2020).

Figure 1. Map of Puffin Patrol search area in Newfoundland, Canada, including all relevant
locations of research activities (Bay Bulls, Lower Pond Beach, Ragged Beach, Saint Michaels)
and Atlantic Puffin breeding colonies (Gull Island, Great Island).

To study the behavioural response of pufflings to light at coarse spatial scales, we

performed an "illuminated beach experiment" (hereafter, beach experiment) in August 2022 at

two sites near Witless Bay, Newfoundland: Lower Pond Beach (47.281464°N, 52.810982°W)

and Ragged Beach (47.262068°N, 52.811644°W; Figure 1). Both of these beaches face Gull

Island: Lower Pond Beach at a distance of approximately 3.1 km, and Ragged Beach at a

distance of approximately 2.5 km.

To study the fine-scale spatial behavioural response of pufflings to lights of differing

spectra, we performed both a Y-maze "choice experiment" and an open field test in August 2021

in Bay Bulls, Newfoundland and Labrador, Canada (47.315178°N, 52.813887°W; Figure 1). In

2021, we conducted the fine-scale experiments in one of two identical "field labs" (storage

trailers, 2.4 x 1.4 x 2.1 m) located adjacent to the headquarters for the Canadian Parks and

Wilderness Society's annual Puffin Patrol rescue program (and the designated drop-off location

for all rescued pufflings). One "lab" was used for the choice experiment, while the other was for

the open field test. In August 2022, we tested one additional choice experiment combination in a

heated building in Saint Michaels (47.187247°N, -52.843990°W; Figure 1).

Response of pufflings to light at a coarse spatial scale

Equipment

We illuminated one of two beaches (Lower Pond beach or Ragged Beach; Figure 1) over

11 nights between 15-27 August 2022 (experiments were not conducted August 17 and 18 due to

inclement weather). At both beaches, an LED light apparatus (Willpower 52-inch, 675-Watt LED

light bar: Xalapa, Veracruz, Mexico) was mounted to a 1.5-metre wooden fence board and placed

on two overturned plastic buckets at the highest point of the beach (approximately 2 metres

above sea level). On each night during the experimental period, we illuminated one beach while

the other remained dark. At the Light beach, the light apparatus was connected to a Discover 12V

35AH/20HR rechargeable deep cycle marine battery (Richmond, B.C., Canada) via a wiring

harness with an in-line fuse and toggle power switch; at the Dark beach the apparatus and

associated equipment was set up but not turned on.

Experimental procedure

A coin was flipped to determine which of the two beaches was illuminated on the first

night of the experiment; thereafter, the Light treatment was alternated between beaches. Because

there was one illuminated beach and one unilluminated beach every night, we have $n = 11$

"Light" and $n = 11$ "Dark" experimental trials. Regardless of treatment (Light or Dark), there

was a team of usually two to three (one instance of four, and one instance of five) biologists and

trained volunteers stationed at each beach to operate the light apparatus and to observe and

capture pufflings for the duration of the experiment. Trials ran for exactly two hours, sometime

between 21:30 and 00:30, depending on the night. On two nights there was an equipment

malfunction in the middle of the trial, but by extending the trial we ensured the total duration of

two hours of light exposure remained the same.

Approximately 10 minutes before the experiment start time, 1-2 team members at each

beach conducted a brief survey to a minimum of 25 metres on either side of the light apparatus,

to ensure no pufflings were already present on the beach prior to the start of the experiment. We

then placed two reflective safety vests 25 metres on either side of the light apparatus and two

vests closer to the water's edge at the same distance (total four vests on each beach) to delineate

the focal observation area ($\sim$750 - 1000 m^2). This distance was chosen based on preliminary tests

of the visual acuity and range of the night vision binoculars used (Bestguarder NV900 4.5-22.5 x

40 HD; Shenzhen, China) and the width of the light beam. There was no way to consistently

delineate the focal observation area over water, but the best effort was made to estimate the

relative distances of pufflings observed.

During the two-hour experiment period on the Light and Dark beaches, observations

principally consisted of one team member at each beach visually scanning the 50-metre-long and

approximately 15- to 20-metre-wide focal beach observation area with night vision binoculars.

The remaining team members scanned the beach, water, and airspace with the unaided eye and

recorded each puffling observation, as well as whether a puffling was suspected or known to be

the same individual as one observed earlier. Observers took turns (of no set duration) using the

night vision binoculars to avoid fatigue.

Any pufflings that became stranded were captured by hand as they were observed and

placed individually into a vented plastic holding container (40 cm long x 30 cm wide x 25 cm

tall) in a quiet, dark and sheltered location. We took care to limit our time on the beach during

capture to a minimum (less than 5 minutes), and only executed a capture attempt when no other

birds were seen nearby (either in the water or on land), so as not to influence their behaviour. At

the end of the evening (maximum time in captivity, three hours), pufflings were measured and

banded by Environment and Climate Change Canada personnel, and released at the water's edge

of the same site where they became stranded (always in dark conditions).

Data extraction

From the recorded observational data, we extracted minimum, maximum, and best-

estimate counts of unique individual pufflings observed each night at each beach. We tabulated

these three separate counts because, despite our best efforts in the field, it was not always

possible to determine if an observed puffling was a "new" individual or one already observed

earlier in the night. This was especially true for swimming individuals. *Minimum counts* for each

beach, on each night were calculated by summing the number of observations known to be of

unique individuals, omitting all observations that were potentially duplicates; this was our most

conservative count. *Maximum counts* were calculated by summing the number of observations

known to be of unique individuals plus all those that were potentially duplicates. *Best-estimate*

counts were calculated by summing the number of observations known to be of unique

individuals plus only the potential duplicates that were (according to qualitative field notes)

likely to be unique individuals; this count is likely the most accurate. We also tabulated on a per-

treatment basis the number of pufflings observed in the water, in the air, and on the beach;

however, it is important to note that because of the difficulty in tracking unique individuals

(especially in the water and in the air, as they are able to quickly move in and out of view), these

are counts of observations rather than estimated counts of unique individuals.

Response of pufflings to light at fine spatial scales

Animal Collection, Housing, and Release

To measure phototactic behaviour and activity levels we mainly used stranded pufflings

provided by the Puffin Patrol between 8-24 August 2021. Puffin Patrol volunteers captured

pufflings by hand using long-handled butterfly nets, placed them individually into vented plastic

holding containers (described above) and then transported them to Puffin Patrol headquarters by

vehicle within 2 hours of capture, but usually sooner. Canadian Parks and Wilderness Society

staff recorded the location and time of capture (if the information was not already submitted

electronically by the volunteer(s)) and placed the crated pufflings in a quiet, dark and sheltered

location.

Each night between ~20:00 and 21:00, we brought the first rescued pufflings in their crates

to a grassy area next to our "field labs," approximately 25 metres away. Pufflings rested in their

crates for a minimum of 10 minutes before being brought into the lab and participating in the

choice experiment described below. Pufflings were then returned to their crates and allowed to

rest outside at ambient temperature for a minimum of 10 minutes before beginning the open field

test in the second field lab. Once the open field test was completed, pufflings were returned to

the Puffin Patrol's designated holding area. All pufflings were then measured, banded, and

released the following morning by Environment and Climate Change Canada personnel.

259   In 2022, we performed one additional choice experiment combination (we did not do an

open field test in 2022). We captured stranded pufflings between 10-14 August 2022 in the same

focal search area as that of the Puffin Patrol, placed individuals in crates, and brought them to

Saint Michaels by vehicle (20 minutes travel). Pufflings rested in their crates in a small, dark,

quiet room (~20°C) for a minimum of 10 minutes before we placed them in the choice

experiment. After the experiment, we returned pufflings to their crates and released them the

following morning from a nearby beach.

Equipment for choice experiment to measure phototactic behaviour

267   We used a Y-maze to measure phototactic behaviour and assess pufflings' preferences for

various light spectra (Figure 2; Figure S1). The metal "acclimation box" opened into the "main

box", a large, opaque, blue Rubbermaid container, via a vertically sliding door. Attached to the

main box were two adjacent and parallel "choice arms" constructed of 15 cm-diameter thick-

walled expandable blue dryer vent hose, each of which led to a small, opaque, blue Rubbermaid

container ("choice box"). In the middle of the terminal end of each choice box we cut a round

hole, approximately 3 cm in diameter and with its centre approximately 8 cm above the ground,

which allowed light from the light source to enter the Y-maze. Except for this hole, we covered

the exterior light-facing side of each choice box with aluminum foil to prevent light from passing

through the wall of the choice box and thereby affecting the spectrum of light visible inside the

Y-maze. Further, we placed a 100 x 50 cm Masonite sheet between the two choice arms and

choice boxes to eliminate crossover of the different light types (except in the Bright HPS vs. Dim

HPS group, where the incident light crossing over from the Bright HPS choice box was used for

the Dim HPS choice box). Another small hole (1 cm in diameter) was cut into the outer-most

sides of both choice boxes to allow an infrared baby monitor camera (VTech, VM5262-2; British

Columbia, Canada) to view the threshold between each choice arm and its corresponding choice

box, allowing us to detect when an individual had made a choice (Figure 2).

Figure 2. View from above the Y-maze apparatus used for our choice experiment. We placed an
individual puffling inside the acclimation box prior to the start of a behavioural trial. The inner
door was then opened, and the bird was allowed to enter and explore the main box, choice arms,
and choice boxes. An infrared camera detected when an individual became visible at the
threshold of a choice box. Note that the diagram is not to scale.

Three different types of lights were used in combination with various filters to produce six

spectral choice combinations (spectra displayed in Figure S1): High Pressure Sodium with a

correlated colour temperature (CCT) of 2100 K ("HPS"; Light EnerG, 400 Watt; manufacturer

location unknown); Warm White Light Emitting Diode (LED) with a CCT of 2700 K (Philips,

5W / 40W-equivalent Soft White LED); and Cool White LED with a CCT of 5000 K (Philips,

5W / 40W-equivalent Daylight LED). In total, there were six choice combinations, with number

of trials for each bracketed: 1) Bright HPS vs Darkness ($n = 15$); 2) Bright HPS vs Dim HPS ($n =$

22); 3) Bright HPS vs Warm White LED ($n = 23$); 4) Warm White LED vs Cool White LED ($n =$

21); 5) Dim Blue LED vs Orange LED ($n = 25$); 6) Bright Blue LED vs Orange LED ($n = 25$, in

2022 only).

When used, LED lights were fixed to a retort stand and positioned approximately 5 cm

from the choice box's distal hole. The HPS light fixture had an attached rectangular metal

reflector on one side, which was bent around the lightbulb to create a cylindrical shape to focus

the light out of the distal end of the cylinder. When HPS light was used, the end of the light

reflector was placed approximately 15 cm away from the hole. These distances were calibrated

with the use of variable neutral density filters (crossed sheets of polyvinyl alcohol-iodine

polarizer films; Alight Polarizers, PF006 linear polarizer; Texas, USA) and diffusers (parchment

paper) to ensure that the amount of light entering the Y-maze at the peak wavelength of each

light type's respective spectrum was uniform and approximately equal across light types, as

measured by a calibrated spectrometer (see Supplementary Material). The two blue LED light

options and the Orange LED light option were created by pairing the Cool White and Warm

White LED lightbulbs, respectively, with neutral density filters and diffusers as well as coloured

acrylic (ePlastics; California, USA) and glass filters (see Supplementary Material). We tested

pufflings in blue and orange LED light in an attempt to distinguish effects of "warm"- and

"cool"-coloured light on phototaxis, following distinctions reported in the literature for other

seabird species (Salamolard et al., 2001 as cited in Minatchy, 2004; Rodríguez et al., 2017b).

Procedure of choice experiment to measure phototactic behaviour

In 2021, 107 pufflings were randomly chosen from those that were stranded and rescued by

the Puffin Patrol and divided among five choice combinations. We tested only one combination

each night via selection out of a hat (i.e., it took five consecutive nights to conduct trials in all

five combinations, after which they were drawn nightly again in random order), except in the last

few days of the field season when we selected the choice combination each night to correct for

sample-size deficiencies. In 2022, 25 pufflings were tested in Bright Blue LED vs Orange LED,

resulting in a total choice experiment sample size of $n = 132$.

Before each trial, the light stimuli were put in their positions and turned on. Each puffling

was brought into the field lab, removed from its holding container, and placed in the acclimation

box of the Y-maze. After precisely 2 minutes the inner sliding door was removed and the puffling

was allowed 13 minutes to make a "choice". If the puffling did not voluntarily exit the

acclimation box by 5 minutes after removal of the inner door, it was gently prodded with the

experimenter's hand reaching through the outer sliding door until the puffling entered the main

box. In total, 20% (27 of 132) of pufflings needed prodding. A choice was considered to have

been made when a bird reached the end of a choice arm and any part of the body was visible on

the monitor (even if it did not fully enter the choice box). The bird was removed from the

apparatus after making a choice or after 15 minutes had elapsed (deemed "No Choice"),

whichever came first, and returned to its holding container in preparation for the open field test.

Between each individual's trial, the entirety of the Y-maze was cleaned with 70% isopropyl alcohol and a disposable cloth to eliminate possible olfactory cues, and the two lights were alternated between the two choice boxes to account for any potential side preference on part of the birds.

Equipment to measure activity levels under various light spectra

The cube-shaped open field test arena was constructed from 6 semi-opaque white twin wall polycarbonate panels (1 m x 1 m), held together by acrylonitrile butadiene styrene (ABS) strips. The floor was left unattached for easy removal and cleaning between trials. A 20 cm x 20 cm grid composed of 25 squares (i.e., 5 squares x 5 squares) was created on the floor using black electrical tape. A flap door (approximately 20 cm wide x 15 cm tall) was cut into the bottom edge of one side of the arena to allow pufflings to be placed into and removed from the apparatus. To eliminate any ambient light, an opaque blackout curtain was placed over the roof and sides of the arena. Two holes were cut in the center of the roof and overlaying curtain: one hole, approximately 3cm in diameter, accommodated the lens of a miniature action camera (Activeon, CCA10W, Activeon Inc.; San Diego, CA, USA) with a wide-angle lens to allow for full view of the floor of the arena. The other hole, approximately the same size and 5 cm away, allowed for a controlled amount of light to enter the arena from an overhanging light source. When in use, LED lights rested directly on the hole, while the HPS light was suspended approximately 20 cm above the hole. This was done to decrease the amount of heat applied to the arena roof (since the HPS light was warmer than the LED lights), and because the HPS light was brighter than the LED lights (qualitatively, this distance resulted in a relatively equal amount of light entering the arena).

Experimental procedure to measure activity levels under various light spectra

To determine whether puffling activity level was affected by light type, we performed an open field test. In 2021, 96 of the 107 Atlantic Puffin fledglings used in the choice experiment were used in the open field test, whereby we exposed them to either darkness or one of three light spectra. Five individuals were removed from the final dataset due to camera malfunction, resulting in a final sample size of $n = 91$. Treatments were: 1) Darkness ($n = 23$); 2) HPS ($n = 19$); 3) Warm White (2700 K) LED ($n = 24$); and 4) Cool White (5000 K) LED ($n = 25$). We did not measure activity in response to blue or orange light. Each night, we used only two light treatments in the open field test, and they were never the same as the two light types used in the Y-maze on the same night. The initial treatment each night was chosen randomly, and treatments were alternated for the rest of the night.

Prior to each open field trial, the appropriate light was set up and turned on (except in the Dark treatment), and a video recording (with audio) was started. The puffling was placed just inside the flap door of the arena, and an electronic timer started. After 10 minutes, the video and audio recording were stopped. The puffling was removed from the arena, replaced in its holding container, and returned to Puffin Patrol headquarters. The floor of the arena was not cleaned between trials unless a puffling defecated during its trial, in which case the affected area was cleaned with 70% isopropyl alcohol and allowed to dry before the next trial. As many trials as possible were completed each night (median 10 trials; range 3-13 trials per night).

Activity level data extraction

Prior to reviewing any open field test videos, we determined the order in which to watch

each video using the random number "Sequence Generator" on random.org (Randomness and

Integrity Services Ltd., Dublin, Ireland). We used behavioural scoring software JWatcher

(Version 1.0, Macquarie University and University of California, Los Angeles) to score two

levels of behaviour for 10 minutes: "mobile" behaviour included walking, running, or flying;

otherwise, birds were considered "immobile". The software automatically calculated the total

amount of time spent mobile and immobile by each puffling in milliseconds, based on the timing

of our key presses, and we subsequently converted each score of time spent mobile into seconds.

In the case of "Dark" treatment videos, only the audio was discernible. Puffins have large,

webbed feet and claws, which allowed us to hear them as they walked over the polycarbonate

floor of the arena. Instead of viewing the file, we listened to its audio and recorded instances

when we heard the bird moving (mobile) or halting (immobile). Billing (tapping or scraping the

arena walls with the bill) and scratching (clawing at the arena floor with the feet) could also be

heard but were not scored as "mobile" behaviour on their own. Each time a key code was

pressed, the previous behaviour was automatically assumed to have ceased.

To determine whether the time spent mobile and immobile in the Dark treatment could be

accurately scored by listening to the audio alone, we selected a subset of 25 videos from the three

lighted conditions (HPS, Warm White LED, Cool White LED) that had already been scored. We

listened to all 25 videos in random order with headphones and scored them audially as normal. A

single observer did all visual and audial scoring.

Statistical Analysis

All analysis was performed in program R (ver. 4.3.0; R Core Team, 2021). We report results to a significance level of $\alpha = 0.05$.

Response of pufflings to light at a coarse spatial scale

We tabulated the *minimum*, *maximum*, and *best-estimate counts* of pufflings seen during the "Light" and "Dark" treatments each night, as well as counts of observations of pufflings in the water, in the air, and on the beach. To control for search effort, we calculated "estimated observation rate" (reported as birds per observer) by dividing the best-estimate counts of pufflings during a given treatment on a given night (over the two-hour experimental period) by the number of observers present. We report medians and interquartile ranges. For a more conservative approach we also compared median estimated observation rates on the beach only (estimated count on the beach divided by number of observers), between Light and Dark. This was because the night vision binoculars worked best while looking at a solid backdrop (i.e., the beach itself) and because our land-based search area at both beaches was standardized. The results of this experiment were definitive, obviating the need for formal statistical analysis.

Response of pufflings to light at fine spatial scales

<u>Phototactic behaviour</u>

In the choice experiment we tested a final sample size of $n = 131$ pufflings. To test whether pufflings displayed evidence for positive phototaxis or preferences for certain light spectra over others we excluded individuals who exhibited No Choice ($n = 23$). For the remaining

individuals, we performed a binomial test (package "*stats*", v. 4.3.0) of the light stimuli chosen in

each of the six choice combinations to determine if choices significantly differed from

hypothesized proportions of 0.5, and we report 95% confidence intervals of the estimated true

proportions. We then conducted a Fisher exact test (package "*stats*", v. 4.3.0) on the number of

pufflings from each combination that were "prodded" to determine if the proportion of prodded

individuals differed among combinations.

<u>Activity levels</u>

To determine if scoring videos aurally may have artificially biased our estimates of time

spent mobile in the "Dark" treatment group (in comparison to scoring the light videos, which

was done visually), we conducted a simple linear regression of time spent mobile scored visually

as a function of time spent mobile scored audially of the 25 selected lighted videos that were

scored both visually and audially. The regression equation was statistically significant ($r^2 =$

0.985; $F(1, 23) = 1478$; $p < 0.001$); audial scores slightly overestimated but nonetheless strongly

predicted visual scores of time spent mobile ($\beta = 0.9228$; $p < 0.001$; Supplementary Figure S2).

We converted the "Dark" data using the regression equation and used these "converted" values in

statistical comparisons with the unmodified visual scores from the other treatment groups.

We identified one outlier in the entire dataset: an individual exposed to Warm White LED

light that spent 353 seconds mobile (a deviation from that treatment group's 75% quartile by

approximately +2.3 times the interquartile range). Because it is unclear what caused this

individual to behave so differently from the rest of the individuals exposed to Warm White LED

light, we removed it from all analyses (both choice experiment and open field test). Retention of

this outlier causes a loss of statistical significance in one out of three comparisons, but the

general trends remain unchanged (to see results with the outlier included, see Supplementary

Material). Because activity level data did not meet assumptions of normality or homogeneity of

variance, we used non-parametric statistics. To test whether individuals differed in the time spent

mobile across treatment groups, we performed a Kruskal-Wallis H test (package "*stats*", v. 4.3.0)

and a post hoc Dunn test (package "*FSA*", v. 0.9.4; Ogle et al., 2023) for multiple comparisons.

Results

Response of pufflings to light at a coarse spatial scale

We calculated a best-estimate total of 136 pufflings that arrived at the beaches during the

Light treatment, and only 2 pufflings during the Dark treatment, over the 11 nights of the

experiment. Similar patterns emerged between the Light and Dark treatments when we expressed

best-estimate count data on a nightly basis (Figure 3A): during the Light treatment the median

best-estimate count was 7.0 (IQR 4.5 – 14.5) individuals per night compared to 0.0 (IQR 0.0 –

0.0) in the Dark. When numbers are compared within the same experimental night, *minimum*,

best estimate, and *maximum* counts of pufflings observed, as well as observation rates that

control for search effort, all tell a similar story: the frequency of pufflings was consistently

higher during Light treatments than during Dark treatments (Supplementary Tables S1, S2).

When we controlled for search effort, median estimated observation rate was 2.0 (IQR 1.2 – 6.0)

birds per observer in the Light and 0.0 (IQR 0.0 – 0.0) birds per observer in the Dark (Figure

3B). Focussing only on the standardized 50-metre-long beach search areas for which we have

greatest confidence, pufflings were observed 54 times during the Light treatment, which greatly

outnumbered the Dark treatment's single observation. Controlling for search effort, a median of

0.7 birds per observer (IQR 0.4 – 2.3) was seen on the beach in the Light treatment, compared to

0.0 birds per observer (IQR 0.0 – 0.0) in the Dark.

Figure 3. Number of fledgling puffins observed per night on artificially lit and dark beaches. A)
Best-estimate total counts of pufflings seen each night during the Dark (black circle) and Light
(yellow circle) treatments, 15-27 August 2022. Note that points overlap on August 26 due to zero
birds being observed during either treatment. B) Observation rates (best-estimate counts of
pufflings observed, divided by the number of observers present) at the two beach sites during
Dark and Light treatments. Each data point represents the observation rate on one experimental
night at each site and points have been offset from one another to improve readability.

Assessment of phototactic behaviour and light preferences

Significantly more individuals chose Bright HPS light than Dark (p = 0.002; estimated true

proportion of Bright HPS, 95% CI: 0.69-1.00; Table 1) in the choice experiment. Although not

statistically significant, more individuals chose Bright HPS over Dim HPS (p = 0.1; estimated

true proportion of Bright HPS, 95% CI: 0.46-0.88). There was no significant preference of any

light type in any other remaining choice combination (Table 1). There was no statistical evidence

for handedness, as pufflings chose Choice Box 1 (n = 57) relatively equally to Choice Box 2 (n =

52; binomial test p = 0.7; estimated true proportion of Choice Box 1, 95% CI: 0.43-0.62). There

was no difference among choice combinations in the number of pufflings prodded out of the

acclimation box after 5 minutes (Fisher Exact Test; p = 0.6).

Table 1. Number of pufflings that chose each of the two light options in each choice
combination. Individual binomial tests assessed evidence for a preference of one light type over
the other in each choice combination (excluding "No choice" individuals), using 0.5 as the
theoretical (null) proportion for each of the two options. We report the 95% confidence intervals
of the true proportion of "Light type 1" as a reference. We also present the number of individuals
that made "No choice" in each combination.

Light type 1	Quantity	Light type 2	Quantity	95% CI	*p*-value	No choice
Bright HPS	10	Dark	0	0.69-1.00	0.002	5
Bright HPS	14	Dim HPS	6	0.46-0.88	0.1	2
Bright HPS	10	Warm White	11	0.26-0.70	1.0	2
Warm White	10	Cool White	10	0.27-0.73	1.0	2
Dim Blue	13	Orange	9	0.36-0.79	0.5	2
Bright Blue	9	Orange	6	0.32-0.84	0.6	10

Assessment of activity levels in different light types

In our open field test, pufflings spent significantly more time mobile in at least one

treatment group compared to the others (Kruskal-Wallis $\chi^2(3)$ = 18.396; p = 0.0004). Pufflings

spent more time mobile in Darkness (median time spent mobile = 120 seconds) and in HPS light

(median 111 seconds) than in Cool White LED light (median 32 seconds; Dunn test p = 0.003

and p = 0.004, respectively; Figure 4). Pufflings also spent more time mobile in Darkness than in

Warm White LED light (median 54 seconds; Dunn test p = 0.045). Pufflings spent more time

mobile, but only marginally, in HPS light compared to Warm White LED light (Dunn test p =

0.050; Figure 4). If the statistical outlier is included (see Methods), the difference between Warm

White LED and Darkness is no longer significant, although the trend remains the same ($p = 0.09$;

see Supplementary Material).

Figure 4. Time spent mobile by fledgling puffins when exposed to each of four light treatment
groups. Figure combines a violin plot and boxplot with overlaid individual data points. Sample
sizes: Warm White LED light ($n = 24$), Cool White LED light ($n = 25$), Darkness ("converted"
data; $n = 23$, see text), and High Pressure Sodium (HPS) light ($n = 19$). The single data point
>300 seconds in the warm LED treatment (outlined in red) was considered a statistical outlier
and was not included in analysis. Individuals were significantly less mobile when exposed to
Cool White LED and Warm White LED than when exposed to Darkness, and also significantly
less mobile in Cool White LED than in HPS.

Discussion

Our results form the first experimental evidence that light attraction behaviour is a causal

mechanism behind the stranding of Atlantic Puffin fledglings (pufflings) in coastal areas during

their initial fledging flights. This evidence emerges at both coarse and fine spatial scales. Even

after controlling for search effort and size of the search area, in our coarse-scale beach

illumination experiiment we found significantly more pufflings stranded during the Light

treatment than during the Dark treatment. Further, consistent with a light attraction hypothesis of

stranding, in our y-maze experiments pufflings significantly preferred light over darkness and

somewhat preferred bright HPS light over dim HPS light. Contrary to our expectation that

pufflings would prefer "cool" light spectra over "warm" light spectra (based on results of

Salamolard et al., 2001 as cited in Minatchy, 2004; Rodríguez et al., 2017b), we found no

preferences when pufflings were simultaneously presented with two different light spectra. Our

open field test experiment, in which we exposed pufflings to either darkness or one of three light

types, revealed that pufflings were more active in Darkness than in any LED light, and more

active in HPS light than under Cool White LED light. Pufflings also tended to be more active in

HPS light than in Warm White LED light, but this difference only bordered on statistical

significance.

We are first to demonstrate a reduction in stranding numbers of an alcid through the

experimental reduction of ALAN. Our finding that more pufflings became stranded at our two

beach sites during the Light treatment compared to the Dark treatment is consistent with similar

studies on procellariiforms (Reed et al., 1985; Miles et al., 2010; Rodríguez et al., 2014; Burt et

al., 2023). Our results therefore provide further support for recommendations to reduce or

eliminate sources of ALAN near breeding colonies of seabirds affected by stranding, especially

where populations are already in decline (e.g., Rodríguez et al., 2017a).

Previously, we expressed concern that human visual perception is more acute, and

therefore the detection of stranded birds is likely more efficient in light conditions compared to

darkness, potentially leading to a bias in detection probability (Brown et al., 2023). As such, in

our coarse-scale light attraction experiment we attempted to reduce the bias between our two

treatments by: 1) using identical night vision binoculars at each of our sites on each night of the

experiment; 2) clearly delineating a search area of the same size (area of approximately 50 m x

15-20 m) on both beaches that took into account the maximum visual range of the binoculars;

and 3) controlling for the number of observers present at each site in our data analysis. Even if

our Light treatment was still subject to bias toward elevated detection probability, it is highly

unlikely that the large difference in number of pufflings found ($n = 54$ in Light vs $n = 1$ in Dark)

is attributable to missing a significant number of pufflings in the Dark treatment. In fact, the

single puffling we detected within the search area during the Dark treatment was on a night with

very low lunar illumination (13.3%), after the moon had already set, and during overcast and

rainy conditions. That pufflings were visible to observers even in the worst of visibility

conditions suggests that if more individuals were indeed present in the Dark treatment, they

would have been detected. Additionally, by alternating each night the site location for Light

treatment, we simultaneously controlled for the potential effects of date, moon phase (see

Wilhelm et al., 2021) and site on the numbers of pufflings we observed.

In addition to the stark difference in number of pufflings observed within the delineated

beach search areas in the Light versus Dark treatment, more individuals were also observed

swimming and flying in the Light treatment ($n = 77$ and $n = 78$, respectively) than in the Dark

treatment ($n = 1$ and $n = 1$).

Support for the light attraction hypothesis of stranding in pufflings is bolstered by the

positive phototactic behaviour pufflings displayed in our choice experiment. Pufflings chose

Bright High Pressure Sodium (HPS) light significantly more often than darkness and chose

Bright HPS light slightly (but insignificantly) more often than Dim HPS. These results agree

with those of a similar study in which Ancient Murrelet chicks chose light significantly more

often than darkness in a T-maze just after leaving their nests (Gaston et al., 1988), but

interestingly are contrary to results of a Y-maze experiment in which Cory's shearwater

fledglings chose darkness over light (Atchoi et al., 2024). Ancient Murrelet chicks leave the nest

at night under parental care on average only two days after hatching (Gaston and Shoji, 2020),

long before they are capable of flight. It is this lack of flight ability, combined with breeding in

mostly remote areas away from human settlement, that presumably precludes them from

stranding in areas with ALAN. That our Atlantic puffin Y-maze results are consistent with those

of Ancient murrelets but not Cory's shearwaters possibly suggests a taxonomic difference in fine-

scale behavioural responses to ALAN between the two main seabird groups affected by light-

induced stranding (i.e., alcids and procellariiforms); however, more research is necessary to

confirm this trend.

Contrary to our expectations, pufflings did not display any preferences for certain light

spectra over others when presented with a choice. We based our expectations on two premises:

first, previous studies have found that attraction behaviour and stranding numbers of fledgling

procellariiforms increase under exposure to short wavelength-dominant (i.e., high CCT) light

types like metal halide (~4500 K CCT) compared to long wavelength-dominant (low CCT) light

types like high pressure sodium (~2000 K CCT; Salamolard et al., 2001 as cited in Minatchy,

2004; Rodríguez et al., 2017b), although this is not always the case (e.g., shielding of long

wavelength light from radiating upward may negate the effect; Urmston et al., 2022; Atchoi et

al., 2024) and evidence is limited. Second, the procellariiform eye is quite sensitive to short

wavelengths (Reed, 1986; Bowmaker et al., 1997; Hart, 2004), implying a potential link between

the wavelengths of highest spectral sensitivity and those that induce the strongest phototactic

responses. Atlantic Puffins likely have similar spectral sensitivities given the similarities in

photoreceptor profiles between auks and procellariiforms (Ödeen and Håstad, 2003), which is

why we expected they would choose blue light over orange and cool white over warm white

LED light in our Y-maze. It is therefore unclear why these choice combinations did not elicit

differences in phototactic behaviour. Perhaps the spectrum of light presented simply has no effect

on the strength of the phototactic response, assuming the spectrum falls enough within the bird's

sensitivity range to be perceived as light at all. Another possible explanation is that the position

of the birds (i.e., standing and walking) in a small experimental apparatus is so contextually

different from flying that it elicits a completely different mode of behaviour as it relates to

lighting conditions. A coarse-scale beach illumination experiment that compares phototactic

responses and stranding numbers across several different light spectra may provide insight in a

more realistic context (as per Rodríguez et al., 2017b).

When we measured the activity levels of pufflings exposed to darkness and various spectra

of artificial light in our open field test, they spent significantly more time active in Darkness than

in Cool or Warm White LED light, and significantly more time active in High Pressure Sodium

(HPS) light than Cool White LED light. It is unclear what factors drove our results, especially in

Darkness. It is possible that in the HPS light treatment the low-level noise of the associated

ballast fan (unlike the LED lights which have no ballast fan) elicited an attractive response,

which has been anecdotally observed in stranded pufflings (Harris, 1982; Harris et al., 1998;

reviewed in Wilhelm et al., 2013). Conversely, it is worth considering the possibility that the

higher activity levels observed in Darkness and in HPS light were typical, and that LED spectra

(which emit more blue light than HPS, even in relatively warm-hued bulb types) have a calming

effect on pufflings that results in reduced activity. Indeed, ducklings reared under blue light are

less active and exhibit a reduced fear response (i.e., tonic immobility) compared with individuals

reared under other light colours (Sultana et al., 2013). Similarly, black-headed buntings

(*Emberiza melanocephala*) experiencing periods of migratory restlessness are less active under

blue light (λ_{peak} = 450 nm) than under red light (λ_{peak} = 640 nm), although these activity patterns

are reversed outside of periods of migratory restlessness (Yadav et al., 2015). Ultimately, the true

cause of the reduced activity levels that we observed in pufflings exposed to LED light,

especially in Cool White LED light, remains a matter of conjecture.

Conservation implications

Globally, there is a concerted effort to reduce the impact of ALAN on wildlife, which

includes efforts to mitigate and prevent the stranding of various seabird species. To date, most

research has focused on procellariiforms, with comparatively little attention paid to the Atlantic

Puffin. We have demonstrated experimentally that the addition of sources of ALAN near Atlantic

Puffin breeding colonies has the potential to drastically increase the number of fledglings that

become stranded, which in turn would almost certainly contribute to decreased juvenile survival.

However, perhaps counter-intuitively, we suggest that light could also be used to lure wayward

puffins away from areas of high mortality risk (i.e., towns and roads) to those of low risk (e.g.,

supervised remote sites) as part of a dynamic rescue effort, and this potential application of

ALAN in a conservation context deserves further research. From fine-scale behavioural

experiments, we demonstrated that Atlantic Puffin fledglings display positive phototactic

behaviour in response to all tested bulb colours, without a preference for either blue- or red-

dominant spectra. This implies that changes to street lighting from historically prevalent High

Pressure Sodium lamps to more energy-efficient Light Emitting Diode, or (if specifically

attempting to reduce attraction) from cool-hued to warm-hued lamps, may not be effective in

reducing strandings in Atlantic Puffins and that the reduction and shielding of ALAN sources

remain the best recommendations. Contrary to what one might have intuitively expected, our

open field test results imply that LED light exposure during captivity (i.e., post-stranding) may

have a calming effect on stranded pufflings, even when compared with darkness. This latter

result has potentially beneficial implications for rescue, rehabilitation, and even husbandry of

puffins and other auks.

Data availability statement: Data are available as Supplementary Material.

References

Atchoi E, M Mitkus, B Machado, V Medeiros, S Garcia, M Juliano, J Bried, and A Rodríguez (2024). Do seabirds dream of artificial lights? Understanding light preferences of Procellariiformes. bioRxiv. https://doi.org/10.1101/2024.03.01.582998

Bowmaker JK, LA Heath, SE Wilkie, and DM Hunt (1997). Visual pigments and oil droplets from six classes of photoreceptor in the retinas of birds. Vision Research 37:2183-2194. https://doi.org/10.1016/S0042-6989(97)00026-6

Brown TM, SI Wilhelm, GF Mastromonaco, and G Burness (2023). A path forward in the investigation of seabird strandings attributed to light attraction. Conservation Science and Practice 5: e12852. https://doi.org/10.1111/csp2.12852

Burt TV, SM Collins, S Green, PB Doiron, SI Wilhelm, and WA Montevecchi (2023). Reduction of coastal lighting decreases seabird strandings. bioRxiv. https://doi.org/10.1101/2023.11.15.567317

Collins SM, A Hedd, WA Montevecchi, TV Burt, DR Wilson, and DA Fifield (2023). Small tube-nosed seabirds fledge on the full moon and throughout the lunar cycle. Biology Letters 19:20230290. http://doi.org/10.1098/rsbl.2023.0290

Davies TW, and T Smyth (2018). Why artificial light at night should be a focus for global change research in the 21st century. Global Change Biology 24:872-882. https://doi.org/10.1111/gcb.13927

Durmus D (2022). Correlated color temperature: use and limitations. Lighting Research & Technology 54:363-375. https://doi.org/10.1177/14771535211034330

Fabian ST, Y Sondhi, PE Allen, JC Theobald, and H-T Lin (2024). Why flying insects gather at artificial light. Nature Communications 15: 689. https://doi.org/10.1038/s41467-024-44785-3

Fayet AL, GV Clucas, T Anker-Nillsen, M Syposz, and ES Hansen (2021). Local prey shortages drive foraging costs and breeding success in a declining seabird, the Atlantic puffin. Journal of Animal Ecology 90:1152-1164. http://doi.org/10.1111/1365-2656.13442

Gaston AJ, and A Shoji (2020). Ancient Murrelet (Synthliboramphus antiquus), version 1.0. In Birds of the World (A. F. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.

Gaston AJ, IL Jones, DG Noble, and SA Smith (1988). Orientation of ancient murrelet, Synthliboramphus antiquus, chicks during their passage from the burrow to the sea. Animal Behaviour 36:300-303. https://doi.org/10.1016/S0003-3472(88)80277-X

Harris MP (1982). Seasonal variation in fledging weight of the puffin *Fratercula arctica*. Ibis 124:100-103.

Harris MP, S Murray, and S Wanless (1998). Long-term changes in breeding performance of puffins *Fratercula arctica* on St Kilda. Bird Study 45(3):371-374. https://doi.org/10.1080/00063659809461109

Hart NS (2004). Microspectrophotometry of visual pigments and oil droplets in a marine bird,
the wedge-tailed shearwater *Puffinus pacificus*: topographic variations in photoreceptor
spectral characteristics. Journal of Experimental Biology 207:1229-1240.

Jägerbrand AK, and K Spoelstra (2023). Effects of anthropogenic light on species and
ecosystems. Science 380: 1125-1130. https://doi.org/10.1126/science.adg3173

Katz C (2023, March 31). *An Icelandic town goes all out to save baby puffins.* Smithsonian
Magazine online. https://www.smithsonianmag.com/science-nature/icelandic-town-goes-
all-out-save-baby-puffins-180981518/

Longcore T, A Rodríguez, B Witherington, JF Penniman, L Herf, and M Herf (2018). Rapid
assessment of lamp spectrum to quantify ecological effects of light at night. Journal of
Experimental Zoology Part A 329:511-521. https://doi.org/10.1002/jez.2184

Lowther PE, AW Diamond, SW Kress, GJ Robertson, K Russell, DN Nettleship, GM Kirwan,
DA Christie, CJ Sharpe, EFJ Garcia, and PFD Boesman (2020). Atlantic Puffin
(*Fratercula arctica*), version 1.0. In Birds of the World (SM Billerman, Editor). Cornell
Lab of Ornithology, Ithaca, NY, USA.

Mercer J (2018, August 26). *N.L. biologist worries offshore lights stranding seabirds.* Saltwire.
https://www.saltwire.com/newfoundland-labrador/news/nl-biologist-worries-offshore-
lights-stranding-seabirds-236694/

Miles W, S Money, R Luxmoore, and RW Furness (2010). Effects of artificial lights and
moonlight on petrels at St Kilda. Bird Study 57:244-251.
https://doi.org/10.1080/00063651003605064

Miles WTS, R Mavor, NJ Riddiford, PV Harvey, R Riddington, DN Shaw, D Parnaby, and JM
Reid (2015). Decline in an Atlantic Puffin population: evaluation of magnitude and
mechanisms. PLoS ONE 10(7): e0131527. http://doi.org/10.1371/journal.pone.0131527

Minatchy N (2004). Stratégie de reduction de la mortalité des petrels induite par les éclairages
publics. Specialized diploma dissertation, University of Réunion, Réunion.

Ödeen A, and Håstad O (2003). Complex distribution of avian color vision systems revealed by
sequencing the SWS1 opsin from total DNA. Molecular Biology and Evolution 20:855-
861. https://doi.org/10.1093/molbev/msg108

Ogle DH, JC Doll, AP Wheeler, and A Dinno (2023). _FSA: Simple Fisheries Stock Assessment
Methods_. R package version 0.9.4. https://CRAN.R-project.org/package=FSA

Pawson SM, and MK-F Bader (2014). LED lighting increases the ecological impact of light
pollution irrespective of color temperature. Ecological Applications 24:1561-1568.
https://doi.org/10.1890/14-0468.1

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reed JR (1986). Seabird vision: spectral sensitivity and light-attraction behavior. PhD thesis,
University of Wisconsin – Madison, Wisconsin, USA.

Reed JR, JL Sincock, and JP Hailman (1985). Light attraction in endangered procellariiform birds: reduction by shielding upward radiation. Auk 102(2):377-383.

Rich C, and T Longcore (Eds.). (2006). *Ecological consequences of artificial night lighting.* Island Press.

Rodríguez A, G Burgan, P Dann, R Jessop, JJ Negro, and A Chiaradia (2014). Fatal attraction of short-tailed shearwaters to artificial lights. PLoS ONE 9(10): e110114. https://doi.org/10.1371/journal.pone.0110114

Rodríguez A, ND Holmes, PG Ryan, K-J Wilson, L Faulquier, Y Murillo, AF Raine, JF Penniman, V Neves, B Rodríguez, JJ Negro, A Chiaradia, P Dann, T Anderson, B Metzger, M Shirai, L Deppe, J Wheeler, P Hodum, C Gouveia, V Carmo, GP Carreira, L Delgado-Alburqueque, C Guerra-Correa, F-X Couzi, M Travers, and M Le Corre (2017a). Seabird mortality induced by land-based artificial lights. Conservation Biology 31(5): 986-1001.

Rodríguez A, P Dann, and A Chiaradia (2017b). Reducing light-induced mortality of seabirds: High pressure sodium lights decrease the fatal attraction of shearwaters. Journal for Nature Conservation 39:68-72.

Salamolard M, F-X Couzi, M Pellerin, and T Ghestemme (2001). Etude complémentaire concernant la protection du Puffin de Baillon dans le cadre de la Route des Tamarins. Rapp. SEOR/DDE-Réunion. 30 pp. + Annexes 4 pp.

Sanders D, E Frago, R Kehoe, C Patterson, and KJ Gaston (2021). A meta-analysis of biological impacts of artificial light at night. Nature Ecology & Evolution 5: 74-81. https://doi.org/10.1038/s41559-020-01322-x

Sultana S, MR Hassan, HS Choe, and KS Ryu (2013). Impact of different monochromatic LED light colours and bird age on the behavioural output and fear response in ducks. Italian Journal of Animal Science 12(4):e94. https://doi.org/10.4081/ijas.2013.e94

Urmston J, KD Hyrenbach, and K Swindle (2022). Quantifying Wedge-tailed shearwater (*Ardenna pacifica*) fallout after changes in highway lighting on southeast O'ahu, Hawai'i. PLoS ONE 17(3): e0265832. https://doi.org/10.1371/journal.pone.0265832

Van Doren BM, DE Willard, M Hennen, KG Horton, EF Stuber, D Sheldon, AH Sivakumar, J Wang, A Farnsworth, and BM Winger (2021). Drivers of fatal bird collisions in an urban center. PNAS 118: e2101666118. https://doi.org/10.1073/pnas.2101666118

Wilhelm SI, JJ Schau, E Schau, SM Dooley, DL Wiseman, and HA Hogan (2013). Atlantic Puffins are attracted to coastal communities in eastern Newfoundland. Northeastern Naturalist 20:624-630.

Wilhelm SI, J Mailhiot, J Arany, JW Chardine, GJ Robertson, and PC Ryan (2015). Update and trends of three important seabird populations in the western North Atlantic using a geographic information system approach. Marine Ornithology 43:211-222.

Wilhelm SI, SM Dooley, EP Corbett, MG Fitzsimmons, PC Ryan, and GJ Robertson (2021). Effects of land-based light pollution on two species of burrow-nesting seabirds in

Newfoundland and Labrador, Canada. Avian Conservation & Ecology 16:12.
https://doi.org/10.5751/ACE-01809-160112

Witherington BE (1997). The problem of photopollution for sea turtles and other nocturnal
animals. In JR Clemmons & R Buchholz (Eds.), *Behavioral approaches to conservation*
in the wild (pp. 303-328). Cambridge University Press.

Yadav G, S Malik, S Rani, and V Kumar (2015). Role of light wavelengths in synchronization of
circadian physiology in songbirds. Physiology & Behavior 140:164-171.
http://dx.doi.org/10.1016/j.physbeh.2014.12.032

Yang Y, Q Liu, C Pan, J Chen, B Xu, K Liu, J Pan, M Lagisz, and S Nakagawa (2024). Species
sensitivities to artificial light at night: a phylogenetically controlled multilevel meta-
analysis on melatonin suppression. Ecology Letters 27: e14387.
https://doi.org/10.1111/ele.14387