1   Navigating the Night: Effects of Artificial Light on the Behaviour of Atlantic Puffin Fledglings

2   Brown, T.M.[a], S.I. Wilhelm[b], A.D. Slepkov[c], K. Baker[d], G.F. Mastromonaco[e], and G. Burness[d]

Author affiliations

4   [a]Environmental and Life Sciences Graduate Program, Trent University

5   [b]Environment and Climate Change Canada

6   [c]Department of Physics & Astronomy, Trent University

7   [d]Department of Biology, Trent University

8   [e]Wildlife Science Division, Toronto Zoo

Name, email, and complete address of the person to whom correspondence should be sent

Taylor M. Brown
taylor.m.brown1@gmail.com
Environmental and Life Sciences Graduate Program
Trent University
Peterborough, ON, Canada
K9L 0G2
+1 519 719 7608

ABSTRACT

Every year in Newfoundland, young Atlantic Puffins (*Fratercula arctica*) departing their nests at night for the first time become stranded in towns near their breeding colonies, a phenomenon thought to be caused by attraction toward artificial light. To test this hypothesis, we conducted three behavioural experiments. First, we illuminated beaches near a breeding colony to determine whether more fledglings would become stranded in illuminated versus dark conditions. Next, we conducted a Y-maze experiment to test stranded fledglings for phototactic behaviour in general and for preferences among High Pressure Sodium (HPS), Warm White Light-Emitting Diode (LED), Cool White LED, Blue LED, and Orange LED light. Lastly, we quantified activity levels of stranded fledglings in an open field test during exposure to several different light types. We found significantly more fledglings stranded when beaches were illuminated, and fledglings significantly preferred light over darkness in the Y-maze, supporting our hypothesis that Atlantic Puffin fledglings become stranded due to light attraction. Fledglings displayed no preferences for certain light types over others in the Y-maze, potentially suggesting that strandings in this species may not be mitigable by changing the streetlight type in stranding-prone towns. Interestingly, fledglings exhibited higher activity levels in darkness and HPS light than in LED light, potentially holding implications for rescue, rehabilitation, and husbandry programs. Overall, our findings demonstrate that the only evidence-based strategy for the reduction of Atlantic Puffin strandings remains to be the reduction of coastal artificial lighting; however, further research is needed to determine if aspects of artificial light besides bulb type may be altered to effectively reduce light attraction in this species.

Keywords: ALAN, alcid, behaviour, fledging, light attraction, phototaxis, seabird

INTRODUCTION

As global human development continues to increase, artificial light at night (ALAN) is of growing concern as a pollutant of dark nightscapes (Jägerbrand and Spoelstra, 2023). In addition, the widespread conversion to energy-efficient Light-Emitting Diode (LED) lighting is expected to worsen the ecological impacts of ALAN through a variety of mechanisms (Pawson and Bader, 2014; Davies and Smyth, 2018). These mechanisms are both physiological (e.g., many organisms are more sensitive to short-wavelength light, which is prevalent in LED outputs, than to the long-wavelength light that dominates light types used previously, such as sodium vapour) and logistical (e.g., greater energy-efficiency allows for increased light production at the same or lower energy cost as other lighting technologies; Pawson and Bader, 2014; Davies and Smyth, 2018) in nature. Across taxa, disruption of natural diel light cycles by the presence of ALAN can

dysregulate circadian rhythms and lead to endocrinological dysfunction, behavioural changes,

and decreased fitness, among other effects (Sanders et al., 2021; Yang et al., 2024). One way in

which ALAN can alter behaviour, especially that of nocturnal animals, is by acting as a

supernormal visual stimulus that induces phototactic behaviour (either positive or negative; Rich

and Longcore, 2006). Positive phototaxis (or photopositivity), which is often ascribed to light

attraction, has been documented in various animal taxa including insects, amphibians, fish, sea

turtles, migratory songbirds, and seabirds, among others (Rich and Longcore, 2006; Rodríguez et

al., 2017a; Jägerbrand and Spoelstra, 2023; Fabian et al., 2024). Positive phototaxis toward

sources of ALAN can result in mortality due to collisions with glass windows, lighted structures,

and vehicles; stranding and dehydration or desiccation in unfamiliar locations; and predation of

stranded individuals following disorientation and/or collision (Witherington, 1997; Rich and

Longcore, 2006; Rodríguez et al., 2017a; Van Doren et al., 2021).

Seabirds of the orders Procellariiformes and Charadriiformes (family Alcidae) are prone to

becoming stranded in human settlements following phototaxis toward and disorientation by

ALAN (Rodríguez et al., 2017a; but there are other non-mutually exclusive hypotheses for

stranding such as simple navigation error during natural dispersal; Brown et al., 2023).

Fledglings of burrow-nesting and night-fledging seabird species appear to be the most

susceptible to light-induced stranding, although adults of some species are also affected

(Rodríguez et al., 2017a; Burt et al., 2024). There is mounting evidence that suppression of

ALAN near colonies of burrow-nesting, night-fledging procellariiform seabirds can reduce the

number of stranded fledglings (Reed et al., 1985; Miles et al., 2010; Rodríguez et al., 2014; Burt

et al., 2024). However, it is not always possible to eliminate or suppress the use of ALAN,

especially at industrial and commercial sites where lighting is necessary for regular operation

and human safety. There is thus a growing interest from industry, non-governmental conservation

organizations, and various levels of government in investigating and implementing alternate

"less attractive" light types where suppression or elimination of ALAN is not possible (e.g.,

Mercer, 2018). Limited evidence suggests that lights with high correlated colour temperature

(CCT; a relatively crude but simple and widely-used metric of the perceived colour of a nominal

white light source; Durmus, 2022) and predominantly blue-violet wavelengths (such as metal

halide types) may be more attractive to fledgling procellariiforms and other wildlife than those

with low CCT and red-orange dominant spectra (such as sodium vapour types; Salamolard et al.,

2001 as cited in Minatchy, 2004; Rodríguez et al., 2017b; Longcore et al., 2018), but other

findings suggest the opposite (Atchoi et al., 2024) and some indicate no difference (Urmston et

al., 2022). An increased attraction toward blue-violet wavelengths may be explained by an

increased sensitivity to these wavelengths by the visual systems of the affected taxa (Reed, 1986;

Bowmaker et al., 1997; Hart, 2004; Pawson and Bader, 2014; Davies and Smyth, 2018).

While research interest in procellariiform light attraction has increased in recent years,

comparatively little attention has been given to other seabirds that are similarly affected by

ALAN. Most notably, the Atlantic Puffin (*Fratercula arctica*) represents the only non-

procellariiform seabird species that gets stranded routinely in coastal communities, with insular

Newfoundland (Newfoundland and Labrador, Canada) and Iceland being the best-known

affected locations (Rodríguez et al., 2017a; Wilhelm et al., 2021; Katz, 2023). Adults are diurnal,

but fledglings leave their burrows on maiden flights to the sea at night, unattended by parents,

leaving them susceptible to the influence of ALAN. Although populations in the western Atlantic

remain robust and are even increasing (Wilhelm et al., 2015), those in the eastern Atlantic are

declining, likely due to low recruitment (Miles et al., 2015; Fayet et al., 2021). Thus, with

increasing coastal development, understanding the extent and spectral attributes through which

ALAN impacts post-fledging behaviour in this species is urgently needed.

In Newfoundland, certain well-lit commercial and industrial locations near Atlantic Puffin

breeding colonies are known to consistently yield stranded puffin fledglings (hereafter,

"pufflings") and have thus been deemed stranding "hotspots" (Canadian Parks and Wilderness

Society – Newfoundland and Labrador Chapter, unpublished data). The existence of such

"hotspots" has been taken as anecdotal evidence in support of the light attraction hypothesis of

stranding in puffins, although it is unclear to what degree this pattern may be driven by a spatial

bias in non-systematic search effort (e.g., toward well-lit areas, where people are also

presumably more comfortable searching at night). Furthermore, while hundreds of puffins are

stranded annually, this represents only a very small proportion of the total number of individuals

fledging from the colonies (Wilhelm et al. 2021), providing support to the natural random

dispersal hypothesis which suggest that young birds become stranded because of their

inexperience in navigating a new environment (Brown et al., 2023). To date, no experimental

study has been conducted to directly test the light attraction hypothesis and confirm the role of

ALAN in causing Atlantic Puffin strandings. To do so would require a coarse-scale experimental

approach wherein, at a minimum, light levels (including specifically a "dark" condition),

geographic location, and search effort are all tightly controlled (Brown et al., 2023).

At small geographic scales, the behavior of pufflings and other stranding-prone seabird

fledglings exposed to ALAN remains largely unknown but can be studied ex- or in-situ (e.g., in a

Y- or T-maze choice experiment). In the only alcid whose fine-scale phototactic behaviour has

been studied to date, Ancient Murrelet (*Synthliboramphus antiquus*) chicks oriented more often

toward a reflected light source than toward darkness (Gaston et al., 1988). Stranded Cory's

shearwater (*Calonectris borealis*) fledglings, however, consistently chose darkness over light in a similar experiment (Atchoi et al., 2024). In addition to testing for the presence of photopositivity in general (i.e., comparing taxic responses to light versus dark), it is also of interest whether stranding-prone seabird fledglings exhibit fine-scale preferential phototactic responses or differential activity levels in response to certain spectra of artificial light over others. For example, Atchoi et al. (2024) found that both fledgling and adult Cory's shearwaters significantly preferred red over blue light when both were provided as options simultaneously. Such fine-scale experiments may not perfectly mimic what occurs in nature, but they nonetheless inform our understanding of behavioural responses to light, especially in ground-based contexts. This information holds implications for rescue efforts in predicting where stranded seabirds are likely to go and how they will behave following stranding, which should allow for higher rescue success.

In this study, we tested the effect of ALAN on puffin behaviour at both a coarse and fine scale. At a coarse scale, we experimentally tested the light attraction hypothesis of stranding in the field by directing a 30 000 lumen LED light toward the Gull Island (Witless Bay) Atlantic Puffin breeding colony, on alternate nights from two different dark beaches during the fledging season. Given the hypothesis that pufflings become stranded by light attraction, we expected that more pufflings would be observed at the beaches when they were illuminated than when the beaches were dark. To test for effects of ALAN on a fine scale, we used a Y-maze to evaluate the propensity of already-stranded pufflings to exhibit phototactic behaviour and preferences for various artificial light spectra. We expected that fledglings would exhibit positive phototaxis in the Y-maze by preferentially choosing light over darkness when both were provided as options. We further expected that when pufflings were exposed to two different artificial light types, they

would preferentially choose those with "cooler" hues (e.g., Cool White LED; Blue LED) over

those with "warmer" hues (e.g., High Pressure Sodium (HPS); Warm White LED; Orange LED).

This reasoning is based on the peak optical sensitivity of closely-related auks to 406 nm violet

light and that of stranding-prone procellariiforms to 402-405 nm violet light (Ödeen and Håstad,

2003). To determine if different spectra of artificial light induce different activity levels in

pufflings, we conducted an open field test. Although we expected to observe differences in

activity levels among the light treatments, particularly when compared with darkness, we could

not predict directionality *a priori*.

METHODS

Ethical Note

All procedures involving the use of animals for research were performed in accordance

with Trent University's Animal Care Protocol guidelines under the Canadian Council on Animal

Care (CCAC), protocols #26223 and #28033, and Canadian Wildlife Service guidelines under

scientific permit # SC4051. All experiments were compliant with current Canadian law. Birds

were captured using handheld mesh nets, were not manipulated beyond simple handling, and

were held in captivity for as short a time as possible. We further attempted to reduce the stress

experienced by birds by housing them individually in quiet and dimly lit conditions for the

duration of captivity.

Study Sites and Species

In Newfoundland, most stranded pufflings are found along a ~15 km section of coastline

between the communities of Bay Bulls and Bauline East (Figure 1). This section forms the focal

search area for the Canadian Parks and Wilderness Society – Newfoundland and Labrador

Chapter's (CPAWS-NL) "Puffin Patrol", a volunteer-based puffling rescue program that

operates each year during the fledging (and stranding) season (Wilhelm et al., 2013, 2021). The

Witless Bay Ecological Reserve, which borders this coastline, hosts close to 400 000 breeding

pairs of Atlantic Puffins, forming the largest concentration in North America (Lowther et al.,

2020; SIW, unpublished data). Stranded pufflings are thought to largely originate from the two

largest breeding colonies on Gull Island (47.262297°N, 52.773389°W, estimated at 168 000

pairs; SIW, unpublished data) and Great Island (47.187182°N, 52.813495°W, estimated at 204

000 pairs; SIW, unpublished data), both part of the reserve and located less than 3 km from shore

at their closest points (Figure 1).

Figure 1. Map of Puffin Patrol search area in Newfoundland, Canada, including all relevant locations of research activities (Bay Bulls, Lower Pond Beach, Ragged Beach, Saint Michaels) and the two largest Atlantic Puffin breeding colonies (Gull Island, Great Island).

To study the behavioural response of pufflings to light at coarse spatial scales, we performed an "illuminated beach experiment" (hereafter, beach experiment) in August 2022 at two sites near Witless Bay, Newfoundland: Lower Pond Beach (47.281464°N, 52.810982°W) and Ragged Beach (47.262068°N, 52.811644°W; Figure 1). Both of these beaches face Gull Island: Lower Pond Beach at a distance of approximately 3.1 km, and Ragged Beach at a distance of approximately 2.5 km.

To study the fine-scale spatial behavioural response of pufflings to lights of differing spectra, we performed both a Y-maze "choice experiment" and an open field test in August 2021

in Bay Bulls, Newfoundland and Labrador, Canada (47.315178°N, 52.813887°W; Figure 1). In

2021, we conducted the fine-scale experiments in one of two identical "field labs" (storage

trailers, 2.4 × 1.4 × 2.1 m) located adjacent to the headquarters for CPAWS-NL's annual Puffin

Patrol rescue program (and the designated drop-off location for all rescued pufflings). One "lab"

was used for the choice experiment, while the other was for the open field test. In August 2022,

we tested one additional choice experiment combination in a building in Saint Michaels

(47.187247°N, 52.843990°W; Figure 1).

Response of Pufflings to Light at a Coarse Spatial Scale

Materials

We illuminated one of two beaches (Lower Pond beach or Ragged Beach; Figure 1) over

11 nights during the fledging season between 15-27 August 2022, coinciding with a transition

from 85% to 0% lunar illumination (moongiant.com). Experiments were not conducted August

17 and 18 due to inclement weather. At both beaches, a 30 000 lumen, 6390 K LED light

apparatus (Willpower 52-inch, 675-Watt LED light bar: Xalapa, Veracruz, Mexico; spectrum

displayed in Appendix Figure A1) was mounted to a 1.5-metre wooden fence board and placed

on two overturned plastic buckets at the highest point of the beach (approximately 2 metres

above sea level) such that the light sat horizontal and parallel to the beach, facing Gull Island. On

each night during the experimental period, we illuminated one beach while the other remained

dark. At the Light beach, the light apparatus was connected to a Discover 12V 35AH/20HR

rechargeable deep cycle marine battery (Richmond, B.C., Canada) via a wiring harness with an

in-line fuse and toggle power switch; at the Dark beach the battery and the light apparatus with

attached wiring harness were set up but not connected.

Experimental procedure

A coin was flipped to determine which of the two beaches was illuminated on the first

night of the experiment; thereafter, the Light treatment was alternated between beaches. Because

there was one illuminated beach and one unilluminated beach every night, we have $N = 11$

"Light" and $N = 11$ "Dark" experimental trials. Regardless of treatment (Light or Dark), there

was a team of usually two to three (one instance of four, and one instance of five) biologists and

trained volunteers stationed at each beach to operate the light apparatus and to observe and

capture pufflings for the duration of the experiment. Trials ran for exactly two hours, sometime

between 2130 and 0030 hours (to coincide with the peak stranding period; CPAWS-NL,

unpublished data), depending on the night. On two nights there was an equipment malfunction in

the middle of the trial, but by extending the Light treatment trial we ensured the total duration of

two hours of light exposure remained the same.

Approximately 10 minutes before the experiment start time, 1-2 team members at each

beach conducted a brief survey to a minimum of 25 metres on either side of the light apparatus

and scanned the water with the unaided eye and night vision binoculars (Bestguarder NV900 4.5-

22.5 × 40 HD; Shenzhen, China) as thoroughly as allowable by the weather conditions, to ensure

no pufflings were already present in the vicinity prior to the start of the experiment. We then

placed two reflective safety vests 25 metres on either side of the light apparatus and two vests

closer to the water's edge at the same distance (total four vests on each beach) to delineate the

focal observation area (50 metres long and varying from 15-20 metres wide = ~750-1000 m^2).

This distance was chosen based on preliminary tests of the resolution and range of the night

vision binoculars and the width of the light beam. There was no way to consistently delineate the

focal observation area over water, but the best effort was made to estimate the relative distances

of pufflings observed.

During the two-hour experiment period on the Light and Dark beaches, observations

principally consisted of one team member at each beach visually scanning the focal beach

observation area with night vision binoculars; however, it should be noted that on most nights,

even the unaided eye could resolve details of the beach topography (and therefore also

presumably resolve the stark white chests of any pufflings present) with ambient light alone.

Observers took turns (of no set duration) using the night vision binoculars to avoid fatigue. The

remaining team members scanned the beach, water, and airspace with the unaided eye and

recorded each puffling observation, including notes on whether a puffling was suspected or

known to be the same individual as one observed earlier. Team members stood next to one

another and visually tracked each individual puffling as long as it was within view, continuously

communicating about positions and movements of birds to keep track of unique individuals.

Whenever visual contact with a puffling was lost, subsequent visual contact with a puffling in the

same general location was judged in its likelihood of being the same individual based on the

amount of time that had elapsed since previous visual contact. If less than ~10 seconds elapsed

between initial and subsequent visual contact, the second observation was considered to be the

same bird; otherwise, it was recorded as a "potential duplicate".

Any pufflings that became stranded were captured by hand as they were observed and

placed individually into a vented plastic holding container (40 cm long × 30 cm wide × 25 cm

tall) in a quiet, dark and sheltered location. We took care to limit our time on the beach during

capture to a minimum (less than 5 minutes), and only executed a capture attempt when no other

birds were seen nearby (either in the water or on land), so as not to influence their behaviour.

During a capture attempt, the remaining observer(s) divided their attention amongst scanning the

beach, water, and airspace with both night vision binoculars and the unaided eye to ensure all

new puffling observations were documented. At the end of the evening (maximum time in

captivity, three hours), pufflings were measured and banded by Environment and Climate

Change Canada personnel, and released at the water's edge of the same site where they became

stranded (always in dark conditions).

Data extraction

From the recorded observational data, we extracted minimum, maximum, and best-

estimate counts of unique individual pufflings observed each night at each beach. *Minimum*

counts for each beach, on each night were calculated by summing the number of observations

known to be of unique individuals, omitting all observations that were potentially duplicates; this

was our most conservative count. *Maximum counts* were calculated by summing the number of

observations known to be of unique individuals plus all those that were potentially duplicates.

Best-estimate counts were calculated by summing the number of observations known to be of

unique individuals plus only the potential duplicates that were (according to qualitative field

notes) likely to be unique individuals; this count is likely the most accurate. On six of eleven

nights, the number of pufflings present at the illuminated beach at any one time was low enough

to facilitate continuous visual tracking of all or nearly all individuals, resulting in few (if any)

potential duplicates and minimum, maximum, and best-estimate counts that vary little from one

another. On the other five nights, numbers were high enough that continuous visual tracking of

all individuals was not possible despite our best efforts; this is when the number of potential

duplicates was highest, and the greater difference between minimum and maximum counts on

these nights reflects this difficulty. We also tabulated on a per-treatment basis the number of

pufflings observed in the water, in the air, and on the beach; however, it is important to note that

because of the difficulty in tracking unique individuals (see above), these are counts of

observations rather than estimated counts of unique individuals.

Response of Pufflings to Light at Fine Spatial Scales

Animal collection, housing, and release

To measure phototactic behaviour and activity levels we mainly used stranded pufflings

provided by the Puffin Patrol between 8-24 August 2021. Although it is possible that stranded

pufflings do not represent a truly random sample of the entire fledgling population, it is this

population in which we are most interested since it appears to be most affected by ALAN and

also the population from which we would expect the greatest response to our experimental light

stimuli. Puffin Patrol volunteers captured pufflings by hand using long-handled butterfly nets,

placed them individually into vented plastic holding containers (described above) and then

transported them to Puffin Patrol headquarters by vehicle within 2 hours of capture, but usually

sooner. CPAWS-NL staff recorded the location and time of capture (if the information was not

already submitted electronically by the volunteer(s)) and placed the crated pufflings in a quiet,

dark and sheltered location.

Each night between ~2000 and 2100 hours, we brought the first rescued pufflings in their

crates to a grassy area next to our "field labs," approximately 25 metres away. Pufflings rested in

their crates for a minimum of 10 minutes before being brought into the lab and participating in

the choice experiment described below. Pufflings were then returned to their crates and allowed

to rest outside at ambient temperature for a minimum of 10 minutes before beginning the open

field test in the second field lab. Once the open field test was completed, pufflings were returned

to the Puffin Patrol's designated holding area. All pufflings were then measured, banded, and

released the following morning by Environment and Climate Change Canada personnel.

In 2022, we performed one additional choice experiment combination (we did not do an

open field test in 2022). We captured stranded pufflings between 10-14 August 2022 in the same

focal search area as that of the Puffin Patrol, placed individuals in crates, and brought them to

Saint Michaels by vehicle (20 minutes travel). Pufflings rested in their crates in a small, dark,

quiet room (~20°C) for a minimum of 10 minutes before we placed them in the choice

experiment. After the experiment, we returned pufflings to their crates and released them the

following morning from a nearby beach.

Materials for choice experiment to measure phototactic behaviour

We used a Y-maze to measure phototactic behaviour and assess pufflings' preferences for

various light spectra (Figure 2; Appendix Figure A2). The metal "acclimation box" (height: 15

cm) opened into the "main box" (height: 22 cm), a large, opaque, blue Rubbermaid container, via

a vertically sliding door. Attached to the main box were two adjacent and parallel "choice arms"

constructed of 15 cm-diameter thick-walled expandable blue dryer vent hose, each of which led

to a small, opaque, blue Rubbermaid container ("choice box"; height: 18 cm). In the middle of

the terminal end of each choice box we cut a round hole, approximately 3 cm in diameter and

with its centre approximately 8 cm above the ground, which allowed light from the light source

to enter the Y-maze. Except for this hole, we covered the exterior light-facing side of each choice

box with aluminum foil to prevent light from passing through the wall of the choice box and

thereby affecting the spectrum of light visible inside the Y-maze. Further, we placed a 100×50

cm Masonite sheet between the two choice arms and choice boxes to eliminate crossover of the

different light types (except in the Bright HPS vs. Dim HPS group, where the incident light

crossing over from the Bright HPS choice box was used for the Dim HPS choice box). Another

small hole (1 cm in diameter) was cut into the outer-most sides of both choice boxes to allow an

infrared baby monitor camera (VTech, VM5262-2; British Columbia, Canada) to view the

doorway between each choice arm and its corresponding choice box, allowing us to detect when

an individual had made a choice (Figure 2).

Figure 2. View from above the Y-maze apparatus used for our choice experiment. We placed an
individual puffling inside the acclimation box prior to the start of a behavioural trial. The inner
door was then opened, and the bird was allowed to enter and explore the main box, choice arms,
and choice boxes. An infrared camera detected when an individual became visible at the
doorway of a choice box. Note that the diagram is not to scale.

Three different types of lights were used in combination with various filters to produce six

spectral choice combinations (spectra displayed in Appendix Figure A2): HPS with a correlated

colour temperature (CCT) of 2100 K (Light EnerG, 400 Watt; manufacturer location unknown);

Warm White LED with a CCT of 2700 K (Philips, 5W / 40W-equivalent Soft White LED); and

Cool White LED with a CCT of 5000 K (Philips, 5W / 40W-equivalent Daylight LED). In total,

there were six choice combinations, with number of trials for each bracketed: 1) Bright HPS vs

Darkness ($N = 15$); 2) Bright HPS vs Dim HPS ($N = 22$); 3) Bright HPS vs Warm White LED (N

$= 23$); 4) Warm White LED vs Cool White LED ($N = 21$); 5) Dim Blue LED vs Orange LED (N

$= 25$); 6) Bright Blue LED vs Orange LED ($N = 25$, in 2022 only).

When used, LED lights were fixed to a retort stand and positioned approximately 5 cm

from the choice box's distal hole. The HPS light fixture had an attached rectangular metal

reflector on one side, which was bent around the lightbulb to create a cylindrical shape to focus

the light out of the distal end of the cylinder. When HPS light was used, the end of the light

reflector was placed approximately 15 cm away from the hole. These distances were calibrated

with the use of variable neutral density filters (crossed sheets of polyvinyl alcohol-iodine

polarizer films; Alight Polarizers, PF006 linear polarizer; Texas, USA) and diffusers (parchment

paper) to ensure that the amount of light entering the Y-maze at the peak wavelength of each

light type's respective spectrum was approximately equal across light types, as measured by a

calibrated spectrometer (see Appendix Figure A2). The two blue LED light options and the

Orange LED light option were created by pairing the Cool White and Warm White LED

lightbulbs, respectively, with neutral density filters and diffusers as well as coloured acrylic

(ePlastics; California, USA) and glass filters (see Appendix Table A3). We tested pufflings in

blue and orange LED light in an attempt to distinguish effects of "warm"- and "cool"-coloured

light on phototaxis, following distinctions reported in the literature for other seabird species

(Salamolard et al., 2001 as cited in Minatchy, 2004; Rodríguez et al., 2017b).

Procedure of choice experiment to measure phototactic behaviour

In 2021, 107 pufflings were randomly chosen from those that were stranded and rescued by

the Puffin Patrol and divided among five choice combinations. We tested only one combination

each night via selection out of a hat (i.e., it took five consecutive nights to conduct trials in all

five combinations, after which they were drawn nightly again in random order), except in the last

few days of the field season when we selected the choice combination each night to correct for

sample-size deficiencies (all choice combinations tested over a minimum of two non-sequential

nights). In 2022, 25 pufflings were tested in Bright Blue LED vs Orange LED, resulting in a total

choice experiment sample size of $N = 132$.

Before each trial, the light stimuli were put in their positions and turned on. Each puffling

was brought into the field lab, removed from its holding container, and placed in the acclimation

box of the Y-maze. After precisely 2 minutes the inner sliding door was removed and the puffling

was allowed 13 minutes to make a "choice". If the puffling did not voluntarily exit the

acclimation box by 5 minutes after removal of the inner door, it was gently prodded with the

experimenter's hand reaching through the outer sliding door until the puffling entered the main

box. In total, 20% (27 of 132) of pufflings needed prodding. A choice was considered to have

been made when a bird reached the end of a choice arm and any part of the body was visible on

the monitor (even if it did not fully enter the choice box). The bird was removed from the

apparatus after making a choice or after 15 minutes had elapsed (deemed "No Choice"),

whichever came first, and returned to its holding container in preparation for the open field test.

Between each individual's trial, the entirety of the Y-maze was cleaned with 70% isopropyl

alcohol and a disposable cloth to eliminate possible olfactory cues, and the two lights were

alternated between the two choice boxes to account for any potential side bias on part of the

birds.

Materials to measure activity levels under various light spectra

The cube-shaped open field test arena was constructed from 6 semi-opaque white twin wall

polycarbonate panels (1 m × 1 m), held together by acrylonitrile butadiene styrene (ABS) strips.

The floor was left unattached for easy removal and cleaning between trials. A 20 cm × 20 cm

grid composed of 25 squares (i.e., 5 squares × 5 squares) was created on the floor using black

electrical tape. A flap door (approximately 20 cm wide × 15 cm tall) was cut into the bottom

edge of one side of the arena to allow pufflings to be placed into and removed from the

apparatus. To eliminate any ambient light, an opaque blackout curtain was placed over the roof

and sides of the arena. Two holes were cut in the center of the roof and overlaying curtain: one

hole, approximately 3 cm in diameter, accommodated the lens of a miniature action camera

(Activeon, CCA10W, Activeon Inc.; San Diego, CA, USA) with a wide-angle lens to allow for

full view of the floor of the arena. The other hole, approximately the same size and 5 cm away,

allowed for a controlled amount of light to enter the arena from an overhanging light source.

When in use, LED lights rested directly on the hole, while the HPS light was suspended

approximately 20 cm above the hole. This was done to decrease the amount of heat applied to the

arena roof (since the HPS light was warmer than the LED lights), and because the HPS light was

brighter than the LED lights (this distance resulted in a relatively equal amount of light entering

the arena). Median illuminance was similar across the three light treatments when measured at

25 different locations at floor level (Kruskal-Wallis test: $H_2 = 1.3696$, $P = 0.5$) using a

spectrometer (SRI-2000 Spectral Light Meter; Allied Scientific Pro; Gatineau, Quebec, Canada).

The median illuminance from HPS was 26.35 lux (IQR 21.08 – 33.18), while that from Warm

White LED was 24.34 lux (IQR 20.30 – 27.80), and that from Cool White LED was 23.88 lux

(IQR 19.46 – 28.62). These median illuminance values are comparable to that of a similar

behavioural study, wherein illuminance was approximately 30 lux (Atchoi et al., 2023).

Experimental procedure to measure activity levels under various light spectra

To determine whether puffling activity level was affected by light type, we performed an

open field test. In 2021, 96 of the 107 Atlantic Puffin fledglings used in the choice experiment

were used in the open field test, whereby we exposed them to either darkness or one of three

light spectra. Five individuals were removed from the final dataset due to camera malfunction,

resulting in a final sample size of $N = 91$. Treatments were: 1) Darkness ($N = 23$); 2) HPS ($N =$

19); 3) Warm White (2700 K) LED ($N = 24$); and 4) Cool White (5000 K) LED ($N = 25$). We did

not measure activity in response to blue or orange light to focus our testing on spectra from light

sources currently used in human settlements (i.e., HPS and LED). Each night, we used only two

light treatments in the open field test, and they were never the same as the two light types used in

the Y-maze on the same night. The initial treatment each night was chosen randomly, and

treatments were alternated for the rest of the night.

Prior to each open field trial, the appropriate light was set up and turned on (except in the

Dark treatment), and a video recording (with audio) was started. The puffling was placed just

inside the flap door of the arena, and an electronic timer started. After 10 minutes, the video and

audio recording were stopped. The puffling was removed from the arena, replaced in its holding

container, and returned to Puffin Patrol headquarters. The floor of the arena was not cleaned

between trials unless a puffling defecated during its trial, in which case the affected area was

cleaned with 70% isopropyl alcohol and allowed to dry before the next trial. As many trials as

possible were completed each night (median 10 trials; range 3-13 trials per night).

Activity level data extraction

Prior to reviewing any open field test videos, we determined the order in which to watch

each video using the random number "Sequence Generator" on random.org (Randomness and

Integrity Services Ltd., Dublin, Ireland). We used behavioural scoring software JWatcher

(Version 1.0, Macquarie University and University of California, Los Angeles) to score two

levels of behaviour for 10 minutes: "mobile" behaviour included walking, running, or flying;

otherwise, birds were considered "immobile". The software automatically calculated the total

amount of time spent mobile and immobile by each puffling in milliseconds, based on the timing

of our key presses, and we subsequently converted each score of time spent mobile into seconds.

In the case of "Dark" treatment videos, only the audio was discernible. Puffins have large,

webbed feet and claws, which allowed us to hear them as they walked over the polycarbonate

floor of the arena. Instead of viewing the file, we listened to its audio and recorded instances

when we heard the bird moving (mobile) or halting (immobile). Billing (tapping or scraping the

arena walls with the bill) and scratching (clawing at the arena floor with the feet) could also be

heard but were not scored as "mobile" behaviour on their own. Each time a key code was

pressed, the previous behaviour was automatically assumed to have ceased. We decided to score

only "Dark" videos audially (rather than scoring all videos audially) because: 1) we reasoned that

visual scoring is both more accurate (i.e., truer to the correct value) and more precise (i.e.,

repeatable) than audial scoring, and is therefore preferable when three of our four treatments can

be scored visually; 2) the noise associated with the HPS light's ballast fan would have made

audial scoring of those videos difficult and therefore introduce a new source of scoring bias; and

3) extra care was taken during the Dark treatment specifically to ensure that noises similar in

quality to those of puffling footsteps were reduced to an absolute minimum so as to increase the

accuracy of those audial scores, whereas only reasonable care was taken to reduce similar noises

in the lighted treatments.

To determine whether the time spent mobile and immobile in the Dark treatment could be

accurately scored by listening to the audio alone, we selected a subset of 25 videos from the three

lighted conditions (HPS, Warm White LED, Cool White LED) that had already been scored. We

listened to all 25 videos in random order with headphones and scored them audially as normal. A

single observer did all visual and audial scoring.

Statistical Analysis

All analysis was performed in program R (ver. 4.3.0; R Core Team, 2021). We report

results to a significance level of $\alpha = 0.05$.

Response of pufflings to light at a coarse spatial scale

We tabulated the *minimum*, *maximum*, and *best-estimate counts* of pufflings seen during

the "Light" and "Dark" treatments each night, as well as counts of observations of pufflings in

the water, in the air, and on the beach. To control for search effort, we calculated "estimated

observation rate" (reported as birds per observer) by dividing the best-estimate counts of

pufflings during a given treatment on a given night (over the two-hour experimental period) by

the number of observers present. We report medians and interquartile ranges. For a more

conservative approach we also compared median estimated observation rates on the beach only

(estimated count on the beach divided by number of observers), between Light and Dark. This

was because the night vision binoculars worked best while looking at a solid backdrop (i.e., the

beach itself) and because our land-based search area at both beaches was standardized. The

results of this experiment were definitive in demonstrating higher numbers of pufflings at the

beaches during the "Light" treatment regardless of the metric used, obviating the need for formal

statistical analysis.

Response of pufflings to light at fine spatial scales

Phototactic behaviour

In the choice experiment we tested a final sample size of $N = 131$ pufflings. To test whether

pufflings displayed evidence for positive phototaxis or preferences for certain light spectra over

others we excluded individuals who exhibited No Choice ($N = 23$) since this behavioural

outcome indicates no preference. For the remaining individuals, we performed a binomial test

(package "*stats*", v. 4.3.0) of the light stimuli chosen in each of the six choice combinations to

determine if choices significantly differed from hypothesized proportions of 0.5, and we report

95% confidence intervals of the estimated true proportions. We then conducted a Fisher exact

test (package "*stats*", v. 4.3.0) on the number of pufflings from each combination that were

"prodded" to determine if the proportion of prodded individuals differed among combinations.

Activity levels

To determine if scoring videos audially may have artificially biased our estimates of time

spent mobile in the "Dark" treatment group (in comparison to scoring the light videos, which

was done visually), we conducted a simple linear regression of time spent mobile scored visually

as a function of time spent mobile scored audially of the 25 selected lighted videos that were

scored both visually and audially. The regression equation was statistically significant ($r^2 =$

0.985, $F_{1,23} = 1478$, $P < 0.001$); audial scores slightly overestimated but nonetheless strongly

predicted visual scores of time spent mobile ($\beta = 0.9228$, $P < 0.001$, Appendix Figure A3). We

converted the "Dark" data using the regression equation and used these "converted" values in

statistical comparisons with the unmodified visual scores from the other treatment groups (see

Appendix). Note that the average difference between raw and converted scores of time spent

mobile in the "Dark" group was a decrease of only eight seconds (maximum difference, -21

seconds).

We identified one outlier in the entire dataset: an individual exposed to Warm White LED

light that spent 353 seconds mobile (a deviation from that treatment group's 75% quartile by

approximately +2.3 times the interquartile range). Because it is unclear what caused this

individual to behave so differently from the rest of the individuals exposed to Warm White LED

light, we removed it from all analyses (both choice experiment and open field test). Retention of

this outlier causes a loss of statistical significance in one out of three comparisons, but the

general trends remain unchanged (to see results with the outlier included, see Appendix).

Because activity level data did not meet assumptions of normality or homogeneity of variance,

we used non-parametric statistics. To test whether individuals differed in the time spent mobile

across treatment groups, we performed a Kruskal-Wallis H test (package "*stats*", v. 4.3.0) and a

post hoc Dunn test (package "*FSA*", v. 0.9.4; Ogle et al., 2023) for multiple comparisons.

RESULTS

Response of Pufflings to Light at a Coarse Spatial Scale

We calculated a best-estimate total of 136 pufflings that arrived at the beaches (observed in

the water, on the beach, and in the air combined) during the Light treatment, and only 2 pufflings

during the Dark treatment, over the 11 nights of the experiment. Similar patterns emerged

between the Light and Dark treatments when we expressed best-estimate count data on a nightly

basis (Figure 3A): during the Light treatment the median best-estimate count was 7.0 (IQR 4.5 –

14.5) individuals per night compared to 0.0 (IQR 0.0 – 0.0) in the Dark. When numbers are

compared within the same experimental night, *minimum*, *best estimate*, and *maximum* counts of

pufflings observed, as well as observation rates that control for search effort, all tell a similar

story: the frequency of pufflings was consistently higher during Light treatments than during

Dark treatments (Appendix Tables A1, A2). When we controlled for search effort, median

estimated observation rate was 2.0 (IQR 1.2 – 6.0) birds per observer in the Light and 0.0 (IQR

0.0 – 0.0) birds per observer in the Dark (Figure 3B). Focussing only on the standardized 50-

metre-long beach search areas for which we have greatest confidence, pufflings were observed

54 times during the Light treatment, which greatly outnumbered the Dark treatment's single

observation. Controlling for search effort, a median of 0.7 birds per observer (IQR 0.4 – 2.3) was

seen on the beach in the Light treatment, compared to 0.0 birds per observer (IQR 0.0 – 0.0) in

the Dark. Behavioural observations indicated that pufflings in flight generally flew at altitudes of

15 metres or less, in straight lines either along the shoreline (in both directions) or overhead

going inland, and pufflings in the water and on the beach approached the light to varying

distances (approximately 15-40 metres) and then either stopped moving (to preen, rest, etc.) or

proceeded back to the water and swam away (if not captured first).

Figure 3. Number of fledgling puffins observed per night at artificially lit and dark beaches. A)
Best-estimate total counts of pufflings (in the water, on the beach, and in the air combined) seen
each night during the Dark (black circle) and Light (yellow circle) treatments, 15-27 August
2022. Note that points overlap on August 26 due to zero birds being observed during either
treatment. B) Observation rates (same best-estimate counts of pufflings observed, divided by the
number of observers present) at the two beach sites during Dark and Light treatments. Each data
point represents the observation rate on one experimental night at each site and points have been
offset from one another to improve readability.

Assessment of Phototactic Behaviour and Light Preferences

Significantly more individuals chose Bright HPS light than Dark ($P = 0.002$; 95% CI of

estimated true proportion of Bright HPS: 0.69-1.00; Table 1) in the choice experiment. Although

not statistically significant, more individuals chose Bright HPS over Dim HPS ($P = 0.1$; 95% CI

of estimated true proportion of Bright HPS: 0.46-0.88). There was no significant preference of

any light type in any other remaining choice combination (Table 1). There was no statistical

evidence for side bias, as pufflings chose Choice Box 1 ($N = 57$) relatively equally to Choice

Box 2 ($N = 52$; binomial test: $P = 0.7$; 95% CI of estimated true proportion of Choice Box 1:

0.43-0.62). There was no difference among choice combinations in the number of pufflings

prodded out of the acclimation box after 5 minutes (Fisher Exact Test: $P = 0.6$).

Table 1. Number of pufflings that chose each of the two light options in each choice
combination. Individual binomial tests assessed evidence for a preference of one light type over
the other in each choice combination (excluding "No choice" individuals), using 0.5 as the
theoretical (null) proportion for each of the two options. We report the 95% confidence intervals
of the true proportion of "Light type 1" as a reference. We also present the number of individuals
that made "No choice" in each combination.

Light type 1	Quantity	Light type 2	Quantity	95% CI	*P*-value	No choice
Bright HPS	10	Dark	0	0.69-1.00	0.002	5
Bright HPS	14	Dim HPS	6	0.46-0.88	0.1	2
Bright HPS	10	Warm White	11	0.26-0.70	1.0	2
Warm White	10	Cool White	10	0.27-0.73	1.0	2
Dim Blue	13	Orange	9	0.36-0.79	0.5	2
Bright Blue	9	Orange	6	0.32-0.84	0.6	10

Assessment of Activity Levels in Different Light Types

In our open field test, there was a significant effect of light type on time spent mobile by

pufflings (Kruskal-Wallis test: $H_3 = 18.396$, $P = 0.0004$). Pufflings spent more time mobile in

Darkness (median time spent mobile = 120 seconds) and in HPS light (median 111 seconds) than

in Cool White LED light (median 32 seconds; Dunn test: $P = 0.003$ and $P = 0.004$, respectively;

Figure 4). Pufflings also spent more time mobile in Darkness than in Warm White LED light

(median 54 seconds; Dunn test: $P = 0.045$). Pufflings spent more time mobile, but only

marginally, in HPS light compared to Warm White LED light (Dunn test: $P = 0.050$; Figure 4). If

the statistical outlier is included (see Methods), the difference between Warm White LED and

Darkness is no longer significant, although the trend remains the same ($P = 0.09$; see Appendix).

Figure 4. Time spent mobile by fledgling puffins when exposed to each of four light treatment
groups. Figure combines a violin plot and boxplot with overlaid individual data points. Sample
sizes: Warm White LED light ($N = 24$), Cool White LED light ($N = 25$), Darkness ("converted"
data; $N = 23$, see text), and High Pressure Sodium (HPS) light ($N = 19$). The single data point
>300 seconds in the warm LED treatment (outlined in red) was considered a statistical outlier
and was not included in analysis. Individuals were significantly less mobile when exposed to
Warm White LED and Cool White LED than when exposed to Darkness, and significantly less
mobile under Cool White LED than under HPS.

DISCUSSION

Our results form the first experimental evidence that light attraction behaviour is a causal mechanism behind the stranding of Atlantic Puffin fledglings (pufflings) in coastal areas during their initial fledging flights. This evidence emerges at both coarse and fine spatial scales. Even after controlling for search effort and size of the search area, in our coarse-scale beach illumination experiment we found significantly more pufflings stranded during the Light treatment than during the Dark treatment. Further, consistent with a light attraction hypothesis of stranding, in our y-maze experiments pufflings significantly preferred light over darkness and somewhat preferred bright HPS light over dim HPS light (although the latter was not statistically significant). Contrary to our expectation that pufflings would prefer "cool" light spectra over "warm" light spectra (based on results of Salamolard et al., 2001 as cited in Minatchy, 2004; Rodríguez et al., 2017b), we found no preferences when pufflings were simultaneously presented with two different light spectra. Our open field test experiment, in which we exposed pufflings to either darkness or one of three light types, revealed that pufflings were more active in Darkness than in any LED light, and more active in HPS light than under Cool White LED light. Pufflings also tended to be more active in HPS light than in Warm White LED light, but this difference only bordered on statistical significance.

We are first to demonstrate a reduction in stranding numbers of an alcid through the experimental reduction of ALAN. Our finding that more pufflings became stranded at our two beach sites during the Light treatment compared to the Dark treatment is consistent with similar studies on procellariiforms (Reed et al., 1985; Miles et al., 2010; Rodríguez et al., 2014; Burt et al., 2024). Our results therefore provide further support for recommendations to reduce or eliminate sources of ALAN near breeding colonies of seabirds affected by stranding, especially

where populations are already in decline (e.g., Rodríguez et al., 2017a). However, it should be

noted that the number of pufflings stranded during our experiment, and indeed during the entire

Puffin Patrol rescue program, is small relative to the estimated total fledging cohort (e.g., in 2019

an estimated $246,069 \pm 25,988$ juveniles successfully fledged from the Witless Bay Ecological

Reserve and 326 were found stranded; Wilhelm et al., 2021). It is possible, then, that navigation

error during natural dispersal movements inevitably brings a small proportion of fledglings close

to land, where they become attracted to and stranded by nearby artificial lights (Brown et al.,

2023).

Previously, we expressed concern that human visual perception is more acute, and

therefore the detection of stranded birds is likely more efficient in light conditions compared to

darkness, potentially leading to a bias in detection probability (Brown et al., 2023). As such, in

our coarse-scale light attraction experiment we attempted to reduce the bias between our two

treatments by: 1) using identical night vision binoculars at each of our sites on each night of the

experiment; 2) clearly delineating a search area of the same size (area of approximately 50 m ×

15-20 m) on both beaches that took into account the maximum visual range of the binoculars;

and 3) controlling for the number of observers present at each site in our data analysis. Even if

our Light treatment was still subject to bias toward elevated detection probability, it is highly

unlikely that the large difference in number of pufflings found within the beach search areas ($N =$

54 in Light vs $N = 1$ in Dark) is attributable to missing a significant number of pufflings in the

Dark treatment. In fact, the single puffling we detected within the search area during the Dark

treatment was on a night with very low lunar illumination (13.3%), after the moon had already

set, and during overcast and rainy conditions. That pufflings were visible to observers even in the

worst of visibility conditions suggests that if more individuals were indeed present in the Dark

treatment, they would have been detected. Additionally, by alternating each night the site

location for Light treatment, we simultaneously controlled for the potential effects of date, moon

phase (see Wilhelm et al., 2021) and site on the numbers of pufflings we observed.

In addition to the stark difference in number of pufflings observed within the delineated

beach search areas in the Light versus Dark treatment, more individuals were also observed

swimming and flying in the Light treatment ($N = 77$ and $N = 78$, respectively) than in the Dark

treatment ($N = 1$ and $N = 1$, respectively).

Support for the light attraction hypothesis of stranding in pufflings is bolstered by the

positive phototactic behaviour pufflings displayed in our choice experiment. Pufflings chose

Bright HPS light significantly more often than darkness and chose Bright HPS light slightly (but

insignificantly) more often than Dim HPS. These results agree with those of a similar study in

which Ancient Murrelet chicks chose light significantly more often than darkness in a T-maze

just after leaving their nests (Gaston et al., 1988), but interestingly are contrary to results of a Y-

maze experiment in which Cory's shearwater fledglings chose darkness over light (Atchoi et al.,

2024). Ancient Murrelet chicks leave the nest at night under parental care on average only two

days after hatching (Gaston and Shoji, 2020), long before they are capable of flight. It is this lack

of flight ability, combined with breeding in mostly remote areas away from human settlement,

that presumably precludes them from stranding in areas with ALAN. That our Atlantic puffin Y-

maze results are consistent with those of Ancient murrelets but not Cory's shearwaters possibly

suggests a taxonomic difference in fine-scale behavioural responses to ALAN between the two

main seabird groups affected by light-induced stranding (i.e., alcids and procellariiforms);

however, more research is necessary to confirm this trend.

Contrary to our expectations, pufflings did not display any preferences for certain light

spectra over others when presented with a choice. We based our expectations on two premises:

first, previous studies have found that attraction behaviour and stranding numbers of fledgling

procellariiforms increase under exposure to short wavelength-dominant (i.e., high CCT) light

types like metal halide (~4500 K CCT) compared to long wavelength-dominant (low CCT) light

types like high pressure sodium (~2000 K CCT; Salamolard et al., 2001 as cited in Minatchy,

2004; Rodríguez et al., 2017b), although this is not always the case (e.g., shielding of long

wavelength light from radiating upward may negate the effect; Urmston et al., 2022; Atchoi et

al., 2024) and evidence is limited. Second, the procellariiform eye is quite sensitive to short

wavelengths (Reed, 1986; Bowmaker et al., 1997; Hart, 2004), implying a potential link between

the wavelengths of highest spectral sensitivity and those that induce the strongest phototactic

responses. Atlantic Puffins likely have similar spectral sensitivities given the similarities in

photoreceptor profiles between auks and procellariiforms (Ödeen and Håstad, 2003), which is

why we expected they would choose blue light over orange and cool white over warm white

LED light in our Y-maze. It is therefore unclear why these choice combinations did not elicit

differences in phototactic behaviour. Most of our light stimuli were calibrated such that the

amount of light irradiated at the peak wavelength of each spectrum was relatively equal; we did

this because the calibration of spectra based on perceived "brightness", although valuable, is

difficult without a complete understanding of spectral sensitivities in the species of interest. To a

puffling's eyes our provided light options were therefore not necessarily perceived as all being of

the same "brightness", and there may thus have been unquantified effects of perceived brightness

on the choice behaviour they exhibited in some of our choice combinations. Another possible

explanation is that the locomotory context of the birds (i.e., standing and walking) and/or the

spatial scale of the experimental apparatus are so contextually different from flying in a three-

dimensional airspace that they elicit a completely different mode of behaviour as it relates to

lighting conditions. A coarse-scale beach illumination experiment that compares phototactic

responses and stranding numbers across several different light spectra (ideally calibrated

according to perceived brightness) may provide insight in a more realistic context (as per

Rodríguez et al., 2017b).

When we measured the activity levels of pufflings exposed to darkness and various spectra

of artificial light in our open field test, they spent significantly more time active in Darkness than

in Cool or Warm White LED light (despite the effect of "converting" the Dark treatment scores

to reduce the significance of these differences), and significantly more time active in HPS light

than in Cool White LED light. It is unclear what factors drove our results, especially in Darkness.

It is possible that in the HPS light treatment the low-level noise of the associated ballast fan

(unlike the LED lights which have no ballast fan) elicited an attractive response, which has been

anecdotally observed in stranded pufflings (Harris, 1982; Harris et al., 1998; reviewed in

Wilhelm et al., 2013). Conversely, it is possible that the greater proportion of cool-hued light in

LED spectra compared to HPS has a calming effect on pufflings that results in reduced activity,

as blue light does in domestic ducks (Sultana et al., 2013) and in Black-headed Buntings

(*Emberiza melanocephala*) experiencing periods of migratory restlessness (but with this pattern

reversed outside these periods; Yadav et al., 2015). A fear-based freezing response toward some

aspect of LED light could also drive the reduced activity levels we observed in that light type,

similar to that observed in some poultry lines when placed in an open field arena (Forkman et al.,

2007; Campbell et al., 2019), although our anecdotal observations of pufflings preening

themselves in the arena during bouts of immobility would appear to contradict this hypothesis.

Ultimately, the true cause of the reduced activity levels that we observed in pufflings exposed to

LED light, especially in Cool White LED light, remains a matter of conjecture. Regardless of the

reason, these differences in activity patterns may hold implications for detection probability

following stranding; for example, pufflings may be easier to find in darkness and under HPS

light if they are more mobile there, whereas comparatively less active birds under LED light may

be less noticeable, all else being equal.

Conservation Implications

Globally, there is a concerted effort to reduce the impact of ALAN on wildlife, which

includes efforts to mitigate and prevent the stranding of various seabird species. To date, most

research has focused on procellariiforms, with comparatively little attention paid to the Atlantic

Puffin. We have demonstrated experimentally that the addition of sources of ALAN near Atlantic

Puffin breeding colonies has the potential to drastically increase the number of fledglings that

become stranded, which in turn would almost certainly contribute to decreased juvenile survival.

However, perhaps counter-intuitively, we suggest that light could also be used to lure wayward

puffins away from areas of high mortality risk (i.e., towns and roads) to those of low risk (e.g.,

supervised remote sites) as part of a dynamic rescue effort, and this potential application of

ALAN in a conservation context deserves further research. From fine-scale behavioural

experiments, we demonstrated that Atlantic Puffin fledglings display positive phototactic

behaviour in response to all tested bulb colours, without a preference for either blue- or red-

dominant spectra. Although there may also be compounding effects of perceived brightness, this

implies that changes to street lighting from historically prevalent HPS lamps to more energy-

efficient LED, or (if specifically attempting to reduce attraction) from cool-hued to warm-hued

lamps, may not be effective in reducing strandings in Atlantic Puffins and that the reduction and

shielding of ALAN sources remain the best recommendations. Our open field test results imply

that post-stranding LED light exposure may have either a calming or startling effect on stranded

pufflings, with potential implications for rescue, rehabilitation, and even husbandry of puffins

and other auks.

Data availability statement: Data are available as Supplementary Material.

Declaration of Generative AI and AI-assisted technologies in the writing process: During the

preparation of this work the authors used ChatGPT to formulate a first draft of the title. After

using this tool, the authors reviewed and edited the title as needed and take full responsibility for

its content.

REFERENCES

Atchoi E, M Mitkus, B Machado, V Medeiros, S Garcia, M Juliano, J Bried, and A Rodríguez (2024). Do seabirds dream of artificial lights? Understanding light preferences of Procellariiformes. bioRxiv. https://doi.org/10.1101/2024.03.01.582998

Bowmaker JK, LA Heath, SE Wilkie, and DM Hunt (1997). Visual pigments and oil droplets from six classes of photoreceptor in the retinas of birds. Vision Research 37:2183-2194. https://doi.org/10.1016/S0042-6989(97)00026-6

Brown TM, SI Wilhelm, GF Mastromonaco, and G Burness (2023). A path forward in the investigation of seabird strandings attributed to light attraction. Conservation Science and Practice 5: e12852. https://doi.org/10.1111/csp2.12852

Burt TV, SM Collins, S Green, PB Doiron, SI Wilhelm, and WA Montevecchi (2024). Reduction of coastal lighting decreases seabird strandings. PLoS ONE 19: e0295098. https://doi.org/10.1371/journal.pone.0295098

Campbell DLM, EJ Dickson, and C Lee (2019). Application of open field, tonic immobility, and attention bias tests to hens with different ranging patterns. PeerJ 7:e8122 http://doi.org/10.7717/peerj.8122

Collins SM, A Hedd, WA Montevecchi, TV Burt, DR Wilson, and DA Fifield (2023). Small tube-nosed seabirds fledge on the full moon and throughout the lunar cycle. Biology Letters 19:20230290. http://doi.org/10.1098/rsbl.2023.0290

Davies TW, and T Smyth (2018). Why artificial light at night should be a focus for global change research in the 21st century. Global Change Biology 24:872-882. https://doi.org/10.1111/gcb.13927

Durmus D (2022). Correlated color temperature: use and limitations. Lighting Research & Technology 54:363-375. https://doi.org/10.1177/14771535211034330

Fabian ST, Y Sondhi, PE Allen, JC Theobald, and H-T Lin (2024). Why flying insects gather at artificial light. Nature Communications 15: 689. https://doi.org/10.1038/s41467-024-44785-3

Fayet AL, GV Clucas, T Anker-Nillsen, M Syposz, and ES Hansen (2021). Local prey shortages drive foraging costs and breeding success in a declining seabird, the Atlantic puffin. Journal of Animal Ecology 90:1152-1164. http://doi.org/10.1111/1365-2656.13442

Forkman B, A Boissy, M-C Meunier-Salaün, E Canali, and RB Jones (2007). A critical review of fear tests used on cattle, pigs, sheep, poultry and horses. Physiology & Behavior 92:340-374. https://doi.org/10.1016/j.physbeh.2007.03.016

Gaston AJ, and A Shoji (2020). Ancient Murrelet (Synthliboramphus antiquus), version 1.0. In Birds of the World (A. F. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.

Gaston AJ, IL Jones, DG Noble, and SA Smith (1988). Orientation of ancient murrelet, Synthliboramphus antiquus, chicks during their passage from the burrow to the sea. Animal Behaviour 36:300-303. https://doi.org/10.1016/S0003-3472(88)80277-X

Harris MP (1982). Seasonal variation in fledging weight of the puffin *Fratercula arctica*. Ibis
124:100-103.

Harris MP, S Murray, and S Wanless (1998). Long-term changes in breeding performance of
puffins *Fratercula arctica* on St Kilda. Bird Study 45(3):371-374.
https://doi.org/10.1080/00063659809461109

Hart NS (2004). Microspectrophotometry of visual pigments and oil droplets in a marine bird,
the wedge-tailed shearwater *Puffinus pacificus*: topographic variations in photoreceptor
spectral characteristics. Journal of Experimental Biology 207:1229-1240.

Jägerbrand AK, and K Spoelstra (2023). Effects of anthropogenic light on species and
ecosystems. Science 380: 1125-1130. https://doi.org/10.1126/science.adg3173

Katz C (2023, March 31). *An Icelandic town goes all out to save baby puffins.* Smithsonian
Magazine online. https://www.smithsonianmag.com/science-nature/icelandic-town-goes-
all-out-save-baby-puffins-180981518/

Longcore T, A Rodríguez, B Witherington, JF Penniman, L Herf, and M Herf (2018). Rapid
assessment of lamp spectrum to quantify ecological effects of light at night. Journal of
Experimental Zoology Part A 329:511-521. https://doi.org/10.1002/jez.2184

Lowther PE, AW Diamond, SW Kress, GJ Robertson, K Russell, DN Nettleship, GM Kirwan,
DA Christie, CJ Sharpe, EFJ Garcia, and PFD Boesman (2020). Atlantic Puffin
(*Fratercula arctica*), version 1.0. In Birds of the World (SM Billerman, Editor). Cornell
Lab of Ornithology, Ithaca, NY, USA.

Mercer J (2018, August 26). *N.L. biologist worries offshore lights stranding seabirds.* Saltwire.
https://www.saltwire.com/newfoundland-labrador/news/nl-biologist-worries-offshore-
lights-stranding-seabirds-236694/

Miles W, S Money, R Luxmoore, and RW Furness (2010). Effects of artificial lights and
moonlight on petrels at St Kilda. Bird Study 57:244-251.
https://doi.org/10.1080/00063651003605064

Miles WTS, R Mavor, NJ Riddiford, PV Harvey, R Riddington, DN Shaw, D Parnaby, and JM
Reid (2015). Decline in an Atlantic Puffin population: evaluation of magnitude and
mechanisms. PLoS ONE 10(7): e0131527. http://doi.org/10.1371/journal.pone.0131527

Minatchy N (2004). Stratégie de réduction de la mortalité des petrels induite par les éclairages
publics. Specialized diploma dissertation, University of Réunion, Réunion.

Ödeen A, and Håstad O (2003). Complex distribution of avian color vision systems revealed by
sequencing the SWS1 opsin from total DNA. Molecular Biology and Evolution 20:855-
861. https://doi.org/10.1093/molbev/msg108

Ogle DH, JC Doll, AP Wheeler, and A Dinno (2023). _FSA: Simple Fisheries Stock Assessment
Methods_. R package version 0.9.4. https://CRAN.R-project.org/package=FSA

Pawson SM, and MK-F Bader (2014). LED lighting increases the ecological impact of light
pollution irrespective of color temperature. Ecological Applications 24:1561-1568.
https://doi.org/10.1890/14-0468.1

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reed JR (1986). Seabird vision: spectral sensitivity and light-attraction behavior. PhD thesis,
University of Wisconsin – Madison, Wisconsin, USA.

Reed JR, JL Sincock, and JP Hailman (1985). Light attraction in endangered procellariiform
birds: reduction by shielding upward radiation. Auk 102(2):377-383.

Rich C, and T Longcore (Eds.). (2006). *Ecological consequences of artificial night lighting.*
Island Press.

Rodríguez A, G Burgan, P Dann, R Jessop, JJ Negro, and A Chiaradia (2014). Fatal attraction of
short-tailed shearwaters to artificial lights. PLoS ONE 9(10): e110114.
https://doi.org/10.1371/journal.pone.0110114

Rodríguez A, ND Holmes, PG Ryan, K-J Wilson, L Faulquier, Y Murillo, AF Raine, JF
Penniman, V Neves, B Rodríguez, JJ Negro, A Chiaradia, P Dann, T Anderson, B
Metzger, M Shirai, L Deppe, J Wheeler, P Hodum, C Gouveia, V Carmo, GP Carreira, L
Delgado-Alburqueque, C Guerra-Correa, F-X Couzi, M Travers, and M Le Corre
(2017a). Seabird mortality induced by land-based artificial lights. Conservation Biology
31(5): 986-1001.

Rodríguez A, P Dann, and A Chiaradia (2017b). Reducing light-induced mortality of seabirds:
High pressure sodium lights decrease the fatal attraction of shearwaters. Journal for
Nature Conservation 39:68-72.

Salamolard M, F-X Couzi, M Pellerin, and T Ghestemme (2001). Etude complémentaire
concernant la protection du Puffin de Baillon dans le cadre de la Route des Tamarins.
Rapp. SEOR/DDE-Réunion. 30 pp. + Annexes 4 pp.

Sanders D, E Frago, R Kehoe, C Patterson, and KJ Gaston (2021). A meta-analysis of biological
impacts of artificial light at night. Nature Ecology & Evolution 5: 74-81.
https://doi.org/10.1038/s41559-020-01322-x

Sultana S, MR Hassan, HS Choe, and KS Ryu (2013). Impact of different monochromatic LED
light colours and bird age on the behavioural output and fear response in ducks. Italian
Journal of Animal Science 12(4):e94. https://doi.org/10.4081/ijas.2013.e94

Urmston J, KD Hyrenbach, and K Swindle (2022). Quantifying Wedge-tailed shearwater
(*Ardenna pacifica*) fallout after changes in highway lighting on southeast O'ahu, Hawai'i.
PLoS ONE 17(3): e0265832. https://doi.org/10.1371/journal.pone.0265832

Van Doren BM, DE Willard, M Hennen, KG Horton, EF Stuber, D Sheldon, AH Sivakumar, J
Wang, A Farnsworth, and BM Winger (2021). Drivers of fatal bird collisions in an urban
center. PNAS 118: e2101666118. https://doi.org/10.1073/pnas.2101666118

Wilhelm SI, JJ Schau, E Schau, SM Dooley, DL Wiseman, and HA Hogan (2013). Atlantic
Puffins are attracted to coastal communities in eastern Newfoundland. Northeastern
Naturalist 20:624-630.

Wilhelm SI, J Mailhiot, J Arany, JW Chardine, GJ Robertson, and PC Ryan (2015). Update and
trends of three important seabird populations in the western North Atlantic using a
geographic information system approach. Marine Ornithology 43:211-222.

Wilhelm SI, SM Dooley, EP Corbett, MG Fitzsimmons, PC Ryan, and GJ Robertson (2021).
Effects of land-based light pollution on two species of burrow-nesting seabirds in
Newfoundland and Labrador, Canada. Avian Conservation & Ecology 16:12.
https://doi.org/10.5751/ACE-01809-160112

Witherington BE (1997). The problem of photopollution for sea turtles and other nocturnal
animals. In JR Clemmons & R Buchholz (Eds.), *Behavioral approaches to conservation*
in the wild (pp. 303-328). Cambridge University Press.

Yadav G, S Malik, S Rani, and V Kumar (2015). Role of light wavelengths in synchronization of
circadian physiology in songbirds. Physiology & Behavior 140:164-171.
http://dx.doi.org/10.1016/j.physbeh.2014.12.032

Yang Y, Q Liu, C Pan, J Chen, B Xu, K Liu, J Pan, M Lagisz, and S Nakagawa (2024). Species
sensitivities to artificial light at night: a phylogenetically controlled multilevel meta-
analysis on melatonin suppression. Ecology Letters 27: e14387.
https://doi.org/10.1111/ele.14387

APPENDIX

Methods

Conversion of "Dark" treatment activity levels

To determine if we should "convert" the audial scores of time spent mobile by pufflings in

the "Dark" treatment (x) to estimated scores of time spent mobile had been scored visually (y),

we calculated 95% confidence intervals of the slope and intercept estimates (that is, 0.92 and

2.30, respectively). We decided *a priori* that if either the slope estimate's 95% confidence

intervals did *not* contain 1.00 or if the intercept estimate's 95% confidence intervals did *not*

contain 0.00, we would convert the "Dark" data using *y = 2.3 + 0.92x*; otherwise, we would use

the raw "Dark" data for analysis. The 95% confidence interval of the slope was [0.87, 0.97] and

that of the intercept was [-5.29, 9.95]. The slope estimate's 95% confidence intervals therefore

did not contain 1.00, so we converted the "Dark" data for analysis.

Results

Assessment of activity levels in different light types when outlier is included

The following results were obtained when the individual identified as an outlier (in the

Warm White LED light treatment of the open field test; spent 353 seconds mobile) was retained

in our analysis. There was a significant effect of light type on time spent mobile by pufflings

(Kruskal-Wallis test: $H_3 = 17.184$, $P < 0.001$); post-hoc testing revealed that pufflings spent more

time mobile in Darkness (median time spent mobile = 120 seconds) and in HPS light (median

111 seconds) than in Cool White LED light (median 32 seconds; $P = 0.004$ for both comparisons;

Figure 4). Pufflings also tended to spend more time mobile in Darkness and in HPS light than in

Warm White LED light (median 56 seconds; $P = 0.09$ and $P = 0.10$, respectively), but these

differences failed to reach significance (Figure 4).

Table A1. Minimum ("Min"), best-estimate ("Est"), and maximum ("Max") counts of pufflings observed at the two beaches during Dark and Light treatments over the course of 11 experimental nights, accompanied by percent moon illumination on each night ("Moon %") and qualitative observations of weather conditions at the sites. All counts for both treatments were conducted over the same experimental duration each night (two hours). Percent moon illumination data were collected from moongiant.com. Weather conditions were not recorded at the sites on night 4; conditions on this night were obtained from Environment and Climate Change Canada (climate.weather.gc.ca).

Night	Moon %	Weather	Dark			Light		
			Min	Est	Max	Min	Est	Max
1	85	Clear	0	0	0	10	14	23
2	76	Cloud & fog	0	0	0	34	44	52
3	47	Cloud & rain	1	1	1	11	12	17
4	38	Drizzle[*]	0	0	0	27	28	30
5	29	Clear	0	0	0	14	15	19
6	20	Clear	0	0	0	6	6	6
7	13	Cloud & rain	1	1	2	5	5	5
8	8	Cloud & wind	0	0	0	4	4	4
9	3	Cloud	0	0	0	6	7	8
10	1	Cloud & wind	0	0	0	0	0	0
11	0	Clear	0	0	0	1	1	1

[*]Weather data obtained from Environment and Climate Change Canada for St. John's International Airport (47.618611°N, 52.752500°W), located 38-40 km from the two beach sites used in this experiment.

Table A2. Minimum ("Min"), best-estimate ("Est"), and maximum ("Max") "observation rates"
(pufflings seen, divided by number of observers) at the beaches during Dark and Light
treatments over the course of 11 experimental nights. All observation rates for both treatments
apply to a two-hour experimental duration.

Night	Dark			Light		
	Min	Est	Max	Min	Est	Max
1	0	0	0	5	7	11.5
2	0	0	0	11.3	14.7	17.3
3	0.5	0.5	0.5	3.7	4	5.7
4	0	0	0	9	9.3	10
5	0	0	0	4.7	5	6.3
6	0	0	0	2	2	2
7	0.5	0.5	1	1.7	1.7	1.7
8	0	0	0	0.8	0.8	0.8
9	0	0	0	1.5	1.75	2
10	0	0	0	0	0	0
11	0	0	0	0.5	0.5	0.5

908 **Table A3.** Materials used to create each of the eight light options used in the choice experiment,
909 including the specific lightbulbs used, whether a neutral-density filter or diffuser were used, and
910 any other materials used.

Light option	Lightbulb	NDF[a]	Diffuser[b]	Other
Dim HPS	2100 K HPS[c,d]	none	yes	none
Bright HPS	2100 K HPS[d]	yes	yes	none
Cool white LED	5000 K LED[e]	yes	yes	none
Warm white LED	2700 K LED[f]	yes	yes	none
Bright blue LED	5000 K LED[e]	none	none	ePlastics #2424 dark blue acrylic; cobalt glass
Dim blue LED	5000 K LED[e]	yes	yes	ePlastics #2424 dark blue acrylic; cobalt glass
Orange LED	2700 K LED[f]	yes	yes	ePlastics #2422 amber orange acrylic
Darkness	none	none	none	light hole covered

[a]A variable neutral density filter (NDF) consisted of two pieces of polyvinyl alcohol-iodine polarizing film,
overlapping at varying degrees of polarization depending on the light option.
[b]Diffuser consisted of one piece of parchment paper.
[c]Incident light only (i.e., crossover light from Bright HPS on other choice box).
[d]High pressure sodium; Light EnerG, 400 Watt
[e]Light-emitting diode; Philips, 5W / 40W-equivalent Daylight LED
[f]Light-emitting diode; Philips, 5W / 40W-equivalent Soft White LED

Figures

Figure A1. Spectrum of the 6390 K, 30 000 lumen LED light apparatus used on two beach sites
to test the responses of pufflings to light at a coarse spatial scale. Note that the y-axis has been
normalized to arbitrary units with a maximum of 1.0.

Figure A2. Spectra of all light options used in the choice experiment: A) Dim HPS; B) Bright
HPS; C) Cool White (5000 K) LED light; D) Warm White (2700 K) LED light; E) Bright Blue
LED light; F) Orange LED light; G) Dim Blue LED light; and H) Darkness. Irradiance is as
measured by a spectrometer with its sensor placed at the entrance of the "choice arm", inside the
"main box" of the Y-maze. * = note the change in y-axis scale.

930 **Figure A3.** Comparison of time spent mobile by 25 pufflings in a 600 second trial in 25 lighted
931 videos (including HPS, Warm White LED, and Cool White LED), as scored by watching the
932 video (y-axis) versus listening to the video (x-axis). Note that there are two overlapping points
933 around (0, 0).