Oncogenic MAPK pathway activation induces a congenital and progressively lethal rare neuropathy in mice

Elise Marechal[1], Daniel Aldea[1,%], Patrice Quintana[1,%,#], Grégoire Mondielli[1], Nathalie Bernard-Marissal[1], Mathias Moreno[1], Claire El Yazidi[1], Natacha Broucqsault[1], Valérie Delague[1], Lauren A. Weiss[2,3], Anne Barlier[1,4], Heather C. Etchevers[1,*]

[1] INSERM, Marseille Medical Genetics, U1251, MarMaRa Institute, Aix Marseille University, Marseille, France
[2] Department of Psychiatry and Behavioral Sciences, University of California, San Francisco, San Francisco, CA 94143, USA
[3] Institute for Human Genetics, University of California, San Francisco, San Francisco, CA 94143, USA
[4] AP-HM, MMG, MarMaRa Institute, La Conception Hospital Laboratory of Molecular Biology, Marseille, France.

% These authors have contributed equally to this work.

Current address: University of Clermont Auvergne, INSERM, U1107, NEURO-DOL, Clermont-Ferrand, France

* Corresponding author

E-mail : heather.etchevers@inserm.fr

Short title: Sustained Braf promotes cellular immaturity

Keywords: rare disease, mosaicism, signaling, neural crest, glia, cancer, development

Abstract

RASopathies, rare congenital syndromes affecting skin, heart, brain and skeletal systems, often include peripheral neuropathy of unknown origin. Although RASopathy gene variants are familiar proto-oncogenes after birth, the causal relationships between timing and specific pathological outcomes remain unclear. This study examined how a mitogen-activated protein kinase (MAPK) effector, Braf, causes peripheral neuropathy. Targeting the oncogenic Braf(V600E) variant, found in mosaic RASopathies, to cells having expressed myelin protein zero (Mpz) in mice triggered a degenerative neuropathy evoking type 1 Charcot-Marie-Tooth syndrome, with pre-weaning onset of hyperplastic nerves, hindlimb weakness, and unexpectedly reduced body size. Constitutively active Braf expanded a Jun+ Schwann cell repair state, impairing myelination by mature Schwann cells. Patient-derived iPSCs bearing the *BRAF* Q257R variant, associated with cardio-facio-cutaneous syndrome, failed to differentiate into mature Schwann cells, instead exhibiting transcriptional profiles associated with progenitor or repair states. These findings suggest that somatic mosaicism may contribute to unresolved genetic heterogeneity in human neuropathies and enlarge the candidate gene list. They also reveal a MAPK-dependent mechanism by which neural crest-derived Schwann cells regulate both body growth and peripheral nerve homeostasis.

Introduction

Many cancers and congenital diseases co-opt and deploy the molecular mechanisms of normal organogenesis in inappropriate contexts. Activation of the mitogen-activated protein kinase (MAPK) intracellular signaling pathway drives dozens of solid and liquid cancers in somatic tissues of humans and animal models [1]. In contrast, RASopathies are heterogeneous developmental syndromes that preferentially affect epidermis and pigmentation; cardiovascular, craniofacial and skeletal systems; and neurological or endocrine functions impacting body size and fertility [2–7]. Variants of at least 24 genes that boost and sustain MAPK signaling to, and nuclear translocation of, the extracellular signal-regulated kinases (ERK1/2) are responsible for pathogenesis [8]. Activating variants may be expressed constitutionally or prenatally restricted to yield lineage-specific mosaic or somatic RASopathies [9]. Most RASopathies are associated with specific cancer predispositions in constitutional and somatic presentations, particularly when syndromic [10,11].

A limited number of specific MAPK pathway variants engender this wide spectrum of overlapping clinical outcomes. Oncogenic variants in *BRAF*, particularly encoding changes to valine (p.V)600, drive aggressive forms of melanoma as well as thyroid, colorectal, lung, ovarian and breast cancers [1,12–14]. BRAF p.V600E has never been observed in germline disease because of early embryonic lethality, observed in mouse models [15], and cellular cytotoxicity through its strong constitutive ERK1/2 activation. BRAF p.V600E yields an active, monomeric kinase, independent of RAS-mediated transduction [16,17]. In contrast, mild pathway-activating *BRAF* variants such as p.Q257R underlie a growing list of congenital RASopathies [10,18–20]. These include the germline RASopathies, cardio-facio-cutaneous syndrome (CFC; ORPHA:1340) and Noonan syndrome with (ORPHA:500) or without (ORPHA:648) multiple lentigines. Their common clinical features include reduced postnatal growth, dysmorphic facial features, melanocytic nevi or other pigmented lesions, congenital heart defects, and tumor predispositions [21–24].

The mechanisms and triggers of neuropathies in the context of RASopathies have not been explored to date. Only recently have multiple CFC and Noonan syndrome patients been reported with neuropathies causing pain, sensory loss and/or muscle wasting [25–27], evoking similar symptoms in Charcot-Marie-Tooth syndrome, type 1 (CMT1; Callegari et al., 2019). *Mek1*, the major MAP kinase kinase (MAP2K) phosphorylating Erk1/2, is known to be necessary for nerve myelination during maturation, while its conditional over-expression or sustained activity in mature Schwann cells (SC; peripheral glia) induces murine neuropathy [29–32]. Although CMT is genetically heterogeneous, with many dozens of causal genes identified,

none are known MAPK effector genes; that said, one in six clinical CMT1 patients still do not

have a molecular diagnosis [33].

Human peripheral nervous system (PNS) architecture and conduction depend fully on SC,

which provide physical and trophic nerve support while myelinating large, fast-conduction

axons [34–36]. The entire PNS, including SC and perineurial stroma of its sensory and autonomic

components, derives from the multipotent embryonic neural crest (NC) population, which also

supplies critical craniofacial and cardiovascular connective tissues and all melanocytes [36,37].

Given the shared origin of numerous cell types affected in germline RASopathies, we

hypothesized that multipotent NC-derived lineages may be particularly vulnerable to increased

MAPK signaling before birth. The ability of NC-derived cells to instruct fate or regulate

homeostasis in neighboring tissues over time may also be compromised.

The goal of this study was to assess systemic and cell-autonomous effects of constitutively

active BRAF signaling on multipotent, NC-derived progenitor differentiation. We found that

conditional expression of *Braf* p.V600E in mouse cells having expressed the SC-specific gene

Mpz (encoding myelin protein zero, or P0) led to a peripheral neuropathy in association with

induction and maintenance of an ineffective repair SC phenotype. Although nerve myelination

initiated after birth as usual, mutant peripheral glia rapidly misregulated hundreds of genes

associated with mature SC function, including myelination, and nerves underwent active

myelin destruction. Unexpectedly, although endocrine tissues had not been intentionally

targeted in this model, mice also underwent developmental delay and remained sexually

ambiguous or immature. Furthermore, human pluripotent cells derived from two teenaged

donors with CFC syndrome, carrying the most frequent postnatal activating *BRAF* missense

variant, p.Q257R, were also incapable of terminal SC differentiation *in vitro*, in contrast to cells

from wildtype donors, after upregulating repair SC transcripts. Our results suggest that

molecularly undiagnosed cases of RASopathies, or endophenotypes such as CMT type 1 or

idiopathic cryptorchidism [38], may be imputable to somatic mosaicism for oncogenic MAPK

pathway gene variants in the NC lineage.

Results

Sustained Schwann cell MAPK signaling in causes juvenile locomotor and growth deficits

In order to model constitutive Braf-mediated signaling in progenitors of Schwann cells *in vivo*,

we used *Mpz-Cre*[+/°] mice, which transcribe *Mpz* (encoding myelin protein zero, or P0) as early

as embryonic day 12 in neural crest cells [39,40]. Crossed to a knock-in floxed *Braf* allele bearing

the hotspot V600E variant (*Braf*[V600E]) (**Fig S1a**), the *Mpz-Cre*[+/°];*Braf*[V600E/+] mice, abbreviated

hereafter as Mpz-*Braf*, were born at expected Mendelian ratios and remained asymptomatic

for the first week. However, from postnatal day (P)7, all Mpz-*Braf* mutants developed signs of peripheral neuropathy including tremors, abnormal hindlimb postural reflexes and trailing legs (**Fig 1a**). They nonetheless showed age-appropriate activity in exploration, feeding and grooming. Correlating with phenotypic onset, mutant sciatic nerves exhibited significantly more Erk1/2 phosphorylation than controls, increasing from early (P5; **Fig 1c, e**) to later (P21; **Fig 1d, e**) stages of SC-mediated myelination, demonstrating congenital and persistent MAPK activation in peripheral glia.

After P10, mutant mice were also visibly smaller than control littermates, with significantly reduced but proportional nose-to-tail length and weight (**Fig 1f**). Proportionately small body size could result from a deficiency in growth hormone (Gh), particularly as activating *BRAF* mutations have been described in the context of human hypopituitarism [2]. To see whether there were any changes in hormonal transcription, pituitaries at P5 were compared between control and Mpz-*Braf* mice by bulk RNA-sequencing (RNAseq). As for all other genes, Gh releasing hormone receptor *Ghrhr* or *Gh* transcripts, were not significantly altered in the two groups at this stage (**Fig S1b**; **Table S1**). Gh binds to peripheral Ghr, a type I cytokine receptor, to regulate the production of its effector insulin-like growth factor 1 (Igf1) in the liver and other sites, exerting thereby anabolic effects on bone and muscle [41]. We therefore compared the transcription of *Ghr* and *Igf1* by RT-qPCR in the livers of P21 Mpz-*Braf* and control littermate mice. While *Ghr* was upregulated on average by 3.3-fold in mutant livers, Igf1 was far more significantly diminished, by 211-fold (**Fig 1g**). The juvenile Mpz-*Braf* growth anomaly was the likely consequence of a near-complete loss of peripheral Igf1 production.

Braf$^{V600E/+}$ in SCs induces widespread peripheral nerve demyelination and axonal loss

Strikingly broad nerves were visible in all Mpz-*Braf* mice upon autopsy at P5 or P21; mutant nerves also lacked structural bands of Fontana [42] (**Fig 2a**; **Fig S1**). Despite their smaller bodies, Mpz-*Braf* sciatic nerves were significantly wider at P10 (0.48 ± 0.07 *vs* 0.31 ± 0.05 mm diameter in controls; **Fig 2b**). This discrepancy stabilized after weaning (0.74 ± 0.09 *vs* 0.39 ± 0.05 mm at P21, and 0.78 ± 0.05 *vs* 0.50 ± 0.04 mm at P35, respectively). Similar enlargement was also evident in all other peripheral nerves of Mpz-*Braf* mice examined at P21, including the sensorimotor peroneal, tibial, sural, trigeminal and ophthalmic nerves and the sensory saphenous and maxillary nerves (**Fig S1c-e**).

To further evaluate myelination dynamics and study the ultrastructural composition of mutant sciatic nerves, we used transmission electron microscopy (EM) to study cross-sections at P5, P21 and P28. Over time, degenerating axons, extracellular collagen, fibroblasts, macrophages and numerous SC nuclei could be identified. Myelin aberrations were frequent in Mpz-*Braf* nerves from P21 on, including examples of tomaculae, infoldings, onion bulbs and

degenerated axons, and focal hypermyelination of small-caliber axons, among other
anomalies (**Fig 2c**; **Fig S2a-d**). These structures were interspersed with abundant extracellular
collagen fibrils and tentacular fibroblasts (**Fig 2c**) and, by P42, axon-associated macrophages
containing myelin debris (**Fig S2e-f**). In contrast, at P5, no significant differences were
observed in average axonal density (52 ± 2 axons/mm^2 in Mpz-*Braf* vs 48 ± 3 axons/mm^2 in
controls) or diameter (1.447 ± 0.026 µm in Mpz-*Braf* vs 1.476 ± 0.072 µm in controls; **Fig 3a**;
Fig S3a-b). Likewise, no differences in immune infiltration were observed, while myelin
thickness as quantified by the *g*-ratio, the ratio of inner to outer diameters of axonal myelin
sheaths [43], remained constant (**Fig 3b** and **Fig S3c-d**) at an average of 0.732 ± 0.019 in Mpz-
Braf vs 0.708 ± 0.028 in controls. Only occasional myelin aberrations were observed in mutant
nerve cross-sections at this early stage (**Fig 2c**). Radial sorting and myelination therefore
began similarly in mutant and control peripheral nerves.

In contrast, axon density by P21 had declined significantly more in Mpz-*Braf* mutants than age-
matched controls (respectively, 4,060 ± 1,378 axons/mm^2 *vs* 36,172 ± 2,726 axons/mm^2; **Fig**
3c). By P28, a mean 7.2% of myelinated axons persisted in mutant sciatic nerves (1,841 ± 465
axons/mm^2 *vs* 25,703 ± 2,812 axons/mm^2 in controls; **Fig 3c**). Residual mutant axons were
significantly smaller on average (**Fig 3d-e**; 1.89 ± 0.23 µm in Mpz-*Braf* vs 2.64 ± 0.07 µm in
controls). To see which calibers of axons were affected, we examined width distributions of
792 and 746 myelinated axons at P28 from mutant and control mice, respectively (**Fig 3f**).
After weaning, there was relative loss of large axons of over 3 µm in diameter (**Fig 3e-f; Fig**
S3e; on average: 8.7 ± 6.7% in Mpz-*Braf* vs 33.3 ± 5.6% in controls).

The residual axons displayed 12% higher *g*-ratios, indicating thinner average myelin sheaths
(**Fig 3f; Fig S3f**; *g*-ratios: 0.66 ± 0.02 in Mpz-*Braf* vs 0.59 ± 0.00 in controls). To see if all axons
were affected equally, results were binned into small (<1 µm), medium (1-3 µm) and large (3
µm) diameter axons. Medium- and large-caliber axons had significantly thinner myelin in
mutants compared to controls (**Fig 3g**; average *g*-ratios: 0.67 ± 0.03 *vs* 0.59 ± 0.01 for medium-
caliber axons; 0.75 ± 0.00 *vs* 0.60 ± 0.01 for large-caliber axons). In contrast, small-caliber
axons exhibited a modest (10.6%) but significant increase in myelination (*g*-ratios of 0.47 ±
0.01 in mutants *vs* 0.52 ± 0.02 in controls; *cf*. **Fig 2c**, **Fig S2d**). In control mice, myelin thickness
increases with axon caliber to plateau at 4 µm and greater [30]. Fast-conduction axons of
motoneurons [44] were thus most severely affected in terms of myelination and persistence,
while some smaller axons were subject to hypermyelination.

Myelin component mislocalization and SC progenitor expansion
Extending these structural findings, we examined distribution of components of the myelinated
axon by teasing apart fixed fibers from P5 and P21 sciatic nerves. Immunofluorescence against

contact-associated protein 1 (Caspr, also called paranodin) demonstrated that in contrast to

controls (**Fig 4a**), residual myelin internodes were irregularly spaced, diffuse or absent along

mutant axons (**Fig 4b, d-g**). Early breakdown of myelin during its establishment in neighboring

axons was already visible in longitudinal view at P5, when F-actin labeling highlighted

successive beads of myelin ovoids along some mutant sciatic fibers (**Fig 4b-d**). By P21 (**Fig**

4e), only Mpz-*Braf* mice displayed some demyelinated segments entirely devoid of myelin

basic protein (Mbp) along individual fibers (**Fig 4f**). Multiple adherent nuclei between residual

Caspr+ nodes (**Fig 4f**), instead of a single SC following normal radial sorting [45], are consistent

with persistence of immature Mpz-*Braf* SCs.

To visualize SCs capable of proliferating, we examined the expression *in situ* of Sox10 (SRY-

box transcription factor 10), present throughout the Schwann cell lifespan [46], and Ki67, a

common indicator of cell cycle progression beyond quiescence [47,48], using co-immunostaining

of whole sciatic nerves at P42 in sagittal section. While Sox10+ nuclei were visible throughout

control nerves, with concomitant Ki67 immunofluoresence (**Fig S4a**), double-positive as well

as unlabeled nuclei were more densely and haphazardly arranged in the hyperplastic Mpz-

Braf nerves (**Fig. S4b**). All Sox10+ nuclei observed were also strongly Ki67+ in mutant nerves,

interspersed with additional Ki67+ Sox10- cells. Mutant sciatic nerves thus show prolonged

hyperplasia rather than hypertrophy, imputable both to probable inflammation and fibrosis but

also to sustained SC proliferation.

Braf$^{V600E/+}$ glia *in vivo* downregulate myelin sheath gene transcription

Bulk RNAseq was undertaken to identify global transcriptional changes associated with loss

of myelin after MAPK pathway activation, comparing Mpz-*Braf* and control sciatic nerves

dissected from four mice at ages P5 and P21 (**Fig 5** and **Table S2**). In contrast to P5 pituitaries,

we identified 2,540 differentially expressed genes (DEGs) between P5 mutant and control

nerves (*p*adj < 0.05 for significance). By P21, 10,409 DEGs were identified (**Fig 5a**).

Constitutive MAPK signaling therefore leads to early, large-scale transcriptional changes in

cells composing the sciatic nerves.

The myelin sheath consists of 70% to 85% lipids and 15 to 30% proteins [49]. By P21, there was

significant downregulation of terms related to cholesterol biosynthesis and metabolism in *Braf*-

mutant sciatic nerves (**Fig 5b, c**). However, the annotation "spinal cord injury" was significantly

enriched at both P5 and P21 (**Fig 5b**). Among strongly downregulated genes at P21 were

Hmgcr (11.5-fold), encoding a rate-limiting enzyme in cholesterol biosynthesis [50], and *Fa2h*

(7.9-fold), encoding fatty acid 2-hydroxylase, a rate-limiting myelin glycosphingolipid synthesis

enzyme [51] (**Table S2**). While *Pmp2* (encoding a fatty acid-binding protein, also called *Fabp8*)

and *Omg* (oligodendrocyte myelin glycoprotein) mRNA levels were significantly upregulated in P5 Mpz-*Braf* sciatic nerves, by P21 they were not significantly different from controls (**Fig 5d**).

Transcription of other abundant myelin proteins encoded by *Mpz*, *Pmp22* (peripheral myelin protein 22*)* or *Mbp* (myelin basic protein) were not significantly altered at P5, but other myelin components were already significantly if modestly reduced, including transcripts for myelin and lymphocyte protein (*Mal*), periaxin (*Prx*), the myelin-associated 2',3'-cyclic nucleotide 3' phosphodiesterase (*Cnp*), myelin-associated glycoprotein (*Mag*), plasmolipin (*Pllp*) and others (**Fig 5d**; **Table S1**). By P21, when sciatic myelination is nearly complete in controls (**Fig 2**), *Mpz*, *Mbp* and *Pmp22* transcripts among others were significantly less expressed than in controls (**Fig 5d**). The observed demyelination and axonal degeneration phenotype in response to sustained MAPK signaling was a thus a result of transcriptional downregulation of myelin components preceding axonal loss.

Transcriptional control of myelination is disrupted by constitutive MAPK signaling

We next examined the modulation of transcription factors (TF) that covaried with those known to impact the process of myelination. At P5, the major transcriptional activators *Egr2* (early growth factor response 2, also called Krox20 [52]), and *Sox10* [53], were unaffected. However, by P21, mutant SCs showed a significant decrease in *Egr2* expression with concomitant ectopic increases in neural crest or multipotency maintenance factors *Sox2, Foxd3, Tfap2a* and *Tfap2b,* and a significant decrease in the melanocyte determinant *Mitf*, with no effect on *Sox10* transcription (**Fig 6a**; **Table S2**). Simultaneously, SC-associated transcriptional repressors of myelination [54] were significantly upregulated in mutant sciatic nerves at P21, including *Jun*, *Egr1*, *Egr3,* and *Zeb2* (**Fig 6a**). The increase in *Jun* translated into significantly more Jun protein in mutant than control nerves after P21, while the reduction in Egr2 protein was only measurable at P42 (**Fig 6b**).

To corroborate the effects of decreased *Egr2* expression in mutant SCs, we reasoned that a distinct model of hypomyelinating neuropathy triggered by germline expression of a homozygous, hypomorphic *Egr2* allele (Le et al., 2005) should show similar transcriptional deregulation. Indeed, GSEA analysis found that the co-variance of 99/110 (90%) significantly increased or 98/107 (91.5%) decreased DEGs between the two animal models itself showed significant correlation (**Fig 6c**; *p* < 0.001). Sciatic nerve hypomyelination is therefore associated with overexpression of Jun and effected by less *Egr2*-driven target transcription.

During remyelination after mouse sciatic nerve injury, axonal Nrg1 type III stimulates SC *Pmp2* expression facilitate fatty acid uptake and increase mitochondrial ATP production [55]. We had observed significant *Pmp2* and massive *Nrg1* overexpression in the Mpz-*Braf* RNAseq results (**Table S2**); the major source of Nrg1 after P5 in this model is the SC rather than the axon itself

[56]. This reactive overexpression to injury has been shown to be responsible for "onion bulb" formation around neuropathic axons and an ultimately detrimental response to chronic repair SC stimulation [57].

Normally, Nrg1-III binds SC-specific ErbB3 to activate MAPK signaling, among other pathways. We hypothesized that the Mpz-*Braf* sciatic nerves at P5 and P21 should show a similar transcriptional response to an *Egr2*-driven gain-of-function *Mek1*DD allele at P8 and P30, which leads to hypermyelination in bypassing the natural decrease of Erbb3 in maturing SC [30]. As *Mek1* functions downstream of *Braf*, there was indeed robust positive correlation ($p < 0.0001$) at both stages of myelination and significant upregulation of *Erbb3* between P5 and P21 (**Fig 6d**; **Table S2**). *Egr2*-Cre-driven depletion of the upstream MAPK pathway phosphatase Shp2 entails, instead, a hypomyelinating neuropathy [30]. Shp2 is encoded by *Ptpn11*, a RASopathy gene responsible for Noonan syndrome [58]. We observed significant anti-correlation of DEG between Mpz-*Braf* and *Egr2^{Cre};Shp2^{f/f}* sciatic transcriptomes at initial myelinization (P5 vs P8), which changes to positive correlation after weaning (P21 vs P30; **Fig 6e**).

Finally, to see if the PNS effects of oncogenic Braf could be recapitulated by *Egr2*-expressing, rather than *Mpz*-expressing SC *in vivo*, we crossed heterozygous LSL-*Braf^{V600E/+}* mice to the same *Egr2*-Cre driver mice. The *Egr2*-Cre allele is expressed by SCs from mid-gestation on (Topilko et al., 1994; **Fig S5A**). However, mutant neonates were never obtained from this breeding, a significant ($p < 10^{-6}$) departure from the expected yield. Broader expression of *Egr2* in other cell types as compared to *Mpz* prevented further comparisons of postnatal myelination processes, due to fully penetrant embryonic lethality (**Fig S5B**).

Uninjured sciatic nerves initiate and sustain an early repair program

Transcripts significantly overexpressed at P5 and sustained through P21 in mutant sciatic nerves may reflect direct rather than indirect effects of MAPK signaling. We therefore examined the 175 genes that were significantly upregulated by two-fold or more at both stages with annotations from Wikipathways, KEGG and REACTOME (**Fig S6a**) and GSEA (**Fig S6b**; **Fig. 7a**). The term "spinal cord injury" (WP2431) remained one of the most significantly enriched even in this subset (cf. **Fig 5b**). In particular, cell adhesion molecule L1 (*Chl1*), one of the most specific transcripts for repair-type SC identified in single-cell RNAseq [34], was highly enriched over time. Comparing the overall transcriptome of Mpz-*Braf* sciatic nerves with that of a 6- to 8-week mouse model of axotomy at seven days post-injury [59] showed a significant positive correlation at P5 (**Fig S6c**) that increased with time (**Fig 7b**). The general molecular response to nerve injury is thus active during physiological myelinogenesis in juvenile Mpz-*Braf* mice.

Following physical injury such as crush or axotomy, SCs undergo extensive reprogramming, re-expressing genes from less differentiated stages and transcribing novel repair-associated genes. Such transcripts are normally either absent or transcribed at low basal levels in both immature and mature SC. For example, the transcription factor Jun was significantly transcribed and translated at P21 in the Mpz-*Braf* sciatic nerve (**Fig 6a-b**). Jun distinguishes repair SCs in injured adult nerves from immature SCs and represses myelination by opposing Egr2 [60].

We next sought to find out if secreted ligands specific to the repair phenotype, described as Jun-dependent, were also strongly upregulated in Mpz-*Braf* sciatic nerves. As early as P5, *Shh* (Sonic hedgehog) and the neurotrophic factor *Gdnf* (glial cell line-derived neurotrophic factor) [61] were significantly upregulated (**Fig 7c**). Cleaved Shh protein is important for multipotent NC maintenance [62] and its precursor is transcribed again in repair SCs upon nerve injury [63]. We confirmed by immunolabeling of fiber bundles (**Fig 7d**) and Western blot (**Fig. 7e**; **Fig S7e**) that Shh and S100b, a calcium-binding intermediate filament common to SCP and their derivatives, [64], were co-localized and significantly more expressed by P21 in Mpz-*Braf* mutant sciatic nerves than control nerves**.**

Nrg1-III and the Gdnf family member Artemin (Artn) are also secreted by Jun-positive repair SCs after injury to promote axon outgrowth [56,65]. While *Nrg1* was strongly upregulated at P5 and P21, *Artn* was only significantly overexpressed at P21 (**Fig. 7c**). An increase in transcription of the Nrg1 receptor *ErbB3*, a hallmark of nerve damage [66], was translated into significantly more protein at P21 and P42 (**Fig 7c, f-g**; **Fig S7b**). More Erbb3 protein immunofluorescence was also observed in mutant than control nerves, complementary to the Pmp2-positive myelin in individual cells (**Fig. 7f**; **Fig S7d**). We also found increased relative levels of the intermediate filament vimentin, encoded by *Vim*, over time in mutant sciatic nerves, phenocopying the local translation demonstrated in experimentally injured axons [67](**Fig. S7c; Table S2**). Despite heightened expression of repair program genes in Mpz-*Braf* sciatic nerves, pronounced axonal degeneration over time demonstrated that sustained MAPK signaling in SC was counterproductive *in vivo*.

Schwann cell differentiation of CFC patient-derived hiPSCs is impaired

The Mpz-*Braf* model allowed us to assess some direct molecular consequences of constitutive MAPK signaling on SC biology *in vivo*. However, $BRAF^{V600E}$ is not viable in the human (or mouse) germline. In order to assess the effects on SC of the heterozygous activating $BRAF^{Q257R}$ variant frequently found in CFC patients, but without the severe cardiac defects preventing postnatal study of existing CFC mouse models [68,69], we differentiated hiPSC lines from two unrelated CFC patients [70] into SC *in vitro* and compared their maturation with that of

two unrelated control hiPSC lines. Cells of the control lines were morphologically similar to each other and to the original publication throughout the directed differentiation protocol [71]. However, one of the mutant cell lines (RMK0138C) required more frequent passages to maintain a similar density and failed to develop the elongated, refringent soma and long projections characteristic of SCPs in the three other lines, even after 90 days *in vitro* and despite the same *BRAF* genotype (**Fig 8a**).

The differentiation dynamics of the four hiPSC-derived SC lines were assessed by RNAseq at two time points. We chose day (d)18, a committed SCP-like stage, and d31, for a stage more analogous to immature SC (**Fig 8a**). Due to their different phenotypes *in vitro*, we analyzed the mutant lines both together and individually for DEGs relative to control lines (**Table S3**). Pathway analyses of 380 DEGs common to both mutant cell lines at D31 using the human Reactome database [72] highlighted their unexpected downregulation of transcripts associated with epidermal cornification, including eight keratin genes (REAC:R-HSA-6809371; **Fig 8c**; **Table S3**). Another significantly downregulated pathway was "EGR2 and SOX10-mediated initiation of SC myelination" (REAC:R-HSA-9619665, **Fig 8c-d**). Among the downregulated transcripts therein were *EGR2* itself*, MBP*, *PRX* and *LAMA2* (encoding the alpha-2 chain of laminin-211, a major component of SC basal lamina).

In contrast, the transcription factor *POU3F2* was commonly upregulated in CFC patient-derived cell lines at both stages analyzed; **Table S3**). Functionally redundant *Pou3f1* and *Pou3f2* are transiently required in early murine SC, peaking in the first week after birth [73]. Annotations associated with all genes upregulated in concert with *POU3F2* in human SC differentiated from each CFC line, compared to the control hiPSC, evoked the multipotency of neural crest-derived SCPs. Significant associations included the term "melanoma" and the proteome of an amelanotic, $BRAF^{V600E}$-mutated melanoma cell line A2058; cell line proteomes from a *P53*-mutated lung neuroendocrine carcinoma (NCI-H446) and a $NRAS^{Q61L}$-mutated neuroblastoma (SK-N-AS), as well as genes transcribed in "visceral" (parasympathetic) autonomic neurons **(Fig 8e)**. These observations suggest that maintenance of MAPK signaling after birth in immature SC of both mice and humans prolongs their co-expression of transcriptional programs typical of other neural crest lineages [74,75]. Both *in vitro* and *in vivo*, the maintenance of otherwise physiologically transient MAPK pathway activation in cycling SC seems to prevent their terminal differentiation.

Discussion

A new model of RASopathy endophenotypes

In this work, we have demonstrated that constitutive activation of BRAF-mediated signaling in only a subpopulation of NC-derived cells is sufficient to phenocopy a subset of systemic effects found across multiple germline RASopathy syndromes. One major outcome was rapid onset of hyperplastic neuropathy, evocative of that described as "hypertrophic" in case reports and small patient series [25,27,76–79], but which had not been reported in systemic RASopathy animal models [80]. Clinical signs of neuropathy should therefore be more systematically sought across all RASopathy patients, as they converge mechanistically on sustained ERK1/2 activation in Schwann cells.

Severe feeding difficulties are frequent in *BRAF*-associated RASopathies, irrespective of the specific variant [25,81–83]. However, despite their motility issues, Mpz-*Braf* mice ate normally (see Fig 1f). No megacolon was observed in our mouse model at any age as a sign of gut dysmotility, while somatic growth delay due to Igf1 deficiency was well underway before weaning. Enteric glia may not have been efficiently targeted by the *Mpz*-Cre allele [84], as *Mpz* is transcribed at a basal level in postmigratory, Sox10-expressing SCP [85] but more strongly upregulated in myelinating SC [40,86]. Remarkably, some transcripts co-expressed specifically in adult mouse enteric glia, such as *Plp1* and *Gfap* [87], are upregulated over time in Mpz-*Braf* sciatic nerves (Supplemental Table S3). As enteric glia are also NC progeny, this observation suggests that the observed repair phenotype reverts SCPs to a state of commitment that precedes specialization of peripheral glial subtypes.

Unexpected endocrine phenotypes

The conditional Mpz-*Braf* mouse, with its pre-weaning growth retardation, also complements a series of RASopathy animal models that fail to thrive in the early postnatal period following germline mutations in *Kras* [88], *Braf* [89,90,68,69], *Raf1* [91], *Ptpn11* [92] or *Sos1* [93]. No other models were mosaics that restricted strong MAPK activation to cells having expressed *Mpz*. It was thus a surprise that somatic growth was affected.

Mpz-*Braf* mice do not display an effective second growth spurt, normally initiated by Gh from 12-25 days after birth [94], despite initially normal *Ghrhr* and *Gh* transcription by the pituitary. Our results implicate secondary, liver-produced Igf1 deficiency as responsible for small body size in both sexes, but not for the male cryptorchidism (undescended testicles). Postnatal *Igf1*-null mice, when they survive (5-15% depending on background), grow to less than half the size of wildtype littermates; no cryptorchidism was described despite infertility [95,96]. Knock-in of a conditional *Igf1* allele into the natural, but inoperative, *Igf1* locus in *Igf1*-null mice, followed by

activation of liver-restricted transcription, is sufficient to restore both body size and fertility to
both sexes [97].

Cryptorchidism affects the majority of males with RASopathies [6,98]; it is less frequent among
germline *BRAF*- or *RAF1*-mutated patients in one series than for other RASopathy genes [99].
This implies that a pathogenic threshold of maximal MAPK activation is less often attained by
typical RAF family gene mutations in the germline than by mutations in the upstream effectors
PTPN11, *RIT1* or *SOS2*. Furthermore, the *Mpz*-Braf model displays a fully penetrant
phenotype, despite *Mpz* not being expressed by germ cells in mice or humans. Surgical section
of the rat genitofemoral nerve even after puberty leads to neuropathic degeneration of the
cremaster muscle, partial testicular ascent and reduced fertility [100]. Our observations therefore
provide indirect support to the hypothesis that peripheral nerve myelination is also required to
preserve testicular descent and fertility [101,102].

Finally, NS patients and the *Ptpn11*$^{D61G/+}$ RASopathy mouse model for NS also show low Igf1
levels during post-weaning growth. Uncoupling Gh stimulation from Igf1 release by sustained
MAPK signaling downstream of Ghr is reminiscent of the uncoupling of Nrg1 stimulation from
Erbb3-mediated SC differentiation to a myelinating phenotype, and suggests that constitutive
somatic activation of MAPK-mediated pathways may lead to other human disorders. Evidence
for this also comes from the identification of septo-optic dysplasia in some CFC patients with
BRAF gain-of-function mutations, phenocopied in terms of preventing terminal differentiation
of the pituitary gland by conditional expression of murine *Braf* V600E/+ or *Braf* Q241R/+ in
Prop1-expressing pituitary progenitors [2].

A form of murine congenital hypomyelinating neuropathy

Dejerine-Sottas syndrome, or CMT3, is a demyelinating congenital peripheral neuropathy
caused by mutations in *MPZ*, *EGR2*, *PMP22* or *PRX* [103]. CMT3 often presents distal sensory
loss, profuse SC proliferation and as per its first report in 1893, enlarged, "gelatinous"
peripheral nerves that are more affected in the legs than the arms [104]. The tremor and
locomotor troubles we observed in Mpz-*Braf* mice also first affected the hindlimbs without
extending to other body areas before 6 weeks of age. Nonetheless, sensory and motor nerves
around the body were visibly enlarged on autopsy. These observations in the absence of
apparent suffering (aversion, unkempt fur, flattened ears, etc.) make the weanling Mpz-*Braf*
mouse a tractable model in which to carry out further behavioral or electrophysiological studies
of early RASopathy-associated neuropathy, or to test interventional therapies [105]. Our findings
support the Schwann cell as the initiator of these neuromuscular outcomes.

Oligopotent Schwann cell (SC) precursors, persisting both in ganglia and along postnatal
nerves, were first identified when they transdifferentiated into nerve-derived melanocytes after

nerve injury [74]. Mosaic expression of an oncogenic *BRAF* p.V600E variant causes a somatic

RASopathy in humans, congenital melanocytic nevus syndrome, that is not associated with

peripheral neuropathy [106,107]. Instead, bipotent progenitors proliferate *in utero* before settling

on a superficially differentiated skin phenotype after birth. In contrast, when induced

conditionally in either murine melanocyte progenitors with a tyrosinase-driven *Cre*, Braf^{V600E}

causes mid-gestational death with complete penetrance due to heart malformations [19].

Tyrosinase is expressed in mid-gestational NC cells before restriction to human postnatal

bipotent Schwann-melanocytes [108], and only expressed by a small minority of postnatal cardiac

melanocytes in mice [109]. It will be interesting to investigate if tyrosinase is inappropriately

maintained in cardiac NC cells after constitutive MAPK activation. Furthermore, *BRAF*

p.V600E oncogenic variants are also found in somatic, segmental lymphatic and vascular

malformations [110,111]. One of the earliest (but recent) case reports of arteriovenous

malformation with this genotype was in conjunction with a large intracranial gliosarcoma,

derived from brain progenitors capable of giving rise to myelinating and non-myelinating glia

[112]. Sustained MAPK signaling in multipotent cells therefore prevents terminal differentiation in

multiple prenatal and postnatal situations. This induces functional deficiencies despite ongoing

progenitor proliferation.

In our assay of human Schwann cell differentiation, the CFC patient cell line least capable of

acquiring a SC phenotype, 138C, instead maintained strong expression at day 31 of *HOX*

family genes of the first six paralogue groups as well as *PHOX2B*, a master transcription factor

for autonomic neurons and SCPs. Cells from both patients showed common significant

upregulation of *HOXB2*, *ARG2* and *SALL1* (**Table S3**). *ARG2* encodes arginase-2, a metabolic

enzyme significantly upregulated in response to tumor macrophage-stimulated ERK1/2 activity

in neuroblastoma [113]. This neural crest-derived congenital cancer derives from multipotent

Schwann/autonomic progenitors and usually presents in early childhood [114,115]. In high-risk

forms, tumor cells maintain an undifferentiated mesenchymal phenotype associated with *JUN*,

at the expense of *PHOX2B* expression [116]. *SALL1* (Spalt-like transcription factor 1) is a widely

transcribed but determinant transcription factor of central nervous system macrophages known

as microglia, but is not present in bone marrow-derived peripheral microglia [117]. Altogether, we

have shown that patient-derived, *BRAF*^{Q257R}-expressing NC cells are at a stochastic

disadvantage in assuming a differentiated SC phenotype *in vitro*, as compared to *BRAF*-

wildtype cells. Patient-derived nerve biopsies taken for molecular diagnosis would be

necessary to confirm these findings with a prospective study in unresolved cases of either

RASopathies or early-onset CMT.

Methods

Ethical statement

This study complies with all relevant ethical regulations. Animal care and experiments were in accordance with French and European legislation and approved by the French ACUC (C2EA-14; project numbers 37-08102012 and 9522-2017040517496865v5). Human iPSC studies adhered to the Declaration of Helsinki [118] and French bioethics law (2021-1017)[119] consistent with our institutional review board protocols.

Mouse breeding and genotyping

Transgenic *P0-Cre*[+/°]; *Braf*[V600E/+] and *Krox20-Cre*[+/°]; *Braf*[V600E/+] mice were generated by crossing *LSL-Braf*[V600E/+] mice [15] (RRID:IMSR_JAX:017928) with P0(Mpz)-*Cre* (RRID:IMSR_JAX:017927) [39,120] or Krox20-*Cre* (RRID:IMSR_JAX:025744) [121] drivers. Genotyping was performed by PCR (see Supplementary Methods). Mice were housed under standard conditions (12-hour light/dark cycles, *ad libitum* access to food/water).

Western Blots

Sciatic nerve lysates were prepared in RIPA buffer with protease inhibitors, separated by 12% SDS-PAGE, and probed with antibodies against phospho-Erk1/2 (RRID:AB_331646), Erk1/2 (RRID:AB_330744), c-Jun (RRID:AB_2130165) and Gapdh (RRID:AB_2107445). Alternately, primary antibodies to Shh (RRID:AB_528466)[122], Erbb3 (RRID:AB_358277), Egr2 (RRID:AB_10862073), alpha-tubulin (RRID:AB_2241126), vimentin (RRID:AB_2881439) or vinculin (RRID:AB_10603627) were used. Signals were detected using relevant secondary antibodies (RRID:AB_330924 and RRID:AB_2099233) followed by chemiluminescence detection and quantified with ImageJ v.1.54f.

Culture and differentiation of human pluripotent stem cells (hiPSC)

Two female patients with cardio-facio-cutaneous (CFC) syndrome bearing a heterozygous germline *BRAF* Q257R variant, or their legal guardians, provided written consent as approved by the UCSF Human Research Protection Program (CHR #10-02794), to produce the human induced pluripotent stem cell (hiPSC) lines, RMK0056C and RMK0138C, as described [70]. The AG08H and HFF15 clones of healthy donor male hiPSC were derived from commercial cell lines consented for research, AG08498 (RRID:CVCL_1Y51) and HFF-1 SCRC-1041 (RRID:CVCL_3285) as described [123]. Clones were initially grown in mTeSR1 (Stemcell) on human ES cell-qualified Matrigel™ before cryopreservation. Thawed hiPSC were cultured on Synthemax II-SC (Corning) at 0.025 mg/mL and maintained with daily changes of StemMACS iPS-Brew XF (Miltenyi Biotec). At 60-80% confluence, they were passed with ReleSR (StemCell Technologies), with 10 µM Y-27632 for the first day. Differentiation of hiPSCs into

491   SC [71] was initiated at 60% confluence and passed every 6 days until day 31. Thereafter, upon

492   90% confluence, they were split 1:2.

RNA sequencing and analyses

Sciatic nerves and pituitary glands from WT or $Braf^{V600E/+}$ animals, and pellets of WT or

$BRAF^{Q257R/+}$ hiPSC-derived SCP and immature Schwann cells (iSC) were stored at -80°C.

Tissues were lysed in the FastPrep-24 homogenizer (MP Biomedicals) and total RNA

extracted with the RNeasy Lipid Tissue Mini (Qiagen) or, for cell pellets, NucleoSpin RNA Plus

S (Macherey-Nagel) kits. RNA concentrations were measured on a NanoDrop TM 1000

Spectrophotometer (ThermoFisher). Libraries were prepared with the KAPA mRNA HyperPrep

kit (Roche) and sequenced on a NovaSeq 6000 (Illumina). Quality was assessed using fastQC

(v0.11.5). Raw sequencing reads were aligned to GRCm39/GRCh38 using STAR (v2.7.2b).

Bam files were indexed and sorted with Samtools (v1.7). After mapping, reads per feature

(Ensembl or GENCODE v34 annotations for mouse or human data, respectively) were

determined using StringTie v2.1.6, and differential expression analyzed with a Wald test in

DESeq2 (v1.34.0; FDR < 0.05). Data visualization made use of R packages Pheatmap and

EnhancedVolcano [124]. Expression-dependent enrichment analyses were conducted with Gene

Set Enrichment Analysis (GSEA) v4.3.2 [125] using mouse Wikipathway and human Reactome

databases, otherwise with g:Profiler [126] or Enrichr-KG [127], interrogating quantitative proteomics

of the Cancer Cell Line Encyclopedia [128] KEGG, Reactome, Gene Ontology datasets.

These were significantly enriched by Enrichr-KG [127], using databases of annotations related to

human disease [129], and a large single-cell atlas of developmental human gene expression [130].

Nerve and fiber immunostaining

Sciatic nerves were fixed in 4% buffered paraformaldehyde and stained with primary

antibodies to Mbp (RRID:AB_94975); Ki-67 (RRID:AB_10854564); Caspr (RRID:AB_869934);

Sox10 (RRID:AB_2941085); Pmp2 (RRID:AB_2166978); Erbb3 (RRID:AB_358277); S100b

(RRID:AB_1856538); or Shh (RRID:AB_528466)[122], followed by secondary antibodies and

DAPI counterstaining. Further details are in the Supplemental Methods. Image stacks of

teased fibers were acquired with a spinning disk confocal microscope (Nikon SoRa) and NIS-

Elements software and nerve images on a Zeiss LSM800 with Airyscan and Zen software v2.3.

Transmission electron microscopy (TEM) and morphometry

Sciatic nerves were fixed in 2% PFA + 2.5% glutaraldehyde, post-fixed with 1% OsO_4, contrasted in 1% uranyl acetate before Epon resin embedding. Semi-thin sections at 1 μm were counterstained with toluidine blue. Ultra-thin sections were imaged on an FEI Morgagni 120 kv microscope. Myelin morphological parameters were analyzed with MyelTracer [43]. Axon density was assessed with ImageJ using the semi-thin sections.

Statistics

Statistical analyses were performed with Prism v10.2.0 (GraphPad), using unpaired Mann-Whitney or two-way ANOVA non-parametric tests. For multiple comparisons, two-way ANOVA followed by Šidák's post-hoc tests, or Student t-tests followed by Benjamini-Hochberg adjustment, were used. Mendelian segregation was assessed with a chi-square contingency test. Data are presented as mean ± SEM; significance: $* p < 0.05$; $** p < 0.01$; $*** p < 0.001$; $**** p < 0.0001$.

Data availability

Raw and processed RNA-sequencing data from this study have been deposited to the Gene Expression Omnibus (GEO) and assigned the accession identifiers GSE262047 and GSE262048.

Websites

"Catalogue Of Somatic Mutations In Cancer" (COSMIC), https://cancer.sanger.ac.uk/cosmic [1].

Rare disease ORPHA codes are maintained at https://www.orpha.net and available on https://www.orphadata.com/ [18].

fastQC (v0.11.5) is available at https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ and EnhancedVolcano at https://bioconductor.org/packages/EnhancedVolcano.

NIH Fiji/ImageJ software (https://imagej.net/ij/) is available from https://imagej.net/ij/.

Author contributions

E.M. wrote the first draft, performed and analyzed experiments; D.A. provided analyses, performed experiments and extensively revised the manuscript; P.Q., G.M. and M.M. performed experiments; N.B.M. and V.D. provided protocols and guidance; C.E.Y. and N.B. provided control hiPSC lines and guidance; L.A.W. provided patient hiPSC lines; A.B. provided constructive criticism and funding; H.C.E. supervised the experiments, provided funding, wrote and revised the article.

Competing interests

The authors declare no competing interests.

Acknowledgements

We are grateful to C. Humbert, C. Castro, N. Trévisiol and C. Aubert of the MMG Genomics and Bioinformatics facility (GBiM) for assistance with RNAseq and analyses; Dr. F. Magdinier and O. Hadadeh of the Marseille Stem Cell (MaSC) core facility for control hiPSC cell lines and their quality control; Dr. T.-T. Mac, Adeline Querdray and Dr. T. Fauquier for their technical assistance and advice; the PiCSL-France BioImaging core facility, supported by ANR-10-INBS-04, and Dr. C. Leterrier and Dr. L. Fourel of the Institute of NeuroPhysiopathology, CNRS-Aix Marseille University, for their assistance with spinning-disk confocal microscopy.

Funding

The authors declare that they have no conflicts of interest. This work was made possible by the Association Française contre les Myopathies (AFM-Téléthon) grant 'MoThARD' to VD and HCE, the National Institutes of Health New Innovator award (1DP2OD007449) to LAW, and Horizon Europe HORIZON-MISS-2021-CANCER-02-03 MELCAYA (101096667) to HCE. EM was supported by the AFM-Téléthon, the Asociación Española de Nevus Gigante Congénito (AsoNevus, Spain), Naevus 2000 France-Europe (France) and the Marseille Rare Diseases Institute (MarMaRa) of Aix-Marseille University.

References

1. Sondka, Z. *et al.* COSMIC: a curated database of somatic variants and clinical data for cancer. *Nucleic Acids Res.* **52**, D1210–D1217 (2024).

2. Gualtieri, A. *et al.* Activating mutations in BRAF disrupt the hypothalamo-pituitary axis leading to hypopituitarism in mice and humans. *Nat. Commun.* **12**, 2028 (2021).

3. Hilal, N., Chen, Z., Chen, M. H. & Choudhury, S. RASopathies and cardiac manifestations. *Front. Cardiovasc. Med.* **10**, 1176828 (2023).

4. Jaeger, Z. J. *et al.* RASopathies. Part II: Cutaneous and extracutaneous manifestations. *J. Am. Acad. Dermatol.* S0190-9622(25)02340–0 (2025) doi:10.1016/j.jaad.2025.01.106.

5. Kang, M. & Lee, Y.-S. The impact of RASopathy-associated mutations on CNS development in mice and humans. *Mol. Brain* **12**, 96 (2019).

6.   Rauen, K. A. The RASopathies. *Annu. Rev. Genomics Hum. Genet.* **14**, 355–369 (2013).

7.   Siano, M. A. *et al.* Endocrine system involvement in patients with RASopathies: A case

series. *Front. Endocrinol.* **13**, (2022).

8.   Tartaglia, M., Aoki, Y. & Gelb, B. D. The molecular genetics of RASopathies: An update

on novel disease genes and new disorders. *Am. J. Med. Genet. C Semin. Med. Genet.*

190, 425–439 (2022).

9.   Carli, D., Resta, N., Ferrero, G. B., Ruggieri, M. & Mussa, A. Mosaic RASopathies: A

review of disorders caused by somatic pathogenic variants in the genes of the

RAS/MAPK pathway. *Am. J. Med. Genet. C Semin. Med. Genet.* **190**, 520–529 (2022).

10.   Kratz, C. P., Rapisuwon, S., Reed, H., Hasle, H. & Rosenberg, P. S. Cancer in Noonan,

Costello, Cardiofaciocutaneous and LEOPARD Syndromes. *Am. J. Med. Genet. Part C*

157, 83–89 (2011).

11.   Peltonen, S. *et al.* Pediatric malignancies in neurofibromatosis type 1: A population-based

cohort study. *Int. J. Cancer* **145**, 2926–2932 (2019).

12.   Davies, H. *et al.* Mutations of the BRAF gene in human cancer. *Nature* **417**, 949–954

(2002).

13.   Kawakami, H. *et al.* Mutant BRAF Upregulates MCL-1 to Confer Apoptosis Resistance

that Is Reversed by MCL-1 Antagonism and Cobimetinib in Colorectal Cancer. *Mol.*

Cancer Ther. **15**, 3015–3027 (2016).

14.   Xing, M. BRAF mutation in thyroid cancer. *Endocr. Relat. Cancer* **12**, 245–262 (2005).

15.   Mercer, K. *et al.* Expression of endogenous oncogenic V600EB-raf induces proliferation

and developmental defects in mice and transformation of primary fibroblasts. *Cancer Res.*

65, 11493–11500 (2005).

16.   Haling, J. R. *et al.* Structure of the BRAF-MEK Complex Reveals a Kinase Activity

Independent Role for BRAF in MAPK Signaling. *Cancer Cell* **26**, 402–413 (2014).

17.   Yao, Z. *et al.* BRAF Mutants Evade ERK-Dependent Feedback by Different Mechanisms

that Determine Their Sensitivity to Pharmacologic Inhibition. *Cancer Cell* **28**, 370–383

(2015).

18.   Aymé, S., Bellet, B. & Rath, A. Rare diseases in ICD11: making rare diseases visible in

health information systems through appropriate coding. *Orphanet J. Rare Dis.* **10**, 35

(2015).

19.   Dhomen, N. *et al.* Inducible expression of (V600E) Braf using tyrosinase-driven Cre

recombinase results in embryonic lethality. *Pigment Cell Melanoma Res.* **23**, 112–20

(2010).

20.   Gripp, K. W. *et al.* Further delineation of the phenotype resulting from BRAF or MEK1

germline mutations helps differentiate cardio-facio-cutaneous syndrome from Costello

syndrome. *Am. J. Med. Genet. A.* **143A**, 1472–1480 (2007).

21.   Koudova, M., Seemanova, E. & Zenker, M. Novel BRAF mutation in a patient with

LEOPARD syndrome and normal intelligence. *Eur. J. Med. Genet.* **52**, 337–340 (2009).

22.   Nava, C. *et al.* Cardio-facio-cutaneous and Noonan syndromes due to mutations in the

RAS/MAPK signalling pathway: genotype-phenotype relationships and overlap with

Costello syndrome. *J. Med. Genet.* **44**, 763–71 (2007).

23.   Nyström, A.-M. *et al.* Noonan and cardio-facio-cutaneous syndromes: two clinically and

genetically overlapping disorders. *J. Med. Genet.* **45**, 500–506 (2008).

24.   Tartaglia, M., Gelb, B. D. & Zenker, M. Noonan syndrome and clinically related disorders.

Best Pract. Res. Clin. Endocrinol. Metab. **25**, 161–179 (2011).

25.   Manci, E. A. *et al.* Cardiofaciocutaneous syndrome (CFC) with congenital peripheral

neuropathy and nonorganic malnutrition: an autopsy study. *Am. J. Med. Genet. A.* **137**,

1–8 (2005).

26.   De Ridder, W., van Engelen, B. & van Alfen, N. Neurological features of Noonan

syndrome and related RASopathies: Pain and nerve enlargement characterized by nerve

ultrasound. *Am. J. Med. Genet. A.* **188**, 1801–1807 (2022).

27.   Draaisma, F. *et al.* Hypertrophic neuropathy: a possible cause of pain in children with

Noonan syndrome and related disorders. *Eur. J. Pediatr.* **182**, 3789–3793 (2023).

28. Callegari, I. *et al.* Mutation update for myelin protein zero-related neuropathies and the
increasing role of variants causing a late-onset phenotype. *J. Neurol.* **266**, 2629–2645
(2019).

29. Ishii, A., Furusho, M. & Bansal, R. Sustained activation of ERK1/2 MAPK in
oligodendrocytes and schwann cells enhances myelin growth and stimulates
oligodendrocyte progenitor expansion. *J. Neurosci. Off. J. Soc. Neurosci.* **33**, 175–186
(2013).

30. Sheean, M. E. *et al.* Activation of MAPK overrides the termination of myelin growth and
replaces Nrg1/ErbB3 signals during Schwann cell development and myelination. *Genes*
Dev. **28**, 290–303 (2014).

31. Belin, S. *et al.* Neuregulin 1 type III improves peripheral nerve myelination in a mouse
model of congenital hypomyelinating neuropathy. *Hum. Mol. Genet.* **28**, 1260–1273
(2019).

32. Ishii, A., Furusho, M., Dupree, J. L. & Bansal, R. Strength of ERK1/2 MAPK Activation
Determines Its Effect on Myelin and Axonal Integrity in the Adult CNS. *J. Neurosci. Off.*
J. Soc. Neurosci. **36**, 6471–6487 (2016).

33. Pipis, M., Rossor, A. M., Laura, M. & Reilly, M. M. Next-generation sequencing in
Charcot–Marie–Tooth disease: opportunities and challenges. *Nat. Rev. Neurol.* **15**, 644–
656 (2019).

34. Eid, S. A. *et al.* Single-cell RNA-seq uncovers novel metabolic functions of Schwann cells
beyond myelination. *J. Neurochem.* **166**, 367–388 (2023).

35. Geren, B. B. The formation from the schwann cell surfaceof myelin in the peripheral
nerves of chick embryos. *Exp. Cell Res.* **7**, 558–562 (1954).

36. Le Douarin, N., Dulac, C., Dupin, E. & Cameron-Curry, P. Glial cell lineages in the neural
crest. *Glia* **4**, 175–84 (1991).

37. Etchevers, H. C., Dupin, E. & Le Douarin, N. M. The diverse neural crest: from
embryology to human pathology. *Development* **146**, (2019).

38.   Kelnar, C. J. H. Noonan syndrome: the hypothalamo-adrenal and hypothalamo-gonadal
664   axes. *Horm. Res.* **72 Suppl 2**, 24–30 (2009).

665   39.   Feltri, M. L. *et al.* A novel P 0 glycoprotein transgene activates expression of lac Z in
666   myelin-forming Schwann cells. *Eur. J. Neurosci.* **11**, 1577–1586 (1999).

667   40.   Hagedorn, L., Suter, U. & Sommer, L. P0 and PMP22 mark a multipotent neural crest-
668   derived cell type that displays community effects in response to TGF-beta family factors.
669   *Development* **126**, 3781–3794 (1999).

670   41.   Lundberg, E., Kriström, B., Jonsson, B., Albertsson-Wikland, K., & study group. Growth
671   hormone (GH) dose-dependent IGF-I response relates to pubertal height gain. *BMC*
672   *Endocr. Disord.* **15**, 84 (2015).

673   42.   Alvey, L. M., Jones, J. F. X., Tobin-O'Brien, C. & Pickering, M. Bands of Fontana are
674   caused exclusively by the sinusoidal path of axons in peripheral nerves and predict axon
675   path; evidence from rodent nerves and physical models. *J. Anat.* **234**, 165–178 (2019).

676   43.   Kaiser, T. *et al.* MyelTracer: A Semi-Automated Software for Myelin g-Ratio
677   Quantification. *eNeuro* **8**, ENEURO.0558-20.2021 (2021).

678   44.   Kozlowski, M. M. *et al.* Pmp2+ Schwann Cells Maintain the Survival of Large-Caliber
679   Motor Axons. *J. Neurosci.* **45**, e1362242025 (2025).

680   45.   Woodhoo, A. & Sommer, L. Development of the Schwann cell lineage: from the neural
681   crest to the myelinated nerve. *Glia* **56**, 1481–1490 (2008).

682   46.   Kuhlbrodt, K., Herbarth, B., Sock, E., Hermans-Borgmeyer, I. & Wegner, M. Sox10, a
683   novel transcriptional modulator in glial cells. *J. Neurosci.* **18**, 237–250 (1998).

684   47.   Gerdes, J. *et al.* Cell cycle analysis of a cell proliferation-associated human nuclear
685   antigen defined by the monoclonal antibody Ki-67. *J. Immunol. Baltim. Md 1950* **133**,
686   1710–1715 (1984).

687   48.   Miller, I. *et al.* Ki67 is a Graded Rather than a Binary Marker of Proliferation versus
688   Quiescence. *Cell Rep.* **24**, 1105-1112.e5 (2018).

689   49.   Poitelon, Y., Kopec, A. M. & Belin, S. Myelin Fat Facts: An Overview of Lipids and Fatty
690   Acid Metabolism. *Cells* **9**, 812 (2020).

50.   Ohashi, K. *et al.* Early embryonic lethality caused by targeted disruption of the 3-hydroxy-

3-methylglutaryl-CoA reductase gene. *J. Biol. Chem.* **278**, 42936–42941 (2003).

51.   Maldonado, E. N., Alderson, N. L., Monje, P. V., Wood, P. M. & Hama, H. FA2H is

responsible for the formation of 2-hydroxy galactolipids in peripheral nervous system

myelin. *J. Lipid Res.* **49**, 153–161 (2008).

52.   Topilko, P. *et al.* Krox-20 controls myelination in the peripheral nervous system. *Nature*

371, 796–799 (1994).

53.   Peirano, R. I., Goerich, D. E., Riethmacher, D. & Wegner, M. Protein zero gene

expression is regulated by the glial transcription factor Sox10. *Mol. Cell. Biol.* **20**, 3198–

3209 (2000).

54.   Jessen, K. R. & Mirsky, R. Negative regulation of myelination: relevance for development,

injury, and demyelinating disease. *Glia* **56**, 1552–1565 (2008).

55.   Hong, J. *et al.* PMP2 regulates myelin thickening and ATP production during

remyelination. *Glia* **72**, 885–898 (2024).

56.   Stassart, R. M. *et al.* A role for Schwann cell-derived neuregulin-1 in remyelination. *Nat.*

Neurosci. **16**, 48–54 (2013).

57.   Heming, M. *et al.* Multi-omic identification of perineurial hyperplasia and lipid-associated

nerve macrophages in human polyneuropathies. *Nat. Commun.* **16**, 7872 (2025).

58.   Serra-Nédélec, A. D. R. *et al.* Noonan syndrome-causing SHP2 mutants inhibit insulin-

like growth factor 1 release via growth hormone-induced ERK hyperactivation, which

contributes to short stature. *Proc. Natl. Acad. Sci.* **109**, 4257–4262 (2012).

59.   Arthur-Farraj, P. J. *et al.* Changes in the Coding and Non-coding Transcriptome and DNA

Methylome that Define the Schwann Cell Repair Phenotype after Nerve Injury. *Cell Rep.*

20, 2719–2734 (2017).

60.   Parkinson, D. B. *et al.* c-Jun is a negative regulator of myelination. *J. Cell Biol.* **181**, 625–

637 (2008).

61.   Piirsoo, M., Kaljas, A., Tamm, K. & Timmusk, T. Expression of NGF and GDNF family

members and their receptors during peripheral nerve development and differentiation of

Schwann cells in vitro. *Neurosci. Lett.* **469**, 135–140 (2010).

62.   da Costa, M. C., Trentin, A. G. & Calloni, G. W. FGF8 and Shh promote the survival and

maintenance of multipotent neural crest progenitors. *Mech. Dev.* **154**, 251–258 (2018).

63.   Hashimoto, M. *et al.* Neuroprotective effect of sonic hedgehog up-regulated in Schwann

cells following sciatic nerve injury. *J. Neurochem.* **107**, 918–927 (2008).

64.   Takahashi, K. *et al.* Immunohistochemical study on the distribution of alpha and beta

subunits of S-100 protein in human neoplasm and normal tissues. *Virchows Arch. B Cell*

Pathol. Incl. Mol. Pathol. **45**, 385–396 (1984).

65.   Huang, L. *et al.* c-Jun gene-modified Schwann cells: upregulating multiple neurotrophic

factors and promoting neurite outgrowth. *Tissue Eng. Part A* **21**, 1409–1421 (2015).

66.   Carroll, S. L., Miller, M. L., Frohnert, P. W., Kim, S. S. & Corbett, J. A. Expression of

neuregulins and their putative receptors, ErbB2 and ErbB3, is induced during Wallerian

degeneration. *J. Neurosci. Off. J. Soc. Neurosci.* **17**, 1642–1659 (1997).

67.   Perlson, E. *et al.* Vimentin-dependent spatial translocation of an activated MAP kinase in

injured nerve. *Neuron* **45**, 715–726 (2005).

68.   Inoue, S.-I. *et al.* New BRAF knockin mice provide a pathogenetic mechanism of

developmental defects and a therapeutic approach in cardio-facio-cutaneous syndrome.

Hum. Mol. Genet. **23**, 6553–6566 (2014).

69.   Moriya, M. *et al.* Adult mice expressing a Braf Q241R mutation on an ICR/CD-1

background exhibit a cardio-facio-cutaneous syndrome phenotype. *Hum. Mol. Genet.* **24**,

7349–7360 (2015).

70.   Yeh, E. *et al.* Patient-derived iPSCs show premature neural differentiation and neuron

type-specific phenotypes relevant to neurodevelopment. *Mol. Psychiatry* **23**, 1687–1698

(2018).

71.   Hörner, S. J. *et al.* hiPSC-Derived Schwann Cells Influence Myogenic Differentiation in

Neuromuscular Cocultures. *Cells* **10**, 3292 (2021).

72. Milacic, M. *et al.* The Reactome Pathway Knowledgebase 2024. *Nucleic Acids Res.* **52**,

D672–D678 (2024).

73. Jaegle, M. *et al.* The POU proteins Brn-2 and Oct-6 share important functions in Schwann

cell development. *Genes Dev.* **17**, 1380–1391 (2003).

74. Adameyko, I. *et al.* Schwann cell precursors from nerve innervation are a cellular origin

of melanocytes in skin. *Cell* **139**, 366–379 (2009).

75. Furlan, A. *et al.* Multipotent peripheral glial cells generate neuroendocrine cells of the

adrenal medulla. *Science* **357**, eaal3753 (2017).

76. Ando, Y. *et al.* A novel KRAS gene mutation of Noonan syndrome with severe peripheral

nerve hypertrophy. *J. Neurol. Sci.* **333**, e605 (2013).

77. DeRoos, S. T., Ryan, M. M. & Ouvrier, R. A. Peripheral neuropathy in

cardiofaciocutaneous syndrome. *Pediatr. Neurol.* **36**, 250–252 (2007).

78. Maridet, C., Sole, G., Morice-Picard, F. & Taieb, A. Hypertrophic neuropathy in Noonan

syndrome with multiple lentigines. *Am. J. Med. Genet. A.* **170**, 1570–1572 (2016).

79. Vizcaino, M. A. *et al.* Localized Hypertrophic Neuropathy as a Neoplastic Manifestation

of KRAS-Mediated RASopathy. *J. Neuropathol. Exp. Neurol.* **79**, 647–651 (2020).

80. Hernández-Porras, I. & Guerra, C. Modeling RASopathies with Genetically Modified

Mouse Models. *Methods Mol. Biol. Clifton NJ* **1487**, 379–408 (2017).

81. Allanson, J. E. *et al.* Cardio-facio-cutaneous syndrome: Does genotype predict

phenotype? *Am. J. Med. Genet. C Semin. Med. Genet.* **157**, 129–135 (2011).

82. Feng, B. *et al.* Molecular and phenotypic spectrum of cardio-facio-cutaneous syndrome

in Chinese patients. *Orphanet J. Rare Dis.* **18**, 284 (2023).

83. Lee, Y. *et al.* Clinical and molecular spectra of BRAF-associated RASopathy. *J. Hum.*

Genet. **66**, 389–399 (2021).

84. Woods, C. *et al.* Neurons populating the rectal extrinsic nerves in humans express

neuronal and Schwann cell markers. *Neurogastroenterol. Motil.* **33**, e14074 (2021).

85. Kunke, M. *et al.* SOX10-Mediated Regulation of Enteric Glial Phenotype in vitro and its

Relevance for Neuroinflammatory Disorders. *J. Mol. Neurosci.* **75**, 26 (2025).

86. Lee, M. J. *et al.* P0 is constitutively expressed in the rat neural crest and embryonic nerves

and is negatively and positively regulated by axons to generate non- myelin-forming and

myelin-forming Schwann cells, respectively. *Mol. Cell. Neurosci.* **8**, 336–350 (1997).

87. Rao, M. *et al.* Enteric glia express proteolipid protein 1 and are a transcriptionally unique

population of glia in the mammalian nervous system. *Glia* **63**, 2040–2057 (2015).

88. Hernández-Porras, I. *et al.* K-RasV14I recapitulates Noonan syndrome in mice. *Proc.*

Natl. Acad. Sci. U. S. A. **111**, 16395–16400 (2014).

89. Urosevic, J. *et al.* Constitutive activation of B-Raf in the mouse germ line provides a model

for human cardio-facio-cutaneous syndrome. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 5015–

5020 (2011).

90. Andreadi, C. *et al.* The intermediate-activity L597VBRAF mutant acts as an epistatic

modifier of oncogenic RAS by enhancing signaling through the RAF/MEK/ERK pathway.

Genes Dev. **26**, 1945–1958 (2012).

91. Wu, X. *et al.* MEK-ERK pathway modulation ameliorates disease phenotypes in a mouse

model of Noonan syndrome associated with the Raf1(L613V) mutation. *J. Clin. Invest.*

121, 1009–1025 (2011).

92. Araki, T. *et al.* Mouse model of Noonan syndrome reveals cell type- and gene dosage-

dependent effects of Ptpn11 mutation. *Nat. Med.* **10**, 849–857 (2004).

93. Chen, P.-C. *et al.* Activation of multiple signaling pathways causes developmental defects

in mice with a Noonan syndrome–associated Sos1 mutation. *J. Clin. Invest.* **120**, 4353–

4365 (2010).

94. Lupu, F., Terwilliger, J. D., Lee, K., Segre, G. V. & Efstratiadis, A. Roles of Growth

Hormone and Insulin-like Growth Factor 1 in Mouse Postnatal Growth. *Dev. Biol.* **229**,

141–162 (2001).

95. Baker, J., Liu, J.-P., Robertson, E. J. & Efstratiadis, A. Role of insulin-like growth factors

in embryonic and postnatal growth. *Cell* **75**, 73–82 (1993).

96. Powell-Braxton, L. *et al.* IGF-I is required for normal embryonic growth in mice. *Genes*

Dev. **7**, 2609–2617 (1993).

97. Stratikopoulos, E., Szabolcs, M., Dragatsis, I., Klinakis, A. & Efstratiadis, A. The hormonal action of IGF1 in postnatal mouse growth. *Proc. Natl. Acad. Sci.* **105**, 19378–19383 (2008).

98. Champion, K. J. *et al.* Germline mutation in BRAF codon 600 is compatible with human development: De novo p.V600G mutation identified in a patient with CFC syndrome. *Clin. Genet.* **79**, 468–474 (2011).

99. Reynolds, G. *et al.* Update on the Clinical and Molecular Characterization of Noonan Syndrome and Other RASopathies: A Retrospective Study and Systematic Review. *Int. J. Mol. Sci.* **26**, 3515 (2025).

100. Gutiérrez-Pérez, O., Rojas-Castañeda, J. C., Chavez-Saldaña, M., Reyes, G. & Vigueras-Villaseñor, R. M. Infertility in rats subjected to genitofemoral nerve section is not associated with testicular damage. *Andrologia* **46**, 151–157 (2014).

101. Prince, F. P. Ultrastructural evidence of indirect and direct autonomic innervation of human Leydig cells: comparison of neonatal, childhood and pubertal ages. *Cell Tissue Res.* **269**, 383–390 (1992).

102. Tanyel, F. C. Commentary: The fate of germ cells in cryptorchid testis. *Front. Endocrinol.* **15**, 1393040 (2024).

103. Pearce, J. M. S. Dejerine-Sottas Disease (Progressive Hypertrophic Polyneuropathy). *Eur. Neurol.* **55**, 115–117 (2006).

104. Déjerine, J. & Sottas, J. Sur la névrite interstitielle hypertrophique et progressive de l'enfance. *Comp Rend Soc Biol Paris* **45**, 63–96 (1893).

105. Duman, M., Jaggi, S., Enz, L. S., Jacob, C. & Schaeren-Wiemers, N. Theophylline Induces Remyelination and Functional Recovery in a Mouse Model of Peripheral Neuropathy. *Biomedicines* **10**, 1418 (2022).

106. Etchevers, H. C. *et al.* Giant congenital melanocytic nevus with vascular malformation and epidermal cysts associated with a somatic activating mutation in BRAF. *Pigment Cell Melanoma Res.* **31**, 437–441 (2018).

107. Salgado, C. M. *et al.* BRAF Mutations are Also Associated with Neurocutaneous

Melanocytosis and Large/Giant Congenital Melanocytic Nevi. *Pediatr. Dev. Pathol.* **18**,

1–9 (2015).

108. Dusart, P. *et al.* A tissue centric atlas of cell type transcriptome enrichment signatures.

2023.01.10.520698 Preprint at https://doi.org/10.1101/2023.01.10.520698 (2023).

109. Yajima, I. *et al.* A subpopulation of smooth muscle cells, derived from melanocyte-

competent precursors, prevents patent ductus arteriosus. *PloS One* **8**, e53183 (2013).

110. Goss, J. A. *et al.* Somatic mutations in intracranial arteriovenous malformations. *PloS*

One **14**, e0226852 (2019).

111. Zenner, K. *et al.* Somatic activating BRAF variants cause isolated lymphatic

malformations. *HGG Adv.* **3**, 100101 (2022).

112. Zeng, X. *et al.* The Coexistence of Gliosarcoma and Arteriovenous Malformation with the

BRAF V600E Mutation. *World Neurosurg.* **134**, 594–597 (2020).

113. Fultang, L. *et al.* Macrophage-Derived IL1β and TNFα Regulate Arginine Metabolism in

Neuroblastoma. *Cancer Res.* **79**, 611–624 (2019).

114. Bedoya-Reina, O. C. *et al.* Single-nuclei transcriptomes from human adrenal gland reveal

distinct cellular identities of low and high-risk neuroblastoma tumors. *Nat. Commun.* **12**,

5309 (2021).

115. Jansky, S. *et al.* Single-cell transcriptomic analyses provide insights into the

developmental origins of neuroblastoma. *Nat. Genet.* **53**, 683–693 (2021).

116. Boeva, V. *et al.* Heterogeneity of neuroblastoma cell identity defined by transcriptional

circuitries. *Nat. Genet.* **49**, 1408–1413 (2017).

117. Buttgereit, A. *et al.* Sall1 is a transcriptional regulator defining microglia identity and

function. *Nat. Immunol.* **17**, 1397–1406 (2016).

118. The World Medical Association, (WMA). WMA Declaration of Helsinki. *Ethical Principles*

for Medical Research Involving Human Subjects https://www.wma.net/policies-post/wma-

declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/

(2022).

119. Journal Officiel, F. *LOI n° 2021-1017 du 2 août 2021 relative à la bioéthique*. (2021).

120. Feltri, M. L. *et al.* P0-Cre transgenic mice for inactivation of adhesion molecules in
858   Schwann cells. *Ann. N. Y. Acad. Sci.* **883**, 116–123 (1999).

859   121. Voiculescu, O., Charnay, P. & Schneider-Maunoury, S. Expression pattern of a Krox-
860   20/Cre knock-in allele in the developing hindbrain, bones, and peripheral nervous system.
861   *Genes. N. Y. N 2000* **26**, 123–126 (2000).

862   122. Ericson, J., Morton, S., Kawakami, A., Roelink, H. & Jessell, T. M. Two critical periods of
863   Sonic Hedgehog signaling required for the specification of motor neuron identity. *Cell* **87**,
864   661–673 (1996).

865   123. Dion, C. *et al.* SMCHD1 is involved in de novo methylation of the DUX4-encoding D4Z4
866   macrosatellite. *Nucleic Acids Res.* **47**, 2822–2839 (2019).

867   124. Blighe, K., Rana, S. & Lewis, M. Blighe K, Rana S, Lewis M (2023). EnhancedVolcano:
868   Publication-ready volcano plots with enhanced coloring and labeling. R package version
869   1.20.0. (2023).

870   125. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
871   interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci.* **102**, 15545–15550
872   (2005).

873   126. Kolberg, L. *et al.* g:Profiler—interoperable web service for functional enrichment analysis
874   and gene identifier mapping (2023 update). *Nucleic Acids Res.* **51**, W207–W212 (2023).

875   127. Evangelista, J. E. *et al.* Enrichr-KG: bridging enrichment analysis across multiple libraries.
876   *Nucleic Acids Res.* **51**, W168–W179 (2023).

877   128. Nusinow, D. P. *et al.* Quantitative Proteomics of the Cancer Cell Line Encyclopedia. *Cell*
878   **180**, 387-402.e16 (2020).

879   129. Grissa, D., Junge, A., Oprea, T. I. & Jensen, L. J. Diseases 2.0: a weekly updated
880   database of disease–gene associations from text mining and data integration. *Database*
881   **2022**, baac019 (2022).

130. Cao, J. *et al.* A human cell atlas of fetal gene expression. *Science* **370**, eaba7721 (2020).

131. Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z. & Zhang, B. WebGestalt 2019: gene set analysis

toolkit with revamped UIs and APIs. *Nucleic Acids Res.* **47**, W199–W205 (2019).

132. Le, N. *et al.* Analysis of congenital hypomyelinating Egr2Lo/Lo nerves identifies Sox2 as

an inhibitor of Schwann cell differentiation and myelination. *Proc. Natl. Acad. Sci. U. S.*

A. **102**, 2596–2601 (2005).

 # Figure legends

Figure 1. Constitutive MAPK activation mediated by a single copy of $Braf^{V600E}$ in murine
Schwann cell progenitors induces early peripheral neuropathy and small body size.

a) By postnatal day (P)21, Mpz-*Braf* mice displayed abnormal hindlimb postural reflexes and
ability to keep hindfeet on the bars of a wire cage lid (arrowheads), indicating a functional
peripheral nerve defect. **b, c)** Western blot analysis of total and phosphorylated (p)Erk1/2
signaling performed on sciatic nerve lysates from **(b)** P5 and **(c)** P21 control and Mpz-*Braf*
mice. Gapdh was detected as a loading control. **d)** Protein densitometry in control (n=4) *vs*
mutant (n=5) sciatic nerve lysates at P5, and control (n=13) *vs* mutant (n=15) lysates at P21.
e) Body weight of both sexes was monitored at P5, P10, P21, P35 and P42 in control and Mpz-
Braf mice, respectively (n = 28 *vs* 8 at P5, n = 15 *vs* 5 at P10, n = 54 *vs* 45 at P21, n = 14 *vs*
10 at P35, n = 5 *vs* 5 at P42). **f**) Quantitative PCR analysis of *Ghr* and *Igf1* expression in control
(n=5) *vs* mutant (n=5) livers at P21. Statistical significance from unpaired Mann-Whitney test
in **d**), two-way ANOVA followed by Šidák's post-hoc test in **f**), and Student's t-test in **g**): ns, not
significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

Figure 2. Myelination initiation is not impaired in *Braf*-mutant sciatic nerves, while
maintenance is compromised.

a) Macroscopic examination of control and mutant sciatic nerve at P5 and P21. Arrows indicate
Fontana bands, present only in control nerves. **b)** Sciatic nerve diameter measurements at P5,
P10, P21, and P35 in both Mpz-*Braf* and control mice (n = 11 *vs* 6 at P5, n = 6 *vs* 4 at P10, n
= 15 *vs* 10 at P21, and n = 5 *vs* 5 at P35; in control *vs* mutant nerves, respectively). Two-way
ANOVA followed by Šidák's post-hoc test; ns, not significant; ** $p < 0.01$; **** $p < 0.0001$. **c)**
Representative transmission electron micrographs of sciatic nerve cross-sections at P5 and
P21 from control and mutant mice. Bars = 10 µm except for magnified axons at P28, 2 µm.
Arrows, myelin aberrations; c, extracellular collagen; f, fibroblast; nm, non-myelinating
Schwann cell; sc, myelinating Schwann cell;.

Figure 3. $Braf^{V600E}$ in Schwann cells triggers demyelination and degeneration of large-
caliber axons

a) Mean axon diameter at P5 (n = 3 per genotype, not significantly [ns] different). **b**)
MyelTracer-obtained *g*-ratios as a function of axon diameter at P5, to assess myelin thickness
in control (n = 803) *vs* mutant (n=811) myelinated fibers (n=3 per genotype). The slopes of a
simple linear regression do not vary significantly across P5 genotypes. **c)** Axon density
decreases physiologically between P21 and P28 in control sciatic nerves but is significantly
reduced to 7-11% of normal levels in mutant nerves at these stages. **d)** Mean axon diameter
per genotype at P28 (n = 2 per genotype) shows that smaller fibers remain on average. **d)**

Binning analysis of *g*-ratios as a function of axon diameter at P28. A total of n = 792 *vs* 746

myelinated axons were analyzed for control *vs* mutant nerves, respectively, from 2 individuals

of each group (correlation significance indicated by asterisks). **f)** *g*-ratios of axonal diameters

at P28, with a significantly different slope (p<0.0001) after simple linear regression, where

mutant *g*-ratios increase most with size. **g**) When binned by diameter ranges (small 0-1 µm,

medium 1-3 µm, and large >3 µm), there is modestly significant hypermyelination in small

axons but more significant hypomyelination at diameters over 1 µm. Statistical significance

was assessed with two-way ANOVA followed by Šidák's post-hoc test; ns, not significant; * *p*

< 0.05; ** *p* < 0.01; *** *p* < 0.001; **** *p* < 0.0001.

Figure 4. Segmental demyelination is preceded by myelin disassembly in Mpz-*Braf* mice.

Teased fibers from sciatic nerves of control (**a, e**) and Mpz-*Braf* mice (**b-d**, **f-g**) at P5 (top

panels) and P21 (bottom panels) were immunostained with antibodies against Mbp and Caspr

and Alexa Fluor-488-conjugated phalloidin. Mpz-*Braf* fibers had degenerating myelin ovoids

(white arrowheads), fewer apparent nodes of Ranvier with diffusing paranodal Caspr (orange

arrowheads), and segmental demyelination, not observed in controls. Nuclear DNA was

stained with DAPI (grey). Maximum intensity z-projection images were generated in **b, e** and **f**

with Nikon NIS-Elements Viewer software. Scale bars: 50 µm (**a, b, f**) or 10 µm (**c, e, g**).

Figure 5. Comparative transcriptomics show downregulation of myelin sheath component transcripts in *Braf*-mutant sciatic nerves.

a) RNA-seq analysis was performed in control and Mpz-*Braf* mutant sciatic nerves (n = 4 per

genotype). Volcano plots depict the DEGs in *Braf*-mutant sciatic nerves relative to controls at

P5 (left panel) and P21 (right panel). An adjusted *p*-value < 0.5 and Log2FC <-1 (blue) or >1

(red) were used as thresholds to define DEGs. The top 15 DEGs are labeled on each graph

according to Manhattan distance. **b)** Volcano plots illustrate the differentially enriched

pathways provided by GSEA through WebGestalt https://www.webgestalt.org/ [131] using the

mouse Wikipathway database. Pathways are significantly enriched when false discovery rate

(FDR) < 0.05 and normalized enrichment score (NES) < -1 (blue) or >1 (red). Circle diameter

is proportional to gene ratio (count of core enrichment genes / count of pathway genes). **c)**

Enrichment plot for "cholesterol biosynthesis", the most significant gene set downregulated in

mutant sciatic nerves (p-value and FDR < 0.001, NES = -2.3). **d)** Dot plot displaying log2 fold-

change (Log2FC) of myelin-associated genes (*Mpz*, *Pmp22*, and *Mbp* in bold, representing

transcripts of the primary myelin sheath proteins).

Figure 6. *Egr2* and its target genes decrease in Mpz-*Braf* nerves with concomitant
increase in expression of negative regulators of myelination

a) Dot plot displaying log2 fold-change (Log2FC) of known positive (*Sox10*, *Egr2*) or negative
(*Jun*, *Sox2*) transcriptional regulators of myelination in RNAseq of mutant *vs* control sciatic
nerves at P5 and P21. **b)** Western blot of Jun and Egr2 in sciatic nerve lysates from P21 and
P42 Mpz-*Braf* and control mice. Gapdh was detected as a loading control. N=13 individuals of
each genotype. Statistical significance was assessed by an unpaired Mann-Whitney test; ns,
not significant; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$. **c)** Enrichment plot for up- or down-
regulated genes (top and bottom panels, respectively) showing that most genes are commonly
deregulated in sciatic nerves from either the *Mpz-Braf* or homozygous *Egr2* hypomorphic allele
mice [132]. **d-e)** Spearman's *rho* to assess significance of correlation was calculated from bulk
RNAseq of Mpz-*Braf* sciatic nerves at P5 (top panels) or P21 (bottom panels), with analogous
datasets at P8 and P30 from Egr2-*Cre*-driven (**d**) overexpression of constitutively active Mek1
(Egr2-*Cre*; *Mek1*DD) or (**e**) knockout of *Shp2* (Egr2-*Cre*; *Shp2*f/f).

Figure 7. Activation of injury-related genes in Mpz-*Braf* sciatic nerves.

a) GSEA analysis of upregulated genes at P21 shows highly significant correlation with the
Wikipathway annotation "Spinal cord injury" (WP2431). **b**) These are also significantly and
positively correlated with genes upregulated in regenerating nerves of young adult mice 7 days
after axotomy. **c**) Transcription of *Shh*, *Gdnf* and *Nrg1* are greatly increased at both P5 and
P21, while Gdnf-family member *Artn* and *Nrg1* receptor ErbB3 are only significantly
upregulated at P21. Teased fibers from sciatic nerves of P21 control and Mpz-*Braf* mice
immunostained against (**d**) S100b (yellow) and Shh (magenta), with DAPI nuclear counterstain
(gray), or (**f**) Pmp2 (orange) and Erbb3 (blue). Scale bars: 50 µm (**d**) or 10 µm (**f**).

Figure 8. Patient-derived immature Schwann cells express inappropriate gene
signatures *in vitro*.

a) Summary of hiPSC-directed differentiation into SC, derived from two unrelated WT (HFF15
shown here and AG08H) and from unrelated adolescent girls, both carrying the *BRAF* p.Q257R
variant (CFC patients RMK0056C, RMK0138C) over time. Representative bright-field images
shown at D13, D19, a day after passage on D32, and at D95-100. Scale bar = 100 µm for all
frames. **b**) Euler diagram showing the number of common/specific dysregulated genes (*p*-
value < 0.05, Log2FC > 1 or < -1) in RMK0056C and RMK0138C mutant cells compared to
control cells (HFF15) at D21 of differentiation. (C) Bubble plot of enrichment analysis of the
common 380 dysregulated genes in RMK0056C and RMK0138C. Gene set names from the
REACTOME database are indicated on the y-axis. Gene sets are ranked according to
significance level (-log adjusted p-value). The size of the dots represents the number of queried

genes present in the corresponding pathway, and the color corresponds to the gene ratio (number of queried genes present in the gene set / gene set size). (D) Enrichment plot for Egr2- and Sox10-mediated initiation of SC myelination using Reactome database, corresponding to the significant gene set downregulated in RMK0056C and RMK0138C cell lines after 30 days of differentiation. (E) Enrichr-KG analysis of the most significantly enriched terms across three additional databases (Descartes Cell Types and Tissue 2021, Jensen DISEASES and CCLE Proteomics 2020) highlight the close lineage relationship between Schwann cell progenitors and either melanocytes ("melanoma" and "skin cancer", the A2058 cell line) or neuroendocrine cells ("visceral neurons in lung/heart", the SKNAS and NCIH446 cell lines).

Figure 1

a Control   Mpz-*Braf*

P21

b **P5** control   Mpz-*Braf*

pErk1
pErk2
—40

Erk1
Erk2
—40

Gapdh
—35

c **P21** control   Mpz-*Braf*

pErk1
pErk2
—40

Erk1
Erk2
—40

Gapdh
—35

d P5   P21

pErk/Erk ratio

Ctrl Mut   Ctrl Mut

e Weight (grams)

Control

Mpz-*Braf*

ns   ns   **** **** ****

P5 P10 P21 P35 P42

Control
Mpz-*Braf*

Mpz-*Braf*
Control

f *Ghr*   *Igf1*

Relative transcription P21

Ctrl Mut   Ctrl Mut

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

a

	d13	d19	d32	>d90
BRAF WT — hFF15				
BRAF Q257R/+ — Patient 56C				
	Patient 138C			

differentiation · expansion · maturation

d0 hiPSC · d6 NCC · d12 committed NCC · SCP · d30 iSC · d95 SC

b

DEG of mutant hiSC re: WT d31

Patient 138C: 3447 | 380 | Patient 56C: 588

Up: *POU3F2, CHGA, CHGB, ALK, KIT*
Down: **EGR2**, *MBP, PRX, LAMA2*

c

380 common DEG genes

- Formation of the cornified envelope
- Type I hemidesmosome assembly
- Cell-cell communication
- Cell junction organization
- Keratinization
- Generation of second messenger molecules
- Extracellular matrix organization
- **EGR2- & SOX10-mediated initiation of SC myelination**
- Signaling by receptor tyrosine kinases
- Collagen formation
- Degradation of the extracellular matrix
- Laminin interactions
- Developmental biology
- Interferon alpha/beta signaling

gene ratio: 0.46, 0.26, 0.06
count: 5.00, 13.42, 36.00

-Log(adj_p_value)

d

'REACTOME_EGR2_AND_SOX10_MEDIATED_
INITIATION_OF_SCHWANN_CELL_MYELINATION'

Enrichment score (ES)

UP in 56C | DOWN in 56C

UP in 138C | DOWN in 138C

Ranking metric

Patient 56C · Patient 138C

e

Melanoma	DISEASES
Skin cancer	DISEASES
Visceral neurons in Lung	Descartes
NCIH446 Lung (Neuroendocrine)	CCLE_Proteomics_2020
SKNAS Autonomic (Neuroblastoma)	CCLE_Proteomics_2020
A2058 Skin (Melanoma)	CCLE_Proteomics_2020
Vascular ECs in Placenta	Descartes
Visceral neurons in Heart	Descartes