Non-classical NMDA receptor subunit GluN3A in a specialized hippocampal region regulates stress-coping strategies in mice

Wei Zhang[1,2,3], Mingyue Zhao[1,2], Jiajie Dai[1,2,4], Linan Zhuo[1,2], Haimou Ye[1,2], Jiesi Wang[1*],

Weiwen Wang[1,2*]

[1]Key Laboratory of Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing, 100101, China

[2]Department of Psychology, University of Chinese Academy of Sciences, Beijing, China

[3]Affiliated Cixi Hospital, Wenzhou Medical University

[4]School of Psychological and Cognitive Sciences, Peking University, Haidian District, Beijing 100871, China

*Correspondence:

Weiwen Wang, Key Laboratory of Mental Health, Institute of Psychology, No. 16 Lincui Road, Chaoyang District,

Beijing, 100101, China.

Email: wangww@psych.ac.cn

Jiesi Wang, Key Laboratory of Mental Health, Institute of Psychology, No. 16 Lincui Road, Chaoyang District,

Beijing, 100101, China.

Email: wangjs@psych.ac.cn

ABSTRACT

The hippocampus is a highly complex structure, defined by profound structural, functional, and molecular heterogeneity. The GluN3A subunit, which forms non-classic N-methyl-D-aspartate receptors (NMDARs), is enriched in the hippocampal CA1 subregion and has a poorly understood functional role, especially in stress-related behavioral regulation. Here, we employed chronic social defeat stress (CSDS) and GluN3A knockout (KO) mouse models combined with behavioral, molecular, genetic, pharmacological, and circuit-level analyses to investigate the role of hippocampal GluN3A in stress coping. We found that CSDS selectively reduced GluN3A expression in the intermediate CA1 (CA1i) region. Decreased GluN3A expression in CA1i, from either stress or genetic deletion, promoted passive coping behaviors, whereas overexpression of GluN3A in CA1i reversed the phenotype under both conditions. Mechanistically, consistent with GluN3A enrichment in pyramidal neurons, c-fos staining and fiber photometry revealed that GluN3A was essential for CA1i pyramidal neuronal activation during coping behaviors. Further chemogenetic manipulations demonstrated that activating CA1i neurons facilitated active coping, while inhibition induced passivity. Moreover, local administration of D-serine, an agonist of GluN3A-containing NMDARs, rapidly shifted coping behavior from passive to active by activating CA1i neurons via GluN3A-dependent mechanisms. Circuit tracing revealed that CA1i neurons project to multiple sub-hippocampal and cortico-limbic regions, and GluN3A overexpression selectively enhanced downstream neuronal activity in CA3 and infralimbic cortex, but not in the nucleus accumbens. Together, these findings identify GluN3A as a key molecule linking hippocampal subregional specialization with adaptive stress coping, and suggest that the rapid, resilience-promoting effects of D-serine are also GluN3A-dependent.

INTRODUCTION

The hippocampus is a structurally and functionally heterogeneous structure that plays essential roles in learning, memory, emotional regulation, and adaptive responses to stress[1-5]. It is broadly divided into multiple subfields such as CA1, CA2/3, and the dentate gyrus (DG)[6]. Beyond these classical anatomical partitions, traditional studies have also revealed prominent structural and functional heterogeneity along the longitudinal axis of the hippocampus[3,4,7]. For example, the dorsal hippocampus is primarily involved in spatial learning and cognition, whereas the ventral hippocampus is more associated with emotional regulation and affective behaviors[3,4,8]. Furthermore, advances in molecular profiling, such as spatial transcriptomics, have revealed an even finer organization, showing that the classical dorsal-ventral dichotomy of subfields CA1, CA3, and DG can be refined into dorsal, intermediate, and ventral portions[4,9]. These finer subregions differ in molecular expression, morphology, and projection patterns[9,10]. However, how these distinct neuronal populations and molecular profiles, particularly within the intermediate subregions, execute specific behavioral functions remains largely unclear.

Notably, gene expression profiling has shown a concentration of GluN3A expression in the intermediate and ventral regions of CA1; this specific expression pattern identifies it as a key molecular marker distinguishing the intermediate from the dorsal CA1 subregion [9]. This subunit forms non-classical N-methyl-D-aspartate receptors (NMDARs) with GluN1/N2, exhibiting unique biophysical and pharmacological properties distinct from those of conventional GluN1/GluN2 NMDARs[11,12]. Unlike canonical GluN2-dependent receptors that require glutamate for activation, GluN1/GluN3A assemblies can be directly activated by sensing ambient extracellular glycine or D-serine, generating excitatory glycine receptor-like currents[13-15]. Beyond being glutamate-insensitive, GluN3A-containing receptors also exhibit low Ca^{2+} permeability, significantly reduced Mg^{2+} blockade, and resistance to classical NMDAR antagonists such as D-AP5 and MK-801[13,16-18]. Consistent with these unique features, GluN3A-containing NMDARs have been shown to regulate synaptic maturation and plasticity in several brain regions, including the hippocampus, and have been implicated in multiple affective behavioral phenotypes[16,17,19-22]. Furthermore, growing evidence from preclinical and clinical studies suggests that exogenous D-serine can exert rapid antidepressant-like effects and modulate emotional and cognitive behaviors, underscoring the GluN3A subunit's potential as a viable target for future clinical translation[23-29]. Collectively, these advances suggest that the subregion-specific expression of GluN3A, combined with the unique properties of the receptors it forms, may represent a key molecular mechanism underlying the hippocampus's broader spatial and functional heterogeneity; however, its specific hippocampal role remains to be identified, especially in stress-related behavioral regulation.

Chronic stress promotes various negative affective states, including passive coping, anhedonia, and anxiety[8,30-32]. Increasing evidence indicates that chronic stress disrupts hippocampal synaptic transmission and intrinsic excitability, contributing to cognitive deficits, emotional dysregulation, and changes in the stress response [2,32-35]. Furthermore, hippocampal subregions, such as the distinct segments of the CA1, play differential roles in mediating these adverse stress-induced outcomes. For example, the dorsal CA1, as traditionally delineated, is primarily linked to deficits of spatial cognition,

working memory, and social function under chronic stress, whereas the ventral CA1 is more closely
associated with affective disturbances and anxiety-related behaviors[36-40]. Although the roles of
canonical GluN2-containing NMDARs in these processes have been extensively studied[41-43], very
little is known regarding the function of non-classical GluN3A-containing NMDARs, which exhibit a
distinct spatial expression pattern within the hippocampus, under chronic stress conditions. Recent
study has shown that GluN3A deletion in the traditional ventral CA1 region, which encompasses both
the CA1 intermediate (CA1i) and CA1 ventral (CA1v) areas, induces anxiety-like behavior and that
these excitatory glycine receptors (eGlyR) contribute to corticosterone-dependent modulation of
neuronal plasticity, including long-term potentiation[15]. These findings hint that GluN3A may influence
stress-related behavioral outcomes and contribute to the functional heterogeneity of CA1. However,
GluN3A serving as a molecular marker distinguishing the CA1 dorsal (CA1d) and CA1i subregions[9],
it remains unclear which specific hippocampal CA1 loci and behavioral domains GluN3A controls,
and how it functionally contributes to stress regulation, particularly under chronic stress exposure.

To address these critical questions, we employed chronic social defeat stress (CSDS) and GluN3A
knockout (KO) mouse models, combining comprehensive approaches encompassing behavior,
molecular biology, genetic manipulation, microscopic imaging, fiber photometry, pharmacology, and
single-cell transcriptomic analysis. This integrative strategy allowed us to delineate the specific
contribution of hippocampal GluN3A to passive coping behavior, clarify its role relative to anhedonia,
social interaction and anxiety, and identify the neural mechanisms and specific loci underlying this
regulation. Our findings provide mechanistic insight into how non-classical NMDARs contribute to
stress adaptation, revealing a previously unrecognized role of hippocampal GluN3A in shaping coping
strategies under adverse conditions.

RESULTS

Effects of CSDS and GluN3A KO on stress-related behaviors and GluN3A expression in the
hippocampus
We systematically evaluated various stress-related behaviors induced by CSDS in mice (Fig. 1a, b).
Regarding the anhedonia indicator, CSDS mice showed a significant reduction in social interaction
ratio (t_{16} = 5.802, $^{***}P < 0.001$, Fig. 1c, d) in the social interaction test (SIT) but no significant change in
sucrose preference ratio ($P > 0.05$, Supplementary Fig. 1a) in the sucrose preference test (SPT)
compared to the control group. In terms of coping behaviors in response to inescapable stress, CSDS
significantly increased the immobile time of mice in the tail suspension test (TST) (t_{14} = 2.877, $^{*}P$ =
0.012, Fig. 1e) and forced swim test (FST) (t_{11} = 2.240, $^{*}P$ = 0.047, Fig. 1f). In locomotor activity,
compared to the control group, CSDS did not affect the total distance traveled by mice ($P > 0.05$, Fig.
1g, h). Regarding anxiety-related indicators, CSDS markedly induced anxiety-like behavior in mice,
evidenced by a decrease in the number of entries of the central area (t_{13} = 2.801, P = 0.015, Fig. 1i)
and the time spent in the central area (Welch-corrected $t_{9.157}$ = 3.290, $^{*}P$ = 0.009) in the open field test
(OFT) (Fig. 1g, j). Additionally, in the elevated plus maze (EPM), CSDS mice exhibited a reduction in
the number of open arm entries (Welch-corrected $t_{11.43}$ = 2.232, $^{*}P$ = 0.047, Fig. 1k, l) and the open
arm time ratio (Welch-corrected $t_{8.989}$ = 2.206, P = 0.055, Fig. 1k, m). These results indicated that
CSDS successfully induced various negative affective states in mice.

Subsequently, we examined the changes in GluN3A expression levels in various emotion-related
brain regions, particularly the hippocampus, in mice following CSDS (Supplementary Fig. 1b, c). The
quantitative real-time PCR (qRT-PCR) results showed a significant reduction in GluN3A expression in
the CA1i region of the hippocampus in CSDS mice compared to the control group (t_{23} = 2.692, $^*P$ =
0.013, Fig. 1n), while there were no significant changes in the other subregions of the hippocampus,
including CA1d, CA1v, CA3, and DG ($P > 0.05$, Fig. 1n). The western blot results similarly
demonstrated a significant decrease of GluN3A protein expression in the CA1i in CSDS mice (t_9 =
2.866, $^*P$ = 0.024), with no significant changes observed in CA1v and CA3 ($P > 0.05$, Fig. 1o, p).
These data indicated that CSDS downregulated GluN3A expression in the hippocampal CA1i region
in mice.

To further examine the relationship between GluN3A and emotion-related behaviors, we conducted
behavioral tests on GluN3A KO mice under physiological conditions (Fig. 1q). On the anhedonia
indicators, GluN3A KO mice did not exhibit an abnormal ratio of social interactions ($P > 0.05$, Fig. 1r)
and sucrose preference ($P > 0.05$, Supplementary Fig. 2a) compared to wild-type (WT) mice. In
contrast, the results showed that GluN3A KO mice exhibited a passive coping phenotype compared to
WT mice, as evidenced by significantly increased immobile time in TST (t_{16} = 2.778, $^*P$ = 0.013, Fig.
1s) and FST (t_{14} = 2.457, $^*P$ = 0.028, Fig. 1t). In addition, GluN3A KO did not influence the locomotor
activity ($P > 0.05$, Fig. 1u) and anxiety-like behavior ($P > 0.05$, Fig. 1v-y).

To ascertain whether GluN3A KO is associated with an increased susceptibility to social defeat stress
(SDS), a systematic assessment was conducted to evaluate the effects of subthreshold SDS on
emotion-related behaviors in GluN3A KO mice. The results showed GluN3A KO mice subjected to
subthreshold SDS exhibited a similar significant increase in immobile time in the FST (Welch-
corrected $t_{2.520}$ = 9.677, $^*P$ = 0.031, Supplementary Fig. 2d). Still, no substantial changes observed in
social interest, sucrose preference, locomotor activity, and anxiety-like behavior ($P > 0.05$,
Supplementary Fig. 2b, c, e, g-i), except a significant reduction in the number of entries of central
area in the OFT (t_{13} = 2.815, $^*P$ = 0.015, Supplementary Fig. 2f). The data indicated that the GluN3A
KO does not influence the susceptibility of mice to SDS.

Taken together, these results suggested that the decreased expression of GluN3A in the brain was
involved in the manifestation of passive coping behaviors, and that the hippocampal CA1i may be a
potential brain region for the regulation of stress-coping behaviors by GluN3A.

Overexpression of the GluN3A decline in CA1i neurons reversed passive coping behaviors
We further investigated the regulatory role of CA1i GluN3A in emotion-related behaviors by using
Adeno-associated virus (AAV) to overexpress recombinant GluN3A in the CA1i of CSDS mice (Fig. 2a,
b). We found that overexpression of exogenous GluN3A did not significantly affect anhedonia
behaviors in CSDS mice (SIT, Fig. 2c; SPT, Supplementary Fig. 3a). In contrast, CA1i-targeted
GluN3A overexpression markedly reversed the passive coping phenotype induced by CSDS.
Specifically, two-way ANOVA for TST showed a significant interaction effect ($F_{(1, 28)}$ = 5.568, $^*P$ =
0.026, Fig. 2d), with post-hoc analysis revealing a significant increase in immobile time in the stress:
AAV-ZsGreen group compared to the control: AAV-ZsGreen group (False discovery rate-corrected $^*P$
= 0.024), while the stress: AAV-Grin3A group showed a significant decrease in immobile time
compared to the stress: AAV-ZsGreen group (FDR-corrected $^*P$ = 0.016). Two-way ANOVA for FST

showed significant main effects of stress ($F_{(1, 28)}$ = 5.010, $^*P$ = 0.033, Fig. 2e) and viral treatment (i.e.,
AAV-ZsGreen or AAV-Grin3A) ($F_{(1, 28)}$ = 12.99, $^{**}P$ = 0.0012) on the immobile time, with post-hoc
analysis showing a significant increase in immobile time in the stress: AAV-ZsGreen group compared
to the control: AAV-ZsGreen group (FDR-corrected $^*P$ = 0.026) and a significant decrease in immobile
time in both the control: AAV-Grin3A group (FDR-corrected $^*P$ = 0.036) and stress: AAV-Grin3A group
(FDR-corrected $^*P$ = 0.016) compared to their respective AAV-ZsGreen groups. Furthermore,
consistent with the previous experiments, CSDS induced the changes in locomotor indicators and
multiple anxiety indicators in OFT and EPM (Fig. 2f-j), but these were not affected by the
overexpression of the exogenous GluN3A subunit. These results indicated that the CA1i GluN3A
modulated CSDS-induced passive coping behaviors in mice.

To further confirm the causal relationship between CA1i GluN3A and stress-coping behaviors, we
overexpress recombinant GluN3A in the CA1i region of GluN3A KO mice using AAV (Fig. 2k).
Consistent with the findings above, GluN3A overexpression normalized passive coping behaviors
without affecting social interaction, locomotor activity, or anxiety-like behavior ($P > 0.05$, Fig. 2l-n and
Supplementary Fig. 3b-d). Specifically, GluN3A KO mice showed a significant increase in immobile
time in TST (t_{12} = 2.212, $^*P$ = 0.047, Fig. 2l) and FST (t_{11} = 2.309, $^*P$ = 0.041, Fig. 2m) compared to
WT mice, while GluN3A KO: AAV-Grin3A mice exhibited a significant restoration of immobile time to
normal levels in both TST (t_{10} = 2.696, $^*P$ = 0.023) and FST (t_{10} = 2.912, $^*P$ = 0.016) compared to
GluN3A KO mice. In summary, hippocampal CA1i GluN3A played a crucial role in regulating stress-
induced coping strategies.

**185 Activity of CA1i pyramidal neurons mediated by GluN3A-dependent regulation of stress-
coping behaviors**
Having established that GluN3A in the hippocampal CA1i region regulates passive coping behavior in
CSDS mice, we next sought to identify the neuronal types that might contribute to this effect. Through
reanalysis of single-nuclei RNAseq (snRNAseq) data from previously published studies[44], we found
that GluN3A is abundantly expressed in excitatory neurons in the hippocampal CA1 region, and its
expression pattern changes after repeated social defeat (RSD) (Fig. 3a, b). Moreover, compared to
the Control: WT group, GluN3A expression significantly decreased in excitatory neurons of WT
subjected to RSD ($^*P$ = 0.037, Fig. 3c, d).

Therefore, we further investigated the functional activity of CA1i pyramidal neurons during stress-
coping behaviors in the CSDS model under conditions of GluN3A modulation. The results of c-Fos
immunostaining showed reduced neuron activity of the pyramidal layer in the CA1i region after the
FST in stressed mice compared to controls (Fig. 3e-g), which was restored by overexpressing
recombinant GluN3A. Specifically, two-way ANOVA analysis of the number of c-Fos-positive cells
showed significant interaction ($F_{(1, 20)}$ = 4.820, $^*P$ = 0.040, Fig. 3f) and stress main effects ($F_{(1, 20)}$ =
6.015, $^*P$ = 0.024), with a marginal significant effect for viral treatment ($F_{(1, 20)}$ = 3.679, P = 0.070).
Post-hoc analysis revealed a significant decrease in the number of c-Fos-positive cells in the stress:
AAV-ZsGreen group compared to the control: AAV-ZsGreen group (FDR-corrected $^{**}P$ = 0.009), while
the stress: AAV-Grin3A group showed a significant increase in the number of c-Fos-positive cells
compared to the stress: AAV-ZsGreen group (FDR-corrected $^{**}P$ = 0.009).

Furthermore, we examined the direct relationship between the GluN3A subunit and the functional

activity of CA1i pyramidal neurons after the FST in GluN3A KO mice. Immunostaining results showed
that the number of c-Fos-positive cells was significantly increased after FST compared with the basal
condition (no FST experience) in WT mice (t_9 = 2.884, $^*P$ = 0.018), and such activation did not occur
in FST-experienced GluN3A KO mice (t_{10} = 3.232, $^{**}P$ = 0.009, Fig. 3h, i), indicating that the activation
of pyramidal neurons in the CA1i region was involved in FST behavior, while the absence of GluN3A
resulted in this kind of functional defect.

Moreover, we validated the necessity of GluN3A in the activation of excitatory neurons in the CA1i
during the TST using fiber photometry (Fig. 3j). During TST, the excitatory-specific GCaMP signal,
which characterizes calcium levels, showed a significant correlation with the active state of the mice.
The rise of signal was behind the transition from immobile state to struggling state (Fig. 3k).
Subsequently, we observed that the average peak amplitude of neuronal responses significantly
increased during events when mice transitioned from immobility to struggling (Fig. 3l, m), with GluN3A
KO mice showing significantly lower struggle related neuronal activation compared to WT mice (t_9 =
3.398, $^{**}P$ = 0.008, Fig. 3n).

Collectively, these results indicated that the activity of CA1i pyramidal neurons mediated the GluN3A-
dependent regulation of stress-coping behavior in mice.

Bidirectional chemogenetic modulation of CA1i pyramidal neurons directly mimicked or
reversed the stress-induced passive coping behaviors
To directly test the causal role of CA1i pyramidal neuron activity in stress-coping behavior, we
employed bidirectional chemogenetic manipulation. We first increased the activity of CA1i neurons
using AAV-hM3D(Gq) in combination with different concentrations of clozapine-N-oxide (CNO) (Fig.
4a). The results showed that saline, 0.05 mg/kg, 0.2 mg/kg, and 0.5 mg/kg CNO did not affect the
locomotor activity and anxiety-like behavior of normal mice in the OFT and rotarod test (RT) ($P>0.05$,
Supplementary Fig. 4a-d). However, one-way ANOVA showed that concentrations of 0.2 mg/kg (FDR-
corrected $^{**}P$ = 0.009, Fig. 4b-c, Supplementary Fig. 4e) and 0.5 mg/kg (FDR-corrected $^{**}P$ = 0.009)
CNO effectively increased the number of c-Fos-positive cells at the injection site ($F_{(2, 3)}$ = 16.89, $^*P$ =
0.023). Therefore, we chose a concentration of 0.2 mg/kg CNO for subsequent chemogenetic
activation experiments (Fig. 4d). The results indicated that the activation of CA1i pyramidal neuronal
activity in CSDS mice partially enhanced social interest in the SIT (Fig. 4e), and reversed passive
coping behavior induced by CSDS in the TST and FST (Fig. 4f, g), but did not alter their locomotor
activity and anxiety-like behavior in the OFT (Fig. 4h-j). Specifically, two-way ANOVA for the social
interaction ratio showed a significant viral treatment main effect ($F_{(1, 20)}$ = 6.983, $^*P$ = 0.016, Fig. 4e).
Two-way ANOVA for immobile time in the TST showed a significant interaction effect ($F_{(1, 22)}$ = 6.923,
$^*P$ = 0.015, Fig. 4f), with post-hoc analysis showing a significant increase in immobile time in the
stress: AAV-mCherry group compared to the control: AAV-mCherry group (FDR-corrected $^*P$ = 0.011),
while the stress: AAV-hM3D(Gq) group exhibited a significant decrease in immobile time to normal
levels compared to the stress: AAV-mCherry group (FDR-corrected $^*P$ = 0.011). Two-way ANOVA for
immobile time in the FST showed significant stress ($F_{(1, 21)}$ = 5.350, $^*P$ = 0.031, Fig. 4g) and viral
treatment main effects ($F_{(1, 21)}$ = 12.34, $^{**}P$ = 0.002), with post-hoc analysis showing a significant
increase in immobile time in the stress: AAV-mCherry group compared to the control: AAV-mCherry
group (FDR-corrected $^*P$ = 0.023), and a significant decrease in immobile time in both the control:
AAV-hM3D(Gq) group (FDR-corrected $^*P$ = 0.026) and stress: AAV-hM3D(Gq) group (FDR-corrected

$^*P$ = 0.009) compared to their respective AAV-mCherry groups. These results indicated that increased

excitatory activity in CA1i neurons can alleviate CSDS-induced passive coping behavior.

On the other hand, we inhibited the activity of CA1i pyramidal neurons in normal mice using AAV-

hM4D(Gi) (Fig. 4k, l). We found that reducing the activity of the neurons significantly increased

immobile time in the TST (t_{12} = 3.261, $^{**}P$ = 0.007, Fig. 4m) and FST (t_{13} = 2.622, $^*P$ = 0.021, Fig. 4n),

but did not induce abnormalities in locomotor activity and anxiety-like behavior in the OFT ($P>0.05$,

Fig. 4o-q), indicating that the inhibition of neuronal activity in CA1i can directly induce passive coping

phenotype.

Overall, our findings suggest that the activity of CA1i pyramidal neurons may underlie GluN3A's

regulation of stress-coping behavior.

Potentiation of GluN3A-mediated CA1i pyramidal neuron activity was crucial for the active

coping strategies induced by D-serine

We further explored whether GluN3A participates in the rapid D-serine–induced shift of stress-coping

strategies within the hippocampus. By injecting D-serine into the bilateral CA1i region of CSDS mice

and GluN3A KO mice before behavioral testing (Fig. 5a, b), we found that D-serine could significantly

reverse the passive coping behavior induced by CSDS (Fig. 5c, d), while it had no significant effect on

locomotor activity and anxiety-like behavior in the OFT in all groups ($P>0.05$, Fig. 5e-g). Specifically,

two-way ANOVA for immobile time in the TST and FST showed significant interaction (Control-

cerebrospinal fluid (CSF), Control-D-serine, Stress-CSF and Stress-D-serine group, TST: $F_{(1, 27)}$ =

6.084, $^*P$ = 0.020; FST: $F_{(1, 27)}$ = 5.106, $^*P$ = 0.032, Fig. 5c, d) and drug treatment main effects (TST: F

$_{(1, 27)}$ = 11.60, $^{**}P$ = 0.002; FST: $F_{(1, 27)}$ = 11.19, $^{**}P$ = 0.002), with post-hoc analysis showing a

significant increase in immobile time in the stress: CSF group compared to the control: CSF group in

both tests (TST: FDR-corrected $^*P$ = 0.017; FST: FDR-corrected $^*P$ = 0.032), while the stress: D-

serine group exhibited a significant decrease in immobile time compared to the stress: CSF group

(TST: FDR-corrected $^{***}P<0.001$; FST: FDR-corrected $^{**}P$ = 0.002). Interestingly, compared to the

control (WT):D-serine group, the GluN3A KO:D-serine group still exhibited significantly increased

immobile time in both the TST (t_{13} = 2.470, ^+P = 0.028) and FST (t_{13} = 3.225, ^{++}P = 0.007), indicating

that GluN3A is required for D-serine–induced behavioral improvement. Additionally, simultaneously

inhibiting the activity of CA1i neurons in CSDS mice while administering D-serine, we found that the

stress-hM4D(Gi) group showed a significant increase in immobile time in both the TST (t_{12} = 4.628,

$^{###}P<0.001$) and FST (t_{12} = 2.671, $^#P$ = 0.020) compared to the Stress-mCherry group, suggesting

that suppressing CA1i pyramidal neuronal activity blocked the active coping effect of D-serine.

Furthermore, we performed immunostaining for c-Fos in the CA1i pyramidal layer neurons of mice

from the groups above after the FST (Fig. 5h). The results showed that D-serine significantly reversed

the decrease in the number of c-Fos-positive cells induced by CSDS in the CA1i pyramidal layer (Fig.

5i, j). Specifically, two-way ANOVA for the number of c-Fos-positive cells showed significant stress

(Control-CSF, Control-D-serine, Stress-CSF, and Stress-D-serine group, $F_{(1, 16)}$ =11.23, $^{**}P$ = 0.004,

Fig. 5j) and drug treatment effects ($F_{(1, 16)}$ =13.78, $^{**}P$ = 0.002). Post-hoc analysis showed a significant

decrease in the number of c-Fos-positive cells in the stress: CSF group compared to the control: CSF

group (FDR-corrected $^*P$ = 0.037), while both the control: D-serine group (FDR-corrected $^*P$ = 0.016)

and stress: D-serine group (FDR-corrected $^*P$ = 0.045) showed a significant increase in the number of

c-Fos-positive cells compared to their respective CSF groups. However, in GluN3A KO:D-serine mice, the number of c-Fos-positive cells in the CA1i pyramidal layer did not rise to the level of the control (WT):D-serine group (t_8 = 2.967, ^+P = 0.018). In summary, D-serine facilitates a rapid shift from passive to active coping behavior by potentiating GluN3A-dependent CA1i pyramidal neuron activity, highlighting GluN3A signaling in this hippocampal subregion as a crucial modulator of adaptive stress responses.

Projections of CA1i neurons and downstream circuits may be involved in GluN3A-mediated regulation of stress-coping behaviors

To identify the downstream targets of hippocampal CA1i neurons potentially involved in GluN3A-mediated behavioral regulation, we first performed projection-mapping analyses using the Allen Brain Atlas database and AAV-mediated tracing experiments. According to the Allen Brain connectivity data, CA1i neurons send prominent projections to several sub-hippocampal and limbic related brain regions, including the CA3, nucleus accumbens (NAc), infralimbic cortex (IL), BLA, and lateral entorhinal cortex (LEC) (Fig. 6a). Consistent with these database results, injection of AAV2/9-syn-ZsGreen into the CA1i region resulted in robust fluorescent projection signals in these downstream areas, particularly in the CA3, NAc core (AcbC) and shell (AcbSh), IL, BLA, and LEC (Fig. 6b), confirming that CA1i pyramidal neurons form extensive efferent connections with emotion-related circuits.

Next, we examined whether overexpression of the GluN3A subunit in CA1i neurons could influence neuronal activation in these projection targets following chronic stress. c-Fos immunostaining was performed in representative downstream regions after the FST in control and CSDS mice receiving either AAV-ZsGreen or AAV-Grin3A injections in the CA1i region (Fig. 6c-h, Supplementary Fig. 5a-d). In the CA3 region, a two-way ANOVA revealed a significant stress × viral treatment interaction ($F_{(1, 21)}$ = 5.641, $^*P$ = 0.027, Fig. 6d) and a significant stress main effect ($F_{(1, 21)}$ = 5.816, $^*P$ = 0.025). Post hoc analysis showed that in the AAV-ZsGreen group, CSDS significantly decreased the number of c-Fos–positive neurons in the CA3 region compared with controls (FDR-corrected $^{**}P$ = 0.009). In contrast, overexpression of GluN3A in CA1i neurons reversed this reduction, restoring c-Fos expression to control levels (FDR-corrected $^*P$ = 0.017). In the IL, both stress ($F_{(1, 15)}$ = 5.368, $^*P$ = 0.035, Fig. 6g) and viral treatment ($F_{(1, 15)}$ = 5.182, $^*P$ = 0.038) exerted significant main effects. Post hoc analysis indicated that in the AAV-ZsGreen group, CSDS significantly reduced c-Fos–positive cell counts in the IL (FDR-corrected $^*P$ = 0.049), whereas GluN3A overexpression markedly increased c-Fos expression levels relative to the stress: AAV-ZsGreen group (FDR-corrected $^*P$ = 0.049). In contrast, in both the AcbSh and AcbC regions, there were no significant effects observed ($P > 0.05$, Supplementary Fig. 5b, c). These findings suggest that GluN3A modulation in the CA1i primarily affects activity in specific CA3 and IL targets, rather than in the NAC.

Together, these results indicate that CA1i pyramidal neurons establish strong anatomical and functional connections with emotion-related structures, and that GluN3A-dependent regulation of stress-coping behaviors may occur through selective modulation of CA1i-CA3 or CA1i-IL circuit activity rather than global changes across all projection targets.

DISCUSSION

In the present study, we systematically investigated a previously unrecognized role of the GluN3A-

containing non-classical NMDARs within the hippocampal CA1i region in regulating stress-coping
strategies. Our results demonstrated that chronic stress induced a specific decrease in GluN3A
expression in the hippocampal CA1i region, and that CA1i GluN3A explicitly regulates the passive
coping behaviors in the TST and FST. Mechanistically, the effects of GluN3A on stress-coping may be
mediated through the activation of CA1i pyramidal neurons. Moreover, we found that D-serine
administration facilitated a rapid shift from passive to active coping by activating GluN3A-dependent
CA1i neurons. Finally, circuit-level analyses revealed that the selective, GluN3A-mediated activation
of CA1i-CA3 and CA1i-IL pathways, but not CA1i-NAc, may serve as a potential circuit substrate for
stress adaptation. Collectively, these findings highlight a critical role for GluN3A in response to chronic
stress, and its ability to shape adaptive coping through hippocampal CA1i pyramidal neurons and
specific projection circuits, and reveal its contribution to the rapid shift in coping strategy induced by
D-serine.

First, by focusing on the non-classical NMDAR subunit GluN3A and its distinctive expression pattern
in refined hippocampal subregions, we found that mice exhibited a selective reduction in GluN3A
expression in the CA1i region, but not in the traditional, more broadly dorsal and ventral subareas,
after CSDS. This finding provides functional significance for the Hippocampal Gene Expression Atlas
(HGEA) constructed by Bienkowski et al., which divides the CA1 into four subregions: CA1d, CA1i,
CA1v, and the ventral tip (CA1vv), and identifies GluN3A as a key molecular marker distinguishing
CA1i from CA1d[9]. Moreover, our causal manipulations (both genetic knockout and selective CA1i
overexpression in CSDS or KO animals) confirmed that CA1i GluN3A bidirectionally regulates passive
coping behaviors in the TST and FST, without affecting social interest or anxiety-like behaviors. This
demonstrates a regionally and behaviorally specific regulatory role for CA1i GluN3A-expressing
neurons in stress coping. This behavioral specificity for coping refines previous reports. For instance,
while one study linked global hippocampal GluN3A influences to both helplessness and anxiety[45],
another suggested that GluN3A in the traditional ventral hippocampus modulates anxiety-like
behaviors and is essential for neurons' response to corticosterone[15]. Our results reveal a more
precise function, identifying GluN3A as a key molecular determinant of stress response and coping
flexibility, and pinpointing CA1i as a critical functional locus linking molecular heterogeneity to
behavioral specificity during stress adaptation.

Coping behavior represents a core dimension of the stress response, reflecting the individual's ability
to adjust strategies dynamically when facing adversity[46,47]. Active coping is associated with
resilience and positive affect, whereas passive coping (helplessness) is linked to depression
vulnerability[48-51]. Clinically, impairments in coping flexibility are a hallmark of major depressive
disorder and other stress-related psychopathologies[52-54]. In human studies, genetic variations in
the GRIN3A locus have been implicated in bipolar disorder and schizophrenia[55,56], lending clinical
relevance to our findings and further suggesting that the CA1i GluN3A-mediated pathway we
identified may contribute to the pathophysiology of related mental illnesses. Thus, identifying CA1i
GluN3A as a regulator of coping strategies may provide novel therapeutic entry points for disorders
characterized by maladaptive stress responses.

Our study further elucidates the neuronal mechanism by which GluN3A modulates stress-coping
strategies, suggesting that modulation of CA1i neuronal excitability, primarily in pyramidal neurons, is
the key mediator. Several convergent lines of evidence support this conclusion. First, snRNAseq data
revealed a significant downregulation of GluN3A expression in CA1 excitatory neurons of CSDS-
exposed mice[44], a finding consistent with the identified neuronal hypoactivity observed in CSDS and
GluN3A KO mice in the current study. Second, both stress exposure and GluN3A KO decreased CA1i
c-Fos expression after coping behaviors, and KO animals further showed attenuated struggling-
related calcium transients in pyramidal neurons during TST. Third, overexpression of CA1i GluN3A
restored c-fos-represented neuronal activity and rescued passive coping behavior. Critically,
Chemogenetic manipulation further established a causal link between CA1i neuron excitability and
coping behavior: activation promoted active coping, while inhibition induced passive coping. Although
we cannot exclude the potential contribution of GluN3A-expressing inhibitory neurons, our results
suggest that pyramidal neurons are the main effector population translating GluN3A signaling into
behavioral outcomes.

One plausible mechanism underlying this regulatory role in CA1i neuron activity may involve the
unique properties of GluN3A-containing NMDARs, which are inherently glutamate-insensitive but
responsive to extracellular glycine or D-serine, enabling CA1i pyramidal neurons to sense glial-
derived glycine or D-serine released under stress conditions [13,15,16]. Chronic stress-induced
downregulation of GluN3A may disrupt this alternative excitatory pathway, thereby dampening
pyramidal neuron excitability and biasing behavioral output toward passive coping. Recent
electrophysiological evidence supports the core premise of this hypothesis, which indicates that
excitatory glycine receptors are present on ventral hippocampal pyramidal neurons and mediate
distinct synaptic dynamics [15]. Furthermore, the proposed link between GluN3A and neuronal
excitability is further strengthened by findings in other brain regions, such as the olfactory bulb, where
GluN3A KO also reduces c-Fos expression and CaMKII phosphorylation[57]. Finally, the causal link
between CA1 pyramidal neuron hypoactivity and passive coping is consistent with previous studies,
as inhibition of these neurons has been shown to promote depression-like behaviors, including stress-
coping behaviors represented by the FST and TST[34,35]. This convergence of evidence strengthens
the hypothesis that this non-classical GluN3A-containing NMDAR pathway modulates CA1i pyramidal
neuron excitability.

Our findings on D-serine further underscore the translational significance of GluN3A signaling. As a
high-affinity ligand for the glycine-binding site of GluN3 subunits[13,58], D-serine and its derivatives
have been reported to exert rapid antidepressant effects in both preclinical and clinical studies, with
the hippocampus being a critical site of action[23,28,59]. Given that passive coping is a behavioral
endophenotype targeted by antidepressant treatment[60-62], we explored the mechanistic link
between D-serine, GluN3A, and stress-coping behaviors. We found that D-serine microinjection into
the CA1i rapidly shifted coping behavior from passive to active, accompanied by increased CA1i
pyramidal neuron activity. Remarkably, this effect was abolished in GluN3A KO mice or when CA1i
neuron activity was suppressed, indicating that GluN3A-dependent excitation of CA1i pyramidal
neurons is essential for D-serine's behavioral efficacy. Given that D-serine itself has also been
reported to have extensive links, including endogenous changes and exogenous administration, to

symptom or cognitive improvements in clinical and pre-clinical settings[63-66], our findings suggest
that the GluN3A-dependent pathway in CA1i is not only critical for adaptive coping but may also offer
a promising and mechanistically targetable path for the development of therapeutic interventions or
cognitive enhancements in related conditions.

Finally, our initial exploration of CA1i projection analysis points toward potential circuit-level
mechanisms to explain the significant role of GluN3A. We demonstrated that CA1i neurons project
extensively to sub-hippocampal and limbic-related regions, including CA3, IL, and NAc. Furthermore,
our results showed that GluN3A overexpression selectively enhanced downstream neuronal activation
in CA3 and IL, but not NAc. These projection circuits, although not specific to CA1i, have previously
been reported to be associated with stress-coping strategies of mice [30,67,68]. This alignment with
previous reports strongly suggests that CA1i-CA3 and CA1i-IL projections may be the functional
circuits underlying GluN3A-dependent regulation of stress coping by CA1i neurons.

In conclusion, our study identifies non-classical GluN1/GluN3A NMDARs in the hippocampal CA1i as
key regulators of adaptive stress responses by selectively activating pyramidal neurons and defined
projection circuits. While further studies are warranted to determine the exact roles of GluN3A-
expressing pyramidal neurons and their downstream circuit targets, our findings bridge hippocampal
structural and transcriptional heterogeneity with behavioral specialization, offering new insight into the
neural basis of coping and resilience. Furthermore, the discovery that D-serine exerts GluN3A-
dependent rapid modulation of coping behavior highlights a translational opportunity for targeting non-
classical NMDARs in clinical usage, such as the treatment of depression and the improvement of
stress-related symptoms.

MATERIALS AND METHODS

Animals

All animal procedures were approved by the Institutional Animal Care and Use Committee of the
Institute of Psychology, Chinese Academy of Sciences. C57BL/6N male mice and CD-1 (ICR) male
mice were purchased from Charles River, Beijing, China. *Grin3A*$^{-/-}$ (GluN3A KO) mice (C57BL/6J
background) were provided by Tifei Yuan's laboratory. This model was established by the construction
of a specific small guide RNA (sgRNA) vector to direct Cas9 to delete exon 2 of the GluN3A gene.
The GluN3A KO and WT littermates used in this study were bred from the *Grin3A*$^{+/-}$ mice in the
individually ventilated cage (IVC). All mice were housed in a specific-pathogen-free (SPF)
environment with a temperature of 22±2℃, relative humidity of 40-70%, and a 12-hour (h) light-dark
cycle (7:00 am-19:00 pm). The experimental mice had free access to food and water throughout the
experimental period.

CSDS model
CSDS was performed as previously described[69,70]. Male CD1 mice, singly housed for 10-24 weeks,
were screened based on aggression intensity. CD1 mice displaying attacks toward unfamiliar
"intruders" within 3 minutes (min) for 3 consecutive days were designated as "residents" for

subsequent stress experiments. 3 days before the CSDS experiment, "resident" CD1 mice were
placed into the CSDS cage to establish territorial awareness. During each day of the 10-day CSDS
period, adult C57BL/6N mice were introduced into the "resident" CD1 mouse home cage and
subjected to physical attacks for 10 min. For the remainder of the day, a transparent perforated PVC
divider was used to separate them while maintaining visual, auditory, and olfactory contact. For 10
successive days, a C57BL/6N mouse was introduced to a novel aggressive CD1 mouse home cage
every day. C57BL/6N control mice were housed in the CSDS cage with a divider to avoid body
contact, and a new cage mate was introduced every 24 hours (h) for 10 days. After the social defeat
stress, the experimental mice were individually housed until all tests were completed.
468
469 **Subthreshold SDS**
470 Experiments were performed with GluN3A KO mice involving three consecutive 5 min physical
471 interactions with a novel CD-1 mice, with a 15 min interval between each interaction[71].
472
Behavioral assessments
This study assessed the anhedonia symptom using the SPT (sensory anhedonia) and the three-
chamber SIT (social anhedonia). Locomotor activity was evaluated through the OFT and RT, while
coping behavior was measured using the TST and FST. Anxiety symptoms were examined through
the OFT and EPM.
All behavioral tests were conducted during the light cycle, with at least a 24h interval between
different tests. 30min before each behavioral test, experimental mice were placed in a dimly lit testing
room to acclimate to the environment. After each mouse completed the behavioral test, the apparatus
was cleaned with 70% alcohol to eliminate residual mouse odors and traces. All test procedures were
video-recorded.
483  **SPT.** During adaptation phase, mice were acclimatized for 48h by simultaneously giving one
bottle of 1% sucrose solution and one bottle of pure water, and the bottle positions were switched
after 24h of acclimatization in order to eliminate the position preference effect. During testing, the
mice were deprived of water and food for 12h, followed by the placement of 1% sucrose solution and
pure water for another 12h, and the amount of the two solutions consumed in testing phase was
measured. The sucrose preference index was calculated as follows: Sucrose preference ratio (%) =
Sucrose solution intake (g) / [Sucrose solution intake (g) + Pure water intake (g)] × 100%.
490  **OFT.** Mice were placed in the center of an open field arena (40×40×40 cm) to freely explore for
491 10min. The central square area (24×24 cm) was designated as the center zone. The total distance
492 traveled in the open field, entries into the center zone, and time spent in the center zone were
493 automatically tracked using Anymaze software.
494  **EPM.** The EPM apparatus consists of a cross 50cm above the ground (single side arm 35cm
495 long, 5cm wide, closed arm 15cm high). Mice were placed in the central area with their heads facing
496 the open arm and allowed to freely explore for 5 min. The number of entries into the open arms and
497 the time spent in the open arms were automatically tracked using Anymaze software. The percentage
498 of time spent in the open arms (Open Arm Time Ratio) was calculated as follows: time spent in open
499 arms (s) / [time spent in open arms (s) + time spent in closed arms (s)] × 100%.
500  **SIT.** The three-chamber test apparatus is an area (40 cm wide, 60 cm long, 20 cm high) divided
into three equal areas by two transparent dividers with a shuttle channel at the bottom of the centre. A
metal mesh cage is placed in the area on each side of the distal end. The testing process comprised

two 10 min phases. During Phase 1 (adaptation phase), the test mouse is placed in the central area
with the shuttle passage closed, allowing it to freely explore and familiarize itself with the environment.
In Phase 2 (social test phase), an unfamiliar conspecific social mouse is placed in a metal cage on a
random side area. The test mouse is then placed in the central area with the shuttle passage open,
allowing free exploration of the three areas. Anymaze software automatically tracked the time the
mouse spent in each area. The social interaction ratio was calculated as follows: Social interaction
ratio (%) = Time spent in the social area (s) / [Time spent in the social area (s) + Time spent in the
empty cage area (s)] × 100%.

TST. During testing, tape is attached to the tail tip of the experimental mouse, 1 cm from the end.
A plastic tube is placed over the mouse's tail to prevent climbing. The mouse's tail is then secured to
the top of the tail suspension apparatus, suspending the mouse approximately 25 cm above the
ground for 6 min. The duration of immobility (s) during the 6 min video recording was recorded and
analyzed.

FST. The experimental mouse is placed in a transparent cylindrical container (10 cm in diameter,
30 cm high) with water (25 cm deep, 25±1°C) to swim for 6 min. The duration of immobility (s) during
the last 4 min of the 6 min video recording was quantified and analyzed.

RT. During the training phase, mice were placed on a rotating rod at a constant speed of 5 rpm
for 5 min, and the mice were placed back on the rod immediately after falling off the rod to minimize
accidental falls from the rod during the test phase. Then mice were rested for 1h. During the test
phase, the rotating rod was set to run at an acceleration of 20 rpm/min from 4 rpm to a maximum of
40 rpm for 5 min, and the mice were placed on the rotating rod at the beginning of the initial speed.
The formal test consisted of 3 replicate trials with a 30 min rest interval between each trial. The
average value of the fall latency (s) of the 3 repeated trials was calculated.

Molecular biology test
This study employed qRT-PCR and Western blot to assess the mRNA and protein expression levels
of GluN3A subunit in different brain regions of mice. Immunofluorescence staining, coupled with
confocal imaging, was utilized for staining analysis of neurons in the CA1 region and their projection
areas. Additionally, the spatial location of viral expression was confirmed by observing the
fluorescence of the injected viruses in each experiment.

qRT-PCR. After rapidly decapitating the experimental mice, the brain was promptly dissected and
placed in a freezing microtome. Specific brain region boundaries were delineated based on the
mouse brain atlas, and target brain regions were extracted using a 1-mm-diameter tissue extraction
needle [72], as shown in the Supplementary fig1b. Total RNA was extracted from the tissue samples
using TRIzol reagent (Cat# 15596018CN, Invitrogen, USA) following the manufacturer's instructions,
and the concentration was analyzed using NanoDrop (Thermo Scientific, Massachusetts, USA).
Subsequently, 500 ng of RNA samples were reverse transcribed using the M-MLV Reverse
Transcriptase (Cat# M1701, Promega, USA) following the reverse transcription kit instructions, with a
two-step reaction condition: 70°C for 5 min, 37°C for 60 min, to generate cDNA. SYBR Green PCR
Master Mix (Cat# CW2601M, Cowin Biotech, China) was used for fluorescence qPCR reactions. The
real-time PCR instrument reaction steps were: a single cycle (95°C/10 min); 40 cycles (95°C/15 s;
60°C/1 min); a single cycle (95°C/15 s; 60°C/30 s; 95°C/15 s). Analysis was performed using the
ΔΔC(t) method, with GAPDH as the reference gene for standardization. The primer sequences used
were: Grin3A: forward: 5'-AGTACCTGAAGAATGATCCAGAGT-3', reverse: 5'-

TCAGATATGTTAGAGGTCAACGG-3'; Gapdh: forward: 5'-GCATGGCCTTCCGTGTTCCT-3', reverse:
5'-CCCTGTTGCTGTAGCCGTAT-3'.

Western blot. The target brain area tissue was homogenized in RIPA lysis buffer containing 1%
PMSF (Cat# P0100, Solarbio, China). The collected protein supernatant was quantified using a BCA
protein quantification kit (Cat# PC0020, Solarbio, China), and then protein samples were mixed with
4× protein loading buffer (Cat# P1015, Solarbio, China) for denaturation. The equal amounts of
samples were loaded onto a 10% SDS-polyacrylamide gel (Cat# CW0022M, Cowin Biotech, China)
for separation and then transferred to a 0.2 µm PVDF membrane (Cat# 03010040001, Millipore,
Germany). The PVDF membrane was blocked in TBST solution containing 5% milk powder and then
incubated overnight at 4°C with diluted primary antibodies (5% milk powder TBST): Grin3A (1:500;
Cat# orb694249, Biorbyt, UK); Gapdh (1:3000; Cat# 2118, Cell Signaling, USA). After washing with
TBST, the membrane was incubated at room temperature for 2 h with secondary antibodies (5% milk
powder TBST) containing horseradish peroxidase: Goat Anti-Rabbit IgG H&L (1:3000; Cat# ZB-2301,
Zhongshan Golden Bridge, China). After washing again, the ECL luminescent liquid (Cat# P90719,
Millipore, Germany) was added to the PVDF membrane, and the immunoblot signal was captured
using an imaging system (FluorChem E, Protein Simple, California, USA).

Immunofluorescence staining. 90 min after the completion of the FST or CNO injection, mice
were intraperitoneally injected with pentobarbital (100 mg/kg) for anesthesia. Following anesthesia,
mice were perfused with 30 ml of phosphate-buffered saline (PBS) through the heart, followed by 30
ml of 4% paraformaldehyde (PFA). The brain was removed, fixed in 4% PFA for 24 h at 4°C, and then
transferred to PBS with 30% sucrose for dehydration. After the brain settled, 40 µm thick coronal brain
slices were obtained using a freezing microtome (CM3050S, Leica, Wetzlar, Germany) and stored at -
80°C in an anti-freezing solution (50% PBS, 30% ethylene glycol, 20% glycerol). For
immunofluorescence, the brain slices to be stained were washed with PBS, placed in PBST (0.1%
TritonX-100, PBS) containing 5% goat serum at room temperature for 1.5 h, and then transferred to
the primary antibody solution for overnight incubation at 4°C. The primary antibodies used were c-Fos
(1:500; Cat# SYSY.226003, Synaptic Systems, Germany) and DYKDDDDK Tag (1:500; Cat#14793,
Cell Signaling, USA). After washing with PBS, the slices were transferred to the secondary antibody
solution at room temperature away from light for 1.5 h. The secondary antibodies used were Goat
Anti-Rabbit IgG H&L (Alexa Fluor® 488) (1:500; Cat# ab150077, Abcam, UK) and Goat anti-Rabbit
IgG (H+L), Alexa Fluor™ 546 (1:500; Cat#A11010, Invitrogen, USA). After washing with PBS, the
slices were incubated in PBS with DAPI (1:500; Cat# C0060, Solarbio, China) at room temperature
away from light for 10 min. Following another PBS wash, the slices were mounted on glass slides
using a mounting medium (Cat# ab104139, Abcam, UK).

Confocal imaging and analyses. Sliced brains or immunofluorescent stained brain slices were
imaged using a high-speed confocal imaging system (SpinSR, Olympus, Tokyo, Japan) and a digital
slice scanning and analysis system (DMI8, Leica, Wetzlar, Germany) under 10× and 20× objectives.
Automated cell counting for c-Fos-positive neurons was performed using the ImageJ Fiji plugin
(http://imagej.nih.gov/ij/). Two to four images were selected in each animal for statistical analysis of
the brain regions.

In vivo manipulation
AAV. AAV2/9-syn-m-Grin3A-3×flag-Null (titer 1.6×1012 vg/mL) and AAV2/9-syn-ZsGreen (titer
1.6×1012 vg/mL), AAV2/9- syn-hM3D(Gq)-mCherry (titer 1.2×1012 vg/mL), AAV2/9-syn-hM4D(Gi)-

mCherry (titer 1.4×1012 vg/mL), and AAV2/9-syn-mCherry (titer 1.5×1012 vg/mL) were purchased
from Hanbio Biotechnology, Shanghai, China. AAV2/PHP.eB-mCaMKIIα-GCaMP8s-WPRE-pA (titer
1.1×1013 vg/mL) was purchased from Taitool Bio, Shanghai, China. The viral vectors were aliquoted
and stored at -80°C.

Stereotaxic surgery. Mice were anesthetized with isoflurane and secured in a stereotaxic
apparatus (RWD Life Science, Shenzhen, China). After exposing the skull, the corresponding virus
was slowly injected into the target location (AP: -2.90 mm; ML ± 3.30 mm; DV: -2.90 mm) using a
Stoelting pump (QSI storage pump, Stoelting, Wisconsin, USA) attached to a Hamilton syringe. After
injection, the syringe was left in place for 10 min before slow withdrawal. The mice were allowed to
recover for 14 days before behavioral testing.

Regulation of GluN3A subunit and virus tracing experiments. 0.15 µL (unilateral) of AAV2/9-
syn-m-Grin3A-3×flag-Null or AAV2/9-syn-ZsGreen was bilaterally injected into CA1i at a rate of 0.05
µL/min.

Chemogenetic experiments. 0.15 µL (unilateral) of AAV2/9-syn- hM3D(Gq)-mCherry or AAV2/9-
syn-hM4D(Gi)-mCherry (or AAV2/9-syn- mCherry) was injected into each hemisphere of CA1i at a
rate of 0.05 µL/min. 14 days later, CNO (Cat# HY-17366, MCE, USA) was dissolved in dimethyl
sulfoxide (DMSO, Cat# D8418, Sigma, Germany) to create a 100 mg/mL stock solution. Before
injection, further dilution was made with 0.9% saline to achieve a CNO injection quality of 0.2 mg/kg
(injection volume: 10 mL/kg) for chemogenetic activation experiments and 2 mg/kg for chemogenetic
inhibition experiments. CNO was intraperitoneally injected 30 min before behavioral testing.

Fiber photometry experiments. AAV2/PHP.eB-mCaMKIIα-GCaMP8s-WPRE-pA was diluted 1:4
in cerebrospinal fluid. 0.2 µL (unilateral) of the viral diluted solution was injected into the bilateral CA1i
at a rate of 0.05 µL/min. Then, a ceramic optical fiber (200 µm in diameter, 3 mm length, 0.37 NA,
Inper Technology Co., Ltd., Hangzhou, China) was implanted above the CA1i region (AP: -2.90 mm;
ML ±3.30 mm; DV: -2.85 mm). The fiber was secured to the skull with dental cement. 14 days later,
power levels of 470 nm excitation light and 410 nm excitation light in the Inper Signal fiber photometry
system (Inper Technology Co., Ltd., Hangzhou, China) were measured using a photometer and
adjusted to 25 µW and 10-15 µW, respectively. Prior to the TST, the bundled fiber was connected to
the optical fiber implant in the mouse's head for a two-min adaptation. The Inper Signal system
recorded Ca^{2+} activity in the CA1i region of mice suspended and during tail suspension tests,
synchronizing with a camera to record the behavioral state. Inper Plot software was used to analyze
and statistically process the fluorescence change data (F) during hanging and struggling events,
calculating the relative fluorescence change $\Delta F/F_0$ ($\Delta F/F_0 = (F-F_0)/F_0$). Mice with fiber implantation
positions confirming the expected target were included in data analysis.

Drug delivery experiments. After injecting chemogenetic viruses into bilateral CA1i, 22-gauge
single-guide cannulas (RWD Life Science, Shenzhen, China) were bilaterally implanted 0.5 mm above
the target positions (AP: -2.90 mm; ML: ±3.30 mm; DV: -2.40 mm). The cannulas were fixed on the
skull surface with dental cement. 14 days later, 10 mg/mL D-serine was prepared in CSF. 30 min
before the behavioral test, mice were anesthetized with isoflurane, and an injection inner tube
exceeding the cannula target position by 0.5 mm was inserted into the mouse's head cannula.
Subsequently, using a dual-channel microinjection pump (RWD Life Science, Shenzhen, China), 0.5
µL (unilateral) of D-serine or CSF was injected into the bilateral CA1i region at a rate of 0.2 µL/min.
After completion, the inner tube was left in place for 5 min before slow withdrawal.

SnRNAseq Analysis

The samples from "Control WT" and "RSD WT" in GSE253687 were reanalyzed using the standard analysis pipeline of Seurat (version 4.4.0)[73]. Briefly, only cells expressing at least 200 genes and genes containing non-zero values as well as associated with at least 3 cells were maintained for subsequent analysis. "NormalizeData" functions were used to normalize data. Highly variable genes were observed using the "FindVariableFeatures" function with default parameters setting, and these highly variable genes were used to perform principal components analysis (PCA). Dimensionality reduction was performed using the "RunUMAP" function, where reduction was set to 'pca' and dims to 1:50. Default resolution was used for clustering, using "FindNeighbors" and "FindClusters" functions. Next, we extracted the cells from the CA1 excitatory neuron subpopulation (Mpped1 positive) and performed the aforementioned analysis again. Due to the lack of effective marker genes for screening the CA1 inhibitory neuron subpopulation, we could not extract and analyze the inhibitory neuron subpopulation of CA1[74]. The R package ggplot2[75] was used for visualization.

Statistical Analysis

All data are presented as mean ± standard error of the mean (M±SEM) and analyzed using GraphPad Prism 9.0 software. Normality distribution was assessed using the Anderson-Darling test. Differences between two groups were evaluated using unpaired two-tailed Student's t-test, and when variances were unequal, Welch's correction was applied. Analysis of variance with one or two factors was used for comparisons among three groups or more, followed by FDR: Benjamini-Krieger-Yekutieli correction for multiple comparisons. A p-value < 0.05 was considered statistically significant.

REFERENCES

Roddy DW, Farrell C, Doolin K, Roman E, Tozzi L, Frodl T, et al. The Hippocampus in Depression: More Than the Sum of Its Parts? Advanced Hippocampal Substructure Segmentation in Depression. Biol Psychiatry. 2019;85(6):487-97.

Anacker C, Luna VM, Stevens GS, Millette A, Shores R, Jimenez JC, et al. Hippocampal neurogenesis confers stress resilience by inhibiting the ventral dentate gyrus. Nature. 2018;559(7712):98-102.

Strange BA, Witter MP, Lein ES, Moser EI. Functional organization of the hippocampal longitudinal axis. Nat Rev Neurosci. 2014;15(10):655-69.

Fanselow MS, Dong HW. Are the dorsal and ventral hippocampus functionally distinct structures? Neuron. 2010;65(1):7-19.

Tzilivaki A, Tukker JJ, Maier N, Poirazi P, Sammons RP, Schmitz D. Hippocampal GABAergic interneurons and memory. Neuron. 2023;111(20):3154-75.

Schultz C, Engelhardt M. Anatomy of the hippocampal formation. Front Neurol Neurosci. 2014;34:6-17.

Ayhan F, Kulkarni A, Berto S, Sivaprakasam K, Douglas C, Lega BC, et al. Resolving cellular and molecular diversity along the hippocampal anterior-to-posterior axis in humans. Neuron. 2021;109(13):2091-105 e6.

Levone BR, Cryan JF, O'Leary OF. Specific sub-regions of the longitudinal axis of the hippocampus mediate behavioural responses to chronic psychosocial stress. Neuropharmacology. 2021;201:108843.

Bienkowski MS, Bowman I, Song MY, Gou L, Ard T, Cotter K, et al. Integration of gene expression and brain-wide connectivity reveals the multiscale organization of mouse hippocampal networks. Nat Neurosci. 2018;21(11):1628-43.

Okuyama T, Kitamura T, Roy DS, Itohara S, Tonegawa S. Ventral CA1 neurons store social memory. Science. 2016;353(6307):1536-41.

Pachernegg S, Strutz-Seebohm N, Hollmann M. GluN3 subunit-containing NMDA receptors: not just one-trick ponies. Trends Neurosci. 2012;35(4):240-9.

Perez-Otano I, Larsen RS, Wesseling JF. Emerging roles of GluN3-containing NMDA receptors in the CNS. Nat Rev Neurosci. 2016;17(10):623-35.

Bossi S, Pizzamiglio L, Paoletti P. Excitatory GluN1/GluN3A glycine receptors (eGlyRs) in brain signaling. Trends Neurosci. 2023;46(8):667-81.

Grand T, Abi Gerges S, David M, Diana MA, Paoletti P. Unmasking GluN1/GluN3A excitatory glycine NMDA receptors. Nat Commun. 2018;9(1):4769.

Pizzamiglio L, Morice E, Cardoso C, Bossi S, Mailhes-Hamon C, Heimendahl MV, et al. Excitatory glycine receptors control ventral hippocampus synaptic plasticity and anxiety-related behaviors. Proc Natl Acad Sci U S A. 2025;122(37):e2501118122.

Bossi S, Dhanasobhon D, Ellis-Davies GCR, Frontera J, de Brito Van Velze M, Lourenco J, et al. GluN3A excitatory glycine receptors control adult cortical and amygdalar circuits. Neuron. 2022;110(15):2438-54 e8.

Conde-Dusman MJ, Dey PN, Elia-Zudaire O, Rabaneda LG, Garcia-Lira C, Grand T, et al. Control of protein synthesis and memory by GluN3A-NMDA receptors through inhibition of GIT1/mTORC1 assembly. Elife. 2021;10.

18 Chatterton JE, Awobuluyi M, Premkumar LS, Takahashi H, Talantova M, Shin Y, et al.
Excitatory glycine receptors containing the NR3 family of NMDA receptor subunits. Nature.
2002;415(6873):793-8.

19 Murillo A, Navarro AI, Puelles E, Zhang Y, Petros TJ, Perez-Otano I. Temporal Dynamics and
Neuronal Specificity of Grin3a Expression in the Mouse Forebrain. Cereb Cortex.
2021;31(4):1914-26.

20 Otsu Y, Darcq E, Pietrajtis K, Matyas F, Schwartz E, Bessaih T, et al. Control of aversion by
glycine-gated GluN1/GluN3A NMDA receptors in the adult medial habenula. Science.
2019;366(6462):250-54.

21 Zhong W, Wu A, Berglund K, Gu X, Jiang MQ, Talati J, et al. Pathogenesis of sporadic
Alzheimer's disease by deficiency of NMDA receptor subunit GluN3A. Alzheimers Dement.
2022;18(2):222-39.

22 Roberts AC, Diez-Garcia J, Rodriguiz RM, Lopez IP, Lujan R, Martinez-Turrillas R, et al.
Downregulation of NR3A-containing NMDARs is required for synapse maturation and
memory consolidation. Neuron. 2009;63(3):342-56.

23 Wang J, Zhang K, Chen X, Liu X, Teng H, Zhao M, et al. Epigenetic Activation of ASCT2 in
the Hippocampus Contributes to Depression-Like Behavior by Regulating D-Serine in Mice.
Front Mol Neurosci. 2017;10:139.

24 MacKay MB, Kravtsenyuk M, Thomas R, Mitchell ND, Dursun SM, Baker GB. D-Serine:
Potential Therapeutic Agent and/or Biomarker in Schizophrenia and Depression? Front
Psychiatry. 2019;10:25.

25 Otte DM, Barcena de Arellano ML, Bilkei-Gorzo A, Albayram O, Imbeault S, Jeung H, et al.
Effects of Chronic D-Serine Elevation on Animal Models of Depression and Anxiety-Related
Behavior. PLoS One. 2013;8(6):e67131.

26 Zhang W, Wang W, Zhou Y, Wang J. A comparative study of blood and hippocampal D-serine
change patterns in drug-naive patients and animal models of depression. Psychiatry Res.
2025;348:116453.

27 Chen Z, Tang Z, Zou K, Huang Z, Liu L, Yang Y, et al. D-Serine produces antidepressant-like
effects in mice through suppression of BDNF signaling pathway and regulation of synaptic
adaptations in the nucleus accumbens. Mol Med. 2021;27(1):127.

28 Malkesman O, Austin DR, Tragon T, Wang G, Rompala G, Hamidi AB, et al. Acute D-serine
treatment produces antidepressant-like effects in rodents. Int J Neuropsychopharmacol.
2012;15(8):1135-48.

29 Wei IH, Chen KT, Tsai MH, Wu CH, Lane HY, Huang CC. Acute Amino Acid d-Serine
Administration, Similar to Ketamine, Produces Antidepressant-like Effects through Identical
Mechanisms. J Agric Food Chem. 2017;65(49):10792-803.

30 Pignatelli M, Tejeda HA, Barker DJ, Bontempi L, Wu J, Lopez A, et al. Cooperative synaptic
and intrinsic plasticity in a disynaptic limbic circuit drive stress-induced anhedonia and
passive coping in mice. Mol Psychiatry. 2021;26(6):1860-79.

31 Lu JJ, Wu PF, He JG, Li YK, Long LH, Yao XP, et al. BNIP3L/NIX-mediated mitophagy
alleviates passive stress-coping behaviors induced by tumor necrosis factor-alpha. Mol
Psychiatry. 2023;28(12):5062-76.

32 Jing W, Zhang T, Liu J, Huang X, Yu Q, Yu H, et al. A circuit of COCH neurons encodes
social-stress-induced anxiety via MTF1 activation of Cacna1h. Cell Rep. 2021;37(13):110177.

33   Duman RS, Aghajanian GK, Sanacora G, Krystal JH. Synaptic plasticity and depression: new
insights from stress and rapid-acting antidepressants. Nat Med. 2016;22(3):238-49.

34   Li Y, Fan C, Wang C, Wang L, Yi Y, Mao X, et al. Stress-induced reduction of Na(+)/H(+)
exchanger isoform 1 promotes maladaptation of neuroplasticity and exacerbates depressive
behaviors. Sci Adv. 2022;8(45):eadd7063.

35   Su T, Lu Y, Fu C, Geng Y, Chen Y. GluN2A mediates ketamine-induced rapid antidepressant-
like responses. Nat Neurosci. 2023;26(10):1751-61.

36   Xia F, Fascianelli V, Vishwakarma N, Ghinger FG, Kwon A, Gergues MM, et al. Understanding
the neural code of stress to control anhedonia. Nature. 2025;637(8046):654-62.

37   Ma YN, Zhang CC, Sun YX, Liu X, Li XX, Wang H, et al. Dorsal CA1 NECTIN3 Reduction
Mediates Early-Life Stress-Induced Object Recognition Memory Deficits in Adolescent
Female Mice. Neurosci Bull. 2025;41(2):243-60.

38   Kim J, Lei Y, Lu XY, Kim CS. Glucocorticoid-glucocorticoid receptor-HCN1 channels reduce
neuronal excitability in dorsal hippocampal CA1 neurons. Mol Psychiatry. 2022;27(10):4035-
49.

39   Ma H, Li C, Wang J, Zhang X, Li M, Zhang R, et al. Amygdala-hippocampal innervation
modulates stress-induced depressive-like behaviors through AMPA receptors. Proc Natl Acad
Sci U S A. 2021;118(6).

40   Maras PM, Molet J, Chen Y, Rice C, Ji SG, Solodkin A, et al. Preferential loss of dorsal-
hippocampus synapses underlies memory impairments provoked by short, multi-modal stress.
Mol Psychiatry. 2014;19(7):745.

41   Huang M, Bao J, Tao X, Niu Y, Li K, Wang J, et al. Ventral Hippocampal CA1 GADD45B
Regulates Susceptibility to Social Stress by Influencing NMDA Receptor-Mediated Synaptic
Plasticity. Neurosci Bull. 2025;41(3):406-20.

42   Diniz CR, Casarotto PC, Joca SR. NMDA-NO signaling in the dorsal and ventral
hippocampus time-dependently modulates the behavioral responses to forced swimming
stress. Behav Brain Res. 2016;307:126-36.

43   Chang CH, Su CL, Gean PW. Mechanism underlying NMDA blockade-induced inhibition of
aggression in post-weaning socially isolated mice. Neuropharmacology. 2018;143:95-105.

44   Goodman EJ, Biltz RG, Packer JM, DiSabato DJ, Swanson SP, Oliver B, et al. Enhanced fear
memory after social defeat in mice is dependent on interleukin-1 receptor signaling in
glutamatergic neurons. Mol Psychiatry. 2024.

45   Zhang M, Kong X, Chen J, Liu W, Liu C, Dou X, et al. Dysfunction of GluN3A subunit is
involved in depression-like behaviors through synaptic deficits. J Affect Disord. 2023;332:72-
82.

46   Commons KG, Cholanians AB, Babb JA, Ehlinger DG. The Rodent Forced Swim Test
Measures Stress-Coping Strategy, Not Depression-like Behavior. ACS Chem Neurosci.
2017;8(5):955-60.

47   Cryan JF, Mombereau C, Vassout A. The tail suspension test as a model for assessing
antidepressant activity: review of pharmacological and genetic studies in mice. Neurosci
Biobehav Rev. 2005;29(4-5):571-625.

48   Zapater-Fajari M, Crespo-Sanmiguel I, Pulopulos MM, Hidalgo V, Salvador A. Resilience and
Psychobiological Response to Stress in Older People: The Mediating Role of Coping
Strategies. Front Aging Neurosci. 2021;13:632141.

49 Andalman AS, Burns VM, Lovett-Barron M, Broxton M, Poole B, Yang SJ, et al. Neuronal
Dynamics Regulating Brain and Behavioral State Transitions. Cell. 2019;177(4):970-85 e20.

50 Kent M, Bardi M, Hazelgrove A, Sewell K, Kirk E, Thompson B, et al. Profiling coping
strategies in male and female rats: Potential neurobehavioral markers of increased resilience
to depressive symptoms. Horm Behav. 2017;95:33-43.

51 Wood SK, Bhatnagar S. Resilience to the effects of social stress: evidence from clinical and
preclinical studies on the role of coping strategies. Neurobiol Stress. 2015;1:164-73.

52 Kato T. Testing of the coping flexibility hypothesis based on the dual-process theory:
Relationships between coping flexibility and depressive Symptoms. Psychiatry Res.
2015;230(2):137-42.

53 Kato T. The impact of coping flexibility on the risk of depressive symptoms. PLoS One.
2015;10(5):e0128307.

54 Kato T. Development of the Coping Flexibility Scale: evidence for the coping flexibility
hypothesis. J Couns Psychol. 2012;59(2):262-73.

55 Rao JS, Harry GJ, Rapoport SI, Kim HW. Increased excitotoxicity and neuroinflammatory
markers in postmortem frontal cortex from bipolar disorder patients. Mol Psychiatry.
2010;15(4):384-92.

56 Crawley O, Conde-Dusman MJ, Perez-Otano I. GluN3A NMDA receptor subunits: more
enigmatic than ever? J Physiol. 2022;600(2):261-76.

57 Lee JH, Wei L, Deveau TC, Gu X, Yu SP. Expression of the NMDA receptor subunit GluN3A
(NR3A) in the olfactory system and its regulatory role on olfaction in the adult mouse. Brain
Struct Funct. 2016;221(6):3259-73.

58 Yao Y, Mayer ML. Characterization of a soluble ligand binding domain of the NMDA receptor
regulatory subunit NR3A. J Neurosci. 2006;26(17):4559-66.

59 Ragguett RM, Rong C, Kratiuk K, McIntyre RS. Rapastinel - an investigational NMDA-R
modulator for major depressive disorder: evidence to date. Expert Opin Investig Drugs.
2019;28(2):113-19.

60 Keck ME, Welt T, Muller MB, Uhr M, Ohl F, Wigger A, et al. Reduction of hypothalamic
vasopressinergic hyperdrive contributes to clinically relevant behavioral and neuroendocrine
effects of chronic paroxetine treatment in a psychopathological rat model.
Neuropsychopharmacology. 2003;28(2):235-43.

61 Carhart-Harris RL, Nutt DJ. Serotonin and brain function: a tale of two receptors. J
Psychopharmacol. 2017;31(9):1091-120.

62 Mahar I, Bambico FR, Mechawar N, Nobrega JN. Stress, serotonin, and hippocampal
neurogenesis in relation to depression and antidepressant effects. Neurosci Biobehav Rev.
2014;38:173-92.

63 Chen L, Jiao J, Lei F, Zhou B, Li H, Liao P, et al. Ezrin-mediated astrocyte-synapse signaling
regulates cognitive function via astrocyte morphological changes in fine processes in male
mice. Brain Behav Immun. 2024;124:177-91.

64 Gomez-Galan M, De Bundel D, Van Eeckhaut A, Smolders I, Lindskog M. Dysfunctional
astrocytic regulation of glutamate transmission in a rat model of depression. Mol Psychiatry.
2013;18(5):582-94.

65   Basu AC, Tsai GE, Ma CL, Ehmsen JT, Mustafa AK, Han L, et al. Targeted disruption of serine
racemase affects glutamatergic neurotransmission and behavior. Mol Psychiatry.
2009;14(7):719-27.

66   Levin R, Dor-Abarbanel AE, Edelman S, Durrant AR, Hashimoto K, Javitt DC, et al.
Behavioral and cognitive effects of the N-methyl-D-aspartate receptor co-agonist D-serine in
healthy humans: initial findings. J Psychiatr Res. 2015;61:188-95.

67   Yoon SH, Song WS, Chung G, Kim SJ, Kim MH. Activity in the dorsal hippocampus-mPFC
circuit modulates stress-coping strategies during inescapable stress. Exp Mol Med.
2024;56(9):1921-35.

68   Bagot RC, Parise EM, Pena CJ, Zhang HX, Maze I, Chaudhury D, et al. Ventral hippocampal
afferents to the nucleus accumbens regulate susceptibility to depression. Nat Commun.
2015;6:7062.

69   Xu H, Wang J, Jing H, Ellenbroek B, Shao F, Wang W. mPFC GABAergic transmission
mediated the role of BDNF signaling in cognitive impairment but not anxiety induced by
adolescent social stress. Neuropharmacology. 2021;184:108412.

70   Golden SA, Covington HE, 3rd, Berton O, Russo SJ. A standardized protocol for repeated
social defeat stress in mice. Nat Protoc. 2011;6(8):1183-91.

71   Menard C, Pfau ML, Hodes GE, Kana V, Wang VX, Bouchard S, et al. Social stress induces
neurovascular pathology promoting depression. Nat Neurosci. 2017;20(12):1752-60.

72   Franklin KBJ, Paxinos G. Paxinos and Franklin's The mouse brain in stereotaxic coordinates.
Fourth edition. ed. Academic Press, an imprint of Elsevier: Amsterdam; 2013.

73   Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell gene
expression data. Nat Biotechnol. 2015;33(5):495-502.

74   Habib N, Li Y, Heidenreich M, Swiech L, Avraham-Davidi I, Trombetta JJ, et al. Div-Seq:
Single-nucleus RNA-Seq reveals dynamics of rare adult newborn neurons. Science.
2016;353(6302):925-8.

75   Wickham H, Wickham H. Data analysis. Springer; 2016.

ACKNOWLEDGEMENTS

We are grateful to the lab of Tifei Yuan (Shanghai Mental health center, Shanghai, China) for providing the GluN3A KO mice. We are grateful to Wenyu Tian (Shandong University, Shandong, China) for database technologies and analysis assistance. We greatly appreciate the excellent work of the technical support staff at the Institutional Center for Shared Technologies and Facilities of Institute of Psychology, Chinese Academy of Sciences.

AUTHOR CONTRIBUTIONS

Methodology, data curation and investigation, writing-original draft&editing: WZ; Research conceptualization, project administration, funding acquisition and editing: JW; Research conceptualization, resources, project administration, funding acquisition, supervision and writing-review&editing: WW; Methodology: MZ, JD, LZ and HY participated.

FUNDING

This work was supported by the National Natural Science Foundation of China [Grant No. 82071517, U21A20364, 31900731], National Key R&D Program of China [Grant No. 2017YFE0126500], and the Scientific Foundation of Institute of Psychology, Chinese Academy of Sciences (No. E2CX4115CX).

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information

FIGURE AND FIGURE LEGENDS

883 **Figure 1. CSDS induces multiple stress-related behaviors and specifically reduces GluN3A**

884 **expression in the CA1i region, while GluN3A KO specifically promotes passive coping.**

a Experimental scheme of CSDS modeling and behavioral tests. **b** Schematic of CSDS. **c** Example trajectory heatmap of SIT in control mice (left) and stress mice (right). **d** Social interaction ratio of SIT (n = 9 per group). **e** Immobility time in the TST (n = 8 per group). **f** Immobility time in the FST (n = 6 in Control, n = 7 in Stress). **g** Example trajectory heatmap of OFT in control mice (left) and stress mice (right). **h** The total distance traveled in OFT (n = 8 in Control, n = 7 in Stress). **i** The number of entries to centre area of OFT. **j** The time in the centre area of OFT. **k** Example trajectory heatmap of EPM in control mice (left) and stress mice (right). **l** The number of entries to open arms of EPM (n = 9 per group). **m** Percentage of time spent in the open arms of EPM. **n** Quantitative data of *Grin3A* mRNA in target brain area from control and stress mice (CA1 d: n = 8 in Control, n = 14 in Stress; CA1 i: n = 10 in Control, n = 15 in Stress; CA1 v: n = 8 in Control, n = 6 in Stress; CA3: n = 7 per group; DG: n = 8 in Control, n = 16 in Stress). **o** Quantitative data of GluN3A protein in target brain area from control and stress mice (CA1 v: n = 6 in Control, n = 7 in Stress; CA1 i: n = 4 in Control, n = 5 in Stress; CA3: n = 7 per group). **p** Representative western blots for dissected target brain area. **q** Experimental scheme of behavioral studies. **r** Social interaction ratio of SIT (n = 8 per group). **s** Immobility time in the TST (n = 9 per group). **t** Immobility time in the FST (n = 8 per group). **u** The total distance traveled in OFT (n = 8 per group). **v** The entries of centre area in OFT. **w** The time in the centre area of OFT. **x** The number of entries to open arms of EPM (n = 8 per group). **y** Percentage of time spent in the open arms of EPM. $* P < 0.05$, $*** P < 0.001$.

Figure 2. Overexpression of GluN3A in CA1i reverses passive coping behavior in CSDS mice
and GluN3A KO mice.

a Experimental scheme of CSDS modeling, virus injection, and behavior tests. **b** Schematic target
locations and immunofluorescence images of ZsGreen or flag expression of Virus in the CA1i. Scale
bar, 1 mm. **c** Social interaction ratio of SIT (n = 7, 8, 7 and 7 in control:AAV-ZsGreen, control:AAV-
Grin3A, stress:AAV-ZsGreen and stress:AAV-Grin3A, respectively). **d** Immobility time in the TST (n =
8 per group). **e** Immobility time in the FST (n = 8, 8, 7 and 9 in control:AAV-ZsGreen, control:AAV-
Grin3A, stress:AAV-ZsGreen and stress:AAV-Grin3A, respectively). **f** The total distance traveled in
OFT (n = 7, 8, 8 and 9 in control:AAV-ZsGreen, control:AAV-Grin3A, stress:AAV-ZsGreen and
stress:AAV-Grin3A, respectively). **g** The number of entries to centre area of OFT. **h** The time in the
centre area of OFT. **i** The number of entries to open arms of EPM (n = 8 per group). **j** Percentage of
time spent in the open arms of EPM. **k** Experimental scheme of virus injection and behavior tests. **l**
Immobility time in the TST (n = 8, 6 and 6 in WT+AAV-ZsGreen, GluN3A KO+AAV-ZsGreen and
GluN3A KO+AAV-Grin3A, respectively). **m** Immobility time in the FST (n = 7, 6 and 6 in WT+AAV-
ZsGreen, GluN3A KO+AAV-ZsGreen and GluN3A KO+AAV-Grin3A, respectively). **n** The total distance

traveled in OFT (n = 8, 6 and 7 in WT+AAV-ZsGreen, GluN3A KO+AAV-ZsGreen and GluN3A
KO+AAV-Grin3A, respectively). * P < 0.05, ** P < 0.01.

Figure 3. Activity of CA1i pyramidal neurons mediates the GluN3A-dependent regulation of
passive coping behavior.
a Identification of the various cell types in hippocampus using UMAP clustering. **b** Visualization of the
distribution of *Grin3A* expression in excitatory neurons in CA1 for each condition: Control WT, RSD
WT. **c** Dot plot shows the expression of *Grin3A* in excitatory neurons of each group. The size of the
circle representing the percentage of neurons expressing the gene. The color of the circle
representing the average level of the gene expression in neurons. **d** Violin plots shows the expression
level of *Grin3A* in excitatory neurons of each group. **e** Statistical region of c-Fos-positive cells in CA1i
(ZsGreen: green, c-Fos: red, Dapi: blue). Scale bar, 1 mm. **f** Quantification of c-Fos-positive cells in
CA1i pyramidal layer neurons of each group of mice. (n = 6, 6, 5 and 7 in control:AAV-ZsGreen,
control:AAV-Grin3A, stress:AAV-ZsGreen and stress:AAV-Grin3A, respectively, 4 slices for each dot).
g-h Representative images of c-Fos-positive cells in the CA1i of each group of mice (c-Fos: red, Dapi:
blue). Scale bar, 100 μm. **i** Quantification of c-Fos-positive cells in CA1i pyramidal layer neurons of
each group of mice. (n = 5, 6 and 6 in WT+ No-FST, WT+ FST and GluN3A KO+FST, respectively, 4
slices for each dot). **j** Schematic illustrating fiber placement and representative images of GCaMP8s
expression. Scale bar, 100 μm. **k** Top, Example fluorescence trace of the GCaMP signal during TST.
Gray shaded area indicates struggling behavior; white area indicates immobile behavior. Bottom,
heatmap of fluorescence intensity. **l** Representative heatmaps of GCaMP8s transient ΔF/F event
locked to struggling. Each row plots one mice, and a total of 11 mice are illustrated. **m** Dynamics of
GCamp8s signaling in mice 5 seconds before and 10 seconds after struggling in the TST (mean ΔF/F
± s.e.m.). **n** Changes in average ΔF/F of WT and KO mice during the struggle (n = 5 in WT, n = 6 in
KO). * P < 0.05, ** P < 0.01.

Figure 4. Bidirectional chemogenetic modulation of CA1i pyramidal neurons directly mimics or reverses CSDS-induced passive coping behavior.

a Schematic target locations and immunofluorescence images of hM3D(Gq)-mCherry or mCherry expression of Virus in the CA1i. Scale bar, 1 mm. **b** Statistical regions of c-Fos-positive cells in CA1i (mCherry: red, c-Fos: green, Dapi: blue). Scale bar, 1 mm. **c** Representative images of c-Fos-positive cells in the CA1i of the saline and 0.2 mg/kg CNO treatment mice (c-Fos: green, Dapi: blue). Scale bar, 200 μm. **d** Experimental scheme of CSDS modeling, virus injection, and behavior tests without or with chemogenetic activation. **e** Social interaction ratio of SIT (*n* = 6 per group). **f** Immobility time in

the TST (n = 6, 6, 6 and 7 in control:AAV-mCherry, control:AAV-hM3D(Gq), stress:AAV-mCherry and

stress:AAV-hM3D(Gq), respectively). **g** Immobility time in the FST (n = 6, 6, 6 and 7 in control:AAV-

mCherry, control:AAV-hM3D(Gq), stress:AAV-mCherry and stress:AAV-hM3D(Gq), respectively). **h**

The total distance traveled in OFT (n = 8, 8, 7 and 8 in control:AAV-mCherry, control:AAV-hM3D(Gq),

stress:AAV-mCherry and stress: AAV-hM3D(Gq), respectively). **i** The number of entries to centre area

of OFT. **j** The time in the centre area of OFT. **k** Experimental scheme of virus injection, and behavior

tests without or with chemogenetic inactivation. **l** Schematic target locations and immunofluorescence

images of hM4D(Gi)-mCherry or mCherry expression of Virus in the CA1i. Scale bar, 1 mm. **m**

Immobility time in the TST (n = 7 per group). **n** Immobility time in the FST (n = 7 in AAV-mCherry, n =

8 in AAV-hM4D(Gi)). **o** The total distance traveled in OFT (n = 9 per group). **p** The number of entries

to centre area of OFT. **q** The time in the centre area of OFT. * $P < 0.05$, ** $P < 0.01$.

Figure 5. GluN3A-dependent potentiation of CA1i pyramidal neuronal activity is essential for the rapid shift in coping strategy by D-serine.

a Experimental scheme of CSDS modeling, virus injection, cannula implantation, and behavior tests without or with D-serine delivery. **b** Schematic target locations injection of D-serine (CSF) and Virus in the CA1i. Scale bar, 1 mm. **c** Immobility time in the TST (n = 7, 8, 7, 9, 7 and 7 in Control-CSF, Control-D-serine, GluN3A KO:D-serine, Stress-CSF, Stress-D-serine and stress-D-serine-hM4D(Gi), respectively). **d** Immobility time in the FST (n = 7, 8, 7, 10, 6 and 8 in Control-CSF, Control-D-serine, GluN3A KO:D-serine, Stress-CSF, Stress-D-serine and stress-D-serine-hM4D(Gi), respectively). **e** The total distance traveled in OFT (n = 9, 9, 7, 9, 8 and 8 in Control-CSF, Control-D-serine, GluN3A

KO:D-serine, Stress-CSF, Stress-D-serine and stress-D-serine-hM4D(Gi), respectively). **f** The number

of entries to centre area of OFT. **g** The time in the centre area of OFT. **h** Statistical regions of c-Fos-

positive cells in CA1i (mCherry: red, c-Fos: green, Dapi: blue). Scale bar, 1 mm. **i** Representative

images of c-Fos-positive cells in the CA1i of each group of mice (c-Fos: green, Dapi: blue). Scale bar,

50 μm. **j** Quantification of c-Fos-positive cells in CA1i pyramidal layer neurons of each group of mice.

(per mm, *n* = 5 per group, 4 slices for each dot). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; $^{+}$ $P < 0.05$, $^{++}$ P

< 0.01; $^{\#}$ $P < 0.05$, $^{\#\#\#}P < 0.001$.

Figure 6. GluN3A-mediated activation of CA1i pyramidal neurons selectively influenced neuronal activity in downstream CA3 and IL circuits.

a Projection map of the hippocampal CA1i region generated from the Allen Brain Atlas database. **b** Representative images of virus tracing results following AAV2/9-syn-ZsGreen injection into the CA1i region. Scale bar, 1000 μm. **c** Statistical region of c-Fos-positive cells in CA3 (ZsGreen: green, c-Fos: red, DAPI: blue). Scale bar, 500 μm. **d** Quantification of c-Fos-positive cells in CA3 neurons of each

group of mice (n = 6, 7, 5 and 7 in control:AAV-ZsGreen, control:AAV-Grin3A, stress:AAV-ZsGreen

and stress:AAV-Grin3A, respectively, 4 slices for each dot). **e** Representative images of c-Fos-

positive cells in the CA3 of each group of mice (c-Fos: red, Dapi: blue). Scale bar, 500 µm. **f**

Statistical region of c-Fos-positive cells in IL (ZsGreen: green, c-Fos: red, DAPI: blue). Scale bar, 500

µm. **g** Quantification of c-Fos-positive cells in IL neurons of each group of mice (n = 4, 5, 5 and 5 in

control:AAV-ZsGreen, control:AAV-Grin3A, stress:AAV-ZsGreen and stress:AAV-Grin3A, respectively,

4 slices for each dot). h Representative images of c-Fos-positive cells in the IL of each group of mice

(c-Fos: red, Dapi: blue). Scale bar, 50 µm. * $P < 0.05$, ** $P < 0.01$.