<u>Conditional nodule-level sanctions provide long-term stability to the Pea-</u>

<u>*Rhizobium* symbiosis</u>

Thomas J. Underwood[1], Beatriz Jorrin[1], Lindsay A. Turnbull[1], Philip S. Poole[1]

[1]Department of Biology, University of Oxford, Oxford, United Kingdom

Author for correspondence:

Philip Poole

Email: philip.poole@biology.ox.ac.uk

Word count: 4342

Number of Figures: 5

Figures to be in colour: 1, 2, 3 & 5

Tables:

1 table providing a list and description of strains used

Supporting information

The following Supporting Information is available for this article:

Table. S1 Data sheet containing counts from flow cytometer of undifferentiated bacteria

and bacteroids from nodules (single occupancy and mixed nodules).

Doc.S2 R markdown file of statistical analysis.

Fig. S3 Representative iBright image of pea inoculated with two strains of differing

effectiveness.

Fig. S4 Representative confocal images of nodule sections of both nodule types from all

three co-inoculation combinations (Fix$^+$ vs Fix$^-$, Fix$^+$ vs Fix^{int}, Fix^{int} vs Fix$^-$) at 28-, 35- and 42-

days post inoculation.

<u>**Summary**</u>

• Legumes sanction root nodules containing rhizobial strains with low nitrogen fixation
rates (ineffective fixers). Pea (*Pisum sativum*) nodules contain both undifferentiated
bacteria and terminally differentiated nitrogen-fixing bacteroids. It is critical to
understand whether sanctions act on both bacteria and bacteroids. As only
undifferentiated bacteria are able to regrow from senescent nodules, if sanctioning
only acts on bacteroids then ineffective strains might evade sanctioning. In addition,
ineffective strains could potentially evade sanctioning by entering the same nodule
as an effective strain i.e., piggybacking. *P. sativum* was co-inoculated with pairwise
combinations of three strains of rhizobia with different effectiveness, to test
whether ineffective strains can evade sanctions in this way.

• We assessed the effect of sanctions on nodule populations of bacteria and
bacteroids using flow cytometry and confocal microscopy.

• We show that sanctioning affects bacteroids and bacteria, altering survival of
ineffective strains by reducing their numbers and hastening nodule senescence. In
nodules containing two strains differing in their nitrogen-fixing ability, both were
sanctioned equally. Thus, peas sanction whole nodules based on their nitrogen
output, but do not sanction at the cellular level.

• Our results demonstrate that conditional sanctioning in peas provides long-term
stability to symbiosis by effectively reducing fitness of ineffective strains.

Key words: Conditional sanctioning, Symbiosis, Mixed nodules, Nitrogen fixation, legumes,
Rhizobium

Introduction

Legumes have overcome nitrogen limitation by establishing a mutually beneficial interaction with nitrogen-fixing bacteria, hosted within nodules along their root systems (Poole et al., 2018). Certain species of legumes, like peas, form indeterminate nodules, in which some bacteria undergo terminal differentiation to become large swollen cells called bacteroids, which carry out the costly process of nitrogen fixation, but cannot resume their free-living existence (Mergaert et al., 2006). The plant provides carbon compounds in the form of photosynthetically-derived dicarboxylates in return for ammonia provided by the bacteroids, and this relationship continues until the plant's nitrogen requirements have been met (Udvardi & Poole, 2013). At this point nodule senescence ensues, and while the bacteroids then die, any undifferentiated bacteria are released into the soil, which can dramatically increase the concentration of the strain in question, potentially providing a significant fitness advantage (Timmers et al., 2000).

However, this interaction presents an evolutionary dilemma. A 'cheating' strain that reduces investment in the costly process of nitrogen fixation, might be able to produce more reproductive bacteria within the nodules, potentially giving it a fitness advantage. Legumes are known to have evolved sanctions to punish cheating bacteria by cutting off their supply of carbon compounds (Kiers et al., 2003; Oono et al., 2011; West et al., 2002; Westhoek et al., 2017) and these sanctions can be applied conditionally, with pea plants known to sanction nodules that contain an intermediate-fixing strain, only when a better strain is available (Westhoek et al., 2021). But such sanctions could potentially be evaded, allowing intermediate and poor fixers to prosper.

Even in legumes that form indeterminate nodules, there are two clear routes by which cheating might persist in the face of plant sanctions. First, natural selection only concerns itself with short-term fitness; therefore, sanctions targeted only at bacteroids (which are the major consumers of plant-derived carbon) might be selected for, even if they did not affect the undifferentiated bacteria within the nodule. If those undifferentiated bacteria were then released back into the soil in large numbers, this would enhance the populations of

cheating strains and so undermine the long-term stability of the symbiosis. Second, poor

fixers may persist by 'piggybacking' on a more effective strain, which can occur because

multiple strains sometimes occupy a single nodule (Mendoza-Suárez et al., 2021). This route

could be avoided if the plant were able to distinguish between different strains within a

nodule and sanction the less effective one, but despite some evidence for so-called cell

autonomous sanctioning, in the form of premature senescence of cells containing an

ineffective strain (Daubech et al., 2017; Regus et al., 2014), metabolic evidence is missing

(Agtuca et al., 2020), so it remains contentious.

To determine whether either route would allow the persistence of poor fixers, we built on

previous work carried out by Westhoek et al. (2021) using pea plants and a set of otherwise

isogenic rhizobia strains that differ in their ability to fix nitrogen. Specifically, we used (1)

flow cytometry to establish whether sanctioned nodules occupied by a less effective strain

contain fewer undifferentiated bacteria; and (2) fluorescence microscopy to establish how

sanctions influence the timing of nodule senescence. Finally, we used the same techniques

on mixed nodules to determine whether or not pea plants carry out cell-autonomous

sanctioning within nodules. Our results show that plant sanctions are effective at reducing

the fitness of poorly fixing strains. These sanctions operate at the level of whole nodules but

not at the cellular level.

<u>Methods</u>

We co-inoculated plants with two rhizobial strains selected from three otherwise isogenic

strains differing only in their ability to fix nitrogen. They were characterised as a good fixer

(Fix$^+$), an intermediate fixer (Fix^{int}) and a non-fixer (Fix$^-$, see *Bacterial strains and culture*

conditions for more details) This gives three pairwise combinations of effective (E) and

ineffective (I) strains; Fix$^+$ (E) vs Fix$^-$ (I), Fix$^+$ (E) vs Fix^{int} (I) and Fix^{int} (E) vs Fix$^-$ (I). i.e., Fix$^+$ is

always effective and Fix$^-$ always ineffective, but Fix^{int} can be either effective or ineffective,

depending on the co-inoculated strain. To measure conditional sanctioning at the whole-

nodule level, we sampled nodules occupied by single strains only (identified by a fluorescent

marker), and passed them through a flow cytometer to quantify the bacteroid and

undifferentiated bacterial populations. Sections of nodules were also examined using

confocal microscopy to assess cell health and morphology at a range of time points. Both of

these techniques allowed us to assess the effectiveness of plant sanctions on bacterial
populations.

Mixed nodules also form at low frequency on plants co-inoculated with two or more strains.
To test the theory of cell-autonomous sanctioning, bacteroids and bacteria from mixed
nodules were isolated, quantified by flow cytometry, and imaged by confocal microscopy.
We set up three to four plants of each of our three strain combinations (Fix$^+$ & Fix$^-$, Fix$^+$ &
Fix^{int} and Fix^{int} & Fix$^-$) and passed between two and six nodules from each plant through the
flow cytometer. Further nodules were also picked for confocal microscopy to inspect the
health of different cell types. These techniques allow us to see whether the plant can
discriminate between strains within a single nodule and apply sanctions accordingly.

Bacterial Strains and Culture Conditions

Rhizobial strains used in this study are all derivatives of a highly effective nitrogen-fixing
strain (Rlv3841) of *Rhizobium leguminosarum* bv. *viciae* (Johnston & Beringer, 1975), that
infects pea (*Pisum sativum* L. cv. Avola). We used mutant strains that differ in their nitrogen-
fixation ability but are otherwise genetically identical (Table 1). They are labelled with a
fluorescent marker (mCherry or GFP) in order to distinguish the nodules formed by each
strain (Table 1). Strains were maintained on tryptone-yeast (TY) agar (Beringer, 1974) with
the appropriate concentrations of antibiotics (Table 1). For long-term storage, strains were
kept at −80 °C in TY with 15 to 20% glycerol. Rhizobial inoculant was grown on TY agar
before dilution to 1×10^8 cells ml^{-1}.

Plant Growth

Before sowing, all pea seeds were surface sterilized (1 minute in 95% ethanol followed by 5
minutes in 2% NaClO), rinsed, and left to germinate on 1% w/v agar plates at room
temperature in the dark. Seedlings were transplanted after five days by transferring them to
sterilized 1 litre Azlon beakers containing a 1:1 mixture of silver sand and fine vermiculite,
150ml sterilized nitrogen-free nutrient solution, and 5×10^7 cells of each of the two rhizobial
strains (see also Westhoek et al., (2017)). Beakers were covered with clingfilm to reduce

aerial contamination, which was slit after a few days to allow seedlings to grow through.

Plants were grown in a growth room (21 °C, 16 h photoperiod) for 28 to 42 days and

watered as necessary from 7 days onwards.

Table 1: Rhizobial strains and their fixation abilities. All strains derived from Rlv3841 and provided with a strain code, resistance markers, short description and reference.

Name	Strain	Antibiotic resistance	Description	Reference
Fix$^+$ mCherry	OPS1341	Streptomycin, Gentamicin	Rlv3841 Tn7-Gm-mCherry False colour: orange	(Westhoek et al., 2021)
Fix$^-$ GFP	OPS2270	Streptomycin, Spectinomycin, Gentamicin	Ω$nifH$ Tn7-Gm-sfGFP (Rlv3841 mutant, ΩSpc cassette in $nifH$)	(Westhoek et al., 2021)
Fix^{int} GFP	OPS2268	Streptomycin, Spectinomycin, Gentamicin	Fix^{int} Tn7-Gm-sfGFP (Rlv3841 mutant, ΩSpc cassette in promoter region of $nifA$)	(Westhoek et al., 2021)
Fix^{int} mCherry	OPS2269	Streptomycin, Spectinomycin, Gentamicin	Fix^{int} Tn7-Gm-mCherry False colour: orange	(Westhoek et al., 2021)

Harvesting

Plants were harvested at 28-, 35- and 42-days post-inoculation by removing the plant from

the sand/vermiculite mixture and washing the roots carefully. Nodules were then imaged

using a LEICA M165 FC fluorescent stereo microscope and an iBright FL1500 imaging system.

Nodule occupants were identified based on their fluorescence (red for Fix$^+$, green for Fix$^-$,

green or red for Fix^{int}). When selecting single occupancy nodules, the largest of each nodule

type was selected.

Confocal imaging

To assess the health of cells containing undifferentiated bacteria and bacteroids within

nodules, we imaged sections using confocal microscopy, which allows the visualisation of

fluorescently tagged strains and discrimination of the two strains within a mixed nodule. We

placed selected nodules in 8 % w/v agar and cut 100 μm sections using a Leica VT1200S

vibratome. Nodule slices were then imaged with a Zeiss LSM 880 Airy Scan confocal

microscope and analysed with ZEN Black software. To visualise fluorescent tags, mCherry

was excited using a 561 nm wavelength laser and emissions detected between 598 and 649

nm, while GFP was excited using a 488 nm wavelength laser and emissions detected

between 498 and 562 nm.

Flow Cytometry

To identify and quantify the population sizes of undifferentiated bacteria and bacteroids

within nodules, we carried out flow cytometry. This was done initially with selected single

occupant nodules from the three co-inoculant combinations (Fix$^+$ vs Fix$^-$, Fix$^+$ vs Fix^{int} and

Fix^{int} vs Fix$^-$). We then carried out flow cytometry on the three types of mixed nodules (Fix$^+$

& Fix$^-$, Fix$^+$ & Fix^{int} and Fix^{int} & Fix$^-$). All nodules were prepared by placing them in a 1.5ml

Eppendorf tube with 300μl of harvest solution (0.9% NaCl, 0.02% SILWET L-77) and crushing

them with an autoclaved microcentrifuge pestle. This solution was passed through a 40μm

filter and diluted tenfold to increase the accuracy of the flow cytometry. Flow cytometry

was conducted with a Amnis® CellStream® flow cytometer (Luminex) equipped with a 488

and 561nm lasers, which were used for excitation of GFP and mCherry, respectively. Flow

rates were set to low speed and high sensitivity (3.66 μL/min). Bacteria events were defined

based on custom gating parameters. Singlets and doublets were gated with a threshold of

0.400 forward scatter (FSC) aspect ratio. Bacterial singlets with emission at 611/631nm,

when excited at 561 nm, above 6,000 arbitrary units were defined as red. Bacterial singlets

with emission at 528/546nm, when excited at 488nm, above 4,000 arbitrary units were

defined as green. Bacteroids were defined as those events with FSC values of over 10,000

arbitrary units, while undifferentiated bacteria were defined as events with values below

10,000. Flow cytometry data available at http://flowrepository.org with experiment codes

FR-FCM-Z6XN and FR-FCM-Z6XX for the single occupancy experiments and the mixed

nodule, experiments respectively.

Statistical Analysis

We first analysed population sizes for undifferentiated bacteria and bacteroids within

single-strain nodules (Table S1). In total, we had bacterial counts from sixty-nine single-

strain nodules collected from nine plants with different pairwise combinations of strains,

which we treated as independent data points. We compared population sizes in effective vs

ineffective nodules, based on the idea that in each pairwise combination we can identify an

effective and an ineffective strain based on known fixation rates from Westhoek et al.,

(2021). Specifically, Fix^+ nodules can always be classed as the effective partner, as can Fix^{int}

nodules when co-inoculated with a Fix^- strain. In contrast, Fix^- nodules can always be classed

as the ineffective partner, as can Fix^{int} nodules when co-inoculated with a Fix^+ strain. To test

for conditional sanctioning at the whole-nodule level, we binned the data into the three

different nodule types: Fix^+, Fix^- and Fix^{int}, and tested whether population sizes were

dependent on the identity of the co-inoculated strain.

To test for the presence of cell autonomous sanctioning within mixed nodules, we used a

one-sample t-test to determine whether the proportion of the two strains differed

significantly from 50% for both undifferentiated bacteria and bacteroid populations. To

compare the absolute numbers of undifferentiated bacteria and bacteroids within mixed

nodules, we used paired t-tests. To test whether the total number of undifferentiated

bacteria and bacteroids in a mixed nodule differs from that in their respective single-strain

nodules, we directly compared all three nodule types using one-way ANOVA followed by a t-

test comparing effective and ineffective nodule data with mixed nodule data.

All statistical tests were carried out using R version 4.2.1 (2022-06-23) and R studio

2022.07.2 and graphs were produced using Graph Pad version 9. An R markdown document

is provided (Doc. S2). In figures, data is presented log10 transformed. However, statistical
analysis was performed on the untransformed data to avoid back-transformations, which
can distort estimates (Lee, 2020). The analysis was repeated on the log10 transformed data
and is available in Doc.S2. There were no qualitative differences in the outcome of
transformed vs untransformed data.

Results

Sanctioning of less effective nodules

As found by Westhoek et al., (2017 & 2021), we observed, on all plants, clear phenotypic
differences between nodules containing an effective strain (i.e., either Fix$^+$ nodules (vs Fix$^-$
(I) or vs Fix^{int} (I)) or Fix^{int} nodules (vs Fix$^-$ (I)) and those containing an ineffective strain (i.e.,
either Fix$^-$ nodules (vs Fix$^+$ (E) or vs Fix^{int} (E)) or Fix^{int} nodules (vs Fix$^+$ (E)) at 28 dpi. Notably,
nodules containing the effective strain were larger and more tubular, while nodules
containing the ineffective strain were smaller and more spherical (Figure S2). These
differences were observed, regardless of whether the ineffective strain was the non-fixer
(Fix$^-$) or the intermediate fixer (Fix^{int}). In tandem, there was a significant reduction in the
number of undifferentiated bacteria in nodules containing the ineffective strain (mean
difference = -119622, SE = 23424, t = -5.085, p = 3.17×10^{-6}, n = 69) (Figure 1A) and a
reduction in the number of bacteroids (mean difference = -602492, SE = 127765, t = -4.716,
p = 1.26×10^{-5}, n = 69) (Figure 1B).

Conditional sanctioning on whole nodules determines bacterial survival

If plants impose conditional sanctions, then we expect the number of undifferentiated
bacteria and bacteroids within nodules to depend on the identity of the co-inoculated strain
for the intermediate fixer only. In support of this, the number of bacteroids does not
depend on the identity of the co-inoculated strain for either Fix$^+$ nodules (mean difference =
97172, SE = 338503, t = 0.284, p = 0.779, n = 24) or Fix$^-$ nodules (mean difference = -14164,
SE = 43638 t = -0.325, p = 0.750, n = 16). However, nodules occupied by the intermediate-

fixing strain contained significantly fewer bacteroids when the co-inoculated strain was Fix$^+$

rather than Fix$^-$ (mean difference = -143208, SE = 53448, t = -2.679, p = 0.0124, n = 29)

(Figure 1C). We found the same pattern for undifferentiated bacteria. Their numbers did not

depend on the identity of the co-inoculated strain for either Fix$^+$ nodules (mean difference =

18176, SE = 57560, t = 0.316, p = 0.755, n = 24), or Fix$^-$ nodules (mean difference = -13185,

SE = 31632, t = -0.417, p = 0.683, n = 16). However, for the intermediate fixing nodules,

there were significantly fewer undifferentiated bacteria when the co-inoculated strain was

Fix$^+$ rather than Fix$^-$ (mean difference = -68824, SE = 19082, t = -3.607, p = 0.00124, n = 29)

(Figure 1D). This demonstrates that conditional sanctioning affects the undifferentiated

bacteria (and hence bacterial fitness) and not just the terminally differentiated bacteroids.

However, the absolute and relative differences between Fix$^+$, Fix^{int} and Fix$^-$ are greater for

bacteroids than for undifferentiated bacteria, implying that sanctioning has more impact on

bacteroid populations at 28 days (compare Figures 1C&D).

Conditional sanctioning precipitates premature nodule senescence

At 28 days post-inoculation, there was no visible difference in the nodule sections between

those containing the Fix$^-$ strain (Figure 2A) and those containing the Fix$^+$ strain (Figure 2B).

However, at 35 days post-inoculation, there was a clear difference between these nodule

types. Nodules containing the Fix$^-$ strain had begun to senesce and the cells within nodules

were undergoing cell death (Figure 2C), while in contrast, the nodules containing the Fix$^+$

strain showed no evidence of senescence and looked indistinguishable from Fix$^+$ nodules

after 28 days (compare Figures 2B&2D). At 42 days post-inoculation, nodules containing the

Fix$^+$ strain had also begun to senesce (Figure 2F), while cells within nodules containing the

Fix$^-$ strains were fully senesced (Figure 2E).

Similar to the Fix$^+$ strain, after 28 days post-inoculation, sections from nodules containing

the Fix^{int} strain looked the same, regardless of whether the co-inoculated strain was Fix$^+$

(Figure 3A) or Fix$^-$ (Figure 3B). However, at 35 days post-inoculation, the status of nodules

containing the Fix^{int} strain was visibly affected by the identity of the co-inoculated strain,

consistent with conditional sanctioning. In particular, if the co-inoculated strain was Fix$^+$,

then senescence had begun and cell death was visible (Figure 3C), whereas, if the co-

inoculated strain was Fix⁻, then there was no sign of senescence, and nodules were not

visibly different from Fix^{int} nodules at 28 days post-inoculation (compare Figures 3B&3D).

After 42 days, all nodules containing a Fix^{int} strain had begun to senesce. In those where the

co-inoculated strain was Fix⁺ (Figure 3E), senescence had progressed to a more advanced

state than in those co-inoculated with a Fix⁻ strain (Figure 3F).

Example nodules from co-inoculated plants of all three combinations (Fix⁺ vs Fix⁻, Fix⁺ vs

Fix^{int}, Fix^{int} vs Fix⁻) were imaged at 28-, 35- and 42-days post-inoculation and examined for

evidence of senescence and changes to cell morphology (Figure S3).

Conditional sanctioning is not cell-autonomous in peas

In mixed nodules, the proportion of bacteroids that belonged to the less effective strain was

not significantly different from 50% (mean = 49.78%, 95%CI = 41.80 & 57.76%, t = -0.058; p

>0.05, n = 18) (Figure 4A); hence, we did not find evidence for cell-autonomous sanctioning.

Indeed, contrary to the predictions of cell-autonomous sanctioning, the proportion of

undifferentiated bacteria that belonged to the effective strain was significantly less than

50% (mean = 37.78%, 95%CI = 47.6 & 27.94%, t = -2.621, p = 0.0178, n = 18) (Figure 4A),

indicating that the less effective strain could even have an advantage within mixed nodules.

The same pattern emerged when we compared absolute numbers. The number of

bacteroids belonging to the effective strain within a mixed nodule was not significantly

different from the number belonging to the ineffective strain (mean difference = -10312,

95%CI = 172698 & 193321, t = -0.119, p >0.05, n = 18). However, the number of

undifferentiated bacteria belonging to the effective strain was significantly lower than that

belonging to the ineffective strain (mean difference = -616752, 95%CI = -156712 & -

1076790, t = -2.8285, p = 0.0116, n = 18).

Mixed nodules are treated as ineffective nodules by the plant

There were significant differences in the total numbers of bacteroids found in nodules of

different types (mixed vs single-strain effective vs single-strain ineffective; $F_{2,84}$ = 15.89, p =

1.40×10^{-6}; Figure 1A & B) and the same was true for the total number of undifferentiated

bacteria ($F_{2,84}$ = 11.62, p = 3.50×10^{-5}). There were significantly more bacteroids in effective

nodules compared with mixed nodules (mean difference = -89474, SE = 140734, t = 4.189, p

= 6.89×10^{-5}; Figure 4B), but no significant difference between mixed and ineffective nodules

(mean difference = -13018, SE = 141435, t = -0.092, p >0.5; Figure 4B). The same pattern

was seen for the undifferentiated bacterial populations. There were more undifferentiated

bacteria in effective nodules compared with mixed nodules (mean difference = 73709, SE =

30076, t = 2.451, p = 0.0163; Figure 4C), but no significant difference between mixed and

ineffective nodules (mean difference = -45913, SE = 30226, t = -1.519, p > 0.05; Figure 4C).

This would suggest that mixed nodules are treated as ineffective nodules and sanctioned

accordingly.

Timing of senescence is not cell-autonomous in pea

Within mixed nodules at 28 days post-inoculation (n = 13 nodules), there was no evidence

of any difference in cell morphology or health between the two strains, as all cells appeared

healthy (Figure 5A:C). However, at 35 days post-inoculation, we observed that a few cells

within some mixed nodules appear to be undergoing apoptosis (Figure 5F). However, cells

belonging to both strains (whether the more or less effective) appeared to be undergoing

this process with equal probability (Figure 5). As such, within nodules at 35 days post-

inoculation (n = 7 nodules), there was no evidence of cell-autonomous sanctioning (Figure

5D:F). At 42 days post-inoculation, all nodules (n = 9 nodules) showed evidence of apoptosis

and programmed cell death with very few intact cells remaining (Figure 5G:I).

Discussion

We show by quantification of undifferentiated bacteria (Figure 1A) and bacteroids (Figure

1B) that peas sanction nodules containing ineffectively fixing strains, which is consistent

with previous studies (Kiers et al., 2003; Oono et al., 2011; West et al., 2002; Westhoek et

al., 2017). It is also consistent with conditional sanctioning, as the fate of nodules infected

by a Fix[int] strain depends on whether the co-inoculated strain is Fix[+] or Fix[-] (Westhoek et al.,

2021). Sanctions affect both bacteroids (Figure 1C) and undifferentiated bacteria (Figure 1D)

within nodules, although for undifferentiated bacteria the magnitude of effects was smaller than for bacteroids. This is consistent with the findings of Westhoek et al., (2021) that the number of viable bacteria has not begun to drop significantly in sanctioned nodules prior to 52 days post inoculation.

We also show that the timing of nodule senescence is influenced by sanctioning but they are separate events and do not occur simultaneously (Figure 2 & 3). While sanctioning is discernible after 16 days (Westhoek et al., 2017) and very marked by 28 days post-inoculation (Figure 1), there is no evidence of cellular senescence in nodules infected by any strain at either time point. It was only after 35 days that nodules containing an ineffective strain began to show signs of senescence, while those containing an effective strain only senesced after 42 days (Figure 5). Thus, while the timing of cellular senescence is influenced by the relative effectiveness of the strain within a nodule, sanctioning at the whole nodule level occurs well before cells start to break down.

It has been proposed that cell-autonomous sanctions might occur within mixed nodules, allowing plants to combat ineffective "cheating" strains that enter nodules alongside a more effective strain. Evidence for such sanctions would include a bias in the proportion of the effective strain within mixed nodules. However, in our study, the ratio of effective to ineffective bacteroids within mixed nodules did not significantly differ from 50% (Figure 4A). The ratio of effective to ineffective undifferentiated bacteria did significantly differ from 50%, but this is because the number of ineffective bacteria was actually higher – the opposite of what we would expect if cell-autonomous sanctions operated. These results therefore show that peas are incapable of cell-autonomous sanctions. When comparing total mixed nodule populations of bacteroids (Figure 4B) and undifferentiated bacteria (Figure 4C) with both single occupancy effective and ineffective nodules, we see that the mixed nodule populations are significantly lower than the effective nodules and not significantly different from those of ineffective nodules. This demonstrates that mixed nodules are treated as ineffective nodules by pea plants.

Cells within mixed nodules underwent senescence simultaneously, regardless of variation in effectiveness (Figure 5). Consistent with the timing of cell senescence not occurring in a cell-

autonomous manner. Although the timing of senescence varied among mixed nodules,
where senescence occurred, it affected both strains equally. It may therefore be
hypothesised that the timing of senescence within mixed nodules is determined by the ratio
of effective to ineffective strains within the nodule. This would be challenging to test,
because sectioning nodules does not give an accurate measure of strain ratio due to the
irregular 3D distribution of strains within nodules.

Our results align with the findings of Agtuca et al., (2020) who found that the metabolic
profile of sectors of mixed nodules containing different strains did not show significant
differences. However, our results contrast with studies which have found evidence for cell
autonomous sanctions based on variation in the timing of senescence, such as Regus et al.,
(2017) and Daubech et al., (2017). The differences may be explained by variation in the
legume-rhizobium system used: Daubech et al. (2017) used *Cupriavidus* taiwanensis –
Mimosa pudica; the Regus et al. (2017) study used both *Acmispon strigosus* –
Bradyrhizobium and *Lotus japonicus – Mesorhizobium*; and the Agtuca et al. (2020) study
used *Glycine max – Bradyrhizobium japonicum*. A comprehensive understanding of
sanctions will therefore require the continued use of multiple experimental systems.

We have shown that the sanctioning of ineffective nodules within the pea-rhizobium
symbioses is conditional in nature and occurs at the nodule level. In Fix^+/Fix^{int} coinfected
plants, nodules containing Fix^{int} but not Fix^+ strains are sanctioned, suggesting conditional
sanctioning requires comparison between nodule-specific and global nitrogen signals. This
might be achieved through the interaction between a nitrate receptor such as the Nitrate
transreceptor NRT1.1, which both transports nitrate as well as detecting nitrate levels
(Zhang et al., 2019) and the NIN-like Proteins (NLPs) 1 and 4, which are essential for the
nitrate-based regulation of nodule maturation (Lin et al., 2021). In high nitrate conditions
NLP1/4 inhibit cytokinin biosynthesis. Cytokinin biosynthesis drives nodule maturation
through the activation of a signal cascade through the Cytokinin responsive element CRE1
(Lin et al., 2021), which acts to promote expression of the *cep* and *cle* genes (Laffont et al.,
2020). However, how a global signalling mechanism is able to detect the nitrogen output of
individual nodules and compare it to global nitrogen status remains unclear. The
mechanism of sanctions on individual nodules may be achieved through nodule-specific

proteins responding to nitrate levels. One possibility is SnRK1 which, when phosphorylated by the DMI2 kinase in response to Nod factor, phosphorylates malate dehydrogenase 1 and 2, leading to increased malate production within the nodule to supply bacteroids (Guo et al., 2023). This is therefore a prime candidate for a pathway in which the legume would be able to reward effective strain-containing nodules, as well as to punish nodules containing ineffectively fixing strains by cessation of malate production and supply. However, these remain speculations about signalling pathways whose elucidation may aid future efforts to engineer symbioses in the selection and use of more effective nitrogen-fixing bacteria.

Having identified two potential routes for persistence of a poor nitrogen-fixing and/or 'cheating' strain, we now have clear evidence that plants can deal with both. First, while sanctions, at least initially, affect nitrogen-fixing bacteroids more severely, undifferentiated bacteria are also greatly reduced in number inside ineffective nodules. Second, while the plant is unable to discriminate between effective and ineffective strains within a mixed nodule, 'piggybacking' on an effective strain does not provide a viable route by which to evade sanctions. Any potential benefit is limited not only by the scarcity of mixed nodules, but also because the plant views mixed nodules as ineffective, and sanctions them accordingly.

449 **References**
450

Agtuca, B. J., Stopka, S. A., Evans, S., Samarah, L., Liu, Y., Xu, D., Stacey, M. G., Koppenaal, D.
W., Paša-Tolić, L., Anderton, C. R., Vertes, A., & Stacey, G. (2020). Metabolomic
profiling of wild-type and mutant soybean root nodules using laser-ablation
electrospray ionization mass spectrometry reveals altered metabolism. *Plant Journal*,
103(5), 1937–1958. https://doi.org/10.1111/tpj.14815
Beringer, J. E. (1974). R Factor Transfer in Rhizobium leguminosarum. *Journal of General*
Microbiology, *84*(1g74), 188–198.
Daubech, B., Remigi, P., Doin de Moura, G., Marchetti, M., cile Pouzet, C., Auriac, M.-C.,
Gokhale, C. S., Masson-Boivin, C., & Capela, D. (2017). Spatio-temporal control of
mutualism in legumes helps spread symbiotic nitrogen fixation. *ELIFE*, 1–21.
https://doi.org/10.7554/eLife.28683.001
Guo, D., Liu, P., Liu, Q., Zheng, L., Liu, S., Shen, C., Liu, L., Fan, S., Li, N., Dong, J., & Wang, T.
(2023). Legume-specific SnRK1 promotes malate supply to bacteroids for symbiotic
nitrogen fixation. *Molecular Plant*, *16*(9), 1396–1412.
https://doi.org/10.1016/j.molp.2023.08.009
Johnston, A. W. B., & Beringer, J. E. (1975). Identification of the Rhizobium Strains in Pea
Root Nodules Using Genetic Markers. *Journal of General Microbiology*, *87*, 343–350.
Kiers, E. T., Rousseau, R. A., West, S. A., & Denlson, R. F. (2003). Host sanctions and the
legume-rhizobium mutualism. *Nature*, *425*(6953), 78–81.
https://doi.org/10.1038/nature01931
Laffont, C., Ivanovici, A., Gautrat, P., Brault, M., Djordjevic, M. A., & Frugier, F. (2020). The
NIN transcription factor coordinates CEP and CLE signaling peptides that regulate
nodulation antagonistically. *Nature Communications*, *11*(1).
https://doi.org/10.1038/s41467-020-16968-1
Lee, D. K. (2020). Data transformation: A focus on the interpretation. *Korean Journal of*
Anesthesiology, *73*(6), 503–508. https://doi.org/10.4097/kja.20137
Lin, J., Roswanjaya, Y. P., Kohlen, W., Stougaard, J., & Reid, D. (2021). Nitrate restricts
nodule organogenesis through inhibition of cytokinin biosynthesis in Lotus japonicus.
Nature Communications, *12*(1). https://doi.org/10.1038/s41467-021-26820-9
Mendoza-Suárez, M., Andersen, S. U., Poole, P. S., & Sánchez-Cañizares, C. (2021).
Competition, Nodule Occupancy, and Persistence of Inoculant Strains: Key Factors in
the Rhizobium-Legume Symbioses. In *Frontiers in Plant Science* (Vol. 12). Frontiers
Media S.A. https://doi.org/10.3389/fpls.2021.690567
Mergaert, P., Uchiumi, T., Alunni, B., Naë Lle Evanno, G., Lique Cheron, A., Catrice, O.,
Mausset, A.-E., Dé Rique Barloy-Hubler, F., Galibert, F., Kondorosi, A., & Kondorosi, E.
(2006). *Eukaryotic control on bacterial cell cycle and differentiation in the Rhizobium-*
legume symbiosis. https://www.pnas.org
Oono, R., Anderson, C. G., & Denison, R. F. (2011). Failure to fix nitrogen by non-
reproductive symbiotic rhizobia triggers host sanctions that reduce fitness of their
reproductive clonemates. *Proceedings of the Royal Society B: Biological Sciences*,
278(1718), 2698–2703. https://doi.org/10.1098/rspb.2010.2193

Poole, P., Ramachandran, V., & Terpolilli, J. (2018). Rhizobia: From saprophytes to
endosymbionts. *Nature Reviews Microbiology*, *16*(5), 291–303.
https://doi.org/10.1038/nrmicro.2017.171
Regus, J. U., Gano, K. A., Hollowell, A. C., & Sachs, J. L. (2014). Efficiency of partner choice
and sanctions in Lotus is not altered by nitrogen fertilization. *Proceedings of the Royal*
Society B: Biological Sciences, *281*(1781). https://doi.org/10.1098/rspb.2013.2587
Timmers, A. C. J., Soupène, E., Auriac, M.-C., De Billy, F., Vasse, J., Boistard, P., & Truchet, G.
(2000). Publication no. M-2000-0914-03R.. 1996; Long 1996) and takes place through a
series of developmental stages. In *Schultze and Kondorosi* (Vol. 13, Issue 11).
Udvardi, M., & Poole, P. S. (2013). Transport and metabolism in legume-rhizobia symbioses.
In *Annual Review of Plant Biology* (Vol. 64, pp. 781–805).
https://doi.org/10.1146/annurev-arplant-050312-120235
West, S. A., Kiers, E. T., Simms, E. L., & Denison, R. F. (2002). Sanctions and mutualism
stability: Why do rhizobia fix nitrogen? *Proceedings of the Royal Society B: Biological*
Sciences, *269*(1492), 685–694. https://doi.org/10.1098/rspb.2001.1878
Westhoek, A., Clark, L. J., Culbert, M., Dalchau, N., Griffiths, M., Jorrin, B., Karunakaran, R.,
Ledermann, R., Tkacz, A., Webb, I., James, E. K., Poole, P. S., & Turnbull, L. A. (2021).
Conditional sanctioning in a legume-Rhizobium mutualism. *Proceedings of the National*
Academy of Sciences of the United States of America, *118*(19), 1–8.
https://doi.org/10.1073/pnas.2025760118
Westhoek, A., Field, E., Rehling, F., Mulley, G., Webb, I., Poole, P. S., & Turnbull, L. A. (2017).
Policing the legume-Rhizobium symbiosis: A critical test of partner choice. *Scientific*
Reports, *7*(1). https://doi.org/10.1038/s41598-017-01634-2
Zhang, X., Cui, Y., Yu, M., Su, B., Gong, W., Baluška, F., Komis, G., Šamaj, J., Shan, X., & Lin, J.
(2019). Phosphorylation-mediated dynamics of nitrate transceptor NRT1.1 regulate
auxin flux and nitrate signaling in lateral root growth. *Plant Physiology*, *181*(2), 480–
498. https://doi.org/10.1104/pp.19.00346

519
520

Acknowledgments

522 This work was supported by the Biotechnology and Biological Sciences Research Council
523 [BB/M011224/1, BB/T001801/1, BB/T006722/1, BB/T008784/1 and BB/W006219/1].

Competing interests

526 The authors declare no competing interests

Author contributions

529 All authors listed have made a substantial direct and intellectual contribution to the work
530 and approved it for publication. TJU, BJ and PSP conceived this study and designed the
531 experiments. TJU and BJ performed the experiments. TJU analysed the data and prepared
532 the manuscript. TJU, BJ, LAT and PSP critically reviewed the manuscript.

Data Availability

Flow cytometry: Data is available at http://flowrepository.org with experiment codes FR-
FCM-Z6XN and FR-FCM-Z6XX for the single occupancy experiments and the mixed nodule
experiments respectively.

<u>Supporting information</u>

The following Supporting Information is available for this article:

Table. S1 Data sheet containing counts from flow cytometer of undifferentiated bacteria

and bacteroids from nodules (single occupancy and mixed nodules).

Doc.S2 R markdown file of statistical analysis.

Fig. S3 Representative iBright image of pea inoculated with two strains of differing

effectiveness.

Fig. S4 Representative confocal images of nodule sections of both nodule types from all

three co-inoculation combinations (Fix^+ vs Fix^-, Fix^+ vs Fix^{int}, Fix^{int} vs Fix^-) at 28-, 35- and 42-

days post inoculation.

Fig. 1. Nodules containing the ineffective strain have fewer bacteria and bacteroids.

Numbers of (A) undifferentiated bacteria and (B) bacteroids in nodules from pea plants co-inoculated with two strains of rhizobia, one effective (either Fix$^+$ or Fix^{int} co-inoculated with Fix$^-$) and the other ineffective (either Fix$^-$ or Fix^{int} co-inoculated with Fix$^+$). (C) and (D) show the same data broken down into different nodule types (Fix$^+$, Fix^{int} or Fix$^-$). The identity of the co-inoculated strain is also shown. In each case, one strain is denoted effective (E) and the other ineffective (I) as determined by their relative ability to fix nitrogen. Significance levels from unpaired t-tests: ****< 0.0001; *** < 0.001; ** < 0.01; * <0 .05; ns, not significant. Nodules harvested at 28 days post-inoculation (*n* = 69).

Fig. 2. Nodules containing a Fix⁻ strain senesce earlier then nodules containing a Fix⁺ strain.
(A), (C) and (E) Fix⁻ nodule labelled with GFP (blue) and (B), (D) and (F) Fix⁺ nodule labelled with mCherry (orange) at 28 dpi, 35 dpi and 42 dpi, respectively. Longitudinal nodule slices (100μm) were imaged by confocal microscopy. Scale bar = 200μm.

Fix^int Nodule
Fix^int (I) vs Fix^+(E)

Fix^int Nodule
Fix^int (E) vs Fix^- (I)

28dpi

35dpi

42dpi

Fig. 3. Fix^int nodules co-inoculated with Fix^+ strain senesce earlier than those co-inoculated with Fix^- strain
(A), (C) and (E) Fix^int nodule labelled with GFP (blue), co-inoculated with Fix^+. (B), (D) and (F) Fix^int nodule labelled with mCherry (orange), co-inoculated with Fix^-. Nodules were harvested at 28 dpi, 35 dpi and 42 dpi, respectively. Longitudinal nodule slices (100μm) were imaged by confocal microscopy. Scale bar = 200μm.

Fig. 4. The percentage of bacteria and bacteroids of the effective strain within a mixed nodule do not exceed 50%, and the combined population size is lower than that of an effective nodule.

(A) Percentage of bacteroids and undifferentiated bacteria within a mixed nodule that belonged to the effective strain (either Fix[+] or Fix[int] co-inoculated with Fix[-]). The total number of (B) bacteria and (C) bacteroids within single-strain effective nodules, mixed nodules (both populations combined) and single-strain ineffective nodules. ns, not significant. Significance levels in (B) and (C) assessed from pairwise comparisons using t-tests: ****< 0.0001 *** < 0.001, ** < 0.01, * <0 .05. Nodules harvested at 28 dpi.

| | 28 dpi | 35 dpi | 42 dpi |

Fig. 5. The two strains within a mixed nodule do not senesce at different times despite varying in relative effectiveness.

(A), (D) and (G) mixed nodule of Fix$^+$ (orange) and Fix$^-$ (blue). (B), (E) and (H) mixed nodule of Fix$^+$ (orange) and Fix^{int} (Blue). (C), (F) and (I) mixed nodule of Fix^{int} (orange) and Fix$^-$ (blue). Nodules were harvested at 28 dpi, 35 dpi and 42 dpi, as indicated. Longitudinal nodule slices (100μm) were imaged by confocal microscopy. Scale bar = 200μm.