Pisum sativum applies sanctions to less effective strains at the nodule level,

but not within mixed nodules

Thomas J. Underwood[1], Beatriz Jorrin[1], Lindsay A. Turnbull[1], Philip S. Poole[1]

[1]Department of Biology, University of Oxford, Oxford, United Kingdom

Author for correspondence:

Philip Poole

Email: philip.poole@biology.ox.ac.uk

Word count: 5247

Number of Figures: 5

Figures to be in colour: 1, 2 & 5

Tables:

1 table providing a list and description of strains used

Supporting information

The following Supporting Information is available for this article:

Fig. S1 Image of flow cytometry output showing the two populations differentiated by size
based on Forward Scatter (FSC)

Table. S2 Flow cytometry data, analysed using custom gating to calculate the number of

undifferentiated bacteria and bacteroids in single occupancy and mixed nodules, and

acetylene reduction assay and nodulation results from fluorescent tag control experiments.

Doc.S3 R markdown file of statistical analysis.

Fig. S4 Representative iBright image of pea inoculated with two strains of differing

effectiveness.

Fig. S5 Representative confocal images of nodule sections of both nodule types from all

three co-inoculation combinations (Fix^+ vs Fix^-, Fix^+ vs Fix^{int}, Fix^{int} vs Fix^-) at 28-, 35- and 42-

days post inoculation.

Fig. S6 Representative images of the three types of mixed nodules at 28, 35 and 42 days

post inoculation

<u>**Summary**</u>

• Legumes sanction root nodules containing rhizobial strains with low nitrogen fixation

rates (less effective fixers). Pea (*Pisum sativum*) nodules contain both

undifferentiated bacteria and terminally differentiated nitrogen-fixing bacteroids. It

is critical to understand how sanctions act on both bacteria and bacteroids and how

they differ. In addition, less effective strains could potentially evade sanctioning by

entering the same nodule as an effective strain i.e., piggybacking. *P. sativum* was co-

inoculated with pairwise combinations of three strains of rhizobia with different

effectiveness, to test whether ineffective strains can evade sanctions in this way.

• We assessed the effect of sanctions on nodule populations of bacteria and

bacteroids using flow cytometry and the effects on nodule ultra-structure using

confocal microscopy.

• We show that sanctioning lowered the bacteroid population and caused a reduction

in the size of bacteria. Sanctions also precipitated an early change in nodule cell

morphology. In nodules containing two strains that differed in their nitrogen-

fixation ability, both were sanctioned equally. Thus, peas sanction whole nodules

based on their nitrogen output, but do not sanction at the cellular level.

• Our results demonstrate that whole-nodule conditional sanctioning in peas provides

long-term stability to the symbiosis by effectively reducing the fitness of ineffective

strains.

Introduction

Legumes have overcome nitrogen limitation by establishing a mutually beneficial interaction

with nitrogen-fixing bacteria, hosted within nodules along their root systems (Poole et al.,

2018). Certain species of legumes, like peas, form nodules in which some bacteria undergo

terminal differentiation to become large swollen cells called bacteroids, which carry out the

costly process of nitrogen fixation, but cannot resume their free-living existence (Mergaert

et al., 2006). The plant provides carbon compounds in the form of photosynthetically-

derived dicarboxylates in return for ammonia provided by the bacteroids, and this

relationship continues until the plant's nitrogen requirements have been met (Udvardi &

Poole, 2013). At this point nodule senescence ensues, and while the bacteroids then die,

any undifferentiated bacteria are released into the soil (Timmers et al., 2000). The presence

of a host leads to an increase in the soil population of rhizobia (Beringer et al., 1979);

therefore, by engaging in the symbiosis, legumes potentially provide a significant fitness

advantage to rhizobia.

However, this interaction presents an evolutionary dilemma. A 'cheating' strain that reduces

investment in the costly process of nitrogen fixation, might be able to produce more

reproductive bacteria within the nodules, potentially giving it a fitness advantage. Legumes

are known to have evolved sanctions to punish cheating bacteria (Kiers et al., 2003; Oono et

al., 2011; West et al., 2002; Westhoek et al., 2017) and these sanctions can be applied

conditionally, with pea plants known to sanction nodules that contain an intermediate-fixing

strain, only when a better strain is available (Westhoek et al., 2021). Intriguingly, while

Westhoek et al (2021) did see a significant drop in carbon transport and nodule size by

twenty-eight days post inoculation, they did not see a significant decrease in the number of

viable bacteria within the sanctioned nodule (although the number of viable bacteria had

plummeted dramatically by fifty-six days post-inoculation). As such, the cause of the

reduction in nodule size at twenty-eight days post-inoculation, while hypothesised to be due

to a drop in the bacteroid population, has yet to be proven.

Poor fixers may be able to evade plant sanctions by 'piggybacking' on a more effective

strain, which can occur because multiple strains can occupy a single nodule, with mixed

nodules sometimes at frequencies of over 20%, (Mendoza-Suárez et al., 2020),. This route

could be avoided if the plant were able to distinguish between different strains within a

nodule and sanction the less effective one, but despite some evidence for so-called cell-

autonomous sanctioning, in the form of premature senescence of cells containing an

ineffective strain (Daubech et al., 2017; Regus et al., 2014), metabolic evidence has

suggested that this does not occur (Agtuca et al., 2020), so it remains contentious.

This study builds on previous work carried out by Westhoek et al. (2021) using pea plants

and a set of otherwise isogenic rhizobia strains that differ in their ability to fix nitrogen.

First, we used flow cytometry to measure fitness characteristics of both undifferentiated

bacteria and nitrogen-fixing bacteroids within nodules infected with a single rhizobial strain

and fluorescence microscopy to establish how sanctions influenced the health and

morphology of the infected plant cells within those nodules. Second, we used the same

techniques on mixed nodules to determine whether or not pea plants carry out cell-

autonomous sanctioning within nodules. Our results show that sanctions reduce the

number of bacteroids within whole nodules as well as reducing the size of the

undifferentiated bacteria. Fluorescence microscopy showed that the breakdown of cells

within the nodule occurred prematurely in sanctioned nodules. Our results also show that

these sanctions operate at the level of the whole nodule but not at the cellular level. As

such, a piggybacking strain cannot be punished independently of the non-cheater within a

mixed nodule; however, mixed nodules were treated as ineffective and sanctioned at the

nodule level. As such, piggybacking is not an effective route by which cheaters can succeed.

Methods

We co-inoculated plants with two rhizobial strains selected from three otherwise isogenic

strains differing only in their ability to fix nitrogen. They were characterised as a good fixer

(Fix$^+$), an intermediate fixer (Fix^{int}) and a non-fixer (Fix$^-$, see *Bacterial strains and culture*

conditions for more details) This gives three pairwise combinations of a more effective (E)

and less effective (L) strains; Fix$^+$ (E) vs Fix$^-$ (L), Fix$^+$ (E) vs Fix^{int} (L) and Fix^{int} (E) vs Fix$^-$ (L). i.e.,

Fix$^+$ is always effective and Fix$^-$ always less effective, but Fix^{int} can be either effective or less

effective, depending on the co-inoculated strain. To measure conditional sanctioning at the

whole-nodule level, we sampled nodules occupied by single strains only (identified by a

fluorescent marker), and passed them through a flow cytometer to count and measure the

bacteroid and undifferentiated bacterial populations. To confirm that the fluorescent

marker had no effect on the symbiotic characteristics of Rlv3841 the fixation rate and

nodulation efficiency of a strain containing each of our fluorescent tags was compared

against each other and a non-fluorescing wild-type strain. Sections of nodules were also

examined using confocal microscopy to assess cell health and morphology at a range of time

points. Both of these techniques allowed us to assess the effectiveness of plant sanctions on

bacterial populations.

Mixed nodules also form on plants co-inoculated with two or more strains. To test the

theory of cell-autonomous sanctioning, the size and number of bacteroids and bacteria from

within mixed nodules were quantified using flow cytometry, and mixed nodules were

imaged using confocal microscopy. We set up five or six plants of each of our three strain

combinations (Fix$^+$ & Fix$^-$, Fix$^+$ & Fix^{int} and Fix^{int} & Fix$^-$) and passed five nodules from each

plant through the flow cytometer. Further nodules were also picked for confocal microscopy

to inspect the health of different cell types. These techniques allow us to see whether the

plant can discriminate between strains within a single nodule and apply sanctions

accordingly.

Bacterial Strains and Culture Conditions

Rhizobial strains used in this study are all derivatives of a highly effective nitrogen-fixing

strain, *Rhizobium leguminosarum* bv. (Rlv) 3841, that infects pea (*Pisum sativum* L. cv.

Avola) (Johnston & Beringer, 1975). The mutant strains thus differ in their nitrogen-fixation

ability but are otherwise genetically identical (Table 1). They are labelled with a fluorescent

marker (mCherry or GFP) in order to distinguish the nodules formed by each strain (Table 1).

Strains were maintained on tryptone-yeast (TY) agar (Beringer, 1974) with the appropriate

concentrations of antibiotics (Table 1). For long-term storage, strains were kept at −80 °C in

TY with 15 to 20% glycerol. Rhizobial inoculant was grown on TY agar and the number of

bacteria on the slope was determined by measuring the OD600 of the washed slope using a

Name	Strain	Antibiotic resistance	Description	Reference

Genesys 150 UV-Visible spectrophotometer. A dilution of $1x10^8$ cells ml^{-1} was then

produced.

Plant Growth

Before sowing, all pea seeds were surface sterilized (1 minute in 95% ethanol followed by 5

minutes in 2% NaClO), rinsed, and left to germinate on 1% w/v agar plates at room

temperature in the dark. Seedlings were transplanted after five days by transferring them to

sterilized 1 litre Azlon beakers containing a 1:1 mixture of silver sand and fine vermiculite,

150 ml sterilized nitrogen-free nutrient solution, and 5 x 10^7 cells of each of the two

rhizobial strains (see also Westhoek et al., (2017)). Beakers were covered with clingfilm to

reduce aerial contamination, which was slit after a few days to allow seedlings to grow

through. Plants were grown in a growth room (21 °C, 16 h photoperiod) for 28 to 42 days

and watered as necessary from 7 days onwards.

Table 1: Rhizobial strains and their fixation abilities. All strains derived from Rlv3841 and provided with a strain code, resistance markers, short description and reference.				
Fix mCherry	OPS1341	Streptomycin, Gentamicin	Rlv3841 Tn7-Gm-mCherry	(Westhoek et al., 2021)
Fix$^+$ GFP	OPS1339	Streptomycin, Genatmicin	Rlv3841 Tn7-Gm-GFP	(Westhoek et al., 2021)
Fix$^-$ GFP	OPS2270	Streptomycin, Spectinomycin, Gentamicin	Ω*nifH* Tn7-Gm-sfGFP (Rlv3841 mutant, ΩSpc cassette in *nifH*)	(Westhoek et al., 2021)
Fix^{int} GFP	OPS2268	Streptomycin, Spectinomycin, Gentamicin	Fix^{int} Tn7-Gm-sfGFP (Rlv3841 mutant, ΩSpc cassette in promoter region of *nifA*)	(Westhoek et al., 2021)
Fix^{int} mCherry	OPS2269	Streptomycin, Spectinomycin, Gentamicin	Fix^{int} Tn7-Gm-mCherry False colour: orange	(Westhoek et al., 2021)

Harvesting

Plants were harvested at 28-, 35- and 42-days post-inoculation by removing the plant from
the sand/vermiculite mixture and washing the roots carefully. Nodules were then imaged
using a LEICA M165 FC fluorescent stereo microscope and an iBright FL1500 imaging system.
Nodule occupants were identified based on their fluorescence. When selecting single
occupancy nodules, the five largest of each nodule type was selected.

Acetylene reduction assay

Fixation rates were measured via an acetylene reduction assay using the method described
in Westhoek et al (2021).

Confocal imaging

To assess the health of cells containing undifferentiated bacteria and bacteroids within
nodules, we imaged sections using confocal microscopy, which allows the visualisation of
fluorescently tagged strains and discrimination of the two strains within a mixed nodule. We
placed selected nodules in 8 % w/v agar and cut 100 µm longitudinal sections through the
centre of the nodule using a Leica VT1200S vibratome. Nodule slices were then imaged with
a Zeiss LSM 880 Airy Scan confocal microscope and analysed with ZEN Black software. To
visualise fluorescent tags, mCherry was excited using a 561 nm wavelength laser and
emissions detected between 598 and 649 nm, while GFP was excited using a 488 nm
wavelength laser and emissions detected between 498 and 562 nm.

Flow Cytometry

To identify and quantify the population sizes of undifferentiated bacteria and bacteroids
within nodules, we carried out flow cytometry. This was done initially with selected single
occupant nodules from the three co-inoculant combinations (Fix^+ vs Fix^-, Fix^+ vs Fix^{int} and
Fix^{int} vs Fix^-). We then carried out flow cytometry on the three types of mixed nodules (Fix^+
& Fix^-, Fix^+ & Fix^{int} and Fix^{int} & Fix^-). All nodules were prepared by placing them in a 1.5ml
Eppendorf tube with 300µl of harvest solution (0.9% NaCl, 0.02% SILWET L-77) and crushing
them with an autoclaved microcentrifuge pestle. This solution was passed through a 40µm
filter and diluted tenfold to increase the accuracy of the flow cytometry. Flow cytometry
was conducted with a Amnis® CellStream® flow cytometer (Luminex) equipped with a 488

and 561nm lasers, which were used for excitation of GFP and mCherry, respectively. Flow

rates were set to low speed and high sensitivity (3.66 µL/min). Analysis of flow cytometry

data was carried out using the CellStream® Analysis software (Version 1.5.17). Bacterial

events were defined based on custom gating parameters. Singlets and doublets were gated

with a threshold of 0.400 forward scatter (FSC) aspect ratio. Bacterial singlets with emission

at 611/631nm, when excited at 561 nm, above 6,000 arbitrary units were defined as red.

Bacterial singlets with emission at 528/546nm, when excited at 488nm, above 4,000

arbitrary units were defined as green. Bacteroids and bacteria were defined based on size,

as in Mergaert et al., (2006), using custom gating of the FSC detection to separate the two

populations (Figure S1). Flow cytometry data available at http://flowrepository.org with

experiment codes (FR-FCM-Z8AA &) for the single occupancy experiments and the mixed

nodule, experiments respectively.

Statistical Analysis

To test the effect of the fluorescent markers on the symbiotic characteristics of Rlv3841, we

compared the fixation rates of Rlv3841, OPS1341 and OPS1399 (wild-type strains with

different fluorescent markers) using one-way ANOVA. We compared the nodulation

competitiveness of the same three strains using paired t-tests.

To test for conditional sanctioning at the whole-nodule level, we subsetted the data into the

three different nodule types: Fix^+, Fix^- and Fix^{int}, and tested whether three relevant

measures of bacterial fitness and plant sanctioning behaviour were dependent on the

identity of the co-inoculated strain. The chosen measures were: (1) the size of the

undifferentiated bacterial population (a key measure of bacterial fitness), (2) the size of the

bacteroid population and (3) the size of individual undifferentiated bacteria. We did not

measure the size of individual bacteroids because the numbers of bacteroids within some

sanctioned nodules was so small as to make this measure unreliable. Both population size

and size of individuals were analysed using mixed-effects models in which individual plant ID

was the random effect and the identity of the co-inoculant was the fixed effect.

To test for the presence of cell autonomous sanctioning within mixed nodules, we used a

one-sample t-test to determine whether the proportion of the two strains differed

significantly from 50% for both undifferentiated bacteria and bacteroid populations. To
compare the absolute numbers of undifferentiated bacteria and bacteroids within mixed
nodules, we used paired t-tests.

All statistical tests were carried out using R version 4.2.1 (2022-06-23) and R studio
2022.07.2 and graphs were produced using Graph Pad version 9. An R markdown document
is provided (Doc. S3). In figures, bacteria and bacteroid population data is presented log10
transformed. Where a log transformation was carried out for statistical analysis to create a
better model this has been noted in the results.

Results

Fluorescently tagging strains does not affect symbiotic characteristics of Rlv3841

We found that there was no significant difference in the nodulation competitiveness of the
two fluorescently tagged strains when co-inoculated against the untagged wild-type
Rlv3841 (Fix$^+$), nor did the two tagged strains differ from each other when co-inoculated
together (Fix$^+$ vs Fix$^+$ mCherry: estimate = 22.750, SE = 13.288, t = 1.712, p > 0.05, df = 6; Fix$^+$
vs Fix$^+$ GFP: estimate = -5.75, SE = 20.84, t = -0.276, p>0.05, df =6; Fix$^+$ GFP vs Fix$^+$ mCherry:
estimate = -12.250, SE = 11.265, t = -1.087, p>0.05, df = 6). In addition, single inoculated
peas with each of the three strains did not differ in fixation rates per nodules (F = 1.387, p >
0.05, n = 4, df = 2) so no further pairwise comparisons were made.

Conditional sanctioning on whole nodules determines bacteroid population size
254

The number of bacteroids does not depend on the identity of the co-inoculated strain for
either Fix$^+$ nodules (estimate = 970769, SE = 2039541, t value = 0.476, p =0.644, n =60) or
Fix$^-$ nodules (Log transformed data, estimate = -0.540, SE = 0.299, t value = -1.806, p = 0.104,
n = 55) . However, nodules occupied by the intermediate-fixing strain contained significantly
fewer bacteroids when the co-inoculated strain was Fix$^+$ rather than Fix$^-$ (Log transformed
data, estimate = -2.209, SE = 0.324, t value = -6.818, p = 7.74x10^{-5}, n = 55). (Figure 1A)
261

Conditional sanctioning on whole nodules determines the size of bacteria

The average size of the undifferentiated bacteria did not depend on the identity of the co-inoculant for Fix[+] nodules (estimate = 561.3, SE = 301.6, t = 1.861, p = 0.0923, n = 60). However, undifferentiated Fix[-] bacteria were significantly smaller when co-inoculated with a Fix[+] strain (Log transformed data, estimate.= -0.112, SE = 0.043, t value = -2.566, p value = 0.03, n = 55), and undifferentiated Fix[int] bacteria were significantly smaller when co-inoculated with a Fix[+] strain (estimate -1417.6, SE = 405.9, t value = -3.492, p value = 0.00, n = 55; Figure 1B)

Conditional sanctioning on whole nodules does not impact bacterial survival after 28 days

The number of undifferentiated bacteria within a nodule did not depend on the identity of the co-inoculant for Fix[+] nodules (estimate = -212437, SE = 472483, t = -0.450, p = 0.663, n = 60), Fix[-] nodules (estimate = 2794843, SE = 2008643, t = 1.391, p = 0.19753, n = 55) or Fix[int] nodules (Log transformed data, estimate = 0.500, SE = 0.5855, t = 0.855, p = 0.415, n = 55; Figure 1C).

Conditional sanctioning causes changes in cell morphology

There is a clear change to the morphology of the whole nodule of sanctioned nodules after twenty-eight days (Figure S4), as seen in Westhoek et al., (2017). Nodules from co-inoculated plants of all three combinations (Fix[+] vs Fix[-], Fix[+] vs Fix[int], Fix[int] vs Fix[-]) were imaged at 28-, 35- and 42-days post-inoculation and examined for evidence of changes to cell morphology (Figure S5 for all pictures).

At twenty-eight days post-inoculation, all sections taken from nodules containing the Fix[int] strain looked similar, regardless of whether the co-inoculated strain was Fix[-] (Figure 2A) or Fix[+] (Figure 2B). However, at 35 days post-inoculation, the status of nodules containing the Fix[int] strain was visibly affected by the identity of the co-inoculated strain, consistent with conditional sanctioning. In particular, if the co-inoculated strain was Fix[+], then the infected cells within Fix[int] nodules had irregular shapes and the active infected region had retracted

away from the nodule edge (Figure 2D). However, if the co-inoculated strain was Fix$^-$, then

most cells retained a round morphology; indeed these nodules were not visibly different

from Fix^{int} nodules at 28 days post-inoculation (compare Figures 2A&2B). At 42 days post-

inoculation, all nodules containing the Fix^{int} strain showed clear changes to cell morphology;

however, when co-inoculated with Fix$^+$ (Figure 2F), these changes were more pronounced

than when co-inoculated with Fix$^-$ (Figure 2E).

Conditional sanctioning is not cell-autonomous in peas

In mixed nodules, the proportion of bacteroids that belonged to the less effective strain was

not significantly different from 50% (mean = 50.57%, 95%CI = 39.92 & 61.22%, t = 0.11; p

>0.05, n = 24) (Figure 3A) and the same was true for the undifferentiated bacteria (mean =

59.62%, 95%CI = 49.72 - 69.51%, t = 2.01, p >0.05, n = 24) (Figure 3C); hence, we did not find

evidence for cell-autonomous sanctioning (although we note the number of

undifferentiated bacteria was not an indicator of sanctioning in our single strain nodule

data). There was also no significant difference in the size of the undifferentiated bacteria

within mixed nodules (95%CI = -139.18 & 897.55, t = 1.513, p >0.05, n = 24) (Figure 3B); The

same pattern emerged when we compared absolute numbers rather than proportions. The

number of bacteroids belonging to the effective strain within a mixed nodule was not

significantly different from the number belonging to the less effective strain (95%CI = -

236167 - 1834023, t = 0.124, p >0.05, n = 24) and the same was true for undifferentiated

bacteria (95%CI = -148145 & 192398, t = 0.269, p > 0.05, n = 24). These results therefore

show no evidence for cell-autonomous sanctioning.

Mixed nodules are moderately sanctioned

When we consider the total population within mixed nodules, we see that the total number

of bacteroids and the average size of undifferentiated bacteria within mixed nodules

occupies an intermediate value between nodules occupied by the two single strains (Figure

4A&B). This would suggest that mixed nodules are treated as less effective nodules and

sanctioned accordingly. This may imply that mixed nodules are treated as a third strain

which is supplying nitrogen at an intermediate value between the two other strains present.

We did not see a clear pattern in the data for the number of undifferentiated bacteria from

the same set of nodules (Figure 4C). However, given that we did not see evidence for

sanctioning within this data set for the single occupant nodules this result neither supports

nor refutes our conclusions.

Changes to cell morphology are not cell-autonomous in pea

Within mixed nodules there was no evidence of any differences in cell morphology between

the more and less effective strain. If the nodule was unsanctioned (Figure 5A) then all of the

cells within the nodule were healthy, regardless of the strain occupying the cell. In contrast,

if a nodule was sanctioned (Figure 5B) then – as for a single occupant nodule (Figure 2D) –

infected cells within the nodule lost their typical spherical morphology and some burst

open. However, this was equally likely to affect cells containing the more or the less

effective strain. Therefore, we have seen no evidence to support the cell-autonomous

application of sanctions. This was true regardless of the strain combination within the

nodule or the time point at which they were harvested (Figure S6)

<u>**Discussion**</u>

We show that after twenty-eight days the size of undifferentiated bacteria (Figure 1B) and

the number of bacteroids (Figure 1A) but not the number of undifferentiated bacteria

(Figure 1C) within a sanctioned nodule have significantly decreased. That peas sanction

nodules containing less effectively fixing strains after twenty-eight days is consistent with

previous studies (Kiers et al., 2003; Oono et al., 2011; West et al., 2002; Westhoek et al.,

2017). Our undifferentiated bacteria data was consistent with the findings of Westhoek et

al., 2021, the number of undifferentiated bacteria did not drop significantly in sanctioned

nodules after twenty-eight days, post-inoculation. Our data is also consistent with

conditional sanctioning, as the fate of nodules infected by a Fix[int] strain depends on whether

the co-inoculated strain is Fix[+] or Fix[-] (Westhoek et al., 2021). As such we propose that the

sanctioned nodule phenotype, particularly their reduced size, is driven by this significant

drop in bacteroid population.

360
Within sanctioned nodules there was also a reduction in the size of the undifferentiated

bacteria. This would suggest that the cessation of the carbon supply to the nodule has

resulted in the starvation of these bacteria. As such, while their numbers remain high, they

are much smaller (Figure 1B), pre-dating the collapse in their population at a later time

point. As with the bacteroid data, the intermediate-fixing bacteria significantly differed in

size depending on the identity of the co-inoculating strain. Although we also found that the

size of undifferentiated Fix⁻ bacteria in single-strain nodules changed with the identity of the

co-inoculant, the estimated size difference was smaller, and close to the significance

threshold, so could be spurious.

370
371

Not only did the number of undifferentiated bacteria within sanctioned nodules not

decrease but the number of undifferentiated bacteria in our Fix⁻ nodules appeared to be

higher than in the unsanctioned nodules (Figure 1C). This result is particularly noteworthy as

only the undifferentiated bacteria are capable of replication once released into the soil. As

such the ability of sanctions to impact the fitness of a less effective strain is reliant on their

ability to reduce the number of undifferentiated bacteria. However, we know from previous

work that the population of undifferentiated bacteria within less effective nodules collapses

at a later time point. Westhoek et al., (2021) showed that while the population is the same

or slightly higher in sanctioned nodules after twenty-eight days it has reduced dramatically

by fifty-six. Crucially therefore, the population has been reduced by the time of nodule

release and the fitness of the sanctioned strain has been reduced.

Our results also cast some doubt over the use of non-fixing strains in studies investigating

sanctioning. Given that the process of conditionally sanctioning an intermediate fixer must

be an active process, unlike in the case of a non-fixing strain where the plant can simply

respond to the presence or absence of fixed nitrogen, as they must compare nitrogen

outputs. The use of non-fixing strains, which may be sanctioned passively through an

alternate system, in studies examining sanctions does not necessarily allow for the

investigation of the more nuanced elements of the mechanisms behind sanctioning. By

comparing our Fix⁻ and Fix^{int} data in figure 1 we see that non fixing and less effective strains

are treated differently. Particularly, the number of the undifferentiated bacteria which is
much higher in Fix⁻ nodules then in a sanctioned Fix^{int} nodule. This result adds weight to the
argument that future studies of sanctioning should be focussed on using strains of varying
fixation effectiveness and should not utilise non-fixing strains.

We also show that there is a change in the morphology of the ultra-structure within
sanctioned nodules. Previous studies (Regus et al., 2017) have made the link between these
morphological changes and the process of nodule senescence, which seems reasonable to
us. By tracking this change as the nodule develops, we have now established an
understanding of the sequence of events that take place throughout sanctioning. The effect
of sanctions on whole nodule morphology, nodules are smaller and spherical, is discernible
by 16 days (Westhoek et al., 2017). This initial nodule change is present at the same time as
a drop in the number of bacteroids and the size of undifferentiated bacteria after 28 days
(Figure 1). However, there is no visible change in the nodule ultra-structure of sanctioned
nodules before 35 days (Figure 2). All of these changes appear to precipitate the collapse in
undifferentiated bacteria population which is observed by 56 days.

It has been proposed that cell-autonomous sanctions might occur within mixed nodules
(Daubech et al., 2017; Regus et al., 2017), allowing plants to combat less effective
"cheating" strains that enter nodules alongside a more effective strain. If a plant can
differentiate among strains in a mixed nodule and target sanctions accordingly, then for two
equally competitive strains, we expect the average proportion of the more effective strain
to be greater than 50%. Therefore, evidence for cell autonomous sanctions would include a
bias in the proportion of the effective strain within mixed nodules. However, in our study,
the ratio of effective to less effective bacteroids within mixed nodules did not significantly
differ from 50% (Figure 3A). As shown in this study (Figure 1B) plant sanctions also reduce
the average size of individual undifferentiated bacteria within single-strain nodules;
therefore we would expect to see the same size reduction within the mixed nodule, if cell-
autonomous sanctions occur. No such difference was found (Figure 3B).

We also found no difference in the numbers of undifferentiated bacteria within mixed
nodules belonging to the two strains. However, as sanctions did not change the numbers of

undifferentiated bacteria in single-strain nodules after twenty-eight days, this seems to be
of limited relevance to the presence, or lack of, cell autonomous sanctions. As such we
found no evidence within our flow cytometry data to support the presence of cell-
autonomous sanctions in pea plants.

By examining the treatment of mixed nodules at the whole nodule level (Figure 4) we have
shown that mixed nodules are sanctioned on the whole nodule level. This sanctioning is less
severe than for single strain nodules containing the less effective strain on the plant. As
seen in Figure 4, therefore, the mixed nodules are moderately sanctioned with their
bacteroid number and bacteria sizes fitting between the values for the single occupant
sanctioned and unsanctioned nodules. As such we have shown that while mixed nodules
are not sanctioned in a cell-autonomous fashion they are sanctioned at the whole nodule
level.

Cells within mixed nodules underwent senescence simultaneously, regardless of variation in
effectiveness (Figure 5), consistent with the timing of cell senescence not occurring in a cell-
autonomous manner. Although the timing of senescence varied among mixed nodules,
where senescence occurred, it affected both strains equally. It may therefore be
hypothesised that the timing of senescence within mixed nodules is determined by the ratio
of effective to less effective strains within the nodule. This would be challenging to test,
because sectioning nodules does not give an accurate measure of strain ratio due to the
irregular 3D distribution of strains within nodules.

Our results align with the findings of Agtuca et al., (2020) who found that the metabolic
profile of sectors of mixed nodules containing different strains did not show significant
differences. However, our results contrast with studies which have found evidence for cell
autonomous sanctions based on variation in the timing of senescence, such as Regus et al.,
(2017) and Daubech et al., (2017). The differences may be explained by variation in the
legume-rhizobium system used: Daubech et al. (2017) used *Cupriavidus* taiwanensis –
Mimosa pudica; the (Regus et al., 2017) study used both *Acmispon strigosus* –
Bradyrhizobium and *Lotus japonicus* – *Mesorhizobium*; and the Agtuca et al. (2020) study

used *Glycine max – Bradyrhizobium japonicum.* A comprehensive understanding of

sanctions will therefore require the continued use of multiple experimental systems.

We have shown that the sanctioning of less effective nodules within the pea-rhizobium

symbioses is conditional in nature and occurs at the nodule level. In Fix^+/Fix^{int} coinfected

plants, nodules containing Fix^{int} but not Fix^+ strains are sanctioned, suggesting conditional

sanctioning requires comparison between nodule-specific and global nitrogen signals. This

might be achieved through the interaction between a nitrate receptor such as the Nitrate

transreceptor NRT1.1, which both transports nitrate as well as detecting nitrate levels

(Zhang et al., 2019) and the NIN-like Proteins (NLPs) 1 and 4, which are essential for the

nitrate-based regulation of nodule maturation (Lin et al., 2021). In high nitrate conditions

NLP1/4 inhibit cytokinin biosynthesis. Cytokinin biosynthesis drives nodule maturation

through the activation of a signal cascade through the Cytokinin responsive element CRE1

(Lin et al., 2021), which acts to promote expression of the *cep* and *cle* genes (Laffont et al.,

2020). However, how a global signalling mechanism is able to detect the nitrogen output of

individual nodules and compare it to global nitrogen status remains unclear. The

mechanism of sanctions on individual nodules may be achieved through nodule-specific

proteins responding to nitrate levels. One possibility is SnRK1 which, when phosphorylated

by the DMI2 kinase in response to Nod factor, phosphorylates malate dehydrogenase 1 and

2, leading to increased malate production within the nodule to supply bacteroids (Guo et al.,

2023). This is therefore a prime candidate for a pathway in which the legume would be able

to reward effective strain-containing nodules, as well as to punish nodules containing less

effectively fixing strains by cessation of malate production and supply. However, these

remain speculations about signalling pathways whose elucidation may aid future efforts to

engineer symbioses in the selection and use of more effective nitrogen-fixing bacteria.

We now have clear evidence that the change in nodule morphology shown by Westheok et

al., (2017) is driven by a reduction in the number of bacteroids and the size of the

undifferentiated bacteria. However, the number of undifferentiated bacteria does not drop

until a later timepoint. These changes precipitate the changes to the morphological

structure of infected plant cells within sanctioned nodules. We have also shown that, while

the plant is unable to discriminate between effective and less effective strains within a

mixed nodule, 'piggybacking' on an effective strain does not provide a viable route by which

to evade sanctions. Any potential benefit is limited, because the plant views mixed nodules

as less effective, and sanctions them accordingly.

<u>References</u>

Agtuca, B. J., Stopka, S. A., Evans, S., Samarah, L., Liu, Y., Xu, D., Stacey, M. G., Koppenaal, D.
W., Paša-Tolić, L., Anderton, C. R., Vertes, A., & Stacey, G. (2020). Metabolomic
profiling of wild-type and mutant soybean root nodules using laser-ablation
electrospray ionization mass spectrometry reveals altered metabolism. *Plant Journal*,
103(5), 1937–1958. https://doi.org/10.1111/tpj.14815

Beringer, J. E. (1974). R Factor Transfer in Rhizobium leguminosarum. *Journal of General*
Microbiology, *84*(1g74), 188–198.

Beringer, J. E., Brewin, N., Johnston, A. W., Schulmanf, H. M., & Hopwood, D. A. (1979). The
Rhizobium-legume symbiosis. *Proc. R' Soc.Lond' B*, *204*, 219–233.
https://royalsocietypublishing.org/

Daubech, B., Remigi, P., Doin de Moura, G., Marchetti, M., cile Pouzet, C., Auriac, M.-C.,
Gokhale, C. S., Masson-Boivin, C., & Capela, D. (2017). Spatio-temporal control of
mutualism in legumes helps spread symbiotic nitrogen fixation. *ELIFE*, 1–21.
https://doi.org/10.7554/eLife.28683.001

Guo, D., Liu, P., Liu, Q., Zheng, L., Liu, S., Shen, C., Liu, L., Fan, S., Li, N., Dong, J., & Wang, T.
(2023). Legume-specific SnRK1 promotes malate supply to bacteroids for symbiotic
nitrogen fixation. *Molecular Plant*, *16*(9), 1396–1412.
https://doi.org/10.1016/j.molp.2023.08.009

Johnston, A. W. B., & Beringer, J. E. (1975). Identification of the Rhizobium Strains in Pea
Root Nodules Using Genetic Markers. *Journal of General Microbiology*, *87*, 343–350.

Kiers, E. T., Rousseau, R. A., West, S. A., & Denlson, R. F. (2003). Host sanctions and the
legume-rhizobium mutualism. *Nature*, *425*(6953), 78–81.
https://doi.org/10.1038/nature01931

Laffont, C., Ivanovici, A., Gautrat, P., Brault, M., Djordjevic, M. A., & Frugier, F. (2020). The
NIN transcription factor coordinates CEP and CLE signaling peptides that regulate
nodulation antagonistically. *Nature Communications*, *11*(1).
https://doi.org/10.1038/s41467-020-16968-1

Lee, D. K. (2020). Data transformation: A focus on the interpretation. *Korean Journal of*
Anesthesiology, *73*(6), 503–508. https://doi.org/10.4097/kja.20137

Lin, J., Roswanjaya, Y. P., Kohlen, W., Stougaard, J., & Reid, D. (2021). Nitrate restricts
nodule organogenesis through inhibition of cytokinin biosynthesis in Lotus japonicus.
Nature Communications, *12*(1). https://doi.org/10.1038/s41467-021-26820-9

Mendoza-Suárez, M., Andersen, S. U., Poole, P. S., & Sánchez-Cañizares, C. (2021).
Competition, Nodule Occupancy, and Persistence of Inoculant Strains: Key Factors in
the Rhizobium-Legume Symbioses. In *Frontiers in Plant Science* (Vol. 12). Frontiers
Media S.A. https://doi.org/10.3389/fpls.2021.690567

Mendoza-Suárez, M., Barney Geddes, B., Sánchez-Cañizares, C., Ramírez-González, R.,
Kirchelle, C., Jorrin, B., & Poole, P. (2020). Optimizing Rhizobium-legume symbioses by

simultaneous measurement of rhizobial competitiveness and N2 fixation in nodules.
PNAS, 117(18).

Mergaert, P., Uchiumi, T., Alunni, B., Naë Lle Evanno, G., Lique Cheron, A., Catrice, O.,
Mausset, A.-E., Dé Rique Barloy-Hubler, F., Galibert, F., Kondorosi, A., & Kondorosi, E.
(2006). *Eukaryotic control on bacterial cell cycle and differentiation in the Rhizobium-*
legume symbiosis. https://www.pnas.org

Oono, R., Anderson, C. G., & Denison, R. F. (2011). Failure to fix nitrogen by non-
reproductive symbiotic rhizobia triggers host sanctions that reduce fitness of their
reproductive clonemates. *Proceedings of the Royal Society B: Biological Sciences,*
278(1718), 2698–2703. https://doi.org/10.1098/rspb.2010.2193

Poole, P., Ramachandran, V., & Terpolilli, J. (2018). Rhizobia: From saprophytes to
endosymbionts. *Nature Reviews Microbiology, 16*(5), 291–303.
https://doi.org/10.1038/nrmicro.2017.171

Regus, J. U., Gano, K. A., Hollowell, A. C., & Sachs, J. L. (2014). Efficiency of partner choice
and sanctions in Lotus is not altered by nitrogen fertilization. *Proceedings of the Royal*
Society B: Biological Sciences, 281(1781). https://doi.org/10.1098/rspb.2013.2587

Regus, J. U., Quides, K. W., O'Neill, M. R., Suzuki, R., Savory, E. A., Chang, J. H., & Sachs, J. L.
(2017). Cell autonomous sanctions in legumes target ineffective rhizobia in nodules
with mixed infections. *American Journal of Botany, 104*(9), 1299–1312.
https://doi.org/10.3732/ajb.1700165

Timmers, A. C. J., Soupène, E., Auriac, M.-C., De Billy, F., Vasse, J., Boistard, P., & Truchet, G.
(2000). Publication no. M-2000-0914-03R.. 1996; Long 1996) and takes place through a
series of developmental stages. In *Schultze and Kondorosi* (Vol. 13, Issue 11).

Udvardi, M., & Poole, P. S. (2013). Transport and metabolism in legume-rhizobia symbioses.
In *Annual Review of Plant Biology* (Vol. 64, pp. 781–805).
https://doi.org/10.1146/annurev-arplant-050312-120235

West, S. A., Kiers, E. T., Simms, E. L., & Denison, R. F. (2002). Sanctions and mutualism
stability: Why do rhizobia fix nitrogen? *Proceedings of the Royal Society B: Biological*
Sciences, 269(1492), 685–694. https://doi.org/10.1098/rspb.2001.1878

Westhoek, A., Clark, L. J., Culbert, M., Dalchau, N., Griffiths, M., Jorrin, B., Karunakaran, R.,
Ledermann, R., Tkacz, A., Webb, I., James, E. K., Poole, P. S., & Turnbull, L. A. (2021).
Conditional sanctioning in a legume-Rhizobium mutualism. *Proceedings of the National*
Academy of Sciences of the United States of America, 118(19), 1–8.
https://doi.org/10.1073/pnas.2025760118

Westhoek, A., Field, E., Rehling, F., Mulley, G., Webb, I., Poole, P. S., & Turnbull, L. A. (2017).
Policing the legume-Rhizobium symbiosis: A critical test of partner choice. *Scientific*
Reports, 7(1). https://doi.org/10.1038/s41598-017-01634-2

Zhang, X., Cui, Y., Yu, M., Su, B., Gong, W., Baluška, F., Komis, G., Šamaj, J., Shan, X., & Lin, J.
(2019). Phosphorylation-mediated dynamics of nitrate transceptor NRT1.1 regulate
auxin flux and nitrate signaling in lateral root growth. *Plant Physiology, 181*(2), 480–
498. https://doi.org/10.1104/pp.19.00346

571
572

573 ## **Acknowledgments**

574 This work was supported by the Biotechnology and Biological Sciences Research Council
575 [BB/M011224/1, BB/T001801/1, BB/T006722/1, BB/T008784/1 and BB/W006219/1].

Competing interests

The authors declare no competing interests

Author contributions

All authors listed have made a substantial direct and intellectual contribution to the work and approved it for publication. TJU, BJ and PSP conceived this study and designed the experiments. TJU and BJ performed the experiments. TJU analysed the data and prepared the manuscript. TJU, BJ, LAT and PSP critically reviewed the manuscript.

Data Availability

Flow cytometry: Data is available at http://flowrepository.org with experiment codes FR-FCM-Z6XN and FR-FCM-Z6XX for the single occupancy experiments and the mixed nodule experiments respectively.

Supporting information

Fig. S1 Image of flow cytometry output showing the two populations differentiated by size based on Forward Scatter (FSC)

Table. S2 Flow cytometry data, analysed using custom gating to calculate the number of undifferentiated bacteria and bacteroids in single occupancy and mixed nodules, and acetylene reduction assay and nodulation results from fluorescent tag control experiments.

Doc.S3 R markdown file of statistical analysis.

Fig. S4 Representative iBright image of pea inoculated with two strains of differing effectiveness.

Fig. S5 Representative confocal images of nodule sections of both nodule types from all three co-inoculation combinations (Fix$^+$ vs Fix$^-$, Fix$^+$ vs Fix^{int}, Fix^{int} vs Fix$^-$) at 28-, 35- and 42-days post inoculation.

Fig. S6 Representative images of the three types of mixed nodules at 28, 35 and 42 days post inoculation

Fig 1. After 28 days sanctions reduce the number of bacteroids and bacterial size, nut not the number of bacteria.

The number of bacteroids (A) and the size of bacteria (B) in nodules from pea plants co-inoculated with two strains of rhizobia. Data is split into different nodule types (Fix[+], Fix[int] or Fix[-]). The identity of the co-inoculant is given in brackets. Black lines are the mean value with error bars showing the standard error of the mean. Significance level from paired t-test *** < 0.001; ** < 0.01; * <0 .05; ns, not significant. Nodules harvested at 28 days post-inoculation.

Fig. 3. The two strains within a mixed nodule are not sanctioned independently of one another

The proportion of the bacteroid (A), average size of the bacteria (B) and the proportion of bacteria (C) within a mixed nodule belonging to the more or less effective of the two strains.

Nodules harvested twenty-eight days post inoculation.

Fig. 4. Mixed nodules are sanctioned, but less severely than single strain nodules containing the less effective strain.

The number of bacteroids (A) and the average size of the undifferentiated bacteria (B) and the number of undifferentiated bacteria (C) within pea nodules. Data is split by co-inoculant combination (Fix^+ vs Fix^{int}, Fix^+ vs Fix^- and Fix^{int} vs Fix^-). The mean value for the single strain nodules of each of the two strains in each combination and the mean value for mixed nodules containing the two strains is given with error bars showing the standard error of the mean. Nodules were harvested twenty-eight days post inoculation.

Unsanctioned **Sanctioned**

Fig. 5. The two strains within a mixed nodule do not senesce at different times despite varying in relative effectiveness.

Thirty-five day post-inoculation unsanctioned (A) and sanctioned (B) mixed nodules containing two strains of differing fixation effectiveness. When unsanctioned (A), all cells of both strains remain spherical and intact (1). When sanctioned (B), some cells of both strains remain intact (2) but they are no longer spherical and other cells of both strains have burst open (3).
Longitudinal nodule slices (100µm) were imaged by confocal microscopy. Scale bar = 200µm.