The *Chlamydia trachomatis* secreted effector CebN targets nucleoporins and Rae1 to antagonize

STAT1 nuclear import

Brianna Steiert[1,2], Cherilyn A. Elwell[3], Xavier Tijerina[1], Madison A. Elliott[4], Yennifer

Delgado[5,6], Jocelyn Ni[5,6], Robert Faris[1], Parker Smith[1], Shelby E. Andersen[1, 6], Paige N.

McCaslin[1], Julia Kevil-Yeager[1], Quinn Eldridge[1], Brian S. Imai[8], Justine V. Arrington[8], Peter M.

Yau[8], Kathleen M. Mirrashidi[3], Jeffrey R. Johnson[9], Erik Verschueren[9], John Von Dollen[9],

Gwendolyn M. Jang[9], Nevan J. Krogan[9,10,11], Mehdi Bouhaddou[5,6], Christine Suetterlin[4], Joanne

N. Engel[3,12], Mary M. Weber[1*]

[1]Department of Microbiology and Immunology, University of Iowa Carver College of Medicine,

Iowa City, Iowa, USA.

[2]Present address: Department of Veterinary Microbiology and Pathology, Washington State

University, Pullman, WA, USA

[3]Department of Medicine, University of California, San Francisco, San Francisco, CA, USA.

[4]Department of Developmental and Cell Biology, University of California, Irvine, California,

USA

[5]Department of Microbiology, Immunology, and Molecular Genetics (MIMG), University of

California, Los Angeles, Los Angeles, CA, USA

[6]Institute for Quantitative and Computational Biosciences (QCBio), University of California, Los

Angeles, Los Angeles, CA, USA

[7]Present address: Department of Immunology and Microbiology, University of Colorado -

Anschutz Medical Campus, Aurora, CO, USA

[8]Protein Sciences Facility, Roy J. Carver Biotechnology Center, University of Illinois

Urbana–Champaign, Urbana, IL, USA

[9]QB3, California Institute for Quantitative Biosciences, San Francisco, CA 94148, USA

[10]Gladstone Institutes, San Francisco, CA 94158, USA

[11]Department of Cellular and Molecular Pharmacology, University of California, San Francisco,

San Francisco, CA 94158, USA

[12]Department of Microbiology and Immunology, University of California, San Francisco, San

Francisco, CA, USA.

*Correspondence: mary-weber@uiowa.edu

Keywords: *Chlamydia trachomatis*, CT584, CebN, nucleoporin, Rae1, T3SS effector, Inc

ABSTRACT

To usurp host defenses and establish a replicative niche, obligate intracellular pathogens

are tasked with remodeling the host cell using a comparatively small repertoire of effector

proteins. For *Chlamydia trachomatis* (*C.t*), discovery of secreted proteins and their host targets

has been particularly challenging due to the bacterium's historical genetic intractability. Using

affinity purification-mass spectrometry, we defined host interaction partners for 21 secreted

effector proteins, providing the first comprehensive type III secretion system (T3SS) effector-

host interactome generated during infection. Among these, we show that the C-terminus of CebN

(CT584) binds multiple nucleoporins and Rae1, host factors previously associated only with viral

immune evasion. Remarkably, we shown that CebN localizes to the nuclear envelope not only in

infected cells but also in uninfected bystander cells. Functionally, CebN is both necessary and

sufficient to perturb STAT1 nuclear import following IFN-γ stimulation and its expression is

critical for *C.t.* survival, as evidenced by reduced bacterial replication and smaller inclusions in

cells infected with a CebN mutant. Together, these finds expand our understanding of chlamydia

effector biology and highlight novel bacterial strategies for manipulating host defenses at the

nuclear pore.

SIGNIFICANCE

Chlamydia trachomatis (*C.t.*) is a leading cause of sexually transmitted infections and

blindness, yet the molecular mechanisms it uses to manipulate host defenses remain poorly

defined. Unlike many pathogens, *C.t.* relies on a limited set of effectors to remodel the host cell

and establish its niche. We identified host targets for 21 *C.t.* effector proteins. Focusing on

CebN, we show that it binds nucleoporins and Rae1, host factors previously linked only to viral

immune antagonism. CebN localizes to the nuclear envelope of both infected and bystander

cells, and is critical for replication, inclusion development, and perturbation of STAT1 nuclear

import following IFN-γ stimulation. These findings uncover a novel strategy by which *C.t.*

manipulates nuclear pore function to evade host defenses and establish infection.

                                          INTRODUCTION

To counteract host defense mechanisms and establish a favorable replicative niche,

intracellular pathogens are tasked with remodeling the host cell using secreted virulence factors,

termed effector proteins. Identification of the host proteins and pathways targeted by secreted

proteins during active infections has been exceedingly difficult for obligate intracellular

pathogens owing to their genetic intractability[1]. Several obligate intracellular pathogens,

including *Chlamydia trachomatis* (*C.t.*), are the etiological agents of important human diseases

for which no vaccines exist[2]. *C.t.* is the leading cause of infectious blindness and is the most

common bacterial sexually transmitted infection worldwide[3]. Untreated infection can result in

severe complications including pelvic inflammatory disease, ectopic pregnancy, sterility, and

increased risk of developing cervical and ovarian cancer[4,5]. The incidence and prevalence of *C.t.*

infections are rapidly rising due to a lack of long-term protective immunity and treatment failure

following antibiotic therapy[2]. Understanding how *C.t.* co-opts the host cell to form its unique

replicative niche is vital to developing new therapies.

All *Chlamydiae* share a biphasic developmental cycle in which the bacteria alternate

between two forms: the infectious elementary body (EB) and the replicative reticulate body

(RB)[3]. Upon contact with a target host cell, the EB delivers a set of pre-synthesized type III

secretion system (T3SS) effector proteins into the eukaryotic cell to drive cytoskeletal

rearrangements and membrane remodeling, triggering endocytosis of the pathogen[6–9]. The

plasma membrane-derived compartment in which the EB resides is rapidly modified by the

pathogen to form a unique replicative niche, termed the inclusion. The inclusion quickly

dissociates from the endolysosomal pathway[10], trafficking along microtubules to the peri-Golgi

region[11] where the EB differentiates into an RB and initiates replication. Following multiple

rounds of division, RBs convert to EBs, and the bacteria are released by extrusion of EBs or host

cell lysis to begin the infection cycle anew[12]. While it is well established that formation of an

intact replicative niche is vital for *C.t.* proliferation and chlamydial disease[13,14], how *C.t.*

accomplishes such feats remains incompletely understood.

C.t. secreted effectors fall into two major classes: inclusion membrane proteins (Incs) and

conventional T3SS (cT3SS) proteins. While substantial progress has been made in identifying

and characterizing Incs[15] , comparatively little is known about cT3SS effectors. Using T3SS

secretion assays in *Chlamydia*, 33 cT3SS effectors have been identified to date[16,17]. Here, we

leveraged chlamydial genetics, in conjunction with large-scale unbiased affinity purification-

mass spectrometry (AP-MS), to comprehensively map the host pathways targeted by cT3SS

effector proteins during infection. We identified high-confidence interacting partners for 21

cT3SS proteins. Intriguingly, we show that CT584, which we have renamed <u>C</u>hlamydia <u>e</u>ffector

<u>b</u>locking <u>n</u>uclear transport (CebN), binds to a subset of Phe-Gly (FG) nucleoporins (NUPs) and

the mRNA export factor Rae1, which are host targets previously only associated with viral

infection[18–22]. Our data indicate that CebN predominately localizes to the nuclear envelope in

both infected and bystander cells and is necessary and sufficient to inhibit STAT1 import into the

nucleus following interferon (IFN)-γ stimulation. Additionally, a CebN deficient strain of

chlamydia exhibits a marked decrease in growth, further emphasizing the importance of this

effector in infection. This work significantly contributes to our understanding of *C.t.* cT3SS

effectors and their host targets, providing a key stepping-stone for elucidating how these

effector-host interactions contribute to the pathogenesis of *C.t.* infections.

MATERIALS AND METHODS

Bacterial and cell culture. All proteins analyzed in this study were derived from *C.t.* L2

but as is convention in the field, we use the *C.t.* D nomenclature. *Chlamydia trachomatis* serovar

L2 (LGV 434/Bu) was propagated in HeLa 229 cells (America Type Culture Collection) and

EBs were purified using a gastrografin density gradient as previously described[23]. HeLa cells

were propagated in RPMI 1640 medium (Thermo Fisher Scientific) supplemented with 10%

Fetal Bovine Serum (Gibco) at 37°C with 5% CO_2. A2EN cells (Kerafast) were grown in

keratinocyte-serum free media (K-SFM) (Thermo Fisher Scientific) supplemented with

0.16 ng/mL epidermal growth factor, 25 µg/mL bovine pituitary extract, 0.4 mM $CaCl_2$, and

gentamicin at 37°C with 5% CO_2 [24].

Plasmid construction. For AP-MS, each cT3SS effector was cloned into

the NotI/KpnI site of pBomb4-tet-mCherry with a FLAG-tag added to the C-terminus of each *orf*

by PCR. The CebN CRISPRi gblock was cloned into pBOMBL12CRia::L2 by GenScript as

previously described[25] . For the sRNA knockdown approach, a 30 nucleotide-long KD sequence

that targets the sequence in the CebN 5' UTR from -41 to -12 in respect to the start codon was

cloned into pBOMB5-tet-CtrR3 by Gibson assembly to generate pBomb5-tet-CebN sRNA, as

previously described[26]. CebN truncations were similarly cloned into pBomb4-tet-mCherry as

FLAG-tagged fusions. For ectopic expression, CebN was cloned into the KpnI/XhoI site of

pcDNA3.1-GFP. The integrity of all constructs was verified by DNA sequencing at McLab. All

primers used in this study are listed in Table S1.

Transformation of *C.t.* *C. trachomatis* serovar L2 (LGV 434/Bu) EBs were transformed

as previously described[27] with minor modifications. Briefly, plasmid DNA (5 µg), fresh *C.t.* EBs

from infected host cell lysates (~2 x10^6 EBs), and 10 µl 5X transformation mix (50 mM Tris pH

7.4 and 250 mM $CaCl_2$) were gently mixed and the final volume was adjusted to 50 µl with

tissue-culture grade water. Mixtures were incubated at room temperature for 30 min. RPMI with

10% FBS (4 ml) was then added to each transformation mix and 2 ml was applied to 2 wells of a

6-well plate containing a confluent HeLa cell monolayer. Plates were centrifuged at 900 x g for

30 min and at 18h post-infection, the media was replaced with RPMI with 10% FBS containing

0.3 µg/ml penicillin G (PenG). Infectious progeny were harvested every 48h and used to infect

fresh HeLa cell monolayers until viable inclusions were evident (~2-3 passages). Expression of

individual FLAG-tagged fusion proteins was induced using 10 ng/ml anhydrous tetracycline

(aTc), added at time of infection, and expression was confirmed by western blotting. For

pBomb4-tet-CebN CRISPRi and pBomb5-tet-CebN sRNA, 5 ng/ml anhydrous tetracycline (aTc)

was added 3h post-infection and knockdown was confirmed by western blotting using 1:2000

anti-CebN antibody (kindly provided by Luís Jaime Mota[28]).

Western blotting. For AP-MS expression verification and subsequent blots, samples

were resolved either using 3-8% Tris-Acetate protein gels with Tris-Acetate SDS running buffer

(proteins with MW >100kDa) or 4-12% Bis-Tris protein gels with MES running buffer (proteins

with MW <100kDa). Proteins were transferred to a PVDF membrane, and following blocking

with 5% milk in Tris-buffered saline with 0.1% Tween 20, were probed using anti-FLAG

(Thermo Fisher Scientific, 701629), anti-GFP (Novus, NB600-597), anti-NUP54 (Proteintech,

16232-1-AP), anti-NUP153 (Novus, NBP1-81725), or anti-NUP214 (abcam, AB70497)

antibodies (Table S2).

Immunofluorescence (IF) microscopy. For visualization of CebN by stimulated

emission depletion (STED) microscopy, HeLa cells were transfected with pcDNA3.1-GFP

empty vector or plasmid encoding for GFP-tagged CebN. Cells were fixed with 2%

formaldehyde and permeabilized with 0.1% Triton-X 100 at 24h post-transfection and stained

with DAPI and NUP specific antibodies: anti-NUP54, anti-NUP153, or anti-NUP214. Images

were captured on a Leica SP8 inverted microscope. Images were deconvoluted using Imaris

Professional Software.

For visualization of CebN during infection, HeLa cells were infected at an MOI of 2 with

WT *C.t.* or *C.t.* strains expressing a FLAG-tagged empty vector, CebN-FLAG, or TmeA-FLAG.

Expression was induced for 24h using 10 ng/ml aTc added at the time of infection. Cells were

fixed with 4% formaldehyde 24h post-infection and stained with DAPI (ThermoFisher

Scientific), anti-FLAG (Cell Signaling, 14994T), and anti-*C.t.* HSP60 (Sigma, MABF2108).

Images were captured on a Nikon A1 Confocal.

Affinity Purification (AP): HeLa cells, in three T175 flasks, were infected at an MOI of

2 with *C.t.* strains expressing a FLAG-tagged effector protein. Expression was induced for 24h

using 10 ng/ml aTc, added at the time of infection. Four h prior to lysis, 10 µM MG132

(Millipore Sigma) was added to the media. Cells were subsequently lysed in eukaryotic lysis

solution (ELS) (50 mM Tris HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, and 1% Triton-X 100)

with Halt protease inhibitor cocktail (Thermo Fisher Scientific). After incubation on ice for 20

min, lysates were centrifuged at 12,000 x g for 20 min, and the supernatants were incubated with

60 µl preclearing beads (mouse IgG agarose, Millipore Sigma) for 2h at 4°C. The precleared

lysate was then incubated with 30 µl FLAG beads (anti-FLAG M2 Affinity Gel, Millipore

Sigma) overnight at 4°C. The beads were washed six times with ELS without detergent. For

mass spectrometry, samples were stored in 50 mM ammonium bicarbonate prior to digestion and

analysis as previously described[29]. For western blotting, proteins were eluted from the beads in

4X NuPAGE LDS Sample Buffer (Thermo Fisher Scientific) and boiled for 5 min.

Mass Spectrometry (MS). MS was performed as previously described[29], with the

following adjustments. Beads containing samples were washed with 25 mM ammonium

bicarbonate and digested with 0.5 micrograms trypsin (Pierce, Thermo Fisher Scientific, MS

Grade) using a CEM microwave reactor for 30 min at 55°C. Digested peptides were extracted

twice using 50% acetonitrile plus 5% formic acid, lyophilized to dryness, and resuspended in 5%

acetonitrile plus 0.1% formic acid. For LC/MS, samples were injected into an UltiMate 3000

UHPLC system coupled online to a Thermo Scientific Orbitrap Fusion Tribrid mass

spectrometer. Peptides were separated by reversed-phase chromatography using a 50-cm

MicroPac Nano C18 column (Thermo Fisher Scientific) with mobile phases of 0.1% formic acid

in water and 0.1% formic acid in acetonitrile; a linear gradient from 4% to 35% Acetonitrile over

the course of 45 min was employed for peptide separations. The mass spectrometer was operated

in a data-dependent acquisition (DDA) mode, employing precursor scans from 300 to 1,500 m/z

(120,000 resolution) followed by collision induced dissociation (CID) of the most intense

precursors over a maximum cycle time of 3 s (35% NCE, 1.6 m/z isolation window, 60-s

dynamic exclusion window). Raw LC-MS/MS data were converted to peak lists using Mascot

Distiller 2.8 and searched against a database containing UniProt_Human and

Chlamydia_trachomatis_L2434Bu using Mascot 2.8 (Matrix Science). Tryptic digestion was

specified with a maximum of two missed cleavages, while peptide and fragment mass tolerances

were set to 10 ppm and 0.6 Daltons, respectively. Label-free Quantitation was performed

utilizing the Mascot Average method on Mascot Distiller 2.8.2.

AP-MS analysis. Mass Spectrometry interaction STatistics (MiST) was used, as

previously described [30], to assign a confidence score to every host protein identified from MS.

Localization, Reactome pathways, biological processes, and molecular functions were

determined for each host prey with a MiST[30] score ≥0.7 using Uniprot and GeneCards. Dot plots

for visualization were generated using R package ggplot2. For the protein-protein interaction

network, R programming language version 4.4.1 and CORUM [31] were used to analyze the host

proteins identified in the AP-MS and compare the dataset with previous published studies[32–34].

Cytoscape (3.10.3)[35] was used to visualize the protein-protein interaction network.

STAT1 and mCherry translocation assays. HeLa cells were transfected with

pcDNA3.1-GFP plasmids containing empty vector, CebN, or TmeA using Lipofectamine LTX

(Thermo Fisher Scientific). For mCherry-NLS translocation, 4h post-GFP-transfection the cells

were transfected with mCherry-NLS (Addgene 58476). At 24h post transfection, the media was

changed, with half the samples receiving normal RPMI media and half with RPMI with 100

U/ml IFN-γ for 1h. To monitor STAT1 nuclear translocation during infection, cells were infected

at an MOI of 1 with or without 5 ng/ml aTc added to the media at the time of infection. At 24h

post-infection, RPMI with 100 U/ml IFN-γ was added to half the samples for 1h. Transfected or

infected cells were fixed with 4% formaldehyde, permeabilized with 0.1% Triton-X, blocked in

3% BSA, and stained with DAPI, anti-STAT1 (Cell Signaling 14994T), and anti-*C.t.* HSP60

antibodies (ThermoFisher Scientific MA3-023). Images were collected on Nikon Eclipse 2

microscope using the same exposure conditions between groups. For all nuclear translocation

experiments (STAT1 and mCherry) nuclear signal was quantified by measuring the fluorescence

intensity of STAT1/mCherry in the nucleus using Fiji with 150 transfected or infected cells per

biological replicate with three replicates.

Growth curve and inclusion measurements: HeLa or A2EN cells were infected at an

MOI of 1 on ice. After 30 min, the media was changed with half the samples receiving 5 ng/ml

aTc. For growth curves, samples were collected at 0 and 48 h, lysed in water and applied to fresh

HeLa cell monolayers to determine the number of infectious forming units as previously

described[14,36,37]. For inclusion area measurements, infected cells were fixed with 4%

formaldehyde at 24h post-infection and stained using anti-*C.t.* HSP60 antibodies. Circles were

drawn around the inclusions in Fiji to measure the inclusion area using 30-45 images across three

coverslips per biological replicate with three replicates total.

Statistics: Statistical analyses were performed using GraphPad Prism 10.5.0 software.

Depending on the dataset, either Welch's *t*-test or one-way ANOVAs followed by Tukey's

multiple comparisons were applied. Statistical significance was defined as $P < 0.05$ (*), $P < 0.01$

(**), and $P < 0.001$ (***), $P < 0.0001$ (****).

RESULTS

Identification of putative host proteins and pathways targeted by cT3SS effector

proteins during *C.t.* infection. A few cT3SS effector proteins have been functionally

characterized and shown to modulate diverse host cell signaling pathways[6,8,9,29,38–41]. However,

the function of most of these putative virulence factors remains unknown. Affinity purification-

mass spectrometry (AP-MS) has emerged as a powerful technique to comprehensively map

protein-protein interactions (PPIs) between bacterial effectors or viral proteins and host proteins,

yielding key mechanistic insights into how these pathogens establish their unique replicative

niches[15,30,42–44]. While informative, most of these studies have been undertaken by

overexpressing a single effector protein in a mammalian cell at non-physiological levels and in

the absence of additional bacterial or viral factors that might promote or hinder PPIs. With the

increasing ease of genetic tractability of *C.t.*, we are poised to evaluate effector-host PPIs in the

context of infection.

Here, we leveraged AP-MS to systematically interrogate host interaction networks of 33

C.t. cT3SS effectors in the context of infection[16,17,45]. Of these, 24 were successfully expressed in

C.t. and the remaining effector proteins were excluded from further analysis due to the inability

to obtain chlamydial transformants or due to the lack of detectable expression by western

blotting. Following AP-MS, only putative host target or "prey" proteins with at least 2 unique

peptides that were present in at least 2 of the replicates were further pursued. CT311 and CT161

were excluded from further analysis due to lack of detection of the bait protein following AP-

MS. To identify high-confidence PPIs, we analyzed the complete data set using Mass

Spectrometry interaction STatistics (MiST) (Table S3), which evaluates prey reproducibility,

abundance, and specificity to generate scores between 0 and 1[30]. Using a cut-off score of ≥0.7,

we identified 241 putative host interacting partners for 21 cT3SS effectors. While CT144 was

detected in the AP-MS, and multiple preys were identified, none of these were predicted high-

confidence interactions when scored by MiST.

To further define the potential function of the *C.t.* cT3SS effector proteins, Gene

Ontology (Genecards) and pathway analysis (Reactome and Uniprot) was performed for each of

the 241 MiST high-confidence interactors (Table S3) (Fig. 1, S1). As shown in Supp. Fig. 1A,

most of the host proteins targeted by *C.t.* effectors reside in the cytoplasm, nucleus, endoplasmic

reticulum, and mitochondria; however, a few host proteins associated with the Golgi apparatus,

ribosomes, cytoskeleton, and plasma membrane were also noted. To further analyze the potential

function of these effector proteins, in conjunction with the above analysis, we employed

Cytoscape to map PPI networks to identify effector proteins associated with multiprotein

complexes or biological pathways (Fig. 1 and S1). Within this subset, we found individual

cT3SS effector proteins that associate with several members of multiprotein complexes,

including the proteosome (CebN), ribosome biogenesis (CT620 and CT691), tRNA processing

(CT627), mRNA processing (CT386), redox homeostasis (CT053), and the nuclear pore complex

(CebN) (Fig. 1). We also noted several instances of two or more cT3SS effector protein

interacting with the same host protein. For example, three cT3SS effectors (CT695/TmeB,

CT712, and CT392) interacted with coatomer subunit alpha (COPA) (Fig. 1), a protein required

for Golgi budding and that is essential for retrograde transport and Golgi structural integrity,

suggesting that these effectors may modulate COPI-dependent trafficking and Golgi architecture.

Three proteins (CT736, CT691, and CT392) interacted with transferrin receptor protein 1

(TRFC), which mediates cellular iron uptake (Fig. 1). Together with prior evidence that

transferrin is recruited to the inclusion in a CpoS-dependent manner[32,37,46,47], these findings

suggests that these effectors might aid in TFRC-dependent trafficking to facilitate iron

acquisition. Two cT3SS effector proteins (CebN and CT053) interacted with nucleolin (NCL), a

major component of the nucleolus. Most striking was the observation that CebN associates with

numerous components of the nuclear pore complex (NPC). In total, we identified 68 high-

confidence interactors (Table S3), many of which mediate ribonucleoprotein import, mRNA

export, or transcription (Fig. 1)

Consistent with proximity-labeling studies that identified inclusion-associated

proteins[33,34] and proteomic analyses of isolated inclusions[32], several host factors including

inverted formin 2 (IF2), COPA, transitional endoplasmic reticulum ATPase (VCP/p97), protein

transport protein Sec61 subunit alpha isoform 1 (SEC61A1), epidermal growth factor receptor

(EGFR), cytoplasmic dynein 1 intermediate chain 2 (DYNC1I2), amongst others, interacted with

cT3SS effector proteins (Fig. 1). These concordant observations provide orthogonal validation of

our AP-MS dataset and support the biological relevance of the effector-host interactome.

Ectopically expressed CebN binds to multiple nucleoporins and Rae1. Transfected

CebN-GFP predominately localized to the nuclear envelope (Fig. S2), a pattern that aligns with it

binding to host proteins involved in nucleocytoplasmic transport. To confirm these interactions,

and to rule out the requirement of additional bacterial proteins that contribute to the CebN

infection interactome, we performed AP-MS on Strep-tagged CebN as previously described[15].

This approach identified 30 high-confidence interactors (MIST $\geq$ 0.7) for CebN (Table S3), of

which 19 overlapped with the infection IP (Table 2, Fig. 2A). Notably, 9 nucleoporins (NUP58,

NUP214, NUP98, NUP54, NUP62, NUP88, NUP153, POM121/NUP121 and

RANBP2/NUP358) and the mRNA export factor Rae1 were present in both the CebN infection

and transfection interactomes.

Nucleoporins (NUPs) are a family of ~30 proteins that form the nuclear pore complex

(NPC) and play an important role in regulating import and export of small molecules into and out

of the nucleus[48]. The NPC is organized into an inner pore ring, the nuclear and cytoplasmic

rings, the nuclear basket, and the cytoplasmic filaments, each of which are enriched for select

NUPs (Fig. 2B)[48]. Intriguingly, while most of the NUPs that CebN binds make up the

cytoplasmic filaments, interactions with NUPs in other subcomplexes of the NPC were noted

(Fig. 2B), suggesting that secreted CebN may play a broad role in modulating NPC function.

Rae1 is an mRNA export factor that binds to NUP98 to aid in the transport of messenger

ribonucleoprotein (mRNP) complexes through the nuclear pore complex[49]. Several viral proteins

target NUPs and Rae1 to promote replication of their genomic information and to dampen the

host response to infection by blocking import of important transcription factors[19–22]. To the best

of our knowledge, no bacterial protein has been identified that targets NUP proteins or Rae1,

making CebN an intriguing effector protein to study.

CebN binds to and co-localizes with NUPs and Rae1. To confirm CebN binding to

NUP proteins and Rae1, we immunoprecipitated FLAG tagged CebN from *C.t.* infected cells and

probed with antibodies specific to NUPs and Rae1. We focused on NUP54, NUP153, and

NUP214 due to their high peptide counts in the AP-MS (Table S3). NUP54, NUP153, NUP214,

and Rae1 IP with CebN but not with vector or TmeA (Fig. 2C), an effector previously shown to

bind to N-WASP[6,8]. Processing of NUP153 and NUP214 was noted on these blots in infected

samples. We determined that CPAF, a broad-spectrum protease produced by *C.t.*, was

responsible for this cleavage, as this processing was absent in lysates derived from HeLa cells

infected with a CPAF mutant[50] (Fig. S3).

To additionally confirm the interaction between CebN and NUPs, HeLa cells were

transfected with GFP-CebN or GFP, fixed, and stained using anti-NUP54, NUP153, or NUP214

antibodies. Imaging by stimulated emission depletion (STED) microscopy confirmed that CebN

colocalized (white) with these specific NUP proteins, whereas no colocalization was noted with

GFP (Fig. 3A, B). Pearson's correlation coefficient was calculated as a measure of

colocalization, and a significant difference was found between GFP and GFP-CebN transfected

cells for each individual NUP (Fig. 3B). Taken together our results indicate that the cT3SS

effector protein CebN localizes to the nuclear envelope where it binds to multiple NUPs and to

Rae1.

CebN localizes to the nuclear envelope of infected and bystander cells. Most cT3SS

effector proteins are not readily visualized by microscopy, and thus their subcellular localization

is generally assessed by transfection of tagged proteins. Due to CebN's unique localization to the

nuclear envelope, we assessed CebN localization directly in *C.t.* infected cells. In line with our

ectopic expression data, CebN-FLAG was found to localize to the nuclear envelope of infected

cells (Fig. 3C). Intriguingly, we also observed CebN on the nuclear envelopes of bystander cells.

(Fig. 3C). While one possibility is that CebN is directly translocated into neighboring cells,

potentially through exosomes or tunneling nanotubes, an alternative explanation is that these

signals represent effectors retained in daughter cells following mitotic division.

The C-terminus of CebN is required for interactions with NUPs and Rae1 as well as

for its localization to nuclear envelope. To delineate the region of CebN that is necessary for

interaction with NUPs and Rae1, we generated 20-40 amino acid sequential truncations from the

C-terminus of the 183 amino acid protein and expressed these truncations as FLAG-tagged

constructs in *C.t.* Immunoprecipitation of these truncations, followed by subsequent western

blotting, showed that the C-terminal 23 amino acids of CebN are necessary for interactions with

NUP54, NUP153, NUP214, and Rae1 (Fig. 4A). We further confirmed the importance of this

region by performing confocal microscopy on HeLa cells infected with *C.t.* strains expressing

the CebN FLAG-tagged deletion constructs (Fig 4B). As noted above, full length CebN-FLAG

localizes to the nuclear envelope of infected cells, as well as the nuclear envelopes of bystander

cells (Fig. 4B). However, none of the truncated versions of CebN localized to the nuclear

envelope in infected or bystander cells (Fig. 4B). In total, the C-terminus of CebN is necessary

for its interaction with NUPs and Rae1 and for its localization to the nuclear envelope in infected

and bystander cells.

CebN is important for chlamydial replication *in vtiro*. To determine the importance of

CebN during *Chlamydia* infection, we generated two conditional CebN mutants, using CRISPRi

and sRNA systems[25,26,51], in which expression of CebN is repressed upon induction with

anhydrous tetracycline. Knockdown of CebN for both mutant strains was confirmed by western

blotting (Fig. 5A). CRISPRi was implemented using a deactivated Cas 12 enzyme that binds

186bp downstream of the *cebN* start site, thereby blocking effective transcription of the gene. In

parallel, the sRNA system employed engineered small RNAs that base-pair with *cebN* mRNA

transcripts to inhibit translation of CebN protein. Together, these complementary approaches

enable conditional knockdown of CebN at both the transcriptional and translational levels. In

both HeLa and A2EN cells, both mutants exhibited significantly reduced growth upon aTc

induction compared to uninduced or wild-type infections (Fig. 5B). The growth defect was

further reflected by a decrease in inclusion size in both cell types (Fig. 5C). Collectively, these

results demonstrate that CebN is required for efficient bacterial replication and inclusion

development.

CebN attenuates STAT1 import into the nucleus following interferon-γ stimulation.
Viral proteins from HIV, SARS-CoV-2, Kaposi's sarcoma-associated herpesvirus, and vesicular

stomatitis virus interact with and remodel the nuclear pore complex to modulate nuclear import

of transcription factors required for the antiviral response[19,20,22,52–58]. Similarly, *C.t.* attenuates

STAT1 nuclear import following IFN-γ stimulation[59]. We hypothesized that CebN interactions

with NUPs and Rae1 could underlie this perturbation. To test this, HeLa cells were transfected

with GFP-CebN, GFP-empty vector, or GFP-TmeA, treated with IFN-γ, and imaged by confocal

immunofluorescence microscopy using an anti-STAT1 antibody (Fig. 6A). We observed a

significant reduction in the proportion of cells with nuclear STAT1 in CebN-transfected cells

compared to those transfected with empty vector or TmeA (Fig. 6B), indicating that CebN is

sufficient to impair STAT1 nuclear translocation.

To test whether CebN is necessary for this process during infection, HeLa cells were

infected with WT or CebN CRISPRi mutant strain, treated with IFN-γ, stained with anti-STAT1

antibody, and nuclear STAT1 signal was quantified. We observed a significant increase in

STAT1 nuclear localization when the CebN mutant strain was induced with aTc, demonstrating

that CebN is required to block STAT1 nuclear translocation (Fig. 6C).

To assess whether this effect was due to a general block in nuclear import, we used a

nuclear localization signal (NLS)-tagged mCherry reporter plasmid. In HeLa cells co-transfected

with mCherry-NLS and GFP-empty vector, GFP-CebN, or GFP-TmeA, we detected no

significant differences in nuclear mCherry signal, suggesting that CebN selectively interferes

with import rather than imparting a general blockade (Fig. S4). Taken together, our results

suggest that CebN, through interactions with nucleoporins and Rae1, plays a key role in

dampening the host response to *C.t.* infection by perturbing nuclear translocation of

transcriptional regulators of the host innate immune response.

DISCUSSION

In this study, we combined *C.t.* genetics with AP-MS to generate the first cT3SS effector-

host interactome. Our approach successfully identified high confidence interacting host partners

for 21 of the 36 uncharacterized cT3SS effectors. Our work is especially valuable as it not only

begins to build a compendium of PPIs during active infection but also provides a launch point for

detailed mechanistic characterization of these effector proteins. Importantly, screens such as AP-

MS, can reveal novel pathways targeted by intracellular bacteria[42–44]. Here we discovered that

CebN targets NUPs to impair STAT1 nuclear import, revealing a potential mechanism by which

C.t. dampens the host innate immune response to survive intracellularly. Altogether, these

studies enhance our understanding of how obligate intracellular pathogens remodel the host to

form their unique replicative niches.

One of the most striking findings from our AP-MS screen was the interaction between

CebN and multiple nucleoporins and Rae1. While our study is, to the best of our knowledge, the

first time a bacterial effector has been shown to interact with host nucleoporin proteins, several

viral proteins have been identified that co-opt NUPs and Rae1. ORF6 of SARS-CoV-2, ORF10

of Kaposi's sarcoma-associated herpesvirus, and M protein of vesicular stomatitis virus all bind

to the NUP98-Rae1 complex, whereas the HIV-1 capsid binds to multiple nucleoporins leading

to altered NUP expression and localization[19,20,22,52–58]. While HIV-1 manipulates NUPs to

facilitate viral import and integration of its genome into the host genome[60], other viral proteins

interact with NUPs to disrupt nucleocytoplasmic transport of key transcription factors such as

STAT1. During SARS-CoV-2 infection, ORF6-Rae1-NUP98 interactions block STAT1 nuclear

import and mRNA export, resulting in a significantly diminished host response[18,20,22,57]. The

ability of CebN to perturb STAT1 import during *C.t.* infection, along with our observation that it

binds NUPs-Rae1, suggests it may subvert host defense mechanisms in a manner reminiscent of

viral proteins such as ORF6. However, given that nucleoporins and Rae1 regulate a broad range

of cellular processes including mRNA export, protein import, transcriptional regulation, and

chromatin organization[48], it remains possible that CebN's primary and physiologically relevant

function involves one or more of these other pathways.

450   Similar to how viruses dampen the immune response, *C.t.* has been shown to antagonize

interferon pathways[61–66]. IFN production activates the JAK-STAT signaling pathway, leading to

phosphorylation and homodimerization of STAT1, which is imported into the nucleus by

karyopherin alpha 1 and karyopherin beta 1 heterodimers[48]. Once in the nucleus, the STAT1

homodimer complex binds to gamma-activated site promoter elements to drive expression of a

subset of ISGs meant to impede the infection. Several studies have shown that following

prolonged IFN-γ stimulation, nuclear translocation of STAT1 is reduced[59,62] and inhibition of the

JAK-STAT pathway correlates with lower mRNA and protein levels of key interferon response

elements in infected cells compared to uninfected controls[67]. Furthermore, this difference was

dependent on *de novo C.t.* protein synthesis, supporting the role of a *C.t.* effector protein in this

process[67]. Our new data demonstrate that CebN is necessary and sufficient to decrease STAT1

import immediately following IFN-γ stimulation (1 h). Importantly, we show that this inhibition

is not global, as CebN does not block import of NLS-tagged mCherry. These findings suggest

that CebN impairs nuclear import of some factors but does not impart a total nuclear blockade.

Consistent with this, prior work by Walsh et al.[65] demonstrated that *C.t.* mutants lacking the Inc

GarD are highly susceptible to IFN-γ-mediated killing due to the activity of the interferon-

stimulated gene RNF213, and that RNF213 is induced at comparable levels in uninfected and

infected cells. Thus, while CebN limits STAT1 import, this may not necessarily result in

decreased expression of all ISGs, likely reflecting redundancy in IFN-γ-activated host defense

pathways. Determining the extent to which CebN-mediated inhibition of STAT1 impacts specific

ISGs is an area of future investigation.

Lining the NPC are intrinsically disordered NUPs that harbor numerous Phe-Gly (FG)

repeats separated by a hydrophilic spacer of 5-30 amino acids[48]. Movement of large cargo across

the NPC requires highly specific interactions between these so-called FG-NUPs and transporters

of the karyopherin family, which enables entry and rapid diffusion of the cargo-transporter

complex through the NPC. Of the 11 NUPs identified as putative binding partners of CebN, 9 are

classified as FG-NUPs (Table 2, S3). Recognition of FG motifs within these select NUPs might

explain how CebN binds to multiple nucleoporins and is able to selectively inhibit nuclear

import.

Crystallization of CebN revealed an N-terminal four-helix bundle (α1-4), followed by a

three-stranded antiparallel β-sheets (β1-3)[68], and a C-terminal kinked antiparallel pair of α-

helices[68]. AlphaFold modeling predicts that the C-terminus of CebN harbors a coiled-coil

domain (amino acids 153-181). Truncation of the last 23 amino acids of CebN would disrupt this

predicted coiled-coil, thus abrogating binding to NUPs and Rae1. While analysis of CebN did

not identify motifs known to be required for interactions with NUPs or Rae1, new motifs are

constantly being discovered, and it is possible that CebN possesses a previously undefined motif.

Short linear motifs (SLiMs), short 3-15 amino acid motifs often embedded in intrinsically

disordered or coiled-coil regions, serve as binding interfaces for structured partners and are

abundant in the human proteome as well as in viral and, less frequently, bacterial effectors[69,70].

Ectopically expressed CebN appeared to concentrate at the nuclear envelope, and we

observed a similar localization in infected cells and unexpectedly also in neighboring bystander

cells. This unique localization in apparently uninfected bystander cells has only ever been

reported once before with the *C. psittaci* cT3SS effector SINC, which similarly targets the

nuclear envelope through interactions with lamins[71]. How effectors reach bystander cells remains

unknown, but proposed mechanisms include packaging into exosomes for release at the cell

surface, tunneling nanotubes, or effectors leftover during cell division prior to inclusion

segregation into one daughter cell. Prior work with *Mycobacterium tuberculosis* has revealed

that mycobacterial proteins are packaged into vesicles and released via calcium-regulated

lysosomal exocytosis. These proteins are then trafficked to uninfected bystander cells[72].

Tunneling nanotubes, on the other hand, are transient cellular connections that play a role in cell-

to-cell communication and facilitate exchange of molecules between cells. Analogous to viruses,

previous work has shown that *C.t.* may spread cell-to-cell by nanotubules[73]. Thus, it is

conceivable that effector proteins may also be transmitted to adjacent cells via this mechanism. If

an infected cell undergoes cell division, the inclusion is partitioned into one cell, leaving the

other daughter cell "uninfected[74]." It is intriguing to consider that secreted effector proteins may

be left behind and continue to function in the absence of infection. Future work will involve

delineating the exact mechanism by which *C.t.* is able to transport effector proteins into

neighboring, uninfected cells and whether effectors besides CebN and SINC can access adjacent

cells.

As with all screens, false positives and negatives can result. To add rigor to our AP-MS

data set analysis, we employed MiST, which combines metrics of reproducibility, specificity,

and abundance across the entire data set to identify putative host binding partners more

accurately and stringently. Using this technique, we identified high confidence targets for 21 of

the secreted effectors tested herein. In our study, we sought to define PPIs for all the previously

uncharacterized cT3SS effector proteins and included CT695 (TmeB) as at the onset of this study

TmeB had no function ascribed to it. Recent work has since shown that ectopically expressed

TmeB targets the ARP2/3 complex[75]. While we did not find components of the ARP2/3 complex

in our infection AP-MS, we did identify an actin-binding protein, inverted formin 2 (INF2).

INF2 belongs to the formin family of proteins, which function to both polymerize and

depolymerize actin filaments[76]. Formins and the ARP2/3 complex act in parallel to regulate the

actin cytoskeleton[77]. Differences in experimental set-up between transfection of TmeB-FLAG

and infection of a *C.t.* strain expressing TmeB-FLAG could contribute to these differences in

identified putative host binding partners. Additionally, our timepoint of 24 hours post infection

may correlate with functions of TmeB beyond invasion.

By combining chlamydial genetics with large-scale screens, we have begun to define the

compendium of putative host proteins and pathways targeted by secreted *C.t.* effector proteins

during infection. Our analysis uncovered a wealth of unique high-confidence host interactors,

laying the groundwork for detailed mechanistic characterization of cT3SS effector proteins. Importantly, our approach has identified targets previously not associated with bacterial infection, including nucleoporins, which are commonly targeted by viral proteins to modulate host defenses[19–22]. We propose a model in which secretion of CebN leads to inhibition of STAT1 nuclear translocation through interactions with NUPs and Rae1 (Fig. 7), a mechanism that may alter host cell transcription and contribute to suppression of the innate immune response in infected and uninfected bystander cells. Moreover, the ability of CebN to translocate to bystander cells may provide a mechanism by which *C.t.* can prime nearby cells for infection. Understanding how CebN co-opts NUPs and Rae1 will not only advance our understanding of how *C.t.* establishes a persistent infection despite a robust host response but may also identify druggable targets applicable to both bacterial and viral infections.

Acknowledgements

We thank Dr. Raphael Valdivia for kindly sharing the *cpa*::*cat* mutant and Dr. Luís Jaime Mota for the CebN antibody. We thank Dr. Scot Ouellette for technical assistance with generation of the CRISPRi mutant. Thanks to Dr. Dominique Limoli for assistance with troubleshooting CebN staining during infection. We would like to thank Jawad Haider for excellent technical assistance and Steven Huang for critical review of this manuscript. We acknowledge grant support from the NIH (M.M.W.: R01 AI150812, R01 AI155434, R61 AI179999; J.E.: R01 AI073770, R01 AI105561; N.J.K: P50 GM082250; PO1 AI090935, PO1 090935, P50 GM081879, PO1 091575, U19 AI106754, U54AI081680, DARPA-10-93-Prophecy-PA-008), University of Iowa Training in Mechanisms of Parasitism T32 AI007511 (B.S. and X.T.), and the University of California San Francisco Pathogenesis and Host Defense T32 AI060537 (K.M.M). We also acknowledge support from the University of Iowa Stead

Family Scholars Program (M.M.W.), the University of Iowa Stinski fellowship (S.E.A), UCSF

Program for Breakthrough in Biomedical Research (J.E. and N.K.), and the American Heart

Association (K.M.M.). The Krogan Laboratory has received research support from Vir

Biotechnology, F. Hoffmann-La Roche, and Rezo Therapeutics. Nevan Krogan has a financially

compensated consulting agreement with Maze Therapeutics. He is the President and is on the

Board of Directors of Rezo Therapeutics, and he is a shareholder in Tenaya Therapeutics, Maze

Therapeutics, Rezo Therapeutics, GEn1E Lifesciences, and Interline Therapeutics.

558
559 Table 1: Host proteins with significant MiST scores (≥0.7) identified in both infection- and

560 transfection-based immunoprecipitations of CebN. Gene name and information regarding

561 subcellular localization, molecular function, and biological process was obtained from Uniprot.

Gene	Subcellular Localization	Molecular Function	Biological Process	Transfection IP MiST	Infection IP MiST
HNRNPR	nucleoplasm	RNA binding	mRNA processing	1.000	0.992
NUP58	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.995	0.993
NUP214	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.994	0.994
NUP98	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.993	0.994
RAE1	nuclear pore	RNA binding	transport: nucleocytoplasmic	0.991	0.992
TCF12	nucleosome	DNA binding	transcription	0.990	0.993
CNOT4	cytoplasm	transferase activity	ubiquitin-dependent protein catabolic process	0.989	0.986
NUP54	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.986	0.989
ZNF326	nuclear matrix	RNA binding	mRNA processing	0.982	0.986
NUP62	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.973	0.993

NUP88	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.972	0.992
POM121	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.971	0.968
AGFG1	nuclear pore	GTPase activator activity	mRNA export	0.963	0.988
NUP153	nuclear pore	structural component of nuclear pore complex	transport: nucleocytoplasmic	0.961	0.994
UNK	cytoplasm	RNA binding	translation	0.946	0.988
ANKRD17	nucleoplasm	protein binding	immune system process	0.911	0.983
TMEM131	membrane	unknown	unknown	0.866	0.994
ANKHD1	cytoplasm	protein binding	immune system process	0.826	0.983
RANBP2	nuclear pore	ligase activity	transport: nucleocytoplasmic	0.690	0.966

Figure 1. Host pathways targeted by *C.t.* secreted effector proteins identified using AP-MS. MiST was used to identify high-confidence interacting partners for each effector screened. Those with a MiST score ≥0.7 were considered significant and used for further analysis using the CORUM database to identify protein complexes targeted by each effector. An R script was used to compare the interactome to data from prior studies indicated by orange squares [32], brown circles [33], or purple diamonds[34] next to the human protein. Red diamonds represent *C.t.* effectors; grey circles represent human proteins; yellow shading highlights protein complexes; blue shading highlights biological pathways. Human-human PPI are shown as grey lines, and bacterial-human PPI interactions as blue lines, with line thickness correlating with MiST score.

Figure 2. CebN interacts with multiple nucleoporins and Rae1. (A) Venn diagram comparing unique and shared hits from infection (purple) and transfection (pink) AP-MS, showing a strong overlap of identified host targets. (B) Schematic of the nuclear pore complex. Bolded and asterisked NUPs were identified as putative CebN targets using AP-MS. (C) IP of *C.t.* expressing

FLAG-tagged vector, CebN, or TmeA from HeLa cells. Blots were probed with antibodies

specific to selected NUPs and Rae1. Data are representative of three replicates.

Figure 3. CebN localizes to the nuclear envelope during both transfection and infection

conditions. (A) STED images of HeLa cells transfected with GFP-tagged empty vector or GFP-

CebN (green) and stained with NUP-specific antibodies (magenta). (B) Quantification of

colocalization (white) was performed in Fiji using Pearson's correlation coefficient. Statistical

significance between GFP and GFP-CebN transfected samples is shown. The graph displays

individual values, the mean (black line), and standard deviation. ****P<0.0001; significance was

determined using one-way ANOVA followed by Tukey's multiple comparisons test. (C) HeLa

cells uninfected or infected at an MOI of 2 with WT L2 or *C.t.* expressing FLAG-tagged vector,

CebN, or TmeA (negative control). Cells were fixed with 4% formaldehyde and stained with

FLAG (orange), *C.t.* HSP-60 (green) and DAPI (blue). Zoomed panels show overexposed

images of the boxed region, highlighting nuclear envelopes of infected and/or bystander cells.

(A, C) Data are representative of three replicates.

Figure 4. The C-terminus of CebN mediates interactions with NUPs and Rae1 and is required

for nuclear envelope localization. (A) Lysates from HeLa cells infected for 24 h with *C.t.*

expressing the indicated C-terminally FLAG-tagged CebN constructs were immunoprecipitated

on FLAG beads and immunoblotted with antibodies specific to selected NUPs and Rae1. (B)

HeLa cells infected for 24 h with *C.t.* expressing the indicated C-terminally FLAG-tagged CebN

constructs were fixed with 4% formaldehyde and stained with antibodies to FLAG (orange), *C.t.*

HSP-60 (green, to visualize bacteria), and with DAPI (blue). Zoomed panels show

overexposures of the boxed region highlighting nuclear envelopes of infected and/or bystander

cells. (A-B) Data are representative of at least two replicates.

Figure 5. CebN is important for intracellular replication and inclusion development. (A) Western

blot confirmation of CebN knockdown in lysates from HeLa cells infected with CRISPRi and

sRNA chlamydial strains with or without aTc induction. (B) Quantification of infectious progeny

at 48 h postinfection in HeLa (left) or A2EN (right) cells infected with WT (grey), CebN

CRISPRi mutant (blue), or CebN sRNA mutant (purple). (C) Quantification of inclusion areas

(μm^2) in HeLa (left) and A2EN (right) cells infected with WT (grey), CebN CRISPRi mutant

(blue), or CebN sRNA mutant (purple) *C.t.* strains. (B-C) Statistical significance was determined

using Welch's t-test. *P<0.05, **P<0.01, ****P<0.0001. Error bars are standard deviation. (A-

C) Data are representative of three replicates.

Figure 6. CebN is necessary and sufficient to perturb IFN-γ-stimulated STAT1 nuclear

translocation. (A) HeLa cells were transfected with GFP-tagged vector, GFP-CebN, or GFP-

TmeA (green) and treated with 600U/ml IFN-γ. Cells were fixed and stained with anti-STAT1

(magenta) antibody and DAPI to demark the nucleus (blue). (B, C) Translocation of STAT1 into

the nucleus of transfected (B) or infected cells (C) was quantified in relative fluorescence units

(RFUs) from 150 cells per biological replicate. Error bars represent standard deviation. Statistical

significance was determined using one-way ANOVA followed by Tukey's multiple comparisons

test. ****P<0.0001. (A-C) Data are representative of three replicates.

Figure 7. Model for CebN-mediated inhibition of STAT1 nuclear translocation. In uninfected cells, IFN-γ stimulation triggers STAT1 translocation to the nucleus through the nuclear pore complex, where STAT1 binds to gamma-activated site (GAS) promoter elements to drive expression of a subset of interferon-stimulated genes that restrict infection. In *C.t.* infected cells, STAT1 translocation is blocked by CebN, which interacts with NUPs and Rae1. We hypothesize this interaction reduces the transcriptional activity of STAT1. In addition, CebN can be translocated into neighboring cells, where it may alter nuclear import and/or export pathways to prime these cells for subsequent infection.

REFERENCES

1. Andersen, S.E., Bulman, L.M., Steiert, B., Faris, R., and Weber, M.M. (2021). Got mutants? How advances in chlamydial genetics have furthered the study of effector proteins. Pathog Dis *79*, ftaa078-. https://doi.org/10.1093/femspd/ftaa078.

2. Maza, L.M. de la, Darville, T.L., and Pal, S. (2021). Chlamydia trachomatis vaccines for genital infections: where are we and how far is there to go? Expert Rev Vaccines *20*, 421–435. https://doi.org/10.1080/14760584.2021.1899817.

3. Elwell, C., Mirrashidi, K., and Engel, J. (2016). Chlamydia cell biology and pathogenesis. Nat Rev Microbiol *14*, 385–400. https://doi.org/10.1038/nrmicro.2016.30.

4. Das, M. (2018). Chlamydia infection and ovarian cancer risk. Lancet Oncol *19*, e338. https://doi.org/10.1016/s1470-2045(18)30421-2.

5. Koskela, P., Anttila, T., Bjørge, T., Brunsvig, A., Dillner, J., Hakama, M., Hakulinen, T., Jellum, E., Lehtinen, M., Lenner, P., et al. (2000). Chlamydia trachomatis infection as a risk factor for invasive cervical cancer. Int. J. Cancer *85*, 35–39. https://doi.org/10.1002/(sici)1097-0215(20000101)85:1<35::aid-ijc6>3.0.co;2-a.

6. Faris, R., McCullough, A., Andersen, S.E., Moninger, T.O., and Weber, M.M. (2020). The Chlamydia trachomatis secreted effector TmeA hijacks the N-WASP-ARP2/3 actin remodeling axis to facilitate cellular invasion. Plos Pathog *16*, e1008878. https://doi.org/10.1371/journal.ppat.1008878.

7. Keb, G., Hayman, R., and Fields, K.A. (2018). Floxed-Cassette Allelic Exchange Mutagenesis
Enables Markerless Gene Deletion in Chlamydia trachomatis and Can Reverse Cassette-Induced
Polar Effects. J. Bacteriol. *200*. https://doi.org/10.1128/jb.00479-18.

8. Keb, G., Ferrell, J., Scanlon, K.R., Jewett, T.J., and Fields, K.A. (2021). Chlamydia
trachomatis TmeA Directly Activates N-WASP To Promote Actin Polymerization and Functions
Synergistically with TarP during Invasion. Mbio *12*, e02861-20.
https://doi.org/10.1128/mbio.02861-20.

9. Ghosh, S., Ruelke, E.A., Ferrell, J.C., Bodero, M.D., Fields, K.A., and Jewett, T.J. (2020).
Fluorescence-Reported Allelic Exchange Mutagenesis-Mediated Gene Deletion Indicates a
Requirement for Chlamydia trachomatis Tarp during In Vivo Infectivity and Reveals a Specific
Role for the C Terminus during Cellular Invasion. Infect. Immun. *88*, 423–428.
https://doi.org/10.1128/iai.00841-19.

10. Scidmore, M.A., Fischer, E.R., and Hackstadt, T. (2003). Restricted Fusion of Chlamydia
trachomatis Vesicles with Endocytic Compartments during the Initial Stages of Infection. Infect.
Immun. *71*, 973–984. https://doi.org/10.1128/iai.71.2.973-984.2003.

11. Grieshaber, S.S., Grieshaber, N.A., and Hackstadt, T. (2003). Chlamydia trachomatis uses
host cell dynein to traffic to the microtubule-organizing center in a p50 dynamitin-independent
process. J Cell Sci *116*, 3793–3802. https://doi.org/10.1242/jcs.00695.

12. Hybiske, K., and Stephens, R.S. (2007). Mechanisms of host cell exit by the intracellular
bacterium Chlamydia. Proc National Acad Sci *104*, 11430–11435.
https://doi.org/10.1073/pnas.0703218104.

13. Sixt, B.S., Bastidas, R.J., Finethy, R., Baxter, R.M., Carpenter, V.K., Kroemer, G., Coers, J.,
and Valdivia, R.H. (2017). The Chlamydia trachomatis Inclusion Membrane Protein CpoS
Counteracts STING-Mediated Cellular Surveillance and Suicide Programs. Cell Host Microbe
21, 113–121. https://doi.org/10.1016/j.chom.2016.12.002.

14. Weber, M.M., Lam, J.L., Dooley, C.A., Noriea, N.F., Hansen, B.T., Hoyt, F.H., Carmody,
A.B., Sturdevant, G.L., and Hackstadt, T. (2017). Absence of Specific Chlamydia trachomatis
Inclusion Membrane Proteins Triggers Premature Inclusion Membrane Lysis and Host Cell
Death. Cell Reports *19*, 1406–1417. https://doi.org/10.1016/j.celrep.2017.04.058.

15. Mirrashidi, K.M., Elwell, C.A., Verschueren, E., Johnson, J.R., Frando, A., Von Dollen, J.,
Rosenberg, O., Gulbahce, N., Jang, G., Johnson, T., et al. (2015). Global Mapping of the Inc-
Human Interactome Reveals that Retromer Restricts Chlamydia Infection. Cell Host Microbe *18*,
109–121. https://doi.org/10.1016/j.chom.2015.06.004.

16. McCaslin, P.N., Andersen, S.E., Icardi, C.M., Faris, R., Steiert, B., Smith, P., Haider, J., and
Weber, M.M. (2023). Identification and Preliminary Characterization of Novel Type III Secreted
Effector Proteins in Chlamydia trachomatis. Infect. Immun., e0049122.
https://doi.org/10.1128/iai.00491-22.

17. Faris, R., Koch, R., McCaslin, P., Challagundla, N., Steiert, B., Andersen, S.E., Smith, P.,
Jabeena, C.A., Yau, P., Rudel, T., et al. (2025). The Chlamydia trachomatis secreted effector
protein CT181 binds to Mcl-1 to prolong neutrophil survival. bioRxiv, 2025.03.16.643443.
https://doi.org/10.1101/2025.03.16.643443.

18. Gao, X., Tian, H., Zhu, K., Li, Q., Hao, W., Wang, L., Qin, B., Deng, H., and Cui, S. (2022).
Structural basis for Sarbecovirus ORF6 mediated blockage of nucleocytoplasmic transport. Nat.
Commun. *13*, 4782. https://doi.org/10.1038/s41467-022-32489-5.

19. Kane, M., Rebensburg, S.V., Takata, M.A., Zang, T.M., Yamashita, M., Kvaratskhelia, M.,
and Bieniasz, P.D. (2018). Nuclear pore heterogeneity influences HIV-1 infection and the
antiviral activity of MX2. eLife *7*, e35738. https://doi.org/10.7554/elife.35738.

20. Kehrer, T., Cupic, A., Ye, C., Yildiz, S., Bouhaddou, M., Crossland, N.A., Barrall, E.A.,
Cohen, P., Tseng, A., Çağatay, T., et al. (2023). Impact of SARS-CoV-2 ORF6 and its variant
polymorphisms on host responses and viral pathogenesis. Cell Host Microbe *31*, 1668-1684.e12.
https://doi.org/10.1016/j.chom.2023.08.003.

21. Hall, R., Guedán, A., Yap, M.W., Young, G.R., Harvey, R., Stoye, J.P., and Bishop, K.N.
(2022). SARS-CoV-2 ORF6 disrupts innate immune signalling by inhibiting cellular mRNA
export. PLoS Pathog. *18*, e1010349. https://doi.org/10.1371/journal.ppat.1010349.

22. Addetia, A., Lieberman, N.A.P., Phung, Q., Hsiang, T.-Y., Xie, H., Roychoudhury, P.,
Shrestha, L., Loprieno, M.A., Huang, M.-L., Gale, M., et al. (2021). SARS-CoV-2 ORF6
Disrupts Bidirectional Nucleocytoplasmic Transport through Interactions with Rae1 and Nup98.
mBio *12*, e00065-21. https://doi.org/10.1128/mbio.00065-21.

23. Faris, R., and Weber, M. (2019). Propagation and Purification of Chlamydia trachomatis
Serovar L2 Transformants and Mutants. Bio-protocol *9*, e3459.
https://doi.org/10.21769/bioprotoc.3459.

24. Buckner, L.R., Schust, D.J., Ding, J., Nagamatsu, T., Beatty, W., Chang, T.L., Greene, S.J.,
Lewis, M.E., Ruiz, B., Holman, S.L., et al. (2011). Innate immune mediator profiles and their
regulation in a novel polarized immortalized epithelial cell model derived from human
endocervix. J Reprod Immunol *92*, 8–20. https://doi.org/10.1016/j.jri.2011.08.002.

25. Ouellette, S.P., Blay, E.A., Hatch, N.D., and Fisher-Marvin, L.A. (2021). CRISPR
Interference To Inducibly Repress Gene Expression in Chlamydia trachomatis. Infect. Immun.
89, e00108-21. https://doi.org/10.1128/iai.00108-21.

26. Ehses, J., Wang, K., Densi, A., Ramirez, C., Tan, M., and Sütterlin, C. (2024). Development
of an sRNA-mediated conditional knockdown system for Chlamydia trachomatis. mBio,
e0254524. https://doi.org/10.1128/mbio.02545-24.

27. Weber, M.M., and Faris, R. (2019). Mutagenesis of Chlamydia trachomatis Using
TargeTron. Methods Mol. Biol. *2042*, 165–184. https://doi.org/10.1007/978-1-4939-9694-0_12.

28. Pais, S.V., Milho, C., Almeida, F., and Mota, L.J. (2013). Identification of Novel Type III
Secretion Chaperone-Substrate Complexes of Chlamydia trachomatis. Plos One *8*, e56292.
https://doi.org/10.1371/journal.pone.0056292.

29. Steiert, B., Icardi, C.M., Faris, R., McCaslin, P.N., Smith, P., Klingelhutz, A.J., Yau, P.M.,
and Weber, M.M. (2023). The Chlamydia trachomatis type III-secreted effector protein CteG
induces centrosome amplification through interactions with centrin-2. Proc National Acad Sci
120, e2303487120. https://doi.org/10.1073/pnas.2303487120.

30. Jäger, S., Cimermancic, P., Gulbahce, N., Johnson, J.R., McGovern, K.E., Clarke, S.C.,
Shales, M., Mercenne, G., Pache, L., Li, K., et al. (2012). Global landscape of HIV–human
protein complexes. Nature *481*, 365–370. https://doi.org/10.1038/nature10719.

31. Steinkamp, R., Tsitsiridis, G., Brauner, B., Montrone, C., Fobo, G., Frishman, G., Avram, S.,
Oprea, T.I., and Ruepp, A. (2024). CORUM in 2024: protein complexes as drug targets. Nucleic
Acids Res. *53*, D651–D657. https://doi.org/10.1093/nar/gkae1033.

32. Aeberhard, L., Banhart, S., Fischer, M., Jehmlich, N., Rose, L., Koch, S., Laue, M., Renard,
B.Y., Schmidt, F., and Heuer, D. (2015). The Proteome of the Isolated Chlamydia trachomatis
Containing Vacuole Reveals a Complex Trafficking Platform Enriched for Retromer
Components. PLoS Pathog. *11*, e1004883. https://doi.org/10.1371/journal.ppat.1004883.

33. Olson, M.G., Widner, R.E., Jorgenson, L.M., Lawrence, A., Lagundzin, D., Woods, N.T.,
Ouellette, S.P., and Rucks, E.A. (2019). Proximity Labeling To Map Host-Pathogen Interactions
at the Membrane of a Bacterium-Containing Vacuole in Chlamydia trachomatis-Infected Human
Cells. Infect. Immun. *87*, e00537-19. https://doi.org/10.1128/iai.00537-19.

34. Dickinson, M.S., Anderson, L.N., Webb-Robertson, B.-J.M., Hansen, J.R., Smith, R.D.,
Wright, A.T., and Hybiske, K. (2019). Proximity-dependent proteomics of the Chlamydia
trachomatis inclusion membrane reveals functional interactions with endoplasmic reticulum exit
sites. PLoS Pathog. *15*, e1007698. https://doi.org/10.1371/journal.ppat.1007698.

35. Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., and Ideker, T. (2003). Cytoscape: A Software Environment for Integrated
Models of Biomolecular Interaction Networks. Genome Res. *13*, 2498–2504.
https://doi.org/10.1101/gr.1239303.

36. Weber, M.M., Noriea, N.F., Bauler, L.D., Lam, J.L., Sager, J., Wesolowski, J., Paumet, F.,
and Hackstadt, T. (2016). A Functional Core of IncA Is Required for Chlamydia trachomatis
Inclusion Fusion. J Bacteriol *198*, 1347–1355. https://doi.org/10.1128/jb.00933-15.

37. Faris, R., Merling, M., Andersen, S.E., Dooley, C.A., Hackstadt, T., and Weber, M.M.
(2019). Chlamydia trachomatis CT229 Subverts Rab GTPase-Dependent CCV Trafficking
Pathways to Promote Chlamydial Infection. Cell Reports *26*, 3380-3390.e5.
https://doi.org/10.1016/j.celrep.2019.02.079.

38. Chellas☐Géry, B., Linton, C.N., and Fields, K.A. (2007). Human GCIP interacts with
CT847, a novel Chlamydia trachomatis type III secretion substrate, and is degraded in a
tissue☐culture infection model. Cell Microbiol *9*, 2417–2430. https://doi.org/10.1111/j.1462-
5822.2007.00970.x.

39. Pennini, M.E., Perrinet, S., Dautry-Varsat, A., and Subtil, A. (2010). Histone Methylation by
NUE, a Novel Nuclear Effector of the Intracellular Pathogen Chlamydia trachomatis. Plos
Pathog *6*, e1000995. https://doi.org/10.1371/journal.ppat.1000995.

40. Hamaoui, D., Cossé, M.M., Mohan, J., Lystad, A.H., Wollert, T., and Subtil, A. (2020). The
Chlamydia effector CT622/TaiP targets a nonautophagy related function of ATG16L1. Proc.
Natl. Acad. Sci. *117*, 26784–26794. https://doi.org/10.1073/pnas.2005389117.

41. Dolat, L., Carpenter, V.K., Chen, Y.-S., Suzuki, M., Smith, E.P., Kuddar, O., and Valdivia,
R.H. (2022). Chlamydia repurposes the actin-binding protein EPS8 to disassemble epithelial
tight junctions and promote infection. Cell Host Microbe *30*, 1685-1700.e10.
https://doi.org/10.1016/j.chom.2022.10.013.

42. Fu, M., Liu, Y., Wang, G., Wang, P., Zhang, J., Chen, C., Zhao, M., Zhang, S., Jiao, J.,
Ouyang, X., et al. (2022). A protein–protein interaction map reveals that the Coxiella burnetii
effector CirB inhibits host proteasome activity. PLoS Pathog. *18*, e1010660.
https://doi.org/10.1371/journal.ppat.1010660.

43. Walch, P., Selkrig, J., Knodler, L.A., Rettel, M., Stein, F., Fernandez, K., Viéitez, C., Potel,
C.M., Scholzen, K., Geyer, M., et al. (2021). Global mapping of Salmonella enterica-host
protein-protein interactions during infection. Cell Host Microbe *29*, 1316-1332.e12.
https://doi.org/10.1016/j.chom.2021.06.004.

44. Penn, B.H., Netter, Z., Johnson, J.R., Dollen, J.V., Jang, G.M., Johnson, T., Ohol, Y.M.,
Maher, C., Bell, S.L., Geiger, K., et al. (2018). An Mtb-Human Protein-Protein Interaction Map
Identifies a Switch between Host Antiviral and Antibacterial Responses. Mol. Cell *71*, 637-
648.e5. https://doi.org/10.1016/j.molcel.2018.07.010.

45. Hower, S., Wolf, K., and Fields, K.A. (2009). Evidence that CT694 is a novel Chlamydia
trachomatis T3S substrate capable of functioning during invasion or early cycle development.
Mol. Microbiol. *72*, 1423–1437. https://doi.org/10.1111/j.1365-2958.2009.06732.x.

46. Ooij, C. van, Apodaca, G., and Engel, J. (1997). Characterization of the Chlamydia
trachomatis vacuole and its interaction with the host endocytic pathway in HeLa cells. Infect.
Immun. *65*, 758–766. https://doi.org/10.1128/iai.65.2.758-766.1997.

47. Rzomp, K.A., Scholtes, L.D., Briggs, B.J., Whittaker, G.R., and Scidmore, M.A. (2003). Rab
GTPases Are Recruited to Chlamydial Inclusions in Both a Species-Dependent and Species-
Independent Manner. Infect Immun *71*, 5855–5870. https://doi.org/10.1128/iai.71.10.5855.

48. Beck, M., and Hurt, E. (2017). The nuclear pore complex: understanding its function through
structural insight. Nat. Rev. Mol. Cell Biol. *18*, 73–89. https://doi.org/10.1038/nrm.2016.147.

49. Pritchard, C.E.J., Fornerod, M., Kasper, L.H., and Deursen, J.M.A. van (1999). RAE1 Is a
Shuttling mRNA Export Factor That Binds to a GLEBS-like NUP98 Motif at the Nuclear Pore
Complex through Multiple Domains. J. Cell Biol. *145*, 237–254.
https://doi.org/10.1083/jcb.145.2.237.

50. Schott, B.H., Antonia, A.L., Wang, L., Pittman, K.J., Sixt, B.S., Barnes, A.B., Valdivia,
R.H., and Ko, D.C. (2020). Modeling of variables in cellular infection reveals CXCL10 levels
are regulated by human genetic variation and the Chlamydia-encoded CPAF protease. Sci Rep-
uk *10*, 18269. https://doi.org/10.1038/s41598-020-75129-y.

51. Ouellette, S.P. (2018). Feasibility of a Conditional Knockout System for Chlamydia Based
on CRISPR Interference. Front. Cell. Infect. Microbiol. *8*, 59.
https://doi.org/10.3389/fcimb.2018.00059.

52. Gong, D., Kim, Y.H., Xiao, Y., Du, Y., Xie, Y., Lee, K.K., Feng, J., Farhat, N., Zhao, D.,
Shu, S., et al. (2016). A Herpesvirus Protein Selectively Inhibits Cellular mRNA Nuclear Export.
Cell Host Microbe *20*, 642–653. https://doi.org/10.1016/j.chom.2016.10.004.

53. Shen, Q., Kumari, S., Xu, C., Jang, S., Shi, J., Burdick, R.C., Levintov, L., Xiong, Q., Wu,
C., Devarkar, S.C., et al. (2023). The capsid lattice engages a bipartite NUP153 motif to mediate
nuclear entry of HIV-1 cores. Proc. Natl. Acad. Sci. *120*, e2202815120.
https://doi.org/10.1073/pnas.2202815120.

54. Dickson, C.F., Hertel, S., Tuckwell, A.J., Li, N., Ruan, J., Al-Izzi, S.C., Ariotti, N., Sierecki,
E., Gambin, Y., Morris, R.G., et al. (2024). The HIV capsid mimics karyopherin engagement of
FG-nucleoporins. Nature *626*, 836–842. https://doi.org/10.1038/s41586-023-06969-7.

55. Quan, B., Seo, H.-S., Blobel, G., and Ren, Y. (2014). Vesiculoviral matrix (M) protein
occupies nucleic acid binding site at nucleoporin pair (Rae1•Nup98). Proc. Natl. Acad. Sci. *111*,
9127–9132. https://doi.org/10.1073/pnas.1409076111.

56. Faria, P.A., Chakraborty, P., Levay, A., Barber, G.N., Ezelle, H.J., Enninga, J., Arana, C.,
Deursen, J. van, and Fontoura, B.M.A. (2005). VSV Disrupts the Rae1/mrnp41 mRNA Nuclear
Export Pathway. Mol. Cell *17*, 93–102. https://doi.org/10.1016/j.molcel.2004.11.023.

57. Miorin, L., Kehrer, T., Sanchez-Aparicio, M.T., Zhang, K., Cohen, P., Patel, R.S., Cupic, A.,
Makio, T., Mei, M., Moreno, E., et al. (2020). SARS-CoV-2 Orf6 hijacks Nup98 to block STAT
nuclear import and antagonize interferon signaling. Proc. Natl. Acad. Sci. *117*, 28344–28354.
https://doi.org/10.1073/pnas.2016650117.

58. Monette, A., Panté, N., and Mouland, A.J. (2011). HIV-1 remodels the nuclear pore complex.
J. Cell Biol. *193*, 619–631. https://doi.org/10.1083/jcb.201008064.

59. Ibana, J.A., Sherchand, S.P., Fontanilla, F.L., Nagamatsu, T., Schust, D.J., Quayle, A.J., and
Aiyar, A. (2018). Chlamydia trachomatis-infected cells and uninfected-bystander cells exhibit
diametrically opposed responses to interferon gamma. Sci. Rep. *8*, 8476.
https://doi.org/10.1038/s41598-018-26765-y.

60. Nunzio, F.D., Fricke, T., Miccio, A., Valle-Casuso, J.C., Perez, P., Souque, P., Rizzi, E.,
Severgnini, M., Mavilio, F., Charneau, P., et al. (2013). Nup153 and Nup98 bind the HIV-1 core
and contribute to the early steps of HIV-1 replication. Virology *440*, 8–18.
https://doi.org/10.1016/j.virol.2013.02.008.

61. Lad, S.P., Fukuda, E.Y., Li, J., Maza, L.M. de la, and Li, E. (2005). Up-Regulation of the
JAK/STAT1 Signal Pathway during Chlamydia trachomatis Infection. J. Immunol. *174*, 7186–
7193. https://doi.org/10.4049/jimmunol.174.11.7186.

62. Fontanilla, F.L., Ibana, J.A., Carabeo, R.A., and Brinkworth, A.J. (2024). Chlamydia
trachomatis modulates the expression of JAK-STAT signaling components to attenuate the type
II interferon response of epithelial cells. mBio *15*, e01834-24.
https://doi.org/10.1128/mbio.01834-24.

63. Haldar, A.K., Piro, A.S., Finethy, R., Espenschied, S.T., Brown, H.E., Giebel, A.M., Frickel,
E.M., Nelson, D.E., and Coers, J. (2016). Chlamydia trachomatis Is Resistant to Inclusion
Ubiquitination and Associated Host Defense in Gamma Interferon-Primed Human Epithelial
Cells. mBio *7*, e01417-16. https://doi.org/10.1128/mbio.01417-16.

64. Giebel, A.M., Hu, S., Rajaram, K., Finethy, R., Toh, E., 1, J.A.B., Morrison, S.G., Suchland,
R.J., Stein, B.D., Coers, J., et al. (2019). Genetic Screen in Chlamydia muridarum Reveals Role
for an Interferon-Induced Host Cell Death Program in Antimicrobial Inclusion Rupture. mBio
10,]e00385-19. https://doi.org/10.1128/mbio.00385-19.

65. Walsh, S.C., Reitano, J.R., Dickinson, M.S., Kutsch, M., Hernandez, D., Barnes, A.B.,
Schott, B.H., Wang, L., Ko, D.C., Kim, S.Y., et al. (2022). The bacterial effector GarD shields
Chlamydia trachomatis inclusions from RNF213-mediated ubiquitylation and destruction. Cell
Host Microbe *30*, 1671-1684.e9. https://doi.org/10.1016/j.chom.2022.08.008.

66. Morrison, R.P. (2000). Differential sensitivies of Chlamydia trachomatis strains to
inhibitory effects   of gamma interferon. Infect Immun *68*, 6038–6040.
https://doi.org/10.1128/iai.68.10.6038-6040.2000.

67. Fontanilla, F.L., Carabeo, R.A., and Brinkworth, A.J. (2024). Chlamydia trachomatis
modulates the expression of JAK-STAT signaling components to attenuate the Type II interferon
response of epithelial cells. bioRxiv, 2024.01.09.574898.
https://doi.org/10.1101/2024.01.09.574898.

68. Barta, M.L., Hickey, J., Kemege, K.E., Lovell, S., Battaile, K.P., and Hefty, P.S. (2013).
Structure of CT584 from Chlamydia trachomatis refined to 3.05 Å resolution. Acta Crystallogr.

Sect. F: Struct. Biol. Cryst. Commun. *69*, 1196–1201.
https://doi.org/10.1107/s1744309113027371.

69. Pha, K., Mirrashidi, K., Sherry, J., Tran, C.J., Herrera, C.M., McMahon, E., Elwell, C.A., and Engel, J.N. (2024). The Chlamydia effector IncE employs two short linear motifs to reprogram host vesicle trafficking. Cell Rep. *43*, 114624. https://doi.org/10.1016/j.celrep.2024.114624.

70. Pittner, N.A., Solomon, R.N., Bui, D.-C., and McBride, J.W. (2023). Ehrlichia effector SLiM-icry: Artifice of cellular subversion. Front. Cell. Infect. Microbiol. *13*, 1150758. https://doi.org/10.3389/fcimb.2023.1150758.

71. Mojica, S.A., Hovis, K.M., Frieman, M.B., Tran, B., Hsia, R., Ravel, J., Jenkins-Houk, C., Wilson, K.L., and Bavoil, P.M. (2015). SINC, a type III secreted protein of Chlamydia psittaci, targets the inner nuclear membrane of infected cells and uninfected neighbors. Mol Biol Cell *26*, 1918–1934. https://doi.org/10.1091/mbc.e14-11-1530.

72. Fleming, A., Sampey, G., Chung, M., Bailey, C., Hoek, M.L., Kashanchi, F., and Hakami, R.M. (2014). The carrying pigeons of the cell: exosomes and their role in infectious diseases caused by human pathogens. Pathog. Dis. *71*, 109–120. https://doi.org/10.1111/2049-632x.12135.

73. Jahnke, R., Matthiesen, S., Zaeck, L.M., Finke, S., and Knittler, M.R. (2022). Chlamydia trachomatis Cell-to-Cell Spread through Tunneling Nanotubes. Microbiol. Spectr. *10*, e02817-22. https://doi.org/10.1128/spectrum.02817-22.

74. Grieshaber, S.S., Grieshaber, N.A., Miller, N., and Hackstadt, T. (2006). Chlamydia trachomatis Causes Centrosomal Defects Resulting in Chromosomal Segregation Abnormalities. Traffic *7*, 940–949. https://doi.org/10.1111/j.1600-0854.2006.00439.x.

75. Scanlon, K.R., Keb, G., Wolf, K., Jewett, T.J., and Fields, K.A. (2023). Chlamydia trachomatis TmeB antagonizes actin polymerization via direct interference with Arp2/3 activity. Front. Cell. Infect. Microbiol. *13*, 1232391. https://doi.org/10.3389/fcimb.2023.1232391.

76. Chhabra, E.S., and Higgs, H.N. (2006). INF2 Is a WASP Homology 2 Motif-containing Formin That Severs Actin Filaments and Accelerates Both Polymerization and Depolymerization*. J. Biol. Chem. *281*, 26754–26767. https://doi.org/10.1074/jbc.m604666200.

77. Truong, D., Copeland, J.W., and Brumell, J.H. (2014). Bacterial subversion of host cytoskeletal machinery: Hijacking formins and the Arp2/3 complex. BioEssays *36*, 687–696. https://doi.org/10.1002/bies.201400038.

A.

Infection MS   Transfection MS

49   19   11

B.

Import of nuclear proteins | Shuttling protein and RNA | Export of nuclear RNAs

NUP107 subcomplex
NUP160
NUP133
NUP107
*NUP98
NUP96
NUP75
NUP43
NUP37
SEH1L
SEC13

Cytoplasmic NUPs
*RANBP2/NUP358
*NUP214
*NUP98
*NUP88
*NUP62

Transmembrane NUPs
NDC1
NUP210
*POM121/NUP121

Nup62 Subcomplex
*NUP62
*NUP58
*NUP54

NUP93 Subcomplex
NUP205
NUP188
*NUP155
*NUP93
NUP35

Nucleoplasmic NUPs
*NUP153
*NUP50
TPR

Import of nuclear proteins | Shuttling protein and RNA | Export of nuclear RNAs

C.

	Inputs				Elutes			

MW (kDa)

Uninfected | CebN-FLAG | TmcA-FLAG | Empty Vector | Uninfected | CebN-FLAG | TmcA-FLAG | Empty Vector

38 — 28 —   anti-FLAG IP

49 —   anti-NUP54

180 — 130 —   anti-NUP153

250 — 180 — 130 —   anti-NUP214

38 —   anti-Rae1

A.

	Uninfected	Full Length CebN	CebN 1-160aa	CebN 1-140aa	CebN 1-100aa	
MW (kDa)

28 —
14 —
anti-FLAG IP

300 —
250 —
180 —
anti-NUP214 Input

300 —
250 —
180 —
anti-NUP214 Elute

180 —
130 —
anti-NUP153 Input

180 —
130 —
anti-NUP153 Elute

49 — anti-NUP54 Input
49 — anti-NUP54 Elute

	Full Length CebN	CebN 1-160aa	CebN 1-140aa	CebN 1-100aa	
MW (kDa)

28 —
14 —
anti-FLAG IP

38 — anti-Rac1 Input

38 — anti-Rae1 Elute

B.

	FLAG	*C.t.* HSP-60	DAPI	Merge	Zoom
Full Length					
1-160 aa					
1-140 aa					
1-100 aa					

Uninfected Cell

WT *C.t.* Infected Cell

Bystander Cell