The tendon differentiation program is activated in human adipose stromal cells embedded

in a 3D-collagen matrix

Maxime Hordé[a], Jonathan Fouchard[a], Xavier Laffray[b], Cédrine Blavet[a], Véronique Béréziat[c],

Claire Lagathu[c], Ludovic Gaut [a], Delphine Duprez[a,1,#] and Emmanuelle Havis[a,1,#]

[a] Sorbonne Université, Institut Biologie Paris Seine, CNRS UMR7622, Developmental Biology

Laboratory, Inserm U1156, F-75005 Paris, France

[b] Université Paris Est Creteil, Glycobiology, Cell Growth and Tissue Repair Research Unit (Gly-

CRRET), EA 4397, F-94010 Creteil, France.

[c] Sorbonne Université, Inserm UMRS938, Centre de Recherche Saint-Antoine (CRSA), Institut

Hospitalo-Universitaire de Cardio-Métabolisme et Nutrition (ICAN), F-75012 Paris, France

[1] Co-senior authors; [#] Co-corresponding authors

Abstract

During development, tendon cell differentiation relies on molecular and mechanical parameters

that control the expression of tendon collagen and associated extracellular matrix proteins.

However, the minimal cues able to initiate tendon differentiation from progenitor cells remains

unknown. We analysed the molecular and mechanical parameters associated with tendon

differentiation in human adipose stromal cells (hASCs) in a 3D-culture system. We generated

3D-hASC constructs by embedding hASCs in a type-I collagen gel under a static uniaxial

geometrical constraint with no additional molecular and mechanical cues, and assessed tendon

matrix gene expression and mechanical properties for up to three weeks in culture. Analysis of

tendon matrix genes revealed a molecular progression consistent with the acquisition of a

tendon phenotype. The analysis of viscoelastic properties of 3D-hASC constructs by nano-

indentation indicated a progressive increase in tissue stiffness up to 10 kPa, concomitant with

a reduced stress relaxation indicative of solid-like mechanical properties. These changes in

mechanical properties were associated with transcription of tendon matrix components. In

summary, we have established that hASCs cultured in 3D-system with minimal mechanical

cues progress into the tendon differentiation program associated with variations of mechanical

properties reminiscent of embryonic tendon development.

Introduction

How a progenitor cell differentiates via the tendon lineage is not fully understood. The molecular and mechanical cues regulating the tendon program have not been completely identified.

Tendon is a key component of the musculoskeletal system that transmits forces from muscle to bone to allow locomotion. To accomplish this function, tendons exhibit specific viscoelastic properties, with a Young's modulus much higher than other soft tissues [1,2]. Tendon is a connective tissue composed of tendon fibroblasts that produce a dense extracellular matrix (ECM) consisting mainly of type I collagen fibrils that display a specific spatial organisation to resist the tensile forces generated by muscle contraction. Compared to other components of the musculoskeletal system, little is known about the mechanisms involved in tendon cell differentiation from undifferentiated cells. There is no identified master gene for the tendon lineage. The tendon program is difficult to follow because of the lack of recognized markers for each step of the tendon differentiation program [3]. Moreover, developmental tendon fibroblast populations are not well characterized and the molecular markers of tendon cell populations are not well identified [4].

In vertebrates, type I collagen, encoded by *COL1A1* and *COL1A2* genes, is the main structural and functional component of tendons. The regulation of *COL1A1* transcription is under the control of several transcription factors at different stages of tendon formation. The main developmental tendon marker, the bHLH transcription factor Scleraxis (SCX) [5] regulates *COL1A1* transcription during development [6]. Two other transcription factors have been shown to regulate *COL1A1* transcription in adult mouse tendons, EGR1 [7] and MKX (Mohawk) [8,9]. In addition to *COL1A1* regulation, SCX is required and sufficient for the expression of the transmembrane glycoprotein *TNMD* (Tenomodulin) in developing limb tendons and tendon cell cultures [6,10,11]; TNMD being involved in tenocyte proliferation and tendon collagen fibril maturation in new-born mice [12,13]. This indicates that SCX acts as a starter of the tendon program, although not being a master gene for the tendon lineage. Tendons are still observed in *Scx* mutant mice, however, force-transmitting tendons display severe alterations with disorganized matrix [6,14]. The transcriptomic analysis of *SCX*-positive cells during mouse limb tendon development provided us with a list of genes enriched in fetal tendon cells [15]. Although the function of all the 100 top genes have not been studied, when studied, gene function analyses show an involvement in tendon collagen fibrillogenesis; exampled with *THBS2*, coding for Thrombospondin 2 a secreted ECM glycoprotein [16], *COL6A3* encoding

the chainα3 of the non-fibrillar type VI collagen [17] and *COL14A1* encoding the chainα1 of

the FACIT (fibril-associated collagen with interrupted triple helices) [18]. Top 100 genes also

include various ECM proteins contributing to tendon biomechanical properties, such as the

small leucine-rich proteoglycan, DCN (Decorin) [19], the secreted matrix protein POSTN

(Periostin) [20,21], the collagen-bonding protein, DPT (Dermatopontin) [22–24] and the lysyl

oxidase enzyme LOX, which catalyzes collagens cross-linking of through oxidative

deamination of lysine residues [25]. All these tendon-associated molecules provide us with

molecular markers to assess the developmental tendon phenotype. In addition to molecular

cues, mechanical parameters are key for tendon development. Axial, head and limb tendon

development is stopped in the absence of muscle contraction in chicken and mouse models

[26,27].

Mesenchymal Stem Cells (MSCs) are a standard in the field of adult stem cell biology,

due to their high differentiation capacities in various cell lineages, including osteoblast,

adipocyte or chondrocyte. Although MSCs can also differentiate into the tendon lineage, the

progression into the tendon lineage is not completely understood. Tendon cell differentiation is

difficult to follow *in vitro*, since the tendon phenotype is not maintained in tendon 2D-cell

cultures. Stem cells require molecular stimuli to induce a tendon phenotype when cultured in

2D [7,28,29]. Compared to 2D-culture systems, 3D-culture systems are more favourable to

trigger tendon differentiation [29–31]. Human, rat, mouse and avian tendon fibroblasts have

been extensively used in 3D-culture systems to study collagen fibrillogenesis [32–37]. In a

fibrin 3D-culture system, chicken embryonic tendon cells were able to produce matrix genes

resembling to native tendons [38]. An alternative approach has been to use MSCs in 3D-culture

systems, which enable the study of the mechanisms by which the tendon lineage is initially

selected [31]. The tendon differentiation potential has been analysed in numerous MSC types,

such as bone marrow-derived Mesenchymal Stem Cells (bmMSCs), human Adipose-derived

Stromal Cells (hASCs)*,* mouse MSCs (with the C3H10T1/2 cell line)*.* However, these 3D-

culture systems made of MSCs require molecular cues [7,30,39–41] or mechanical loading

[7,42–45] to reach and maintain a tendon phenotype.

Here, we developed a 3D-culture system with hASCs in a type I collagen hydrogel

maintained between two anchor points to assess whether hASCs were capable of undergoing a

tendon differentiation program with no external molecular cues and mechanical stimulation for

three weeks. The increased expression of tendon genes, tendon-associated matrix genes,

combined with increased stiffness and reduced relaxation of 3D-hASC constructs, show that

hASCs are directed towards a tendon differentiation program. This shows that hASCs undergo towards a tendon phenotype in a minimal 3D-culture system with no external molecular or mechanical cues.

Results

Development of a 3D-culture system mimicking tendon formation from human Adipose Stromal Cells.

Since adipose tissue is an attractive alternative source of MSCs [46,47], we developed a 3D-culture system with hASCs isolated from the stromal vascular fraction of human adipose tissue collected after liposuction surgery [48,49]. Because collagen hydrogels recapitulate the native biophysical microenvironment of tendon tissues, we chose collagen hydrogel as biomaterial. During embryonic development, tendon primordia are high-cell-density linear structures anchored between bone and muscle in formation. To mimic tendon embryonic development, hASCs were homogenously embedded in a collagen gel deposited between two anchor points using the Flexcell technology (Supplementary Figure 1). This technology consists in a holder with 24 cylindrical moulds conferring a homogeneous and regular shape to the constructs. Each mould is perforated with vertical holes through which the vacuum is applied, allowing the silicon elastic surface of the well to acquire the shape of the mould (Fig.1A). At the extremities of each cylinder, two nylon anchor stems allowed the maintenance of static tension in each construct (Fig. 1A). After isolation and amplification, hASCs were embedded in a 3.5% bovine type I collagen solution and seeded between the two anchor stems. Vacuum was maintained for 2 hours to generate a construct of a cylindrical shape. 3D-hASC constructs (which we define as Day -2) were then left for 48 hours to allow cell development under static tension by condensing the surrounding matrix. 3D-constructs were analysed starting from this time point, defined as Day 0 (Fig. 1B). Once generated, the 3D-constructs were maintained under static conditions for 21 days (Fig. 1B). We did not apply external forces and the 3D-constructs were only subjected to the tension forces exerted between the two anchoring points. At Day 0, 3D-hASC constructs exhibited diameter values of 2.08+/-0.08 mm (Fig. 1D). The diameters of 3D-hASC constructs decreased drastically from Day 0 to Day 7 (0.67+/-0.02 mm at Day 7) and then displayed a slow decrease from Day 7 to Day 21 (0.59+/-0.02 mm at Day 14, 0.55+/-0.04 mm at Day 21) (Fig. 1D). Notably, because the length of the construct is fixed, we could estimate a volume reduction of 3D-constructs of about 15-fold in the course of the 21 Days of culture. 3D-no cell constructs not seeded with hASC exhibit constant diameter values during the 3 weeks of culture, 4.46+/-0.48 mm at Day 0, 4.41+/-0.45 mm at Day 7, 4.5+/-0.45 mm at Day 14 and 4.46+/-0.48 mm at

Day 21 (Supplementary Fig. 1D-E). Similarly, the cross-sectional areas of 3D-no cell constructs
remain constant and significantly higher than those of 3D-hASC constructs (Supplementary
Fig. 1F). The total number of cells in the constructs remained relatively stable between Day 0
and Day 21, with a slight increase at Day 7 and a slight decrease between Day 7 and Day 21
(Fig. 1D). We did not observe any apoptosis features between Day 7 and Day 14
(Supplementary Fig. 2).

At a cellular level, hASCs in this 3D-culture system displayed a fusiform shape and
spindle-shaped nuclei, aligned in the direction defined by the two anchor points (Fig. 1E),
reminiscent of cell organisation in native tendons, while hASCs exhibited a star shape in 2D
cultures (Supplementary Fig. 1B). In addition, cells progressively aligned along the tension axis
as the angle between the cell axis and the axis of the 3D-hASC constructs decreased throughout
culture (Fig. 1F). RT-qPCR analysis of the expression of the main tendon collagen *COL1A1*
show a progressive increase from Day 0 to Day 21 (Fig. 1G). In situ hybridization revealed that
COL1A1 was highly expressed in tendon constructs until Day 14 (Fig. 1H-I). However, at Day
21, *COL1A1* appeared to be more highly expressed in central cells than in ring cells (Fig. 1I).
We used this 3D-culture system to analyse the tendon differentiation potential of hASCs.

hASCs are prone to tendon differentiation in a 3D-culture system.
In order to assess the tendon differentiation potential of hASCs cultured in the 3D-conditions
described above, we analysed tendon gene expression at the transcript levels with RT-qPCR
and in situ hybridization experiments. We started the analyse with recognized tendon markers
at different time points during the culture. The mRNA expression of the following genes was
first analysed: the recognized tendon markers *SCX*, *MKX*, *TNMD* and *THSB2*. *SCX* expression
was detected, although at low level, both by in situ hybridization and RT-qPCR (Fig. 2A-B).
SCX expression slightly increased up to Day 7, remained constant up to Day 14 and then
decreased up to Day 21 (Fig. 2A). *MKX* expression was not detected at Day 0 and Day 2, but
started to be expressed on Day 4 and increased up to Day 21 (Fig. 2C). Interestingly, *TNMD*
was expressed in all 3D-hASC constructs on Day 4 but was not detected or at low levels at the
other timepoints (Fig. 2D). *THBS2* mRNA expression was significantly increased during human
3D-hASCs construct development (Fig. 2E). We also analysed the expression of other genes
identified in the Top100 genes (ordered with the highest enrichment score) of *SCX*-positive
cells during mouse limb development [15], such as *TM4SF1* (Transmembrane 4 superfamily
member 1, ranked 4th), *ANXA1* (Annexin A1, ranked 19th) and *S100A10* (member of the S100
family of cytosolic Ca^{2+}-binding proteins, ranked 73rd). *TM4FS1* expression was detected from

Day 0 and increased significantly in 3D-hASC constructs overtime (Fig. 2G). In contrast,

ANXA1 expression was first decreased at Day 2 and slightly increased at Day 21 (Fig. 2H). The

S100A10 expression levels were stable in 3D-hASC constructs until Day 14 and slightly

decreased between Day 14 and Day 21 (Fig. 2I). Consistent with the activation of the tendon

program, 3D-hASC constructs did not express osteogenic or adipogenic markers and did not

contain any adipocytes or osteocytes (Supplementary Fig. 3B-D). It has to be noted that hASCs

displayed the ability to differentiate into adipocytes and osteocytes in appropriate culture media

in 2D-culture conditions (Supplementary Fig. 3A-C) [47,50]. The cartilage-specific collagen

gene *COL2A1* was not detected 3D-hASC constructs (Fig. 2J), indicating that hASCs had not

differentiated into cartilage cells.

We conclude that hASCs undergo towards a tendon program when cultured in 3D-

conditions in a collagen gel maintained between two anchor points.

The Young's modulus of 3D-hASC constructs increases over time.

In order to assess whether the activation of the tendon program in hASCs was correlated with

a stiffening of the 3D-constructs, we measured the Young's modulus of 3D-hASC constructs

in the course of the 21 Days of culture. We used nanoindentation [51] to probe the mechanical

properties of the 3D-hASC constructs at the local scale (micron scale), with the application of

controlled deformations on the surface of the sample. To maintain the 3D-hASC constructs in

an immobile position during the nanoindentation experiments, they were detached from their

anchor points and deposited onto a support composed of grooves whose length and diameters

were adjusted according to constructs geometry. The support was maintained in a petri dish

with agarose gel, while 3D-hASC constructs were fixed to the support using dissection pins

inserted through the anchor stems and in the agarose gel (Fig. 3A). Indentation was performed

by moving the base of the cantilever at a constant speed of $10\mu m.s^{-1}$, with beads of diameters

ranged from 50 to 100 μm (Fig. 3B-D). The relationship between applied force and deformation

provides information about 3D-hASCs construct mechanical properties. In particular, tissue

effective Young's modulus E_{eff} was calculated by fitting the force-indentation curve over a one-

micron indentation range using the Hertz model (Fig. 3E, see methods). The effective Young's

modulus increased significantly during the 21 Days of construct culture starting from 1.3 kPa

at Day 0 to 9.2 kPa at Day 21 (Fig. 3F). Notably, E_{eff} increased by two-fold factor every seven

days. The effective Young's modulus of 3D-no cell constructs not seeded with cells remains

constant for 21 days, at 0.4 kPa (Fig. 3F).

We conclude that 3D-hASC constructs display an increased stiffness over time.

Extracellular matrix gene expression is correlated with changes in viscoelastic properties of
3D-hASC constructs.

Tendon stiffness relies on the synthesis of a dense and organized extracellular matrix generated by tendon cells. We thus analysed the expression of transcripts encoding tendon-associated collagens and fibrillogenesis-associated proteins during 3D-hASC constructs formation. Similarly to *COL1A1* (Fig. 1G), the mRNA levels of the tendon-associated collagens *COL6A3* and *COL14A1* were detected at Day 0, increased until Day 7 and remained higher at Day 21 compared to Day 0 (Fig. 4A-B). We also analysed the expression of genes encoding fibrillogenesis-associated proteins, *DPT*, *DCN*, *POSTN* and *LOX* which exhibit an elevated enrichment score in E14.5 compared to E11.5 *SCX*-positive cells during mouse limb development [52]. The transcript levels of *DPT*, *DCN*, *POSTN* and *LOX* were significantly increased during human 3D-hASC construct formation (Fig. 4C-F). The persistent increase of tendon collagen and collagen fibrillogenesis gene expression was indicative of matrix production and parallel the increase of effective Young's modulus (Fig. 3F) during 3D-hASCs construct formation. In addition, the increased expression of *DPT*, *DCN*, *POSTN*, *THBS2* and *LOX*, regulators of ECM network, also suggested that the structure and organisation of the ECM was modified during the course of 3D-hASC construct formation. We thus tested whether ECM transcript expression was accompanied with a modification of the viscoelastic properties of 3D-hASC constructs along time. To do this, we carried out stress relaxation experiments using nanoindentation (Fig. 5). In particular, we calculated the percentage of relaxation r_{relax} from the peak force F_{max} measured at the end of the indentation ramp and the force $F_{t=10s}$ measured after 10 seconds of relaxation ($r_{relax} = (F_{max}-F_{t=10s})/F_{max}$). The higher this ratio is, the more liquid-like the material. Oppositely, a smaller ratio indicates more solid, elastic-like viscoelastic properties. On Day 0, the percentage of relaxation of 3D-hASCs was 35% and decreased along time down to 24% on Day 21 (Fig. 5E), in parallel with the increase of ECM transcript levels. Interestingly, the percentage of relaxation of 3D-no cell constructs on Day 0 was 23%, decreased to 15% on Day 7 and remained constant (Fig. 5E).

We conclude that the 3D-hASC constructs acquire solid-like properties, consistent with the expression of ECM transcripts during the time course of the cultures.

3D-hASC constructs organize into two cell populations.

In order to analyse cell organisation, we performed longitudinal and transverse sections of 3D-hASC constructs (Fig. 1E, Supplementary Fig. 4, Fig. 6). We identified two cell populations,

one at the centre and the other one at the periphery as a ring, as soon as Day 2 (Fig. 6B). Phalloidin staining showed that the peripheral cell population formed a ring composed of 3 to 4 cell layers from Day 4 to Day 21, which was separated from the central region by a sharp interface (Fig. 6B). Cell density at the ring and the centre was maintained from Day 4 to Day 21 (Fig. 6C). Interestingly, the expression of ECM genes evolved in the two cell populations in a different manner (Fig. 1I, 2F). *COL1A1* was expressed in both centre and ring cells of 3D-hASC constructs with in situ hybridization from Day 7 to Day 14 (Fig. 1I). However, at Day 21, *COL1A1* displayed stronger expression in centre cells compared to ring cells (Fig. 1I). By contrast, *THBS2* was visualized in both centre and ring 3D-hASCs construct cells along the 21 days of culture (Fig. 2F). In addition, a marker of peripheral cells of adult rat tendons *COL3A1* [53] was detected both in centre and ring cells of 3D-hASC constructs, suggesting that the cells at the periphery were not mature peritenon cells (Supplementary Fig. 5).

We conclude that 3D-hASC constructs self-organize into cell populations with different location.

Discussion

This study showed that hASCs embedded into a cylindrical scaffold of type I-collagen attached to two anchor points leads to the expression of transcription factors and matrisome genes specific of tendon phenotype, and to the acquisition of viscoelastic properties reminiscent of tendon embryonic development.

Molecular progression of the tendon phenotype in hASCs in 3D-cultures

With no external molecular cues other than the culture medium, hASCs mixed in a 3D-collagen gel expressed tendon genes encoding transcription factors regulating *COL1A1* transcription and tendon matrix genes. The molecular progression in 3D-hASCs is reminiscent of tendon gene expression during development. The transcriptional regulators of *COL1A1* transcription, *SCX* and *MKX* are sequentially expressed during 3D-hASC constructs formation, as during mouse tendon development [8,9]. The expression pattern of *SCX* displays a bell shape, while the tendon matrix gene expression increases overtime in 3D-hASC constructs, consistent with SCX function as a starter of the developmental tendon program. The sequential expression of *SCX* and *TNMD* and the decrease of *SCX* and *TNMD* at Day 21 in the 3D-hASC constructs are consistent with *TNMD* regulation by SCX [6,11]. The two newly-identified genes, *TM4SF1* and *ANXA1,* coding for a Tetraspanin transmembrane protein and AnnexinA1, respectively, display

a regular increase overtime as the *LOX* gene (lysyl oxidase proteins). Interestingly, the *TM4SF1*, *ANXA1* and *LOX* genes have been identified in a fibroblast cluster for good prognosis in bladder cancer [54] and are seen as therapeutic targets for fibrosis [55–57]. ANXA1 and S100A10 belong to two families of proteins known to form hetero-tetramers [58], suggesting that ANXA1 interacts with S100A10 to maintain the tendon phenotype of 3D-hASC constructs. The mRNA expression levels of the tendon-associated collagen genes (*COL1A1*, *COL6A3*, *COL14A1)* and genes encoding proteins involved in tendon collagen fibrillogenesis (*THBS2*, *DCN, POSTN, DPT)* rise sharply during the formation of 3D-hASC constructs, and were maintained at high levels at the end of culture. These results indicate that the reduced expression of *SCX* in 3D-hASCs does not affect the molecular identity of tendon ECM in the constructs.

Our study shows that 3D-hASCs activate the tendon differentiation program, without dynamic mechanical stimulation or molecule addition, mimicking the development of embryonic tendons.

Progression of mechanical properties of hASCs in 3D-cultures

During the 21-day culture period, the 3D-hASCs showed reduced diameter, increased stiffness and reduced relaxation. The reduction in the diameter of 3D-hASC constructs was already observed in 3D-constructs using human dermal fibroblasts (HDF01035) or calf patellar tendon cells embedded in a collagen gel [59,60]. Because the diameters of 3D-no cell constructs without any cell remained constant, we conclude that hASCs were responsible for the collagen compaction of 3D-hASC constructs.

There is a linear increase of the stiffness of 3D-hASC constructs during the time of culture. Notably, E_{eff} increased by two-fold factor every seven days. This indicates that this long time-scale stiffening is unlikely to be due to ECM compaction, which occurred almost exclusively during the first week of culture. We rather believe that the expression of genes encoding specific ECM components would drive collagen compaction. After three weeks of cell culture, 3D-hASC constructs exhibited a Young's modulus values comprised between 5.6 and 18.3 kPa. Interestingly, measurements of nanoscale elastic modulus of chicken embryonic calcaneus tendons calculated similarly through Hertz contact theory gave results of same order of magnitude (5-17 kPa) [2,61]. This suggests that 3D-hASC constructs mimic the progression of mechanical properties observed during embryonic tendon development. In studies using human adult tendon fibroblasts (not MSCs) to generate 3D constructs, Young's modulus measurements lead to a value close to that found by tensile tests in adult tendons, of the order of MPa [32,35].

Link between the progression of ECM expression and mechanics in 3D-hASC constructs

The increase of expression of ECM-encoding genes parallels the stiffness increase and concomitant relaxation reduction of 3D-hASC constructs. We see this correlation as a bidirectional interaction between mechanical properties and ECM components during the time of the culture. It is recognized that changes of physical properties act on ECM composition and conversely, ECM interferes with mechanical properties [62]. We found that hASCs progressively acquired a spindle-shaped form associated with elongation of the cell shape during the course of the 3D-culture, as observed in native tendons. The elongated shape of cells observed in native tendons and 3D-hASC constructs is probably involved in tendon matrix production, since nuclear response upon deformation is recognized to include changes in transcriptional activity [63].

3D-hASC constructs self-organize into two cell populations, one being at the centre of the construct and the other forming a ring at the periphery. Such organisation has been previously described in 3D-constructs generated with the Flexcell technology either with hASCs [30] or avian tendon internal fibroblasts [64]. This cell organisation could be an effect of the geometry of the constructions defined by the anchoring points, which may orient cell shape and ECM. The peripheral cells layer of 3D-hASC constructs is reminiscent of peritenon surrounding native tendon. This is consistent with the lower *COL1A1* expression in the ring cells compared to centre cells in 3D-hASC constructs at Day 21 (Fig. 1I) observed in equine and rat peritenon cells versus tendon core cells [53,65]. *COL3A1* was identified recently as a peritenon marker in adult rat patellar tendons [53]. However, *COL3A1* was detected both at the centre and the periphery of 3D-hASC constructs (Supplementary Fig. 5), indicating that *COL31A1* expression does not discriminate the peritenon identity between ring and center cells in 3D-hASC constructs. Interestingly, a peripheral layer has been identified surrounding developing tendons in mouse and chicken embryos [4,15,66], indicating that future peritenon cells are present early during the tendon program.

Although the 3D-hASC constructs display a tendon matrisome gene expression profile at 3 weeks, we do not exclude that external mechanical cues would improve the tendon phenotype of 3D-hASC constructs as already reported [45,67]. During development, muscle contractions are recognized to maintain *SCX* expression and the tendon phenotype in developing tendons [26]. An interesting hypothesis is that application of external mechanical cues would prevent *SCX* and *TNDM* down-regulation in 21-day-old 3D-hASC constructs.

In summary, we have established that hASCs, cultured in 3D-system with minimal molecular and mechanical cues, progress into the tendon differentiation program along three weeks of culture. The molecular tendon program is associated with variations of mechanical properties reminiscent of embryonic tendon development.

Experimental procedures

Cell sample collection

Human abdominal subcutaneous adipose tissues were obtained from healthy donors after liposuction surgery (AP-HP Saint-Antoine Hospital, Paris, France). Human Adipose stromal cells (hASCs) were isolated from 3 healthy donors (one woman and two men).

All donors provided their written, informed consent to the use of their tissue specimens for research purposes. The study was performed in compliance with the principles of the Declaration of Helsinki and was approved by an institutional review board. The study was approved by the French regulatory authorities (Codecoh DC2023-5617). Cells were cryopreserved in 2mL cryotubes (2x106 cells per vial) stored in liquid nitrogen tanks.

Cell amplification

For each experiments, cell vials were thawed 2-3 min in a 37°C water bath and transferred in T25 cell culture flasks (Gibco) with 5ml of hASCs culture medium containing Minimum Essential Mediun (αMEM; Gibco; ref 22571-020) complemented with 10% Fetal Calf Serum (FCS; Gibco; ref 35-079-CV), 1% HEPES (1M, sigma; ref H0887-100ml), 1% glutamine (100X, 200mM, Gibco; ref 25030-024), 1% Penicillin/Streptomycin (Gibco; ref 15140-122) and 2.5 ng/mL FGF2 (PeproTech, Rocky Hill, NJ; ref 100-18B). Confluent cells were submitted to 2 passages, at 1:9 and 1:3 ratio. For each passage, we used 0.05% Trypsin (Gibco; ref 25300-054) for 5 minutes.

Preparation of 3D-hASC constructs

3D-hASC constructs were prepared as described [42]. Briefly, sub confluent cells were detached in 5 ml of 0.05% trypsin for 5 minutes, then suspended with hASC culture medium (described below) and counted with a Malassez counting chamber. 16.2 million cells (600,000 cells per 3D-hASC construct) were centrifuged at 1200 rpm for 5 minutes and resuspended in 4.05 mL of Type I collagen solution containing 70% Purecoll (Sigma-Aldritch, Oakville, Canada; ref 5074-35ml), 10% FCS and 20% αMEM brought to 7.4 pH with 0.1 N NaOH

immediately before use. 150 µl of hASCs-collagen mix was pipetted into each well of an

untreated TissueTrain© plate (FlexCell International, Hillsborough NC; ref TTLC-4001C)

according to the manufacturer's instructions. The collagen hydrogel was allowed to set for 2

hours, and the 3D-hASC constructs were covered with 2 ml of hASC culture medium completed

with 0.25 mM ascorbic acid. The medium was replaced every 2-3 Days.

3D-hASC construct sections

3D-hASC constructs were fixed in 1% paraformaldehyde (PFA, Gibco Sigma; ref 158127-

100G) overnight at 4°C, washed in 1X PBS (Gibco; ref :14190-094) and put in a 15% sucrose

solution at 4°C for 3 Days. Before inclusion, 3D-hASC constructs were incubated in a 7,5%

gelatin-15% sucrose solution in a 37°C water bath for 1 hour. Included 3D-hASC constructs

were then frozen at -80°C in isopentane for 1 minute and 12µm longitudinal and transversal

cryosections were generated with a CM3050 S Cryostat (Leica Biosystems).

Cell angle measurements

Longitudinal sections were stained with DAPI (D9542, Sigma-Aldrich) to visualize cell nuclei

and phalloidin (A12380 Alexa Fluor Phalloidin, Invitrogen/ ThermoFisher scientific) to

visualize cytoskeletal F-actin. Fluorescent images were captured using a Zeiss Axio Observer

Z1 microscope equipped with a Zeiss Apotome 2 and an Axiocam 506 monochrome camera.

Cell/nucleus angles relative to the axis of the 3D-hASC constructs were measured using Fiji

Image J software.

RNA isolation

Entire 3D-hASC constructs were put in 500µL RLT (Qiagen; ref 74106) and homogenized

using a mechanical disruption device (Lysing Matrix A, Fast Prep MP1, 4×30 s, 6 m.s^{-1}). Total

RNA was isolated using the RNeasy mini kit (Qiagen; ref 74106), according to the

manufacturer's instructions, including a 15 min of DNase I (Qiagen; ref 79254) treatment.

Reverse-Transcription and quantitative real time PCR

500ng of total RNAs extracted from each 3D-hASCs construct, for each timepoint, were

Reverse Transcribed using the High Capacity Retro-transcription kit (Applied Biosystems; ref

4387406). Quantitative PCR analyses were performed using primers listed in Supplementary

Table 1 and SYBR Green PCR Master Mix (Applied Biosystems; ref 4385614). The relative

mRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method [68]. The cycle threshold (C_T)

corresponds to the number of cycles required for the fluorescent signal to cross the threshold

(to exceed background level). The ΔC_Ts were obtained from C_T normalized to *YWHAZ* C_T level

in each sample. The primers used for RT-qPCR experiments are listed in Supplementary table

1.

In situ hybridization

3D-hASCs construct 12μm cryosections were subject to a standard in situ hybridization

protocol [69]. *COL1A1, SCX, THBS2 and COL3A1* and probes were generated using a

Roboprobe kit (Promega; ref P1460), according to manufacturer's instructions. Images were

captured using a Nanozoomer SQ Digital Slide Scanner with NDP.view2 software (Hamamatsu

Photonics). The primers used for probe synthesis are listed in Supplementary table 1.

Nanoindentation measurements

Live 3D-hASC constructs were detached from their anchor points and immediately fixed onto

a 3D-printed support. To avoid sliding of the construct during mechanical testing, the support

was designed with three half-cylindrical grooves at its surface whose diameters were adjusted

to the typical 3D-hASC constructs diameters. In order to conserve the original length of the 3D-

hASCs construct, the length of the support was also calibrated to the original distance between

the anchor points in the bioreactor. The support was then embedded in an agarose gel in a Petri

dish, and the samples were immersed in culture medium. Mechanical testing was performed at

room temperature, using a Piuma nano-indenter (Optics11 Life), as described in the main text.

Samples were positioned under the probe using a magnifying glass. Five measurements were

performed along the 3D-hASCs construct length and their mean value was calculated to extract

effective Young's modulus and percentage of stress relaxation for each sample.

The effective Young's modulus E_{eff} was calculated from a fit of the force versus indentation

curve over a 1μm indentation course and using a Hertzian contact model as follows:

444
$$F = \frac{4}{3} E_{eff} \sqrt{R_i} \cdot h^{3/2}$$

445 where F is the force, Ri the tip radius, and h the indentation.

447 *Statistical analyses*

448 Data was analysed using the non-parametric Mann-Whitney test with Graphpad Prism V10.

449 Results are shown as means ± standard deviations. The p-values are indicated either with the

450 value or with * or #.

Funding sources

 This work was supported by the CNRS, INSERM, Sorbonne University and ANR_TENORS.

Acknowledgements

We would like to thank Mathieu Hautefeuille for proofreading the manuscript.

References

[1] J. Peltonen, N.J. Cronin, L. Stenroth, T. Finni, J. Avela, Viscoelastic properties of the achilles tendon in vivo, Springerplus 2 (2013) 1–8. https://doi.org/10.1186/2193-1801-2-212.

[2] B.K. Connizzo, A.J. Grodzinsky, Tendon exhibits complex poroelastic behavior at the nanoscale as revealed by high-frequency AFM-based rheology, J Biomech 54 (2017) 11–18. https://doi.org/10.1016/j.jbiomech.2017.01.029.

[3] A.H. Huang, J.L. Galloway, Current and emerging technologies for defining and validating tendon cell fate, Journal of Orthopaedic Research 41 (2023) 2082–2092. https://doi.org/10.1002/jor.25632.

[4] E. Hirsinger, C. Blavet, M.A. Bonnin, L. Bellenger, T. Gharsalli, D. Duprez, Limb connective tissue is organized in a continuum of promiscuous fibroblast identities during development, IScience 27 (2024). https://doi.org/10.1016/j.isci.2024.110305.

[5] R. Schweitzer, J.H. Chyung, L.C. Murtaugh, A.E. Brent, V. Rosen, E.N. Olson, A. Lassar, C.J. Tabin, Analysis of the tendon cell fate using Scleraxis, a specific marker for tendons and ligaments, Development 128 (2001) 3855–3866. https://doi.org/10.1242/dev.128.19.3855.

[6] N.D. Murchison, B.A. Price, D.A. Conner, D.R. Keene, E.N. Olson, C.J. Tabin, R. Schweitzer, Regulation of tendon differentiation by scleraxis distinguishes force-transmitting tendons from muscle-anchoring tendons, Development 134 (2007) 2697 LP – 2708. https://doi.org/10.1242/dev.001933.

[7] M.-J. Guerquin, B. Charvet, G. Nourissat, E. Havis, O. Ronsin, M.-A. Bonnin, M. Ruggiu, I. Olivera-Martinez, N. Robert, Y. Lu, K.E. Kadler, T. Baumberger, L. Doursounian, F. Berenbaum, D. Duprez, Transcription factor EGR1 directs tendon differentiation and promotes tendon repair, Journal of Clinical Investigation 123 (2013). https://doi.org/10.1172/JCI67521.

[8] Y. Ito, N. Toriuchi, T. Yoshitaka, H. Ueno-Kudoh, T. Sato, S. Yokoyama, K. Nishida, T. Akimoto, M. Takahashi, S. Miyaki, H. Asahara, The Mohawk homeobox gene is a critical regulator of tendon differentiation, Proc Natl Acad Sci U S A 107 (2010) 10538–10542. https://doi.org/10.1073/pnas.1000525107.

[9] W. Liu, S.S. Watson, Y. Lan, D.R. Keene, C.E. Ovitt, H. Liu, R. Schweitzer, R. Jiang, The Atypical Homeodomain Transcription Factor Mohawk Controls Tendon Morphogenesis, Mol Cell Biol 30 (2010) 4797–4807. https://doi.org/10.1128/mcb.00207-10.

[10] C. Shukunami, A. Takimoto, M. Oro, Y. Hiraki, Scleraxis positively regulates the expression of tenomodulin, a differentiation marker of tenocytes, Dev Biol 298 (2006) 234–247. https://doi.org/10.1016/j.ydbio.2006.06.036.

[11] C. Shukunami, A. Takimoto, Y. Nishizaki, Y. Yoshimoto, S. Tanaka, S. Miura, H. Watanabe, T. Sakuma, T. Yamamoto, G. Kondoh, Y. Hiraki, Scleraxis is a transcriptional activator that regulates the expression of Tenomodulin, a marker of

mature tenocytes and ligamentocytes, Sci Rep 8 (2018) 3155.

https://doi.org/10.1038/s41598-018-21194-3.

[12] D. Docheva, E.B. Hunziker, R. Fässler, O. Brandau, Tenomodulin is necessary for

tenocyte proliferation and tendon maturation, Mol Cell Biol 25 (2005) 699–705.

https://doi.org/10.1128/MCB.25.2.699-705.2005.

[13] S. Dex, P. Alberton, L. Willkomm, T. Söllradl, S. Bago, S. Milz, M. Shakibaei, A.

Ignatius, W. Bloch, H. Clausen-Schaumann, C. Shukunami, M. Schieker, D. Docheva,

Tenomodulin is Required for Tendon Endurance Running and Collagen I Fibril

Adaptation to Mechanical Load, EBioMedicine 20 (2017) 240–254.

https://doi.org/10.1016/j.ebiom.2017.05.003.

[14] A.H. Huang, S.S. Watson, L. Wang, B.M. Baker, H. Akiyama, J. V Brigande, R.

Schweitzer, Requirement for scleraxis in the recruitment of mesenchymal progenitors

during embryonic tendon elongation, Development 146 (2019) dev182782.

https://doi.org/10.1242/dev.182782.

[15] E. Havis, M.-A. Bonnin, I. Olivera-Martinez, N. Nazaret, M. Ruggiu, J. Weibel, C.

Durand, M.-J. Guerquin, C. Bonod-Bidaud, F. Ruggiero, R. Schweitzer, D. Duprez,

Transcriptomic analysis of mouse limb tendon cells during development, Development

141 (2014) 3683–3696. https://doi.org/10.1242/dev.108654.

[16] T.R. Kyriakides, Y.H. Zhu, L.T. Smith, S.D. Bain, Z. Yang, M.T. Lin, K.G. Danielson,

R. V. Iozzo, M. LaMarca, C.E. McKinney, E.I. Ginns, P. Bornstein, Mice that lack

thrombospondin 2 display connective tissue abnormalities that are associated with

disordered collagen fibrillogenesis, an increased vascular density, and a bleeding

diathesis, Journal of Cell Biology 140 (1998) 419–430.

https://doi.org/10.1083/jcb.140.2.419.

[17] T.C. Pan, R.Z. Zhang, D. Markova, M. Arita, Y. Zhang, S. Bogdanovich, T.S.

Khurana, C.G. Bönnemann, D.E. Birk, M.L. Chu, COL6A3 protein deficiency in mice

leads to muscle and tendon defects similar to human collagen VI congenital muscular

dystrophy, Journal of Biological Chemistry 288 (2013) 14320–14331.

https://doi.org/10.1074/jbc.M112.433078.

[18] H.L. Ansorge, X. Meng, G. Zhang, G. Veit, M. Sun, J.F. Klement, D.P. Beason, L.J.

Soslowsky, M. Koch, D.E. Birk, Type XIV collagen regulates fibrillogenesis:

Premature collagen fibril growth and tissue dysfunction in null mice, Journal of

Biological Chemistry 284 (2009) 8427–8438. https://doi.org/10.1074/jbc.M805582200.

[19] G. Zhang, Y. Ezura, I. Chervoneva, P.S. Robinson, D.P. Beason, E.T. Carine, L.J.

Soslowsky, R. V. Iozzo, D.E. Birk, Decorin regulates assembly of collagen fibrils and

acquisition of biomechanical properties during tendon development, J Cell Biochem 98

(2006) 1436–1449. https://doi.org/10.1002/jcb.20776.

[20] J.E. Ackerman, S.N. Muscat, E. Adjei-Sowah, A. Korcari, A.E.C. Nichols, M.R.

Buckley, A.E. Loiselle, Identification of Periostin as a critical niche for myofibroblast

dynamics and fibrosis during tendon healing, Matrix Biology 125 (2024) 59–72.

https://doi.org/10.1016/j.matbio.2023.12.004.

[21] H. Li, A. Korcari, D. Ciufo, C.L. Mendias, S.A. Rodeo, M.R. Buckley, A.E. Loiselle,

G.S. Pitt, C. Cao, Increased Ca2+ signaling through CaV 1.2 induces tendon

hypertrophy with increased collagen fibrillogenesis and biomechanical properties,

FASEB J 37 (2023) e23007. https://doi.org/10.1096/fj.202300607R.

[22] O. Okamoto, S. Fujiwara, Dermatopontin, a Novel Player in the Biology of the

Extracellular Matrix, Connect Tissue Res 47 (2006) 177–189.

https://doi.org/10.1080/03008200600846564.

[23] A.J. de Micheli, J.B. Swanson, N.P. Disser, L.M. Martinez, N.R. Walker, D.J. Oliver,

B.D. Cosgrove, C.L. Mendias, Single-cell transcriptomic analysis identifies extensive

heterogeneity in the cellular composition of mouse Achilles tendons, Am J Physiol Cell
Physiol 319 (2020) C885–C894. https://doi.org/10.1152/ajpcell.00372.2020.

[24] T. Omoto, D. Yimiti, Y. Sanada, M. Toriyama, C. Ding, Y. Hayashi, Y. Ikuta, T.
Nakasa, M. Ishikawa, M. Sano, M. Lee, T. Akimoto, C. Shukunami, S. Miyaki, N.
Adachi, Tendon-Specific Dicer Deficient Mice Exhibit Hypoplastic Tendon Through
the Downregulation of Tendon-Related Genes and MicroRNAs, Front Cell Dev Biol 10
(2022) 1–14. https://doi.org/10.3389/fcell.2022.898428.

[25] P.K. Nguyen, K. Hall, I. Holt, C.K. Kuo, Recombinant lysyl oxidase effects on
embryonic tendon cell phenotype and behavior, Journal of Orthopaedic Research 41
(2023) 2175–2185. https://doi.org/10.1002/jor.25655.

[26] L. Gaut, D. Duprez, Tendon development and diseases, Wiley Interdiscip Rev Dev Biol
5 (2016) 5–23. https://doi.org/10.1002/wdev.201.

[27] E. Havis, M.-A. Bonnin, J. Esteves de Lima, B. Charvet, C. Milet, D. Duprez, TGFβ
and FGF promote tendon progenitor fate and act downstream of muscle contraction to
regulate tendon differentiation during chick limb development, Development 143
(2016) 3839–3851. https://doi.org/10.1242/dev.136242.

[28] K. Otabe, H. Nakahara, A. Hasegawa, T. Matsukawa, F. Ayabe, N. Onizuka, M. Inui,
S. Takada, Y. Ito, I. Sekiya, T. Muneta, M. Lotz, H. Asahara, Transcription factor
mohawk controls tenogenic differentiation of bone marrow mesenchymal stem cells in
vitro and in vivo, Journal of Orthopaedic Research 33 (2015) 1–8.
https://doi.org/10.1002/jor.22750.

[29] L. Gaut, M.A. Bonnin, C. Blavet, I. Cacciapuoti, M. Orpel, M. Mericskay, D. Duprez,
Mechanical and molecular parameters that influence the tendon differentiation potential
of C3H10T1/2 cells in 2D- And 3D-culture systems, Biol Open 9 (2020).
https://doi.org/10.1242/bio.047928.

[30] G. Yang, B.B. Rothrauff, H. Lin, R. Gottardi, P.G. Alexander, R.S. Tuan, Enhancement
of tenogenic differentiation of human adipose stem cells by tendon-derived
extracellular matrix, Biomaterials 34 (2013) 9295–9306.
https://doi.org/10.1016/j.biomaterials.2013.08.054.

[31] D. Jaiswal, L. Yousman, M. Neary, E. Fernschild, B. Zolnoski, S. Katebifar, S.
Rudraiah, A.D. Mazzocca, S.G. Kumbar, Tendon tissue engineering: Biomechanical
considerations, Biomedical Materials (Bristol) 15 (2020). https://doi.org/10.1088/1748-
605X/ab852f.

[32] A. Herchenhan, M.L. Bayer, R.B. Svensson, S.P. Magnusson, M. Kjær, In vitro tendon
tissue development from human fibroblasts demonstrates collagen fibril diameter
growth associated with a rise in mechanical strength, Developmental Dynamics 242
(2013) 2–8. https://doi.org/10.1002/dvdy.23896.

[33] A. Herchenhan, F. Uhlenbrock, P. Eliasson, M. Weis, D. Eyre, K.E. Kadler, S.P.
Magnusson, M. Kjaer, Lysyl oxidase activity is required for ordered collagen
fibrillogenesis by tendon cells, Journal of Biological Chemistry 290 (2015) 16440–
16450. https://doi.org/10.1074/jbc.M115.641670.

[34] Z. Kapacee, S.H. Richardson, Y. Lu, T. Starborg, D.F. Holmes, K. Baar, K.E. Kadler,
Tension is required for fibripositor formation, Matrix Biology 27 (2008) 371–375.
https://doi.org/10.1016/j.matbio.2007.11.006.

[35] A.M. Avey, O. Valdez, K. Baar, Characterization of an in vitro engineered ligament
model, Matrix Biol Plus 21 (2024). https://doi.org/10.1016/j.mbplus.2023.100140.

[36] M.L. Bayer, C.Y.C. Yeung, K.E. Kadler, K. Qvortrup, K. Baar, R.B. Svensson, S.
Peter Magnusson, M. Krogsgaard, M. Koch, M. Kjaer, The initiation of embryonic-like
collagen fibrillogenesis by adult human tendon fibroblasts when cultured under tension,
Biomaterials 31 (2010) 4889–4897. https://doi.org/10.1016/j.biomaterials.2010.02.062.

[37] N.S. Kalson, D.F. Holmes, Z. Kapacee, I. Otermin, Y. Lu, R.A. Ennos, E.G. Canty-
Laird, K.E. Kadler, An experimental model for studying the biomechanics of
embryonic tendon: Evidence that the development of mechanical properties depends on
the actinomyosin machinery, Matrix Biology 29 (2010) 678–689.
https://doi.org/10.1016/j.matbio.2010.08.009.
[38] C.Y.C. Yeung, L.A.H. Zeef, C. Lallyett, Y. Lu, E.G. Canty-Laird, K.E. Kadler, Chick
tendon fibroblast transcriptome and shape depend on whether the cell has made its own
collagen matrix, Sci Rep 5 (2015). https://doi.org/10.1038/srep13555.
[39] C.F. Hsieh, Z. Yan, R.G. Schumann, S. Milz, C.G. Pfeifer, M. Schieker, D. Docheva,
In vitro comparison of 2D-cell culture and 3D-cell sheets of scleraxis-programmed
bone marrow derived mesenchymal stem cells to primary tendon stem/progenitor cells
for tendon repair, Int J Mol Sci 19 (2018). https://doi.org/10.3390/ijms19082272.
[40] Z. Kapacee, C.Y.C. Yeung, Y. Lu, D. Crabtree, D.F. Holmes, K.E. Kadler, Synthesis of
embryonic tendon-like tissue by human marrow stromal/mesenchymal stem cells
requires a three-dimensional environment and transforming growth factor β3, Matrix
Biology 29 (2010) 668–677. https://doi.org/10.1016/j.matbio.2010.08.005.
[41] L. Gaut, N. Robert, A. Delalande, M.A. Bonnin, C. Pichon, D. Duprez, EGR1 regulates
transcription downstream of mechanical signals during tendon formation and healing,
PLoS One 11 (2016) 1–16. https://doi.org/10.1371/journal.pone.0166237.
[42] A. Scott, P. Danielson, T. Abraham, G. Fong, A. V. Sampaio, T.M. Underhill,
Mechanical force modulates scleraxis expression in bioartificial tendons, Journal of
Musculoskeletal Neuronal Interactions 11 (2011) 124–132.
[43] S. Pentzold, B. Wildemann, Mechanical overload decreases tenogenic differentiation
compared to physiological load in bioartificial tendons, J Biol Eng 16 (2022) 1–10.
https://doi.org/10.1186/s13036-022-00283-y.
[44] C.K. Kuo, R.S. Tuan, Mechanoactive tenogenic differentiation of human mesenchymal
stem cells, Tissue Eng Part A 14 (2008) 1615–1627.
https://doi.org/10.1089/ten.tea.2006.0415.
[45] H. Park, S.N. Nazhat, D.H. Rosenzweig, Mechanical activation drives tenogenic
differentiation of human mesenchymal stem cells in aligned dense collagen hydrogels,
Biomaterials 286 (2022) 121606. https://doi.org/10.1016/j.biomaterials.2022.121606.
[46] P. Zuk, Adipose-Derived Stem Cells in Tissue Regeneration: A Review, Int Sch Res
Notices 2013 (2013) e713959. https://doi.org/10.1155/2013/713959.
[47] L.E. Kokai, K. Marra, J.P. Rubin, Adipose stem cells: Biology and clinical applications
for tissue repair and regeneration, Translational Research 163 (2014) 399–408.
https://doi.org/10.1016/j.trsl.2013.11.009.
[48] V. Béréziat, C. Mazurier, M. Auclair, N. Ferrand, S. Jolly, T. Marie, L. Kobari, I.
Toillon, F. Delhommeau, B. Fève, A.K. Larsen, M. Sabbah, L. Garderet, Systemic
dysfunction of osteoblast differentiation in adipose-derived stem cells from patients
with multiple myeloma, Cells 8 (2019) 1–16. https://doi.org/10.3390/cells8050441.
[49] J. Gorwood, T. Ejlalmanesh, C. Bourgeois, M. Mantecon, C. Rose, M. Atlan, D.
Desjardins, R. Le Grand, B. Fève, O. Lambotte, J. Capeau, V. Béréziat, C. Lagathu,
SIV Infection and the HIV Proteins Tat and Nef Induce Senescence in Adipose Tissue
and Human Adipose Stem Cells, Resulting in Adipocyte Dysfunction, Cells 9 (2020).
https://doi.org/10.3390/cells9040854.
[50] M. Waldner, W. Zhang, I.B. James, K. Allbright, E. Havis, J.M. Bliley, A. Almadori,
R. Schweizer, J.A. Plock, K.M. Washington, V.S. Gorantla, M.G. Solari, K.G. Marra,
J.P. Rubin, Characteristics and immunomodulating functions of adipose-derived and
bone marrow-derived mesenchymal stem cells across defined human leukocyte antigen
barriers, Front Immunol 9 (2018). https://doi.org/10.3389/fimmu.2018.01642.

[51]   H. Van Hoorn, N.A. Kurniawan, G.H. Koenderink, D. Iannuzzi, Local dynamic
mechanical analysis for heterogeneous soft matter using ferrule-top indentation, Soft
Matter 12 (2016) 3066–3073. https://doi.org/10.1039/c6sm00300a.

[52]   E. Havis, M.-A. Bonnin, I. Olivera-Martinez, N. Nazaret, M. Ruggiu, J. Weibel, C.
Durand, M.-J. Guerquin, C. Bonod-Bidaud, F. Ruggiero, R. Schweitzer, D. Duprez,
Transcriptomic analysis of mouse limb tendon cells during development, Development
(Cambridge) 141 (2014). https://doi.org/10.1242/dev.108654.

[53]   D. Steffen, M. Mienaltowski, K. Baar, Spatial gene expression in the adult rat patellar
tendon, Matrix Biol Plus 19–20 (2023). https://doi.org/10.1016/j.mbplus.2023.100138.

[54]   J. Wang, Z. Tan, Y. Huang, C. Li, P. Zhan, H. Wang, H. Li, Integrating single-cell
RNA-seq to identify fibroblast-based molecular subtypes for predicting prognosis and
therapeutic response in bladder cancer, 2024. www.aging-us.comwww.aging-us.com.

[55]   Z. Yan, X. Cheng, T. Wang, X. Hong, G. Shao, C. Fu, Therapeutic potential for
targeting Annexin A1 in fibrotic diseases, Genes Dis 9 (2022) 1493–1505.
https://doi.org/10.1016/j.gendis.2022.05.038.

[56]   L. Chen, S. Li, W. Li, LOX/LOXL in pulmonary fibrosis: potential therapeutic targets,
J Drug Target 27 (2019) 790–796. https://doi.org/10.1080/1061186X.2018.1550649.

[57]   J. Zhao, C. Jing, R. Fan, W. Zhang, Prognostic model of fibroblasts in idiopathic
pulmonary fibrosis by combined bulk and single-cell RNA-sequencing, Heliyon 10
(2024). https://doi.org/10.1016/j.heliyon.2024.e34519.

[58]   S. Réty, D. Osterloh, J.-P. Arié, S. Tabaries, J. Seeman, F. Russo-Marie, V. Gerke, A.
Lewit-Bentley, Structural basis of the Ca 2+-dependent association between S100C
(S100A11) and its target, the N-terminal part of annexin I, 2000.

[59]   D. Schulz Torres, T.M. Freyman, I. V. Yannas, M. Spector, Tendon cell contraction of
collagen-GAG matrices in vitro: Effect of cross-linking, Biomaterials 21 (2000) 1607–
1619. https://doi.org/10.1016/S0142-9612(00)00051-X.

[60]   M.B. Vaughan, G. Xu, T.L. Morris, P. Kshetri, J.X. Herwig, Predictable fibroblast
tension generation by measuring compaction of anchored collagen matrices using
microscopy and optical coherence tomography, Cell Adh Migr 13 (2019) 303–314.
https://doi.org/10.1080/19336918.2019.1644855.

[61]   J.E. Marturano, J.D. Arena, Z.A. Schiller, I. Georgakoudi, C.K. Kuo, Characterization
of mechanical and biochemical properties of developing embryonic tendon,
Proceedings of the National Academy of Sciences 110 (2013) 6370 LP – 6375.
https://doi.org/10.1073/pnas.1300135110.

[62]   A. Saraswathibhatla, D. Indana, O. Chaudhuri, Cell-extracellular matrix
mechanotransduction in 3D HHS Public Access, Nat Rev Mol Cell Biol 24 (2023)
495–516. https://doi.org/10.1038/s415XX-XXX-XXXX-X.

[63]   Y.A. Miroshnikova, S.A. Wickström, Mechanical Forces in Nuclear Organization,
Cold Spring Harb Perspect Med 14 (2022).
https://doi.org/10.1101/cshperspect.a039685.

[64]   J. Garvin, J. Qi, M. Maloney, A.J. Banes, Novel System for Engineering Bioartificial
Tendons and Application of Mechanical Load, Tissue Eng 9 (2003) 967–979.
https://doi.org/10.1089/107632703322495619.

[65]   M.Y. Pechanec, T.N. Boyd, K. Baar, M.J. Mienaltowski, Adding exogenous biglycan
or decorin improves tendon formation for equine peritenon and tendon proper cells in
vitro, BMC Musculoskelet Disord 21 (2020). https://doi.org/10.1186/s12891-020-
03650-2.

[66]   S. Eloy-Trinquet, H. Wang, F. Edom-Vovard, D. Duprez, Fgf signaling components are
associated with muscles and tendons during limb development, Developmental
Dynamics 238 (2009) 1195–1206. https://doi.org/https://doi.org/10.1002/dvdy.21946.

[67] T. Wang, P. Chen, M. Zheng, A. Wang, D. Lloyd, T. Leys, Q. Zheng, M.H. Zheng, In
vitro loading models for tendon mechanobiology, Journal of Orthopaedic Research 36
(2018) 566–575. https://doi.org/10.1002/jor.23752.
[68] K.J. Livak, T.D. Schmittgen, Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−∆∆CT Method, Methods 25 (2001) 402–408.
https://doi.org/10.1006/meth.2001.1262.
[69] M.A. Bonnin, C. Laclef, R. Blaise, S. Eloy-Trinquet, F. Relaix, P. Maire, D. Duprez,
Six1 is not involved in limb tendon development, but is expressed in limb connective
tissue under Shh regulation, Mech Dev 122 (2005) 573–585.
https://doi.org/10.1016/j.mod.2004.11.005.
[70] M. Bléher, B. Meshko, I. Cacciapuoti, R. Gergondey, Y. Kovacs, D. Duprez, A.
L'Honoré, E. Havis, Egr1 loss-of-function promotes beige adipocyte differentiation
and activation specifically in inguinal subcutaneous white adipose tissue, Sci Rep 10
(2020) 1–15. https://doi.org/10.1038/s41598-020-72698-w.
712

714   **Figure Legends**

Figure 1: Generation of 3D-hASC constructs using a Flexcell bioreactor. A) Human
adipose stromal cells are isolated from lipoaspirates and amplified before seeding. The cells are
embedded in a 3.5% collagen hydrogel and deposited between the two anchor points. The
polymerization is performed by maintaining the vacuum and at 37°C for 2 hours. The vacuum
is then broken and the 3D-hASC constructs are kept in the incubator for 48 hours before starting
the experiments. Scale bar 100µm. B) 3D-hASC constructs were cultured for 21 days and C)
Diameters were measured from constructs at Day 0 (n=11), Day 7 (n=15), Day 14 (n=14) and
Day 21 (n=19). Each color represents a set of experiments. 4 independent experiments were
performed with 3<n<8 biological replicates for each experiment. The p-values were obtained
using the Mann-Whitney test. Asterisks * indicate the p-values of 3D-hASCs construct diameter
compared to each following stage, ****P<0.001. Scale bars 500µm. D) 3D-hASC construct
total cell number was counted at Day 0 (n=9), Day 7 (n=9), Day 14 (n=9) and Day 21 (n=9).
Each color represents a set of experiments. 3 independent experiments were performed with
n=3 biological replicates for each experiment. The p-values were obtained using the Mann-
Whitney test. Asterisks * indicate the p-values of 3D-hASCs construct cell number compared
to each following stage, ** P<0.01, ****P<0.0001. E) Longitudinal sections of 3D-hASC
constructs were performed at Day 4 and stained with DAPI/Phalloidin to visualize cellular and
nuclear shape. Scale bars: 50 µm. F) Longitudinal sections of 3D-hASC constructs were
performed on Day 0, Day 2, Day 4, Day 7, Day 14 and Day 21. The angles of 15 cells were
measured on a minimum of 3 constructs per time point and compared to the axis of tension.
Each dot represents the mean of 15 values per 3D-hASC construct. Asterisks * indicate the p-
values of 3D-hASCs construct cell number compared to each following stage, * P<0.05,
**P<0.01. G) After 0, 2, 4, 7, 14 and 21 Days of culture, 3D-hASC constructs were used for
mRNA purification. mRNA expression levels of *COL1A1* were analysed by RT-qPCR. H) 3D-
hASC constructs at Day 4 were transversally cryo-sectioned. 12 µm sections were hybridized
with the DIG-labeled antisense probes for *COL1A1*. Scale bar: 250µm. I) 3D-hASC constructs
at Day 4 Day 0, Day 7, Day 14 and Day 21 were transversally cryo-sectioned. 12 µm sections
were hybridized with the DIG-labeled antisense probes for *COL1A1*. Scale bars: 50µm.
743

744 **Figure 2: 3D-hASC constructs express tendon genes.** A) After 0, 2, 4, 7, 14 and 21 Days of
745 culture, 3D-hASC constructs were used for mRNA purification. mRNA expression levels of
746 *SCX* were analysed by RT-qPCR. B) 3D-hASC constructs at Day 0, Day 7, Day 14 and Day 21

were transversally cryo-sectioned. 12 μm sections were hybridized with the DIG-labeled antisense probes for *SCX*. Scale bars: 50μm. After 0, 2, 4, 7, 14 and 21 Days of culture, 3D-hASC constructs were used for mRNA purification. mRNA samples were analysed by RT-qPCR for C) *MKX* D) *TNMD* and E) *THBS2*. F) 3D-hASC constructs at Day 0, Day 7, Day 14 and Day 21 were transversally cryo-sectioned. 12 μm sections were hybridized with the DIG-labeled antisense probes for *THBS2*. Scale bars 50μm. mRNA expression levels of G) *TM4FS1*, H) *ANXA1*, I) *S100A10*, J) *COL2A1* were analysed by RT-qPCR. Transcripts are shown relative to the level of *YWHAZ* transcripts. The relative mRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method, with control being normalized to 1. Each color represents a set of experiments. 8 independent experiments were performed with 14 < n < 29 3D-hASCs construct generated for each step (n=29 at Day 0, n=19 at Day 2, n=18 at Day 4, n=25 at Day 7, n=17 at Day 14 and n=16 at Day 21). The C_T values shown for tendon genes and *YWHAZ* represent the average of all C_T values obtained for each time point. Error bars represent the mean + standard deviations. The p-values were obtained using the Mann-Whitney test. Asterisks * indicate the p-values of gene expression levels in 3D-hASC constructs compared to the first Day of gene detection *P<0.05, **P<0.01, *** P<0.001, **** P<0.0001.

Figure 3: 3D-hASC construct Young's modulus increased along time. A) 3D-hASC constructs are fixed on a 3D-printed support maintained in agarose gel. B) 3D-hASC constructs are placed in an appropriate channel according to their diameter and fixed using dissection pins inserted into the anchor stems and agarose. C) The one-piece optical probe of the nanoindenter is composed of an optical fiber, a cantilever and a spherical tip. After an approach phase, the tip touches the sample and indent it. The cantilever bending is detected by the optical fiber. D) Force as a function of cantilever base displacement for a typical indentation. Inset: Cantilever base displacement is applied at constant rate of 10μm.s-1. E) Force as a function of indentation for a typical indentation curve. Hertz contact fit (red) is applied over a 1 μm indentation. F) Evolution of effective Young's modulus along culture time in 3D-constructs without cells and seeded with hASCs. For 3D-no cell constructs without cells, each dot represents the mean of five measurements performed on one construct. Indentation experiments were performed at Day 0 (n=8), Day 7 (n=6), Day 14 (n=6) and Day 21 (n=12). Each color represents a set of experiments. 2 independent experiments were performed with 3<n<6 biological replicates for each experiment. For 3D-hASC constructs, each dot represents the mean of five measurements performed on one 3D-hASCs construct. Indentation experiments were performed at Day 0 (n=17), Day 7 (n=17), Day 14 (n=13) and Day 21 (n=14). Each color represents a set of experiments. 5 independent experiments were performed with 3<n<8 biological replicates for each experiment. The p-values were obtained using the Mann-Whitney test. Asterisks * indicate the p-values of Young's modulus levels (kPa) in 3D-hASC constructs compared to each following stage, ***P<0.001, ****P<0.0001. [#] indicates the p-values of Young's modulus levels (kPa) in 3D-no cell constructs compared to 3D-hASC constructs, [###]P<0.001, [####]P<0.0001.

Figure 4: Extra-cellular matrix gene expression was highly increased during 3D-hASC construct development. After 0, 2, 4, 7, 14 and 21 Days of culture, 3D-hASC constructs were used for mRNA purification. mRNA expression levels of tendon-associated collagens A) *COL6A3*, B) *COL14A1* and extracellular matrix protein coding genes C) *DCN*, D) *POSTN,* E) *DPT* and F) *LOX* were analysed by RT-qPCR. Transcripts are shown relative to the level of *YWHAZ* transcripts. The relative mRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method, with control being normalized to 1. Each color represents a set of experiments. 8 independent experiments were performed with 14 < n < 29 3D-hASCs construct generated for each step (n=29 at Day 0, n=19 at Day 2, n=18 at Day 4, n=25 at Day 7, n=17 at Day 14 and n=16 at Day

797 21). The C_T values shown for tendon genes and *YWHAZ* represent the average of all C_T values
798 obtained for each time point. Error bars represent the mean + standard deviations. The p-values
799 were obtained using the Mann-Whitney test. Asterisks * indicate the p-values of gene
800 expression levels in 3D-hASC constructs compared to the first Day of gene detection *P<0.05,
801 **P<0.01, *** P<0.001, **** P<0.0001.

Figure 5: Viscoelastic properties of human 3D-hASC constructs change over the course
of the culture. A) Schematic of a force relaxation experiment. In open loop, the base of the
cantilever is fixed while relaxation takes place. B) Typical curve showing force as a function
of indentation during loading, relaxation and unloading phase. Inset: Indentation as a function
of time. C) Force as a function of cantilever base displacement for a typical indentation. Inset:
Cantilever base displacement is applied at constant rate of 10µm.s-1. D) Force as a function
time. Force peaks at a value Fmax and reaches is close to a plateau after 10s of relaxation.
Frelax is defined as the force at this instant. E) Percentage of stress relaxation ((Fmax – Ft=10
sec) / Fmax)) x100 during the first 10 seconds of relaxation, after different periods of cell
culture. For 3D-no cell constructs without cells, each dot represents the mean of five
measurements performed on one construct. Indentation experiments were performed at Day 0
(n=8), Day 7 (n=6), Day 14 (n=6) and Day 21 (n=12). Each color represents a set of
experiments. 2 independent experiments were performed with 3<n<6 biological replicates for
each experiment. For 3D-hASC constructs, indentation experiments were performed at Day 0
(n=17), Day 7 (n=17), Day 14 (n=13) and Day 21 (n=14). Each color represents a set of
experiments. 5 independent experiments were performed with 3<n<8 biological replicates for
each experiment. The p-values were obtained using the Mann-Whitney test. Asterisks *
indicates the p-value of 3D-hASCs construct relaxation compared to each following stage;
*P<0.05; [@] indicates the p-value of 3D-no cell construct relaxation compared to each following
stage, [@]P<0.05; [#] indicates the p-value of in 3D-no cell constructs compared to 3D-hASC
constructs, [###]P<0.001, [####]P<0.0001.

Figure 6: Cellular organization in 3D engineered 3D-hASCs. A) 3D-hASCs transverse
section representation. B) Transverse sections of 3D-hASC constructs were performed on Day
0, Day 2, Day 4, Day 7, Day 14 and Day 21 of culture and stained with DAPI/Phalloidin to
visualize cell organization within the 3D-hASC constructs. From Day 2, 3D-hASC constructs
are composed of two cell populations, one at the centre and the other at the periphery, forming
a ring. D) Cell density was counted at the centre and the ring of 3D-hASC constructs at Day 0
(n=18), Day 2 (n=12), Day 4 (n=10), Day 7 (n=15), Day 14 (n=12) and Day 21 (n=9). The p-
values were obtained using the Mann-Whitney test compared to each following stage. Asterisks
* indicate the p-values of cell number at the centre, *** P <0.001, **** P<0.0001; # indicate
the p-values of cell number at the ring, #### P <0.0001; Scale bars 100µm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6