A complete map of specificity encoding enables reprogramming of a dynamic protein interaction

Taraneh Zarin[1], Cristina Hidalgo-Carcedo[1], and Ben Lehner[1,2,3,4]

[1]Centre for Genomic Regulation (CRG), Barcelona Institute for Science and Technology (BIST), Barcelona, Spain
[2]Wellcome Sanger Institute, Cambridge, UK
[3]Universitat Pompeu Fabra (UPF), Barcelona, Spain
[4]Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

Abstract:

Thousands of human proteins function by binding short linear motifs embedded in intrinsically disordered regions that can sample different conformations. How affinity and specificity are encoded in these regions and the relative contribution of structured and dynamic binding modes is not completely understood. Here we present a complete map of specificity encoding in a PDZ domain binding to a structurally heterogeneous disordered ligand and use this map to reprogram binding to new ligands. Quantifying >200,000 energetic interactions between the PDZ domain and its ligand identifies 17 residues that control specificity for six sites in the ligand through both direct and allosteric mechanisms. Most sites where specificity can be controlled through the domain are in the structured part of the ligand, but dynamic residues additively contribute to binding affinity and communicate with structured ligand residues to enable specificity changes. Combining mutations in each of the six PDZ domain modules allows specificity to be quantitatively reprogrammed to recognize new ligands. This approach of quantifying the specificities of thousands of proteins in a single experiment should allow many different types of molecular interaction to be understood and interpretably reprogrammed.

Introduction:

Specific physical interactions between proteins underlie nearly all aspects of biology from transcription and signaling to mechanics and neuronal information processing[1]. Many protein-protein interactions are mediated by intrinsically disordered regions (IDRs) of proteins which do not form stable secondary and/or tertiary structures[2,3]. While some IDRs adopt conditional structures upon binding to their partners[4–6], others remain dynamic upon binding, forming so-called 'fuzzy' complexes[7,8]. The ensemble of conformations afforded by dynamic binding has been shown to facilitate diverse regulatory functions in eukaryotes [7,9–11] but how dynamic binding and conformational heterogeneity contribute to binding affinity and/or specificity is not well understood[8,12–15].

One of the most frequent modes of protein-protein interaction in human cells is the binding of a globular domain to a short linear motif embedded within an IDR[3,16,17]. These peptide recognition

domains typically exist in large protein families and have affinity for many potential binding targets[18,19]. How binding specificity is encoded in these domains and how these large protein families evolve without causing interaction cross-talk is a major area of interest[20] and is crucial for understanding human disease and drug development[21–23]. PDZ (postsynaptic density 95, PSD-95; discs large, Dlg; zonula occludens-1, ZO-1) domains, for example, are the largest family of human protein interaction domains with more than 270 PDZ domains in 155 human proteins[19]. These domains typically bind to IDRs at the C-termini of proteins[18,19,21]. Upon binding, the last four amino acids of a PDZ ligand adopt a well-defined structure[24] and have historically been used to classify PDZ domains into distinct groups based on the identity of the P0 (C-terminal) and P-2 positions[25]. The adjacent residues of the IDR however typically remain dynamic in the complex[26], with X-ray structures revealing progressively increasing motion in the bound complex moving away from the C-terminus (Fig. 1a). While these adjacent N-terminal residues have not been nearly as well-studied, there is evidence that they are also important for binding[21,26,27], consistent with an emerging theme that the context around motif consensus sites is important for function[28,29]. Moreover, NMR (nuclear magnetic resonance), statistical coupling, and double mutant cycle experiments have quantified dynamic allosteric networks within PDZ domains[23,30–34], showing for example that the presence of an extra-domain alpha helix in one PDZ domain increases binding affinity for the ligand by 21-fold without any evidence for direct contact between the ligand and the alpha helix[30]. Several mechanisms[23,30,35] have been proposed for this change in affinity. The PDZ domain-peptide interaction thus presents an elegant model system in which to understand not only how affinity and specificity are encoded, but also how this encoding is distributed between structured versus dynamic binding modes.

Here we present a complete map of how binding affinity and specificity are encoded in a globular protein domain interacting with a disordered peptide. Our map reveals that specificity encoding is modular, with distinct residues determining specificity for each position in the ligand by both direct and allosteric mechanisms. The dynamic tail of the ligand is more robust to mutation but can be used to additively tune affinity and interacts with the stable part of the peptide to enable a specificity change. Combining mutations in the specificity modules allows reprogramming of the PDZ domain to bind a ligand with all six ligand positions mutated to different amino acids.

Results:

Quantifying a combinatorial genetic landscape for peptide recognition

The third PDZ domain (PDZ3) from PSD-95 binds to the C-termini of proteins matching the consensus motif -X-S/T-X-Φ-COOH (where X is any amino acid, aa, and Φ is a hydrophobic residue)[22,36,37]. The bound peptide is structured at the C-terminus but increasingly dynamic before the last four residues (Fig. 1b), a general property of PDZ domains bound to short ligands (Fig. 1a). To better define the binding specificity of PDZ3 for the canonical structured C-

terminal part of the ligand, we quantified its binding to >100,000 variants of a 9 aa peptide from the CRIPT protein in which the sequence of each of the last four aa (positions 0 to -3) was fully randomized to any of the 20 aa (Fig. 1d, Fig. S1). Binding was quantified using a highly-validated protein fragment complementation assay (PCA)[38] (Fig. 1c) and binding scores were very reproducible between replicate experiments (Fig. S1b-c, e, h-i). The position-weight matrix for the top 1% of binders matched the reported consensus[22,36,37] from a previous selection experiment[22] (Fig. 1f). The distribution of binding fitness (Fig. 1e) and proportion of peptides binding to PDZ3 (Fig. S1k) sharply decreased with an increasing number of substitutions. However, because of the exponential increase in the size of sequence space when combining mutations (Fig. S1j), there are actually more peptides with 4 aa changes that bind PDZ3 than peptides with one mutation (179 vs 73; 4.8-fold more, Fig. 1g). Position-weight matrices for peptides containing between one and four mutations show that the preference for T/S at the -2 position and hydrophobic residues (including cysteine) at the 0 position is maintained with increasing mutation order (Fig. 1f-g).

Additive energy models accurately predict peptide binding

Precisely quantifying the binding of PDZ3 to >100,000 peptides provides an opportunity to evaluate the extent to which binding to each of the four C-terminal residues is independent of sequence variation at the other three sites. We used MoCHI[39] to fit a two-state thermodynamic model to our binding data. The model accounts for the non-linear relationship between the Gibbs free energy of binding (dG) and the fraction of ligand bound to PDZ3 but otherwise assumes that the energetic effects of mutations (ddG) combine additively with no pairwise or higher order energetic couplings between mutations at different sites (Fig. 2a and Methods). Fitted to a balanced set of >7,000 binding and non-binding genotypes, the model provides good predictive performance (R^2=0.75 evaluated by 10-fold cross validation, Fig. 2b-iii). Not accounting for the non-linear relationship between the fraction of ligand bound and the binding energy resulted in worse predictive performance (R^2=0.56, Fig. S3c) and systematically biased predictions (Fig. S3c residuals). Moreover, allowing energetic couplings between mutations in different sites did not improve predictive performance (Fig. S3c). The additive energy model represents a very large compression of the binding data (>1000-fold, 110,000 genotypes/76 model coefficients) and formally shows that the effects of mutations in the C-terminus of the ligand have largely energetically independent effects on binding.

Dynamic residues are important for binding affinity

We next focused on the N-terminal residues in the CRIPT peptide (position -4 to -7) which display a progressive increase in movement in the PDZ3-CRIPT complex (Fig. 1a-b). To test the importance of these more dynamic residues to binding, we quantified the binding of >90,000 variants of CRIPT in which the four aa before the consensus binding motif were randomized to all other aa (positions -4 to -7) (Fig. 1d-e). Binding measurements were well correlated across independent experiments as with the C terminus (Fig. S1a, d, f-g). The distribution of binding scores was, however, very different to that of the C-terminus library (Fig. 1e), with a more gradual decrease in binding with an increasing number of substitutions (Fig. S1k). Thus,

although mutations in these four aa typically have smaller effects on PDZ3 binding than mutations in the C-terminal four aa, the region still makes an important contribution to binding affinity, with a clear preference for positively charged and aromatic residues at the -4 and -5 positions (Fig. 1f), and the combined effects of multiple substitutions frequently being very detrimental (Fig. 1g, S1k).

Fitting an additive thermodynamic model to the data, we again found that mutations have largely independent energetic effects (R^2=0.73 by 10-fold cross-validation, Fig. 2b-i). Ignoring the non-linear relationship between free energy (dG) and binding again resulted in biased predictions (Fig. S2c). Allowing energetic couplings between mutations provided almost the same predictive performance as the additive model (R^2=0.76 Fig. S2c). Therefore, as for mutations in the C-terminus, the effects of mutations in this more dynamic region of the ligand are largely energetically independent.

The mutational energy matrix for PDZ3 binding

The combined energy matrices for positions 0 to -3 and -4 to -7 provide a complete description of the energetic effects of substitutions in all eight positions of the PDZ3 ligand (Fig. 2c, Fig. S2d-e). Substitutions in positions 0 and -2 are most detrimental for binding, consistent with the description of the consensus motif as X-S/T-X-Φ-COOH[22,25,36]. However, not all substitutions in positions 0 and -2 have the same energetic effects and many changes at positions -1 and -3 also cause large changes in binding energy. Moreover, mutations at position -4, which is outside of the canonical motif, have the third largest energetic effects and multiple substitutions in positions -5 and -7 cause detrimental (ddG>0) energy changes (Fig. 2c).

The energy matrix also reveals that CRIPT harbors sites where many mutations are energetically favorable (ddG<0), particularly in the first five positions (Fig. 2c). For example, E is energetically favored at position -3; W, F, R and H are all favored at position -5; K and R favored at -6; and R is favored at -7 (Fig. 2c). Indeed, in the first four positions but not in the C-terminal four positions, the net charge of the peptide is a strong predictor of binding strength (Spearman's rho=0.57 for N, -0.04 for C, Fig. 1h, Fig. S1l), with positively charged peptides having higher binding scores. The ability of many substitutions in the N-terminal region of the peptide to increase affinity further emphasizes the importance of this region for binding, with the longer range of electrostatic interactions[40] compared to other non-covalent interactions consistent with a dynamic or 'fuzzy' contribution[8] to affinity.

Interactions between dynamic and structured regions of a disordered ligand

We next quantified binding of a library with all possible double mutants across the entire peptide to test whether the dynamic tail interacts energetically with the structured peptide region (Fig. 3a). This library allowed us to quantify >8,000 energetic couplings between mutations in all positions of the peptide (Fig. 3b-c, Fig. S3e-f). Once again, we found excellent reproducibility

across replicate experiments and libraries (Fig. S3a-c) and used shared variants to normalize fitness measurements across experiments (Fig. S3d).

To quantify interactions between positions, we measured the residuals to an additive two-state thermodynamic model for binding (Fig. 3d, Fig. S3e-f). In total, we identified 373 non-zero energetic couplings between pairs of mutations across the CRIPT peptide (Z-test, Benjamini-Hochberg FDR<0.1, Fig. S3g). We define a positive intra-peptide energetic interaction, or change in specificity, when the observed binding has a positive and significant (Z-test, FDR<0.1) residual to our additive energy model (Fig. S3i) and the observed binding score is significantly different from the binding score distribution of STOP codon variants) (Fig. S3h-i). We then identified pairs of sites in CRIPT that are enriched for these positive energetic couplings (Fig. 3e). This identified a significant coupling (FDR<0.1, hypergeometric test) between the structured and dynamic parts of the peptide, between the -3 and -5 positions (Fig. 3e). This pair of sites contains more than 20 significant (FDR<0.1, Z-test) positive energetic couplings (Fig. 3f), most of which are linked to a change in position -3 to a G residue. Substitution to a G at -3 changes the peptides bound by PDZ3 so that any residue that is not aromatic at the -5 position is now more strongly bound than expected (Fig. 3f).

A change in the sequence in the structured region of the peptide therefore changes the binding preference at a more dynamic region. Reciprocally, this energetic coupling means that specific mutations in a more dynamic residue alter the binding at a structured residue of the peptide.

Quantifying >200,000 energetic interactions between a PDZ domain and its ligand

We next designed an experiment to understand how the binding specificity of PDZ3 is encoded in the globular protein domain (Fig. 4a-b, Fig. S4). We performed a comprehensive set of double mutant cycles, mutating every position in PDZ3 and CRIPT to every other aa alone and in *trans*-double mutant combinations and measured binding (Fig. 4c). The binding measurements were highly correlated with measurements from the combinatorial CRIPT mutagenesis libraries (R^2=0.937 and 0.875 for N and C, n=36 and n=68, respectively, Fig. S1f-i), allowing us to normalize binding scores across independent experiments (Methods). Consistent with previous data[41,42], mutations in PDZ3 detrimental for binding are strongly enriched in the binding interface with CRIPT (Fig. 4d-e).

In total this dataset precisely defines the binding specificity of >1,800 globular proteins: we quantified the binding specificity of nearly every mutation in PDZ3 for nearly every mutation at every position in the ligand (Fig. S4f). Formally, changes in specificity are identified when there is an energetic coupling (or genetic interaction) between a mutation in the PDZ domain and a mutation in the ligand i.e. when binding is not well predicted by an additive energy model (Fig. 4f). The binding scores in this much larger experiment were also extremely well correlated across three replicate selections (Pearson's r>0.91, Fig. S4d), as were the inferred free energy changes (ddGs, Fig. S4g), allowing us to quantify >200,000 energetic couplings between mutations in PDZ3 and its ligand.

Hundreds of mutations in a PDZ domain change its binding specificity

In total, we identified over 600 non-zero energetic couplings between mutations in PDZ3 and mutations in the ligand (Z-test, Benjamini-Hochberg FDR<0.1, Fig. S5a). Once again, we define a specificity-changing mutation as one where the observed binding has a positive and significant (Z-test, FDR<0.1) residual to our additive energy model (Fig. 4f, Fig. S5c) and the observed binding score is significantly different from the binding score distribution of STOP codon variants) (Fig. S5b-c). These non-additive energetic interactions involve 340 distinct mutations in 73 different positions in the PDZ domain, and 114 distinct mutations in all 8 residues of the ligand (Fig. 4g). Mutations in PDZ3 can therefore alter its specificity for all 8 positions within the ligand.

Major specificity encoding residues and the modular encoding of binding specificity

We next identified the residues in PDZ3 most important for encoding specificity for each ligand position. In total, 20 pairs of PDZ3-ligand residues are enriched for specificity-changing mutations (hypergeometric test, FDR<0.1, Fig. 5a). We refer to the PDZ3 residues within these pairs as major specificity encoding residues. There are two major specificity-encoding residues for ligand position 0, six for position -1, five for position -2, three for position -3, three for position -4, and one for position -5 (Fig. 5a, Fig. S5e).

Strikingly, the major specificity encoding residues are largely distinct for each ligand position (Fig. 5a, Fig. S5d). There are 17 unique PDZ positions across the 20 pairs of positions. 14/17 PDZ positions only act as a major specificity determinant for a single ligand position, with 3 acting as major specificity determining sites for two ligand positions (Fig. S5d). The encoding of specificity in the PDZ domain is thus highly modular, with specificity for each ligand residue largely encoded by a distinct subset of residues in the domain.

The three exceptions are positions N326 and S339, which are both major specificity-encoding residues for positions -1 and -3, and G330, which defines the specificity for positions -2 and -4. N326 and S339 contact each other and -3 (N326 contacts -1 as well) (Fig. S5g). Similarly, G330 contacts H372 (Fig. S5g), which is a contact of position -2. G330 is also adjacent to E331, which is a contact of -4 (Fig. S5g). These positions therefore form small networks of specificity-defining residues, with pairs of residues in PDZ3 interacting with pairs of residues in the ligand.

Eleven of these sites have been previously reported to be important for specificity in different PDZ domains[21,22,41] (Fig. S5h, Table S1) but nine are novel sites that have not been previously shown to alter specificity (Fig. S5h). Four of these nine novel sites (L379 with V0, F325 and N326 with S-1, and E373 with K-4) are dynamic[26] or non-dynamic contacts[36] (Fig. S5f). Nine of the 20 sites (3 of the 9 novel sites) have previously been found to show signs of positive epistasis within the PDZ domain (as assessed by quantifying the energetic effects of double

mutants in cis in the PDZ domain), thereby showing evidence of coupling of these sites to the active site[43].

The major specificity changing mutations that we find in the PDZ domain (293/305 mutations in major specificity encoding residues for which we obtained abundance measurements[44]) do not change abundance of the protein as assessed by abundance PCA[42,44,45] (Fig. S5i). Furthermore, when we use the abundance data in combination with binding to fit a first order 3-state thermodynamic model (Fig. S5j), the performance is nearly identical to that of fitting the binding data without abundance (R^2=0.946 with abundance, R^2=0.947 without abundance). Quantifying binding residuals to the model with abundance as we did with the model without abundance (Fig. 4f), we find that the energetic couplings remain nearly identical (Fig. S5j, R^2=0.987 for >200,000 energetic couplings with and without abundance) as do the number of specificity-changing mutations (R^2=0.994 for 784 pairs of residues, Fig. S5k).

Direct and local allosteric encoding of specificity

Visualizing the 20 energetically coupled pairs of residues on the structure of the PDZ3-CRIPT complex (Fig. 5c-j), shows that not only are the major specificity encoding residues distinct for each ligand position, but they are also spatially clustered. Indeed 11/17 of the major specificity encoding residues constitute the PDZ3 binding interface (Fig. 5a) and out of the 20 major energetically coupled pairs of residues, nine are direct contacts (<5Å apart and/or predicted to be contacts via hydrogen bond, salt-bridge, or van der Waals force) Fig. 5c-j, Fig. S5f), suggesting local energetic coupling as the mechanism of action.

A further 11 pairs of major specificity encoding residues are not structural contacts. These residues must therefore encode specificity either via creating new contacts or allosterically. Three of these 11 pairs involve PDZ3 residues at the binding interface that contact other residues in the ligand (Fig. 5c-j, red-filled circles with asterisks). The other eight PDZ3 residues are all contacts of PDZ3 residues that contact the ligand (Fig. S4g).

In summary, specificity is encoded both modularly and locally within the PDZ domain. Mutations in a discrete set of spatially clustered sites encode specificity for each ligand residue.

A comprehensive map of specificity-changing mutations

For each pair of positions enriched in specificity-changing mutations, we considered how the coupled mutations in the domain and ligand relate to each other (Fig. 6 for three examples, full map in Fig. S6a, 20x20 maps for each pair of residues in Fig. S7), as well as how energetic couplings relate to changes in binding (compared to the wildtype preference) and raw binding scores (Fig. S6a). The majority of mutations that we identify in major specificity encoding residues change specificity such that the mutated ligand residue is preferred over wildtype, similar to 'class-switching' phenotypes[22,41] (Fig. S6a, middle panel). Others bind the mutated and wildtype ligand with similar affinity, acting similarly to 'class-bridging' specificity changes[22,41].

Interestingly, the diversity of specificity-changes that could be accommodated at each position in the ligand is higher for the less dynamic positions, except for position 0 (Fig. S6b). The most diverse specificity changes are observed for mutations at position -2, followed closely by -1 and -3 (Fig. S6b). This is consistent with previous reports that found specificity "modulators" of PDZ domains at non-canonical motif positions in the ligand[46]. It is also consistent with a study in which all positions common to the binding site across PDZ domains (N=10 sites) were mutated and found that only mutations in one of these 10 sites alters the specificity for position 0[21], suggesting that this is a general result across PDZ domains.

Overall, we find a large diversity of mutational couplings. While some positions are clearly dominated by charge-charge interactions (Fig. 6, S6), others appear to be mediated by gain/loss of other sidechain interactions or other molecular mechanisms. We highlight some of the coupled sites and mutations below for three of the six modules defining specificity for each ligand residue.

Position -2 binding specificity is encoded by H372 (a contact of T-2) and A376 (a contact of H372 and K380, contacts of T-2 and V0, respectively) (Fig. 6). It is also encoded by G329 (a contact of K-4 and H372), G330, and I336, all in the B2 and B3 beta-strands that contain residues directly involved in binding. Our comprehensive map finds all positions that were previously found to change specificity for T-2[21,22,41] across multiple studies (Fig. S5h). In addition, since we have couplings for every possible amino acid, we find hundreds of new specificity-changing mutations within these sites that have not been previously characterized.

The PDZ3 residue most strongly energetically coupled to position -2 is H372 (FDR<0.001, hypergeometric test, black rectangle Fig. 6). Indeed, PDZ3 position 372 and ligand position -2 are the most strongly coupled pair of positions in the dataset (table S2). This directly contacting position pair is connected by more than 70 energetic couplings. H372A was previously found to be a "class-switching" mutation[22,41], changing the preference at the T-2 position to an aromatic residue (F). As noted in a previous study, it is not only H372A that achieves this change in specificity, but many other mutations as well[22,41]. We gain resolution into the -2 preference and find that it can also be broadened to include other aromatic residues (W/Y), as well as to hydrophobic and positively charged residues (Fig. 6). For example, H372V/I/L/G/P/A change the position -2 specificity to aromatic, hydrophobic, and positively charged residues. H372C/T have more specific preference to aromatic or hydrophobic (but not positively charged) residues, and H372D shifts specificity to aromatic residues only. Interestingly, a fourth class of specificity change (T-2D, to an acidic residue) can be achieved with a A376C/G mutation. Based on a full hierarchically clustered map of specificity-changing residues across all sites (not just major specificity changing residues), several other mutations in PDZ can also change specificity of T-2 to T-2D (Fig. S8, blue cluster).

In terms of residues that are not direct contacts, hydrophobic and aromatic mutations in G329 change -2 specificity to hydrophobic residues (M/I/L). Similarly, mutations in the adjacent residue, G330, to V or A change specificity to I, and G330W changes specificity to I/C/V/F.

The main specificity-encoding positions for -3 are N326 (a contact of Q-3 and S-1), S339 (a contact of Q-3), and F340 (a contact of N326 and S339), meaning that the network of specificity-changing mutations around position -3 is fully connected by hydrogen bonds (Fig. 6, Fig. S5f-g). N326 mutations to negatively charged residues (D/E) change -3 specificity to positively charged residues (R/K) and there is a similar pattern with S339, where mutations to D/E (or Q, and, more weakly, C) change preference at -3 to R/K. The same strong charge-coupling pattern can be seen for F340, where mutations to D/E create a preference for Q-3K/R.

Position -5 has a single major specificity determinant: V328 (Fig. 6). Once again, we find charge-based couplings: V328E (negatively charged) enables binding to positively charged R/H, and V328R (positively charged) enables binding to negatively charged D/E.

Combinatorial quantitative reprogramming of binding specificity

Complete mutagenesis of the PDZ domain suggests a simple genetic architecture where binding specificity for six different ligand positions is encoded by six different modules of PDZ domain residues. However, in these experiments only a single mutation is made in the PDZ domain and only a single mutation in the ligand. What happens when multiple specificity-affecting mutations in the PDZ domain and/or the ligand are combined?

To address this question, we designed an experiment in which mutations in all six specificity residues of the PDZ domain and mutations in all six ligand sites are combined in all possible combinations. In the PDZ domain we included a strong specificity-changing mutation in each of the six modules (Fig. 7a) and programmed them as all single, double, triple, quadruple, quintuple and sextuple combinations (Fig. 7b-d). In the ligand we combined a total of 18 different matching specificity changes in the six positions, again as all single, double, triple, quadruple, quintuple and sextuple combinations (Fig. 7b-d). In total, there are 63 PDZ genotypes of different mutation orders paired with 2519 CRIPT genotypes of different mutation orders (Fig. S9a). Together these combine to a total of >165,000 designed genotypes when including STOP and synonymous variants. In total we quantified the binding of 150,367 of these combinatorial mutants using PCA (>90%, Fig. 7e).

Strikingly, fitting a thermodynamic model to this data shows that the binding of these combinatorial mutants is very well predicted by a simple model in which mutations have fixed energetic effects with a small contribution from pairwise energetic couplings (R^2=0.82, Fig. 7f, Fig. S9c). Interestingly, accurate prediction of binding for combinatorial mutants requires not only the pairwise energetic interactions between the PDZ domain and ligand—i.e. the previously measured specificity changing mutations (Fig 7h) — but also pairwise energetic couplings within the PDZ domain and within the ligand (Fig S9 d-i).

To illustrate how specificity mutations can be combined to re-programme binding to a new ligand we consider the ligand KNDERWWL. This ligand has mutations in all six specificity positions (Y-5D, K-4E, Q-3R, T-2W, S-1W, V0L). KNDERWWL is not bound by the wild-type

PDZ domain, but it is bound as predicted by the energy model by a PDZ domain with a mutation in each of the six specificity determining sites (Fig. 7i). Moreover, the energy model also accurately predicts the binding to this ligand of all 62 intermediate combinatorial genotypes i.e. the six single mutants, 15 double mutants, 20 triple mutants, 15 quadruple mutants and 6 quintuple mutants (R^2=0.94, Fig. 7j). Similarly, the energy model accurately predicts the quantitative binding of this six mutant PDZ domain across 2323 different ligands with between one and six different mutations (R^2=0.71, Fig. 7k).

Discussion:

To our knowledge, our data provide the first comprehensive map of how binding specificity is encoded in a globular protein domain. The map required the binding specificities of >1,800 proteins to be experimentally measured and provides a comprehensive and interpretable energy model describing how mutations throughout a protein domain reprogram its binding specificity.

In total we quantified >200,000 energetic couplings between mutations in a PDZ domain and its ligand to identify hundreds of mutations that alter the binding specificity of the domain. These mutations are, however, organized into six modules, with the specificity for each peptide residue encoded in a spatially clustered set of residues and by a mixture of direct and local allosteric effects. Our data further show that this simple and modular energetic architecture allows specificity to be accurately reprogrammed by combining mutations in each of the six modules (Fig. 8), with the binding of single, double, triple, quadruple, quintuple and sextuple mutants in the specificity-determining residues accurately predicted for both PDZ domain and ligand.

Our data also allow a direct comparison of the encoding of affinity and specificity in structured and dynamic molecular recognition – whilst specificity is primarily encoded in the structured region of the ligand, mutations in the dynamic region can be used to additively tune binding affinity. Moreover, interactions between the dynamic and structured region of the ligand – particularly between positions -5 and -3 – show that a dynamic region can contribute to binding specificity.

We believe that the approach that we have taken here – high-throughput quantification of energetic couplings between a protein and a ligand in a single pooled experiment – can be quite widely applied to understand how specificity and affinity are encoded across the diversity of molecular recognition[47]. It should be possible to apply this approach to both protein-protein and protein-nucleic acid interactions[48] to predictably and interpretably reprogram binding of structurally diverse proteins.

References:

1. Diss, G. Towards attaining a quantitative and mechanistic model of a cell. *Nat. Rev. Mol. Cell Biol.* **21**, 301–302 (2020).

2. Van Der Lee, R. *et al.* Classification of intrinsically disordered regions and proteins. *Chem. Rev.* **114**, 6589–6631 (2014).

3. Tompa, P., Davey, N. E., Gibson, T. J. & Babu, M. M. A Million peptide motifs for the molecular biologist. *Mol. Cell* **55**, 161–169 (2014).

4. Katuwawala, A., Peng, Z., Yang, J. & Kurgan, L. Computational Prediction of MoRFs, Short Disorder-to-order Transitioning Protein Binding Regions. *Comput. Struct. Biotechnol. J.* **17**, 454–462 (2019).

5. Wright, P. E. & Dyson, H. J. Linking folding and binding. *Curr. Opin. Struct. Biol.* **19**, 31–38 (2009).

6. Alderson, T. R., Pritišanac, I., Kolarić, Đ., Moses, A. M. & Forman-Kay, J. D. Systematic identification of conditionally folded intrinsically disordered regions by AlphaFold2. *Proc. Natl. Acad. Sci.* **120**, e2304302120 (2023).

7. Tompa, P. & Fuxreiter, M. Fuzzy complexes: polymorphism and structural disorder in protein–protein interactions. *Trends Biochem. Sci.* **33**, 2–8 (2008).

8. Fuxreiter, M. Fuzzy protein theory for disordered proteins. *Biochem. Soc. Trans.* **48**, 2557–2564 (2020).

9. Mittag, T. *et al.* Dynamic equilibrium engagement of a polyvalent ligand with a single-site receptor. *Proc. Natl. Acad. Sci.* **105**, 17772–17777 (2008).

10. Tuttle, L. M. *et al.* Gcn4-Mediator Specificity Is Mediated by a Large and Dynamic Fuzzy Protein-Protein Complex. *Cell Rep.* **22**, 3251–3264 (2018).

11. Borgia, A. *et al.* Extreme disorder in an ultrahigh-affinity protein complex. *Nature* **555**, 61–66 (2018).

12. Skriver, K., Theisen, F. F. & Kragelund, B. B. Conformational entropy in molecular recognition of intrinsically disordered proteins. *Curr. Opin. Struct. Biol.* **83**, 102697 (2023).

13. Teilum, K., Olsen, J. G. & Kragelund, B. B. On the specificity of protein-protein interactions in the context of disorder. *Biochem. J.* **478**, 2035–2050 (2021).

14. Brodsky, S. *et al.* Intrinsically Disordered Regions Direct Transcription Factor In Vivo Binding Specificity. *Mol. Cell* 1–13 (2020) doi:10.1016/j.molcel.2020.05.032.

15. Baughman, H. E. R. *et al.* An intrinsically disordered transcription activation domain increases the DNA binding affinity and reduces the specificity of NFκB p50/RelA. *J. Biol. Chem.* **298**, 102349 (2022).

16. Nguyen Ba, A. N. *et al.* Proteome-wide discovery of evolutionary conserved sequences in disordered regions. *Sci. Signal.* **5**, rs1 (2012).

17. Davey, N. E. *et al.* Attributes of short linear motifs. *Mol BioSyst* **8**, 268–281 (2012).

18. Nourry, C., Grant, S. G. N. & Borg, J.-P. PDZ Domain Proteins: Plug and Play!

19. Amacher, J. F., Brooks, L., Hampton, T. H. & Madden, D. R. Specificity in PDZ-peptide interaction networks: Computational analysis and review. *J. Struct. Biol. X* **4**, 100022 (2020).

20. Laub, M. T. & Goulian, M. Specificity in Two-Component Signal Transduction Pathways. *Annu. Rev. Genet.* **41**, 121–145 (2007).

21. Tonikian, R. *et al.* A Specificity Map for the PDZ Domain Family. *PLoS Biol.* **6**, e239 (2008).

22. Raman, A. S., White, K. I. & Ranganathan, R. Origins of Allostery and Evolvability in Proteins: A Case Study. *Cell* **166**, 468–480 (2016).

23. Liu, X. & Fuentes, E. J. Emerging Themes in PDZ Domain Signaling. in *International Review of Cell and Molecular Biology* vol. 343 129–218 (Elsevier, 2019).

24. Doyle, D. A. *et al.* Crystal structures of a complexed and peptide-free membrane protein-binding domain: Molecular basis of peptide recognition by PDZ. *Cell* **85**, 1067–1076 (1996).

25. Songyang, Z. *et al.* Recognition of Unique Carboxyl-Terminal Motifs by Distinct PDZ Domains. *Science* **275**, 73–77 (1997).

26. Mostarda, S., Gfeller, D. & Rao, F. Beyond the binding site: The role of the β2 - β3 loop and extra-domain structures in PDZ domains. *PLoS Comput. Biol.* **8**, (2012).

27. Saro, D. *et al.* A thermodynamic ligand binding study of the third PDZ domain (PDZ3) from the mammalian neuronal protein PSD-95. *Biochemistry* **46**, 6340–6352 (2007).

28. Ivarsson, Y. & Jemth, P. Affinity and specificity of motif-based protein–protein interactions. *Curr. Opin. Struct. Biol.* **54**, 26–33 (2019).

29. Jordan, D. F., Dubé, A. K., Dionne, U., Bradley, D. & Landry, C. R. Residues Neighboring an SH3-Binding Motif Participate in the Interaction *In Vivo*. Preprint at https://doi.org/10.1101/2024.05.13.593936 (2024).

30. Petit, C. M., Zhang, J., Sapienza, P. J., Fuentes, E. J. & Lee, A. L. Hidden dynamic allostery in a PDZ domain. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 18249–18254 (2009).

31. Gianni, S. *et al.* Demonstration of Long-Range Interactions in a PDZ Domain by NMR, Kinetics, and Protein Engineering. *Structure* **14**, 1801–1809 (2006).

32. Gautier, C., Visconti, L., Jemth, P. & Gianni, S. Addressing the role of the α-helical extension in the folding of the third PDZ domain from PSD-95. *Sci. Rep.* **7**, 12593 (2017).

33. Lockless, S. W. & Ranganathan, R. Evolutionarily conserved pathways of energetic connectivity in protein families. *Science* **286**, 295–299 (1999).

34. Laursen, L., Kliche, J., Gianni, S. & Jemth, P. Supertertiary protein structure affects an allosteric network. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 24294–24304 (2020).

35. Kumawat, A. & Chakrabarty, S. Hidden electrostatic basis of dynamic allostery in a PDZ domain. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E5825–E5834 (2017).

36. Niethammer, M. *et al.* CRIPT, a Novel Postsynaptic Protein that Binds to the Third PDZ Domain of PSD-95/SAP90. *Neuron* **20**, 693–707 (1998).

37. Skelton, N. J. *et al.* Origins of PDZ Domain Ligand Specificity. *J. Biol. Chem.* **278**, 7645–7654 (2003).

38. Tarassov, K. *et al.* An in vivo map of the yeast protein interactome. *Science* **320**, 1465–1470 (2008).

39. Faure, A. J. & Lehner, B. MoCHI: neural networks to fit interpretable models and quantify energies, energetic couplings, epistasis and allostery from deep mutational scanning data. Preprint at https://doi.org/10.1101/2024.01.21.575681 (2024).

40. Zhou, H.-X. & Pang, X. Electrostatic Interactions in Protein Structure, Folding, Binding, and Condensation. *Chem. Rev.* **118**, 1691–1741 (2018).

41. McLaughlin, R. N., Poelwijk, F. J., Raman, A., Gosal, W. S. & Ranganathan, R. The spatial architecture of protein function and adaptation. *Nature* **491**, 138–142 (2012).

42. Faure, A. J. *et al.* Mapping the energetic and allosteric landscapes of protein binding domains. *Nature* **604**, 175–183 (2022).

43. Nedrud, D., Coyote-Maestas, W. & Schmidt, D. *A Survey of Pairwise Epistasis Supports an Outside-in Hierarchy of Clade-Specifying and Function-Defining Residues in PSD95 PDZ3.* http://biorxiv.org/lookup/doi/10.1101/2020.06.26.174375 (2020) doi:10.1101/2020.06.26.174375.

44. Hidalgo-Carcedo, C., Faure, A. J., Martí-Aranda, A. & Zarin, T. Allosteric and Energetic Remodeling by Protein Domain Extensions.

45. Levy, E. D., Kowarzyk, J. & Michnick, S. W. High-resolution mapping of protein concentration reveals principles of proteome architecture and adaptation. *Cell Rep.* **7**, 1333–1340 (2014).

46. Amacher, J. F., Cushing, P. R., Brooks, L., Boisguerin, P. & Madden, D. R. Stereochemical Preferences Modulate Affinity and Selectivity among Five PDZ Domains that Bind CFTR: Comparative Structural and Sequence Analyses. *Structure* **22**, 82–93 (2014).

47. Forman-Kay, J. D. & Mittag, T. From sequence and forces to structure, function, and evolution of intrinsically disordered proteins. *Struct. Lond. Engl. 1993* **21**, 1492–9 (2013).

48. Li, X. & Lehner, B. TF-MAPS: fast high-resolution functional and allosteric mapping of DNA-binding proteins.

49. Diss, G. & Lehner, B. The genetic landscape of a physical interaction. *eLife* **7**, 1–31 (2018).

50. Grant, B. J., Skjærven, L. & Yao, X. The Bio3D packages for structural bioinformatics. *Protein Sci.* **30**, 20–30 (2021).

51. Weng, C., Faure, A. J., Escobedo, A. & Lehner, B. The energetic and allosteric landscape for KRAS inhibition. *Nature* https://doi.org/10.1038/s41586-023-06954-0 (2023) doi:10.1038/s41586-023-06954-0.

52. Faure, A. J., Schmiedel, J. M., Baeza-Centurion, P. & Lehner, B. DiMSum: an error model and pipeline for analyzing deep mutational scanning data and diagnosing common experimental pathologies. *Genome Biol.* **21**, 207 (2020).

53. Wagih, O. Ggseqlogo: A versatile R package for drawing sequence logos. *Bioinformatics* **33**, 3645–3647 (2017).

54. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview Version 2-A multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–1191 (2009).

55. Bolognesi, B. *et al.* The mutational landscape of a prion-like domain. *Nat. Commun.* **10**, (2019).

56. de Hoon, M. J. L., Imoto, S., Nolan, J. & Miyano, S. Open source clustering software. *Bioinformatics* **20**, 1453–1454 (2004).

57. Saldanha, A. J. Java Treeview — extensible visualization of microarray data. **20**, 3246–3248 (2004).

58. Kim, G. *et al.* Easy and accurate protein structure prediction using ColabFold. *Nat. Protoc.* **20**, 620–642 (2025).

59. Meng, E. C. *et al.* UCSF ChimeraX : Tools for structure building and analysis. *Protein Sci.* **32**, e4792 (2023).

Acknowledgements:

Funding: This work was funded by European Research Council (ERC) Advanced grant (883742), Wellcome (220540/Z/20/A), the Spanish Ministry of Science and Innovation (LCF/PR/HR21/52410004, EMBL Partnership, Severo Ochoa Centre of Excellence), the Bettencourt Schueller Foundation, the AXA Research Fund, Agencia de Gestio d'Ajuts

Universitaris i de Recerca (AGAUR, 2017 SGR 1322), and the CERCA Program/Generalitat de Catalunya. T.Z. was funded by a EMBO Long-term Postdoctoral Fellowship (ALTF 525-2021) and a Marie Skłodowska-Curie Postdoctoral Fellowship (GPIDR, 101068134).

We acknowledge all members of the Lehner lab for feedback during the project, particularly Toni Beltran, Albert Escobedo, Andre Faure, Taylor Mighell, Michael Thompson, Magda Topolska, Max Stammnitz and Chenchun Weng for feedback/advice with experiments and data analysis. We thank Júlia Domingo for sharing plasmid vectors pGJJ211 and pGJJ215. We acknowledge the CRG Genomics Unit for DNA sequencing services. Molecular graphics and analyses were performed with UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

Author contributions:

T.Z. and B.L. conceptualized the study. T.Z. performed all formal analysis, data curation, methodology, software, validation, and visualization. T.Z. performed all experiments except for the abundancePCA of PDZ3, which was performed by C. H-C. T.Z. and B.L. acquired funding. B.L. provided resources and supervision. T.Z. and B.L. wrote the original draft and reviewed and edited the manuscript.

Data availability:

All raw DNA sequencing data and associated processed data files (binding fitness measurements for all datasets) have been deposited in the Gene Expression Omnibus under the accession number GSE265816. All other data required to reproduce analyses are available at https://zenodo.org/records/11048045.

Code availability:

All custom scripts required to reproduce analyses are available at https://github.com/lehner-lab/fuzzy_specificity.

Competing interests:

B.L. is a founder and shareholder of ALLOX.

List of supplementary materials:

Materials and Methods

Figs. S1 to S9
Tables S1 to S5

Figures:

Figure 1. Combinatorial mutagenesis of a peptide ligand. a) Normalized b-factor vs position of residue for PDB structures of PDZ3 crystallized with its ligand b) PDZ3-CRIPT structure (PDB ID: 5heb[22]) where CRIPT is coloured by b-factor (normalized to be on the same scale as all other available PDZ3 structures with their ligands). K-7 is represented in white and dashed/moving lines since it is absent from the crystal structure. c) libraries of variants are transformed into yeast and selected via a protein complementation assay that quantifies protein binding, in this case

PDZ3-CRIPT[38,49] (see Methods). The input library (pre-selection for binding) and the output library are sequenced and a fitness score is derived representing the enrichment or depletion of variants upon selection as compared to the wildtype sequence (see Methods). d) library design and pipeline to determine binding fitness for all possible mutants of dynamic N and structured C terminus of CRIPT e) density distribution of binding fitness for N, C, and STOP variants as compared to wildtype. f) Position weight matrix (PWM) based on binding fitness scores for top 1% of variants in N vs C. g) Number and percent of top 1% of variants stratified by AA Hamming distance and their associated PWMs. h) Increasing net charge of residues in N results in incrementally increased fitness, but not in C.

a) 2-state MoCHI model

$$phenotype_b = \frac{1}{1 + e^{\Delta G_b/RT}}$$

$$\Delta G_b = -RTlog(K_b c)$$

$$\hat{y} = h(g(f_b(x)))$$

PDZ3 + CRIPT ⇌ PDZ3 CRIPT
unbound   bound
        K_b

b)

N — dynamic
C — structured

CRIPT
K-7 N-6 Y-5 K-4 Q-3 T-2 S-1 V0

2-state MoCHI model (1st order)

i) R^2 0.731 | N = 76893
Predicted fitness vs Observed fitness
count: 250 200 150 100 50

iii) R^2 0.749 | N = 7405
Predicted fitness vs Observed fitness
count: 500 400 300 200 100

ii) Observed fitness vs Additive trait
count: 400 300 200 100

iv) Observed fitness vs Additive trait
count: 30 20 10

c)

Mutation (W Y F Q N T S G P D E K R H C I L V M A) vs Position (K-7 N-6 Y-5 K-4 Q-3 T-2 S-1 V0)

ddG: 2 1 0 -1 □ WT

Mean

Position

Figure 2. Energetic couplings in dynamic and structured portions of CRIPT. a) Two-state MoCHI thermodynamic model to transform binding fitness scores of mutations into energetic terms (ddG) of binding. K_b is the binding equilibrium constant, phentoype$_b$ is the fraction bound, ΔGb is the Gibbs free energy of binding, R is the gas constant, T is the temperature in Kelvin, c is the standard reference concentration. ŷ is the output variable (predicted binding fitness), h is a linear activation function transforming the molecular phenotype to the units of the assay, g is a function for global epistasis describing the nonlinear relationship between the energetic effects of mutation and the binding

phenotype, and f(x) is the additive trait map. b) Performance of model and additive trait coefficients of the dynamic N (left, panel i and ii) and structured C (right, panel iii and iv) portions of CRIPT. c) Heatmap of energetic binding terms for the dynamic N (left) and structured C (right) portions of CRIPT for all possible mutations (y-axis, coloured and ordered by physicochemical property).

Figure 3. Measuring >8,000 energetic couplings between CRIPT residues reveals specificity-encoding positions within the dynamic peptide. a) Design of CRIPT cis double mutant library b) Density distributions of fitness and c) barplot of amino acid hamming distance for all variants d) Performance of first order MoCHI model fit on the CRIPT cis double mutant data e) Heatmap of the number of specificity-changing mutations per pair of CRIPT positions, with one significantly enriched coupled site outlined in black (FDR<0.1) f) Heatmap of residuals for each pair of mutations in the significantly coupled CRIPT sites Q-3 and Y-5. Mutations on axes are coloured by physicochemical property.

Figure 4. Measuring >200,000 energetic couplings between PDZ3 and CRIPT allows identification of specificity-encoding sites between the domain and peptide. a) Design of PDZ3-CRIPT trans library in which a library of all single substitutions in CRIPT is combined with a library of all single substitutions in PDZ3. Both libraries also contained designed STOP mutations and synonymous substitutions to comprise a total of 320k possible genotypes. b) AA Hamming distance and c) density distribution of binding fitness across the library for variants that pass quality thresholds. d) PDZ3-CRIPT structure (PDB ID: 5heb[22]) coloured by the average binding fitness effect for single substitutions. The canonical/constrained binding motif sites in CRIPT are labelled and have the strongest average binding fitness defects, as expected. e) heatmaps showing binding fitness of single substitutions in CRIPT (left) and PDZ3. STOP mutations are designed to only be in the N-terminus of CRIPT and two blocks of PDZ3. Most widely deleterious effects are at the binding interface (also seen in d)). f) Performance of first order 2-state MoCHI model on >240k variants (left) and residuals to the model (right) where those hexbins that pass FDR threshold of 10% (significantly different z-score from background) are outlined in yellow. g) Number of unique specificity-changing mutations (FDR<0.1) across CRIPT (left) and PDZ3 (right).

Figure 5. Major specificity encoding residues for each residue of a peptide. a) Heatmap showing number of specificity-changing mutations across every position-pair in PDZ3 and CRIPT. Outlined boxes mark those that pass FDR thresholds for enrichment as shown (black FDR <0.01, dark red FDR 0.01-0.1) and comprise the major specificity encoding sites. PDZ annotations for binding interface (< 5Å) and secondary structure are shown along x axis). b) Number of specificity-changing mutations across each position pair versus distance between the position pair in question. Circles are coloured by FDR of enrichment, showing as in a) that most position pairs are not enriched in specificity-changing mutations and that position-pairs that are enriched in specificity-changing mutations tend to be closer to each other in the structure. c) – j) PDZ3-CRIPT (PDB ID: 5heb[22]) structure coloured by number of specificity-changing mutations, split by the identity of the CRIPT residue (highlighted in orange [V0, S-1, T-2, Q-3, K-4, Y-5, N-6, K-7] in separate panels). Number of specificity-changing mutations is thresholded between 1 and 10 to more clearly show range for significant (FDR<0.1) sites. Only positions in PDZ3 that pass the FDR threshold of 0.1 are labelled.

Figure 6. Examples of specificity changes. Energetic couplings are quantified as binding residuals for the three examples of coupled residues colored on the crystal structure[22] (Y-5 and V328 lilac, Q-3 and N326 magenta, T-2 and H372 orange) along with their sidechains (where available). Full specificity map in Figure S6.

Figure 7. Quantitative reprogramming of binding specificity. a) Top specificity-changing positions in PDZ coloured to show how they are paired to each module/ligand position for which they change specificity in CRIPT. b) Design of reprogramming library, where the top specificity-changing mutation at each domain position is encoded along with the

specificity changes it enables at each position in CRIPT. All possible single, double, triple, quadruple, quintuple, and sextuple mutations are encoded in the designed library. c) Two example specificity-changing mutations in PDZ are shown along with the mutations in the ligand that they are energetically coupled to (i.e. the specificity changes they enable shown via quantification of energetic couplings). The colour of the box around the mutation corresponds to the colour of the mutation in panel b. d) Crystal structure for wildtype PDZ3[22] shown alongside predicted structures for PDZ3 domains with varying degrees of combined specificity-changing mutations. Physicochemical property of residue identity is coloured as per legend in panel c and corresponding ligand sequence for each structure is shown at the bottom left. Colouring of residues on structure matches panel a, denoting how the residues in the ligand and domain are coupled. e) Distribution (top) and sample size (bottom) of designed (white) and recovered (blue) library of variants with amino acid hamming distance (or stop variant) denoted by gradient of colours (purple, 1 amino acid hamming distance to yellow, 12 amino acid hamming distance). f) Performance of two-state thermodynamic model fit with MoCHI on entire reprogramming design library considering all pair-wise interactions between mutations (top), additive trait coefficients extracted from model vs. observed fitness (bottom). g) dddG (2^{nd} order energetic couplings) between PDZ specificity-changing mutations (x axis) and designed mutations in CRIPT (y axis), with corresponding specificity changes based on trans-double mutant library outlined in black. Negative dddG is a favourable interaction, blue is a detrimental interaction, units are kcal/mol. h) Linear correlation between energetic couplings inferred from trans double mutant library for all designed mutation pairs in reprogrammed library (x-axis) versus dddG energetic couplings inferred from fitting the thermodynamic model in panel f to reprogrammed library (y-axis). Dots are coloured by mutation-pair identity corresponding to panel b, and mutation pairs that are direct structural contacts are denoted by an asterisk. i) Binding fitness distribution of all variants matching one of the fully redesigned ligands (KNDERWWL) grouped by the number of mutations in PDZ3 (or STOP codon mutations in CRIPT). j) Performance of 2-state thermodynamic model in d for the designed KNDERWWL ligand with dots coloured by varying number of mutations in domain k) Performance of 2-state thermodynamic model on domain with 6 designed mutations with dots coloured by varying number of mutations in ligand.

Figure 8. Modular encoding of ligand specificity. Number of specificity changing mutations in each of the 17 major specificity determining sites (left). The sites in each module (-5,-4,-3,-2,-1,0) are colored differently (center). Combining mutations from the different modules allows quantitative reprogramming of specificity (right).

Materials and Methods:

Media

- LB: 10 g/L Bacto-tryptone, 5 g/L Yeast extract, 10 g/L NaCl. Autoclaved 20 min at 120ºC.
- YPD: 20 g/L glucose, 20 g/L Peptone, 10 g/L Yeast extract. Autoclaved 20 min at 120ºC.
- SORB: 1 M sorbitol, 100 mM LiOAc, 10 mM Tris pH 8.0, 1 mM EDTA.
- Filter sterilized (0.2 mm Nylon membrane, ThermoScientific).
- Plate mixture: 40% PEG3350, 100 mM LiOAc, 10 mM Tris-HCl pH 8.0, 1 mM EDTA pH 8.0. Filter sterilized.
- Recovery medium: YPD (20 g/L glucose, 20 g/L Peptone, 10 g/L Yeast extract) +0.5 M sorbitol. Filter sterilized.
- SD -URA: 6.7 g/L Yeast Nitrogen base without amino acid, 20 g/L glucose, 0.77 g/L complete supplement mixture drop-out without uracil. Filter sterilized.
- SD -URA/ADE: 6.7 g/L Yeast Nitrogen base without amino acid, 20 g/L glucose, 0.76 g/L complete supplement mixture drop-out without uracil, adenine and methionine. Filter sterilized.
- MTX competition medium: SD –URA/ADE + 200 ug/mL methotrexate (BioShop Canada Inc., Canada), 2% DMSO.
- DNA extraction buffer: 2% Triton-X, 1% SDS, 100mM NaCl, 10mM Tris-HCl pH8, 1mM EDTA pH8.

B-factor analysis

We obtained all 214 available PDZ domain family entries bound to a ligand from the PDB. We used the Bio3D[50] R package to read in the entries, filtering for those with 2 unique chains in the structure (n = 61), further filtered for those that had B-factors available (n = 58). We normalized the B-factors by the minimum value within that chain in order to be able to compare B-factors across crystal structures. We filtered the length of chain B to be in the first quartile of the dataset (in order to roughly filter for PDZ domains bound to a peptide ligand that was 10 aa or less) and present the 14 PDZs bound to a short ligand and their respective normalized B-factors in Fig. 1a).

Library construction

We designed five libraries to probe different questions about the PDZ3-CRIPT binding interaction. All libraries had the same backbone bindingPCA vector structure of DLG4-PDZ3 (aa 303-402) N-terminally fused to DHFR3 and CRIPT (0 to -8) N-terminally fused to DHFR1,2 as in[42]. All plasmids used in the study are described in table S3. The library design for measuring PDZ3 abundance (Fig. S5i) is as described in[44].

Combinatorial N- and C-terminal CRIPT libraries were each ordered as an NNK degenerate oligo from IDT and cloned into the bindingPCA vector containing PSD95-PDZ3 (aa 303-402) fused to DHFR3 and no bait fused to DHFR1,2 (hence "empty"). The library transformation was bottlenecked in *E. coli* such that the combinatorial space of variants was reduced from the possible 32^4 total variants (32 possible codons at 4 positions) to 20^4 in order to facilitate greater downstream sequencing depth for each variant.

The CRIPT "cis" double mutant library was ordered as an IDT pool of NNK oligos with 28 NNK degenerate oligos encoding all different combinations of double mutants across CRIPT positions 0 to -7. The library was transformed into the same empty vector as the CRIPT N and CRIPT C libraries described above.

The PDZ3-CRIPT "trans" double mutant library was ordered as 3 separate TWIST oligo pools. PDZ3 (aa 303-402) was split into two non-overlapping blocks for mutagenesis: Block 1 encoding the first 50 aa in the domain and Block 2 encoding the latter 50 aa in the domain. We designed 1000 variants encoding the following for each block: wildtype, all possible single mutants, half of all possible synonymous mutants, and half of all possible single STOP codon variants starting at the N-terminus of each block. Variants were designed such that each possible single mutant codon was firstly scored by the number of nucleotide substitutions away from wildtype (such that 2-3 substitutions were preferred, followed by 1, followed by 0) and then by the most optimal yeast codon (based on the *S. cerevisiae* codon usage table). This design strategy enabled us to use non-overlapping reads to sequence the long amplicons that encoded PDZ3 at one end and CRIPT at the other (schematic in Fig. S4a, designed library and results in Fig. S4b). The CRIPT single mutant library was designed in the same way but ordered as a single block due to its short length. Each library was amplified using distinct primers and transformed separately into the same backbone/bindingPCA vector (PGJJ001 as in [42]). The PDZ3 portion of the Block 1 and Block 2 vectors were then cloned in two separate reactions into the vector with the CRIPT single mutant library to produce 2 double mutant library vectors (PDZ3 Block 1 single mutants + CRIPT single mutants, PDZ3 Block 2 single mutants + CRIPT single mutants).

The combinatorial reprogramming library was ordered as 2 separate TWIST oligo pools: one for PDZ3 and one for CRIPT. The PDZ3 pool contained all the 6 target specificity-changing mutations (N326D, V328R, E331N, S339D, H372G, L379G within aa 326-379, Fig. 7b) as single, double, triple, quadruple, quintuple, and sextuple mutations, as well as wildtype, synonymous, and STOP codon variants (at the most N-terminal position [N326] for the latter). The PDZ mutations were chosen as follows: the residual values for major specificity changing mutations were ordered first by the FDR associated with the paired position (Table S2), then sorted by residual quantile, then by abundance score quantile, and finally by residual value. The CRIPT pool contained 18 mutations at all 6 corresponding positions in CRIPT where the chosen PDZ3 mutations changed specificity (i.e. they had a positive and significant [FDR<10%] energetic coupling based on results from the trans double mutant experiment). These 18 CRIPT mutations (Fig. 7b-c) were again included as single, double, triple, quadruple, quintuple and sextuple mutations along with wildtype, synonymous, and STOP codon variants. As with the trans double mutant library, the pool of PDZ variants was combined with the pool of CRIPT mutants (all-by-all) by cloning the libraries sequentially into the bindingPCA vector (PGJJ001) to encode 63*2519 variants (Fig. S9a). We show the distribution of designed variants recovered in the assay (>90%) in Fig. 7e.

Large-scale transformations of libraries into yeast and competition assays

We transformed each library of variants into *S. cerevisiae* in 3 replicates at a large volume scaled to the size of the library in order to ensure that all variants were present in multiple (at least 100) copies as in previous work[42,51] to prevent bottlenecking the library. We grew the cultures to saturation in synthetic complete media with 2% glucose as a carbon source. We harvested these cultures as the "input" replicates to our selection assay, and subjected the input library to selection. The selection experiment is based on a well-described protein complementation assay[38] wherein the yeast are grown in synthetic complete media with added methotrexate (MTX), a drug that requires dihydrofolate reductase (DHFR) for metabolization. In

the presence of MTX, CRIPT (or the bait protein) must be bound to PDZ3 (the prey protein) to bring together the split fragments of DHFR and enable cell growth (Fig. 1c). We harvested these cultures as the "output" from the selection assay. We extracted the DNA from each replicate using a standard phenol chloroform procedure as in[42]. To prepare the DNA for Illumina sequencing, we used two PCR steps to obtain the amplicon for each library (table S4). In PCR1, we added frameshifting oligonucleotides (table S5) and amplified the regions of interest (amplicon with constant region and mutated region) from the extracted DNA with 5 cycles. In PCR2 we added Illumina sequencing barcodes with PCR using the minimum number of cycles necessary to reach amplification plateau for each sample based on a qPCR run.

Next Generation Sequencing and analysis of sequencing data (read counts to fitness scores)

We use sequencing as a quantitative readout for binding between PDZ3 and CRIPT (Fig. 1c). We obtained reliable sequencing data for more than half a million variants across all five libraries, obtaining a fitness score (and associated error) for each variant using DimSum[52]. Parameters used to filter sequencing reads for DimSum required an input count of 10 reads in at least one replicate, and we filtered each dataset for the specific design of each library (i.e. libraries made with NNK degenerate oligos were filtered for NNK design and the custom pdz3-cript trans and combinatorial reprogramming libraries were filtered to only keep designed variants).

Normalization of fitness scores across experiments

The CRIPT N and C combinatorial, double mutant CRIPT, and PDZ3-CRIPT trans mutant libraries contained overlapping variants that had highly correlated fitness (binding) scores when processed independently (Fig. S1f,h, Fig. S3c). In order to make all scores comparable across the study, we used a linear transformation based on these highly correlated shared variants to normalize each library (Fig. S1g,i, Fig. S3d).

Position-weight matrices

All position-weight matrices were constructed using ggseqlogo[53] with a custom Zappo-based[54] color scheme to mark physicochemically related amino acids.

Calculation of physicochemical properties

We calculated features from a curated list of amino acid property scales (n=386) (http://www.genome.jp/aaindex/) as in[55] to quantify the correlation of these scores with the binding scores for the combinatorial N and C libraries in Fig. 1. We present those features that had a | Spearman's r | > 0.4 for either the N or C libraries in Fig. S1l.

Modeling phenotype to free energy with MoCHI

To translate the fitness scores, which capture the phenotypic effects of mutations, into free energy terms, we used the MoCHI package[39] to model the fitness with a two-state thermodynamic model for protein binding. Briefly, MoCHI takes as input amino acid sequences of each variant and predicts their fitness while correcting for global non-linearities (non-specific epistasis). Using the coefficients extracted from the model, we obtain the change in free energy associated with each mutation for the phenotype in question (in our case, binding). We used default parameters for a two-state model with one phenotype (binding) for all datasets, except for the model that includes abundance in Fig. S5i. For all models, we used L1 and L2

regularization with a lambda of 10^{-6}. We evaluated the model using the held-out "fold" from the 10 times that the model was run on the dataset.

The CRIPT C combinatorial library contained an overabundance (>90%) of non-binding variants, so we balanced the dataset by sampling the distribution of >100k variants without replacement so that the variants with the largest distance from the peak of non-binding variants (specified by 2 s.d. away from mean of STOP codon binding distribution) had a higher probability of being sampled (Fig. S2a-b). We used a distance function based on a power-law distribution to weight the sampling probability as follows:

$$W_i = k^{1 - \frac{x_i}{d}}$$

Where W_i is the sampling weight of each variant binding score x_i, d is the mean binding score for the non-binding/dead mode, and k is a constant (=10^{10}, chosen to balance the dataset towards a reasonable percentage of binding variants [~25%], Fig. S2b).

To test the reproducibility of the model results, we repeated the sampling procedure 10 times and performed the three implementations of MoCHI modeling (linear model with no global epistasis, two-state thermodynamic model with first order terms, two-state thermodynamic model with second order terms) (Fig. S2f). We found the modeling results to be extremely stable across 10 iterations and therefore show one representative iteration in Fig. 2, Fig. S2 and Fig. S4g.

Analysis of residuals to quantify energetic couplings

For both PDZ3-CRIPT trans and cis CRIPT double mutant libraries we quantified the residuals from the observed vs. mean predicted binding fitness for each variant in the respective dataset. We converted the residuals to Z-scores (using the error derived from DimSum as the denominator) and performed a Z-test to derive p-values for each variant as the Z-scores were normally distributed. We corrected the p-values for multiple testing (Benjamini-Hochberg) and report the FDR values associated with residuals where relevant. Residuals (energetic couplings) that are significant by the multiple test-corrected Z-test, >0 and pertain to variants that pass the non-binding threshold (i.e. are binding) are classified as specificity-changing. To test for enrichment of specificity-changing mutations in PDZ3-CRIPT pairs, we did a test for enrichment (hypergeometric test) of specificity-changing mutations across all pairs of sites. We again performed multiple test correction (Benjamini-Hochberg) on these to identify the major specificity encoding residues/coupled sites as those that pass an FDR threshold of 0.1. We highlight these major energetically coupled sites in Fig. 5 and Fig. S6a, and list them in table S1 and S2.

Unsupervised clustering of binding residuals

We clustered the vector of binding residuals to all possible CRIPT variants for each PDZ3 variant that had at least one specificity-changing mutation (N=340), filtered to include those PDZ3 variants with >80% data present (N=290). We used Cluster 3.0[56] to perform unsupervised hierarchical clustering with a weighted cosine distance, average linkage and otherwise default parameters. We used JavaTreeview[57] to visualize the full cluster plot and several manually highlighted clusters in Fig. S8.

Protein contact determination

We used getContacts (https://getcontacts.github.io/) to predict contacting residues using get_static_contacts.py with the 5heb.pdb[22] structure, --itypes option set to "all", and otherwise default parameters. We also used the Bio3D[50] package to calculate inter-residue distances between specific residues in PDZ3 and CRIPT based on the 5heb[22] structure from PDB.

Prediction of protein structures with multiple mutations

To predict protein structures for the PDZ3-CRIPT interaction with mutations in PDZ3 and/or CRIPT, we used ColabFold v1.5.5[58] using template mode: pdb100 and otherwise default parameters. We show the predicted structures with 2 and 12 mutations in Fig. 7c where indicated.

Visualization of protein structures

All protein structures were based on the PDZ3-CRIPT structure with PDB ID 5heb[22]. In Fig. 1b we present the full crystalized structure, but since the K-7 position is missing, we added this residue using ChimeraX[59] v1.4 and therefore present it as a cartoon with motion lines around it and no associated B-factor. All other represented protein structures include only those residues in PDZ3 and CRIPT that we had in our mutagenesis design, i.e. aa 303-402 in PDZ3 and 0 to -7 in CRIPT. All quantitative visualization on structures was performed with the color by attribute function in ChimeraX.

Supplementary Figures:

Figure S1. Overview of CRIPT combinatorial N and C data and quality. a) replicate fitness correlations (Pearson's r) for all N-terminal CRIPT variants b) replicate fitness correlations (Pearson's r) for all C-terminal CRIPT variants and c) top 1% of variants d) replicate count correlations (Pearson's r) for N-terminal CRIPT and e) C-terminal CRIPT. f-i)

normalization of fitness via a linear transformation using shared variants between CRIPT combinatorial libraries and the pdz3-cript trans double mutant library. j) Number of variants in N vs C for each AA hamming distance away from wildtype sequence k) Hamming distance vs. fitness of N vs C Top physicochemical features that correlate (Spearman's rho > 0.4) with binding fitness scores in N or C dataset.

Figure S2. Additional MoCHI modeling results for combinatorial CRIPT N and CRIPT C datasets a) distribution of all variants in CRIPT C combinatorial library, marking percentage of variants that are "alive"/binding (i.e. 2 s.d. away from the mean of the distribution of STOP codons) b) distribution of subsampled variants that were used to balance

the dataset (see Methods) c) performance of several models on CRIPT N (left) and C (right) datasets, including a linear model that does not account for global epistasis (top), residuals of a 2-state MoCHI model shown in Fig. 2 (middle), and a 2-state MoCHI model that takes into account interactions between mutations (bottom). d) The relationship between ddG values and error, with cutoff of 1 kcal/mol indicated by dashed line to represent confident ddGs – all ddG values with high error values are in the V0 and T-2 positions that also have the highest ddG values. e) correlation between ddGs from 1^{st} order vs 2^{nd} order MoCHI model for CRIPT N and C datasets. Error bars for x and y axis indicate respective 95% confidence interval in kcal/mol. f) Results from running all models on 10 iterations of subsampling the CRIPT C dataset. Mean of model performance (i.e. explained variance, R^2 observed vs. predicted fitness) via 10-fold cross-validation for the 10 models is shown with 95% confidence intervals (2*s.d.) for each model.

Figure S3. Overview of CRIPT cis double mutant library data and quality. a) replicate count correlations and b) replicate fitness correlations for all variants. c) correlation and d) normalization using linear transformation of shared variants across CRIPT cis double mutant library (y axis) and PDZ3-CRIPT double mutant trans library e) additive trait coefficients and f) residuals from fitting MoCHI two-state model g) distribution of FDR of residuals quantified from fitting MoCHI two-state model with 1% and 10% FDR cutoffs show h) distribution of variants with STOP codons and justification for non-binding/dead variants threshold (2 s.d. away from mean of STOP variants) i) Performance of two-state MoCHI model with data points coloured by the FDR of their residuals to the model fit. Dotted line shows dead threshold based on fitness distribution of STOP codon variants. Specificity-changing mutations are demarcated.

Figure S4. Overview of PDZ3-CRIPT trans library data and quality. a) sequencing strategy of PDZ3-CRIPT double mutant library where R1 and R2 denote forward and reverse sequencing reads, respectively. b) PDZ3-CRIPT library sequenced variants (_seq_>10 to reflect more than 10 input reads in the sample) reflect library design to maximize

number of nucleotide substitutions for each encoded variant along with the most optimal yeast codon (see methods) c) replicate count correlations for PDZ3-CRIPT library d) replicate fitness correlations for PDZ3-CRIPT library e) percent of reads in input sample at each position (out of total reads in input) for each of the 100 positions assayed across the PDZ domain – even sequencing expectation is 1%, denoted by dashed red line, showing that most positions are evenly sampled except for the missing tail end of block 2. f) all-by-all heatmap of binding fitness for PDZ3-CRIPT double mutants. Mutation physicochemical properties are represented by colours as shown in Fig. 1), positions are coloured from N to C (dark to light gradient). g) correlation between ddGs from two-state MoCHI model trained on combinatorial CRIPT libraries vs. PDZ3-CRIPT libraries with corresponding 95% confidence intervals as error bars on x and y axes.

Figure S5. Determination of specificity-changing mutations and association with structural contacts. a) FDR of binding fitness residuals to MoCHI 2-state model for PDZ3-CRIPT mutants with different cutoffs b) distribution of STOP codon binding fitness and threshold for "dead"/non-binding threshold (2 s.d. away from STOP codon variant fitness

mean). c) definition of a specificity-changing mutation d) the number of CRIPT positions for which each major specificity encoding residue in PDZ encodes specificity, showing that the vast majority encode specificity for only one CRIPT position e) Number of major specificity encoding residues (in PDZ3) for each CRIPT position (-6 and -7 have zero) f) PDZ3 position vs. CRIPT position showing position pairs enriched in specificity-changing mutations (major specificity-encoding sites) outlined by FDR value (solid vs dashed line representing <1% vs 1-10% respectively) and the contact type of each pair as classified by getContacts (see Methods); vdw = van der waal's, hbbb = hydrogen bond backbone-backbone, hbsb = hydrogen bond sidechain-backbone, hbss = hydrogen bond sidechain-sidechain, sb = salt bridge, wb = water bond, hp = hydrophobics g) PDZ3 positions of major specificity-encoding residues (y axis) and their contacts in PDZ3 as determined by getContacts. CRIPT residue for which each PDZ3 position encodes specificity is represented by filled circles. h) Venn diagram of all 20 major specificity encoding sites discovered in this study, and whether or not they were previously found in 3 other studies that experimentally examined change in specificity of a ligand upon mutation of a PDZ domain (see Methods for details). i) Distribution of abundance measurements for the PDZ domain (STOP= STOP codon variants, all other variants=non-STOP codon variants), with vertical dark lines indicating where the 305/323 major specificity changing mutations for which abundance measurements are available are located on the abundance fitness distribution. 293/305 of these sites are abundant (2 s.d. away from blue STOP codon variant distribution). The wildtype is indicated by a blue dotted line at 0. j) Three-state first order thermodynamic model fit to the binding measurements for trans mutations between PDZ and CRIPT as in Fig. 4.f, but also including abundance measurements k) Pearson correlation between energetic couplings (binding residuals) derived from thermodynamic model fit with and without abundance measurements included in the model l) Pearson correlation between the number of specificity-changing mutations per pair of residues with and without abundance measurements included in the thermodynamic model

Figure S6. A complete map of all specificity-changing mutations across major specificity encoding sites in PDZ3-CRIPT a) Three columns of heatmaps show the energetic coupling (binding fitness residual), change in binding (binding fitness of mutant vs. wildtype), and raw binding fitness scores for all major specificity-changing PDZ3

mutations. Mutations are colored by physicochemical property and positions in PDZ are colored by their position in PDZ (from N to C-terminus) as well as their contact/allosteric classification for the CRIPT residue that they are coupled to. b) Number of observed unique specificity changes in CRIPT vs b-factor for each residue.

Figure S6. (continued)

Figure S7. Clustered heatmaps of all 20 PDZ3-CRIPT major pairs of specificity encoding positions (i.e. significantly enriched FDR<0.1 for specificity-changing mutations).

Figure S8. Clustered heatmap of binding residuals for all PDZ mutations (y-axis, N=290) that have at least one associated specificity-changing mutation in CRIPT (x axis, N=152). Outlined boxes represent manually-chosen clusters that point to PDZ positions enriched for specificity-changing mutations (H372, N326, S339) as shown in detail in Fig. 6 but also other sets of positions that are seemingly unrelated (orange, purple, blue clusters) with corresponding structural annotations of these sites to the right of each cluster.

Figure S9. Data quality and first and second energetic terms for reprogramming library. a) library design showing number of single, double, triple, quadruple, quintuple and sextuple mutations for individual PDZ3 and CRIPT libraries that were combined to make up the reprogramming library. b) replicate correlations (Pearson's r) for bindingPCA of

reprogramming library. c) additive thermodynamic 2-state model fit using MoCHI on reprogramming library d) first order energetic terms for the ligand from the two-state thermodynamic model shown in Fig. 7f e) second order energetic terms (dddG) between mutations in the ligand f) strongest negative interaction between mutations in the ligand shown on the structure g) first order energetic terms for PDZ3 based on the two-state thermodynamic model shown in Fig. 7f h) second order energetic terms (dddG) between mutations in the domain i) strongest negative interaction between mutations in the domain and the positively interacting mutations with which it interacts j) Distribution of binding fitness effects for any designed ligand k) distribution of binding fitness effects for designed ligands omitting negative interactions between T-2R and S-1W.

Supplementary tables:

PDZ residue	Secondary Structure	Secondary Structure (alt)	Secondary structure class	CRIPT residue	Number of specificity changes (this study)	**Major specificity change (this study)**	Tonikian et al. 2008[21]	McLaughlin et al. 2012[41]	Raman et al. 2016[22]
A343			loop	V0	7	1			
L379	A2-8	AB	alpha_helix	V0	9	1			
G324	B1:B2-7		loop	S-1	16	1			
F325	B2-1	BB	beta_sheet	S-1	6	1			
N326	B2-2	BB	beta_sheet	S-1	57	1			
S339	B3-4	BC	beta_sheet	S-1	8	1	1		
L342			loop	S-1	8	1	1		
G345	A1-2		alpha_helix	S-1	8	1			
G329	B2-5	BB	beta_sheet	T-2	7	1		1	
G330	B2:B3-1		loop	T-2	5	1		1	1
I336	B3-1	BC	beta_sheet	T-2	5	1		1	1
H372	A2-1	AB	alpha_helix	T-2	74	1	1	1	1
A376	A2-5	AB	alpha_helix	T-2	8	1	1		1
N326	B2-2	BB	beta_sheet	Q-3	12	1	1		
S339	B3-4	BC	beta_sheet	Q-3	32	1	1		

F340	B3-5	BC	beta_sheet	Q-3	5	1	1		
G330	B2:B3-1		loop	K-4	5	1			
E331	B2:B3-2		loop	K-4	8	1			
E373	A2-2	AB	alpha_helix	K-4	5	1			
V328	B2-4	BB	beta_sheet	Y-5	6	1	1		

Table S1. Comparison of energetically coupled sites between PDZ3 and CRIPT in our study versus others that used a comparable approach. We include the positions in PDZ with standard names based on the secondary structure fold to facilitate comparison across PDZ domains.

Rank	Pair ID	CRIPT N vs. C	FDR (enrichment in number of spec. changing mutations across all pairs)	Number of spec. changing mutations	PDZ3 residue	CRIPT residue	Distance between residues (Å)
1	372_-2	C	2.03E-125	74	H372	T-2	2.715445
2	326_-1	C	1.49E-83	57	N326	S-1	3.593695
3	339_-3	C	1.22E-38	32	S339	Q-3	2.543977
4	324_-1	C	7.82E-14	16	G324	S-1	7.03372
5	326_-3	C	1.32E-08	12	N326	Q-3	3.08644
6	379_0	C	7.58E-06	9	L379	V0	4.245281
7	376_-2	C	3.74E-05	8	A376	T-2	3.924201
8	329_-2	C	0.000116	7	G329	T-2	9.165152
9	331_-4	N	0.000116	8	E331	K-4	3.126404
10	339_-1	C	0.000116	8	S339	S-1	8.3229
11	345_-1	C	0.000116	8	G345	S-1	11.18858
12	342_-1	C	0.000121	8	L342	S-1	7.746873
13	343_0	C	0.000607	7	A343	V0	12.59954
14	325_-1	C	0.007777	6	F325	S-1	7.572008
15	328_-5	N	0.007777	6	V328	Y-5	8.612266
16	330_-2	C	0.025456	5	G330	T-2	9.862768
17	336_-2	C	0.028273	5	I336	T-2	8.526768
18	340_-3	C	0.033814	5	F340	Q-3	4.862758
19	330_-4	N	0.057264	5	G330	K-4	7.177915
20	373_-4	N	0.076377	5	E373	K-4	2.670947

Table S2. Major (FDR<0.1, hypergeometric test) energetically coupled pairs of sites that control specificity

plasmid	description	link	Use	source
pGJJ001	bindingPCA vector with DHFR3 and DHFR1,2 fused to nothing (empty bait and prey sites)	https://benchling.com/s/seq-FRMVd55qhXRSdnCerIB4?m=slm-HFeYB6ebmq6hKEVIXJ7S	Vector for CRIPT single mutant library followed by PDZ3 block 1 and block 2 libraries	Faure, Domingo, Schmiedel et al. Nature 2022

pGJJ215	DHFR3-wildtype PDZ3, empty "bait" for inserting CRIPT libraries	https://benchling.com/s/seq-YgJTHoCogB748M3lvrz1?m=slm-96kiUpsaHHKMnzOCiKvl	Vector for CRIPT N and C combinatorial libraries, CRIPT cis double mutant library	Lehner lab
pGJJ211	DHFR3-wildtype PDZ3 "mutagenesis" plasmid for cloning libraries	https://benchling.com/s/seq-ZLKW8KZ0LU1BsoN88fBn?m=slm-pK7yOttj3EJwRkSJzK98	Vector for PDZ3 block 1 and block 2 libraries	Lehner lab
pGJJ518	bindingPCA vector with wildtype PDZ3, wildtype CRIPT	https://benchling.com/s/seq-5EQ0SO1NzUXy4soExGvc?m=slm-maaM0f6pwkJWq60vFxQR	Wildtype control (used to obtain empirical estimates of sequencing error, quantify amount of plasmid in genomic DNA extractions and test bindingPCA assay)	This work

Table S3. Plasmid sequences used in this study

amplicon	link	notes
CRIPT	https://benchling.com/s/seq-ZjVNfyB4q0j7ghLbVvos?m=slm-tekXiMuhgKyP7qCWdIhJ	Reference amplicon for CRIPT N, CRIPT C combinatorial libraries and CRIPT CIS double mutant library
PDZB1_CRIPT	https://benchling.com/s/seq-m7a9OYYNBzp1HrSv0cgQ?m=slm-j3wlHLONBYjdpZy7uZVO	Reference amplicon for PDZ3-CRIPT -- block 1 of PDZ3
PDZB2_CRIPT	https://benchling.com/s/seq-5krvszGJdAW94riW4Pmi?m=slm-b0Ku0YyzgXMeTFhjQ54N	Reference amplicon for PDZ3-CRIPT TRANS double mutant library -- block 2 of PDZ3
PDZ_repro_CRIPT	https://benchling.com/s/seq-xu1tTD1vsGCAgReFo1vW?m=slm-SRKT4rhimKjvpagPyDxD	Reference amplicon for PDZ3-CRIPT combinatorial reprogramming library

Table S4. Amplicon sequences used in this study

name	nuc_seq	forward_reverse	library_1	library_2	library_3
oTZ132	ACACTCTTTCCCTACACGACGCTCTTCCGATCTAGGTGGAGGCGGATCCACC	forward_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ134	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNAGGTGGAGGCGGATCCACC	forward_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ135	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNNAGGTGGAGGCGGATCCACC	forward_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ136	ACACTCTTTCCCTACACGACGCTCTTCCGATCTCNBAGGTGGAGGCGGATCCACC	forward_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE

oTZ137	ACACTCTTTCCCTACACG ACGCTCTTCCGATCTTBH YAGGTGGAGGCGGATCC ACC	forward_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ138	ACACTCTTTCCCTACACG ACGCTCTTCCGATCTGAH YYAGGTGGAGGCGGATCC CACC	forward_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ133	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTtc gaaggctttaatttgaCTAGTCTA	reverse_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ139	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTNt cgaaggctttaatttgaCTAGTCT A	reverse_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ140	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTN Ntcgaaggctttaatttgacta CTA	reverse_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ141	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTR NWtcgaaggctttaatttgaCTAG TCTA	reverse_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ142	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTC RSNtcgaaggctttaatttgaCTA GTCTA	reverse_CRIPT	CRIPT_N	CRIPT_C	CRIPT2XA3_CRIPT_CIS_DOUBLE
oTZ154	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTA GGTGGAGGCGGATCCAC C	reverse_CRIPT	PDZ3CRIPT _TRANS_D OUBLE	PDZ3_RE PRO_CRI PT_REPR O	NA
oTZ155	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTN AGGTGGAGGCGGATCCA CC	reverse_CRIPT	PDZ3CRIPT _TRANS_D OUBLE	PDZ3_RE PRO_CRI PT_REPR O	NA
oTZ156	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTN NAGGTGGAGGCGGATCC ACC	reverse_CRIPT	PDZ3CRIPT _TRANS_D OUBLE	PDZ3_RE PRO_CRI PT_REPR O	NA
oTZ157	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTC NBAGGTGGAGGCGGATC CACC	reverse_CRIPT	PDZ3CRIPT _TRANS_D OUBLE	PDZ3_RE PRO_CRI PT_REPR O	NA
oTZ158	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTT BHYAGGTGGAGGCGGAT CCACC	reverse_CRIPT	PDZ3CRIPT _TRANS_D OUBLE	PDZ3_RE PRO_CRI PT_REPR O	NA
oTZ159	GTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTG AHYYAGGTGGAGGCGGA TCCACC	reverse_CRIPT	PDZ3CRIPT _TRANS_D OUBLE	PDZ3_RE PRO_CRI PT_REPR O	NA

oTZ160	ACACTCTTTCCCTACACGACGCTCTTCCGATCTTCGGGAGGTGGAGCTAGC	forward_PDZ3B1	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ161	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNTCGGGAGGTGGAGCTAGC	forward_PDZ3B1	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ162	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNNTCGGGAGGTGGAGCTAGC	forward_PDZ3B1	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ163	ACACTCTTTCCCTACACGACGCTCTTCCGATCTCNWTCGGGAGGTGGAGCTAGC	forward_PDZ3B1	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ164	ACACTCTTTCCCTACACGACGCTCTTCCGATCTAWBRTCGGGAGGTGGAGCTAGC	forward_PDZ3B1	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ165	ACACTCTTTCCCTACACGACGCTCTTCCGATCTTBANWTCGGGAGGTGGAGCTAGC	forward_PDZ3B1	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ166	ACACTCTTTCCCTACACGACGCTCTTCCGATCTGCAGACCTCAGTGGGGAG	forward_PDZ3B2	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ167	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNGCAGACCTCAGTGGGGAG	forward_PDZ3B2	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ168	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNWGCAGACCTCAGTGGGGAG	forward_PDZ3B2	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ169	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNNWGCAGACCTCAGTGGGGAG	forward_PDZ3B2	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ170	ACACTCTTTCCCTACACGACGCTCTTCCGATCTCSNWGCAGACCTCAGTGGGGAG	forward_PDZ3B2	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ171	ACACTCTTTCCCTACACGACGCTCTTCCGATCTTANNTGCAGACCTCAGTGGGGAG	forward_PDZ3B2	PDZ3CRIPT_TRANS_DOUBLE	NA	NA
oTZ200	ACACTCTTTCCCTACACGACGCTCTTCCGATCTTCCACGGGCCTGGGCTTC	forward_PDZrepro	PDZ3_REPRO_CRIPT_REPRO		
oTZ201	ACACTCTTTCCCTACACGACGCTCTTCCGATCTNTCCACGGGCCTGGGCTTC	forward_PDZrepro	PDZ3_REPRO_CRIPT_REPRO		

oTZ202	ACACTCTTTCCCTACACG ACGCTCTTCCGATCTNNT CCACGGGCCTGGGCTTC	forward_PDZrepro	PDZ3_REP RO_CRIPT_ REPRO		
oTZ203	ACACTCTTTCCCTACACG ACGCTCTTCCGATCTCNW TCCACGGGCCTGGGCTT C	forward_PDZrepro	PDZ3_REP RO_CRIPT_ REPRO		
oTZ204	ACACTCTTTCCCTACACG ACGCTCTTCCGATCTAWB RTCCACGGGCCTGGGCT TC	forward_PDZrepro	PDZ3_REP RO_CRIPT_ REPRO		
oTZ205	ACACTCTTTCCCTACACG ACGCTCTTCCGATCTGBA NKTCCACGGGCCTGGGC TTC	forward_PDZrepro	PDZ3_REP RO_CRIPT_ REPRO		

Table S5. List of frameshifting oligonucleotides used for PCR1