Abstract

Natural history museum collections harbour a record of wild species from the past centuries, providing a unique opportunity to study animals as well as their infectious agents. Thousands of great ape specimens are kept in these collections, and could become an important resource for studying the evolution of DNA viruses. Their genetic material is likely to be preserved in dry museum specimens, as reported previously for monkeypox virus genomes from historical orangutan specimens. Here, we screened 209 great ape museum specimens for 99 different DNA viruses, using hybridization capture coupled with short-read high-throughput sequencing. We determined the presence of multiple viruses within this dataset from historical specimens and obtained several near-complete viral genomes. In particular, we report high-coverage (>18-fold) hepatitis B virus genomes from one gorilla and two chimpanzee individuals.

Keywords

Museomics; Great Apes; Target-enrichment capture; Viruses; Hepatitis B virus

Screening great ape museum specimens for DNA viruses

Authors

Michelle Hämmerle[1,2], Meriam Guellil[1,2], Olivia Cheronet[1,2], Susanna Sawyer[1,2], Irune Ruiz-Gartzia[3], Esther Lizano[4,3], Aigerim Rymbekova[1,2], Pere Gelabert[1,2], Paolo Bernardi[1], Sojung Han[1,2], Lovro Trgovec-Greif[5,6], Thomas Rattei[5], Verena J. Schuenemann[1,2,7,8], Tomas Marques-Bonet[3,4,9,10], Katerina Guschanski[11,12], Sebastien Calvignac-Spencer[13,14], Ron Pinhasi[1,2], Martin Kuhlwilm[1,2]

Affiliations

1. Department of Evolutionary Anthropology, University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria
2. Human Evolution and Archaeological Sciences (HEAS), University of Vienna, Austria
3. Institute of Evolutionary Biology (UPF-CSIC), PRBB, Dr. Aiguader 88, 08003 Barcelona, Spain
4. Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, Barcelona, SpainCatalan Institution of Research and Advanced Studies (ICREA), Barcelona, Spain
5. Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria
6. Doctoral School of Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria
7. Institute of Evolutionary Medicine, University of Zurich, Zurich, Switzerland
8. Department of Environmental Sciences, University of Basel, Basel, Switzerland
9. National Center for Genomic Analysis (CNAG), Baldiri i Reixac 4, 08028 Barcelona, Spain
10. Institució Catalana de Recerca i Estudis Avançats (ICREA) and Universitat Pompeu Fabra. Pg. Luís Companys 23, 08010, Barcelona, Spain
11. Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh, Edinburgh, UK
12. Department of Ecology and Genetics, Animal Ecology, Uppsala University, SE-75236 Uppsala, Sweden
13. Helmholtz Institute for One Health, Helmholtz-Centre for Infection Research (HZI), 17489 Greifswald, Germany
14. Faculty of Mathematics and Natural Sciences, University of Greifswald, 17489 Greifswald, Germany

corresponding author: Martin Kuhlwilm (martin.kuhlwilm@univie.ac.at)

Introduction

The extensive collections of fossils and specimens preserved in natural history museums document the diversity of life forms, ecosystem dynamics, and the transformative processes that have shaped our planet over millions of years. They are a critical resource for scientific research. By employing high-throughput DNA sequencing techniques tailored to ancient or historical specimens, scientists can reconstruct ancient genomes[1] opening a new dimension in the field of museomics. The integration of genomic technologies with traditional museum collections presents unprecedented opportunities to address consequential questions in ecology and evolutionary biology[2]. Comparing genomic data from museum specimens with modern populations allows for the reconstruction of a more comprehensive tree of life. It facilitates assessments of environmental change impacts on genetic diversity and population

structure. This information is essential not only for understanding biodiversity and the forces
that have shaped it in the past but also for developing effective conservation strategies to
preserve genetic diversity and mitigate anthropogenic disturbances to natural ecosystems[3].

An intriguing avenue of investigation for museomics is the study of infectious diseases. Natural
history museum specimens have rarely been used for this purpose, but archaeological
specimens have demonstrated the immense potential of ancient microbial genetic material in
shedding light on the evolution and spread of infectious agents[4]. This is notably true for
research on viruses with a DNA genome (DNA viruses). Hundreds to thousands of years old
DNA viral genomes reconstructed from archeological specimens have unveiled key aspects of
their recent evolutionary histories. For example, researchers have shown complete lineage
turnover of hepatitis B viruses (HBV; family *Hepadnaviridae*) in European human populations
around the Bronze Age[5,6]. Genomic pathways to increased host adaptation and virulence have
also been clarified for variola viruses (*Poxviridae*) in medieval human populations[7], for Marek's
disease virus (*Herpesviridae*) in 19th/20th century poultry[8], or myxoma virus (*Poxviridae*) in
rabbits[9].

In parallel, efforts geared toward a better understanding of the determinants of health in our
closest relatives, the nonhuman great apes (orangutans – *Pongo pygmaeus*, *P. tapanuliensis*
and *P. abelli*, gorillas – *Gorilla gorilla* and *G. beringei*, chimpanzees – *Pan troglodytes* – and
bonobos – *Pan paniscus*), have revealed that these species host a broad range of DNA viruses.
This includes enzootic viruses belonging to the families *Adenoviridae*, *Anelloviridae*,
Circoviridae, *Hepadnaviridae*, *Herpesviridae*, *Papillomaviridae*, *Parvoviridae* and
Polyomaviridae, as well as emerging viruses such as members of the family *Poxviridae*[10,11]. Co-
phylogenetic analyses have shown that DNA viruses have often been stably associated with
their hominid hosts for extensive periods of time. While co-divergence with their hosts is
frequent, rare cross-species transmission events between hominids have contributed to
shaping all hominid DNA viromes. For example, the herpes simplex virus 2 (*Herpesviridae*) that
causes genital herpes in humans likely arose from the cross-species transmission of a virus
infecting members of the gorilla lineage, several million years ago[12]. Conversely, patterns of
genomic variation suggest that HBV was transmitted from humans to nonhuman great apes
over the last few thousand years[6]. As we learn more about nonhuman great ape viromes, we
will uncover more of the complex origins of their DNA viruses and those infecting humans.

Explorations of nonhuman great ape viromes are ongoing and should proceed in the
framework of long-term studies of wild populations, but they could be efficiently
complemented by analyses of the thousands of specimens kept in natural history museums
around the world. Not only do these collections provide immediate access to animal tissues
(as opposed to the noninvasive samples that constitute the bulk of biological sampling from
modern populations), they also open a direct window on 200 years of increased anthropogenic
disturbance characterized by dwindling, increasingly fragmented populations of these
animals[13]. These processes have likely altered nonhuman DNA viromes significantly, and
possibly resulted in the extinction of viral lineages[14]. Moreover, emerging viruses constitute a
threat to wild great ape populations[11,15], and they are often of human origin[16]. Therefore,
studying the DNA viromes of nonhuman great ape museum specimens can provide insights
into both extinct and contemporary viral diversity. This may also allow to monitor changes in
viromes of wild populations with the potential to intervene should the populations be exposed
to novel viruses (e.g. from humans or other species) they have not co-evolved with.

Here, we present high-throughput sequencing data from 209 nonhuman great ape museum
specimens which were sampled and analysed in search for DNA viruses. These specimens cover
all nonhuman great ape species, most subspecies and large parts of their recent geographical
distribution. A majority originated in the wild, but our sample set also includes captive
individuals from European zoos (13%, n=28). Obtaining data from ancient specimens can be

challenging due to DNA degradation and damage, as well as potential contamination during storage and handling[1]. While museum specimens are younger than typical archaeological remains, high variability in DNA quality has been observed[17], and handling in specific clean laboratory facilities, following strict protocols is necessary[18], and was applied for this study.

In addition, we also developed and implemented an efficient customised enrichment strategy, in-solution hybridization capture with RNA baits[19] to account for the extremely low abundance of viral genetic material. Although metagenomic detection of viruses is possible, it is particularly challenging for DNA viruses that, with a few exceptions, usually do not replicate as intensely as RNA viruses[20]. Hybridization capture offers considerable flexibility, because it allows us to multiplex baits targeting many different viruses, as well as enrich even significantly divergent targets (up to 58% divergence tolerance according to some authors[21]). However, probably many DNA viruses evolve relatively slowly[22]. We designed and used a bait set that covered the genomes of 99 viral lineages representing 13 families. For capture, 8-10 samples were pooled for a more cost-effective screening and sequenced on an Illumina platform.

Results

Sequencing data
The median number of raw reads per library was 1,122,746 reads, with a high variability of yield (SD=3,556,238). After adapter trimming, a median of 956,296 reads was retained, with a reduction due to the removal of adapter-multimers between 0.11 and 93.38%. The rates of duplicated reads ranged between 21.35% and 88.67%, leaving a median of 338,560 high-quality unique reads across the 214 libraries (SD=971,276; Fig. 1A-B).

Figure 1. Distribution of sequencing reads per library. A) Numbers of raw reads, reads after trimming and unique reads (after BBmap clumpify) across 214 libraries. B) Same as in A) but cut at 4 million reads (without tail). C) Number of reads per library assigned to *Homo*, Bacteria, or Viruses using Kraken2. We note that mapping great ape sequencing data to the human genome is commonly performed in genomic studies to avoid reference bias[23]. D) Same as C), but number of reads on log-scale. E) Percentage of assigned reads per library.

The sequence length of the raw reads was 100 bp, as defined by the sequencing setup, and after filtering and trimming, the median fragment length per library ranged from 40 to 100 bp. This suggests that the extracted DNA may not have been as much fragmented as ancient DNA[1], but likely contained a large proportion of fragments longer than 100 bp. This might be expected for museum specimens, which are younger than archaeological specimens.

Results from virus capture

Fragments from targeted virus strains are likely absent or very rare in the libraries. Given the low expected abundance of any viral DNA, a large degree of amplification led to high duplication rates (see above). Furthermore, as we chose a lower hybridization temperature and a prolonged incubation time for the already highly amplified libraries, more unspecific binding resulted from our approach. We observed a median assignment of 21.95% to *Homo sapiens*, 13.91% to bacterial sequences and solely 0.1% to viruses (Fig. 1C-E). Mapping to

human DNA likely reflected endogenous great ape DNA (and possibly human contamination), and bacterial DNA likely resulted from post-mortem colonization of the specimen. While many viral reads were assigned to bacteriophages (viruses infecting bacteria, not animals), many libraries contained at least one read assigned to a viral family known to infect animals (Table S3, Fig. 2A-B). Numerous libraries contained at least a small number of reads (25 or more) assigned to a viral family (Fig. 2C). A surprising amount of libraries apparently comprised poxvirus reads, which likely represented wrong assignments. When restricting to virus strains that were included in the capture design, we observe fewer instances (n=30) of libraries with potential virus fragments (Fig. 3), which likely reflects a more accurate picture.

Figure 2. Summaries of reads assigned to different virus domains and families. A) Proportion of virus-assigned reads based on kraken2 across 214 libraries, stratified by realm and by family (for DNA viruses and *Retroviridae*). B) Number of libraries with any read assigned to virus families. C) Number of libraries with at least 25 reads assigned to virus families.

Figure 3. Heatmap of positive reads per viral species. The figure depicts all libraries that have at least 25 hits assigned to one of the 99 viruses in the capture kit (with merged numbers for Cytomegalovirus strains).

Henceforth, we focused on libraries with more than 500 assigned reads, which are the most likely to reflect true viral infection. We found six such instances, with assignments either to the family *Poxviridae* (n=3; all orangutan specimens) or the family *Hepadnaviridae* (n=3; 1 gorilla and two chimpanzee specimens). The only library with more viral than bacterial or human-mapping sequences is L1949 (19.13% viral reads), obtained from an orangutan specimen. We note that further rounds of enrichment capture might have provided higher on-target coverage[1,24], but at substantially higher costs.

Positive specimens

To validate the results from classifying reads, we attempted to reconstruct the corresponding viral genomes. Using reference-based mapping, we managed to obtain low-coverage monkeypox virus (MPXV; family *Poxviridae*) genomes from three libraries (up to 3.6-fold coverage on the reference genome sequence KJ642614) (Table 1). Further investigation of these specimens, including deeper sequencing to obtain high-coverage genomes, was published in another study[25], where we present their origin from a zoo outbreak in 1965. Similar efforts aimed at assembling HBV genomes from the three most promising HBV-positive specimens resulted in one high-coverage genome from a wild gorilla, and two from wild chimpanzees (Table 1), sufficient for generating consensus sequences (Methods). In other samples, putative viral sequences were at a much lower abundance (Table S3).

Library ID (genus)	Individual ID (museum)	Reference	Mapped Reads	Mean ED	Mean DOC	1X	5X	10X
L1946 (*Pongo*)	ZFMK_MAM1965-0547 (Bonn)	KJ642614 (monkeypox virus)	7,931	0.51	3.83	93.04	33.25	2.58
L1947 (*Pongo*)	ZFMK_MAM1965-0545 (Bonn)	KJ642614 (monkeypox virus)	4,659	0.5	2.29	84.45	12.04	0.09
L1949 (*Pongo*)	ZFMK_MAM1965-0546 (Bonn)	KJ642614 (monkeypox virus)	2,352	0.61	1.13	63.83	1.03	0
L3064 (*Gorilla*)	96550 (Frankfurt)	AJ131567 (hepatitis B virus)	2,969	0.82	97.07	99.62	99.56	99.43
L3863 (*Pan*)	ZMB_Mam_83617 (Berlin)	ON706349 (hepatitis B virus)	517	1.82	18.19	94.44	73.60	53.80
L3859 (*Pan*)	ZMB_Mam_11638 (Berlin)	AF305327 (hepatitis B virus)	919	1.98	31.27	92.36	75.83	61.56

251 **Table 1. Statistics for virus genomes identified in this study.** Mapping metrics for the six viral
252 genomes discovered in this study, including proportion of the reference genome covered by at
253 least 1, 5 or 10 reads. Mapped reads = unique reads with MQ >30. DOC = Depth of coverage. ED =
254 Edit distance.

Discussion

256
In this study, we screened a total of 209 great ape museum samples with a myBaits® capture
kit including 99 viral strains. We obtained six complete virus genomes, three HBV (from one
gorilla and two chimpanzees), and three MPXV genomes (from three orangutans).
The high-coverage Hepatitis B genome obtained in this study is most similar to one sequenced
from the blood serum of a wild-caught western lowland gorilla from Cameroon. The serum
was part of the primate serum bank stored at the Zoological and Botanical Garden Wilhelma,
Stuttgart, Germany. The individual in our study is also a gorilla (without subspecies
identification) from Cameroon. The two other HBV genomes are from chimpanzees and have
a higher edit distance, indicating a deeper divergence between the genomes sequenced here,
and these closest identified reference genomes. Only the human version of HBV was included
in the capture kit, while the great ape-infecting lineages are nested within the diversity of
human HBV lineages. Hence, the gorilla and chimpanzee versions were identified and
successfully retrieved through enrichment in this study. HBV is endemic in wild gorillas[26] with
a rate of at least 11 to 30 % of past HBV infections[27], and finding the genetic legacy of a gorilla
HBV in this study demonstrates that such infections can be detected from wild individuals from
museum collections.
According to the museum metadata from, the orangutan specimens yielding MPXV reads were
from Sumatra. However, MPXV infections in the wild have not been observed outside of Africa.
A search in the museum archive yielded a letter from the wildlife trader, stating that those
were zoo individuals, without specifying the zoo from which they were taken. We identified
an MPXV outbreak in Rotterdam Zoo in 1964/65[28], during which six out of nine infected
orangutans died. Mappings to an MPXV genome from this outbreak revealed a close match to
the ones we obtained, leading to the conclusion that we sampled orangutans that died during
this outbreak. A comprehensive study, including more data, has been published separately[25].

In this study, we were able to screen a large number of samples (209 specimens), with the
inclusion of all great ape species. As great apes are endangered nowadays, these specimens
provide a unique insight into a time when human influence on their populations was less
pronounced. By pooling eight to ten samples together, it was feasible to perform an effective
screening. However, limitations are the availability of viral strains within databases and the
use of target enrichment, which limits our detection of undescribed historical strains, as well
as the fact that the targeted viruses were likely absent in some libraries. Further rounds of
enrichment capture may have resulted in higher on-target rates of coverage[1,29], but this would
have significantly reduced the number of samples that could be screened.
Despite the low specificity, we successfully determined the presence of multiple DNA viruses
in this sample set, including the detection of more than 500 reads of MPXV in three
orangutans, as well as HBV in two chimpanzees and one gorilla, respectively. For these
specimens, the reconstruction of near-complete viral genomes was possible. Our work
demonstrates the feasibility of viral DNA enrichment and detection from museum specimens
of great apes.
296
297
298

Methods

300
Samples
For this project, we collected a total of 209 great ape specimens (Fig. 4), from which 214
sequencing libraries were produced. Of these libraries, 66 were from gorillas, 84 from
chimpanzees (10 of these as *Pan* sp., but most likely *Pan troglodytes* ssp.), eight from bonobos,
and 56 from orangutans, including different subspecies. We note that two separate extracts
and sequencing libraries were prepared for five specimens. The samples were obtained from
specimens housed in European natural history museums, namely in Germany in Berlin (n =
28), Bonn (n = 28), Dresden (n = 26), Frankfurt (n = 63) and Stuttgart (n = 6), in the Czech
Republic in Prague (n = 24), and in Austria in Salzburg (n = 22) and Vienna (n = 12).
Approximately 92% of libraries were obtained from teeth (n = 196), 17 libraries from soft
tissues (Table S1), and two specimens were phalanges. According to the museum metadata,
the oldest specimen was from 1838, and the most recent (a captive individual) was from 2014.
Some individuals were held in captivity, mostly in zoos, whereas others were wild-caught.
More detailed information concerning the individuals and the libraries can be found in Table
S1.
The museum identifiers of the specimens are as follows: Senckenberg Forschungsinstitut und
Naturmuseum Frankfurt/M.: 10325, 1110, 1111, 1112, 1113, 1114, 1115, 1118, 1119, 1120,
1121, 1126, 1132, 1134, 1576, 1579, 15792, 15817, 16180, 17826, 17961, 24510, 2495, 2538,
2638, 2639, 2654, 3221, 4103, 4104, 4106, 4107, 4108, 4109, 45713, 5277, 5532, 59140,
59147, 59158, 59296, 59297, 59298, 59299, 59301, 59303, 59304, 6716, 6779, 6782, 6785,
6992, 89780, 89781, 92953, 94796, 94797, 94799, 96255, 96550, 97029, 97143, ZIH9;
Naturhistorisches Museum Vienna: NMW 1779, NMW 20516, NMW 25124, NMW 3081, NMW
3105/ST 663, NMW 3106, NMW 3107, NMW 3111/ST 665, NMW 3119, NMW 3948, NMW
7136, NMW 793/ST 1647; Přírodovědecké muzeum Prague: NMP 09605, NMP 10588, NMP
10784, NMP 22891, NMP 22892, NMP 22893, NMP 23283, NMP 23284, NMP 23295, NMP
23296, NMP 23297, NMP 24474, NMP 24475, NMP 46815, NMP 46816, NMP 46816-b, NMP
47007, NMP 47656, NMP 49711, NMP 50432, NMP 94205, NMP 94564, NMP 94957, NMP
95098; Senckenberg Naturhistorische Sammlungen Dresden: MTD B11877, MTD B12034, MTD
B12062, MTD B12099, MTD B12101, MTD B12177, MTD B12178, MTD B1384-A.S. 1289, MTD
B14244, MTD B15789, MTD B1607-A.S. 1690, MTD B247-A.S. 216, MTD B249-A.S. 214, MTD
B251-A.S. 231, MTD B253-A.S. 221, MTD B266-A.S. 211, MTD B281-A.S. 239, MTD B287-A.S.
200, MTD B288-A.S. 198, MTD B3686, MTD B4188, MTD B4786, MTD B4788, MTD B4789, MTD

B4793, MTD B61-A.S. 244; Museum für Naturkunde Berlin: ZMB_Mam_108652,
ZMB_Mam_11637, ZMB_Mam_11638, ZMB_Mam_12799, ZMB_Mam_14644,
ZMB_Mam_17011, ZMB_Mam_24838, ZMB_Mam_30755, ZMB_Mam_31617,
ZMB_Mam_31621, ZMB_Mam_37523, ZMB_Mam_45130, ZMB_Mam_48173,
ZMB_Mam_83519, ZMB_Mam_83522, ZMB_Mam_83547, ZMB_Mam_83606,
ZMB_Mam_83607, ZMB_Mam_83617, ZMB_Mam_83642, ZMB_Mam_83643,
ZMB_Mam_83647, ZMB_Mam_83648, ZMB_Mam_83653, ZMB_Mam_83675,
ZMB_Mam_83681, ZMB_Mam_83682, ZMB_Mam_83685; Museum Koenig Bonn:
ZFMK_MAM1938-0136, ZFMK_MAM1957-0003, ZFMK_MAM1957-0004, ZFMK_MAM1962-
0131, ZFMK_MAM1963-0660, ZFMK_MAM1965-0544, ZFMK_MAM1965-0545,
ZFMK_MAM1965-0546, ZFMK_MAM1965-0547, ZFMK_MAM1965-0550, ZFMK_MAM1976-
0410, ZFMK_MAM1994-0482, ZFMK_MAM1997-0070, ZFMK_MAM1997-0076,
ZFMK_MAM2012-0036, ZFMK_MAM2015-0479, ZFMK_MAM2019-0404, ZFMK_MAM2019-
0405, ZFMK_MAM2019-0407, ZFMK_MAM2019-0408, MAM2019-0410, ZFMK_MAM2019-
0415, ZFMK_MAM2019-0416, ZFMK_MAM2019-0417, ZFMK_MAM2019-0418,
ZFMK_MAM2019-0419, ZFMK_MAM2019-0420, ZFMK_MAM2019-0421; Haus der Natur
Salzburg: HNS-Mam-S-0073, HNS-Mam-S-0075, HNS-Mam-S-0076, HNS-Mam-S-0077, HNS-
Mam-S-0078, HNS-Mam-S-0079, HNS-Mam-S-0082, HNS-Mam-S-0084, HNS-Mam-S-0085,
HNS-Mam-S-0086, HNS-Mam-S-0519, HNS-Mam-S-0524, HNS-Mam-S-0525, HNS-Mam-S-
0530, HNS-Mam-S-0531, HNS-Mam-S-0532, HNS-Mam-S-0533, HNS-Mam-S-0534, HNS-Mam-
S-0535, HNS-Mam-S-0536, HNS-Mam-S-0550, HNS-Mam-S-0742; Staatliches Museum für
Naturkunde Stuttgart: SMNS-Z-MAM-001687, SMNS-Z-MAM-001750, SMNS-Z-MAM-002012,
SMNS-Z-MAM-045995, SMNS-Z-MAM-046000, SMNS-Z-MAM-048948.
356

357
358 **Figure 4. Number of libraries per great ape species included in this study.**
359

DNA extraction and library preparation
All steps from grinding to indexing except the qPCR were performed in laboratories designed
and dedicated only to ancient DNA (aDNA) research while wearing protective clothing and
following aDNA laboratory best practices. DNA was directly extracted from soft tissue. Bone
and teeth were treated with a sandblaster and ground to bone powder using a *MixerMill*
(Retsch). For each specimen, 50 mg of powder was collected. DNA was extracted using an
established protocol used for aDNA[30]. Single-stranded DNA libraries were prepared[31], followed
by a clean-up with the *QIAGEN MinElute PCR Purification Kit*. We performed a quantitative
PCR for calculating the cycle number in the indexing PCR. Indexing was performed in
quadruplicates using *NEBNext Q5U*, followed by a clean-up using the *NucleoMag® NGS Clean-*
up and Size Select kit. Indexes are listed in Table S1. We assessed quantity and quality using an

Invitrogen QubitTM 4 Fluorometer and an *Agilent 4150 TapeStation*. The grinding step for all samples was performed in the Vienna Ancient DNA laboratory of the University of Vienna, while the DNA extraction, library and QC steps were performed for a subset of libraries (n=97, Table S1) in the ancient DNA laboratory at the Universitat Pompeu Fabra in Barcelona, following the exact same protocols and best practices.

To maximise economic efficiency, we constructed pools of 8-10 libraries, except of four, where a pathological condition of the individual might have influenced the specimen condition. If DNA concentration was below 5ng/µl per sample, we performed another amplification, to preserve sufficient amounts of library. If the concentration was above 25 ng/µl, a dilution was required. Between 8 and 10 libraries were pooled by equal concentration, whereby it was crucial to avoid pooling those with overlapping P5 or P7 adapters. In total, 210 libraries were pooled in 24 pools. We used the same concentration threshold for un-pooled libraries.

Hybridization capture

As aDNA or historical DNA does not only contain host and host-associated microbiome DNA, but often an overwhelming abundance of bacterial and environmental DNA, shotgun sequencing is usually economically infeasible for viruses[32], and target-specific approaches can help in enrichment of sequences of interest[18]. We designed a capture set containing 99 different viruses from 13 families, whereby 49 were human-infecting, 18 great ape-infecting, and the remaining were isolated from other primate species. The design was based on reference genomes available in NCBI, and commercially produced by *myBaits®*, where a BLAST search against the human reference genome was performed to exclude sequences with any hits to it; baits were designed to be 80 nucleotides long with a 2X tiling. The viruses and the NCBI reference sequence are reported in Table S2. We followed the protocol for the capture provided by the manufacturer (Version 5.03, as ordered in July 2021). Briefly, after adding the blockers and the hybridization mix with the RNA baits, the libraries were incubated for approximately 40 hours at 60°C, to increase the efficiency of the capture reaction and to allow for higher sequence deviation from the baits. DNA was eluted from the beads in 30 µl Buffer E, and the supernatant was kept. A qPCR of the capture product was performed in order to estimate the yield, and another PCR to amplify the capture product. Libraries were pooled to 20ng total DNA, and single-end sequencing was performed on the *Illumina NovaSeq 6000 (SP SR100 XP workflow)* at the Vienna BioCenter. The targeted amount of sequencing reads per library was around one million reads.

Bioinformatic processing

Adapters were trimmed from the fastq files with trimmomatic[33] (version 0.39), and BBmap (version 39.01) clumpify was used to remove duplicates introduced by PCR amplification[34]. To determine the metagenomic composition of the sequenced libraries, taxonomic classification via Kraken2 using the standard database was performed[35]. This database contains the strains included in the capture design. Heatmaps were plotted via a customized python3 script for the target taxa at species and genus levels.

Where Kraken2 assigned more than 500 reads to one of the reference genomes included in the capture kit, we performed a mapping with BWA[36] (version 0.7.17), using bwa aln (with parameters "-n 0.04 -l 1000"). We performed SNP calling to the respective reference genomes presented in Table 1 using freebayes[37], and used bcftools consensus[38] (with parameters '-a "N" --exclude 'FILTER="LOWQUAL"') to obtain consensus sequences reported in Supplementary Material SM3, with sequences for the monkeypox virus based on results with higher coverage in a separate study[25]. Mapping coverage along the reference genome was visualized using aDNA-BAMplotter[39], and inspected individually. In case of unequal coverage along the genome, we performed a literature search for alternative genomes obtained from great apes. Summary statistics were calculated using these best reference genomes, including edit distance, mapping quality, mapping quality ratio, and the percentage for 1-, 2- , and 10-fold

423　coverage, using a customized python3 script. Other figures were created with the R package
424　ggplot2[40].
425
426
427

References

1.　Orlando, L. *et al.* Ancient DNA analysis. *Nat. Rev. Methods Prim.* **1**, 14 (2021).

2.　Raxworthy, C. J. & Smith, B. T. Mining museums for historical DNA: advances and
challenges in museomics. *Trends Ecol. Evol.* **36**, 1049–1060 (2021).

3.　Blair, M. E. Conservation museomics. *Conserv. Biol.* **n/a**, (2023).

4.　Duchêne, S., Ho, S. Y. W., Carmichael, A. G., Holmes, E. C. & Poinar, H. The Recovery,
Interpretation and Use of Ancient Pathogen Genomes. *Curr. Biol.* **30**, R1215–R1231
(2020).

5.　Mühlemann, B. *et al.* Ancient hepatitis B viruses from the Bronze Age to the Medieval
period. *Nature* **557**, 418–423 (2018).

6.　Arthur, K. *et al.* Ten millennia of hepatitis B virus evolution. *Science (80-.).* **374**, 182–
188 (2021).

7.　Mühlemann, B. *et al.* Diverse variola virus (smallpox) strains were widespread in
northern Europe in the Viking Age. *Science (80-.).* **369**, eaaw8977 (2020).

8.　Fiddaman, S. R. *et al.* Ancient chicken remains reveal the origins of virulence in Marek's
disease virus. *Science (80-.).* **382**, 1276–1281 (2023).

9.　Alves, J. M. *et al.* Parallel adaptation of rabbit populations to myxoma virus. *Science
(80-.).* **363**, 1319–1326 (2019).

10.　Calvignac-Spencer, S., Düx, A., Gogarten, J. F., Leendertz, F. H. & Patrono, L. V. Chapter
One - A great ape perspective on the origins and evolution of human viruses. in (eds.
Kielian, M., Mettenleiter, T. C. & Roossinck, M. J. B. T.-A. in V. R.) vol. 110 1–26
(Academic Press, 2021).

11.　Calvignac-Spencer, S., Leendertz, S. A. J., Gillespie, T. R. & Leendertz, F. H. Wild great
apes as sentinels and sources of infectious disease. *Clin. Microbiol. Infect.* **18**, 521–527
(2012).

12.　Wertheim, J. O. *et al.* Discovery of Novel Herpes Simplexviruses in Wild Gorillas,
Bonobos, and Chimpanzees Supports Zoonotic Origin of HSV-2. *Mol. Biol. Evol.* **38**,
2818–2830 (2021).

13.　van der Valk, T., Díez-del-Molino, D., Marques-Bonet, T., Guschanski, K. & Dalén, L.
Historical Genomes Reveal the Genomic Consequences of Recent Population Decline
in Eastern Gorillas. *Curr. Biol.* **29**, 165-170.e6 (2019).

14.　Beatrix, K. *et al.* Local Virus Extinctions following a Host Population Bottleneck. *J. Virol.*
89, 8152–8161 (2015).

15.　Fontsere, C. *et al.* The genetic impact of an Ebola outbreak on a wild gorilla population.
BMC Genomics **22**, 735 (2021).

16.　Tan, C. C. S., van Dorp, L. & Balloux, F. The evolutionary drivers and correlates of viral
host jumps. *Nat. Ecol. Evol.* (2024) doi:10.1038/s41559-024-02353-4.

17.　van der Valk, T., Lona Durazo, F., Dalén, L. & Guschanski, K. Whole mitochondrial
genome capture from faecal samples and museum-preserved specimens. *Mol. Ecol.
Resour.* **17**, e111–e121 (2017).

18.　Spyrou, M. A., Bos, K. I., Herbig, A. & Krause, J. Ancient pathogen genomics as
an emerging tool for infectious disease research. *Nat. Rev. Genet.* **20**, 323–340 (2019).

19.　Furtwängler, A. *et al.* Comparison of target enrichment strategies for ancient pathogen
DNA. *Biotechniques* **69**, 455–459 (2020).

20.　Moustafa, A. *et al.* The blood DNA virome in 8,000 humans. *PLOS Pathog.* **13**, e1006292
(2017).

21.　Wylie, T. N., Wylie, K. M., Herter, B. N. & Storch, G. A. Enhanced virome sequencing

475    using targeted sequence capture. *Genome Res.* **25**, 1910–1920 (2015).

476 22. Patterson Ross, Z. *et al.* The paradox of HBV evolution as revealed from a 16th century
477    mummy. *PLOS Pathog.* **14**, e1006750 (2018).

478 23. De Manuel, M. *et al.* Chimpanzee genomic diversity reveals ancient admixture with
479    bonobos. *Science (80-.).* **354**, (2016).

480 24. Fontsere, C. *et al.* Maximizing the acquisition of unique reads in non-invasive capture
481    sequencing experiments. *Mol. Ecol. Resour.* 1755–0998.13300 (2020)
482    doi:10.1111/1755-0998.13300.

483 25. Hämmerle, M. *et al.* Link between Monkeypox Virus Genomes from Museum
484    Specimens and 1965 Zoo Outbreak. *Emerg. Infect. Dis. J.* **30**, 815 (2024).

485 26. Njouom, R., Mba, S. A. S., Nerrienet, E., Foupouapouognigni, Y. & Rousset, D. Detection
486    and characterization of hepatitis B virus strains from wild-caught gorillas and
487    chimpanzees in Cameroon, Central Africa. *Infect. Genet. Evol.* **10**, 790–796 (2010).

488 27. Bonvicino, C. R., Moreira, M. A. & Soares, M. A. Hepatitis B virus lineages in mammalian
489    hosts: Potential for bidirectional cross-species transmission. *World J. Gastroenterol.*
490    *WJG* **20**, 7665 (2014).

491 28. Nakazawa, Y. *et al.* A Phylogeographic Investigation of African Monkeypox. *Viruses* **7**,
492    2168–2184 (2015).

493 29. Fontsere, C. *et al.* Maximizing the acquisition of unique reads in noninvasive capture
494    sequencing experiments. *Mol. Ecol. Resour.* **21**, 745–761 (2021).

495 30. Dabney, J. *et al.* Complete mitochondrial genome sequence of a Middle Pleistocene
496    cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci.* **110**,
497    15758–15763 (2013).

498 31. Kapp, J. D., Green, R. E. & Shapiro, B. A Fast and Efficient Single-stranded Genomic
499    Library Preparation Method Optimized for Ancient DNA. *J. Hered.* **112**, 241–249 (2021).

500 32. Gaudin, M. & Desnues, C. Hybrid Capture-Based Next Generation Sequencing and Its
501    Application to Human Infectious Diseases. *Front. Microbiol.* **9**, (2018).

502 33. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
503    sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

504 34. Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner.

505 35. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
506    *Genome Biol.* **20**, 257 (2019).

507 36. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler
508    transform. *Bioinformatics* **26**, 589–595 (2010).

509 37. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read
510    sequencing. *arXiv Prepr. arXiv1207.3907* 9 (2012) doi:arXiv:1207.3907 [q-bio.GN].

511 38. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008
512    (2021).

513 39. Guellil, M. MeriamGuellil/aDNA-BAMPlotter: aDNA-BAMPlotter. (2021)
514    doi:10.5281/zenodo.5676093.

515 40. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York,
516    2009).

Acknowledgements

520 The computational results of this work have been achieved using the Life Science Compute
521 Cluster (LiSC) of the University of Vienna. We are grateful to the Zoologisches
522 Forschungsmuseum A. Koenig, Leibniz-Institut zur Analyse des Biodiversitätswandels in Bonn,
523 in particular Eva Bärmann, Jan Decher, and Christian Montermann at the Section Theriology;
524 to Irina Ruf and Katrin Krohmann at the Mammalogy collection at Senckenberg
525 Forschungsinstitut und Naturmuseum Frankfurt/M.; to Frank Zachos and Alexander Bibl at
526 Naturhistorisches Museum Wien; to Stefan Merker at Zoology at Staatliches Museum für

Naturkunde Stuttgart; to Petr Benda at Department of Zoology at National Museum (Natural
History) Prague; to Clara Stefen and Jens Jakobitz at Mammalogie at Senckenberg
Naturhistorische Sammlungen Dresden; to Robert Lindner at Haus der Natur – Museum für
Natur und Technik in Salzburg; to Christiane Funk and Frieder Mayer at the mammalian
collection at Museum für Naturkunde/Leibniz-Institut für Evolutions- und
Biodiversitätsforschung in Berlin.
533
534   This project has been funded by the Vienna Science and Technology Fund (WWTF)
535   [10.47379/VRG20001] and by the Austrian Science Fund (FWF) [FW547002] to M.K. K.G.
536   received support from the Swedish Research Council (VR) through grant 2020-03398.
537
538
539

Author contributions

M.K. conceived the topic of the study, supervised data analysis and wrote the manuscript.
M.H. performed experiments and data analysis and wrote the manuscript. R.P. conceived the
study and supervised the experimental work. M.G. supervised data analysis. S.C.-S. wrote the
manuscript. S.S., E.L. and O.C. supervised experimental work. I. R.-G. performed experiments
and contributed to the design of the dataset. P.B. performed experiments. A.R., P.B. and L.G.
provided help in analysing data. P.G., T.R., V.J.S., T.M.-B. and K.G. provided help in writing the
manuscript.
548

Data Availability Statement

A custom hybridization capture design for great ape DNA viruses is reported in this publication.
We report the bait design for 99 virus strains of potential relevance to great apes. The final
design can be found as Supplementary Material SM2, the underlying NCBI identifiers are
reported in Table S2, and the NCBI sequences used in fasta format as Supplementary Material
SM1. Raw sequencing data after capture for the 214 individual libraries has been uploaded to
the Short Read Archive under the accession ID PRJEB75038. Consensus sequences of the
discovered viruses are included as Supplementary Material SM3.
557

Additional information

559   The authors declare no competing interests.
560   Usage notes: Data can be reprocessed using the tools described in the Methods section,
561   namely kraken2 for metagenomic classification, bwa for mapping to reference genomes,
562   following the documentation in the associated repository. Custom code for processing and
563   visualisation is available under https://github.com/admixVIE/Great-Ape-DNA-Virome.
564
565
566