Urbanization Affects Nest Material Composition and Use of Anthropogenic Materials in Blue Tit Nests

Joseph Roy, Davide Dominoni

Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow,
UK

<u>Abstract</u>

Due to urbanisation the nest material availability changes and birds are shown to change their
nesting behaviour. The study on passerines about the change in nest material use and its implications
is scarce. We studied the changes in the nest composition of blue tits as a comparison between
urban and rural habitats and assessed its implications on their breeding success. There was a
significant decrease in the number of natural materials and an increase in anthropogenic materials in
urban nests. The amount of moss acts as a strong predictor of fledgling success. Earlier urban nests
showed a higher amount of anthropogenic material, which could be an indication of parent quality.
In the rural habitats, the presence of anthropogenic materials along with feathers might show
preferential selection as an extended sexual phenotype, while in urban habitats, the anthropogenic
materials is often present as the structural component of the cup. Quantifying the availability of
natural materials is needed to allow a clearer interpretation of the preferential selection of
anthropogenic materials in urban habitats.

    Introduction

The process of urbanization can have various impacts on ecosystems, such as the elimination of

natural vegetation and the construction of artificial and impermeable structures (Sepp et al., 2018),

the introduction of pollutants, including light and chemicals (Dominoni et al., 2013), and the

increasing levels of human disturbance (Sol et al., 2013). There may be relaxation in ecologically

significant factors such as predation pressure (Eötvös et al., 2018), food availability (such as an

increase in lower quality food) (Marzluff, 2001) and temperature (due to the urban heat island

effect) (Puppim de Oliveira et al., 2014). Whether a species can thrive in an urban environment

depends on how it perceives the changes brought about by its new habitat. Some animals can adapt

and take advantage of the opportunities presented become exploiters, and adapt and survive.

Unfortunately, some animals may experience a population decline and avoid the area altogether.

(Kark et al., 2007; González-Lagos & Quesada, 2017; Corsini et al., 2021). The urban adapters might

have to change their life histories to adapt to the conditions and they will have to utilize the available

resources. Urban dwellers adapt and adjust their life histories to cope with changes in their

environment. Changes in circadian rhythm, foraging behaviour, communication strategies, vigilance

behaviour and changes in predator response are some examples (Marzluff, 2001; Sol et al., 2013;

Tätte et al., 2019).

The way birds build their nests is a crucial part of their life, and many studies have shown that

nesting behaviour can change in urban areas. These changes can involve using less natural materials,

altering the structure of the nest, and incorporating man-made materials (Antczak et al., 2010; Votier

et al., 2011). Numerous similar observations have been documented, indicating the widespread

impact of human actions on a global level (Jagiello et al., 2019). The term "anthropogenic nest

material" refers to items not naturally found in nature but created by human actions. Materials such

as plastic, string, and coir are examples of anthropogenic nest materials (Briggs et al., 2023). The

incorporation of human-derived materials is shown to have a positive correlation with the human

footprint index (Z. Jagiello et al., 2019). In urban environments, the availability of natural nest

materials is often reduced, leading birds to use debris as a replacement (Wang et al., 2009).

Synanthropic birds (those that thrive in environments modified by human activity, such as cities) may

incorporate more debris into their nests than non-synanthropic birds (those that do not adapt well to

human-modified environments) (Z. Jagiello et al., 2019). This raises the question of why birds use

anthropogenic materials and how it affects their life histories.

Jagiello et al., 2022 provided three major hypotheses that explain anthropogenic material use in bird

nests. 1) The availability hypothesis, which states that the availability of anthropogenic materials and

the absence of natural materials causes the increased use of anthropogeic materials. A study that

looked at the preference for nest material in Pied Flycatchers found that when available, the birds

preferred natural material over human-derived material (Briggs et al., 2023). 2) The age hypothesis,

which puts forward the concept that the age of parent birds and their experience is connected to the

use of anthropogenic material, where old and experienced birds use it often as an extended sexual

phenotype (Sergio et al., 2011). 3) The functional hypothesis, which states that the functional

advantage of the anthropogenic materials leads to their use in nests.

Several advantages have been reported for the use of human-derived materials in bird nests.

Undoubtedly, plastic strings have their advantages, as they play a crucial role in enhancing the nest

structure, which is a key factor in safeguarding the eggs and fledglings from harsh weather elements

like the wind (Deeming, 2004). Manipulating these materials during nest building could be

significantly easier than using natural alternatives (Esquivel et al., 2020). Shrikes, for instance, are

known for impaling various objects and using them for construction. Another benefit of having many

man-made materials readily available in the area is that it can lower the effort and time spent

searching for nesting materials, leading to faster nest building (Antczak et al., 2010). Cigarette butts

are found to have a repelling effect comparable to green materials, which could decrease the

number of ectoparasites found in city bird nests (Suárez-Rodríguez et al., 2013). A study in Black kites

showed that older birds showed more tendency to incorporate plastic material in their nests up to an

age and they stopped using plastic after senescence. This suggests that the use of plastic might be

used as a signal of sexual quality (Sergio et al., 2011).

Even though there are advantages to the use of anthropogenic materials, there are several potential

negative impacts. Modifications of nests might alter their structural and functional properties

including integrity, camouflage and thermoregulation (Mainwaring et al., 2014; Lopes et al., 2020).

Suárez-Rodríguez et al., 2013 looked at the effects of cigarette butts and its effects on breeding

success and found that even though butts have ectoparasite-repellent effects, the number of

cigarette butts in the nest was positively associated with the level of female genotoxic damage in

house finches during breeding. The study also found that the negative effects of cigarette butts on

the breeding success of urban birds outweigh any potential benefits. Entanglement of chicks in the

plastic debris is reported in a study of a colonial seabird Northern Gannet (Morus bassanus) (Votier

et al., 2011). Plastic waste can also cause chemical harm through the release of toxins, leachates, and non-degraded persistent organic pollutants which can enter food webs via a process known as trophic transfer (Teuten et al., 2009). As mentioned, camouflage is an important characteristic to prevent nest predation. Predation is a major cause of nest failure and several studies have shown the effects of artificial nesting materials on predation rates (Corrales-Moya et al., 2023). In their 2010 study, Antczak et al. noted in their study of great grey shrikes (*Lanius excubitor*), two nests that had entire white plastic strings, made them easier to be detected by predators. In a study on clay-coloured thrush (*Turdus grayi*), the proportion of the outer layer of the nest covered by artificial materials was negatively correlated with the nest's daily survival rate, indicating that the more exposed artificial material on the nest, the higher the chance of it being predated (Corrales-Moya et al., 2023). The study analysed whether the internal or external dimensions of the nest had any impact on the nest survival rate. The findings revealed that variations in the nest dimensions did not affect predation events while the only factor that had an effect was exposed anthropogenic material. Although plastic is considered a thermal insulator, in hotter climates this might act negatively causing thermal risk for the chicks by increasing the temperature in the nests. In American crow (*Corvus brachyrhynchos)*; breeding success was significantly low in nests with higher amounts of plastic. A major reason for the reduction was the entanglement of the chicks, but even in un-entangled nests, the success was low owing to the negative effects of plastic in nests (Townsend & Barker, 2014). A similar case of strangling was reported in Great Gray Shrikes (*Lanius excubitor*) and it is suggested that both adults and the chicks are prone to strangling (Antczak et al., 2010).

Although several studies have looked into the effect of anthropogenic materials in breeding success, most of the studies have focused on marine habitats and sea birds and only few on passerines (Jagiello et al., 2019). A study on blue tits and great tits found that the presence of anthropogenic materials was negatively correlated to breeding success in blue tits but not in great tits (Jagiello et al., 2022). In the current stydy we aim to undestand the effect of urbanization the availability of different nest materials and how it these changes affects the breeding success.

We collected 64 blue tit (*Cyanistes caeruleus*) nests and breeding data from the city of Glasgow (n=31) and the nearby forests of Loch Lomond and Trocssachs National park (n= 33), UK. Nest dissection was carried out to collect the nest material data while the environmantal data was extracted using geographcal information system. Analysis was carried out on the changes on nest materials as a function of urbanization and the effects of the nest material change in breeding success.

Methods

Monitoring of Breeding Success

The study was carried out in the urban-rural gradient at the University of Glasgow, taking 9 sites from the extremes of the 20-site gradient spanning 35 miles from urban green spaces in the city and the Ross woods of the Scottish Centre for Ecology and the Natural Environment located at the east banks of Loch Lomond. Over the past few decades, approximately 500 Schwegler Woodcrete nest boxes with 32mm entrance diameters have been placed along this gradient and monitored for hole-nesting birds (Reid, 2020). From April 1st to June 15th, 2023, breeding success was monitored across all sites. Once the birds started nesting, all the boxes were checked once a week to assess the progress of the nest building and incubation. Close to the expected hatching time we started to check the active nest every other day until chicks were found. Once the chicks were 10 days old, they were ringed by BTO licenced ringers, and morphological measures (wing length, tarsus length and weight) were collected. Once the chicks were fledged the data for breeding success was collected by counting the number of unhatched eggs and dead chicks in the nest boxes and the nests were collected in plastic zip-lock bags.

Site Name	Location (central point)		Number of Nests Collected	Habitat
	Latitude	Longitude		
GCU	55.8669	-4.2506	4	Urban
St Mungo Avenue	55.8655	-4.2465	3	Urban
Kelvingrove Park	55.8700	-4.2859	14	Urban
Old Station Park	55.8803	-4.3059	1	Urban
Dawsholm Park	55.8985	-4.3152	4	Urban
Garscube	55.9043	-4.3186	5	Urban
Cashel	56.1093	-4.5779	8	Rural
Sallochy	56.1238	-4.6008	8	Rural
SCENE	56.1299	-4.6171	17	Rural

140 Table 1- Table showing the sites, their coordinates, the number of nests collected from each site and

141 their respective habitats. From 6 urban sites and 3 rural sites, a total of 64 nests were collected.

143 <u>Nest dissection</u>

144 Once the nests were collected, they were frozen for at least 12 hours to reduce the activity of

145 arthropods and they were kept frozen until the time of dissection. Before dissection, the nests were

146 dried in a heat chamber at a temperature of 50 degrees Celsius for at least 12 hours to prevent

147 moisture in the nests. Once dried the nests were dissected and each category was measured up to an

148 accuracy of 0.001g. The materials were grouped into four categories: moss and grass (natural

149 materials), feathers, wool and hair (naturally derived), and anthropogenic material (together they

150 will be referred to as nest materials from now on). As these materials are intertwined intricately,

151 maximum efforts were taken to separate the materials into each class.

154 *Figure 1- Picture showing different categories of nest materials.*

Figure 2- Images of nests with anthropogenic materials. On the left- a nest with the whole cup made of human-derived fibres. The right image shows a multicoloured nest.

Environmental Variables

Habitat was selected as an environmental variable with two levels (urban and rural) to represent that

the nests were collected from the far extremes of the urban-rural gradient. We also extracted the

impervious surface area of around 100m, which was calculated as a percentage (IS100 from now on)

of the nest box from available land cover maps (Imperviousness — Copernicus Land Monitoring

Service, 2018). This variable was used by Z. Jagiello et al., 2022 in a similar study that looked at

anthropogenic materials. The impervious surface area was selected as a proxy for urbanisation.

Statistical Analysis

The statistical analysis was performed using version.string R version 4.3.3 (2024-02-29 ucrt)-- "Angel

Food Cake". The analysis was conducted in two major parts.

Nest Materials and Urbanization; As our sites were focused on the two extremes of the gradient, we

looked at each material between urban and rural habitats. Boxplots were created using the ggplot2

package to visualize these differences. To assess the environmental effect of the amount of nest

materials, we used Linear Mixed-effect Models (LMMs) with each nest material as the response

variable, and the site as a random variable. Two sets of models were run with the following fixed

explanatory variables

1) habitat (urban, rural) and first egg date

2) IS100 and first egg date.

The interactions between explanatory variables were considered and kept in the model if they were
significant; if not, only linear terms were kept. The predictive visualizations were created using the
plot_model() function in the sjPlot package.

Breeding Success and Nest materials: From the breeding dataset collected during the spring, clutch
size, hatching success and fledgling success were extracted. Generalized Linear Mixed-effect Models
(GLMMs) were used to model clutch size with a Poisson distribution. Hatching success was modelled
as percentages using binomial GLMMs, and fledgling success was also modelled using binomial
GLMMs. In all these models nest materials and the first egg date were added as fixed explanatory
variables and the site as a random variable. The predictive visualizations were created using the
plot_model() function in the sjPlot package.

Results

Nest Materials Overview

In the analysis, I focused on four key variables: total weight, anthropogenic materials, moss and
grass, feathers, wool, and hair, with a total of 64 observations. For the total weight, the values span
from 7.45 to 37.01g, with a median of 19.45 and a mean of approximately 20.35g. Urban nests had
the highest nest weight and had more variability than the rural nests which were around the mean
for total weight. I observed anthropogenic material ranging from 0.00 to 7.89g. The median was just
0.008g, pointing to a skewed distribution with a mean of around 0.45g. Moss and grass, values
ranged from 4.83 to 28.98g, with a mean of 14.38g and a median slightly below the mean at 14.10g.
In the case of feathers, the values spanned from 0.035 to 7.02g. The mean was around 1.06g with a
median of 0.81g. Finally, wool and hair values ranged from 0.00 to 11.44g, showing a mean of 2.29g
and a median of 1.41g. 71.87% of all the nests had the presence of anthropogenic material in them.

Figure 3- Pie charts showing the presence of anthropogenic materials in the nests. There was
anthropogenic material in half of rural nests while 90% of urban nests had anthropogenic materials
in them. In total three-quarters of the nests had anthropogenic materials. As the rural habitats had
ample availability of natural resources, this might point towards an intentional selection of
anthropogenic materials.

Assessing the data set as Urban (N=31) and Rural(N=33), the urban habitat had heavier nests with a
mean of 22 while the rural nests had a mean of 18. Anthropogenic materials were near zero with a
mean of 0.02 and a median of 0.00 for rural while the mean and median for urban nests were 0.92
and 0.40 respectively. Rural habitats had more the weight of moss and grass with a mean value of
15.4 while urban nests had a mean of 13.3. Feather was higher in the urban habitat with a mean of
1.20g while the rural habitat had 0.94g. The combined weight of wool and hair was also higher in the
urban habitat with a mean of 3.41g while the rural habitat had a mean of 1.25g In the rural habitat,
45.45% of nests contained anthropogenic materials while the urban nests had 90.32%. In total

71.88% of nests, I had the presence of anthropogenic materials. However, the mean weight of

anthropogenic materials in the rural habitats is considerably lower (0.02) than in the urban nests

(0.92) while the median is zero as shown in Table 2.

Characteristic	Rural, N = 33[1]	Urban, N = 31
Total	18, (18), (5)	22, (21), (7)
Anthro	0.02, (0.00), (0.05)	0.92, (0.40), (1.70)
Moss+ Grass	15.4, (14.5), (5.1)	13.3, (12.3), (4.9)
Feather	0.94, (0.55), (1.16)	1.20, (1.00), (1.20)
Wool+ Hair	1.25, (0.30), (1.80)	3.41, (3.03), (3.05)

1Mean, (Median), (SD)

226   *Table 2- Table representing the weight of each nesting material considered between habitats. All the*
227   *parameters other than Moss and Grass were high in urban habitats. Only anthropogenic materials*
228   *show a statistically significant variation at habitat level.*

Figure 4- Representation of Nest Materials from urban-rural habitats. [A] Shows the total weight of nests, Rural habitat has fewer heavy nests while their weights showed uniformity around 20g while urban nest weights are variable. Anthropogenic materials were significantly higher in urban habitats and other materials did not show statistical significance.

Effect of Urbanization on Nest Materials

In the models with the nest materials as response variables and the interaction between first egg date and habitat (urban/rural) as fixed explanatory variables, models showed no significant effect on all the materials other than anthropogenic materials. When interactions were not significant, only linear terms were kept in the model. For Anthropogenic materials, urban habitat showed a positive correlation (estimate= 12, p-value= 0.05) and the interaction between urban habitat and first egg date also showed a marginal significance with a slight negative estimate (estimate= -0.09 p-value= 0.07). The random effect site showed a variance of 0.23. Even though non-significant, the total weight and weight of moss also showed a similar trend with the urban nest's weights declining as the days progressed further while the nest weight stayed relatively stable in the rural habitats. Summary tables are added in the supplementary material.

Figure 5- Predictive graph of anthropogenic materials and total nest weight predicted by first egg
date in both urban and rural habitats. The solid red line represents fitted linear regression and the
shaded area represents 95% confidence intervals. In urban habitats, both anthropogenic materials
and total weight drastically reduce as the first egg date progresses. Anthropogenic materials show a
marginal significance.

For assessing urbanization as a continuous variable, the interaction between IS 100 and the first egg

date was selected as an explanatory variable and the site was added as a random variable. No

interactions were significant and hence only linear terms were kept in the final model. For

anthropogenic materials, the first egg date showed a statistically significant negative trend

(estimate= - 0.42, p-value= 0.01). In the case of Moss and Grass, IS100 showed a significant negative

trend with the increase in impervious surface area (estimate= -2, p-value= <0.01).

Figure 6- The solid red line represents fitted linear regression and the shaded area represents 95%
confidence intervals, Left; is the predictive graph of anthropogenic materials (g) predicted by the first
egg date. There is a statistically significant reduction in the weight of anthropogenic materials with
the change in the first egg date. Right; the predictive graph of the change in the weight of moss with
an increase in impervious surface area in 100m. There is a statistically significant reduction in the
weight of moss with the increase in the amount of impervious surface area.

Breeding Success and Nest Materials

The Generalized Linear Mixed Models (GLMMs) were run for clutch size, hatching success and

fledging success. The models with nest materials and first egg date showed no statistically significant

results for clutch size and hatching success. However, for fledgling success, the model with nesting

materials and the first egg date showed a significant result for moss. By model selection using the

Akaike information criterion, only moss was retained in the model, and moss showed a positive

effect on fledgling success with a marginal significance (estimate= 0.26, p-value= 0.06). The detailed

results and model selection tables are added in supplementary material.

Figure 7 - Predictive graphs on fledgling success and the variation in the weight of moss. The solid red
line represents fitted linear regression and the shaded area represents 95% confidence intervals.
Fledging success is increased with the increase in the weight of Moss and Grass. (Summary tables
added in supplementary material)

Discussion

The findings from the present study shed light on the influence of nest materials and urbanisation on

bird populations and their breeding success. The study had 33 nests from the rural environment and

31 nests from the urban environment. In summary, the rural nests were significantly heavier and had

more natural materials than their urban counterparts; more anthropogenic materials were seen in

the urban nests, and feathers and higher wool and hair did not show any significant results.

Looking at the individual materials, the combination of moss and grass (natural materials) was the

only material that showed a significant reduction with an increase in impervious surface area. Moss

and grass emerge as significant predictors of fledgling success which indicates the essential role of

natural materials in nest construction, by providing better insulation or structural support. Also, the

total nest weight is driven by the weight of moss and grass, while other materials only add

incremental differences. In these models, anthropogenic materials did not show a significant effect

on breeding success when it was assessed along with other materials. This underlines the

importance of natural materials in the breeding outcome.

Even though anthropogenic material was not significant, there was a notable increase in the amount

of anthropogenic materials in the urban nests. It is to be noted that 72.8% of all the nests had the

presence of anthropogenic material in them. We have noted that in urban nests, anthropogenic

material was mostly used in cups to provide structural support instead of grass (J.Roy pers obs). The

reduced availability of grass might be the reason for birds to use anthropogenic materials in their

nests. The anthropogenic materials were mostly fibres from tennis balls, fabrics and plastic threads

which serve the purpose of providing structural integrity. The selective picking of nesting material

was demonstrated by Briggs et al., 2023 and it was observed in several previous studies. Even though

urbanization can be considered as a proxy for the reduction of materials like moss and grass, it can

only be confirmed if these materials are selected out of choice or out of need. The amount of

anthropogenic material was predicted to increase the number of unhatched eggs Jagiello et al.,

2022; James Reynolds et al., 2019. It is to be noted that this study mostly focused on anthropogenic

material while in our results moss had the most effect on breeding success. James Reynolds et al.,

2019 suggest city-specific differences in nesting behaviours, the changes we observed can be due to

differences in the availability of materials to the different micro-habitat which is offered by nesting

locations. Suárez-Rodríguez et al., 2013, in a similar study, reported cigarette butts in blue tit nests as an insect repellent, no nests in our sample had cigarette butts in them.

Several concepts try to explain the reasons why birds incorporate anthropogenic materials in their nests. When provided with natural materials, birds showed a preference for natural materials over anthropogenic nest materials (Lee et al., 2015). However, in our study, the three rural sites were oak woodlands with ample amounts of natural materials and the anthropogenic materials were scarce and would need an active search to find them (J.Roy pers obs). Still, 54.5% of rural nests had anthropogenic materials in them (*Figure*) even though the weights were considerably lower than in urban nests as shown in *Table* . The presence of anthropogenic materials in the rural nests, where natural materials are readily available, might show a preferential selection towards anthropogenic material in rural habitats as observed by Briggs et al., 2023. As previous studies have suggested, birds tend to use anthropogenic materials as an extended sexual phenotype. In the case of rural habitats, the anthropogenic materials were mostly used along with feathers on the lining which might play a decorative role in the nests. Such a preferential selection of anthropogenic materials could show a selective preference and potentially use anthropogenic materials as an extended sexual phenotype. Feathers in nests are associated with increased thermal regulation and resistance against microbial and ectoparasite infections (Järvinen & Brommer, 2020). On these accounts, Jagiello et al., 2022 suggested that the chicks from feather-rich nests have a higher chance of recruitment to the breeding population as the number of feathers shows female reproductive quality. As presence of anthropogenic materials in nests might affect mate choice which can in turn affect breeding success.

In urban habitats, anthropogenic materials were often incorporated into the structural parts of the nests instead of grass which could mean that birds are using these materials for stabilizing the nest structure. In urban habitats, 90.3% of nests had the presence of anthropogenic materials, also the weight of anthropogenic materials was significantly high in comparison to the rural habitats. The reason for urban birds to use anthropogenic nests cannot be concluded without quantifying the availability of natural nesting material in urban habitats. Although there is a marked increase in the amount of anthropogenic materials, breeding success was mostly driven by the total weight of the natural materials (moss and grass). Also, the earlier nests had a significantly higher weight of anthropogenic material in the city while in the rural areas, the amount of anthropogenic materials stayed low throughout. Earlier laying date is considered to show parent quality and our study has also shown a significantly higher clutch size in earlier nests. As anthropogenic materials are higher in

earlier nests, it is interesting to note that anthropogenic materials might be selected by higher-
quality parents, especially in urban habitats.

Figure 7- A graphical representation of the study results. Early birds have heavier nests, higher
anthropogenic materials, and higher clutch sizes.

The can be a species-specific difference in the functional and structural aspects of the use of
anthropogenic materials. We have noted a higher number of anthropogenic materials in the great tit
nests than in blue tit nests (J.Roy pers.obs.). As seen in *Figure 2*, we found a great tit nest with a
complete cup made of anthropogenic fibres. Jagiello et al., 2022 did not find an effect of
anthropogenic material on great tit breeding success, yet it is interesting to assess how such drastic
changes affect the breeding outcome. Considering predation, both the nests shown in *Figure 2* are
examples of how anthropogenic materials affect the cryptic nature of the nests. As Corrales-Moya et
al., 2023 reported that the predation rates were directly connected to the exposed anthropogenic
material, these nests might increase the risk of predation by exposing the nests to predators.

In the study design moss and grass were weighed together considering them as natural materials.
While moss was present in both the cup and base, the highest percentage was present in the base.
On the other hand, the grass is seen only in the cup and was used as a structural material to keep the
nest together. Mostly anthropogenic material, especially in urban habitats was used in cups instead

of grass to provide structural integrity for the nest. Even though, the combination of combination of

moss and grass showed a significant effect on breeding success it is important to find which item is

contributing to this effect. Further studies are needed to confirm the effect specific to moss, grass

and anthropogenic materials on breeding success. Also, adding the availability of the materials in the

surroundings to the study can be used to assess the selective preference of nesting materials.

Conclusion

Urbanization has a clear impact on the nesting materials chosen by birds. While anthropogenic

materials are being used more frequently in nests, especially in urban environments, they do not

necessarily predict breeding success as significantly as natural materials like moss and grass. This

study emphasizes the importance of understanding how changing environments, especially

urbanized ones, influence avian nesting behaviour and breeding outcomes. Further research is

essential to delve deeper into the specific roles of individual nesting materials and the choices birds

make in different habitats.

Acknowledgements

References

Antczak, M., Hromada, M., Czechowski, P., Tabor, J., Zabłocki, P., Grzybek, J., & Tryjanowski, P. (2010).

394   A new material for old solutions—The case of plastic string used in Great Grey Shrike nests.

395   *Acta Ethologica*, *13*(2), 87–91. https://doi.org/10.1007/s10211-010-0077-2

396   Briggs, K. B., Deeming, D. C., & Mainwaring, M. C. (2023). Plastic is a widely used and selectively

397   chosen nesting material for pied flycatchers (Ficedula hypoleuca) in rural woodland habitats.

398   *Science of The Total Environment*, *854*, 158660.

399   https://doi.org/10.1016/j.scitotenv.2022.158660

400   Corrales-Moya, J., Barrantes, G., Chacón-Madrigal, E., & Sandoval, L. (2023). A potential consequence

401   for urban birds' fitness: Exposed anthropogenic nest materials reduce nest survival in the

402   clay-colored thrush. *Environmental Pollution*, *326*, 121456.

403   https://doi.org/10.1016/j.envpol.2023.121456

404   Corsini, M., Schöll, E. M., Di Lecce, I., Chatelain, M., Dubiec, A., & Szulkin, M. (2021). Growing in the

405   city: Urban evolutionary ecology of avian growth rates. *Evolutionary Applications*, *14*(1), 69–

406   84. https://doi.org/10.1111/eva.13081

407   Deeming, D. C. (Ed.). (2004). *Avian incubation: Behaviour, environment, and evolution* (Reprinted).

408   Oxford Univ. Press.

409   Dominoni, D., Quetting, M., & Partecke, J. (2013). Artificial light at night advances avian reproductive

410   physiology. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1756), 20123017.

411   https://doi.org/10.1098/rspb.2012.3017

412   Eötvös, C. B., Magura, T., & Lövei, G. L. (2018). A meta-analysis indicates reduced predation pressure

413   with increasing urbanization. *Landscape and Urban Planning*, *180*, 54–59.

414   https://doi.org/10.1016/j.landurbplan.2018.08.010

415   Esquivel, C., De La O, J. M., Sánchez Vargas, S., Paniagua, S., Esquivel Cambronero, A., Núñez, D., &

416   Quesada Ávila, G. (2020). Anthropogenic materials used by birds to nest in an urban

landscape of Costa Rica. *UNED Research Journal*, *12*(2), e3124. https://doi.org/10.22458/urj.v12i2.3124

González-Lagos, C., & Quesada, J. (2017). Stay or Leave? Avian Behavioral Responses to Urbanization in Latin America. In I. MacGregor-Fors & J. F. Escobar-Ibáñez (Eds.), *Avian Ecology in Latin American Cityscapes* (pp. 99–123). Springer International Publishing. https://doi.org/10.1007/978-3-319-63475-3_6

Jagiello, Z., Corsini, M., Dylewski, Ł., Ibáñez-Álamo, J. D., & Szulkin, M. (2022). The extended avian urban phenotype: Anthropogenic solid waste pollution, nest design, and fitness. *Science of The Total Environment*, *838*, 156034. https://doi.org/10.1016/j.scitotenv.2022.156034

Jagiello, Z., Dylewski, Ł., Tobolka, M., & Aguirre, J. I. (2019). Life in a polluted world: A global review of anthropogenic materials in bird nests. *Environmental Pollution*, *251*, 717–722. https://doi.org/10.1016/j.envpol.2019.05.028

James Reynolds, S., Ibáñez-Álamo, J. D., Sumasgutner, P., & Mainwaring, M. C. (2019). Urbanisation and nest building in birds: A review of threats and opportunities. *Journal of Ornithology*, *160*(3), 841–860. https://doi.org/10.1007/s10336-019-01657-8

Järvinen, P., & Brommer, J. E. (2020). Lining the nest with more feathers increases offspring recruitment probability: Selection on an extended phenotype in the blue tit. *Ecology and Evolution*, *10*(23), 13327–13333. https://doi.org/10.1002/ece3.6931

Kark, S., Iwaniuk, A., Schalimtzek, A., & Banker, E. (2007). Living in the city: Can anyone become an ?urban exploiter'? *Journal of Biogeography*, *34*(4), 638–651. https://doi.org/10.1111/j.1365-2699.2006.01638.x

Lee, K., Jang, Y. C., Hong, S., Lee, J., & Kwon, I. K. (2015). Plastic Marine Debris Used as Nesting Materials of the Endangered Species Black-Faced Spoonbill Platalea minor Decreases by Conservation Activities. *Journal of the Korean Society for Marine Environment & Energy*, *18*(1), 45–49. https://doi.org/10.7846/JKOSMEE.2015.18.1.45

Lopes, C. S., De Faria, J. P., Paiva, V. H., & Ramos, J. A. (2020). Characterization of anthropogenic

materials on yellow-legged gull (Larus michahellis) nests breeding in natural and urban sites

along the coast of Portugal. *Environmental Science and Pollution Research*, *27*(29), 36954–

36969. https://doi.org/10.1007/s11356-020-09651-x

Mainwaring, M. C., Hartley, I. R., Lambrechts, M. M., & Deeming, D. C. (2014). The design and

function of birds' nests. *Ecology and Evolution*, *4*(20), 3909–3928.

https://doi.org/10.1002/ece3.1054

Marzluff, J. M. (2001). Worldwide urbanization and its effects on birds. In J. M. Marzluff, R. Bowman,

& R. Donnelly (Eds.), *Avian Ecology and Conservation in an Urbanizing World* (pp. 19–47).

Springer US. https://doi.org/10.1007/978-1-4615-1531-9_2

Puppim de Oliveira, J. A., Doll, C. N. H., Moreno-Peñaranda, R., & Balaban, O. (2014). Urban

Biodiversity and Climate Change. In B. Freedman (Ed.), *Global Environmental Change* (pp.

461–468). Springer Netherlands. https://doi.org/10.1007/978-94-007-5784-4_21

Reid, R. (2020). *The relationship between habitat characteristics, food availability and the*

reproductive success of blue tit (Cyanistes caeruleus) and great tit (Parus major) along a

rural-urban gradient. 49.

Sepp, T., McGraw, K. J., Kaasik, A., & Giraudeau, M. (2018). A review of urban impacts on avian

life-history evolution: Does city living lead to slower pace of life? *Global Change Biology*,

24(4), 1452–1469. https://doi.org/10.1111/gcb.13969

Sergio, F., Blas, J., Blanco, G., Tanferna, A., López, L., Lemus, J. A., & Hiraldo, F. (2011). Raptor Nest

Decorations Are a Reliable Threat Against Conspecifics. *Science*, *331*(6015), 327–330.

https://doi.org/10.1126/science.1199422

Sol, D., Lapiedra, O., & González-Lagos, C. (2013). Behavioural adjustments for a life in the city.

Animal Behaviour, *85*(5), 1101–1112. https://doi.org/10.1016/j.anbehav.2013.01.023

Suárez-Rodríguez, M., López-Rull, I., & Macías Garcia, C. (2013). Incorporation of cigarette butts into

nests reduces nest ectoparasite load in urban birds: New ingredients for an old recipe?

Biology Letters, *9*(1), 20120931. https://doi.org/10.1098/rsbl.2012.0931

Tätte, K., Ibáñez-Álamo, J. D., Markó, G., Mänd, R., & Møller, A. P. (2019). Antipredator function of

vigilance re-examined: Vigilant birds delay escape. *Animal Behaviour*, *156*, 97–110.

https://doi.org/10.1016/j.anbehav.2019.08.010

Teuten, E. L., Saquing, J. M., Knappe, D. R. U., Barlaz, M. A., Jonsson, S., Björn, A., Rowland, S. J.,

Thompson, R. C., Galloway, T. S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C., Viet, P. H.,

Tana, T. S., Prudente, M., Boonyatumanond, R., Zakaria, M. P., Akkhavong, K., … Takada, H.

(2009). Transport and release of chemicals from plastics to the environment and to wildlife.

Philosophical Transactions of the Royal Society B: Biological Sciences, *364*(1526), 2027–2045.

https://doi.org/10.1098/rstb.2008.0284

Townsend, A. K., & Barker, C. M. (2014). Plastic and the Nest Entanglement of Urban and Agricultural

Crows. *PLoS ONE*, *9*(1), e88006. https://doi.org/10.1371/journal.pone.0088006

Votier, S. C., Archibald, K., Morgan, G., & Morgan, L. (2011). The use of plastic debris as nesting

material by a colonial seabird and associated entanglement mortality. *Marine Pollution*

Bulletin, *62*(1), 168–172. https://doi.org/10.1016/j.marpolbul.2010.11.009