1   ,

Caspar specifies primordial germ cell count and identity in *Drosophila melanogaster*

Subhradip Das[1,#], Sushmitha Hegde[1,#], Neel Wagh[1], Jyothish Sudhakaran[1], Adheena Elsa Roy[1],

Girish Deshpande[1,2,*] & Girish S Ratnaparkhi[1,*]

[#]Equal Contribution

[1]Department of Biology, Indian Institute of Science Education & Research, Pune, INDIA

[2] Department of Molecular Biology, Princeton University, Princeton, NJ 08544, USA Princeton

University,

[*]Author(s) for Correspondence: girish@iiserpune.ac.in (GR), gdeshpande@princeton.edu (GD)

ORCID IDs: SH (0000-0003-0777-962X); SD (0009-0002-4553-966X); AR (0009-0008-6000-
9572); NW (0009-0006-0639-6116), JS (0000-0001-5828-5182); GR (0000-0001-7615-3140),
GD (0000-0002-5200-7090)

Running Title. Caspar specifies germ cells

Figures: 10

Suppl. Material: 4

Abstract

Repurposing of pleiotropic factors during execution of diverse cellular processes has emerged as a regulatory paradigm. Embryonic development in metazoans is controlled by maternal factors deposited in the egg during oogenesis. Here, we explore maternal role(s) of Caspar (Casp), the *Drosophila* orthologue of human Fas-associated factor-1 (FAF1) originally implicated in host-defense as a negative regulator of NF-κB signaling. Maternal loss of either Casp or it's protein partner, Transitional endoplasmic reticulum 94 (TER94) leads to partial embryonic lethality correlated with aberrant centrosome behavior, cytoskeletal abnormalities, and defective gastrulation. Although ubiquitously distributed, both proteins are enriched in the primordial germ cells (PGCs), and in keeping with the centrosome problems, mutant embryos display a significant reduction in the PGC count. Moreover, the total number of pole buds is directly proportional to the level of Casp. Consistently, it's 'loss' and 'gain' results in respective reduction and increase in the Oskar protein levels, the master determinant of PGC fate. To elucidate this regulatory loop, we analyzed several known components of mid-blastula transition and identify the translational repressor Smaug, a zygotic regulator of germ cell specification, as a potential critical target. We present a detailed structure-function analysis of Casp aimed at understanding its novel involvement during PGC development.

Keywords: Fas-associated factor 1, Germ plasm, Valosin Containing Protein, Maternal, Pole cell, Vasa, P97, MZT, ZGA.

 Introduction

As a model organism, *Drosophila melanogaster*, has been instrumental in establishing

and advancing several developmental paradigms underlying embryonic development.

Drosophila has also emerged as a relatively simple yet tremendously useful model, to analyze

the underpinnings of the immune response (MEDZHITOV 2001; LEMAITRE AND HOFFMANN 2007).

Critically, despite the small size and modest cellular complexity, the insect immune system

shares many fundamental traits with the higher vertebrates, including the humoral and innate

arms and the dialogue between the two (HULTMARK 1993; ADAMS *et al.* 2000). Furthermore,

several factors that contribute to the proper functioning of the insect immune system are highly

conserved and perform analogous functions across evolution (HOFFMANN AND REICHHART 2002;

LEMAITRE AND HOFFMANN 2007).

Interestingly, almost all the protein components of the insect immune system are highly

pleiotropic and involve diverse activities during developmental progression. For example, the

Toll class of proteins belonging to a larger family of pattern recognition receptors are essential

while mounting robust immune response against the microorganismal invasion (BRENNAN AND

ANDERSON 2004; WANG AND LIGOXYGAKIS 2006). The same proteins are also deployed earlier

during embryonic patterning and morphogenesis (GOVIND 1999; HOFFMANN AND REICHHART

2002; VALANNE *et al.* 2011). Evidently, in several instances, different pathway components or

modules used during early development are repurposed to mediate immunity both in the insect

and the mammalian context (BELVIN AND ANDERSON 1996; GOVIND 1999; ROTH 2023).

The involvement of early embryonic morphogen, Dorsal, during humoral response in

flies is a canonical example of the context-specific and diverse functions of immune system

components (GOVIND 1999; ROTH 2023). Maternal loss of function of genes involved in

Toll/Dorsal signaling affect Dorso-ventral patterning (BELVIN AND ANDERSON 1996). A nuclear-

cytoplasmic gradient of Dorsal, set up by asymmetric Toll signalling establishes cell fate across

the Dorsal-ventral axis in the syncytial embryo (NUSSLEIN-VOLHARD 2022) . Pathogenic invasion

of insects induces both the humoral and cellular immune response (BELVIN AND ANDERSON

1996; WILLIAMS 2007). The humoral response consists of the production of antimicrobial

peptides by the fat bodies, which serve as a first line of defence (BELVIN AND ANDERSON 1996;

IMLER AND HOFFMANN 2000). Two Rel family member proteins, Dorsal and Dif, homologous to

the mammalian NF-kappa B, induce the expression of defense peptides (BELVIN AND ANDERSON

1996; GOVIND 1999; BUCHON *et al.* 2014). NF-kappa B is essential for differentiating

lymphocytes, which engineer the acute-phase response. Altogether, undertaking functional

analysis of proteins, in a temporally distinct developmental context, has proven to be highly

informative and insightful. Notably, however, thus far, such analysis has focused on only

activator proteins. Here we have investigated the embryonic function of Caspar (Casp), a

protein involved in inhibiting the immune response (KIM *et al.* 2006).

Casp is an intracellular negative regulator discovered in a genetic screen to identify

suppressors of antibacterial immunity (KIM *et al.* 2006). Flies mutant for *casp* were identified due

to their high rates of melanization, an innate immune response that leads to encapsulation of

pathogens in the gut and fat body. Interestingly, *casp* mutants isolated in this study were

resistant to Gram-negative bacterial infections due to elevated expression of the antimicrobial

peptide (AMP) diptericin; consequently, infected flies survived longer than their wild-type

counterparts. Strikingly, *casp* overexpression inhibited nuclear localization of Rel in response to

infection in the fat body. Excess levels of Casp result in the cytoplasmic retention of Rel in its

uncleaved, inactive form, presumably due to inhibition of the protease, Dredd (KIM *et al.* 2006).

Intriguingly, sequence analysis of Casp indicated a high degree of similarity to the

mammalian Fas-associated factor 1 (FAF1) protein (KIM *et al.* 2006; TENDULKAR *et al.* 2022)

(Fig.1A). FAF1 is evolutionarily conserved and was initially discovered as an interactor of Fas, a

pro-apoptotic member of the tumor necrosis factor receptor family (CHU *et al.* 1995). FAF1 also

interacts with the components of the death-inducing signaling complex (DISC) such as the Fas-

associated death domain (FADD) and Caspase-8 proteins (RYU *et al.* 2003). These interactions

are mediated by the death effector domains (DED) in FADD and Caspase-8 and the DED-

interacting domain (DEDID) in FAF1 (RYU *et al.* 2003). Both 'loss' and 'gain' of function

experiments indicated that FAF1 is crucial in promoting cell death via transduction of the

apoptotic signal (RYU AND KIM 2001; DE ZIO *et al.* 2008). FAF1, like its *Drosophila* ortholog, is a

negative regulator of NFκB signalling (MIN-YOUNG *et al.* 2004; PARK *et al.* 2007).

FAF1's myriad cellular functions can be attributed to its multiple protein-interaction

domains that allow its participation in ubiquitin-related processes such as protein degradation

(SONG *et al.* 2005; MENGES *et al.* 2009; ZHANG *et al.* 2011). Consistently, FAF1 harbors a

Ubiquitin-associated (UBA) domain, a UAS domain and a Ubiquitin-like regulatory X (UBX)

domain (Fig. 1A). Furthermore, the N-terminal UBA domain recruits polyubiquitinated proteins,

leading to their accumulation (SONG *et al.* 2009). The C-terminal UBX domain, on the other

hand, interacts with the molecular chaperone, valosin-containing protein (VCP/p97) bound to

the Npl4-Ufd1 heterodimeric complex (SCHUBERTH AND BUCHBERGER 2008; KLOPPSTECK *et al.*

2012; EWENS *et al.* 2014). FAF1 complexed with VCP-Npl4-Ufd1 and polyubiquitinated proteins

is known to assist endoplasmic reticulum-associated degradation (ERAD) (LEE *et al.* 2013). The

UAS domain, a domain of unknown function, is involved in interaction with long-chain fatty

acids, which is thought to promote the polymerization of FAF1 (KIM *et al.* 2013).

While Casp has a well-established role in the immune response, modENCODE RNAseq

and proteomics data suggest that *casp* is also highly expressed in the 0-3 hour old embryo

(BROWN *et al.* 2014; CASAS-VILA *et al.* 2017). Consistently, a snapshot of *casp* staining in a

high-throughput RNA *in-situ* experiment indicates ubiquitous expression of maternally deposited

casp in the *Drosophila* embryo (WEISZMANN *et al.* 2009). We thus wondered if Casp expression

in early embryos is functionally relevant for proper developmental progression. To explore the

possible developmental function of Casp, we first assessed if Casp function is needed for

viability. Analysis of a hypomorphic allele of *casp* demonstrated that Casp is indeed required

maternally for embryonic development. Consequently, roughly half of the embryos maternally

compromised in *casp*, fail to undergo gastrulation. Furthermore, such embryos display

developmental defects including aberrant cytoskeletal network starting from early blastoderm

stages. Interestingly, Casp is expressed strongly in primordial germ cells (PGCs). Consistent

with the enrichment, maternal reduction of *casp* significantly affects the total number of pole

cells. Here, we present an analysis of *casp* function during early embryonic development and

its role in the formation and/or specification of PGCs. We show that Casp activity regulates

Oskar levels and centrosome function, two critical determinants of PGC fate in *Drosophila*

embryo. Upon loss of *casp*, the total amount of Oskar and Smaug changed reciprocally to

influence the PGC count. Ubiquitin-based protein degradation is critically involved during early

embryonic events, including the maternal-zygotic transition. We present a model explaining the

involvement of Casp and its protein partner, TER94, during germ cell development, considering

their influence on the clearance of Smaug, a critical regulator of maternal to zygotic transition

(MZT).

    Results

Casp^c04227 is a loss of function allele of casp.

To better understand embryonic function of Casp we decided to first characterize a *casp*
allele, *w^1118; pBac(PB)casp^c04227* (Bloomington stock number:11373) (Supplementary Fig. 1A;
labelled as S1A). This allele is induced by a piggyBac insertion in the 5' regulatory region of
CG8400/FBgn0034068/Casp locus which is situated at 2R, region 52D14-15 (THIBAULT *et al.*
2004). As summarized in Fig. 1, our data indicate that *casp^c04227* is a strong hypomorphic allele
and is referred to as a 'loss of function' allele (*casp^lof*) here onwards.

Fig. 1. *casp* is a maternal effect gene. (A) Comparison between human FAF1 and Casp shows conserved protein
domains, which are described in the text. (**B**) Embryos laid by homozygous *casp^lof* females show ~70% lethality,
irrespective of the paternal genotype (*w^1118* or *casp^lof*), suggesting a strictly maternal function of *casp* (**C**) The use of
a Deficiency in the *casp* locus validates the lethal phenotype associated with the mutation, which drops to ~40%. In

208   panels B and C, the parental genotype is listed on the X-axis, with percent embryonic lethality plotted as a bar graph.
209   N = 3, ordinary one-way ANOVA/ unpaired t-test, (***) P <0.001, (*) P <0.05. (**D**) Immunofluorescence images of
210   0-3 h embryos derived from w^{1118} and $casp^{lof}/Df$ females, stained with Hoechst and Casp (**E**) Casp protein levels
211   were assessed in 0-3 embryos laid by w^{1118} and $Casp^{lof}/Df$ animals, evaluated via western blotting. Tubulin is used as
212   a loading control.
213

~70% embryos laid by the homozygous $casp^{lof}$ females failed to hatch into larvae (Fig.

1B). Moreover, the number of embryos that failed to hatch did not change substantially when

$casp^{lof}$ virgin females were mated either with w^{1118} males or $casp^{lof}$ males. That the extent of

viability was independent of the paternal genotype established that $casp$ is a maternal effect

gene and its activity is required for viability (Fig. 1B). To rule out the possibility that the lethality

was induced by non-specific background mutations, we procured multiple deficiency (Df) lines

spanning the 52D locus and confirmed that $casp^{lof}/Df$ mothers also displayed significant

embryonic lethality (40%) (Fig. 1C). Lastly, as in the case of $casp^{lof}$ females, $casp^{lof}$ males are

also semi-fertile which was assessed by mating the mutant males with w^{1118} virgin females (Fig.

1B).

To better understand the function of Casp, we first decided to analyze how $casp$ RNA

and Casp protein are expressed during embryogenesis. Consistent with the maternal effect

lethality, modENCODE (MUERS 2011; CHEN et al. 2014) data suggests that $casp$ mRNA is

deposited maternally, being highly expressed in the 0-2 hours embryo, with a drop in expression

2-4 hours post fertilization (Fig. S1B)(BROWN et al. 2014). Publicly available In-Situ databases

such as FlyAtlas, Fly-FISH and BDGP (WEISZMANN et al. 2009) (TOMANCAK et al. 2002;

LECUYER et al. 2007) also confirmed maternal deposition of $casp$ mRNA, with ubiquitous

expression seen in stage 1-3 embryos. Fly-FISH data (LECUYER et al. 2007; WILK et al. 2016)

further suggests that expression in the pole cells persists while maternally deposited somatic

transcripts are selectively degraded from stage 4 onwards. We immunostained 0-3 hours old

embryos derived from w^{1118} mothers with the anti-Casp antibody (TENDULKAR et al. 2022) and

found that Casp protein was predominantly cytoplasmic, and as in the case of RNA, exhibited a

relatively ubiquitous distribution in the somatic compartment (Fig. 1D). Again, consistent with the

in-situ hybridization pattern, Casp-specific antibody staining appeared to be enriched in the

posteriorly localized primordial germ cells (PGCs) compared to the surrounding soma (Fig. 1D,

arrow). To establish the specificity of the anti-Casp antibodies in the embryonic context, we

stained the embryos laid by both the $casp^{lof}/ casp^{lof}$ and $casp^{lof}/Df$ mothers. As expected, Casp

protein was nearly absent in embryos of both these genotypes suggesting that Casp protein is

significantly reduced in the 0-3 hours old $casp^{lof}$ mutant embryos. Analysis of the embryonic

lysates using Western blotting, suggested that trace amounts of Casp protein (estimated 5-10%

of w^{1118}), were present in embryos laid by $casp^{lof}/casp^{lof}$ and $casp^{lof}/Df$ mothers, supporting the

conclusion that $casp^{lof}$ is a strong hypomorphic allele and not a null (Fig.1E). Interestingly,

mature egg chambers of wild-type animals had Casp expression in somatically derived follicle

cells, but Casp protein could not be readily detected in the egg itself (Fig. S1C). This

observation suggested that, unlike the RNA, only trace amounts of Casp protein may be

maternally deposited. Thus, the protein detected in the young embryos is likely generated post-

fertilization, by the translation of maternally deposited *casp* mRNA.

Casp function is required for early embryonic development.

Larval cuticle patterns are an excellent readout for major patterning defects in

Drosophila (NUSSLEIN-VOLHARD AND WIESCHAUS 1980). To better understand the function of

Casp protein during embryonic development, we analyzed the cuticles of *casp* mutant embryos.

0-3 hours old embryos laid by $casp^{lof}/Df$ mothers were collected and aged for 22 hours to

prepare cuticles (see Material and Methods). The cuticular patterns were visualized under a

dark field microscope. While ~60% of larvae displayed cuticular patterns comparable to wild

type larvae, development in the remaining 40% embryos appeared to have stalled before they

could deposit cuticle. These data were consistent with the extent of maternal effect lethality.

Next, to define the stage of developmental arrest, we performed time-lapse live Bright-field

imaging (CAVEY AND LECUIT 2008). These data (Movies, Fig. S2A) confirmed that 40% of

embryos did not proceed to gastrulation and showed developmental arrest before germ band

extension (around stage 6; Fig. S2, panels C4, D4). Such embryos displayed irregular,

uncoordinated morphogenetic movements with blebbing of the plasma membrane possibly

inhibiting both cephalic furrow formation, and germband elongation (Fig. S2 panels C2-C3, D2-

D3).

Compromising casp activity leads to centrosomal abnormalities in early embryos.

To trace the defects during mid-embryogenesis including failure of gastrulation, we

sought to visualize $casp^{lof}$ embryos at blastoderm stage. We labelled the 0-3 hours old $casp^{lof}$

and control embryos with the nuclear dye Hoechst and the cytoskeletal F-actin marker phalloidin

(Fig. 2A, B). $casp^{lof}$ embryos displayed significant structural abnormalities that are reflected in an

irregular actin network that lacks stereotypical organization, which is partially interrupted in

several places (Fig. 2B1, compared to Fig. 2A1). Similarly, unlike the age-matched control

embryos, regular nuclear spacing is disrupted, and nuclei are unevenly distributed across the

embryo with occasional instances of nuclear fusion and possibly mitotic asynchrony (Fig. 2B vs

Fig. 2A).

Fig. 2. A Significant proportion of embryos (~45%) laid by *casp^{lof}/Df* **mothers display nuclear division and cytoskeletal defects**. (**A-B**) Confocal images of nuclear cycle 13/14 embryos (single sections) of the indicated genotypes stained with Hoechst, phalloidin, and gamma-tubulin. Both the regular arrangement and uniform density of nuclei are disrupted in the mutants, as indicated by a yellow asterisk. F-actin, marked with phalloidin, shows a regular, hexagonal compartmentalization in *w^{1118}* (panel A2), while disorganized F-actin (white arrows; panel B3) is observed in the mutant. Defective centrosomes marked with gamma-tubulin can also be observed in the *Casp* mutant (inset of B3, compare to inset A3), indicated by yellow arrows. The extent of defects is quantified in nuclear cycle(nc) 9/10 (**C**) and nc 12/13 (**D**). (**E-F**) Confocal images of sections of nc 13/14 embryos of the indicated genotypes stained with Hoechst, phalloidin, and gamma-tubulin. The nuclear, cytoskeletal, and centrosomal defects are rescued (panel E) when wild-type *casp* is expressed in the *casp^{lof}/Df* background (compare panels **G-H** to C-D).

330 Overexpression of *casp* on its own (panel F) does not appear to affect nuclear or cytoskeletal architecture. The bar
331 charts represent the percentage of defective centrosomes in nuclear cycle 9/10 and 12/13 embryos. Number of
332 embryos imaged ~15.
333

The nuclear and cytoskeletal defects observed in *casp* embryos prompted us to analyze

the centrosomes in embryos deficient in *casp* function. Centrosomes function as the microtubule

organizing centers, and actomyosin-based cytoskeletal defects have been correlated with

aberrant centrosome activity ((WU AND AKHMANOVA 2017; BLAKE-HEDGES AND MEGRAW 2019)

and references therein). In wild-type nuclei, centrosome duplication occurs simultaneously with

the initiation of the nuclear division cycle. After completion of duplication, centrosomes separate

and migrate along the nuclear membrane to reach the opposite sides of the nucleus (reviewed

by (LATTAO *et al.* 2017; WU AND AKHMANOVA 2017; BLAKE-HEDGES AND MEGRAW 2019)). In the

embryos maternally compromised for *casp*, several characteristic aspects of centrosome

behavior during mitotic divisions are altered (Compare Fig. 2B3 to Fig. 2A3). At times,

centrosomes appeared to divide even without a nucleus (or DNA), and many 'orphan'

centrosomes devoid of nuclear DNA were observed (Fig. 2B3, inset). By contrast, in some

instances, centrosomes were duplicated but remained in proximity indicating failed or

incomplete migration (see arrows in Figure 2B3, inset). Quantitation of the centrosomal

abnormalities in *casp^{lof}* embryos (Fig. 2C, D) further revealed that the behavior of the

centrosomes, especially their ability to separate and migrate to the opposite poles correctly

deteriorated progressively (Compare Fig. 2D to Fig. 2C), as nuclear division cycles advanced.

To confirm the specificity of the phenotypic consequences induced by the maternal loss

of *casp*, we overexpressed *UAS-casp^{wt}* transgene in *casp^{lof}/Df* embryos using *nos Gal4*, a

maternal Gal4 driver. Embryos derived from such mothers were stained using nuclear dye

Hoechst, phalloidin and anti-Gamma tubulin antibodies that mark the centrosomes. As shown in

Fig. 2E, maternal overexpression of *casp* substantially rescues centrosomal and cytoskeletal

abnormalities seen in *casp^{lof}* (compare panels E1-E3 to A1-A3). This rescue is quantitated in

Fig. 2 G, H (compare with Fig. 2C, D). The *casp^{lof}/Df; nos Gal4/UASp-casp^{wt}* showed a rescue

of lethality of 20-25% as compared to 40-45% for *casp^{lof}/Df*.

Taken together, these data showed that Casp plays an essential role in early embryonic

development in *Drosophila*, and loss of <u>casp</u> results in conspicuous nuclear and cytoskeletal

defects that correlate with incomplete gastrulation movements and developmental arrest, for

~45% of embryos. Moreover, such defects do not appear to be region-specific or localized

within an embryo arguing in support of ubiquitous function across the embryo for Casp protein.

Maternal Casp protein is enriched in pole cells and controls total pole cell count in
blastoderm embryos.

Primordial germ cells (PGCs) are precursors of the gametes—sperm and eggs. Their
specification is essential for the development of the germline, which in turn passes genetic
information to the next generation. During early embryonic development, PGCs are specified
and segregated from somatic cells. In *Drosophila*, the PGCs are formed in the posterior end of
the embryo. As the early embryonic syncytial nuclear division cycles progress, a few nuclei and
centrosomes associated with them invade posteriorly localized and anchored pole plasm ahead
of the rest of the somatic nuclei (RAFF AND GLOVER 1989). The precocious entry of the
centrosomes results in the release and microtubule-dependent transport of pole plasm, which is
sequestered in newly cellularized PGCs (RAFF AND GLOVER 1989; LERIT *et al.* 2017). Thus, the
PGCs or the germline stem cell precursors are set aside during early embryogenesis. Moreover,
centrosome behavior and dynamics are crucial for proper cellularization and mitotic cell
divisions of early germ cells (LERIT AND GAVIS 2011; LERIT *et al.* 2017). Centrosomes, the
microtubule (MT)-organizing centers, ensure the faithful segregation of germ plasm, a reservoir
of germ cell determinants, into PGCs. Taken together with the observation that Casp protein is
readily detectable in early PGCs, we wondered if it could influence the total pole cell count. To
examine this possibility, stage 5 embryos derived from *cs* and *casp^{lof}/ Df* mothers were
immunostained for the pole cell marker Vasa and visualized under a confocal fluorescence
microscope. Interestingly, while *cs* and *nos-Gal4* embryos had an average of ~30 pole cells
(Fig 3, panels A1-B3, for quantitation see panel F) *casp^{lof}/ Df* showed a drastic reduction in total
PGC count to ~10 (Fig. 3 panels C1-3, F). By contrast, stage 5 embryos derived from
nosGal4>UAS-casp mothers, where *casp* was overexpressed (Fig. 3G), displayed considerably
elevated total number of PGCs (Fig. 3, panels E1-E3, F). To confirm that the loss of *casp* is
specifically responsible for the reduction in the total number of PGCs, we simultaneously
overexpressed *casp* in the *casp^{lof}/Df* mothers in a germline-specific manner, which resulted in a
significant rescue (~26 pole cells; Fig.3, panels D1-D3, F). Taken together, these data
suggested that the total pole cell number in late syncytial /cellular blastoderm embryos
specifically depends on *casp* function. Furthermore, increasing *casp* levels maternally, is
sufficient to elevate the total pole cell number substantially.

442
Fig. 3. Casp influences total PGC count. (**A-E**): Shown are the Confocal images of the posterior terminii of
nuclear cycle 13/14 embryos of the following genotypes: *cs* (A1-A3), *nos gal4* (B1-B3), *casp^{lof}/Df* (C1-C3),
casp^{lof}/Df; *nos gal4>casp^{wt}* (D1-D3) and *nos gal4> casp^{wt}* (E1-E3). Embryos were immunostained with Vasa (1:50)
antibody. Hoechst marks the nuclei. (**F**) The total number of germ cells marked with Vasa were quantified and
plotted as bar graphs. N (embryos)= 30. Ordinary one-way ANOVA, (****) P <0.0001. (**G**) Casp protein levels
were assessed in 0-3 embryos via western blotting. Rabbit anti-Casp (1:10,000) and rabbit anti-HA (1:2000)
antibodies were used to probe the blot. Ponceau is used as a loading control. N = 3.
450

Casp interacts with TER94 in the early embryo.

Data presented in the previous section demonstrated that both the 'loss' and 'gain' of *casp*

activity exert reciprocal influence on total PGC numbers in early embryos. As very few germ

plasm components have been shown to display this trait (JONGENS *et al.* 1994), we decided to
explore the phenomenon further. We have previously reported that Casp protein physically
associates with both transitional endoplasmic reticulum ATPase (TER94; also called valosin-
containing protein, VCP or p97) and vesicle-associated membrane protein-associated protein B
(VAPB/VAP33A)(TENDULKAR *et al.* 2022). Furthermore, based on interaction studies and
biochemical analysis, we proposed that Casp may act as an adapter that either directly or
indirectly mediates the physical association between VAPB and TER94 (TENDULKAR *et al.*
2022). To arrive at this conclusion, we had primarily relied on S2 cell lysates as well as adult
and fly head total protein extracts. To specifically identify the major protein partners of Casp in
the early embryos, we performed similar immunoprecipitations using anti-Casp antiserum in
early (0-3 hour) embryonic lysates and interactors were identified via mass spectrometry (Fig.
S3A). Peptides from 122 proteins were recovered in the Casp IP but not the IgG IP and thus,
were considered significant interactors. The top 23 interactors are presented in Fig. S3A.
Interestingly, as in the case of brain and S2 cell lysates, TER94 and VAPB were the top hits,
with germ cell determinants also identified as interactors (Fig. S3C; Discussed in a subsequent
section). These observations suggested that a functional protein complex between Casp,
TER94 and VAPB likely exists in many different tissue/cellular contexts, including early
embryos.
472
Maternal requirement of TER94.
Mammalian VCP/p97 is an essential chaperone for proteostasis that modulates several
ubiquitin-associated processes (PETERS *et al.* 1990; DAI *et al.* 1998; MEYER *et al.* 2000; MEYER
2005; YE 2006; JENTSCH AND RUMPF 2007; MEYER *et al.* 2012). Its *Drosophila* counterpart
TER94 has been studied mostly in the context of neurodegeneration (GRICIUC *et al.* 2010;
CHANG *et al.* 2011; AZUMA *et al.* 2014; KUSHIMURA *et al.* 2018; TENDULKAR *et al.* 2022;
THULASIDHARAN *et al.* 2024). Loss of TER94 was shown to ameliorate polyQ-induced eye
degeneration. Moreover, the overexpression of TER94 promoted the apoptosis of neuronal cells
(HIGASHIYAMA *et al.* 2002). TER94 has been uncovered in a genetic screen for maternal
proteins that are phosphor-regulated (ZHANG *et al.* 2018) and have roles in oogenesis and early
development.
TER94 is one of the significant interactors of Casp protein in the embryonic context as
suggested by the Casp interactome (Fig. S3A) and has been implicated in ER-associated
degradation (CHANG *et al.* 2011; TENDULKAR *et al.* 2022). Thus, we wondered if TER94 also has
a maternal function and whether embryos maternally compromised for *TER94* display similar

phenotypes as *casp*. To achieve this, we employed VALIUM 20/22 maternal RNAi lines (*UAS-TER94i*). First, we used a *mat-α4tubulin:VP16-Gal4* (*Mat-αtubGal4*) driver to deplete *TER94* activity in the late stages of oogenesis (Fig. 4).

Fig. 4. Maternal requirement of *TER94* during early embryogenesis. (A) Viability of embryos derived from *w^1118* and *Mat-αtubGAL4 > TER94i* (referred to as *TER94 RNAi* henceforth) mothers is represented as a bar graph. N = 3, ordinary one-way ANOVA, (***) *P* <0.001. (**B**) Western blot analysis indicates the knockdown efficiency of TER94 in the 0-3 h embryo, estimated to be ~90%. , levels remain unaffected in *casp^lof/Df* embryos. Tubulin was used as a loading control. N = 3. (**C, D**) Confocal images of sections of nuclear cycle 13/14 embryos of the indicated genotypes stained with Hoechst, phalloidin, and gamma-tubulin. Nuclei and F-actin are visualized with Hoechst and phalloidin respectively. Nuclear disruption (D, asterisks) and F-actin aggregates (D2, arrows) are observed in the

540 mutant. Defective centrosomes (D3, arrows, inset) can be observed in the *TER94 RNAi* embryos. (**E, F**) The
541 percentage of defective centrosomes in nuclear cycles 9/10 and 12/13 are represented as bar charts. n = 10.
542

We sought to observe the consequence of *TER94* knockdown both in the adult females
and early embryos derived from such females. In the females compromised for *TER94*, egg-
laying behavior remained largely unaffected. Interestingly, however, the viability of such
embryos was severely impaired, with >95% of embryos failing to hatch (Fig 4A), in agreement
with earlier studies (ZHANG *et al.* 2018).
A western blot of lysates derived from *Mat-GAL4>UAS-TER94i* mothers probed with
anti-TER94 antiserum indicated a robust knockdown of TER94 as compared to control (Fig. 4B).
Phenotypic analysis of embryos derived from *Mat-GAL4> TER94i* females revealed that >70%
of *Mat-GAL4/ TER94i* embryos failed to progress to the syncytial blastoderm stage, with even
fewer reaching Stage 5. The late-stage syncytial/ cellular blastoderm embryos were further
assessed for cell cycle defects. As in the case of *casp^{lof}/Df*, *Mat-GAL4>TER94i* embryos also
displayed defects ranging from irregular nuclear distribution to perturbed F-actin localization, as
assessed by HOECHST and phalloidin staining, respectively (Compare Fig. 4D1-D3 with Fig.
4C1-C3).
We also examined if maternal loss of *TER94* activity results in centrosome aberrations
like those observed in *casp^{lof}* embryos. Indeed, embryos deficient in *TER94* function showed
both the characteristic phenotypes observed in the *casp* embryos including a) inefficient
separation of duplicated centrosomes and b) multiple instances of 'orphan' centrosomes devoid
of nuclear DNA (Fig. 4E, F). An earlier study (ZHANG *et al.* 2018) had found that TER94 RNAi
leads to multi-polar spindles with supernumerary centrosomes.
Taken together, these data demonstrated that embryos maternally compromised for
either Casp or TER94 functions share several phenotypes, suggesting that both the proteins
likely perform essential and likely related functions during syncytial nuclear cycles in *Drosophila*
embryos. Furthermore, the phenotypic consequences observed upon maternally compromising
casp and *TER94* overlap but are not identical, with TER94 embryos showing higher penetrance
in terms of phenotypes and embryonic lethality. Future experiments will be necessary to resolve
the functional distinction between the two.
570

TER94, a known component of pole plasm, is detectible in PGCs.
Maternal loss of TER94 function mimicked both the cytoskeletal and corresponding
centrosome aberrations observed upon similar loss of *casp* activity. Both centrosomes and
germ plasm, are essential for the proper specification and formation of PGCs in *Drosophila*

embryos (LERIT AND GAVIS 2011; LERIT *et al.* 2017). This prompted us to examine the Casp

interactome for the possible enrichment of germ plasm components and proteins that may

regulate centrosome dynamics and/or behavior. Fig. S3B lists Casp interactors that are

important constituents of the germplasm, including eIF4A, me31B, and TER94, known

constituents of the polar granules (THOMSON *et al.* 2008).

Among the different components of the germ plasm, Oskar serves as the principal

determinant of the PGC fate (KIM-HA *et al.* 1991; SNEE AND MACDONALD 2004; VANZO *et al.*

2007; LEHMANN 2016). Supporting the conclusion, the loss and gain of function of *oskar* leads to

reciprocal phenotypes. Compromising *oskar* activity maternally, leads to reduction in total

number of PGCs whereas anterior ectopic expression of *oskar* using the *bicoid* mRNA

localization signal induces pole cell formation at the anterior (EPHRUSSI AND LEHMANN 1992).

The germ plasm is supplemented with mitochondria and polar granules, which consist of

ribonucleoprotein complexes. Downstream of *oskar*, *vasa* and *tudor* are two genes that are

essential for the assembly of pole plasm (BREITWIESER *et al.* 1996; ARKOV *et al.* 2006; JONES

AND MACDONALD 2007). Biochemical proteomic analysis of Vasa (VAS) and Tudor (TUD)

containing polar granule complexes identified eIF4A, me31B, and TER94. This indicated that

the germ plasm consists of the components of translational machinery and the endoplasmic

reticulum assembly (THOMSON *et al.* 2008).

TER94 is a component of pole plasm and is also physically associated with Casp.

Furthermore, proteomic analysis indicated that Casp could be part of a complex comprising

several germ plasm proteins including Tudor and Me31B. To confirm that, as in the case of

Casp, TER94 is also detected in PGCs. We used antibodies generated against Casp (Fig. S3D)

and TER94 (Fig. S3E) to label wild-type embryos. Embryos were also co-immunostained for

germ cell specific marker, Vasa. As can be seen, both Casp and TER94 proteins are found in

PGCs, colocalized with Vasa.

Casp and TER94 regulate embryonic germ cell formation.

As we were specifically interested in investigating possible similarities and distinctions

between the respective functions of TER94 and Casp during early embryonic germ cell

development, we decided to examine their possible functions during PGC formation (also

referred to as pole cell budding). This seemed especially pertinent as centrosome behavior and

dynamics are crucial for proper formation and cellularization of PGCs. Evidently, a germ plasm

component, Germ-cell-less (CINALLI AND LEHMANN 2013) was shown to be necessary for proper

separation of daughter centrosomes in the dividing pole buds (LERIT *et al.* 2017). Similarly, loss

of centrosome components such as Centrosomin results in partial loss of early embryonic

PGCs. As maternal loss of both *casp* and *TER94* results in aberrant centrosomes in the

surrounding somatic nuclei in early blastoderm embryos, we sought to investigate if posteriorly

positioned centrosomes in the vicinity of pole plasm also display similar problems. Indeed,

maternally compromising *casp* led to significant loss of pole buds and corresponding defective

centrosome behavior (Fig.5).

615
616
617

Fig. 5. *casp and TER94* **regulate pole bud formation. (A-C)** Confocal sections of the posterior termini of nuclear cycle 9/10 embryos derived from *w^1118^* (A1-A3), *casp^lof^/Df* (B1-B3), and *TER94 RNAi* (C1-C3) mothers immunostained with gamma-tubulin. γ-tubulin labels centrosomes of pole buds, identified based on Vasa staining. Nuclei are stained with Hoechst. The inset(s) for B3 and C3 highlight the centrosomal defects (arrow) when compared with A3. Defects were quantified and plotted as bar graphs in (**D**) and (**E**) respectively. n = 20.

Especially these pole buds display defective centrosome separation and/or orphan centrosomes (Fig. 5B1-B2,D,E), as observed in the somatic nuclei. Intriguingly, while *casp^lof^* embryos show discernible loss of pole buds, *TER94i* embryos displayed an increased number of round-shaped, 'bud-like' cells at the posterior of the stage 3 embryos (Fig 5 C1-C2). Typically, such ectopically localized pole buds also showed inadequate separation of centrosomes and could be identified as pole buds due to the enrichment of Vasa. However, Vasa distribution and accumulation in *TER94i* buds was variable and non-uniform. Since TER94 is required for *Osk* mRNA localization in the oocyte, the defects in pole cell budding including their ectopic positioning could be a consequence of inappropriate segregation of *oskar.* Moreover, the early nuclear division cycles and nuclear migration defects seen in are defective in *TER94i* embryos could also contribute to PGC formation/ cellularization. Notably, although the pole bud count is elevated in *TER94i* embryos, presumably, these pole buds don't survive through subsequent nuclear cycles, and consequently, late blastoderm *TER94i* embryos almost completely lack PGCs.

Casp activity is needed for the accumulation of Oskar protein at the embryonic posterior pole.

Taken together, our data demonstrate that change in Casp levels leads to corresponding alteration in the total PGC count. Moreover, PGC formation is affected by maternal loss of Casp, although the precise nature of its involvement in this process remains to be determined. As Oskar is the master determinant of germ cell fate, we sought to determine if Oskar levels (KIM-HA *et al.* 1991; EPHRUSSI AND LEHMANN 1992) are correspondingly altered upon change in Casp. To assess this possibility, we stained wild-type control embryos and embryos maternally compromised for *casp* using anti-Oskar antibodies (Compare Fig 6B to 6A). Simultaneously, we stained embryos derived from *nosGal4>UASp-casp* mothers (Fig 6C1-C3) that display significantly elevated number of PGCs, presumably due to excess levels of Casp protein in the germline.

Fig. 6. Casp regulates Oskar levels and modulates cell division
(A-C) Confocal microscopy images of stage 1/2 embryo laid by w^{1118}, $casp^{lof}/Df$, and *nos gal4> casp^{wt}* females and immunostained with Oskar antibodies. Replicate (1-3) are shown to highlight variable reduction in Oskar levels. In addition, as compared to controls (w^{1118}) embryos (A1-A3), spread of Oskar appears to be restricted in $casp^{lof}/Df$ (B1-B3) whereas it is expanded in *nos gal4> casp^{wt}* (C1-C3).
(D) Volume occupied by Oskar, per embryo is measured and plotted for all three genotypes (w^{1118}, $casp^{lof}/Df$, and *nos gal4> casp^{wt}*), n = 8. Ordinary one-way ANOVA, (**) $P <0.01$, (*) $P <0.05$.
(E) Number of pole buds (nc 9/10) and pole cells (nc 11/12, nc 12/13) in 0-3h old *nos gal4, casp^{lof}/Df and nos gal4> casp^{wt}* embryos at different nuclear cycles. Two-way ANOVA, (*) $P <0.05$, (****) $P <0.0001$. n = 10
(F) Graph showing the distribution of actively dividing pole cells as indicated by the presence of phospho-histone 3 (Ser10) antibody. Quantitative analysis was performed using nuclear cycle 12-14 in 0-3h old, *nos gal4, casp^{lof}/Df, nos gal4> casp^{WT} and casp^{lof}/Df;nos gal4> casp^{WT}* embryos.

Satisfyingly, the change in Oskar protein levels is consistent with the alteration in the Casp

levels in the maternal germline (Fig 6D). Embryos derived from *casp* mutant mothers show

considerably diminished levels of Oskar as compared to the control whereas overexpression of

Casp in the maternal germline results in elevation in the Oskar protein amount (Fig 6D).

669   Furthermore, in all instances, Oskar protein appears to be anchored to the posterior pole as the

670   wild-type control embryos and the only discernible change is observed in its levels.

672   ***Casp levels influence total number of phosphor-histone3 (pH3) positive pole buds and***

673   ***PGCs.***

674   Changes in Casp protein levels influence the accumulation of Oskar protein. Oskar is

necessary and sufficient to assemble pole plasm at the posterior pole which controls the total

number of pole buds in an embryo. To directly evaluate the influence of Casp on pole bud

formation, total number of pole buds and pole cells were quantitated upon maternal loss and

gain of Casp against the control embryos at the same stage (Fig. 6E). Consistent with the data

presented in the previous sections, total number of pole buds (nc 9 and 10) were reduced in

casp^{lof} embryos whereas the number was elevated in the presence of excess Casp (Fig 6E).

Similar quantitation was performed in older embryos in two stages (nc11-12 and NC12-13), and

a progressive increase was observed in the total number of PGCs upon the gain of Casp

function (Fig. 6E). By contrast, the total PGC count for *casp^{lof}/Df* embryos did not increase

appreciably at nc11-12 and nc12-13.

685   Primordial germ cell number is determined by limited mitotic cell divisions that each pole

bud undergoes in a stochastic and asynchronous manner. As these are mitotic divisions, the

dividing PGCs can be identified using antibodies against the miotic marker phosphoHistone3

(pH3). As the total number of pole buds and PGCs change per Casp levels, we wanted to

examine if this is also reflected in the total number of pH3-positive germ cells simultaneously

labeled with the anti-Vasa antibodies. We decided to focus on syncytial blastoderm embryos

between NC12-14. As anticipated, maternal overexpression of Casp led to a corresponding

increase. In control (*nos gal4*) embryos, roughly 20% PGCs display either 1 or >1 pH3 positive

PGCs whereas, 70% of *nos-gal4>Casp^{WT}* embryos show 1 or >1 pH3 positive PGCs (Fig. 6F).

Conversely, less than 10% of *casp^{lof}* embryos showed either 1 or >1 pH3 positive PGCs and

pole cells with 3 or >3 pH3 positive cells were completely absent in this background as opposed

to 25% present in the control embryos (Fig. 6F). Importantly, maternal expression of the rescue

construct in the *casp^{lof}* embryos ameliorated the loss of pH3 positive PGCs seen in just the

mutant (*casp^{lof}*) PGCs.

Does Casp function affect canonical MBT regulators?

Our analysis thus far has revealed two related yet distinct phenotypes associated with

maternal loss of *casp*. First, it can influence proper PGC formation and specification during

early embryonic development via its effect on the accumulation of germ cell determinant, Oskar.

Subsequently, it helps orchestrate cellular movements leading up to gastrulation during mid-

embryogenesis. While the first activity is likely germ cell autonomous, the second relates to its

role in the somatic cells/nuclei. Moreover, both the somatic and germline compartments of

casp^{lof} embryos share centrosomal as well as cytoskeletal aberrations. We thus wondered

whether the two activities are mechanistically connected, and if the possible connection relates

to the maternal-to-zygotic transition (MZT). One aspect of the MZT that has been highlighted in

recent years is the active degradation of maternal proteins (CAO *et al.* 2020). These proteins

comprise 2% of the maternal proteome and are degraded abruptly, at the end of the MZT. In

Drosophila, ubiquitin-proteasome-based degradation of three repressor proteins, namely Smaug

(Smaug), Trailer hitch (Tral), Maternal expression at 31B (Me31B), marks the MZT (CAO *et al.*

2020). We thus decided to probe, using antibodies generated against these proteins (Fig. 7) if

the degradation of any of these proteins is affected due to maternal loss of Casp/*casp* during

the initial hours of embryonic development (0-1, 1-2 & 2-3 hours). TER94 protein is unchanged

(Fig. D1, D2) as is the α-tubulin loading control (Fig. 7E1, E2). Furthermore, pattern of

degradation of Me31B and TRAL proteins in the *casp^{lof}* embryos is relatively unaffected, when

compared to the *w^{1118}* control samples.

Fig. 7. Reduction in Casp activity specifically affects Smaug degradation during the MZT. At the MZT, a few maternal proteins are actively degraded (CAO *et al.* 2020). Embryos from mothers with genotype *w[1118]* **(A1-E1)** and *casp[lof]/Df* **(A2-E2)** were collected using three time intervals (0-1, 1-2, 2-3 Hrs), and embryonic lysates were separated on SDS-PAGE gels. The Western blots were probed with antibodies against ME31B (A1-A2), Tral (B1-B2) and Smaug (C1-C2) to assess the extent of protein degradation, as part of the MZT. α-tubulin (E1-E2) was used as a loading control, while TER94 (D1-D2) was used as a negative control as it is a maternal protein that does not undergo degradation at MZT. **(G-J)** Quantitation of the intensity of change of protein band (normalized to 1), for Tral (G), Smaug (H), ME31B (I) and tubulin (J). Each data point is an average of over 5 western blots. The intensities at each time point for *nos gal4* vs *casp[lof]/Df* are statistically not significant.

By contrast, Smaug (Fig, C1-C2) is an interesting exception. In control, 0–1hr old embryos, low

levels of Smaug protein were observed, which increased in the 1-2hr old embryo extract but

decreased the 2-3 hr time-window (Fig. 7C), data that agrees with previously published work

(CAO *et al.* 2020). However, in the *casp^{lof}/Df* embryos levels of Smaug protein are modestly

elevated from the start (0-1 and 1-2 hr old extract samples respectively; Fig. 7C2). Critically,

unlike control embryos, Smaug protein persists in 2-3hr old *casp^{lof}/Df* embryonic extracts (Fig

7C2, *). Altogether, these data argue that Smaug degradation is specifically adversely

influenced in the *casp^{lof}/Df* embryos at the end of the MZT.

Germ cell specifc Smaug levels are influenced by casp activity.

Since Smaug levels appear to be elevated in embryonic lysates in the MZT, we decided

to test if the Smaug levels are elevated specifically in *casp^{lof}/Df* pole cells. To this end, pre-

syncytial as well as early syncytial blastoderm embryos of both the genotypes (control and

casp^{lof}/Df) were stained with anti-Smaug and anti-Vasa antibodies. Aligned with the Western

blot data presented earlier, levels of Smaug protein are significantly elevated in pre-syncytial

embryos (Fig. 8, compare panel B3 with A3). Also, in *casp^{lof}/Df* embryos, Smaug protein seems

to accumulate in discernible large puncta which are barely visible at this stage in control

embryos (Fig. 8, B3 vs A3). At the syncytial blastoderm stage (Stage 4), however, the Smaug

positive puncta appear larger and more numerous in the few surviving PGCs from the *casp^{lof}/Df*

embryos (Fig. 8, D3) when compared to controls (C3).

Taken together, our data suggest that Smaug protein levels are appreciably elevated in

pole cells from *casp^{lof}/Df* embryos. The specific increase in Smaug levels may, in part, be due to

inappropriate accumulation and/or stabilization of Smaug. Smaug protein is necessary for

translational control of the posterior determinant *nanos*. Early reports indicated that unlocalized

nos RNA is translationally repressed by Smaug which binds to Smaug Response Elements

(SREs) within 3'UTR of *nos* RNA. In addition to *nos*, it also regulates *hsp83* translation. Smaug

participates in multiple, overlapping mechanisms including interaction with the components of

the translation machinery as well as deadenylation to regulate translation/localization of the

target RNAs (DAHANUKAR *et al.* 1999; ZAESSINGER *et al.* 2006).

Fig. 8. Decrease in Casp correlates with an increase in Smaug levels in the pole buds and PGCs. Confocal microscopy images of the posterior termini of stage 1-2, and 4 embryos derived from *nos gal4* and *casp^lof^/Df* females. Embryos were immunostained with Vasa (green) and Smaug (red) antibodies. Hoechst marks the nuclei (A-D). Smaug expression was quantified across the two genotypes and plotted as bar graphs (E-G). N=5 embryos (Stage 2) and 10 embryos (Stage 4).

Intriguingly, recent data from Lipshitz lab has, in fact, shown that Smaug protein accumulates in germ granules (SIDDIQUI *et al.* 2023). Moreover, in addition to its canonical role in regulating *nos* translation, it can also repress *oskar* at the translational level (see Discussion).

886   As the embryonic PGC count depends upon Oskar levels, *Smaug* mutant embryos show an

887   increase in the total number of PGCs. These data fit nicely with our observations and are

888   entirely consistent with a model incorporating the downregulation of Smaug mediated by Casp.

890   ***Functional analysis of different protein domains within Casp.***

891   To decipher the specific functions of the individual protein domains of Casp during pole

cell formation and division we followed an experimental strategy involving rescue of the loss of

function phenotype. We compared the extent of rescue observed in the presence of individual

deletion constructs to the full length wild-type control. To this end, fly lines expressing domain

deletions of Casp under the *pUASp* promoter were generated (Fig. 9A). The *pUASp-casp$^{\Delta UBA}$*,

pUASp-casp$^{\Delta UBX}$ and *pUASp-casp$^{\Delta UAS}$* construct represent deletions of the UBA, UBX, and UAS

domains, respectively. Additionally, we also generated fly lines that express the *pUASp-*

casp$^{\Delta UAS\Delta UBX}$ which deletes both the N and C-terminal domains simultaneously. A full-length

coding sequence of Casp was also cloned, and this *pUASp-casp^{WT}* line was used as a positive

control carrying the 'rescue' construct. A western blot confirmed the expression of these

constructs in adult animals with different domain deletions (Δ), *UAS-casp$^{\Delta}$* driven by *nos-GAL4*

driver (Fig. 9B). To assess the domain-specific functions of Casp in isolation without the

confounding effects of endogenous Casp, the fly lines were balanced with a *casp^{lof}* allele on the

second chromosome. These balanced lines were crossed to a fly line with the maternal driver

nos-GAL4 on the third chromosome and an allele with a deficiency for *casp (Df)*. Embryos laid

by mothers of the genotype *casp^{lof}/Df; nos-GAL4/pUASp-casp$^{\Delta}$* were used to determine

maternal effects of the domain deletions (Fig. 9C).

Fig. 9. Structure-function analysis of Casp protein domains in regulating pole cell number. (A) Schematic representation of different domain deletion variants of Casp. As shown in the schematic, the WT Casp protein consists of different functional domains including UBA, UAS, UBX and the FFAT-like motif. **(B)** Western blot of the truncated Casp proteins, deficient in the domains indicated, probed with rabbit anti-casp (1:10,000) and rabbit anti-HA (1:2000) antibody. *nos gal4* served as a control. Ponceau staining was used as a loading control. Asterisk (*) denotes non-specific antibody binding. **(C)** The total number of germ cells from nc 13/14 embryos derived from mothers expressing different *casp* domain deletion constructs in the *casp*^{lof}/*Df* background were marked with Vasa antibodies, quantified and plotted as a bar graph. n = 30, Ordinary one-way ANOVA, (****) *P* <0.0001, (*) *P<0.05*. **(D)** The total number of germ cells from nc 13/14, derived from mothers expressing casp domain deletion, in an otherwise *wild-type* background, were stained with anti-Vasa antibodies. Vasa positive cells (i.e. PGCs) were quantified and plotted as a bar graph. n = 30, Ordinary one-way ANOVA, (****) *P* <0.0001.

Embryos were collected from females of different genotypes and stained using anti-Vasa

antibodies. As can be seen by the comparison shown in the bar graph (Fig. 9C), in terms of the

rescue of PGC numbers, deletion of the UBA domain does not affect the rescuing activity of the

Casp protein, the UBX domain deletion can rescue, but is weaker than the UBA domain

deletion, whereas deletion of the UAS domain is the least effective with regards to the rescue.

To extend these observations, we overexpressed individual Casp deletion mutants in an

otherwise wild-type background and counted the total number of PGCs. Neither the maternal

overexpression of *pUASp-casp^{wt}* nor any of the *pUASp-casp^{Δ}* increased lethality in terms of

decrease in hatching of embryos, in all cases the hatching was in the range 90-95%. Again, as

seen before, overexpression of *casp^{wt}* under these conditions led to a significant increase in the

total number of PGCs in blastoderm embryos, with average PGC's at 40. The ability to enhance

the PGC division is retained in the absence of the UBA domain as, in this particular background,

the mean count of the total number of germ cells was similar to embryos that expressed full-

length protein. By contrast, deletion of the UAS domain is unable to support the (increased)

germ cell division, and the total PGC count dropped to the same level as *nos Gal4* embryos.

Interestingly, deletion of UBX domain, yet again, displayed intermediate activity, with the *ΔUBX-*

ΔUBA domain deletion also being unable to drive the increase in PGCs.

Proteomic analysis of the embryonic extracts suggested that in addition to

TER94 protein, Casp is also associated with VAP (Fig. S4A; (TENDULKAR *et al.* 2022)). We thus

sought to test if physical interaction between Casp and VAP is relevant for pole cell formation.

To assess this possibility, we used a deletion variant of Casp, Casp^{ΔFFAT}, which cannot interact

with VAPB (TENDULKAR *et al.* 2022). Expression of *casp^{ΔFFAT}* under the control of maternal

driver, in a *casp^{lof}*/Df mother showed hatching (~60%) frequencies at par with wild type (Fig.

S4B), with expression levels equal to other *casp* constructs (Fig. S4C). Importantly, *nos-Gal4-*

dependent maternal overexpression *of UAS-casp^{ΔFFAT}* led to an increase in the total number of

PGCs in syncytial blastoderm embryos, comparable to full-length *UAS-casp^{wt}* (average PGC

~23). Lastly, expression of *casp^{ΔFFAT}* could rescue the PGC loss upon maternal loss of *casp*

(compare Fig. S4 F1-3 to Fig. 9C). These data suggested that interaction between Casp and

VAP is either not essential or partially redundant for Casp function in the context of determining

PGC. Altogether, the functional analysis of different protein domains within Casp underscores

the importance of UAS and UBX domains, especially in embryonic contexts, including PGC

development.

 Discussion

Proper development of biological systems in an organismal context involves complex
interactions between various individual pathways. The qualitative nature of the interaction(s)
between pathway components ultimately determines how different pathways intersect in a
context-specific manner. Curiously however, there are only a defined number of signaling
pathways/circuits that have been elucidated thus far. Consistently, the entire molecular
cassettes that constitute a given pathway or a select number of pathway components, are
reiteratively used in a variety of biological contexts even within a lifecycle of the same organism.
Moreover, while such repurposing is relatively frequent, the corresponding biological outcomes
are diverse in nature and insightful in ways more than one. Our data detailing the functional
involvement of Casp and TER94 in the PGCs is a case in point. Casp function was initially
characterized in the context of *Drosophila* immune response (KIM *et al.* 2006). However, the
original study did not investigate the developmental roles of Casp and its interactors. Here, we
report novel activities of *Drosophila melanogaster* immune components Casp and TER94 during
early embryonic development with a specific focus on germline.

Early embryonic patterning is a dynamic process in metazoans which is primarily
regulated by the deposition of maternal gene products including RNAs and proteins. Such
maternal determinants are localized in a spatially restricted manner (SCHIER 2007). On many
instances, the unique pattern of localization of specific factors underlies their activities during
embryo patterning. To execute their functions properly, patterning determinants heavily rely
upon the members of the housekeeping machinery that carefully calibrate the synthesis,
stability, transport, and degradation of diverse regulatory components. Here, we have explored
novel embryonic functions for the FAF1 ortholog, Casp in *Drosophila*. Initially, we uncovered
that maternal loss of *casp* activity results in partially penetrant embryonic lethality. The *casp*
allele was determined to be a strong loss of function, but not a null. The partial penetrance could
also be due to redundancy. Consistent with either possibility, quantitation of the individual
phenotypes yielded significant but variable penetrance.

Our data demonstrate Casp's involvement during cellular movements that lead to
gastrulation, a likely cause underlying the lethality. Surprisingly, Casp protein is enriched in the
PGCs, which are specified at the posterior pole under the control of the master germ cell
determinant, Oskar. PGC formation and specification in a young *Drosophila* embryo depends on
the posteriorly anchored specialized cytoplasm (or pole plasm) enriched in RNA and protein
components essential to determine germ cell identity and behavior. Two important traits
distinguishing early pole cells from the surrounding somatic nuclei include precocious budding

and limited mitotic self-renewal. Consistent with its functional involvement in both these

processes, maternal loss of either Casp or its protein partner TER94 resulted in a reduced

number of buds and subsequent loss of PGCs. While qualitatively similar, the severity of

phenotypic consequence due to loss of Casp and TER94 regarding PGC loss, is not identical.

Interestingly, however, both proteins appear to influence centrosome behavior and dynamics

that are of paramount significance in forming pole cells in *Drosophila* embryos.

Thus far, Germ cell-less (Gcl) is the only protein shown to control PGC formation in a

similar manner (JONGENS *et al.* 1992; CINALLI AND LEHMANN 2013). Importantly Gcl activity

depends on its ability to influence proper separation of centrosomes and elaboration of astral

microtubules in dividing pole buds (LERIT *et al.* 2017). Aberrant centrosome behavior in *gcl*

mutant embryos adversely affects PGC budding and equitable distribution of germ plasm

among daughter cells. Intriguingly, these phenotypic traits can be recapitulated by simply

engineering defective centrosome separation (LERIT *et al.* 2017). It is thus noteworthy that

maternal loss of Casp and TER94 leads to similar defects and future experiments will reveal

mechanistic details underlying roles of these two proteins in regulating centrosome dynamics in

PGCs and their possible interaction with Gcl and its protein partners. Of note, a germplasm

interactome (THOMSON *et al.* 2008) included TER94 along with other important germ plasm

proteins including Vasa, Tudor, and others.

TER94, an ER protein, is a major Casp interactor. Our data also indicate that in addition

to PGC formation, TER94 and Casp regulate early nuclear division cycles in syncytial

blastoderm embryos and subsequent cell divisions in the somatic compartment cell cycle

processes in the gastrulating embryos. (Casp and TER94 are distinct compared to Gcl in this

regard which has a strictly germ cell specific function). Taken together, these observations

suggest that Casp and TER94 contribute to critical early developmental functions leading to

mid-blastula transition which precedes gastrulation and germband extension. In the mammalian

context, the FAF1-VCP interaction is mediated by the UBX domain. Thus, it is unsurprising that

deletion of the protein domain crucial for association between Casp and TER94 resulted in a

somewhat diminished function as compared to a full-length version. Curiously however, deleting

UAS fragment, a protein domain of unknown function, implicated in self-association, resulted in

substantially compromised activity as compared to native protein. Future experiments will focus

on the specific molecular interactions this (and other) individual domain(s) are involved in. It will

be also important to determine how protein degradation especially engineered via ubiquitin

modification contributes to Casp stability and function.

Intriguingly our data also argue that regulation mediated by *casp* is likely critical for

maintenance of PGC fate via maintenance of Oskar levels. Importantly, several independent

observations suggest that this influence is likely zygotic. First, barely detectable amount of

Casp protein is deposited in the egg and bulk of the Casp protein is generated by translation

post-fertilization. Second, Casp protein controls overall levels of Smaug at the post-

transcriptional level including in the early embryonic PGCs. It was recently reported that Smaug

protein accumulates in the germ granules where it controls Oskar (and Bruno) translation

negatively by binding to Smaug response elements present in the untranslated regions within

the respective RNAs (SIDDIQUI *et al.* 2023). Consistently loss of function mutations in *Smaug*

result in a modestly elevated PGC count in stage 4-5 embryos whereas a reciprocal phenotype

is observed in *casp^{lof}* embryos. Taken together with the increase in Smaug levels in *casp^{lof}*

embryos, it would be reasonable to propose that PGC-specific phenotypes observed upon loss

of *casp* are, in part, mediated by excess accumulation of Smaug. Moreover, these authors also

suggest that Smaug dependent regulation of *oskar* RNA likely has a significant zygotic i.e.

embryonic component which aligns well with our data. Future experiments will be necessary to

elucidate the mechanism underlying regulation of Smaug levels by Casp. It will also be

interesting to examine if Smaug and Casp regulate centrosome behavior reciprocally to control

the final PGC count.

Degradation of maternally deposited RNAs and proteins constitutes an important

transition during embryonic development. Zygotic Genome Activation (ZGA) and turning over of

maternal determinants (Both RNAs and proteins) happen almost simultaneously. Together

these two events constitute MZT which is delayed in the germ cell compartment as opposed to

soma. Nonetheless, recent data have suggested that the two events possibly occur in a

coordinated manner possibly via shared components (COLONNETTA *et al.* 2023). Many of the

embryonic phenotypes that *casp^{lof}* embryos display (aberrant nuclear migration, cellularization

defects, defective gastrulation etc.), are shared by mutations in gene products either directly or

indirectly involved in one of these processes. It will be important to elucidate new interactors of

Casp as its function likely impacts protein degradation and/or stability of the target proteins.

Recent reports have indicated potential involvement of ZGA regulators such as Zelda and

CLAMP in germline/soma distinction. Critically this function of the components of ZGA depends

on proper anchoring, release and transmission of posteriorly anchored pole plasm that involves

centrosome function. Maternally compromising ZGA components resulted in inappropriate

release and transmission of pole plasm RNAs, a phenotype that is partially recapitulated in

casp^{lof} and *TER94i* embryos.

While our data have clearly established the involvement of Casp and TER94 during PGC

formation and specification, further studies will be necessary to elucidate their precise molecular

function(s) underlying this activity. Several observations are noteworthy in this regard and will

guide the course of future investigation. The first set of results points to the possible

participation of both Casp and TER94 during ubiquitin-dependent protein degradation.

A regulatory network incorporating Bru1, Cup, Oskar and Smaug are key to PGC

specification. The Skp Cullin and F-box (SCF) containing complex marks proteins such as

Smaug for degradation by functioning as an E3 ubiquitin ligase ((CAO *et al.* 2020; CAO *et al.*

2022); Fig. 10A). The ubiquitinated Smaug is then degraded by the proteasome. In the

mammalian context, the Casp/TER94 orthologs FAF1/VCP (in *green* font; Fig. 10A) have been

found to interact with the SCF complex. Our data supports a similar scenario in *Drosophila*.

1136

Figure 10. A model for Casp function in PGCs.

(**A**) <u>Smaug is marked for degradation by the SCF complex</u>. Specific pole plasm components including Cup, Oskar

and Smaug are actively degraded. *oskar* is regulated by Bru1 and Cup (*ref 1-3*; (NAKAMURA *et al.* 2004; KIM *et al.*

2015; BAYER *et al.* 2023)). Oskar (*ref 4-6*; (MAHOWALD 2001; HUYNH AND ST JOHNSTON 2004; LEHMANN 2016)) is

the master determinant of PGC fate, and hence the stability of the Oskar:Smaug complex (*ref 7*;(KUBIKOVA et al.

2023)) is a key to PGC determination and proper specification. Oskar also regulates posterior cell fate by regulating

nos translation which is modulated by Smaug (*ref 8-10;* (DAHANUKAR et al. 1999; ZAESSINGER et al. 2006; JESKE et

al. 2011)). The SCF complex is a multi-protein E3 ubiquitin ligase complex, and Smaug is one of its prominent

targets (*ref 11-12;* (CAO et al. 2020; CAO et al. 2022)). Smaug appears to repress the translation of *oskar* and *Bruno*

1 mRNA (*ref 13*; (SIDDIQUI *et al.* 2023)). In mammals (marked with green font), FAF1 modulates the SCF complex

(*ref 14*; (MORAIS-DE-SA *et al.* 2013)) with VCP/p97 assisting in the degradation of ubiquitinated Smaug (*ref 15-16;*
(LI et al. 2014; REIM et al. 2014)), suggesting potential conservation of the activities in the fly orthologs, Casp and
TER94.
(**B**) Casp/TER94 interact with an overlapping set of proteins. TER94 has earlier been demonstrated to be associated
with Oskar (RUDEN *et al.* 2000). Our data points to an association between *Drosophila* Casp and TER94 (this study
and *ref 17;* (TENDULKAR et al. 2022)) that participates in protein degradation. In a proteomic analysis of germ cell
components, TER94 was identified (this study and *ref 18*; (THOMSON *et al.* 2008)), along with other bona-fide germ
cell constituents including Cup, Tral, Bel, eIF4A, Tud and Vasa (*ref 18* and *19* (THOMSON *et al.* 2008; DEHAAN *et*
al. 2017)). Both, Casp and TER94 are thus enriched in the pole cells and interact with proteins that specify PGC
fate.
(**C**). The SCF Complex, Gcl and Centrosome integrity. The SCF complex is localized to the centrosome. Gcl
interacts with Cullin 3 to degrade Torso receptor to promote PGC fate (*ref 20*; (PAE *et al.* 2017)). The SCF complex
regulates the cell cycle (*ref 21-22*; (MARGOTTIN-GOGUET *et al.* 2003; ROGERS *et al.* 2009)) and is a known
regulator of centrosomal integrity (*ref 23-26*;(WOJCIK *et al.* 2000; MURPHY 2003; PHUONG THAO *et al.* 2006;
CUNHA-FERREIRA *et al.* 2009)), thereby influencing PGC specification (LERIT *et al.* 2017).
(**D**) Casp and TER94 assist in the degradation of Smaug. The SCF complex works as a ubiquitin E3 ligase to mark
Smaug for Degradation and PolyUb-Smaug is degraded by the action of Casp and TER94 during the late-MZT.
1164
1165
It would be reasonable to propose that Casp recruits TER94 via Casp's UBX domain

(Fig. 10B). Independent proteomic studies ((THOMSON *et al.* 2008; DEHAAN *et al.* 2017;

TENDULKAR *et al.* 2022); Fig 10 B) suggest multiple overlaps between Casp/TER94 and germ-

cell specific protein complexes (Fig. 10B), again suggesting functional relationships between

Casp/TER94 and germ cell determinants. In addition, the SCF complex contributes to cell cycle

progression and integrity of centrosomes ((WOJCIK *et al.* 2000; MURPHY 2003; PHUONG THAO *et*

al. 2006); Fig 10C). Total PGC count in early embryos depends on centrosome function and

efficient mitotic divisions. So SCF complex based degradation may have multiple targets that

participate during germ cell formation and specification (Fig. 10C).

Casp likely functions as an adapter protein that works in conjunction with the SCF

complex ((CAO *et al.* 2020; CAO *et al.* 2022); Fig. 10D) and TER94 (Fig. 10D). The binary

complex between TER94 and Casp subsequently targets ubiquitinated proteins such as Smaug

which are present in a complexed form with other proteins. Homeostatic maintenance of Smaug

levels via Casp-mediated proteolytic degradation could explain many of our observations.

Consistently, increased Smaug levels, due to diminished Casp function leads to loss of PGCs

whereas, *smaug* mutant embryos show the opposite phenotype (SIDDIQUI *et al.* 2023).

Intriguingly, even though Smaug and Oskar are members of a complex, the reduction in Casp

activity results in the stabilization of Smaug. By contrast, Oskar levels are diminished resulting

in loss of PGCs. The mechanism underlying the substrate-dependent divergent activities is not

known at this point.

The similarity between the phenotypic consequences due to loss of ZGA components

and Casp/TER94 also needs a special mention. The only study implicating TER94 during early

embryogenesis (Zᴇɴɢ *et al.* 2014) suggested that TER94 can potentiate BMP signaling.

Interestingly, *decapentaplegic* (*dpp*), a BMP ligand which is one of the important targets of ZGA

regulators including *Zelda*, can also influence the specification of embryonic PGCs

(Cᴏʟᴏɴɴᴇᴛᴛᴀ *et al.* 2023). PGCs need BMP signals (Dpp) to maintain their identity.

Furthermore, the exposure to BMP signals needs careful calibration as excess BMP signaling in

the surrounding soma leads to PGC loss. Importantly, components of protein degradation

machinery including those involved in ubiquitination (Smurf, a Ubiquitin ligase) and sumoylation

(Ubc9) appear to be involved in fine-tuning the signaling (Dᴇsʜᴘᴀɴᴅᴇ *et al.* 2014). Thus, it will

be of interest to determine if the aberrant germ cell specification observed due to loss of *casp*

and the components of BMP signaling pathway is mechanistically connected. This seems

especially relevant in the light of their respective dependence on the components of the protein

degradation machinery. It will be of considerable interest to investigate whether and how,

zygotic activity of Smaug, one of important canonical MBT regulators, fits into this picture. In

sum, while the specific details await detailed examination, it is apparent that possible

recapitulation of maternal regulation in the zygotic context may be a recurrent theme rather than

an isolated anomaly.

1204

1205

1206

1207

Fly husbandry and stocks: Flies were raised on standard cornmeal agar at 25 °C unless

stated otherwise. *Casp^lof* (11373), *Casp Df* (23691), *nos Gal4* (4937), *Mat α4-tubulinGal4:VP16*

(7063) and *TER94 RNAi* (32869) lines were procured from the Bloomington *Drosophila* Stock

Centre, with numbers in brackets indicated Stock numbers.

Cloning of *casp* deletion constructs and generation of transgenics: UASp-casp$^\Delta$ constructs

(*pUASp casp^ΔUBA, pUASp casp^ΔUAS, pUASp casp^ΔUBX, pUASp casp^ΔUASΔUBX*) were generated,

starting from a *pUASp-attB-Casp* construct (TENDULKAR *et al.* 2022), with the design including

an N-terminal HA tag. The constructs were injected into a *w^1118; Attp2* embryo in the NCBS-

CAMP transgenic injection facility (https://www.ncbs.res.in/research-facilities/drosophila-

services), and stable lines were generated by balancing against a *w^1118; TM3Sb/TM6Tb* animal.

For cloning and amplification, a functional N-terminal HA tag was introduced with the 5'-

forward primers and a 3'-homology arm was introduced using the 3'-reverse primers. The

exception was the *pUASp Casp^ΔUAS* construct where two fragments, one upstream and one

downstream of the UAS domain were amplified separately with 22 nucleotide homologous

overhang between them to mediate homologous recombination. The upstream fragment

contained the HA sequence while the downstream fragment contained the 3'-homology arm.

The sequences of the primers used are as follows. *TGTTCCAGATTACGCTGGCGGC* was

used as the 5'-forward primer for the *pUASp casp^wt, pUASp casp^ΔUBX, pUASp casp^ΔUASΔUBX* and

the upstream fragment of the *pUASp casp^ΔUAS* constructs.

TGTTCCAGATTACGCTGGCGGCCCCCATCCTATCCTGGTGCC was used as the 5'-forward

primer for the *pUASp casp^ΔUBA* construct.

CGGATGATGAGATAAGTGGCTCCACGGAAACATGCGAAATGTTTGAGGAGCAG was used

as forward primer for amplifying downstream fragment of the *pUASp casp^ΔUAS* construct.

ACCATGGGTTTAGGTATAATGTTATCAAGCTCC was the reverse primer for *pUASp casp^wt,*

pUASp casp^ΔUBA, and downstream fragment of *pUASp casp^ΔUAS*.

TTAGGTATAATGTTATCAAGCTCCTCATTCGGACGGCTCCTGAGGTAG was the reverse

primer for *pUASp casp^ΔUBX*.

TTAGGTATAATGTTATCAAGCTCCTCACGTGGAGCCACTTATCTCATCATCCG was used as

the reverse primer for *pUASp casp^ΔUASΔUBX*. *CGTGGAGCCACTTATCTCATCATCCG* was the

reverse primer for upstream fragment of *pUASp casp^ΔUAS*. PCR products were further amplified

with a common 5'-primer

ATAGGCCACTAGTGGATCTGATGTACCCATACGATGTTCCAGATTACGCTGGCGGC and

their respective reverse primers to introduce the 5'-homology arm. Inserts were recombined into

pUASp AttB vector linearised at the BamH1 site, using a variation of the SLiCE cloning method

(ZHANG *et al.* 2012; ZHANG *et al.* 2014). In short, E.coli DH10B expressing the optimised λ-

prophage red recombinase system (PPY cells) was cultured in the presence of arabinose to

induce recombinase expression and subsequently used for preparing competent cells. DNA

fragments containing homologous overhangs were co-transformed into the competent bacteria

and resultant colonies were screened through PCR to identify proper recombinants and

subsequently confirmed through sequencing.The pUASp *casp^{ΔFFAT}* was PCR amplified from

pRM-HA:casp ^{ΔFFAT} (TENDULKAR *et al.* 2022), cloned into pUASp-AttB, sequenced for validation

and injected into a *w^{1118}; Attp2* animal.

Embryonic lethality*:* 0-3 hr embryos were collected, transferred to a fresh sugar-agar plate,

and unhatched larvae were scored after 48 hours to determine viability.

Immunoprecipitation*:* 0–3-hour embryos were lysed in Co-IP Lysis Buffer (20mM Tris pH 8.0,

137mM NaCl, 1% IGEPAL, 2mM EDTA, 1X PIC) using a Dounce homogenizer, and centrifuged

at 21,000 g for 30 minutes. 3 mg of total lysate was incubated with 5 µg of primary antibody (Rb

anti-Casp) and 5 µg of Normal Rabbit IgG overnight at 4 °C. Antigen-antibody complexes were

captured using 50 µL of BioRad SureBeads Protein A (1614013) at 4 °C for 4 hours. Beads

were washed six times with Co-IP Lysis Buffer and protein complexes eluted by boiling in 1X

Laemmli Sample Buffer. Eluted proteins were resolved on a 10% polyacrylamide gel followed by

western blotting or in-gel trypsin digestion, described in the following sections.

Western blot analysis*:* Embryos collected at varied time points (0-3, 0-1, 1-2, 2-3 hours) were

lysed in RIPA buffer (50 mM Tris-Cl, 150 mM NaCl, 0.1% SDS, 0.01% Sodium azide, 0.5%

sodium deoxycholate, 1 mM EDTA, 1% Triton X-100, 1X PIC) with a pellet pestle (Kontes).

Lysates were cleared by centrifugation at 21,000 g at 4 °C for 30 minutes. Protein concentration

was estimated using a BCA assay (Pierce) and 30-40 µg of total protein was loaded onto the gel

after boiling in 1X Laemmli Sample Buffer. Proteins separated by 10% SDS-PAGE were

transferred onto a PVDF membrane (Immobilon-E, Merck) and blocked in 5% milk in Tris-Buffer

Saline (TBS) with 0.1% Tween 20 (TBS-T) for an hour. Blots were then incubated overnight with

primary antibody diluted in 5% milk in TBS-T, at 4 °C. Following three washes with TBS-T, blots

were incubated with secondary antibodies diluted in 5% milk in TBS-T, for 1 hour at room

temperature. Blots were washed thrice with TBS-T and visualized on a LAS4000 Fuji imaging

system after incubating with Immobilon Western Chemiluminescent HRP substrate (Merck). The

following antibodies were used: Rabbit anti-VAP, 1:10000 (TENDULKAR *et al.* 2022), Mouse anti-

α-Tubulin, 1:10000 (T6074, Sigma), Mouse anti-Ubiquitin, 1:1000 (P4D1, Santa Cruz

Biotechnology), Mouse anti-HA, 1:5000 (H3663 HA-7, Sigma-Aldrich), Rabbit anti-Casp,

1:10000 (TENDULKAR *et al.* 2022), 1:2000 (04-902 DW-2, Sigma-Aldrich) Goat anti-rabbit HRP

and Goat anti-mouse HRP secondary antibodies, each at 1:10000 (Jackson ImmunoResearch).

Rabbit anti-me31B, 1:1000, Rabbit anti-Smaug, 1:500, Rat anti-Tral, 1:1000 were a kind gift

from Elmar Whale (CAO *et al.* 2020).

In-gel Trypsin Digestion and LC-MS/MS Analysis*:* Before in-gel trypsin digestion of the Co-IP

eluate, the antibody was crosslinked to the SureBeads using DMP (Sigma) according to the

NEB crosslinking protocol to avoid elution of the antibody. After crosslinking 10µg Casp

antibody, Co-IP was performed as described above. In-gel trypsin digestion was carried out as

previously described (SHEVCHENKO et al. 2006). Briefly, Coomassie-stained bands on the gel

were excised and cut into 1 mm cubes. Gel pieces were transferred to a clean microcentrifuge

tube and destained with buffer containing 50% acetonitrile in 50 mM Ammonium bicarbonate.

Reduction and alkylation were carried out on the destained gel pieces by incubating with 10mM

dithiothreitol (DTT) followed by incubating with 20mM iodoacetamide. Gel pieces were saturated

with sequencing grade Trypsin (Promega) at a concentration of 10 ng/µL and incubated

overnight at 37 °C. Peptides were extracted by sequential addition of 100 µl of 0.4%

Trifluoroacetic acid (TFA) in 10% ACN, 100 µl of 0.4% TFA in 60% ACN and 100 µl of ACN.

The pooled extract was dried in a vacuum centrifuge and reconstituted with 50 µl of 0.1% TFA.

The peptides in TFA were purified using the StageTip protocol (RAPPSILBER et al. 2007). LC–

MS/MS analysis was performed on the Sciex TripleTOF6600 mass spectrometer interfaced with

an Eksigent nano-LC 425. Tryptic peptides (1 µg) were loaded onto an Eksigent C18 trap (5 µg

capacity) and subsequently eluted with a linear acetonitrile gradient on an Eksigent C18

analytical column (15 cm × 75-µm internal diameter). A typical LC run lasted 2 h post loading

onto the trap at a constant flow rate of 300 nL/min with solvent A consisting of water + 0.1%

formic acid and solvent B consisting of acetonitrile. The gradient schedule for the LC run was

5% (vol/vol) B for 10 min, a linear gradient of B from 0% to 80% (vol/vol) over 80 min, 80%

(vol/vol) B for 15 min and equilibration with 5% (vol/vol) B for 15 min. Data was acquired in an

information-dependent acquisition (IDA) mode over a mass range of 300–2,000 m/z. Each full

MS survey scan was followed by MS/MS of the 15 most intense peptides. Dynamic exclusion

was enabled for all experiments (repeat count 1; exclusion duration 6 s). Peptides were

identified and quantified using the SCIEX ProteinPilot software at a false discovery rate (FDR)

of 1%. A RefSeq *Drosophila* protein database (release 6) was used for peptide identification.

Proteins that were identified in two or more replicates were tabulated.

Fixation, immunostaining, and imaging of embryos: 0-3 hr embryos were collected in a

sieve and dechorionated in 4% sodium hypochlorite for 90 seconds. After thorough washes with

distilled water, embryos were fixed in a 1:1 heptane:4% PFA solution for 20 minutes. The PFA

layer was removed, and embryos in heptane were re-constituted with an equal volume of

methanol. Embryos were devitellinized by vigorous shaking in the 1:1 heptane:methanol

mixture. The heptane layer and the interphase containing non-devitellinized embryos were

carefully removed. Devitellinized embryos in the bottom methanol phase were washed twice

with methanol and stored at -20 °C till they were ready to be imunostained. For phalloidin and

Smaug staining, embryos fixed in heptane: 4% PFA were hand de-vitellinized, after which the

standard immunostaining procedure was followed. For immunostaining, embryos were

rehydrated by washing thrice with 0.3% PBS-TritonX 100 (PBS-T) for 15 minutes each.

Embryos were blocked in 2% BSA in 0.3% PBS-T for 1 hour at room temperature (RT).

Embryos were incubated at 4 °C overnight with primary antibodies diluted in 2% BSA in 0.3%

PBS-T at the appropriate dilutions. Following three 15-minute washes with 0.3% PBS-T,

embryos were incubated in the appropriate secondary antibodies for 1 hour at room

temperature. The following antibodies were used: Rabbit anti-Casp, 1:1000 (TENDULKAR *et al.*

2022); Rat anti-α-vasa; 1:50 (DSHB); Rabbit anti-VCP, 1:200 (#2648, Cell Signaling

Technology); Rabbit anti-Oskar, 1:1000 (generated by Mandy Jeske, Anne Ephrussi lab); Rabbit

anti-Smaug, 1:50; Mouse anti-γ-tubulin, 1:1000 (T6557, Sigma), Rabbit P-Histone H3 (S10)

1:300 (#9701, Cell Signalling); Alexa Fluor 568 Phalloidin, 1:1000 (Invitrogen) The following

secondary antibodies were used, goat anti-mouse Alexa488/ goat anti-rabbit Alexa568/ goat

anti-rat Alexa647/ goat anti-rabbit Alexa647/ goat anti-rabbit Alexa480, 1:1000 (DSHB).

Embryos were washed thrice with 0.3% PBS-T, and DAPI/Hoechst (1:500) was added in the

penultimate wash. Embryos were mounted in 70% glycerol and observed under a Leica sp8

confocal microscope with a 20X objective.

Live imaging of embryos: Embryos at the appropriate stage were washed and dechorionated

as described previously. A 2-well Nunc™ Lab-Tek™ II Chamber Slide™ System (ThermoFisher

Scientific) was affixed with a 3M Scotch double-sided tape, and dechorionated embryos with

intact vitelline membranes were mounted on the tape under a dissecting microscope.

Halocarbon oil 200 was used to cover the embryos to prevent dehydration during imaging.

Time-lapse imaging of embryos was performed on an inverted LSM confocal system (Zeiss

multiphoton 710) at 20X for ~8 hours. Embryo images were acquired at 5 min intervals.

1340

Quantification of western blots: Western blots were quantified using ImageJ. Each protein

was normalized to the tubulin loading control, and the highest signal for each was set to 1 and

all other tubulin band intensities were normalized against it. For MZT experiments, western blots

were used to determine protein degradation dynamics. 3-5 biological replicates, representing 3-

5 independent embryo collection were assayed. Band intensities were quantified using ImageJ

and normalized to α-tubulin as loading control. For each replicate intensities were normalized to

the first time point (0-1 hour). Significance thresholds are presented in the figures and figure

captions.

Quantification of the Smaug Immunostaining: All quantification of Smaug stained images

were done on ImageJ. ROIs were made for a single pole cell through the z-stacks, thereby only

analysing one pole cell at a time. For each embryo five such pole cells were measured. To

quantify intensity, mean intensity was measured for each ROI per stack and added to get the

final intensity. Each data point on the graph represents an average of all five cell intensities per

embryo. For Smaug punctae measurement, a combined ROI for each cell was taken and

duplicated. Each of the stack was then analysed using 3D objects counter, where thresholding

for all images was kept within the range of 80-90 with a minimum size filter kept at zero. The

number of objects counted was taken as the total number of Smaug punctae per cell. Five such

pole cells were analysed per embryo and the average was plotted as a single data point on the

graph. For quantification of poleplasm volume in the embryos, embryos were stained with

Oskar. An ROI (kept same for all images) encompassing the posterior part of the embryo was

drawn. Gaussian blur 3D (X sigma = 2, Y sigma = 2 and Z sigma = 2) was applied to each

image. The volume of pole plasm was analysed by using 3D objects counter after appropriate

thresholding.

1364
1365
1366

1367   **Competing interests**

1368   The authors declare no competing or financial interests.

1369

Funding

Science & Engineering Research Board (SERB) grant CRG/2018/001218 to GR. Pratiksha

Trust Extra-Mural Support for Transformational Aging Brain Research grant

EMSTAR/2023/SL03 to GR, facilitated by the Centre for Brain Research (CBR), Indian Institute

of Science, Bangalore. GD's visits to IISER Pune (2023-2025) are supported by the Ministry of

Education (MoE) Scheme for the promotion of Academic & Research Collaboration (SPARC),

Grant ID SPARC-1587, managed by IIT Kharagpur, which facilitates collaboration between

Grant ID SPARC-1587, managed by IIT Kharagpur, which facilitates collaboration between

IISER Pune and Princeton University. The IISER *Drosophila* media and Stock centre are

supported by the National Facility for Gene Function in Health and Disease (NFGFHD) at IISER

Pune, which in turn is supported by an infrastructure grant from the DBT, Govt. of India

(BT/INF/22/SP17358/2016). JS, NW and AR are undergraduates supported by INSPIRE/KVPY

fellowships. SD and SH are graduate students supported by research fellowships from the

Council of Scientific & Industrial Research (CSIR), Govt. of India.

1383

1384 **Contributions**

1385 SH and SD performed most of the experiments, with contributions from NW, JS, and AR. GR

1386 and GD conceptualised the project with input from SH and SD. SH, SD, GR, and GD analyzed

1387 and interpreted the data. GD, GR, SH, and SH wrote the manuscript. GR and GD supervised

1388 the project and acquired funding.

1389

1390 **Acknowledgements**

1395

1396

References

Adams, M. D., S. E. Celniker, R. A. Holt, C. A. Evans, J. D. Gocayne *et al.*, 2000 The genome
sequence of Drosophila melanogaster. Science 287**:** 2185-2195.
Arkov, A. L., J. Y. Wang, A. Ramos and R. Lehmann, 2006 The role of Tudor domains in germline
development and polar granule architecture. Development 133**:** 4053-4062.
Azuma, Y., T. Tokuda, M. Shimamura, A. Kyotani, H. Sasayama *et al.*, 2014 Identification of
ter94, Drosophila VCP, as a strong modulator of motor neuron degeneration induced by
knockdown of Caz, Drosophila FUS. Hum Mol Genet 23**:** 3467-3480.
Bayer, L. V., S. Milano, S. K. Formel, H. Kaur, R. Ravichandran *et al.*, 2023 Cup is essential for
oskar mRNA translational repression during early Drosophila oogenesis. RNA Biol 20**:**
573-587.
Belvin, M. P., and K. V. Anderson, 1996 A conserved signaling pathway: the Drosophila toll-
dorsal pathway. Annu Rev Cell Dev Biol 12**:** 393-416.
Blake-Hedges, C., and T. L. Megraw, 2019 Coordination of Embryogenesis by the Centrosome in
Drosophila melanogaster. Results Probl Cell Differ 67**:** 277-321.
Breitwieser, W., F. H. Markussen, H. Horstmann and A. Ephrussi, 1996 Oskar protein interaction
with Vasa represents an essential step in polar granule assembly. Genes Dev 10**:** 2179-
2188.
Brennan, C. A., and K. V. Anderson, 2004 Drosophila: the genetics of innate immune recognition
and response. Annu Rev Immunol 22**:** 457-483.
Brown, J. B., N. Boley, R. Eisman, G. E. May, M. H. Stoiber *et al.*, 2014 Diversity and dynamics
of the Drosophila transcriptome. Nature 512**:** 393-399.
Buchon, N., N. Silverman and S. Cherry, 2014 Immunity in Drosophila melanogaster--from
microbial recognition to whole-organism physiology. Nat Rev Immunol 14**:** 796-810.
Cao, W. X., S. Kabelitz, M. Gupta, E. Yeung, S. Lin *et al.*, 2020 Precise Temporal Regulation of
Post-transcriptional Repressors Is Required for an Orderly Drosophila Maternal-to-Zygotic
Transition. Cell Rep 31**:** 107783.
Cao, W. X., A. Karaiskakis, S. Lin, S. Angers and H. D. Lipshitz, 2022 The F-box protein Bard
(CG14317) targets the Smaug RNA-binding protein for destruction during the Drosophila
maternal-to-zygotic transition. Genetics 220.
Casas-Vila, N., A. Bluhm, S. Sayols, N. Dinges, M. Dejung *et al.*, 2017 The developmental
proteome of Drosophila melanogaster. Genome Res 27**:** 1273-1285.
Cavey, M., and T. Lecuit, 2008 Imaging cellular and molecular dynamics in live embryos using
fluorescent proteins. Methods Mol Biol 420**:** 219-238.
Chang, Y. C., W. T. Hung, Y. C. Chang, H. C. Chang, C. L. Wu *et al.*, 2011 Pathogenic
VCP/TER94 alleles are dominant actives and contribute to neurodegeneration by altering
cellular ATP level in a Drosophila IBMPFD model. PLoS Genet 7**:** e1001288.
Chen, Z. X., D. Sturgill, J. Qu, H. Jiang, S. Park *et al.*, 2014 Comparative validation of the D.
melanogaster modENCODE transcriptome annotation. Genome Res 24**:** 1209-1223.
Chu, K. T., X. H. Niu and L. T. Williams, 1995 A Fas-Associated Protein Factor, Faf1, Potentiates
Fas-Mediated Apoptosis. Proceedings of the National Academy of Sciences of the United
States of America 92**:** 11894-11898.
Cinalli, R. M., and R. Lehmann, 2013 A spindle-independent cleavage pathway controls germ cell
formation in Drosophila. Nat Cell Biol 15**:** 839-845.
Colonnetta, M. M., P. Schedl and G. Deshpande, 2023 Germline/soma distinction in Drosophila
embryos requires regulators of zygotic genome activation. Elife 12.

Cunha-Ferreira, I., A. Rodrigues-Martins, I. Bento, M. Riparbelli, W. Zhang *et al.*, 2009 The
SCF/Slimb ubiquitin ligase limits centrosome amplification through degradation of
SAK/PLK4. Curr Biol 19**:** 43-49.
Dahanukar, A., J. A. Walker and R. P. Wharton, 1999 Smaug, a novel RNA-binding protein that
operates a translational switch in Drosophila. Mol Cell 4**:** 209-218.
Dai, R. M., E. Chen, D. L. Longo, C. M. Gorbea and C. C. Li, 1998 Involvement of valosin-
containing protein, an ATPase Co-purified with IkappaBalpha and 26 S proteasome, in
ubiquitin-proteasome-mediated degradation of IkappaBalpha. J Biol Chem 273**:** 3562-
3573.
De Zio, D., E. Ferraro, M. D'Amelio, V. Simoni, M. Bordi *et al.*, 2008 Faf1 is expressed during
neurodevelopment and is involved in Apaf1-dependent caspase-3 activation in proneural
cells. Cell Mol Life Sci 65**:** 1780-1790.
DeHaan, H., A. McCambridge, B. Armstrong, C. Cruse, D. Solanki *et al.*, 2017 An in vivo
proteomic analysis of the Me31B interactome in Drosophila germ granules. FEBS Lett
591**:** 3536-3547.
Deshpande, G., E. Willis, S. Chatterjee, R. Fernandez, K. Dias *et al.*, 2014 BMP signaling and the
maintenance of primordial germ cell identity in Drosophila embryos. PLoS One 9**:** e88847.
Ephrussi, A., and R. Lehmann, 1992 Induction of germ cell formation by oskar. Nature 358**:** 387-
392.
Ewens, C. A., S. Panico, P. Kloppsteck, C. McKeown, I. O. Ebong *et al.*, 2014 The p97-FAF1
protein complex reveals a common mode of p97 adaptor binding. J Biol Chem 289**:** 12077-
12084.
Govind, S., 1999 Control of development and immunity by rel transcription factors in Drosophila.
Oncogene 18**:** 6875-6887.
Griciuc, A., L. Aron, M. J. Roux, R. Klein, A. Giangrande *et al.*, 2010 Inactivation of VCP/ter94
suppresses retinal pathology caused by misfolded rhodopsin in Drosophila. PLoS Genet 6.
Higashiyama, H., F. Hirose, M. Yamaguchi, Y. H. Inoue, N. Fujikake *et al.*, 2002 Identification
of ter94, Drosophila VCP, as a modulator of polyglutamine-induced neurodegeneration.
Cell Death Differ 9**:** 264-273.
Hoffmann, J. A., and J. M. Reichhart, 2002 Drosophila innate immunity: an evolutionary
perspective. Nat Immunol 3**:** 121-126.
Hultmark, D., 1993 Immune reactions in Drosophila and other insects: a model for innate
immunity. Trends Genet 9**:** 178-183.
Huynh, J. R., and D. St Johnston, 2004 The origin of asymmetry: early polarisation of the
Drosophila germline cyst and oocyte. Curr Biol 14**:** R438-449.
Imler, J. L., and J. A. Hoffmann, 2000 Signaling mechanisms in the antimicrobial host defense of
Drosophila. Curr Opin Microbiol 3**:** 16-22.
Jentsch, S., and S. Rumpf, 2007 Cdc48 (p97): a "molecular gearbox" in the ubiquitin pathway?
Trends Biochem Sci 32**:** 6-11.
Jeske, M., B. Moritz, A. Anders and E. Wahle, 2011 Smaug assembles an ATP-dependent stable
complex repressing nanos mRNA translation at multiple levels. EMBO J 30**:** 90-103.
Jones, J. R., and P. M. Macdonald, 2007 Oskar controls morphology of polar granules and nuclear
bodies in Drosophila. Development 134**:** 233-236.
Jongens, T. A., L. D. Ackerman, J. R. Swedlow, L. Y. Jan and Y. N. Jan, 1994 Germ cell-less
encodes a cell type-specific nuclear pore-associated protein and functions early in the
germ-cell specification pathway of Drosophila. Genes Dev 8**:** 2123-2136.

Jongens, T. A., B. Hay, L. Y. Jan and Y. N. Jan, 1992 The germ cell-less gene product: a posteriorly
localized component necessary for germ cell development in Drosophila. Cell 70: 569-584.
Kim, G., C. I. Pai, K. Sato, M. D. Person, A. Nakamura *et al.*, 2015 Region-specific activation of
oskar mRNA translation by inhibition of Bruno-mediated repression. PLoS Genet 11:
e1004992.
Kim, H., H. Zhang, D. Meng, G. Russell, J. N. Lee *et al.*, 2013 UAS domain of Ubxd8 and FAF1
polymerizes upon interaction with long-chain unsaturated fatty acids. J Lipid Res 54: 2144-
2152.
Kim, M., J. H. Lee, S. Y. Lee, E. Kim and J. Chung, 2006 Caspar, a suppressor of antibacterial
immunity in Drosophila. Proceedings of the National Academy of Sciences of the United
States of America 103: 16358-16363.
Kim-Ha, J., J. L. Smith and P. M. Macdonald, 1991 oskar mRNA is localized to the posterior pole
of the Drosophila oocyte. Cell 66: 23-35.
Kloppsteck, P., C. A. Ewens, A. Forster, X. Zhang and P. S. Freemont, 2012 Regulation of p97 in
the ubiquitin-proteasome system by the UBX protein-family. Biochim Biophys Acta 1823:
125-129.
Kubikova, J., G. Ubartaite, J. Metz and M. Jeske, 2023 Structural basis for binding of Drosophila
Smaug to the GPCR Smoothened and to the germline inducer Oskar. Proc Natl Acad Sci
U S A 120: e2304385120.
Kushimura, Y., T. Tokuda, Y. Azuma, I. Yamamoto, I. Mizuta *et al.*, 2018 Overexpression of
ter94, Drosophila VCP, improves motor neuron degeneration induced by knockdown of
TBPH, Drosophila TDP-43. Am J Neurodegener Dis 7: 11-31.
Lattao, R., L. Kovacs and D. M. Glover, 2017 The Centrioles, Centrosomes, Basal Bodies, and
Cilia of Drosophila melanogaster. Genetics 206: 33-53.
Lecuyer, E., H. Yoshida, N. Parthasarathy, C. Alm, T. Babak *et al.*, 2007 Global analysis of mRNA
localization reveals a prominent role in organizing cellular architecture and function. Cell
131: 174-187.
Lee, J. J., J. K. Park, J. Jeong, H. Jeon, J. B. Yoon *et al.*, 2013 Complex of Fas-associated factor 1
(FAF1) with valosin-containing protein (VCP)-Npl4-Ufd1 and polyubiquitinated proteins
promotes endoplasmic reticulum-associated degradation (ERAD). J Biol Chem 288: 6998-
7011.
Lehmann, R., 2016 Germ Plasm Biogenesis--An Oskar-Centric Perspective. Curr Top Dev Biol
116: 679-707.
Lemaitre, B., and J. Hoffmann, 2007 The host defense of Drosophila melanogaster. Annu Rev
Immunol 25: 697-743.
Lerit, D. A., and E. R. Gavis, 2011 Transport of germ plasm on astral microtubules directs germ
cell development in Drosophila. Curr Biol 21: 439-448.
Lerit, D. A., C. W. Shebelut, K. J. Lawlor, N. M. Rusan, E. R. Gavis *et al.*, 2017 Germ Cell-less
Promotes Centrosome Segregation to Induce Germ Cell Formation. Cell Rep 18: 831-839.
Li, J. M., H. Wu, W. Zhang, M. R. Blackburn and J. Jin, 2014 The p97-UFD1L-NPL4 protein
complex mediates cytokine-induced IkappaBalpha proteolysis. Mol Cell Biol 34: 335-347.
Mahowald, A. P., 2001 Assembly of the Drosophila germ plasm. Int Rev Cytol 203: 187-213.
Margottin-Goguet, F., J. Y. Hsu, A. Loktev, H. M. Hsieh, J. D. Reimann *et al.*, 2003 Prophase
destruction of Emi1 by the SCF(betaTrCP/Slimb) ubiquitin ligase activates the anaphase
promoting complex to allow progression beyond prometaphase. Dev Cell 4: 813-826.
Medzhitov, R., 2001 Toll-like receptors and innate immunity. Nat Rev Immunol 1: 135-145.

Menges, C. W., D. A. Altomare and J. R. Testa, 2009 FAS-associated factor 1 (FAF1) Diverse
functions and implications for oncogenesis. Cell Cycle 8**:** 2528-2534.
Meyer, H., M. Bug and S. Bremer, 2012 Emerging functions of the VCP/p97 AAA-ATPase in the
ubiquitin system. Nat Cell Biol 14**:** 117-123.
Meyer, H. H., 2005 Golgi reassembly after mitosis: the AAA family meets the ubiquitin family.
Biochim Biophys Acta 1744**:** 108-119.
Meyer, H. H., J. G. Shorter, J. Seemann, D. Pappin and G. Warren, 2000 A complex of mammalian
ufd1 and npl4 links the AAA-ATPase, p97, to ubiquitin and nuclear transport pathways.
EMBO J 19**:** 2181-2192.
Min-Young, P., M. Ji-Hyun and K. Eunhee, 2004 Fas associated factor 1, FAF1, impairs NF-
kappaB activation by interacting with the IkappaB kinase complex. Molecular Biology of
the Cell 15**:** 2544-2549.
Morais-de-Sa, E., A. Vega-Rioja, V. Trovisco and D. St Johnston, 2013 Oskar is targeted for
degradation by the sequential action of Par-1, GSK-3, and the SCF(-)Slimb ubiquitin
ligase. Dev Cell 26**:** 303-314.
Muers, M., 2011 Functional genomics: the modENCODE guide to the genome. Nat Rev Genet 12**:**
80.
Murphy, T. D., 2003 Drosophila skpA, a component of SCF ubiquitin ligases, regulates
centrosome duplication independently of cyclin E accumulation. J Cell Sci 116**:** 2321-
2332.
Nakamura, A., K. Sato and K. Hanyu-Nakamura, 2004 Drosophila cup is an eIF4E binding protein
that associates with Bruno and regulates oskar mRNA translation in oogenesis. Dev Cell
6**:** 69-78.
Nusslein-Volhard, C., 2022 The Toll gene in Drosophila pattern formation. Trends Genet 38**:** 231-
245.
Nusslein-Volhard, C., and E. Wieschaus, 1980 Mutations affecting segment number and polarity
in Drosophila. Nature 287**:** 795-801.
Pae, J., R. M. Cinalli, A. Marzio, M. Pagano and R. Lehmann, 2017 GCL and CUL3 Control the
Switch between Cell Lineages by Mediating Localized Degradation of an RTK. Dev Cell
42**:** 130-142 e137.
Park, M. Y., J. H. Moon, K. S. Lee, H. I. Choi, J. Chung *et al.*, 2007 FAF1 suppresses I kappa B
kinase (IKK) activation by disrupting the IKK complex assembly. Journal of Biological
Chemistry 282**:** 27572-27577.
Peters, J. M., M. J. Walsh and W. W. Franke, 1990 An abundant and ubiquitous homo-oligomeric
ring-shaped ATPase particle related to the putative vesicle fusion proteins Sec18p and
NSF. EMBO J 9**:** 1757-1767.
Phuong Thao, D. T., H. Ida, H. Yoshida and M. Yamaguchi, 2006 Identification of the Drosophila
skpA gene as a novel target of the transcription factor DREF. Exp Cell Res 312**:** 3641-
3650.
Raff, J. W., and D. M. Glover, 1989 Centrosomes, and not nuclei, initiate pole cell formation in
Drosophila embryos. Cell 57**:** 611-619.
Reim, G., M. Hruzova, S. Goetze and K. Basler, 2014 Protection of armadillo/beta-Catenin by
armless, a novel positive regulator of wingless signaling. PLoS Biol 12**:** e1001988.
Rogers, G. C., N. M. Rusan, D. M. Roberts, M. Peifer and S. L. Rogers, 2009 The SCF Slimb
ubiquitin ligase regulates Plk4/Sak levels to block centriole reduplication. J Cell Biol 184**:**
225-239.

Roth, S., 2023 Neofunctionalization of Toll Signaling in Insects: From Immunity to Dorsoventral Patterning. Annu Rev Cell Dev Biol 39**:** 1-22.

Ruden, D. M., V. Sollars, X. Wang, D. Mori, M. Alterman *et al.*, 2000 Membrane fusion proteins are required for oskar mRNA localization in the Drosophila egg chamber. Dev Biol 218**:** 314-325.

Ryu, S. W., and E. Kim, 2001 Apoptosis induced by human Fas-associated factor 1, hFAF1, requires its ubiquitin homologous domain, but not the Fas-binding domain. Biochem Biophys Res Commun 286**:** 1027-1032.

Ryu, S. W., S. J. Lee, M. Y. Park, J. Jun, Y. K. Jung *et al.*, 2003 Fas-associated factor 1, FAF1, is a member of Fas death-inducing signaling complex. Journal of Biological Chemistry 278**:** 24003-24010.

Schier, A. F., 2007 The maternal-zygotic transition: death and birth of RNAs. Science 316**:** 406-407.

Schuberth, C., and A. Buchberger, 2008 UBX domain proteins: major regulators of the AAA ATPase Cdc48/p97. Cell Mol Life Sci 65**:** 2360-2371.

Siddiqui, N. U., A. Karaiskakis, A. L. Goldman, W. V. I. Eagle, C. A. Smibert *et al.*, 2023 Smaug regulates germ plasm synthesis and primordial germ cell number in Drosophila embryos by repressing the oskar and bruno 1 mRNAs. bioRxiv.

Snee, M. J., and P. M. Macdonald, 2004 Live imaging of nuage and polar granules: evidence against a precursor-product relationship and a novel role for Oskar in stabilization of polar granule components. J Cell Sci 117**:** 2109-2120.

Song, E. J., S. H. Yim, E. Kim, N. S. Kim and K. J. Lee, 2005 Human Fas-associated factor 1, interacting with ubiquitinated proteins and valosin-containing protein, is involved in the ubiquitin-proteasome pathway. Mol Cell Biol 25**:** 2511-2524.

Song, J., J. K. Park, J. J. Lee, Y. S. Choi, K. S. Ryu *et al.*, 2009 Structure and interaction of ubiquitin-associated domain of human Fas-associated factor 1. Protein Sci 18**:** 2265-2276.

Tendulkar, S., S. Hegde, L. Garg, A. Thulasidharan, B. Kaduskar *et al.*, 2022 Caspar, an adapter for VAPB and TER94, modulates the progression of ALS8 by regulating IMD/NFkappaB-mediated glial inflammation in a Drosophila model of human disease. Hum Mol Genet 31**:** 2857-2875.

Thibault, S. T., M. A. Singer, W. Y. Miyazaki, B. Milash, N. A. Dompe *et al.*, 2004 A complementary transposon tool kit for Drosophila melanogaster using P and piggyBac. Nat Genet 36**:** 283-287.

Thomson, T., N. Liu, A. Arkov, R. Lehmann and P. Lasko, 2008 Isolation of new polar granule components in Drosophila reveals P body and ER associated proteins. Mech Dev 125**:** 865-873.

Thulasidharan, A., L. Garg, S. Tendulkar and G. S. Ratnaparkhi, 2024 Age-dependent dynamics of neuronal VAPB(ALS) inclusions in the adult brain. Neurobiol Dis. 196**:** 106517.

Tomancak, P., A. Beaton, R. Weiszmann, E. Kwan, S. Shu *et al.*, 2002 Systematic determination of patterns of gene expression during Drosophila embryogenesis. Genome Biol 3**:** RESEARCH0088.

Valanne, S., J. H. Wang and M. Ramet, 2011 The Drosophila Toll signaling pathway. J Immunol 186**:** 649-656.

Vanzo, N., A. Oprins, D. Xanthakis, A. Ephrussi and C. Rabouille, 2007 Stimulation of endocytosis and actin dynamics by Oskar polarizes the Drosophila oocyte. Dev Cell 12**:** 543-555.

Wang, L., and P. Ligoxygakis, 2006 Pathogen recognition and signalling in the Drosophila innate
immune response. Immunobiology 211**:** 251-261.
Weiszmann, R., A. S. Hammonds and S. E. Celniker, 2009 Determination of gene expression
patterns using high-throughput RNA in situ hybridization to whole-mount Drosophila
embryos. Nat Protoc 4**:** 605-618.
Wilk, R., J. Hu, D. Blotsky and H. M. Krause, 2016 Diverse and pervasive subcellular distributions
for both coding and long noncoding RNAs. Genes Dev 30**:** 594-609.
Williams, M. J., 2007 Drosophila hemopoiesis and cellular immunity. J Immunol 178**:** 4711-4716.
Wojcik, E. J., D. M. Glover and T. S. Hays, 2000 The SCF ubiquitin ligase protein slimb regulates
centrosome duplication in Drosophila. Curr Biol 10**:** 1131-1134.
Wu, J., and A. Akhmanova, 2017 Microtubule-Organizing Centers. Annu Rev Cell Dev Biol 33**:**
51-75.
Ye, Y., 2006 Diverse functions with a common regulator: ubiquitin takes command of an AAA
ATPase. J Struct Biol 156**:** 29-40.
Zaessinger, S., I. Busseau and M. Simonelig, 2006 Oskar allows nanos mRNA translation in
Drosophila embryos by preventing its deadenylation by Smaug/CCR4. Development 133**:**
4573-4583.
Zeng, Z., D. J. de Gorter, M. Kowalski, P. ten Dijke and O. Shimmi, 2014 Ter94/VCP is a novel
component involved in BMP signaling. PLoS One 9**:** e114475.
Zhang, L., F. Zhou, T. van Laar, J. Zhang, H. van Dam *et al.*, 2011 Fas-associated factor 1
antagonizes Wnt signaling by promoting beta-catenin degradation. Mol Biol Cell 22**:** 1617-
1624.
Zhang, Y., U. Werling and W. Edelmann, 2012 SLiCE: a novel bacterial cell extract-based DNA
cloning method. Nucleic Acids Res 40**:** e55.
Zhang, Y., U. Werling and W. Edelmann, 2014 Seamless Ligation Cloning Extract (SLiCE)
cloning method. Methods Mol Biol 1116**:** 235-244.
Zhang, Z., A. R. Krauchunas, S. Huang and M. F. Wolfner, 2018 Maternal Proteins That Are
Phosphoregulated upon Egg Activation Include Crucial Factors for Oogenesis, Egg
Activation and Embryogenesis in Drosophila melanogaster. G3 (Bethesda) 8**:** 3005-3018.
1656

A

Source: FlyBase (flybase.org)

B

Source: Brown et. al. (2014), *Nature* 512: 393.

C

Fig. S1. **(A)** Schematic of the the chromosomal location of the *casp* locus and the associated transposon insertion that generates the *casp^{c04227}* allele. This allele is referred to as *casp^{lof}* hereafter. **(B)** RNA expression data from the modENCODE database, plotted as a bar chart **(C)**. DAPI marks the nuclei, orb marks the oocyte. Casp marks the follicle cells, but is not significantly enriched in the germline (egg, nurse cells).

Fig. S3. TER94 and germline components are major interactors of Casp. (**A**) shows a list of the 24 proteins enriched after a Casp IP, followed by mass spectrometry, listed in the order of their peptide counts. The peptide counts are averaged from 4 biological replicates. A subset of interactors with known functions or expression in the germline are tabulated in (**B**) A schematic of the interaction of Casp with both TER94 and VAPB (Tendulkar et al., 2022). TER94 interacts with the Casp UBX domain, while VAPB with the FFAT motif (**C**) List of proteins that are associated with Casp and are previously identified as pole-cell components (**D-E**) Immunostaining of wild-type nuclear cycle 13/14 embryos with Casp (D2) and TER94-specific (E2) antibodies indicates expression in the primordial germ cells (C1-D4). Hoechst marks nuclei and Vasa marks the germ cells.

Fig S4. Casp:VAP interaction is not required for defining pole cell number. (**A**) Casp interacts with VAPB via its FFAT-like motif (**B**) represents embryonic lethality at 25° C plotted as a bar graph. The genotype of the mother is listed on the X-axis. N = 3. n = 200, ordinary one-way ANOVA, (****) P <0.0001. (**C**) Casp protein levels were assessed in 0-3h embryos via western blotting. Rabbit anti Casp (1:10,000) and rabbit anti HA (1:2000) antibodies were used to probe the blot. Mouse anti Tubulin (1:10,000)is used as a loading control. (**D-F**) Confocal microscopy images of the posterior of nuclear cycle 13/14 embryos from mated *nos gal4* (D1-D3), *nos gal4> casp$^{\Delta FFAT}$* (E1-E3), and *casp^{lof}/Df; nos gal4> casp$^{\Delta FFAT}$* (F1-F3) females immunostained with vasa antibody (1:50). Hoechst marks the nuclei. (**G**) The germ cells at the posterior marked by vasa quantified and plotted as bar graphs. n = 30-40. Ordinary one-way ANOVA, (****) P <0.0001