The T3SS structural and effector genes of *Chlamydia trachomatis* are expressed in distinct phenotypic cell forms.

Authors:
Nicole A. Grieshaber*, Cody Appa*, Megan Ward*, Alorah Grossman*, Sean McCormik*, Brendan S. Grieshaber*, Travis Chiarelli*, Hong Yang**, Anders Omsland**, and Scott S. Grieshaber*+

*Department of Biological Sciences, University of Idaho, Moscow, ID
**Allen School for Global Health, Washington State University, Pullman, WA

+ Corresponding Author: Scott S. Grieshaber, scottg@uidaho.edu

Abstract:
Bacteria in the chlamydiales order are obligate intracellular parasites of eukaryotic cells. Within this order, the genus *Chlamydia* contains the causative agents of a number of clinically important infections of humans. Biovars of *C. trachomatis* are the causative agents of trachoma, the leading cause of preventable blindness worldwide, as well as sexually transmitted infections with the potential to cause pelvic inflammatory disease and infertility. Irrespective of the resulting disease, all chlamydial species share the same obligate intracellular life cycle and developmental cell forms. They are reliant on an infectious cycle consisting of at least three phenotypically distinct cell forms termed the reticulate body (RB), the intermediate body (IB) and the elementary body (EB). The EB is infectious but does not replicate. The RB replicates in the host cell but is non-infectious, while the IB is an intermediate form that transitions to the EB form. In this study, we ectopically expressed the transcriptional repressor Euo, the two nucleoid-associated proteins HctA and HctB, and the two component sensor kinase CtcB in the RB. Transcriptional analysis using RNA-seq, differential expression clustering and fluorescence *in situ* hybridization analysis show that the chlamydial developmental cycle is driven by three distinct regulons corresponding to the RB, IB or EB cell forms. Moreover, we show that the genes for the T3SS were cell type restricted, suggesting defined functional roles for the T3SS in specific cell forms.

Importance:
Chlamydia trachomatis, a sexually transmitted bacterial infection, poses a significant global health threat, causing over 100 million infections annually and leading to complications like ectopic pregnancy and infertility. This study investigates the gene expression patterns of *Chlamydia trachomatis* during its unique life cycle within human cells. As an obligate intracellular parasite, *C. trachomatis* transitions through distinct developmental stages - one for infection and dissemination, another for replication, and a third for transitioning back to the infectious form. By analyzing gene expression profiles at each stage, we identified key genes involved in these processes. Interestingly, our research also reveals the presence of two separate T3SS (Type III Secretion System) translocons expressed in distinct stages, suggesting their crucial roles in specific functions during the infection cycle.

Introduction:

Many bacterial species undergo dramatic phenotypic changes to adapt to different environments or to generate cells with specific physiological functions. All the bacteria in the Chlamydiales order are obligate intracellular parasites of eukaryotic cells that undergo a developmental cycle with both non-replicating and actively replicating cell forms [1,2]. Chlamydial species are important pathogens of humans. *C. psittaci* causes zoonotic infections resulting in pneumonia, while *C. pneumoniae* is a human pathogen that causes respiratory disease. Biovars of *C. trachomatis (Ctr)* are the causative agents of trachoma, the leading cause of preventable blindness worldwide, as well as sexually transmitted infections with the potential to cause pelvic inflammatory disease, ectopic pregnancy and infertility [3–5].

Success of a chlamydial infection depends on the completion of a complex intracellular developmental cycle, consisting of multiple cell forms; the elementary body (EB), the reticulate body (RB) and the intermediate body (IB) [1,2]. Although the timing of cell type conversion may differ, the broad strokes of this cycle are conserved in all the Chlamydiaceae [6,7]. Our current understanding of the developmental cycle as determined through promoter reporter strains, single inclusion kinetics, single cell gene expression and agent based modeling has led to a clearer picture of the cycle [8,9]. The EB, characterized by its condensed nucleoid and small size (~0.2 nm diameter), initiates infection of the host through the use of a type III secretion system (T3SS) and pre-formed effectors [10,11]. These effectors promote pathogen phagocytosis and entry into the targeted cell. After entry, the EB form resides in an endocytic vesicle termed the inclusion that is modified through chlamydial gene expression [10,11]. The EB completes EB to RB differentiation and becomes replication competent at ~10 hpi (*Ctr* serovar L2) [12]. The RB, which is phenotypically characterized as larger than the EB (~1 nm diameter) and containing a dispersed nucleoid, then undergoes several rounds of amplifying replication before maturing to produce IB cells that then progress to the infectious EB, a process that takes place over ~8-10 hours after IB formation [8,9]. The mature RBs continue to produce IB cells, acting as a stem cell population [9]. This developmental program results in a growth cycle that does not act like a typical bacterial growth culture (lag, log, stationary phase) but instead asynchronously progresses through the RB, IB and EB cell type transitions until cell lysis or inclusion extrusion [1,2,8,9].

The current understanding of the regulation of the developmental cycle comes primarily from population level studies that frame the cycle in terms of time, treating the chlamydial population as a time dependent uniform culture. Population level gene expression data has been determined for chlamydial infections and has been described according to time after infection. These studies include RT-qPCR, microarray and RNA-seq studies and contribute to the canonical early (~0-10 hpi, EB to RB differentiation), mid-cycle (~10-18 hpi, RB replication) and late (~18 hpi-cell lysis, EB formation) gene expression paradigm [13–16]. The reliance on population level data of a mixed cell population has confounded the understanding of gene expression as it pertains to the specific chlamydial cell forms.

Here we sought to define the transcript profiles of the cell forms that underpin the observed growth cycle by investigating the effects of ectopic expression of four transcriptional regulatory proteins in *Chlamydia trachomatis* (*Ctr*): Euo, HctA, HctB and CtcB. Euo (Early Upstream ORF) is among the earliest genes expressed post EB to RB differentiation during chlamydial infection [17]. Current evidence suggests that Euo is a DNA binding protein that acts to repress a handful of late cycle genes [17,18].

The HctA protein is a small DNA binding protein that is expressed transiently in the IB cell type ~8-10 hrs before HctB [8,9]. HctA has been shown to bind DNA and to repress transcription broadly across chromosomes of both *Ctr* and, when ectopically expressed, *E. coli* [19–21]. HctB is a second small positively charged protein that has limited homology to the H1 histone family and is thought to contribute to condensation of the EB nucleoid [22]. Our data show that unlike HctA, HctB is expressed late in EB development, during the final stages of EB formation [8,9]. In addition to these DNA binding proteins, *Ctr* contains a single cytosolic two component regulatory system (TCS) consisting of CtcB/CtcC (histidine kinase/response regulator) [23]. The *Ctr* TCS is actively transcribed during RB to EB development, and the protein products are functional with respect

to phosphotransfer [23]. Additionally, expression of the ATPase effector domain of the response regulator, CtcC,
resulted in the up regulation of the sigma54 regulon which included many developmentally regulated genes [24].
We expect that ectopic expression of CtcB would phosphorylate CtcC and amplify the signaling and activation
process of sigma 54 gene expression.
We have shown that Euo, HctA and HctB promoter activities help define the RB, IB and EB cell forms
[8,9]. Therefore, along with CtcB we determined the effects of ectopic expression of these regulatory factors on
the transcriptome of *Ctr* using RNA-seq. Our data produced gene regulation profiles consistent with cell form-
specific transcriptomes. This allowed us to bin the expression of a large fraction of the chlamydial genome into
RB, IB and EB-specific transcript categories. Within our cell form expression prediction groups was a number
of T3SS genes. Using fluorescence *in situ* hybridization (FISH), we showed that components of the T3SS were
expressed in specific cell forms.
104
Results:
Ectopic expression of Euo, HctA, CtcB, and HctB resulted in disruption of the developmental cycle.
To determine the effects of the ectopic expression of Euo, HctA, HctB and CtcB on gene expression and the
developmental cycle, we expressed these proteins and the GFP protein Clover as a control from the
chlamydial plasmid. We chose to induce expression at 15 hpi in order to evaluate the effects of these proteins
on the RB to EB stage of the developmental cycle. At 15 hpi the vast majority of the chlamydial cells would be
in the RB form [9].
We expressed each of these constructs in *Ctr* using the native plasmid, the T5 promoter (*E. coli*
sigma70 constitutive promoter) and the E riboswitch for conditional translational expression [25]. The ORFs of
each of the four transcriptional regulatory proteins (*euo*, *hctA*, *ctcB*, and *hctB*) were cloned upstream of a C-
terminal FLAG tag and transformed into *Chlamydia trachomatis* serovar L2 434/bu (*Ctr*) creating the strains L2-
E-euo-FLAG, L2-E-hctA-FLAG, and L2-E-ctcB-FLAG. As described previously, we could not successfully
transform *Ctr* with the E-hctB-FLAG construct, therefore we developed a tet-riboJ-E promoter system that
combines both transcriptional and translational control to hctB-FLAG expression, creating the strain L2-tet-J-E-
hctB-FLAG [25].
Cells were infected with each strain and expression of Euo-FLAG, HctA-FLAG, CtcB-FLAG, HctB-
FLAG and Clover-FLAG was induced at 15 hpi. We tested for the production of infectious progeny (EBs) using
a reinfection assay at 48 hpi. Ectopic expression of all four transcriptional regulatory proteins resulted in a
significant inhibition of EB production as compared to the Clover-FLAG controls (Fig. 1A). In addition, infected
cells were imaged using transmission electron microscopy (TEM). Cells were infected with L2-E-euo-FLAG,
L2-E-hctA-FLAG, L2-E-ctcB-FLAG, L2-tet-J-E-hctB-FLAG and L2-E-clover-FLAG, induced for expression at 15
hpi and fixed and prepared for TEM analysis at 30 hpi (Fig. 1B). The induced and uninduced Clover-FLAG
control samples were indistinguishable. The inclusions for both samples had similar ratios of RBs (large cell
forms) and EBs (small electron dense forms). Cells ectopically expressing Euo had inclusions with very few
visible EBs (small electron dense forms) and most cells appeared RB like (large less electron dense cell
forms). In the HctB and HctA expressing bacteria, the inclusions contained few abnormal RB like forms as well
as cells with dense structures resembling condensed nucleoids (electron dense regions inside cells). The
inclusions of the CtcB expressing bacteria contained both RB and EB-like cells and an increase in intermediate
forms, i.e. RB sized cells with condensed nucleoids (Fig. 1B). These data suggested that ectopic expression of
all four of these transcriptional regulatory proteins resulted in an aborted developmental cycle as indicated by
both the IFU studies and the dysregulated cell forms seen by EM.
136
RNA-seq of the ectopic protein expressing chlamydial strains. To better understand the effects of the
ectopic expression of each regulatory protein, we used RNA-seq to characterize the corresponding
transcriptomes. We infected host cells with each strain (L2-E-clover-FLAG, L2-E-euo-FLAG, L2-E-hctA-FLAG

L2-E-ctcB-FLAG and L2-tet-J-E-hctB-FLAG), induced expression at 15 hpi and harvested RNA for library
construction at 18 hpi and 24 hpi. We chose to investigate gene expression three hours after induction (18 hpi)
to capture potential immediate effects on the developmental cycle. We also investigated gene expression at 9
hours after induction (24 hpi) as we reasoned that this time point would capture potential effects of inhibition of
the advancement of the cycle. This later time point allowed for the detection of changes between the
advancement of the cycle in control samples vs potential inhibition of the cycle by ectopic expression of the
regulatory proteins.

We compared the transcriptome of each sample in triplicate using principal component analysis (PCA).
Each of the triplicate samples clustered closely together suggesting little differences between the biological
replicates (Fig. 2A). As expected, the Clover control samples clustered in distinct groups depending on
isolation timepoint. The 18 hpi samples clustered together in a grouping and the 24 hpi samples clustered in a
separate distinct grouping. For the Euo induced samples, all the 18 hpi and 24 hpi samples clustered closely
together suggesting only small differences between both the time point samples. The timepoint experimental
samples were also clustered together for the HctA 18 hpi and 24 hpi samples. For the HctB 18 hpi and 24 hpi
experimental samples, each time point clustered separately but the two clusters were closer to each other than
any of the other experimental conditions. The CtcB 18 hpi and 24 hpi experimental samples were similar, the
replicates for each time point clustered tightly together and the 18 hpi and 24 hpi samples clustered closer to
each other than to the samples from the other experimental conditions.(Fig. 2A). Together, these data suggest
that each ectopically expressed protein generated a unique gene expression pattern.

The RNA-seq data sets were compared to the induced Clover controls; 18 hpi Clover to 18 hpi
experimental sample, and 24 hpi Clover to 24 hpi experimental samples. In addition to the ectopic data, we
used the gene expression data from our published data set (NCBI's Sequence Read Archive (SRA),
submission number GSE276506) and determined the differential gene expression between *Chlamydia* from
the 18 hpi sample and the 24 hpi sample, capturing changes in late gene expression (RB to IB and EB). We
compared this differential gene expression pattern to the differential gene expression patterns of the ectopic
expression experimental data. We generated a hierarchically-clustered heatmap using the Seaborn clustering
algorithm [26] (Fig. 2B and C). The 24 hpi samples for Euo and HctB were used for clustering analysis as these
proteins acted as inhibitors and blocked cycle progression (Fig. 2B and C). The 24 hpi samples allowed more
time for accumulated changes as the Clover controls progressed to the production of late genes while the Euo
and HctB expressing samples did not. The 18 hpi data from the HctA and CtcB ectopic expression experiments
were used for cluster analysis as they both acted as inducers (Fig. 2B and C). These changes were most
obvious in the 18 hpi samples as the Clover controls had not begun to express late genes.

Clustering produced two dominant groups (Fig. 2B and C). Both groups featured genes that were
dramatically up-regulated between 18 hpi vs 24 hpi during the wt infection (Fig. 2B and C, HeLa_24h)
suggesting that all of these genes would be considered late genes [14,15]. The major difference between the two
cluster groups is the changes in gene expression induced by HctA and CtcB ectopic expression (Fig. 2B and
C). One of the clusters featured genes that were not dramatically induced upon HctA-FLAG or CtcB-FLAG
ectopic expression (Fig. 2B). The other cluster showed the opposite with all the genes upregulated by the
ectopic expression of HctA-FLAG and CtcB-FLAG (Fig. 2C). Ectopic expression of Euo-FLAG and HctB-FLAG
led to a down-regulation of both of these sets of genes as compared to the Clover control (Fig. 2B and C).
Many of the genes in the first cluster (Fig. 2B) have been shown to be expressed mid-cycle or late-cycle [14,15],
while most of the genes in the second cluster (Fig. 2C) have been identified to be expressed late in the
developmental cycle [14,15]. Additionally, many of the second cluster genes have been identified as
sigma54/ctcB-ctcC regulated genes [24,27]. These data suggest that the late expressed genes can be divided into
two distinct categories.

Gene expression cluster groups map to cell type specific gene expression profiles. Using the clustering

data observations, we created a selection criteria to categorize the RNA-seq data into three gene expression
categories (Table S1). The first group were genes for which we observed little to no change after Euo ectopic
expression when compared to the Clover control and had little to no change between 18 hpi and 24 hpi during
a wt infection. We separated the late genes into two groups. The first group was defined as genes whose
expression increased between 18 hpi and 24 hpi in the wt infection but were not induced by HctA, CtcB or Euo
ectopic expression. The second group of late genes were defined as genes whose expression was increased
from 18 hpi to 24 hpi in the wt infection and were up-regulated by CtcB and HctA ectopic expression but not
increased by Euo ectopic expression. Based on the observation that *euo* was a member of the first group, we
defined these genes as RB genes (Table S1). For the two late gene groups we noticed that *hctA*, an IB gene
[8,9], was a member of the first group and therefore designated this group as IB genes (Table S1). The second
late gene group was designated as EB genes, i.e., the regulon likely involved in the final stage of generating
EB infectivity. This group contains the hctB, tarp, and scc2 genes that we have previously shown to be
expressed late in the IB to EB developmental stage [8,9].

We next used volcano plots to visualize the individual effects of each of the ectopic expression
constructs on changes in gene expression of all *Ctr* genes. The expression changes were plotted (log2fold
change) vs statistical significance (-log of the p value) and the RB, IB and EB gene groups designated in Table
S1 were highlighted (Fig. 3). We plotted changes in gene expression from 18 hpi to 24 hpi from a wt infection,
which as expected indicated that the genes from Table S1 designated as RB genes were largely unchanged in
gene expression between 18 hpi and 24 hpi, while both the designated IB and EB genes showed increased
expression. RB gene expression was for the most part unchanged when Euo was ectopically expressed in
contrast to dramatic reductions in the expression of both IB and EB designated genes (Fig. 3, Euo 24hpi vs
Clover 24hpi). The ectopic expression of both HctA and CtcB dramatically increased the expression of EB
genes (Fig. 3, HctA and CtcB 18hpi vs Clover 18hpi). HctB ectopic expression also resulted in a dramatic
repression in the expression of both IB and EB genes while also increasing the relative expression of a subset
of RB genes relative to Clover (Fig. 3, HctB 24hpi vs Clover 24hpiE). We noticed that many of the genes that
showed increased expression when HctB was ectopically expressed were ribosomal protein genes (Fig. 3,
HctB 24hpi vs Clover 24hpi). We interpreted this effect as HctB inhibiting the expression of most genes with
the exception of the ribosomal genes, perhaps allowing them to be expressed early during EB to RB
differentiation upon the next infection cycle.

Fluorescence *in situ* hybridization verification of cell type gene expression. To verify the association of
binned genes with specific cell forms we used fluorescence *in situ* hybridization (FISH) to visualize gene
expression in cells expressing GFP and RFP from developmental stage-specific promoters. To this end, we
constructed two dual promoter reporter constructs to delineate gene expression from the *euo*, *hctA* and *hctB*
promoters which we have shown to be associated with RB, IB and EB cells forms respectively. We generated
the strains L2-*hctB*prom_Scarlet-*euo*prom_neongreen (L2-BsciEng) and L2-*hctA*prom_Scarlet-
*euo*prom_neongreen (L2-AsciEng) which express the RFP mScarlet-I (sci) from either the *hctB* promoter or
hctA promoter along with the GFP protein Neongreen (ng) driven by *euo* promoter. To validate our system, the
mRNA expression of *euo*, *hctA* and *hctB* was visualized in each strain using custom FISH probes from
Molecular Instruments (Fig. 4). Cells were infected with L2-AsciEng and L2-BsciEng and processed for each
FISH probe at 24 hpi. Although dual promoter strains were used in these experiments, the data is presented in
a single promoter format to simplify presentation. *Euo* and *hctA* data was processed from L2-AsiEng samples
and *hctB* data was processed from L2-BsciEng samples. As expected, *euo* mRNA was observed primarily in
the *euo*prom+ (RB) cells and not in either of the *hctA*prom+ (IB) or *hctB*prom+ (EB) cells (Fig. 4A). The *hctA*
mRNA was observed in a subset of cells that had overlap with the *hctA*prom signal but not *euo*prom or
*hctB*prom signal (Fig. 4B). The *hctB* mRNA was observed in a subset of cells with overlap with *hctA*prom+ and
*hctB*prom+ cells but not *euo*prom+ cells (Fig. 4C).

To quantify the FISH results, we used the TrackMate plugin in Fiji [28] to identify and measure both the
mRNA signal and promoter reporter signal for each chlamydial cell in five inclusions from each infection. Cells
were identified by their promoter reporter signal (green) and separately by their mRNA fluorescence signal
(pink). The fluorescence intensity was measured and plotted for both channels (FISH and promoter reporter) in
both identified populations. Therefore, each plot represents two identified cell populations per reporter, mRNA+
(pink) population and their corresponding promoter reporter fluorescence intensity, and the promoter reporter+
(green) population and their corresponding FISH signal. This analysis was performed for all three promoter
reporters (*euo*p, *hctA*p and *hctB*p) for each FISH mRNA probe (*euo*, *hctA*, and *hctB*) (Fig. 4).

RB: *euo* FISH. Using Fiji and TrackMate, we identified individual chlamydial cells expressing the *euo*
mRNA in host cells that were infected with the promoter reporter strains expressing fluorescent proteins from
the *euo*prom, *hctA*prom, and *hctB*prom. The analysis indicates that the *euo* mRNA+ cell population (pink)
when plotted for *euo*prom fluorescence and *euo* mRNA fluorescence were primarily double positive with high
levels of both *euo* FISH signal and *euoprom* signal. These *euo* mRNA+ cells from *hctAprom* infections were
primarily single positive. This was also observed for the *hctB*prom infections; the *euo* mRNA+ cells when
plotted for *euo* mRNA FISH signal intensity against *hctB*prom signal intensity were primarily single positive
(Fig. 4A).

We also used TrackMate to identify the promoter reporter positive cell populations (green) and plotted
the expression intensities of the promoter reporter signals against the *euo* mRNA FISH signal. The *euo*prom+
cells were predominantly double positive (high *euo* mRNA signal, high *euo*prom signal), while the *hctA*prom+
cells and *hctB*prom+ cell populations (green) were predominantly single positive (low *euo* mRNA signal high
*hctA*prom or *hctB*prom signal) (Fig. 4A).

IB: *hctA* FISH. The *hctA* mRNA positive population was identified in the *euo*prom, *hctA*prom and
*hctB*prom infected cells using TrackMate (magenta) and the intensities of the *hctA* mRNA FISH signal was
plotted against each of the promoter reporter intensity signals. The *hctA* mRNA+ cells from the *euo*prom
infection were mostly single positive but had carryover Neongreen signal from RB *euo*prom expression (Fig.
4B). The *hctA* mRNA+ cell population (magenta) was mostly double positive when compared to the *hctA*prom
signal. Additionally, the *hctA* mRNA+ population had little to no *hctB*prom signal (Fig. 4B).

We identified the promoter reporter positive chlamydial cells (green) and plotted both the promoter
reporter signals and the FISH signal. The *euo*prom+ cell population (green) was mostly single positive with a
small population of double positive cells presumably from the long lived Neongreen protein carry-over (Fig.
4B). The *hctA*prom+ cell population (green) demonstrated a large double positive sub-population as well as a
single positive sub-population that again was likely due to the long half-life of the mScarlet-I protein carried
over into the EB population (Fig. 4B). The *hctB*prom+ cell population was almost entirely single positive with
little *hctA* mRNA signal (Fig. 4B).

EB: *hctB* FISH. We performed the same analysis for the *hctB* mRNA+ cells (magenta). The *hctB*
mRNA+ chlamydial cells were mostly single positive when compared to the *euo*prom signal with some
detected long-lived *euo*prom signal (Fig. 4C). The *hctB* mRNA+ cells were generally double positive for the
*hctA*prom signal and double positive for the *hctB*prom signal (Fig. 4C). For the promoter reporter cells (green),
the *euo*prom+ cells had a significant single positive population and a smaller double positive sub-population,
while the *hctA*prom+ cells were both single and double positive for the *hctB* mRNA signal. In the *hctB*prom+
cells there was both a single and double positive population (Fig. 4C).

The apparent disconnect between mRNA expression and cognate fluorescent protein fluorescence for
euo, hctA and hctB FISH results are as expected as the fluorescent proteins have a much longer half life than
mRNA. Additionally, fluorescence from mScarlet-I and Neongreen proteins lags mRNA expression as the
proteins must be translated and then folded into the mature fluorescent state. Together, these data indicate
that, as expected, euo mRNA is expressed in RBs (double positive for euoprom signal and euo mRNA signal),
while hctA mRNA is expressed in IBs (hctAprom+, hctA mRNA+ and hctBprom negative) and hctB mRNA is

expressed in late IB/EBs (hctBprom+ and hctB mRNA+). Therefore, we used this workflow to interrogate cell
form gene expression predicted by the RNA-seq clustering, binning and volcano plot analysis.
283
Validation of *porB* as an IB gene. We next sought to determine if a gene predicted to be expressed in a
particular cell form by the RNA-seq clustering experiment could be verified. The porin gene *porB* [29] which
clustered with the IB gene group (Fig. 2B) as well as with hctA (a proven IB gene) was selected for the
analysis. Cells were infected with either L2-AsciEng or L2-BsciEng, fixed at 24 hpi and probed for the *porB*
mRNA. Confocal images were taken and viewed as z projections (Fig. 5A). The *porB* mRNA signal (magenta)
did not completely overlap with the euoprom signal (green) and appeared to be expressed in a subset of cells
(Fig. 5A). The *porB* mRNA had significant but not complete overlap with the *hctA*prom+ cells (green) and
almost no overlap with the *hctB*prom+ cells (Fig. 5A). Using the TrackMate protocol described above we
identified the promoter reporter expressing populations, *euo*prom, *hctA*prom, and *hctB*prom (green) and then
separately, the *porB* mRNA+ population (magenta) and measured the fluorescence intensity of each channel
(fluorescent reporter proteins and mRNA signal within each population). The *porB* mRNA+ population
(magenta) in the *euo*prom channel experiment were a mix of single and double positive cells. This was also
true for the euoprom+ population (green). In comparison, the *porB* mRNA+ population (magenta) in the
*hctA*prom channel experiment were primarily double positive, i.e. high *porB* mRNA and high *hctA*prom activity.
However, the *hctA*prom+ population (green) was a mix of single positive and double positive cells (Fig. 5A). In
the *hctB*prom channel experiment, the *porB* mRNA signal+ population (magenta) were primarily single positive
and did not have appreciable *hctB*prom fluorescence. Conversely the *hctB*prom+ population (green) was
primarily single positive with low *porB* mRNA signal. Taken together, the *porB* mRNA expression pattern was
similar to that of the *hctA* mRNA expression pattern strongly suggesting *porB* is expressed primarily in the IB
cell form.
We also evaluated the kinetics of the activity of the *porB* promoter. The *hctA* promoter from AsciEng
was replaced with the promoter region of *porB* (-137bp to +30bp) and transformed into *Ctr* to create L2-
PsciEng. We used live cell imaging to measure the expression of Neongreen driven by *euo*prom and mScarlet-
I driven by *porB*prom. Cells were infected with PsciEng at an MOI ~0.3 and imaged for both the Neongreen
and mScarlet-I fluorescence every 30 minutes from 10 hpi for 55 hours. For comparisons, L2-AsciEng and L2-
BsciEng strains were imaged in parallel. As previously published, *euo* promoter activity is detected at ~12 hpi
followed by the *hctA* promoter and finally the *hctB* promoter [8,9] (Fig. 5B). The kinetics of L2-PsciEng mirrored
that of L2-AsciEng. *Euo*prom activity was detected at ~12 hpi followed by the activity of the *porB*prom at ~20
hpi followed by *hctB*prom activity starting at ~26 hpi. Taken together these data suggest that *porB*, as
predicted by the RNA-seq clustering experiment, can be considered an IB gene.
314
Predicted cell type expression of T3SS genes. We noticed an intriguing expression pattern with the type III
secretion system (T3SS) structural genes in the clustering data that suggested cell form specific expression.
To explore this observation, we plotted the effects of ectopic expression on the T3SS operons. We used our wt
RNA-seq data [30], operon prediction software [31] and the RT-PCR data published by Hefty et. [32] to annotate the
T3SS operons (Table S2) and plotted the expression data using volcano plots. These plots showed that the
majority of the T3SS operons were regulated in an IB like pattern, i.e. up-regulated between 18 hpi and 24 hpi,
repressed by Euo and HctB ectopic expression, but not induced by CtcB or HctA ectopic expression (Fig. 6).
The exception to this pattern were the two operons for the T3SS translocons, (*CTL0238, lcrH, copB_2,*
copD_2) and (*scc2, CTL0840, copB, copD*). The *CTL0238* operon (*CTL0238*-op) was regulated like an RB
gene while the *scc2* operon (*scc2*-op) was regulated like an EB gene (Fig. 6).
325
Cell type expression of T3SS genes by FISH. We next investigated the cell type expression of two of the
T3SS operons predicted to be expressed in the IB, the *sctU* operon and the *sctL* operon. The *sctU* operon

(*sctU*-op) encodes the genes *sctU*, *sctV*, *lcrD*, *copN*, *scc1*, and *malQ*, while the *sctL* operon (*sctL*-op) includes
the genes *sctL*, *sctR*, *sctS*, and *sctT* We used custom FISH probes encompassing *sctU* through *lcrD* for the
detection of the *sctU-op* mRNA and *sctL* to *sctR* for the detection of *sctL*-op mRNA. Cell monolayers were
infected with L2-AsciEng and L2-BsciEng at an moi ~ 0.3 and processed for FISH staining at 16 and 24 hpi
(Fig. 7 and Fig. S2). FISH signal was not observed in the RB cells (*euo*prom+) at 16 hpi for either *sctU*-op (Fig.
7A) or *sctL*-op (Fig. S2A). In the infections fixed at 24 hpi, the FISH staining for both operons was observed in
cells distinct from *euo*prom+ and *hct*Bprom+ cells (Fig. 7B and Fig. S2B). However, both T3SS operon mRNAs
were detected in a subset of the *hctA*prom+ cell population (Fig. 7B and Fig. S2B). We next used our
TrackMate workflow to quantitate these data. Cells were identified by their promoter reporter signal (green) and
separately by their mRNA fluorescence signal (magenta). The fluorescence intensity was measured and
plotted for both channels (FISH (magenta) and promoter reporter (green)) in both identified populations as
described in Fig. 4.

Expression in RB cells. We identified the mRNA+ cell population (magenta) and quantified both the
*euo*prom signal intensity and the FISH signal intensity. For both the *sctU*-op and the *sctL*-op there was
primarily a double positive population (high mRNA signal and high *euo*prom signal). We also quantified the
mRNA expression in RBs by identifying the *euo*prom+ cells (green) and measuring the *sctU*-op (Fig. 7C) and
sctL-op (Fig. S2C) FISH signal and plotted this against the *euo*prom signal intensity. The *euo*prom+ population
(green) were both single and double positive for both operons (Fig. 7C and Fig. S2C, *euo*p). These data
suggest that neither operon was expressed in RBs. As there was no mRNA signal in the 16 hpi RB cells, it is
likely that the mRNA+ double positive cell population at 24 hpi were IBs with carryover Neongreen signal, while
the *euo*prom+ single positive population were the RBs.

Expression in IB cells. The mRNA+ cell population (magenta) was identified and both the *hctA*prom
signal intensity and the FISH signal intensity was quantified and plotted. For both the s*ctU*-op (Fig. 7C) and
sctL-op (Fig. S2C) mRNA+ cells (magenta), the majority were double positive for mRNA signal and *hctA*prom
signal. For the *hctA*prom+ cell population (green) there was both a single (*hctA*prom) and double positive
(*hctA*prom and mRNA) population (Fig. 7C and Fig. S2C, *hctA*p). These data further suggest that the *sctU*-op
and *sctL*-op were expressed in the IB cell type. It's likely that the *hctA*prom+ single positive population are late
IB/EB cell forms that are becoming EBs and have repressed *sctU*-op and *sctL*-op expression.

Expression in EB cells. In contrast, the mRNA+ cell population for both the *sctU*-op and the *sctL*-op in
the hctBprom expressing cells were distinct single positive (mRNA signal) populations (Fig. 7C and Fig. S2C,
*hctB*p). Additionally, the *hctB*prom+ cell population was also primarily single positive (*hctB*prom). These data
suggest that the sctU-op and the sctL-op were not expressed in the EB cell forms. Combined, these overall
expression patterns of the *sctU* and *sctL* operons were very similar to that of the *hctA* mRNA and *porB* mRNA
FISH, supporting an IB like gene expression pattern.

Cell type expression of the T3SS translocon genes by FISH. As mentioned above, the *Ctr* genome
encodes two operons for the T3SS translocon each of which contain four genes, the *CTL0238*-op and the
scc2-op [33]. This duplication is conserved in all the vertebrate infecting chlamydial species. The expression
profiles from our clustering data and volcano plots suggested that the two translocon operons are expressed in
different cell types; the *CTL0238*-op in RBs and the *scc2*-op in EBs (Fig. 6). To verify differential cell type
expression, host cells were infected with L2-BsciEng, fixed at 24 hpi and probed with custom FISH probes
designed against C*TL0238*-op and *scc2*-op. These infected and FISH stained cells were imaged by confocal
microscopy. The confocal micrographs showed that the mRNA FISH signal for *CTL0238*-op heavily overlapped
with the *euo*prom channel but was distinct from *hctB*prom+ cells (Fig. 8A, *CTL0238*-op mRNA). In contrast, the
mRNA signal for the *scc2*-op was distinct from the *euo*prom+ cells but almost completely overlapped the
*hctB*prom+ cells (Fig. 8B, *scc2*-op mRNA).

We again quantified this expression pattern using our TrackMate workflow. Chlamydial cells were
identified by their promoter reporter signal (green) and separately by their mRNA fluorescence signal
(magenta). The fluorescence intensity was measured and plotted for both channels, FISH (magenta) and
promoter reporter (green).

Expression in RB cells. We identified the CTL0238-op mRNA+ cell population (magenta) and
quantified both the euoprom signal intensity and the FISH signal intensity. For the CTL0238-op mRNA+ cells,
there was primarily a double positive population (high mRNA signal and high euoprom signal) (Fig. 8A). We
also quantified the mRNA expression in RB cells (*euo*prom+ cells, green) The CTL0238-op FISH signal was
plotted against the *euo*prom signal intensity and the euoprom+ cells (green) were mostly double positive with
an additional single positive population (Fig. 8A). This single positive (euoprom+, CTL0238-op mRNA-)
population is likely due to the long half life of the GFP protein.

Expression in EB cells. In contrast, the CTL0238-op mRNA+ cell population (megenta) when plotted
for mRNA signal and the hctBprom signal was a distinct single positive population (CTL0238-op mRNA+,
hctBprom-) (Fig. 8A). We also identified the hctBprom+ cell population (green) and plotted the mRNA signal
and hctBprom signal. This population was also primarily single positive (hctBprom+, CTL0238-op mRNA-).

For the scc2-op mRNA FISH quantification the results were opposite. The scc2-op+ cells (megenta)
were primarily single positive when plotted against the euoprom signal and double positive when plotted
against the hctBprom signal (Fig. 8B). While the euoprom+ cell population (green) was also primarily single
positive and the hctBprom+ cells (green) were primarily double positive (Fig. 8B).

To further highlight the differential expression of *CTL0238*-op mRNA and *scc2*-op mRNA, we infected
cells with L2 AsciEng and processed the samples for FISH at 16 hpi when most of the chlamydial cells are
RBs. At 16 hpi essentially all the cells were green RBs (*euo*prom+) with little to no red IBs (*hctA*prom+) cells.
The green *euo*prom+ cells were all positive for the *CTL0238*-op FISH signal (Fig. 8C, *CTL0238*-op). In
contrast, the *scc2*-op FISH signal was undetectable in *euo*prom+ cells (Fig. 8D, *scc2*-op). Taken together,
these data support the observation that the two translocon operons are differentially regulated and are
expressed in distinct cell forms. The *scc2*-op is expressed in late IB/EB cells while the *CTL0238*-op is
expressed in RB cells.

Predicted cell type expression of T3SS effectors. In general, T3SS translocons are involved in interacting
with host membranes to facilitate the secretion of T3SS effectors into target cells [34]. During infection, *Ctr*
secretes effectors into/through two membrane systems, the host cell plasma membrane and, once inside the
cell, the inclusion membrane. Additionally, the chlamydial T3SS is known to secrete different kinds of effectors,
soluble proteins as well as the integral membrane inclusion (Inc) proteins [35–38]. Using volcano plots we asked
how the genes encoding the soluble effector proteins (Table S3) and Inc proteins (Table S4) were regulated by
the ectopic expression of Euo, HctA, HctB and CtcB. The vast majority of the soluble T3SS effector genes
were regulated like EB genes; higher in 18-24 hpi and induced by HctA and CtcB ectopic expression, but down
regulated by Euo and HctB ectopic expression (Fig. 9A). In contrast, most of the *incs* were regulated like RB
genes with the exception of *incM* and *incV,* which were regulated like EB genes (Table S4 and Fig. 9B).

Cell type expression of *inc* genes by FISH. The volcano plots suggested that the majority of the *inc* effector
genes were expressed in an RB like pattern. However, two Inc genes (*incV* and *incM)* stood out as potential
EB genes (Fig. 9B). To determine if the putative late Inc, *incV,* was expressed late in RBs or was a bona fide
EB gene, we compared mRNA expression of a known RB expressed Inc, *incD* to *incV* using FISH. Cells
infected with L2-BsciEng were probed for the expression of *incD* and *incV* at 16 hpi (mostly RBs) and 24 hpi
(all three cell forms). Confocal microscopy revealed that, as expected, *incD* was expressed in *euo*prom+ RB
cells at both 16 hpi and 24 hpi and not in *hctB*prom+ EBs present at 24 hpi (Fig. 10A). Conversely, *incV* mRNA
could not be detected at 16 hpi and was expressed exclusively in *hctB*prom+ EBs (Fig. 10B).

Again, we quantified expression of each gene in RBs and EBs from inclusions from the 24 hpi
experiments using our TrackMate workflow. Chlamydial cells were identified by their promoter reporter signal
(green) and separately by their mRNA fluorescence signal (magenta) and the signal for both populations was
plotted.
Expression in RB cells. We identified the incD mRNA+ cell population (magenta) and plotted both the
euoprom signal intensity and the FISH signal intensity (Fig. 10C). For the incD mRNA+ cells there was
primarily a double positive population (high incD mRNA signal and high euoprom signal). We also quantified
the mRNA expression in RB cells (*euo*prom+ cells, green) The incD FISH signal was plotted against the
euoprom signal intensity and the euoprom+ cells (green) were mostly double positive with an additional small
single positive population. This single positive (*euo*prom+, incD mRNA-) population is likely due to the long half
life of the GFP protein allowing it to also label early IB cells forms.
Expression in EB cells. In contrast, the incD mRNA+ cell population (magenta) when plotted for
mRNA signal and the hctBprom signal was a distinct single positive population (incD mRNA+, hctBprom-) (Fig.
10C). We also identified the hctBprom+ cell population (green) and plotted the mRNA signal and hctBprom
signal. This population was also primarily single positive (hctBprom+, incD mRNA-).
For the incV FISH signal the results were opposite. The incV+ cells (magenta) were primarily single
positive when plotted against the euoprom signal and double positive when plotted against the hctBprom signal
(Fig. 10D). While the euoprom+ cell population (green) was also primarily single positive and the hctBprom+
cells (green) were primarily double positive (Fig. 10D). These data support the hypothesis that incD is indeed
an RB gene and that incV, and likely incM, are EB genes.
441
Discussion:
The chlamydial developmental cycle has traditionally been defined by the timeline of the infection. The
infectious EB invades the host cell and begins EB to RB differentiation into the RB cell form that then begins to
divide. The genes involved in this process have been described as the early genes. Sometime after EB to RB
differentiation and the initiation of replication transitionary forms appear. The gene expression associated with
this timeframe is usually considered the chlamydial midcycle. At ~24 hpi until cell lysis EBs accumulate in the
inclusion. Genes upregulated at this time point and forward are considered late genes [14–16]. We have dissected
the developmental cycle and developed a model based on cell type transitions [9]. Our model suggests that the
developmental cycle is best described by a programmed cell production model [9]. In this model, the EB enters
the host cell (through the use of premade effectors) and initiates immediate early protein synthesis (EB to RB
differentiation genes) to begin the EB to RB differentiation process. The EB to RB differentiation process takes
~ 10 hours to complete. The completion of EB to RB differentiation is defined by the first division of the nascent
cell resulting in RB cells. At this stage the RBs expand in number through cell division, amplifying the infection.
Our model suggests the RBs mature during this amplification stage, ultimately producing daughter cells with
asymmetric fates. One daughter cell becomes the IB cell form while the other remains an RB. Our model
defines the IB cell form as the committed cell to EB formation. The mature RBs at this stage continue to
replicate producing one IB and one RB. The IBs never re-enter the cell cycle and instead transition into the
infectious EB form which takes ~ 10 hours to complete [9].
In this study, we ectopically expressed four transcriptional regulatory proteins that all blocked the
progression of the developmental cycle. The effects of expression of these regulatory proteins using RNA-seq
was determined and compared using a clustering algorithm which resulted in three distinct regulation patterns.
The first cluster contained genes that were unaffected by the ectopic expression of Euo and were not
upregulated between 18 hpi and 24 hpi of a *Ctr* L2 infection. The second cluster consisted of genes whose
expression increased from 18 hpi to 24 hpi of a wt infection and were not induced by ectopic expression of
HctA and CtcB. The third cluster of genes were upregulated between 18 hpi and 24 hpi and by the ectopic
expression of both HctA and CtcB. These groups fit well into the major cell categories in our model; RBs, IBs
and EBs. Using the clustering observation, we created selection criteria based on changes in gene expression

from our RNA-seq experiments. We were able to bin 639 out of 902 genes (~70%) into one of the RB, EB, and
IB categories. The genes that could not be assigned were either expressed at levels too low to have
confidence in the expression pattern, or had a unique expression pattern that did not fit into the three
categories suggesting potential unique roles in chlamydial biology. This study focused on determining gene
expression through measuring mRNA and it remains to be determined if any of these genes are translationally
regulated as well.

The RB cell is the replicating cell form leading to expansion of cell numbers. Based on the changes in
gene expression after ectopic expression of Euo, HctA, CtcB and HctB we found that 532 genes were
regulated as RB genes. This category included cell replication genes, genes involved in protein synthesis,
genes for many of the Inc proteins and the *euo* gene. Based on our selection criteria this group likely
encompasses both potential constitutive genes (expressed in RBs, IBs and potentially early EBs) as well as
RB specific genes such as *euo*, *incD* and the *CTL0238*-op which were expressed only in the RB cell form.

The IB cell type is the transitional form between the RB and the EB and is currently poorly defined. We
define the IB cell type as the committed step to EB formation; the IB is the cell form that exits the cell cycle and
begins the program to transition into the infectious EB [9]. Our data identified 67 genes that are likely expressed
specifically in the IB cell type. The functions of these genes vary widely. We identified two porin genes (*porB*
and *CLT0626*), two disulfide isomerases (*CTL0149* and *CTL0152*) and six polymorphic outer membrane
proteins (*pmpB*, *C, E*, *F*, *G*, and *H*) as IB genes suggesting dramatic changes to the outer membrane of the IB
as it transitions into the EB.

The EB cell is the infectious cell form that is "terminally" differentiated. Once formed in the inclusion, the
EB maintains an infectious phenotype through active metabolism but has very low levels of protein expression
as shown by click chemistry and the EBs ability to maintain infectivity in the presence of protein synthesis
inhibitors while inside the inclusion, and by the requirement of nutrients for the EB to maintain infectivity in the
presence of protein synthesis inhibitors after cell exit [30]. Here, we define the EB regulon as the genes
expressed during the late IB to EB maturation phase resulting in the infectious form. Of the 46 EB genes, 18
had been previously shown to be directly regulated by the sigma54 alternative sigma factor and 4 were
reported to be sigma28 regulated genes [24,27]. The regulation of the remaining 24 genes is unknown. As both
HctA ectopic expression and the ectopic expression of CtcB induce the expression of the EB genes, the EB
regulon is likely regulated by a complex shift in gene expression and activation of the sigma54 and sigma28
regulons is a part of this shift.

We tested one of the predicted IB genes, *porB*, and showed that its regulation, both by promoter
specific gene expression in chlamydial cells and by its developmental kinetics, matched that of *hctA.* This was
further confirmed using FISH to demonstrate cell type gene expression matched that of the IB gene, *hctA*. We
have previously published the kinetics of the *euo*, *hctA* and *hctB* promoters and showed that the promoter
activities fit into the RB, IB and EB model [8,9]. Here we combined these promoter reporter strains with FISH and
demonstrated that the *euo* mRNA was expressed primarily in RBs, that *hctA* mRNA was expressed in IBs, and
that *hctB* mRNA was expressed in EBs demonstrating the usefulness of FISH for identifying cell type specific
gene expression.

Overall, these data support a model that includes (at least) three dominant cell forms: the RB, the IB
and the EB. These cells have dramatically different gene expression profiles and phenotypes. The EB has
been well characterized as it is the infectious form, does not replicate and has a dramatically condensed
nucleoid. The nucleoid structure is due in part to the binding of the two histone like proteins, HctA and HctB to
the chromosome [19,22]. Our data indicates that the construction of the compact nucleoid occurs in two distinct
and temporally separated steps [8,9]. HctA is expressed as an IB gene and, when ectopically expressed, resulted
in the expression of the EB genes suggesting that HctA expression is an important regulator of the IB to EB
transition. HctB on the other hand is expressed as an EB gene and, when ectopically expressed, resulted in
the inhibition of the expression of most genes with the exception of the ribosomal protein genes. An intriguing

hypothesis is that the ribosomal protein genes are potentially free of inhibition and that protein synthesis could
be rapidly reinitiated upon EB to RB differentiation, before complete removal of HctA and HctB from the
chromosome. Taken together, these data suggest that the transition from the IB to EB occurs in two steps; 1)
HctA chromosomal binding turns off RB and IB genes, allowing EB genes to become expressed, and 2) HctB
is expressed late in EB formation creating the final condensed nucleoid that potentially spares the ribosomal
genes.

Volcano plots of the effects of ectopic expression of the four regulatory genes support the
categorization of most chlamydial genes into the RB, IB and EB categories. We specifically focused on the
expression of the T3SS operons and observed that the majority of the operons for the structural components
were IB-like in their regulation. This was verified using FISH for both the *sctL* operon (*sctL, sctR, sctS, and*
sctT) and the *sctU* operon (*sctU, sctV, lcrD, copN, scc1, and malQ*). While the majority of the T3SS structural
operons were expressed as IB genes, the two translocon operons (*CTL0238, lcrH, copB_2, copD_2*) and
(s*cc2, CTL0840, copB, copD*) were predicted by clustering and volcano plots to be expressed in RB and EB
cells respectively. This prediction was again verified by FISH in the context of dual promoter reporter strains.

The observation that the two translocons were expressed in distinct cell forms (*CTL0238*-op in RBs and
scc2-op in EBs) prompted us to determine the expression of the T3SS effectors. *Ctr* encodes two classes of
effectors, soluble and inclusion membrane embedded proteins (Incs) [36,39,40]. The data from this study showed
that the majority of the Inc protein effectors (28 out of 36) were expressed as RB genes while the majority of
the soluble T3SS effectors (17 out of 23) were expressed as EB genes and that none of the soluble effectors
were expressed as RB genes. This pattern suggests an intriguing hypothesis; the *scc2*-op translocon
translocates soluble effectors as the EB contacts host cells and mediates entry events, while the *CTL0238*-op
is expressed early during the EB to RB differentiation process in the nascent inclusion and translocates the
transmembrane Inc effectors. Whether this separation is temporal or whether the two translocons are
specialized for the translocation of soluble vs. inclusion membrane effectors is currently unknown. Interestingly,
although the majority of the Inc proteins were expressed as RB genes there were two Incs that were
determined to be expressed in EBs, *incV* and *incM*. In addition to its regulation pattern, we also verified that
incV was an EB gene using FISH. Both IncV and IncM are involved in the establishment of early inclusion
functions [41,42]. We hypothesize that these "pre-loaded" Inc proteins are among the first to be secreted from
internalized *Ctr* after the *CTL0238*-op is deployed.

Ctr communicates and reprograms the host cell to create and maintain its intracellular replication niche
in part through the use of the T3SS. We were surprised that the majority of the T3SS operons for the structural
components of the system were expressed as IB genes. This expression pattern along with the cell type
specific expression of the translocons (one in the RB and one in the EB) and effectors suggests that the T3SS
is constructed, deployed and secretes effectors in a cell type specific manner that is likely a critical component
of the complex developmental cycle and host cell reprogramming.

Taken together, these data support our overall model that the chlamydial developmental cycle should
be viewed as consisting of discrete phenotypic cell forms that are produced by an underlying developmental
program consisting of RBs that amplify the infection, RBs that produce IBs and IBs that transition into the
infectious EB [8,9] (Fig. 11). This process creates an infection with mixed phenotypic forms throughout the
infection. The data also suggest that each cell type has a distinct gene expression profile (RB, IB and EB) and
that the relative numbers of each of these cell forms creates the time based expression patterns reported in
population based studies [13–16]. Our model suggests that the EB binds to and enters cells in part through the
deployment of soluble effectors and the *scc2*-op translocon expressed during EB development. After entry, EB
to RB differentiation begins and the RB genes are expressed including the *CTL0238*-op translocon which
deploys the Inc proteins for the creation of the inclusion replication niche and the genes required for chlamydial
replication leading to RB amplification. After an amplification period, the RB matures into a stem cell-like cell
form and begins to produce IBs [9]. The IB synthesizes the T3SS structural components and matures into the

EB form over ~8-10 hours. That the IB and not the RB expresses the genes for the construction of the T3SS
suggests that the T3SS apparatus that entered on the EB cells remains on the RBs and is diluted with every
replication. It is unclear if the secretion system gets partitioned equally or is retained in a subset of RBs.
Intriguingly, this supports a proposed role of T3SS dilution in cell form maturation/development put forth
previously [37,43,44].
The IB also expresses the histone like DNA binding protein, HctA. Previous studies have shown that
when expressed in *E. coli*, HctA can alter gene expression in a gene specific manner [19]. Our data suggest that
HctA has an important role in shifting gene expression from the IB pattern to the EB genes. This is likely in
conjunction with the CtcB/C two component regulatory system and the sigma54 alternative sigma factor [24,27].
The EB genes as previously mentioned include the majority of the soluble T3SS effectors and the *scc2*-op
translocon as well as the HctB DNA binding protein. We hypothesize that EB gene expression loads the EB
with the invasion-related proteins and HctB shuts down the majority of gene expression creating the dense
nucleoid, the final step of EB formation. This prepares the EB for the initiation of the next round of infection
(Fig. 11). Our data has highlighted three categories of gene expression that define the three major phenotypic
cell forms, RB, IB and EB. However, future studies are needed to define the regulatory circuits and DNA
elements that create these cell form expression patterns.
DNA replication is tightly controlled during the *Ctr* developmental cycle; only the RB cell form replicates
the chromosome and the IB and EB cells contain a single fully replicated chromosome [45]. The role of the
control of DNA replication in regulating gene expression is currently unknown. However, it is intriguing to
speculate that DNA replication could contribute to changes in DNA supercoiling which has been shown to play
a role in gene expression during the chlamydial developmental cycle [46–48]. Our identification of cell type gene
expression of a large percentage of the chlamydial genome will aid the determination of the functions of the
many hypothetical genes encoded in the chlamydial genome. Understanding the function of many of these
genes has been hampered by the mixed cell environment of the chlamydial inclusion. Additionally, with the
emerging genetic tools available to investigate the functional roles of genes during infection, knowing in which
cell type they are expressed will improve the interpretation of the results.
589
590   **Materials and Methods**
591
Cell Culture
Cell lines were obtained from the American Type Culture Collection. Cos-7 cells (CRL-1651) were grown in
RPMI-1640, supplemented with 10% FBS and 10 µg/mL gentamicin (Cellgro). *Chlamydia trachomatis* serovar
L2 (LGV Bu434) was grown in Cos-7 cells. Elementary Bodies (EBs) were purified by density gradient (DG)
centrifugation essentially as described [49] following 48 h of infection. EBs were stored at -80ºC in Sucrose
Phosphate Glutamate (SPG) buffer (10 mM sodium phosphate [8mM K2HPO4, 2mM KH2PO4], 220 mM
sucrose, 0.50 mM l-glutamic acid, pH 7.4) until use.
599
Vector Construction
All constructs used p2TK2-SW2 [50] as the backbone and cloning was performed using the In-fusion HD EcoDry
Cloning kit (FisherScientific). Primers and geneblocks (gBlocks) were ordered from Integrated DNA
Technologies (IDT) and are noted in Table S5. The construction and characterization of p2TK2-SW2-T5-E-
clover-3xFLAG (E-clover), p2TK2-SW2-Tet-J-E-clover-3xFLAG (Tet-JE-clover) and p2TK2-SW2-Tet-J-E-hctB-
3xFLAG (Tet-JE-hctB) has been described previously [25].
E-hctA-3xFLAG (E-hctA), E-euo-3xFLAG, (E-euo) and E-ctcB-3xFLAG (E-ctcB),
The *hctA*, *euo* and *ctcB* ORFs were amplified from *Ctr* L2(434) using the primers indicated in Table S5.
Each fragment was used to replace Clover in p2TK2-SW2-T5-E-clover-3xFLAG.

*hctB*prom_Scarlet-*euo*prom_neongreen (BsciEng), *hctA*prom_Scarlet-*euo*prom_neongreen (AsciEng) and
*porB*prom_Scarlet-*euo*prom_neongreen (PsciEng).

The gBlock mScartlet-I (Table S5) was cloned into BMELVA [9] to replace mKate which created the
construct BscieLVA. The degradation tag LVA was then removed from neongreen using the primers indicated
to create BsciEng. The *hcA* and *porB* promoters were amplified and used to replace the *hctB* promoter using
the primers indicated to create AsciEng and PsciEng respectively.

Chlamydial Transformation and Isolation.
Transformation of Ctr L2 was performed essentially as previously described [8]. Briefly, 1×10^8 EBs + >2µg
DNA/well were used to infect a 6 well plate. Transformants were selected over successive passages with
1U/ml penicillin G or 500µg/ml spectinomycin as appropriate for each plasmid. The new strain was clonally
isolated via successive rounds of inclusion isolation (MOI, <1) using a micromanipulator. Clonality of each
strain was confirmed by isolating the plasmid, transforming into *E. coli* and sequencing six transformants.

Replating Assay.
Ctr were isolated by scraping the infected monolayer into media and pelleting at 17200 rcfs. The EB pellets
were resuspended in RPMI via sonication and seeded onto fresh monolayers in a 96-well microplate in a 2-fold
dilution series. Infected plates were incubated for 24 hours prior to fixation with methanol and stained with 4′,6-
diamidino-2-phenylindole (DAPI) and *Ctr* MOMP Polyclonal Antibody, FITC (Fishersci). The DAPI stain was
used for automated microscope focus and visualization of host-cell nuclei, and the anti-*Ctr* antibody was used
to visualize the *Ctr* to identify and count inclusions. Inclusions were imaged using a Nikon Eclipse TE300
inverted microscope utilizing a scopeLED lamp at 470nm and 390nm, and BrightLine band pass emissions
filters at 514/30nm and 434/17nm. Image acquisition was performed using an Andor Zyla sCMOS in
conjunction with µManager software. Images were analyzed using ImageJ software and custom scripts.
Statistical comparisons between treatments were performed using an ANOVA test followed by Tukey's Honest
Significant Difference test.

Transmission Electron Microscopy
For analysis of the structure of *Ctr* upon ectopic protein expression, cell monolayers were infected with the
indicated strain at an moi of 0.5 and induced with 0.5mM Theophylline at 15 hpi. Infected cells were released
from the plate with Trypsin-EDTA at 30 hpi, rinsed with 1xPBS and the pellet was fixed with EM fixative
(2%PFA, 2% Glutaraldehyde, 0.1M Phosphate Buffer, pH 7.2) overnight at 4ºC. Fixed pellets were rinsed and
dehydrated before embedding with Spurr's resin and cross sectioned with an ultramicrotome (Riechert Ultracut
R; Leica). Ultra-thin sections were placed on formvar coated slot grids and stained with uranyl acetate and
Reynolds lead citrate. TEM imaging was conducted with a Tecnai G2 transmission electron microscope (FEI
Company; Hillsboro, OR).

RNA-Seq
Expression of each of the ectopic proteins was induced at 15 hpi with either 0.5 mM theophylline and 30ng/ml
anhydrotetracycline (HctB) or 0.5 mM theophylline (Clover, HctA, CtcB and Euo). Total RNA was isolated from
the infected cells for each treatment from triplicate experiments. Briefly, the infected monolayer was scraped
into ice cold PBS, lysed using a Dounce homogenizer and the *Ctr* enriched over a 30% MD-76R pad. Total
RNA was isolated using TRIzol reagent (Life Technologies) following the protocol provided, and genomic DNA
was removed (TURBO DNA-free Kit, Invitrogen). Both prokaryotic and eukaryotic rRNAs were depleted using
Illumina Ribo-Zero Plus. The enriched RNA samples were quantified and the libraries built and barcoded by
the IBEST Genomics Resources Core at the University of Idaho. The libraries were sequenced by University of
Oregon sequencing core using the Illumina NovaSeq platform. The chlamydial specific reads were analyzed by

aligning to the published *Ctr* L2 Bu 434 genome using the Bowtie2 aligner software [51]. The aligned chlamydial
reads were quantified for each chlamydial ORF using HT-seq. Statistical analysis and normalization of read
counts was accomplished using DESeq2 in R [52]. Log2fold change and statistics were also calculated using
DESeq2. Heatmaps and hierarchical clustering were generated and visualized using Python with Pandas and
the Seaborn visualization package [26]. The raw reads and HT-seq counts are accessible from the NCBI's Gene
Expression Omnibus with the accession number of GSE262652 and GSE276506. Volcano plots were
constructed from the log2fold change data using Python and the Bokeh plotting library (Bokeh Development
Team).
664
Total RNA was isolated from cells infected with the indicated strains. Expression of each protein was induced
at 15 hpi with either 0.5 mM theophylline and 30ng/ml anhydrotetracycline (HctB) or 0.5 mM theophylline
(Clover, HctA, CtcB and Euo) and the *Ctr* isolated at 18 and 24 hpi on ice. Briefly, the infected monolayer was
scraped into ice cold PBS, lysed using a Dounce homogenizer and the *Ctr* isolated over a 30% MD-76R pad.
Total RNA was isolated using TRIzol reagent (Life Technologies) following the protocol provided and genomic
DNA removed (TURBO DNA-free Kit, Invitrogen). The enriched RNA samples were quantified and the libraries
built and barcoded by the IBEST Genomics Resources Core at the University of Idaho. The libraries were
sequenced by University of Oregon sequencing core using the Illumina NovaSeq platform. RNA-seq reads
were aligned to the published *Ctr* L2 Bu 434 genome using the Bowtie2 aligner software [51]. Reads were
quantified using HTseq . Statistical analysis and normalization of read counts was accomplished using
DESeq2 in R [52]. Log2fold change and statistics were also calculated using DESeq2. Heatmaps and
hierarchical clustering were generated and visualized using Python with Pandas and the Seaborn visualization
package [30]. Aligned reads are accessible from the NCBI's Gene Expression Omnibus with the accession
number of GSE262652. Volcano plots were constructed from the log2fold change data using Python and the
Bokeh plotting library (Bokeh Development Team).
680
RNA fluorescence in situ hybridization (RNA-FISH)
All FISH probes were designed by Molecular Instruments (Los Angeles, CA) using the sequence indicated in
(Tabel S6). Cos7 monolayers seeded on coverslips were infected with the indicated strains at an MOI ~ 0.3.
Infected cells were fixed at the indicated times in 4% paraformaldehyde (PFA) for 10 min at RT at 24 hpi,
washed 2x with 1XPBS and dehydrated overnight at -20ºC in 70% EtOH. Samples were probed and the signal
was amplified as described by the protocol provided by Molecular Instruments with the exception that DAPI
was added to the final wash to visualize DNA. Coverslips were mounted on a microscope slide with MOWIOL®
mounting solution (100 mg/mL MOWIOL® 4-88, 25% glycerol, 0.1 M Tris pH 8.5).
Fluorescence images were acquired using a Nikon spinning disk confocal system with a 60x oil-
immersion objective, equipped with an Andor Ixon EMCCD camera, under the control of the Nikon elements
software. Images were processed using the image analysis software ImageJ (http://rsb.info.nih.gov/ij/).
Representative confocal micrographs displayed in the figures are maximal intensity projections of the 3D data
sets, unless otherwise noted.
694
Live cell imaging
Monolayers of Cos7 cells were grown in a glass bottom 24 well plate and infected with the promoter reporter
strains L2-BsciEng, L2-AsciEng and L2-PsciEng. Live cell imaging of the developing inclusions was started at
8 hpi using an automated Nikon epifluorescent microscope equipped with an Okolab (http://www.oko-
lab.com/live-cell-imaging) temperature controlled stage and an Andor Zyla sCMOS camera
(http://www.andor.com). Multiple fields of view from each well were imaged every fifteen minutes. The
fluorescence intensity of each inclusion over time was tracked using the ImageJ plugin TrackMate [28] and the
results were averaged and plotted using python and matplotlib [53].

Figure 1. Ectopic expression of Euo, HctA, CtcB and HctB resulted in inhibition of the developmental cycle. **(**A) Cos-7 cells were infected with the four strains and a Clover control strain and induced for ectopic expression at 15 hpi. EBs were harvested at 48 hpi. IFU production was dramatically reduced by the ectopic expression of Euo, HctA, CtcB and HctB but not by the expression of the Clover protein. * = P < 0.01. (B) Transmission EM of Cos-7 cells infected with *Ctr* expressing Clover, Euo, HctB, HctA or CtcB. Ectopic expression was induced at 15 hpi and the cells were fixed and stained at 30 hpi. The bacteria in the Clover induced and uninduced chlamydial infections looked similar with inclusions of both samples containing large RB like cells as well as electron dense EB like cells. The *Ctr* in the Euo expressing inclusions were primarily RB like cells while very few cells were electron dense EB cells. The chlamydial cells in the HctB expressing inclusions were abnormal looking, some with apparent condensed nucleoids. The HctA expressing *Ctr* also appeared abnormal with condensed nucleoids. Many of the CtcB expressing *Ctr* cells were target-like RB sized cells with a condensed nucleoid.

Figure 2. RNA-seq analysis of *Ctr* ectopically expressing Euo, HctA, HctB, or CtcB. **(**A) For each of the induced samples (n=3) the RNA-seq PCA profiles clustered within the same ectopic expression group but each group had a distinct profile as visualized by plotting the first and second principal component. (B and C) The cluster plots revealed two distinct late gene regulation groups. (B) The IB gene cluster was defined as those that were upregulated between wt *Ctr* infections at 18 hpi and 24 hpi but were not upregulated by the ectopic expression of HctA and CtcB as compared to the Clover control. (C) The EB gene cluster was defined as genes that were upregulated between wt *Ctr* infections at 18 hpi and 24 hpi and were induced by ectopic expression of HctA and CtcB as compared to the Clover control.

Figure 3. Effects of ectopic expression of Euo, HctA, CtcB and HctB on the gene expression of every *Ctr* gene. RNA-seq differential expression was determined for each gene using DESeq2 for native infection at 18 hpi vs 24 hpi, Euo-FLAG expression, HctA-FLAG expression, CtcB-FLAG expression and HctB-FLAG expression vs the control Clover-FLAG. Volcano plots show that the expression of RB designated genes (orange) were largely unchanged from 18 hpi to 24 hpi while IB (green) and EB (red) designated genes were dramatically up regulated. For Euo-FLAG expression the RB genes (orange) were largely unchanged while IB genes (green) and EB genes (red) were all down regulated. Ectopic expression of HctA-FLAG resulted in the repression of many of the RB genes (orange) and up regulation of the EB genes (red) but had little impact on the expression of the IB genes (green). CtcB-FLAG expression had very little effect on RB genes (orange) but dramatically upregulated EB genes (red). The ectopic expression of HctB-FLAG resulted in the down regulation of both IB (green) and EB (red) genes but up-regulated many RB genes (orange). Additionally, HctB-FLAG expression resulted in the upregulation of many of the ribosomal protein genes (purple).

Figure 4. Cell type expression of representative genes from the three gene categories (RB, IB and EB) correspond to the three chlamydial cell forms. Cos-7 cells infected with L2-AsciEng or L2-BsciEng, fixed at 24 hpi and stained using FISH probes for *euo* mRNA, *hctA* mRNA and *hctB* mRNA. (A) Z-projection confocal micrographs showing *euo* mRNA localization in comparison to *euo*prom, *hctA*prom and *hctB*prom activity. Individual chlamydial cells with *euo* mRNA signal from 5 inclusions were identified using TrackMate and the fluorescence intensity for each channel (mRNA and promoter reporter) was plotted (magenta dots). Individual chlamydial cells positive for *euo*prom, *hctA*prom or *hctB*prom signal from 5 inclusions were also identified using TrackMate and their expression intensity for each channel was plotted (green dots). (B) Z-projection confocal micrographs showing *hctA* mRNA localization in comparison to *euo*prom, *hctA*prom and *hctB*prom activity. Individual chlamydial cells with *hctA* mRNA signal from 5 inclusions were identified using TrackMate and the fluorescence intensity for each channel (mRNA and promoter reporter) was plotted (magenta dots). Individual

chlamydial cells positive for *euo*prom, *hctA*prom or *hctB*prom signal from 5 inclusions were also identified using
TrackMate and their expression intensity for each channel was plotted (green dots). (C) Z-projection confocal
micrographs showing *hctB* mRNA localization in comparison to *euo*prom, *hctA*prom and *hctB*prom activity.
Individual chlamydial cells with *hctB* mRNA signal were identified from 5 inclusions using TrackMate and the
fluorescence intensity for each channel (mRNA and promoter reporter) was plotted (magenta dots). Individual
chlamydial cells positive for *euo*prom, *hctA*prom or *hctB*prom signal from 5 inclusions were also identified using
TrackMate and their expression intensity for each channel was plotted (green dots). Size bar = 5µm.
757
Figure 5. PorB gene expression is consistent with being classified as an IB gene. **(**A) FISH analysis of *porB*
mRNA expression in comparison to *euo*prom, *hctA*prom and *hctB*prom activity at 24 hpi. Confocal micrographs
of L2-AsciEng and L2-BsciEng infected cells probed for *porB* mRNA expression from 5 inclusions using
Molecular instruments FISH probes. TrackMate was used to identify the *porB* mRNA+ cell population and
measure the FISH fluorescent signal as well as the *euo*prom, *hctA*prom and *hctB*prom fluorescent signal. The
intensity for both channels for each cell was plotted (magenta dots). The *euo*prom, *hctA*prom and *hctB*prom+
cell population were also identified using TrackMate and the signal from the FISH channel and fluorescent
protein channels were plotted on the same graphs (green dots). (B) Cells were infected with L2-PsciEng and
the developmental gene expression kinetics were compared to those of L2-AsciEng, and L2-BsciEng. The
*euo*prom expression kinetics were comparable for all strains with expression first detected at ~15 hpi.
*PorB*prom expression kinetics were nearly identical to *hctA*prom expression kinetics, first detected at ~20 hpi,
while *hctB*prom expression was initiated at ~26 hpi. Error cloud for fluorescent reporters represents SEM.
n > 20 inclusions per strain.
771
Figure 6. The effects of ectopic expression of Euo, HctA, CtcB and HctB on T3SS structural genes. The
log2fold change RNA-seq differential expression data from the ectopic expression experiments were plotted
against the -log of the p value (-log padj) and the operons for the T3SS were highlighted. For the wt *Ctr* 18 hpi
vs 24 hpi samples most of the T3SS structural genes were upregulated while for the Euo ectopic expression
experiment most of these operons were downregulated. Again, like the IB genes in the HctA and CtcB ectopic
expression experiments, most of the structural genes were downregulated or unchanged. Two operons did not
follow this pattern, the *CTL0238*-op and *scc2*-op. Both operons encode the components of the T3SS
translocon. The *CTL0238*-op (gold circle) was regulaged like an RB gene while the *scc2*-op (purple circle) was
regulated like an EB gene.
781
Figure 7. IB cell type expression of the T3SS structural operon SctU. (A) Cells were infected with L2-AsciEng
for 16 hpi and fixed and stained using a FISH probe (*sctU-lcrD*) to the mRNA for the T3SS structural operon
sctU-op, the RB control *euo* and the IB control *hctA*. All cells were positive for *euo*prom expression (green).
The FISH stained cells were only positive for *euo* mRNA (magenta) and were negative for *hctA* mRNA
(magenta) and *sctU-op* mRNA (magenta). (B) Cells were infected with L2-AsciEng and L2-BsciEng for 24 hpi
and fixed and stained using FISH for the *sctU*-op mRNA. For the *euo*prom sample, the *sctU*-op FISH signal
(magenta) was present in a distinct subset of cells and not in the majority of the *euo*prom+ cells (green). (C)
TrackMate was used to identify the *sctU*-op mRNA+ cells from 5 inclusions from 24 hpi samples and the signal
for *euo*prom and FISH were quantified for each *sctU*-op+ cell and plotted (magenta dots). The converse was
also performed, the *euo*prom+ cells were identified and the *euo*prom signal and FISH signal was quantified for
each *euo*prom+ cell and plotted (green dots). The FISH signal was also compared to the *hctA*prom expression
pattern and showed subsets of cells that were stained for both *sctU*-op mRNA and *hctA*prom expression as
well as non overlapping populations. The *sctU*-op mRNA+ cells were again identified using TrackMate and the
signal for *hctA*prom and FISH were quantified for each *sctU*-op+ cell and plotted (magenta dots). Each
*hctA*prom+ cell was also identified and the FISH and *hctA*prom signal was determined and plotted (green

dots). The *sctU*-op FISH staining was also compared to the expression from the *hctB*prom reporter. The *sctU*-
op mRNA FISH staining was again present in a subset of cells but showed little overlap with the *hctB*prom
fluorescent signal. The FISH signal and *hctB*prom signal were measured in both cell populations (*sctU* mRNA+
cells and *hctB*prom+ cells) and plotted, *sctU* mRNA+ cells magenta dots and *hctB*prom+ cells green dots. Both
populations were primary single positive, either *sctU*-op mRNA high or *hctB*rpom high but rarely both. Size bar
= 5µm.

Figure 8. Cell type expression of the two T3SS translocons. A) Cos-7 cells were infected with L2-BsciEng for
24 hpi and stained for the mRNA expression of the *CTL0238*-op using FISH (magenta), *euo*prom expression
(green) and *hctB*prom expression (green). Individual chlamydial cells with *CTL0238*-op mRNA signal from 5
separate inclusions were identified using TrackMate and the fluorescence intensity for each channel (mRNA
and promoter reporter) was plotted (magenta dots). Individual chlamydial cells positive for *euo*prom, or
*hctB*prom signal from 5 separate inclusions were also identified using TrackMate and the expression intensity
for each channel (mRNA and promoter reporter) was plotted (green dots). B) Cos-7 cells were infected with L2-
BsciEng for 24 hpi and stained for the mRNA expression of the *scc2*-op using FISH. Scc2-op FISH signal in
magenta, *euo*prom and *hctB*prom signal in green. Individual chlamydial cells positive for *scc2*-op mRNA signal
from 5 inclusions were identified using TrackMate and the fluorescence intensity for each channel (mRNA and
promoter reporter) was plotted (magenta dots). Individual chlamydial cells positive for *euo*prom, or *hctB*prom
signal from 5 inclusions were also identified using TrackMate and the expression intensity for each channel
(mRNA and promoter reporter) was plotted (green dots). Host cells infected with AsciEng and fixed at 16 hpi
were probed for *CTL0238*-op mRNA (C) and scc2-op mRNA (D) expression using FISH. *Euo*prom expression
(green) had significant overlap with *CTL0238*-op mRNA signal. For the *scc2*-op the FISH signal was
undetected. Size bar = 5µm.

Figure 9. Effects of ectopic expression of Euo, HctA, CtcB and HctB on the expression of T3SS effectors.
RNA-seq differential expression data (log2fold change) plotted vs the - log of the P value (-log padJ) for *Ctr*
ectopically expressing Euo, HctA, CtcB and HctB. **(A)** The T3SS effectors are highlighted in purple. (B) All the
inc protein genes are highlighted in purple while the genes for the *incD-G* operon are highlighted in green, *incV*
and *incM* are highlighted in red and orange respectively.

Figure 10. Cell type expression of *incD* and *incV*. Cos-7 cells infected with L2-BsciEng for 16 and 24 hpi and
stained for *incD* and *incV* mRNA expression using custom FISH probes. (A) The *incD* mRNA (magenta) was
visibly expressed in the *euo*prom+ (green) RB cells at 16 hpi while *hctB*prom signal was not detected. At 24 hpi
the *incD* mRNA signal (magenta) overlapped with the *euo*prom signal (green) but was separate from the
*hctB*prom+ cells (yellow). (B) The *incV* mRNA signal (magenta) was undetected at 16 hpi. At 24 hpi the *incV*
mRNA signal showed overlap with the *hctB*prom signal (yellow) but not the *euo*prom signal (green). C)
Individual chlamydial cells positive for *incD* mRNA signal were identified from 5 separate inclusions at 24 hpi
using TrackMate and the fluorescence intensity for each channel (mRNA and promoter reporter) was plotted
(magenta dots). Individual chlamydial cells positive for *euo*prom, or *hctB*prom signal were also identified using
TrackMate and the expression intensity for each channel (mRNA and promoter reporter) was plotted (green
dots). D) Individual chlamydial cells positive for *incV* mRNA signal from 5 separate inclusions at 24 hpi were
identified using TrackMate and the fluorescence intensity for each channel (mRNA and promoter reporter) was
plotted (magenta dots). Individual chlamydial cells positive for *euo*prom or *hctB*prom signal were also identified
using TrackMate and the expression intensity for each channel (mRNA and promoter reporter) was plotted
(green dots). Size bar = 5µm.

Figure 11. Model of cell type specific deployment of the T3SS. In this model the *scc2*-op translocon secretes
effectors across the plasma membrane (PM) for host cell entry. The *scc2*-op translocon is replaced in the RB
with the *CTL0239*-op translocon for the secretion of the Inc proteins across the inclusion membrane (IM). The

structural components of the T3SS are then reconstructed during the IB to EB maturation phase. The Euo+,
HctA+, CtcB+ and HctB+ nomenclature indicates in which cell form the protein is active.
848

849 Supplemental table legends
850 Table S1. Chlamydial genes designated as RB, IB, EB expressed genes and their regulation from ectopic
851 expression of Euo, HctA, CtcB and HctB. Gene expression time categories: Immediate early genes, Mid-late
852 genes and Late genes from Soules et al and Belland et al. Serovar D annotation included just for reference
853 and does not indicate experimentally measured values.
854

855 Table S2. T3SS structural operons.
856

857 Table S3. T3SS effector gene regulation from ectopic expression of Euo, HctA, CtcB and HctB. Serovar D
858 annotation included for reference and does not indicate experimentally measured values.
859

860 Table S4. Inc gene regulation from ectopic expression of Euo, HctA, CtcB and HctB. Serovar D annotation
861 included just for reference and does not indicate experimentally measured values.
862

863 Table S5. List of primers used to construct plasmids
864

865 Table S6. List of FISH probes and the location on the Ctr genome

References:

1. Abdelrahman, Y.M., and Belland, R.J. (2005). The chlamydial developmental cycle. FEMS Microbiol. Rev. *29*, 949–959. https://doi.org/10.1016/J.FEMSRE.2005.03.002.

2. Rockey, D.D., and Matsumoto, A. (2000). The chlamydial developmental cycle. Prokaryotic Dev., 403–425.

3. Bakken, I.J. (2007). Chlamydia trachomatis Infection and the Risk for Ectopic Pregnancy. Sex. Transm. Dis. *34*. https://doi.org/10.1097/01.OLQ.0000253171.99295.48.

4. Ohman, H., Tiitinen, A., Halttunen, M., Lehtinen, M., Paavonen, J., and Surcel, H.-M. Cytokine polymorphisms and severity of tubal damage in women with Chlamydia-associated infertility. J. Infect. Dis. *199*, 1353–1359. https://doi.org/10.1086/597620.

5. Reekie, J., Donovan, B., Guy, R., Hocking, J.S., Kaldor, J.M., Mak, D., Preen, D., Ward, J., Liu, B., and Chlamydia and Reproductive Health Outcome Investigators (2019). Risk of Ectopic Pregnancy and Tubal Infertility Following Gonorrhea and Chlamydia Infections. Clin. Infect. Dis. *69*, 1621–1623. https://doi.org/10.1093/cid/ciz145.

6. Schachter, J. (1988). Chlamydiaceae: The Chlamydiae. In, pp. 847–863. https://doi.org/10.1007/978-1-4612-3900-0_43.

7. Everett, K.D., Bush, R.M., and Andersen, A.A. (1999). Emended description of the order Chlamydiales, proposal of Parachlamydiaceae fam. nov. and Simkaniaceae fam. nov., each containing one monotypic genus, revised taxonomy of the family Chlamydiaceae, including a new genus and five new species, and standards for the identification of organisms. Int. J. Syst. Evol. Microbiol. *49 Pt 2*, 415–440. https://doi.org/10.1099/00207713-49-2-415.

8. Chiarelli, T.J., Grieshaber, N.A., Omsland, A., Remien, C.H., and Grieshaber, S.S. (2020). Single-Inclusion Kinetics of Chlamydia trachomatis Development. MSystems *5*. https://doi.org/10.1128/MSYSTEMS.00689-20.

9. Chiarelli, T.J., Grieshaber, N.A., Appa, C.R., and Grieshaber, S.S. (2023). Computational Modeling of the Chlamydial Developmental Cycle Reveals a Potential Role for Asymmetric Division. mSystems *8*, e00053-23. https://doi.org/10.1128/msystems.00053-23.

10. Clifton, D.R., Dooley, C.A., Grieshaber, S.S., Carabeo, R.A., Fields, K.A., and Hackstadt, T. (2005). Tyrosine phosphorylation of the chlamydial effector protein Tarp is species specific and not required for recruitment of actin. Infect. Immun. *73*, 3860–3868. https://doi.org/10.1128/IAI.73.7.3860-3868.2005.

11. Chen, Y.-S., Bastidas, R.J., Saka, H.A., Carpenter, V.K., Richards, K.L., Plano, G.V., and Valdivia, R.H. (2014). The Chlamydia trachomatis type III secretion chaperone Slc1 engages multiple early effectors, including TepP, a tyrosine-phosphorylated protein required for the recruitment of CrkI-II to nascent inclusions and innate immune signaling. PLoS Pathog. *10*. https://doi.org/10.1371/JOURNAL.PPAT.1003954.

12. Abdelrahman, Y., Ouellette, S.P., Belland, R.J., and Cox, J.V. (2016). Polarized Cell Division of Chlamydia trachomatis. PLoS Pathog. *12*. https://doi.org/10.1371/JOURNAL.PPAT.1005822.

13. Wurihan, W., Zou, Y., Weber, A.M., Weldon, K., Huang, Y., Bao, X., Zhu, C., Wu, X., Wang, Y., Lai, Z., et al. (2021). Identification of a GrgA-Euo-HrcA Transcriptional Regulatory Network in Chlamydia. mSystems *6*, e0073821. https://doi.org/10.1128/mSystems.00738-21.

14. Belland, R.J., Zhong, G., Crane, D.D., Hogan, D., Sturdevant, D., Sharma, J., Beatty, W.L., and Caldwell, H.D. (2003). Genomic transcriptional profiling of the developmental cycle of Chlamydia trachomatis. Proc. Natl. Acad. Sci. *100*, 8478–8483. https://doi.org/10.1073/PNAS.1331135100.

15. Shaw, E.I., Dooley, C.A., Fischer, E.R., Scidmore, M.A., Fields, K.A., and Hackstadt, T. Three temporal classes of gene expression during the Chlamydia trachomatis developmental cycle. Mol. Microbiol. *37*, 913–925.

16. Nicholson, T.L., Olinger, L., Chong, K., Schoolnik, G., and Stephens, R.S. (2001). Global Stage-Specific Gene Regulation during the Developmental Cycle of Chlamydia trachomatis. J. Bacteriol. *185*, 3179–3189. https://doi.org/10.1128/JB.185.10.3179-3189.2003.

17. Hakiem, O.R., Rizvi, S.M.A., Ramirez, C., and Tan, M. Euo is a developmental regulator that represses late

916 genes and activates midcycle genes in *Chlamydia trachomatis*. mBio *0*, e00465-23.
917 https://doi.org/10.1128/mbio.00465-23.
918 18. Rosario, C.J., Hanson, B.R., and Tan, M. The transcriptional repressor EUO regulates both subsets of
919 Chlamydia late genes. Mol. Microbiol. *94*, 888–897. https://doi.org/10.1111/mmi.12804.
920 19. Barry, C.E., Brickman, T.J., and Hackstadt, T. (1990). Hc1-mediated effects on DNA structure: a potential
921 regulator of chlamydial development. Mol. Microbiol. *9*, 273–283. https://doi.org/10.1111/j.1365-
922 2958.1993.tb01689.x.
923 20. Grieshaber, N.A., Grieshaber, S.S., Fischer, E.R., and Hackstadt, T. A small RNA inhibits translation of the
924 histone-like protein Hc1 in Chlamydia trachomatis. Mol. Microbiol. *59*, 541–550.
925 https://doi.org/10.1111/j.1365-2958.2005.04949.x.
926 21. Grieshaber, N.A., Fischer, E.R., Mead, D.J., Dooley, C.A., and Hackstadt, T. (2004). Chlamydial histone-
927 DNA interactions are disrupted by a metabolite in the methylerythritol phosphate pathway of isoprenoid
928 biosynthesis. Proc. Natl. Acad. Sci. U. S. A. *101*, 7451—7456. https://doi.org/10.1073/pnas.0400754101.
929 22. Brickman, T.J., Barry, C.E. 3rd, and Hackstadt, T. (1991). Molecular cloning and expression of hctB
930 encoding a strain-variant chlamydial histone-like protein with DNA-binding activity. J. Bacteriol. *175*.
931 23. Koo, I.C., and Stephens, R.S. (2003). A developmentally regulated two-component signal transduction
932 system in Chlamydia. J. Biol. Chem. *278*, 17314–17319. https://doi.org/10.1074/JBC.M212170200.
933 24. Soules, K.R., LaBrie, S.D., May, B.H., and Hefty, P.S. (2020). Sigma 54-Regulated Transcription Is
934 Associated with Membrane Reorganization and Type III Secretion Effectors during Conversion to
935 Infectious Forms of Chlamydia trachomatis. MBio *11*. https://doi.org/10.1128/MBIO.01725-20.
936 25. Grieshaber, N.A., Chiarelli, T.J., Appa, C.R., Neiswanger, G., Peretti, K., and Grieshaber, S.S. (2022).
937 Translational gene expression control in Chlamydia trachomatis. PLoS ONE *17*.
938 https://doi.org/10.1371/JOURNAL.PONE.0257259.
939 26. Waskom, M.L. (2021). seaborn: statistical data visualization. J. Open Source Softw. *6*, 3021.
940 https://doi.org/10.21105/joss.03021.
941 27. Hatch Nathan D. and Ouellette Scot P. (2023). Identification of the alternative sigma factor regulons of
942 Chlamydia trachomatis using multiplexed CRISPR interference. mSphere *8*, e00391-23.
943 https://doi.org/10.1128/msphere.00391-23.
944 28. Tinevez, J.-Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E., Bednarek, S.Y.,
945 Shorte, S.L., and Eliceiri, K.W. (2014). TrackMate: An open and extensible platform for single-particle
946 tracking. Methods *115*, 80–90. https://doi.org/10.1016/j.ymeth.2016.09.016.
947 29. Kubo, A., and Stephens, R.S. (2000). Characterization and functional analysis of PorB, a Chlamydia porin
948 and neutralizing target. Mol. Microbiol. *38*, 772–780. https://doi.org/10.1046/J.1365-2958.2000.02167.X.
949 30. Grieshaber, S., Grieshaber, N., Yang, H., Baxter, B., Hackstadt, T., and Omsland, A. (2018). Impact of
950 Active Metabolism on Chlamydia trachomatis Elementary Body Transcript Profile and Infectivity. J.
951 Bacteriol. *200*. https://doi.org/10.1128/JB.00065-18.
952 31. Price, M.N., Huang, K.H., Alm, E.J., and Arkin, A.P. (2005). A novel method for accurate operon
953 predictions in all sequenced prokaryotes. Nucleic Acids Res. *33*, 880–892.
954 https://doi.org/10.1093/nar/gki232.
955 32. Hefty, P.S., and Stephens, R.S. (2007). Chlamydial type III secretion system is encoded on ten operons
956 preceded by sigma 70-like promoter elements. J. Bacteriol. *189*, 198–206.
957 https://doi.org/10.1128/JB.01034-06.
958 33. Stephens, R.S., Kalman, S., Lammel, C., Fan, J., Marathe, R., Aravind, L., Mitchell, W., Olinger, L.,
959 Tatusov, R.L., Zhao, Q., et al. (1998). Genome sequence of an obligate intracellular pathogen of humans:
960 Chlamydia trachomatis. Science *282*, 754–759. https://doi.org/10.1126/SCIENCE.282.5389.754.
961 34. Dey, S., Chakravarty, A., Guha Biswas, P., and De Guzman, R.N. (2019). The type III secretion system
962 needle, tip, and translocon. Protein Sci. Publ. Protein Soc. *28*, 1582–1593. https://doi.org/10.1002/pro.3682.
963 35. Betts-Hampikian, H.J., and Fields, K.A. (2010). The Chlamydial Type III Secretion Mechanism: Revealing
964 Cracks in a Tough Nut. Front. Microbiol. *1*. https://doi.org/10.3389/FMICB.2010.00114.
965 36. Dehoux, P., Flores, R., Dauga, C., Zhong, G., and Subtil, A. Multi-genome identification and

characterization of chlamydiae-specific type III secretion substrates: the Inc proteins. BMC Genomics *12*.
https://doi.org/10.1186/1471-2164-12-109.

37. Peters, J., Wilson, D.P., Myers, G., Timms, P., and Bavoil, P.M. (2005). Type III secretion à la Chlamydia.
Trends Microbiol. *15*, 241–251. https://doi.org/10.1016/j.tim.2007.04.005.

38. Ferrell, J.C., and Fields, K.A. A working model for the type III secretion mechanism in Chlamydia.
Microbes Infect. Inst. Pasteur *18*, 84–92. https://doi.org/10.1016/j.micinf.2015.10.006.

39. Lutter, E.I., Martens, C., and Hackstadt, T. Evolution and conservation of predicted inclusion membrane
proteins in chlamydiae. Comp. Funct. Genomics *2012*. https://doi.org/10.1155/2012/362104.

40. Rockey, D.D., Scidmore, M.A., Bannantine, J.P., and Brown, W.J. (2002). Proteins in the chlamydial
inclusion membrane. Microbes Infect. *4*, 333–340. https://doi.org/10.1016/S1286-4579(02)01546-0.

41. Stanhope, R., Flora, E., Bayne, C., and Derré, I. (2017). IncV, a FFAT motif-containing Chlamydia protein,
tethers the endoplasmic reticulum to the pathogen-containing vacuole. Proc. Natl. Acad. Sci. *114*, 12039–
12044. https://doi.org/10.1073/PNAS.1709060114.

42. Luís, M.P., Pereira, I.S., Bugalhão, J.N., Simões, C.N., Mota, C., Romão, M.J., and Mota, L.J. (2023). The
Chlamydia trachomatis IncM Protein Interferes with Host Cell Cytokinesis, Centrosome Positioning, and
Golgi Distribution and Contributes to the Stability of the Pathogen-Containing Vacuole. Infect. Immun. *91*,
e0040522. https://doi.org/10.1128/iai.00405-22.

43. Wilson, D.P., Timms, P., McElwain, D.L.S., and Bavoil, P.M. (2006). Type III secretion, contact-dependent
model for the intracellular development of chlamydia. Bull. Math. Biol. *68*, 161–178.
https://doi.org/10.1007/s11538-005-9024-1.

44. Hoare, A., Timms, P., Bavoil, P.M., and Wilson, D.P. (2006). Spatial constraints within the chlamydial host
cell inclusion predict interrupted development and persistence. BMC Microbiol. *8*.
https://doi.org/10.1186/1471-2180-8-5.

45. Grieshaber, N.A., Runac, J., Turner, S., Dean, M., Appa, C., Omsland, A., and Grieshaber, S.S. (2021). The
sRNA Regulated Protein DdbA Is Involved in Development and Maintenance of the Chlamydia trachomatis
EB Cell Form. Front. Cell. Infect. Microbiol. *11*. https://doi.org/10.3389/FCIMB.2021.692224.

46. Shen Li, Gao Leiqiong, Swoboda Abigail R., and Ouellette Scot P. (2024). Targeted repression of topA by
CRISPRi reveals a critical function for balanced DNA topoisomerase I activity in the Chlamydia
trachomatis developmental cycle. mBio *15*, e02584-23. https://doi.org/10.1128/mbio.02584-23.

47. Niehus, E., Cheng, E., and Tan, M. (2008). DNA supercoiling-dependent gene regulation in Chlamydia. J.
Bacteriol. *190*, 6419–6427. https://doi.org/10.1128/JB.00431-08.

48. Cheng Eric and Tan Ming (2012). Differential Effects of DNA Supercoiling on Chlamydia Early Promoters
Correlate with Expression Patterns in Midcycle. J. Bacteriol. *194*, 3109–3115.
https://doi.org/10.1128/jb.00242-12.

49. Howard, L., Orenstein, N.S., and King, N.W. (1970). Purification on renografin density gradients of
Chlamydia trachomatis grown in the yolk sac of eggs. Appl. Microbiol. *27*, 102–106.

50. Derré, I., Swiss, R., and Agaisse, H. (2011). The lipid transfer protein CERT interacts with the Chlamydia
inclusion protein IncD and participates to ER-Chlamydia inclusion membrane contact sites. PLoS Pathog. *7*.
https://doi.org/10.1371/JOURNAL.PPAT.1002092.

51. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat. Methods *9*, 357–
359. https://doi.org/10.1038/nmeth.1923.

52. Love, M.I., Huber, W., and Anders, S. (2012). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol *15*. https://doi.org/10.1186/s13059-014-0550-8.

53. Shukla, X.U., and Parmar, D.J. (2014). Python – A comprehensive yet free programming language for
statisticians. J. Stat. Manag. Syst. *19*, 277–284. https://doi.org/10.1080/09720510.2015.1103446.