Title: Predicting gene expression responses to environment in *Arabidopsis thaliana* using natural

variation in DNA sequence

Authors: Margarita Takou[1], Emily S Bellis[1,2,] & Jesse R Lasky[1]

Affiliations:

1 Pennsylvania State University, University Park, 16802, USA

2 Department of Computer Science, Arkansas State University, Jonesboro AR

email addresses:

Margarita Takou: mtakou1@uni-koeln.de

Emily S Bellis: embellis@gmail.com

Jesse R Lasky: jrl35@psu.edu

Corresponding author:

Margarita Takou: mtakou1@uni-koeln.de

Abstract:

The evolution of gene expression responses is a critical component of adaptation to variable environments. Predicting how DNA sequence influences expression is challenging because the genotype to phenotype map is not well resolved for *cis* regulatory elements, transcription factor binding, regulatory interactions, and epigenetic features, not to mention how these factors respond to environment. We tested if flexible machine learning models could learn some of the underlying *cis*-regulatory genotype to phenotype map. We tested this approach using cold-responsive transcriptome profiles in 5 diverse *Arabidopsis thaliana* accessions. We first tested for evidence that *cis* regulation plays a role in environmental response, finding 14 and 15 motifs that were significantly enriched within the up- and down-stream regions of cold-responsive differentially regulated genes (DEGs). We next applied convolutional neural networks (CNNs), which learn *de novo cis*-regulatory motifs in DNA sequences to predict expression response to environment. We found that CNNs predicted differential expression with moderate accuracy, with evidence that predictions were hindered by biological complexity of regulation and the large potential regulatory code. Overall, approaches to predict DEGs between specific environments based only on proximate DNA sequences require further development, and additional information may be required.

Key words: low frequency variants, gene expression prediction, machine learning, regulatory elements, evolution, genotype to phenotype map

Introduction

Adaptation is often a complex process that occurs via selection on traits and the underlying genetic mechanisms (1). The genetic basis of ecologically important quantitative traits often involves mutations with effects on gene expression. In variable environments, the evolution of gene expression responses to environment (expression plasticity) are likely a critical component of adaptation (2). These processes are often studied using evidence from variation in DNA sequence and mRNA abundance. For instance, the incorporation of transcriptomic information into genome wide association studies (GWAS) can help identify genes controlling a specific phenotype (3). Association or linkage mapping approaches can be used to map expression quantitative trait loci (eQTL), and allele-specific expression can determine the presence of *cis* or *trans* acting eQTL (4–6). However, the integration of sequence and expression data in evolutionary quantitative genomics is hindered by gaps in our understanding of how DNA sequence influences expression. In particular, understanding the genetic basis of expression plasticity might be beneficial to applied biology given the importance of predicting organismal responses to environmental changes(Keagy et al. 2023).

Gene expression responses to the environment differ among genes in the genome and among different natural genotypes due to DNA sequence variation. Differences in gene expression among genotypes may be driven by sequence differences at other loci in the genome, which regulate expression of a given gene (*trans* effects). At the same time, differences among genes and genotypes may arise from differences in sequence proximal to the gene that determine how its expression is regulated (*cis* effects). Researchers have documented cases where mutations in genes involved in environmental sensing (8,9) or in environmentally responsive genes that bind DNA sequences (10,11) have downstream effects on expression of other genes. Additionally,

researchers have characterized how some *cis* mutations influence DNA binding by

environmentally responsive transcription factors, ultimately altering environmental responsive

expression (12–14). Transcription factors that tend to show expression responses to environment

have been documented, and the DNA sequence motifs they bind to have been roughly

characterized (15). However, across the genome there are many transcription factors that often

respond to a single environmental stressor and interact, and this knowledge is currently too

limited to predict how genetic variation in DNA sequences influence genome-wide expression

responses to environment.

The large number of *cis*-regulatory motifs allows the investigation of genomic and evolutionary

properties, which are associated with their distribution across the genome (16–18). Genome-wide

expression profiling in association mapping or linkage mapping populations as well as allele-

specific expression studies have revealed that *cis* expression quantitative trait loci (*cis*-eQTL) are

abundantly segregating within species. *Cis*-regulatory mutations have been of particular

importance for evolutionary biology because some have suggested they may be less likely to be

deleterious than mutations in transcription factors, which are expected to be more pleiotropic

(19,20). Relatedly, the possibility of easier dissection of the genotype to phenotype map for *cis*-

eQTL (versus *trans*-acting mutations likely influencing multiple genes' expression) combined

with the abundance of *cis*-eQTL may be an opportunity to build a more integrative understanding

of quantitative trait genetics.

Using machine learning methods in fields such as genomics and population genetics can help

navigate increasingly bigger datasets and reveal complex patterns (21). Especially in gene

regulation and evolution, deep learning approaches can have an advantage over traditional

methods in decoding enhancer grammar of gene regulation, as these models can learn complex *cis*-regulatory rules in a precise manner not biased by current knowledge (22) and can outperform more traditional methods, like clustering by kmer or linear regression in predicting gene expression (23). For example, training a machine learning model on the imputed *cis* haplotypic information of RNA expression, Giri et al. (2021) managed to exclude the impact of trans effects on gene expression variation in maize. The prediction was less variable than those using SNPs and had an increased accuracy for predicting within population variance (24). Random forest models successfully classified protein coding vs non-protein coding genes in maize (25). Moore et al. (2021) showed how known regulatory sites of Arabidopsis thaliana and transcription factors (TFs) interactions can be used in machine learning to predict clusters of gene co-expression related to stress response (26). Similarly, in Arabidopsis thaliana, information on TF binding sites has been harnessed to predict differential gene expression under stress such as response to Fe (27), high salinity (28) and combined heat and drought stress (29). However, in all of these cases, the authors used additional -omics information, such as chromatin accessibility, or co-expression clusters, during deep learning. There is increasing evidence that genetic diversity on the *cis* elements of plants is under selection (30). Deep learning algorithms could be used for genome wide identification of deleterious and adaptive mutations, which is a prerequisite for the genetic improvement of crops (31). Using upstream and downstream genetic coding information could improve the prediction of the role of *cis* regulatory elements in differential gene expression, in contrast to using only upstream regions in maize (32), especially when more than one genotype was included in the analysis (33). However, those methods do not capture distal regulatory elements well if at all (34) and the source of variability can impact their efficiency (35).

In *Arabidopsis thaliana,* whose importance in studying evolution and ecology patterns is

growing over the years (36), the conserved noncoding sequences are mostly concentrated close

to the coding region and impact gene regulation (37), as *cis*-regulatory variants are usually in

linkage with the expressed transcript (16,38,39). Even though there are conserved gene reactions

between species and populations, the increased genetic diversity within populations or species

causes phenotypic variance. Genes, whose expression is affected by genotype x environment

interactions (GxE), tend to show allele frequency correlations with climate and have elevated

genetic variation in stress responsive transcription factor binding sites (40). Known TF binding

site motifs of *A. thaliana* have been used to identify putative binding sites and then train a deep

model for predicting *cis* changes in tomato (41).

Our aim was to use genetic variation among genes within individuals and among genotypes in

the regions upstream and downstream of genes to predict expression response to environment in

A. thaliana. We aimed to predict in a both a more general sense than other studies by using the

sequences instead of known TFBS, and in a more restrictive sense by classifying gene expression

in specific environments within one species. For the analysis we used a published dataset on

differentially expressed genes (DEGs) during cold acclimation of diverse *A. thaliana* genotypes

(42). Response to cold stress not only has some well characterized signaling pathways,

transcription factors and transcription factor binding sites, but also has been connected to

adaptation to local environments (43–45). Hannah et al. (2006) had identified approximately

1,500 up regulated DEGs and down regulated DEGs out of around 20,000 genes for multiple

accessions as a response to cold acclimation and stress (S1 Table). Here, we focus on the five *A.*

thaliana accessions *Can, Col-0, Cvi, Ler* and *Rsch,* for which the pseudogenomes from whole

genome resequencing data are published (46). We included multiple genotypes in the analysis, so

as to capture some of the natural genetic diversity and potential GxE responses.

We first tested whether there are specific sequence motifs enriched along those regions of the

differentially expressed genes that would suggest *cis* regulation drives some aspect of cold

expression response. We determined if these motifs were related to known stress-responsive

transcription factors (TFs). Then, we tested whether several naïve deep learning models could

predict expression response, without a priori designation of important motifs, by using only the

sequences upstream and downstream of genes.

Material and Methods

Transcriptome and genome data

We used the publicly available dataset of differentially gene expression during cold acclimation

by (42). In brief, the authors used multiple accessions of *A. thaliana* to test their cold acclimation

potential. They found that approximately 4,000 genes respond to cold acclimation, indicating

regulatory pathways being involved in their control. Out of the accessions used in the original

study, we used the five accessions, *Col-0, Ler, Cvi, Can* and *Rsch* (S1 Table) whose

pseudogenomes are available in https://1001genomes.org/. For comparison with a potentially

more simple transcriptomic change, we used the differential gene expression of 1,280 genes in

the *Col-0* background of a *tt8* knock out line. The *tt8* gene regulates biosynthesis of

anthocyanins, which we reasoned would be a narrower regulatory network than that involved in

environmental responses (47).

Searching for sequence motifs using STREME

Transcription factors bind DNA in a sequence specific manner, but not all sequences are known nor is the exact relationship between sequence and binding affinity known. For this reason we began with a *de novo* search for sequence motifs in the upstream and downstream regions of the coding sequence, as those often contain such signals. We used samtools faidx v1.10 (48) to extract the 1kbp upstream and downstream regions of each gene present in the dataset, including the first and last part of the coding region as TFBS often bind there. Specifically, we extracted the 500bp upstream and 500bp downstream of the gene's TSS for the upstream regulatory coding region, and the 500bp upstream and 500bp downstream of the gene's end transcriptional site for the downstream regulatory coding region. We then used the software STREME (49) to calculate significantly enriched ungapped motifs within the set of up- and down- regulated DEGs in relation to their presents in non-DEGs. For both groups we used as a control a randomly down-sampled set of the non-DEGs and stopped the search for motifs when we failed to identify motifs with high enough confidence (e-value > 0.05) three times in a row. This way, the statistical significance of each motif was determined based on whether it can classify sequences as DEGs or non-DEGs.

Because sequence motifs may have a variety of causes and functions, and not all are likely TFBS, we next sought to determine whether motifs discovered *de novo* by STREME specifically correspond to known transcription factor binding sites. We searched for significant overlaps of at least 5bp between the discovered enriched motifs and the JASPAR 2022 CORE (50) plant transcription factor binding sites database with MEME suite's online tool TOMTOM (51). Significant overlap was determined by scoring the match of all possible alignments between

sequence and known motif and summing those scores to produce p values, q values, and e

values.

Random forest analysis

To determine whether known transcription factor binding sites regulate the response to cold, we

attempted to predict up regulated or down regulated DEGs vs non DEGs based on their presence

or absence in the regulatory regions. We used a dataset (52) of TF putative binding sites 1000 bp

upstream of each gene, which were predicted from *A. thaliana* using JASPAR2018 Bioconductor

package (JASPAR) and then matched to the aligned regions of *A. thaliana*. We treated the count

of each TF binding site per gene as a separate feature and additionally determined the total

number of TF binding sites per gene. Taken together, these descriptors of genomic variation

provided a total of 413 features, of which we could assess the predictive impact on gene

regulation.

We trained a random forest model for *Col-0* using the ranger library in R (54) with up-regulated

DEGs in response to cold and non-DEGs as variable and all compiled features of gene regulation

as predictor variables. We set the number of trees of the forest to 500, the number of variables to

possibly split at each node (mtry) to 200, and the impurity mode to impurity_corrected. From the

trained random forest, we extracted the relative importance of each predictor variable as well as

the out-of-bag prediction error, which is informative about the predictive power of the model.

Finally, we estimated a *p* value for each predictor variable using the "Altmann" method with 100

permutations. We repeated the random forest analysis for all down-regulated DEGs. We

estimated the prAUC in R library using the function prauc and extracted its significance by

permuting 100 times the gene identity for each set of variables.

Training Convolutional Neural Networks

We employed a more naive machine learning approach to determine whether we can predict

DEG using regulatory regions. We trained convolutional neural networks (CNNs) to identify a

model that would predict whether each gene was differentially expressed in the stress conditions

relative to the control with high accuracy. We prepared the dataset by separating the genes into a

training and test set for each accession, incorporating information about the gene families that

they belonged to (55). Because sequences may carry spurious associations with expression that

are due to shared evolutionary history, we followed the method of Washburn et al. (2019) and

split genes into gene families and then tested predictions in gene families not used for training.

In the end, we used 68,979 (80%) genes, from which the 20% (13,796 genes) was always

retained as the validation set, and 17,475 (20%) as the test set out of the total 86,454 genes

across all five genotypes. Each unique gene could be present multiple times in the dataset,

representing each accession. The proportion of up- and down-regulated DEGs to the non DEGs

was approximately 1:11 in both sets (S1 Table). For each gene, we used samtools faidx v1.10

(48) to extract the 1.5kbp upstream and downstream regions of each gene from the published

pseudogenomes, using custom scripts. Specifically, we extracted the 1,000 upstream and 500 bp

downstream of the gene's transcriptional starting site (TSS) for the upstream regulatory coding

region, and the 500 bp upstream and 1,000 bp downstream of the gene's transcriptional ending

site for the downstream regulatory coding region. Those sequences were converted into one hot

encoding of each nucleotide in python v3.6, to be used as input for the training and testing of the

CNN models. We then created the labels needed for binary classification of DEG during training

by scoring the up-regulated DEGs as 0 and the non-DEGs as 1. As the number of up-regulated

DEGs was much smaller than non-DEGs, we oversampled with replacement the up-regulated

DEGs to a ratio of 1:1 between the two groups. This process was repeated for training models to

classify genes as down-regulated DEGs and non-DEGs.

We used the prepared inputs and labels of the training set to perform a grid search of tuning

parameters, as we wanted to identify both the models with the best prediction of our dataset, as

well as how different tuning parameters influence model accuracy using the regulatory regions of

multiple accessions. The grid search was performed on a total of 1,344 combinations of

hyperparameters (55) by training each model for 50 epochs using the python library tensorflow-

gpu v2 (56). The tuning parameters included: the number of convolutional layers, the filter of

each convolutional layer, the width of each convolutional layer, the pool width and stride, the

number of dense layers and their units, as well as the dropout rate. A list of the specific values

used to create all the different combinations is provided in S2 Table. During each training phase,

we used the prAUC values of the validation set to tune, select and save the best trained model for

further evaluation. For each trained model, we extracted the loss, accuracy, and the probability-

based accuracy under the curve (prAUC) of the training, validation and testing of each model to

compare the accuracy of the models. Moreover, we estimated the proportion of correct

predictions of each class within the test set so that we could assess the potential overfitting of

each model to predicting one class or the other. We estimated this within class accuracy as the

number of times that the class was predicted correctly divided by the total number of genes

belonging in this class. In this context, class refers to up-regulated or non-DEG genes. The

absolute difference of within class accuracies, referred to in the text as prediction accuracy

between the two classes, was also estimated. We repeated this process for all down-regulated

DEGs.

Determining factors influencing accuracy of CNNs

To determine the degree to which predictions relied on upstream sequence versus both up and

downstream sequences, we repeated the above analysis described for the regulatory regions of all

accessions by using only the upstream regions.

Because genetic variation among genotypes in expression can be driven by *trans*-effects (from

other loci far from the gene), we sought to control for some of this variation. In particular, we

focused on known loss of function variation at large effect CBF transcription factors that are

involved in adaptation to different temperatures, first by training and evaluating CNN models on

those lines (*Can* and *Cvi),* which have all 3 CBFs functional copies differentially expressed

between the experimental conditions (45,57) and also in those lines (*Rsch, Col-0* and *Ler*), which

have no intronic or other SNP differences. The CBFs, or *C-REPEAT BINDING FACTORs* are a

set of 3 transcription factors that regulate cold acclimation and segregate for loss of function

variation at high frequency in *Arabidopsis thaliana*, with loss of function variants associated

with warmer climates (45,58). Hannah et al. (2006) previously showed with our dataset that

intact CBF function explained a substantial portion of variation in cold response transcriptomes

and Des Marais et al. (2017) showed that genetically variable cold responses were enriched in

central locations of coexpression networks.

To determine the role that genetic variation among genotypes, versus only among genes within a

genotype, we trained models with only *Col-0* genes as the set. We repeated the training while

also oversampling the dataset to be the same size as for all five accessions combined to control

for underpowered training due to the smaller dataset.

To assess whether models may have been hindered by factors such as model architecture or data

complexity we introduced a "spike" of 5bp in the upstream region of up-regulated DEGs. This

way we created a positive control that can help validate the capability of the models to recognize

the up regulated DEG group. We replaced the first 5bp of the up-regulated DEGs with the

sequence AAGGG. Then, we evaluated each trained model in two dimensions; both based on

their accuracy of predicting the evaluated set, but also based on their accuracy when predicting

non spiked data.

Evaluating and interpreting model training

We compared the area under the precision-recall curve (prAUC), cluster accuracy, as well as the

difference of the two classes accuracy with a Kolmogorov Smirnov test using the function

'ks.test' in R v4.1.2. To see which hyperparameters had a significant influence on those CNN

performance metrics of the spiked and non-spiked dataset, we ran a generalized linear mixed

model with all the hyperparameters as fixed effect and the models' loss as random effect. We

dropped from the model all the fixed effects that had no significant impact on the prAUC.

Thusly, the final model for the absolute difference between class prediction accuracy was(60)

(61)

$$\text{diff} \sim \text{cl} * \text{dl} + \text{pl} + \text{du} + (1|\text{fl})$$

where diff refers to the absolute difference between class prediction accuracy, cl stands for

number of convolutional layers, dl for number of dense layers, pl is the pool stride, du dense

units as fixed effects and fl the first convolutional layers filter. The residuals followed a Gamma

distribution. The final model for the prAUC of the test set was

273   diff ~ cl * dl * pl + dr + (1|fl)

with dr being the dropout rate, and the residuals' distribution was binomial. We tested the same

models in the original complete dataset. Generalized linear mixed models were performed using

the package lme4's function glmer (60) and *p* values were extracted using the function Anova

from the R package car (61).

For each type of CNN training we selected the model with the best performance during the

testing phase in order to understand the architectures that could offer the best prediction of both

classes .We selected models that had a prAUC value on the test set of at least 0.7, and that they

did not overfit into predicting one class of DEGs. To do so, we estimated the absolute difference

of the accuracy of predicting each class in the dataset. We used the same criteria for all cases.

Once we had selected the model parameters that comprise the architecture which best predicted

the DEG class, we evaluated the significance of the result and excluded the possibility that is the

result of learning from spurious sequences by permuting the label identity of each gene 100

times and re-train using the same model architecture. We estimated the *p* value by counting the

cases that had better performance based on both the prAUC and the class accuracy difference

and divided by 100.

All results were visualized in R v4.1.2 (59).

Results

The presence of enriched motifs in the up regulated DE gene sequences suggests their

potential contribution to environmental response.

We searched for motifs enriched along the putative regulatory regions of the up-regulated DEGs

in comparison to the non-DEGs to assess the possibility of patterns in the regulatory regions

associating with the differential expression status of each gene. We used STREME (49) to search

for de novo enriched motifs within the upstream and downstream regions of all genes of interest

(Figure 1a). When we searched for enriched motifs only within the sequences of up-regulated

DEGs for the reference genotype, *Col-0,* we discovered 9 motifs. In contrast, we identified 14

motifs significantly enriched within the up-regulated DEGs of all 5 accessions, suggesting

greater power with more genotypes ($p < 0.05$). The median discovered motif length was 10bp

(S3 Table) and the majority of the discovered motifs were significantly enriched in high

frequency in our dataset. Specifically, five of the fourteen discovered motifs were present in

more than 98% of the DEGs' regulatory regions tested, even though a few rare ones with less

than 1% frequency were also identified (Figure 1b). When we looked at which genotypes each

discovered motif was significantly enriched for at least once, the average number of motifs per

up-regulated DEG was similar between the genotypes (approximately 6 motifs per gene per

genotype; Figure 1c).

There was a similar number of enriched motifs among the down-regulated DEGs. In total 15

motifs were discovered that were significantly enriched in all genotypes ($p < 0.05$; S4 Table).

Five of those motifs were present in more than 99% of tested DEG sequences for enrichment

(Figure 1d), with an average of 6 discovered motifs per tested DEG sequence (Figure 1e).

Figure 1: Significantly enriched motifs within the regulatory regions of the up- and down-

regulated DEGs. **a.** For each up- or down- regulated gene across all five *A. thaliana* accessions,

we extracted the 500 bp upstream (blue block) and downstream regions (grey block). We also

extracted within the first 500 bp after the transcriptional starting site and 500 bp before the

transcriptional ending (yellow blocks) for a total of 1k bp per gene to include in the motif search.

b,d. The discovered motifs in the upstream and downstream regions of up- or down- regulated

DEGs and how frequent they are in the DEG sequences with STREME. **c,e** The distribution of

discovered motifs in the upstream region of up- DEGs or down- DEGs per accession.

When we cross referenced all the motif sequences we identified as enriched in up- and down-regulated DEGs in a database of experimentally defined TFBSs (50), the discovered motifs had a significant overlap with 135 and 143 known TFBS, respectively ($p < 0.05$). The most common TFBS across the discovered motifs were members of the DOF family, with CDF5, DOF1.7 and DOF5.1 binding sites being present in 3 and 4 of the total discovered motifs in the up-regulated and down-regulated DEGs respectively. The most common discovered sequences overlapping with the members of the DOF family were ARAACAGA in up-regulated and CAAAAAAAA in down-regulated DEGs. Members of the DOF family of transcription factors have been characterized in being involved in cold response(62). Moreover, *PISTILLATA* (PI), member of the MADS-box factors family MIKC, and the C2H2 zinc finger factor SGR5 binding sites, were present in 3 and 4 of the discovered motifs in the up -regulated DEGs, respectively. There was an overlap of the most common TFBS across the discovered motifs of the up-regulated DEGs and the down-regulated DEGs (Table 1). Therefore, within the promoter regions of the up- and down-regulated DEGs, there are motifs present that could be potentially informative for classifying the gene expression.

Table 1: The top known transcription factor binding sites (TFBS) that significantly overlap with the enriched discovered motifs along the upstream and downstream regions of up and down-regulated regions of up and down DEGs.

TFBS	Number of discovered motifs	Gene	Family
MA1159.1	4	SGR5	C2H2 zinc finger factors (class)

MA1268.1	3	CDF5	DOF
MA0556.1	3	DOF1.7	DOF
MA0558.1	3	FLC	MIKC
MA0559.1	3	(PISTILLATA) PI	MIKC
MA0940.1	3	AP1	MADS box factors (AP1)
MA1085.1	3	WRKY40	WRKY (class)
MA1281.1	3	DOF5.1	DOF
MA1367.1	3	AT1G76870	Trihelix
MA1380.1	3	TCX6	CPP
MA1182.1	5	RVE8	Myb-related
MA1184.1	5	RVE1	Myb-related
MA1380.1	5	TCX6	CPP
MA1277.1	4	DOF1.7	DOF
MA1281.1	4	DOF5.1	DOF
MA1268.1	4	CDF5	DOF
MA1278.1	4	DOF3.4	DOF
MA1279.1	4	DOF1.5	DOF
MA1190.1	4	RVE5	Myb-related
MA0933.1	4	AHL20	HMGA factors
MA1267.1	4	DOF5.8	DOF

Known transcription factor binding sites within *Col-0* do not accurately predict the expression response to cold using random forests.

Given the many known and documented transcription factor binding site motifs that exist for different transcription factors (many of which are known to be involved in cold response) and given that we found TFBS motifs in the cold DEGs, we asked if these TFBS could predict genome-wide expression response to cold. We used a dataset of known TFBS in putative promoters 1kbp upstream of genes in the *A. thaliana* accession *Col-0* (52) to test whether they can predict DEGs using random forests classification trees (S1 Figure). For this analysis, we also split the genes based on up-regulated and down-regulated and compared each group to all non-DEGs. Even though both random forests models correctly predicted the gene regulation status for approximately 80% of the genes, the prAUC of each of those models was low. The probability AUC-ROC (prAUC) for predicting up-regulated DEGs was 0.5013 (permutation based p = 0.17) and the prAUC for predicting down-regulated DEGs was 0.499 (permutation based *p* = 0.22). Therefore, RF models based on known TFBS involved in temperature related reactions were not able to predict DEGs with high accuracy within this dataset. This, taken together with the various discovered motifs within the upstream and downstream regions of the genes, indicates that a method that incorporates more genetic information within the code might improve the prediction of regulatory status better within specific conditions.

Predicting gene expression regulation based on regulatory regions using naive methods is possible

We next investigated whether a currently commonly used machine learning method could learn

cis regulatory motifs simply from sequence and DE data alone. If *de novo* motifs can be

identified in DE genes, even though known TFBS were not good predictors of expression in the

RF model, we might obtain better predictions learning *de novo* important motifs. To this end, we

trained convolutional neural networks (CNNs) that incorporate the upstream and downstream

genetic sequence as input layer via one hot encoding (Figure 2a; see Material and Methods). We

trained 1,344 models with different combinations of the hidden layer's parameters to find the

model architecture that best predicts the genes DE (S2 Table). We trained the models using the

information from all 5 accessions that had available pseudogenomes to predict up- or down-

regulated DEGs separately.

Figure 2: Training Convolutional Neural Networks to predict up-regulated or down-regulated DEGs between specific environments. a. Description of the different type of input data used to train CNNs to predict up- or down-regulated DEGs vs non DEGs. The analysis "up-regulated DEGs" included the five *A.thaliana* accessions and both upstream and downstream regions for up-regulated DEGs. The analysis "spiked upstream" included training on up-regulated DEGs with the first 5bp being replaced consistently with the same sequence. "With active CBFs" refers to the training of models using the two accessions with all *3 CBF* genes active. The analysis "upstream regions" included only the upstream regions of the up-regulated DEGs as input. "Only *Col-0*" was used to train models, as well as when it was five times oversampled ("Col-0x5"). Finally, the results under the category "down regulated DEGs" show the outcome of training in all **b.** prAUC values of the test set in the different analysis. **c.** The difference in accurately predicting the up/down-regulated DEGs to the non DEGs in all different analysis. **d.** The within class prediction accuracy of down- and up- regulated DEGs in the test set in all analysis. The within class prediction accuracy was estimated as the proportion of correctly predicted genes of this class over the total number of genes in this class in the test set. **e**. The within non-DEGs prediction accuracy of the test set in all analysis.

The tested CNN model architectures have had a wide variety of accuracy in predicting the DEG of each gene (S5 Table). We extracted the prAUC of each model during training and moreover, we estimated the prAUC of the test set for each trained CNN model. The median and mean prAUC values during training for predicting up-regulated DEGs vs non-DEGs were 0.5013 and 0.5017, respectively, indicating that many tuning parameter settings did not lead to substantial learning. We also estimated the median within class accuracy in the test set as 0.776 and 0.227 for non DEGs and up-regulated DEGs, respectively. Note that the test set is not over sampled, but it is the original ratio which is mostly non-DEGs. Overall, the median prAUC in the test set of the models was 0.229, with a mean of 0.354 (S2 Figure). We identified 15.6% of the models with prAUC in the test set values above 0.80.

To gain insight into whether tuning parameters had any consistent effects on model learning, we used a generalized linear mixed on the prAUC of the test set using the first layer's filters as

random effect and the number of convolutional and dense layers, with pool stride and dropout

rate as fixed effects. Both the number of convolutional and dense layers had a significant

negative impact on the prAUC of the test set, with $p = 0.000754$ and $p = 3.9708e-16$,

respectively, suggesting more complex models were more poorly trained. Additionally, the pool

stride ($p = 2.918e-10$), which indicates the number of base pairs that are considered together in

each filter, and the dropout rate, or how many model nodes are used each time to predict, had a

significant negative impact of the prAUC of the test set. However, none of the interactions

between the fixed effects were significant, suggesting little importance of specific combinations

of the tuning parameters on learning.

The trained models for predicting down-regulated DEG or non-DEGs (S3 Figure) were very

similar with the results observed for the up-regulated DEGs vs non-DEGs. The median prAUC

of the models during training was 0.5009 and the mean prAUC was 0.5012. The median prAUC

of the test set for the down-regulated DEGs vs non DEGs was 0.253 and the mean 0.363 (S6

Table). The same hyperparameters as for predicting up-regulated DEGs were identified to have a

significant negative effect with *p-values* of 1.511e-07, 0.0168, 4.068e-08 and 0.0485 for number

of convolutional layers, number of dense layers, pool stride and dropout respectively.

We identified the best models that could predict the up-regulated DEGs. For that purpose, we

selected a model that had prAUC in the test set above 0.7. Moreover, in order to understand the

models' potential to predict each class separately we estimated for each class a prediction

accuracy as the number of correct predictions of the class divided by the total number of the

genes in the class tested. So, after this prAUC cutoff, we picked a model whose prediction

accuracy difference was the lowest. The best model for predicting up regulated DEGs had a

prAUC in the test set of 0.701, and prediction accuracy in the test set was 0.368 for up regulated

and 0.693 for non DEG (Table 2). The class prediction accuracy difference was 0.325. The best

model for predicting up-regulated DEGs had 2 convolutional layers and 2 dense layers, a dropout

rate of 0.25 and pool stride 4. Using the same criteria, we were able to identify the best model in

predicting the down-regulated DEGs. The prAUC of the test set for this model is 0.7128 and the

difference in the class prediction accuracy is 0.319, with per class accuracy being 0.365 for

down-regulated DEGs and 0.665 for non-DEGs. The model had 3 convolutional and 2 dense

layers, pool stride of 8 and dropout rate of 0.25.

We assessed whether these models performed better than randomness by permuting the input

labels and re-train the selected model 100 times (S4 Figure), so as to understand to what degree

the input labels are used for tuning by the CNN architecture. For each permutation we estimated

the prAUC of the test set and the prediction accuracy between the two predicted classes using the

un-permuted test set. Then, we compared the values of the original trained model against the

distribution of the metrics estimated with the permuted labels. The significantly higher prAUC

for the up-regulated DEGs ($p < 2.2e\text{-}16$) and down-regulated DEGs ($p < 2.2e\text{-}16$) than the

permuted dataset indicate that our picked models perform better than randomly picking genes.

Table 2: The best model for each training performed are given. For each type of analysis run
(Figure 2), we selected the best performing model based on the prAUC value of the test set and
the the difference in accurately predicting the DEGs vs non DEGs. For each model, the model
parameters, the training, validation and testing prAUC, loss and accuracy of predicting both
DEG classes is given.

	Up regulated DEGs	Down regulated DEGs	Spiked up regulated DEGs	Upstream regions of up DEGs	Up regulated DEGs of only	Up regulated DEGs of upsamp	Genotypes with all *CBF* as

					Col-0	led Col-0	DEGs.
1st conv. filters	128	64	128	64	128	128	128
2nd conv. filters	128	64	None	128	None	128	128
3rd conv filters	None	128	None	128	None	None	64
Conv width	4	4	4	4	4	8	8
Pool width	4	4	8	4	4	4	4
Pool stride	4	8	4	4	8	4	8
Dropout	0.25	0.25	0.25	0.1	0.1	0.1	0.25
1st dense layer units	64	128	128	128	128	128	64
2nd dense units	128	None	128	None	128	128	None
Number of conv. layers	2	3	1	3	1	2	3
Number of dense layers	3	2	3	2	3	3	2
Learning rate	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
Loss	0.692	0.691	0.691	0.691	0.691	0.692	0.691
Accuracy	0.621	0.682	0.673	0.666	0.692	0.636	0.672
Accuracy of DEGs	0.368	0.365	0.480	0.339	0.347	0.368	0.341
Accuracy of non DEGs	0.693	0.684	0.665	0.671	0.697	0.681	0.681
prAUC test set	0.701	0.712	0.704	0.701	0.7207444	0.703	0.711
prAUC training	0.499	0.502	0.522	0.499	0.4971713	0.505	0.499

prAUC of validation set	0.501	0.502	0.578	0.509	0.4851478	0.517	0.506
Loss of the validation set	0.693	0.693	0.692	0.693	0.693379	0.693	0.693
Epoch of best model	37	50	49	29	49	48	49
Difference in accuracy of DEG classes	0.325	0.319	0.185	0.332	0.3495906	0.312	0.340

Number of discovered motifs can impact the correct prediction of up-regulated DEGs

We investigated the profile of the genes that are accurately predicted within the up regulated

DEGs by the selected model with the highest achieved performance. Out of the 4,564 up DEGs

across all five genotypes in the test set, 592 were accurately predicted. The correctly predicted

genes had a significantly higher (wilcoxon rank sum test $p = 0.018$) average number of

discovered motifs in their regulatory regions (median of 12 motifs) by the STREME analysis

than the genes that were incorrectly predicted (mean of 10; S5a Figure). Interestingly, when we

repeated this analysis for the down-regulated DEGs, we did not detect a different accumulation

of TFBS within the correctly predicted genes. Both the correctly and wrongly predicted genes

had an median of 11 motifs per gene and the accumulation of the TFBS upstream and

downstream of the regions was not significantly different (wilcox sum rank test $p = 0.9446$; S5b

Figure). Moreover, the gene expression level of the genes that were correctly predicted vs not

correctly predicted was not significantly different (based on Kolmogorov Smirnov test $p > 0.05$)

for both down-regulated DEGs and up-regulated DEGs.

Up until this point, the discovered motif diversity, the CNN training results, and even the best

model's behavior was very similar for up-regulated DEGs versus down-regulated DEGs. For the

rest of the analyses, we focused on training CNN models for predicting up regulated DEGs

versus non DEG, as we do not expect that altering the input of the CNNs training will alter the

observed pattern between the two regulatory groups.

CNNs imperfectly identify a simple artificial signal within the regulatory regions.

Because many tuning parameter values resulted in models that learned little (training prAUC

near 0.5), we sought to assess potential sources of constraints on these models. We investigated

whether the tested model architectures can learn a simple motif within the sequences that

perfectly predicts DEGs, a motif we refer to as a "spike". If these spiked models are perfectly

able to predict DEGs in this setting, it suggests that the limitations we observed with real data

arise from a lack of such good, simple predictors in the sequence. If models have limited learning

in this setting, it suggests that the model architecture is limited in its ability to find signals in the

data were using, perhaps due to the complexity and abundance of sequence data (2,000 bp per

gene).

To test these hypotheses, we included a spike sequence of 5bp in the place of the first 5bp of all

up-regulated DEGs (Figure 2a). The spike's sequence (AAGGG) does not overlap with the

sequence of any of the discovered enriched sequences with the up DEGs. The expectation was

that the accuracy of the models would increase as they have a perfect sequence to use for

predicting the up-DEG group. During training the prAUC values had median of 0.5018 and mean

of 0.5046. The median prAUC in the test set of the spiked models was 0.36, with a mean of

0.266, which was on average higher than when trained without the spikes (S7 Table). The

distributions of the prAUC values in the test set between the spiked and non-spiked models were

significantly different (Kolmogorov Smirnoff test, $p < 2.2e\text{-}16$; Figure 2b). Generally, the spiked

models had fewer overfitted models than the non-spiked models for both up-regulated and non-

DEG classes (Figure 2c; S6 Figure). In total 188 spiked models with prAUC values of the test set

above 0.8 were detected and 561 models with prAUC values of the test set below 0.2, and thus,

in total, 55.7% of the models tested are overfitted to one of the two classes. In contrast, among

the non-spiked models, 61.5% of the models are overfitted.

When we used the same criteria to pick the best performing models for the spiked data

(permutation $p < 2.2e\text{-}16$) as for the non-spiked data, we identify a model that predicts the two

classes better than the non-spiked best model. prAUC of the test set (0.704) is only marginally

higher by 0.03. In contrast, the difference in the prediction accuracy of the two classes in the test

set is 0.185, which is 1.75 times lower than the best model predicting non-spiked up-regulated

DEGs. This improvement is due to the higher accurate prediction of the up-regulated DEGs

(0.48) in the spiked best model than the non-spiked best model (0.36).

There is no significant correlation between the prAUC values in the test set of the spiked and

non-spiked models under the same values of tuning parameters ($p=0.96$, $\rho =0.0012$), indicating

that the inclusion of the spikes did not improve models with good tuning parameters for the real

data, but instead were best captured with distinct tuning parameters (Figure 3). When we further

investigated the different hyperparameters' impact on the prediction accuracy of each class, we

noticed that the spiked models with 1 convolutional layer had higher accuracy and prAUC in the

test set and simultaneously smaller difference between the accurate prediction of both classes

than the models with 2 and 3 convolutional layers. (Figure 3a). The fact that the models with less

convolutional layers performing better that with more of them was also true for the spiked

models. A generalized mixed model including the number of convolutional layers, dense layers,

pool stride and dense units as fixed effects indicated that the number of convolutional layers had

a significant effect on the difference of the prediction accuracy of the two classes (p = 8.091e-

10). Fewer layers were predicting more accurately, as the model's estimates were negative. This

observed pattern is not as extreme as for the non-spiked models (p = 0.01023; Figure 3a). Other

parameters that were found to have a significant negative impact on the spiked models'

difference in the prediction accuracy of the two classes were the number of dense layers (p =

9.036e-05), the pool stride (p = 0.002137) , the first dense layer's units (p=0.048), as well as the

interaction of the number of convolutional and dense layers (p = 0.00983). All these factors,

except for the interaction of the number of dense and convolutional layers, had a significant

effect on the difference between the accuracy of the two classes for the non-spiked models.

Specifically, the number of dense layers, the pool stride, and the number of convolutional layers

had p values of 6.191e-06 and 0.0102 respectively (Figure 3b-c). The estimates for all

parameters were negative in both spiked and non-spiked data, which means that generally

models with fewer layers were able to predict with higher per class accuracy in the test set. This

result suggests that although different specific sets of tuning parameters for good for spiked data

versus the non-spiked, the general effect of tuning parameters, and specifically model

complexity, was consistent. Therefore, a consistent signal within the up-regulated DEGs can

improve the model performance, indicating that the regulatory complexity among genotypes is

quite high, that we lack information in the sequence data for prediction, or that the DEG response

is highly stochastic.

Figure 3: Simple spiked models can accurately predict an introduced spike of 5bp at the beginning of the up-regulated DEGs promoter region. Comparison of the prAUC in the test set and the difference between the accurate prediction of the two classes for non- spiked (left) and spiked (right) input data. Comparisons are done for a. number of convolutional layers, b. number of dense layers and c. pool stride. The p values stand for the significance level of the KS test when comparing the two distributions shown within each plot.

Predicting only among genotypes sharing alleles for major *trans* eQTL does not improve the prediction of the up-regulated DEGs.

Since the results with the sequence spike indicate some data complexity limited our CNN predictions, we investigated whether we could simplify the prediction task by homogenizing known large-effect *trans* mutations (CBFs). These transcription factors show variation in functionality among genotypes, and in the other natural genotypes tested at least some copies are non-functional. We approached this analysis in two ways. First, we trained models predicting up-regulated DEGs vs non DEGs in *Rsch, Cvi* and *Ler,* which do not have intronic SNP differences between them in the two accessions that have   all three major well-studied cold-response *CBF* transcription factors, including *CBF2*. Then, we trained models to predict up-regulated vs non DEGs on *Can* and *Cvi,* which have differential expression of all 3 *CBFs* during the experiment.

The median and mean prAUC during training on the set with no SNP differences, was 0.5001 and 0.5 respectively, while the prAUC of the test set had mean value of 0.47 and median 0.095 (S7 Figure; S8 Table). This is also reflected in the percentage of model architectures, which have been overfitted for predicting the upregulated DEGs; 52.5%. of the models in contrast to the 45.5% overfitted in predicting the non DEGs, with only 1.7% of the models not showing any sign of overfitting. The best model trained on only the three accessions with similar copies of *CBF* functional had prAUC in the test set of 0.705 and difference in predicting accuracy of the

two DEG groups of 0.47. Similarly, the median prAUC during training on the set with

differentially expressed all 3 *CBFs*, was 0.5014, while the prAUC of the test set had median

value of 0.268 (S7 Figure; S8 Table). The difference in the accurate prediction of the up-

regulated DEGs and non DEGs within the test set has a median of 0.835 and mean 0.701, which

is on average higher than the difference in prediction accuracy of the models applied to all

genotypes. The best model had prAUC in the test set of 0.7 and difference in predicting accuracy

of the two DEG groups of 0.31. Overall, those models did not perform better than the original

models including all the *A. thaliana* accessions. This suggests that the lack of a major *trans*

acting expression variant did not hinder predictability in our bigger set of diverse genotypes here.

Including both upstream and downstream regions improves the prediction of up-regulated

DEG class

As often the *cis* elements are mostly identified in the upstream regions (64), we tested whether

including the downstream coding region has limited the prediction accuracy, perhaps by

incorporating noise. Noise might be introduced by incorporating long sequences that do not

usually have a lot of TFBS, such as the downstream regions. Therefore, we retrained all the

models by including only the 2kbp upstream regions for all 5 accessions, compared to the 1 kb

up and 1kb downstream of the coding region in the approach presented above (Figure 2a). The

median prAUC of the training set was 0.5 and the median prAUC (0.307) in the test set of the

models trained only using the 2kb upstream sequences was slightly higher than including both

the 1kb upstream and 1kb downstream regions, by 0.048 (S8 Figure; S9 Table). The distribution

of the prAUC results in the test set of the two analysis was again significantly different (KS test

$p < 2.2e-16$). The within class prediction accuracy of up-regulated DEGs (0.671) in the test set

was higher than for non-DEGs (0.328). Moreover, 29.8% of the models had prAUC values in the

test set above 0.8. The same pattern was observed for models that had low prAUC in the test set.

Approximately 41% of the models had prAUC in the test set below 0.2, indicating overfitting to

predicting all genes as up-regulated DEGs, to a total of 71.1% of the models overfitting to either

class. In comparison, including both upstream and downstream regions leads to overfitting of

61.5% of the models. (Figure 2c, Figure 2d). Therefore, excluding the downstream regions from

the analysis leads to more cases when the models get fixed in recognizing only one class.

The best model in predicting up-regulated DEGs and non DEGs trained only on putative

promoter regions had 3 convolutional layers, 3 dense layers and a dropout of 0.1 (Table 2). The

prAUC in the test set was 0.7004 and the difference in accurately predicting up-regulated DEGs

and no DEGs is 0.332, which is a similar performance to the best model learning using both

upstream and downstream regions (Table 2). Its performance is better than models with the same

architecture trained on random input ($p < 2.2e\text{-}16$). Therefore, including only the 2kb upstream

regulatory regions of the genes during training does not significantly alter the predictive ability

compared to including 1kb upstream and 1kb downstream.

Genetic diversity does not improve accuracy of differentially expressed genes.

Genetic diversity is an important dimension of natural populations. Incorporating multiple

genotypes in the STREME analysis has had a positive impact into discovering more enriched

motifs. If we consider the introduced spike in the upstream region of up-regulated DEGs as a

large effect mutations due to its 100% frequency within the group, we also showed that the

majority of the information that the models learn is based on small effect mutations. Small effect

mutations can be better identified when incorporating more genetic diversity into any kind of

analysis, as the less frequent variants are easier to identify. We finally compared how effective

CNNs were when including multiple genotypes in the training set, by training all the different

model architectures including only the most commonly studied *A. thaliana* accession, *Col-0*

(Figure 2a). We repeated this twice, once that had the original number of genes in *Col-0* and

once that we up sampled the *Col-0* genes enough times to have the same size of training set as

for the complete set of the genotypes. In both cases, we oversampled the up regulated DEGs to a

1:1 ratio with the no DEGs. The training of the non-oversampled and oversampled sets wielded

median prAUC values of 0.5004 and 0.501 respectively.

When compared to the models that have been trained on the *Col-0* dataset only, we see that the

median AUC of the test set is higher, as it is at 0.488, for the multiple genotypes model (Figure

2b, S9 Figure; S10 Table). The accuracy was significantly impacted by the number of replicates,

as the models trained on the up sampled *Col-0* set had median prAUC of the test 0.508 (KS test

p = 8.8e-16; S11 Table; S10 Figure). During training, both datasets had a high proportion of

overfitted models. Within the up sampled dataset, 50.3% of the prAUC values in the test set were

either above 0.8 or below 0.2. Training on the up-sampled dataset of *Col-0* had fewer overfitted

models than training on the non-oversampled dataset of *Col-0,* in which 77.4% of the models

were overfitting in either direction. The better performance of the oversampled dataset of *Col-0*

is due to the more appropriate size of the dataset; the original *Col-0* dataset has only 34,002 data

points when the up-regulated DEGs are oversampled to 1:1 ratio (65,66). Based on this

observation, below we compare only the models trained on the oversampled dataset of *Col-0* to

the models trained on all five accessions.

The models trained on the *Col-0* dataset have fewer overfitted models (50.3%) than the models

trained on all five accessions (61.5%). The best model in predicting both up-regulated and non-

DEGs on the *Col-0* dataset has similar architecture and performance statistics as the best model

trained on all five accessions. The prAUC value in the test set trained on the oversampled *Col-0*

was 0.7034 and the difference in the accurate prediction between the two classes was 0.3122. It

performed better than when trained on a permuted dataset ($p < 2.2e-16$). Both the prAUC in the

test set (0.7008) and the difference in accurate prediction of both classes (0.3252) indicate that

training on all five accessions was performing not as well as the model trained on the *Col-0*

dataset only. Therefore, including more accessions may either marginally increase the noise and

not improve the accuracy of the models by a lot, or not be variable enough to capture more

contrasts between alleles rather than genes.

Finally, because the models of cold response had only modest success, we tried to test if the

CNN models could accurately predict DEGs in a simpler transcriptomic change (1,282 DEGs),

based on a single transcription factor knockout known to control anthocyanin biosynthesis.

However, even these CNN predictions had modest success (Supplemental Material), highlighting

the challenge in predicting transcriptomic dynamics from proximate DNA sequences.

Discussion

We used the genomic sequence of upstream and downstream regions of *Arabidopsis thaliana*

genes to identify motifs related to the differential expression in cold and control conditions and

also to predict the gene expression response to an abiotic stressor. Many *cis* regulatory elements

have been documented (13,38,63) and they are often involved in gene evolution and adaptation

to novel environments (4,67). Here, within both the up-regulated and down-regulated DEGs we

found motifs of variable length that are enriched within them in comparison to the non-DEGs.

The discovered motifs in *cis* regulatory regions significantly overlap with known transcription

factor binding sites that are involved in the regulation of gene expression under cold stress

(42,62,68). Knowledge of specific transcription factor binding sites can aid the prediction of

gene expression under different types of abiotic and biotic stress (26,33,69,70). The five

accessions used in this study are regulated by differential function of *CBF* and *ZAT12* regulons,

leading to their variable acclimation capacity to cold (42). The three *CBF* genes control the

acclimation to cold conditions and reaction to freezing stress (58,71). It has been shown that

genetic diversity within the gene and its network is connected to adaptation to different

temperatures and latitudes (9,45,72). We identified a frequent overlap with known TFBS, which

are known to be involved in controlling responses to cold or freezing stress such as the RVEs

(73), members of the DOF family of transcription factors (58,62,74,75) and FLC (76). We also

identified frequent binding sites for members of the DREAM complex, which are involved in

repressing growth in response to DNA damage (77), as well as SGR5, which is differentially

spliced under heat stress (78). This indicates the possibility to predict gene expression regulation

based not only on factors with biological meaning, but also with genetic diversity, as the

incorporation of more accessions than *Col-0*, yielded more discovered motif sequences.

However, as often the available information to the researcher is only DNA and mRNA sequence

(and not experimentally verified transcription factor binding sites), we investigated the

possibility of using the genetic sequence to predict gene expression. The need for this was also

evident by trying to predict up- or down-regulated DEGs using random forest analysis. The

random forest used known presence or absence of specific TFBS upstream and downstream of

the genes to predict DEGs. However, neither for up-regulated DEGs nor for down-regulated

DEGs, the prediction accuracy was better than chance. In *A. thaliana,* using known transcription

factor binding sites has not yielded very good prediction of differential gene expression (26),

while the most enriched *cis* regulatory elements did not also predict the most differentially

expressed genes in maize (32). Training CNNs, with the 2kbp upstream and downstream regions

of the genes wielded better prediction accuracy. The model that had both high accuracy and had

the least difference in predicting accuracy per class had prAUC in the test set of 0.7. This value

is slightly lower than what has been reported in similar studies in other species, which identified

models with prAUC between 75% to 85% (55,79). In our case, higher prAUC values showed

signs of overfitting; non DEGs, which comprise more than 80% of the dataset were then almost

exclusively correctly predicted, inflating the accuracy and prAUC. Surprisingly, the inclusion of

both upstream and downstream regulatory regions did not improve the prediction accuracy of

DEGs, as in maize (80). This could be attributed to genomic differences between the species; for

instance downstream regulatory regions have differences in nucleotide composition between

maize and *A. thaliana* (81). Additionally, maize has a different genomic composition than *A.*

thaliana, such as higher transcription element density, with 85% in contrast to 21% in *A.*

thaliana. Authors have suggested the expanded genome of maize could enhance the landscape

and availability of *cis* regulatory adaptations (82). Factors such as this can explain the difficulty

of successfully transferring machine learning protocols generated from one species to another.

Phenotypic responses to environment (plasticity) can be adaptive, which are ultimately fixed by

populations in the specific environment (85). Phenotypic plasticity can also be genetically

variable (i.e. GxE), with gene expression being an important example. Our hypothesis was that

incorporating more genotypes in this study would improve the prediction accuracy as more

genetic variation in *cis* sequences and expression would be incorporated during training.

Incorporating genetic variability between species has been successfully used to train models with

simple architecture to predict gene network relationships (86). Within maize, when more than

one genotype was used for predicting gene expression, then the results were more accurate (33).

We did not observe the same pattern here; in fact an oversampled dataset of *Col-0* upstream and

downstream regions to the same size as the dataset including the multiple genotypes, yielded a

model with higher prAUC and more comparable prediction of the two DEG classes. We do not

believe that this is simply an issue of power due to the dataset being on the smaller side of what

is required for machine learning. A potential contributing factor could be that the *cis* -regulatory

genetic variation might not be of large enough frequency between genotypes but is stronger

between genes. Additionally, the highest prAUC in the test set was achieved when the accessions

with differentially expressed all 3 *CBF* genes were only included during training, suggesting

trans effects in high frequency could limit our predictions across multiple genotypes. This

pattern was confirmed when a high frequency motif was incorporated in the upstream regions of

the up-regulated DEGs; the spiked dataset could be used to train models with higher accuracy.

Within *A. thaliana,* the CBF/DREBs genes of the cold signaling pathway have been linked with

cold adaptation within Europe (44,87) as the major eQTLs controlling acclimation. Taken

together with the fact that more than half of the enriched discovered motifs within the up- and

down-regulated DEGs are present in less than 50% of the genes across all genotypes and thus not

very frequent, it indicates that the large number of *cis* regulatory variants across the genome in

low frequency can decrease the accuracy in prediction of gene expression differences, in contrast

to the presence of a potential large effect eQTL controlling the response to cold. The putative

promoter region likely captures the many *cis*-regulatory elements, which have been related to

large impacts on gene expression divergence and polygenic selection but are often each

individually mutations of small effect (63,88). Incorporating a spike within the upstream region,

which represents a single factor of large effect, supports this hypothesis, as in overall the spike

improved the prediction accuracy of the models. Our findings are consistent with machine

learning predictions of protein folding, where models that accurately predict variation among

genes perform worse on predicting effects of individual amino acid substitutions (89), a task that

was better suited to a distinct prediction approach (90). During training, the models were able to

detect the spike and increase the accuracy of the up-regulated DEGs. Therefore, even though

including multiple accessions could slightly increase the accuracy of predicting GxE, care has to

be taken to which accessions are selected for the training.

One explanation for the remaining variation in expression unpredicted by CNNs in this study is

that these models overlook much of the biological complexity driving gene expression response

to environments. Genes are expressed as a part of complicated regulatory networks that *cis*

regulatory elements only partially capture. We have currently ignored epigenetic states and the

sequence of the coding region. In a sorghum species, it has been more successful to predict

DEGs as response to cold as models incorporated information about the coding region sequence

and methylation information (79). Alongside specific transcription factor binding sites, their

positions on the genome and the chromatin state have been useful to predict the transcription

factor's target genes, with information that is also transferable with high accuracy between

datasets (84). Prediction accuracy of gene expression differences and the impact of *cis* variants

were improved when large up- and downstream regions, as well as when open methylated

regions were included in the analysis (32,33). Trained bimodal CNNs based on chromatin state

and DNA sequence for different cell types can differentially predict the transcription factor

binding specificity, which performs better than neural networks based only on DNA sequence

information (83). During training, therefore, using only *cis* regulatory regions to predict gene

expression response to environment within *A. thaliana* may be of limited accuracy without

incorporating additional data types. However, when that biological information is missing and it

is challenging to acquire, as would be the case in many non-model systems, the accuracy of

imperfect CNN models might still be of use to the evolutionary researcher.

Interpretation of what might have been having an impact on the training could be difficult to

understand. Even models with the best performance do not necessarily have the best interpretable

structure of layers (91). Looking into the statistics of all the different models trained during each

grid search can give insights into this. First, the distribution of the prAUC during testing can give

clear indications of how easy it is to overfit to the model, and therefore how complex this

training task can be. Despite these limitations, using machine learning methods can enable

researchers to draw predictions and conclusions about reactions to specific environments within

species, albeit perhaps not CNN models. We believe that further development of other models

such as language models, which can use less biological information to predict gene expression

responses, would be useful for systems with less information about their (molecular) biology

than the model species.

Acknowledgments

The Pennsylvania State University ACI cluster and the HPC Ramses of University of Cologne

have provided us with computational support. We would like to thank S. Mahony and H.

Dittberner for providing valuable feedback in running convolutional neural network analysis and

M.G. Stetter, T.S. Winkler, A. Singh and J. Floret for feedback on the manuscript.

References

1. Kawecki TJ, Ebert D. Conceptual issues in local adaptation. Ecol Lett.
2004;7(12):1225–41.

2. Romero IG, Ruvinsky I, Gilad Y. Comparative studies of gene expression and the
evolution of gene regulation. Nat Rev Genet. 2012 Jul;13(7):505–16.

3. Yang Z, Xu G, Zhang Q, Obata T, Yang J. Genome-wide mediation analysis: an
empirical study to connect phenotype with genotype via intermediate
transcriptomic data in maize. Genetics. 2022 Apr 23;iyac057.

4. Josephs EB, Lee YW, Stinchcombe JR, Wright SI. Association mapping reveals the
role of purifying selection in the maintenance of genomic variation in gene
expression. Proc Natl Acad Sci. 2015 Dec 15;112(50):15390–5.

5. Josephs EB, Lee YW, Wood CW, Schoen DJ, Wright SI, Stinchcombe JR. The
Evolutionary Forces Shaping Cis- and Trans-Regulation of Gene Expression
within a Population of Outcrossing Plants. Mol Biol Evol. 2020 Aug 1;37(8):2386–
93.

6. Mack KL, Square TA, Zhao B, Miller CT, Fraser HB. Evolution of spatial and
temporal cis-regulatory divergence in sticklebacks. Mol Biol Evol. 2023 Feb
20;msad034.

7. Keagy J, Drummond CP, Gilbert KJ, Grozinger CM, Hamilton J, Hines HM, et al.
Landscape transcriptomics as a tool for addressing global change effects across
diverse species. Mol Ecol Resour [Internet]. 2023 [cited 2023 Apr 11];n/a(n/a).
Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/1755-
0998.13796

8. Huo H, Wei S, Bradford KJ. DELAY OF GERMINATION1 (DOG1) regulates both
seed dormancy and flowering time through microRNA pathways. Proc Natl Acad
Sci. 2016 Apr 12;113(15):E2199–206.

9. Mateos JL, Tilmes V, Madrigal P, Severing E, Richter R, Rijkenberg CWM, et al.
Divergence of regulatory networks governed by the orthologous transcrip-tion

factors FLC and PEP1 in Brassicaceae species. Proc Natl Acad Sci U S A.
2017;114(51):11037–46.

10. Thomashow MF. Molecular Basis of Plant Cold Acclimation: Insights Gained from
Studying the CBF Cold Response Pathway. Plant Physiol. 2010 Oct 1;154(2):571–
7.

11. Adrian J, Farrona S, Reimer JJ, Albani MC, Coupland G, Turck F. cis-Regulatory
Elements and Chromatin State Coordinately Control Temporal and Spatial
Expression of FLOWERING LOCUS T in Arabidopsis. Plant Cell. 2010 May
1;22(5):1425–40.

12. Cubillos FA, Stegle O, Grondin C, Canut M, Tisné S, Gy I, et al. Extensive cis-
Regulatory Variation Robust to Environmental Perturbation in Arabidopsis. Plant
Cell. 2014 Nov 1;26(11):4298–310.

13. Wittkopp PJ, Haerum BK, Clark AG. Evolutionary changes in cis and trans gene
regulation. Nature. 2004 Jul;430(6995):85–8.

14. Lovell JT, Schwartz S, Lowry DB, Shakirov EV, Bonnette JE, Weng X, et al.
Drought responsive gene expression regulatory divergence between upland and
lowland ecotypes of a perennial C4 grass. Genome Res. 2016 Apr 1;26(4):510–8.

15. Schmitz RJ, Grotewold E, Stam M. Cis-regulatory sequences in plants: their
importance, discovery, and future challenges. Plant Cell. 2021 Nov 22;koab281.

16. de Meaux J. An adaptive path through jungle DNA. Nat Genet. 2006
May;38(5):506–7.

17. Meaux J de. Cis-regulatory variation in plant genomes and the impact of natural
selection. Am J Bot. 2018;105(11):1788–91.

18. Erwin DH, Davidson EH. The evolution of hierarchical gene regulatory networks.
Nat Rev Genet. 2009 Feb;10(2):141–8.

19. Wray GA, Hahn MW, Abouheif E, Balhoff JP, Pizer M, Rockman MV, et al. The
Evolution of Transcriptional Regulation in Eukaryotes. Mol Biol Evol. 2003 Sep
1;20(9):1377–419.

20. Brown KE, Kelly JK. Genome-wide association mapping of transcriptome variation
in Mimulus guttatus indicates differing patterns of selection on cis- versus trans-
acting mutations. Genetics. 2022 Jan 1;220(1):iyab189.

21. Korfmann K, Gaggiotti OE, Fumagalli M. Deep Learning in Population Genetics.
Genome Biol Evol. 2023 Feb 1;15(2):evad008.

22. Raicu AM, Fay JC, Rohner N, Zeitlinger J, Arnosti DN. Off the deep end: What can
deep learning do for the gene expression field? J Biol Chem. 2023 Jan
1;299(1):102760.

23. Chen Y, Li Y, Narayan R, Subramanian A, Xie X. Gene expression inference with
deep learning. Bioinformatics. 2016 Jun 15;32(12):1832–9.

24. Giri A, Khaipho-Burch M, Buckler ES, Ramstein GP. Haplotype associated RNA
expression (HARE) improves prediction of complex traits in maize. PLOS Genet.
2021;17(10):e1009568.

25. Sartor RC, Noshay J, Springer NM, Briggs SP. Identification of the expressome by
machine learning on omics data. Proc Natl Acad Sci. 2019 Sep 3;116(36):18119–
25.

26. Moore BM, Lee YS, Wang P, Azodi C, Grotewold E, Shiu SH. Modeling temporal
and hormonal regulation of plant transcriptional response to wounding. Plant
Cell. 2021 Dec 3;koab287.

27. Schwarz B, Azodi CB, Shiu SH, Bauer P. Putative cis-Regulatory Elements Predict
Iron Deficiency Responses in Arabidopsis Roots1 [OPEN]. Plant Physiol. 2020
Mar 1;182(3):1420–39.

28. Uygun S, Azodi CB, Shiu SH. Cis-Regulatory Code for Predicting Plant Cell-Type
Transcriptional Response to High Salinity1 [OPEN]. Plant Physiol. 2019 Dec
1;181(4):1739–51.

29. Azodi CB, Lloyd JP, Shiu SH. The cis-regulatory codes of response to combined
heat and drought stress in Arabidopsis thaliana. NAR Genomics Bioinforma. 2020
Sep 1;2(3):lqaa049.

30. Marand AP, Eveland AL, Kaufmann K, Springer NM. cis-Regulatory Elements in
Plant Development, Adaptation, and Evolution. Annu Rev Plant Biol.
2023;74(1):111–37.

31. Wang H, Cimen E, Singh N, Buckler E. Deep learning for plant genomics and crop
improvement. Curr Opin Plant Biol. 2020 Apr 1;54:34–41.

32. Benoit M. Hot 'n cold: Applying the cis-regulatory code to predict heat and cold
stress response in maize. Plant Cell [Internet]. 2021 Nov 8 [cited 2021 Nov 11];
(koab269). Available from: https://doi.org/10.1093/plcell/koab269

33. Zhou P, Enders TA, Myers ZA, Magnusson E, Crisp PA, Noshay J, et al. Prediction
of conserved and variable heat and cold stress response in maize using cis-
regulatory information. Plant Cell [Internet]. 2021 Nov 4 [cited 2021 Nov 11];
(koab267). Available from: https://doi.org/10.1093/plcell/koab267

34. Karollus A, Mauermeier T, Gagneur J. Current sequence-based models capture
gene expression determinants in promoters but mostly ignore distal enhancers.
Genome Biol. 2023 Mar 27;24(1):56.

35. Sasse A, Ng B, Spiro AE, Tasaki S, Bennett DA, Gaiteri C, et al. Benchmarking of
deep neural networks for predicting personal gene expression from DNA
sequence highlights shortcomings. bioRxiv. 2023 Sep 28;2023.03.16.532969.

36. Takou M, Wieters B, Kopriva S, Coupland G, Linstädter A, De Meaux J. Linking
genes with ecological strategies in Arabidopsis thaliana. J Exp Bot. 2019 Feb
20;70(4):1141–51.

37. Yocca AE, Lu Z, Schmitz RJ, Freeling M, Edger PP. Evolution of Conserved
Noncoding Sequences in Arabidopsis thaliana. Mol Biol Evol. 2021 Jul
1;38(7):2692–703.

38. He F, Arce AL, Schmitz G, Koornneef M, Novikova P, Beyer A, et al. The Footprint
of Polygenic Adaptation on Stress-Responsive *Cis* -Regulatory Divergence in the
Arabidopsis Genus. Mol Biol Evol. 2016 Aug;33(8):2088–101.

39. Steige KA, Laenen B, Reimegård J, Scofield DG, Slotte T. Genomic analysis
reveals major determinants of *cis*- regulatory variation in *Capsella grandiflora*.
Proc Natl Acad Sci. 2017 Jan 31;114(5):1087–92.

40. Lasky JR, Des Marais DL, Lowry DB, Povolotskaya I, McKay JK, Richards JH, et al.
Natural Variation in Abiotic Stress Responsive Gene Expression and Local
Adaptation to Climate in Arabidopsis thaliana. Mol Biol Evol. 2014 Sep
1;31(9):2283–96.

41. Akagi T, Masuda K, Kuwada E, Takeshita K, Kawakatsu T, Ariizumi T, et al.
Genome-wide cis-decoding for expression design in tomato using cistrome data
and explainable deep learning. Plant Cell. 2022 Mar 8;koac079.

42. Hannah MA, Wiese D, Freund S, Fiehn O, Heyer AG, Hincha DK. Natural Genetic
Variation of Freezing Tolerance in Arabidopsis. Plant Physiol. 2006 Sep
1;142(1):98–112.

43. Zhen Y, Dhakal P, Ungerer MC. Fitness Benefits and Costs of Cold Acclimation in
Arabidopsis thaliana. Am Nat. 2011 Jul 1;178(1):44–52.

44. Oakley CG, Savage L, Lotz S, Larson GR, Thomashow MF, Kramer DM, et al.
Genetic basis of photosynthetic responses to cold in two locally adapted
populations of Arabidopsis thaliana. J Exp Bot. 2018 Jan 23;69(3):699–709.

45. Oakley CG, Ågren J, Atchison RA, Schemske DW. QTL mapping of freezing
tolerance: links to fitness and adaptive trade-offs. Mol Ecol. 2014;23(17):4304–
15.

46. 1001 Genomes Consortium. 1,135 Genomes Reveal the Global Pattern of
Polymorphism in Arabidopsis thaliana. Cell; 2016.

47. Rai A, Umashankar S, Rai M, Kiat LB, Bing JAS, Swarup S. Coordinate Regulation
of Metabolite Glycosylation and Stress Hormone Biosynthesis by TT8 in
Arabidopsis. Plant Physiol. 2016 Aug 1;171(4):2499–515.

48. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–
9.

49. Bailey TL. STREME: accurate and versatile sequence motif discovery.
Bioinformatics. 2021 Sep 15;37(18):2834–40.

50. Sandelin A, Alkema W, Engström P, Wasserman WW, Lenhard B. JASPAR: an
open-access database for eukaryotic transcription factor binding profiles. Nucleic
Acids Res. 2004 Jan 1;32(suppl_1):D91–4.

51. Gupta S, Stamatoyannopoulos JA, Bailey TL, Noble WS. Quantifying similarity
between motifs. Genome Biol. 2007 Feb 26;8(2):R24.

52. Takou M, Balick DJ, Steige KA, Dittberner H, Göbel U, Schielzeth H, et al.
Strength of stabilizing selection on the amino-acid sequence is associated with
the amount of non-additive variance in gene expression [Internet]. bioRxiv; 2022
[cited 2022 Nov 3]. p. 2022.02.11.480164. Available from:
https://www.biorxiv.org/content/10.1101/2022.02.11.480164v2

53. JASPAR - A database of transcription factor binding profiles [Internet]. [cited
2021 Nov 22]. Available from: https://jaspar.genereg.net/

54. Wright MN, Ziegler A. ranger: A Fast Implementation of Random Forests for High
Dimensional Data in C++ and R. J Stat Softw. 2017 Mar 31;77(1):1–17.

55. Washburn JD, Mejia-Guerra MK, Ramstein G, Kremling KA, Valluru R, Buckler ES,
et al. Evolutionarily informed deep learning methods for predicting relative
transcript abundance from DNA sequence. Proc Natl Acad Sci. 2019 Mar
19;116(12):5542–9.

56. Abadi M, Agarwal A, Barham P, Brevdo E, Chen Z, Citro C, et al. TensorFlow:
Large-Scale Machine Learning on Heterogeneous Distributed Systems.

57. Lee G, Sanderson BJ, Ellis TJ, Dilkes BP, McKay JK, Ågren J, et al. A large-effect
fitness trade-off across environments is explained by a single mutation affecting
cold acclimation. Proc Natl Acad Sci. 2024 Feb 6;121(6):e2317461121.

58. Fowler SG, Cook D, Thomashow MF. Low Temperature Induction of Arabidopsis
CBF1, 2, and 3 Is Gated by the Circadian Clock. Plant Physiol. 2005 Mar
1;137(3):961–8.

59. R Core Team. R: A language and environment for statistical   computing.
[Internet]. R Foundation for Statistical Computing, Vienna, Austria.; 2018.
Available from: http://www.R-project.org/

60. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models
Using lme4. J Stat Softw. 2015 Oct 7;67(1):1–48.

61. Fox J, Weisberg S. An R Companion to Applied Regression. [Internet]. Third.
Sage; 2019 [cited 2019 Jul 14]. Available from:
https://socialsciences.mcmaster.ca/jfox/Books/Companion/

62. Yanagisawa S. Chapter 12 - Structure, Function, and Evolution of the Dof
Transcription Factor Family. In: Gonzalez DH, editor. Plant Transcription Factors
[Internet]. Boston: Academic Press; 2016 [cited 2023 Jun 30]. p. 183–97.
Available from:
https://www.sciencedirect.com/science/article/pii/B9780128008546000129

63. Metzger BPH, Wittkopp PJ, Coolon JD. Evolutionary Dynamics of Regulatory
Changes Underlying Gene Expression Divergence among Saccharomyces
Species. Genome Biol Evol. 2017 Apr 1;9(4):843–54.

64. Lenhard B, Sandelin A, Carninci P. Metazoan promoters: emerging
characteristics and insights into transcriptional regulation. Nat Rev Genet. 2012
Mar 6;13(4):233–45.

65. Wei Q, Jr RLD. The Role of Balanced Training and Testing Data Sets for Binary
Classifiers in Bioinformatics. PLOS ONE. 2013 Jul 9;8(7):e67863.

66. Buda M, Maki A, Mazurowski MA. A systematic study of the class imbalance
problem in convolutional neural networks. Neural Netw. 2018 Oct 1;106:249–59.

67. Fraser HB, Babak T, Tsang J, Zhou Y, Zhang B, Mehrabian M, et al. Systematic
Detection of Polygenic cis-Regulatory Evolution. PLOS Genet.
2011;7(3):e1002023.

68. Chawade A, Bräutigam M, Lindlöf A, Olsson O, Olsson B. Putative cold
acclimation pathways in Arabidopsis thaliana identified by a combined analysis
of mRNA co-expression patterns, promoter motifs and transcription factors. BMC
Genomics. 2007 Sep 2;8(1):304.

69. Li W, Yin Y, Quan X, Zhang H. Gene Expression Value Prediction Based on
XGBoost Algorithm. Front Genet [Internet]. 2019 [cited 2022 Dec 19];10.
Available from: https://www.frontiersin.org/articles/10.3389/fgene.2019.01077

70. Smet D, Opdebeeck H, Vandepoele K. Predicting transcriptional responses to
heat and drought stress from genomic features using a machine learning
approach in rice. Front Plant Sci [Internet]. 2023 [cited 2023 Jul 31];14. Available
from: https://www.frontiersin.org/articles/10.3389/fpls.2023.1212073

71. Fowler S, Thomashow MF. Arabidopsis Transcriptome Profiling Indicates That
Multiple Regulatory Pathways Are Activated during Cold Acclimation in Addition
to the CBF Cold Response Pathway[W]. Plant Cell. 2002 Aug 1;14(8):1675–90.

72. Fagny M, Austerlitz F. Understanding the adaptation of polygenic traits: the importance of gene regulatory networks. 2021;22.

73. Kidokoro S, Konoura I, Soma F, Suzuki T, Miyakawa T, Tanokura M, et al. Clock-regulated coactivators selectively control gene expression in response to different temperature stress conditions in Arabidopsis. Proc Natl Acad Sci. 2023 Apr 18;120(16):e2216183120.

74. Corrales AR, Carrillo L, Lasierra P, Nebauer SG, Dominguez-Figueroa J, Renau-Morata B, et al. Multifaceted role of cycling DOF factor 3 (CDF3) in the regulation of flowering time and abiotic stress responses in Arabidopsis. Plant Cell Environ. 2017;40(5):748–64.

75. Fornara F, de Montaigu A, Sánchez-Villarreal A, Takahashi Y, Ver Loren van Themaat E, Huettel B, et al. The GI–CDF module of Arabidopsis affects freezing tolerance and growth as well as flowering. Plant J. 2015;81(5):695–706.

76. Kim HJ, Hyun Y, Park JY, Park MJ, Park MK, Kim MD, et al. A genetic link between cold responses and flowering time through FVE in Arabidopsis thaliana. Nat Genet. 2004 Feb;36(2):167–71.

77. Lang L, Pettkó-Szandtner A, Elbaşı HT, Takatsuka H, Nomoto Y, Zaki A, et al. The DREAM complex represses growth in response to DNA damage in Arabidopsis. Life Sci Alliance [Internet]. 2021 Dec 1 [cited 2024 Jan 12];4(12). Available from: https://www.life-science-alliance.org/content/4/12/e202101141

78. Kim JY, Ryu JY, Baek K, Park CM. High temperature attenuates the gravitropism of inflorescence stems by inducing SHOOT GRAVITROPISM 5 alternative splicing in Arabidopsis. New Phytol. 2016;209(1):265–79.

79. Meng X, Liang Z, Dai X, Zhang Y, Mahboub S, Ngu DW, et al. Predicting transcriptional responses to cold stress across plant species. Proc Natl Acad Sci. 2021 Mar 9;118(10):e2026330118.

80. Zhou P, Enders TA, Myers ZA, Magnusson E, Crisp PA, Noshay JM, et al. Prediction of conserved and variable heat and cold stress response in maize using cis-regulatory information. Plant Cell. 2021;34(1):514–34.

81. Gorjifard S, Jores T, Tonnies J, Mueth NA, Bubb K, Wrightsman T, et al. Arabidopsis and maize terminator strength is determined by GC content, polyadenylation motifs and cleavage probability. Nat Commun. 2024 Jul 12;15(1):5868.

82. Mei W, Stetter MG, Gates DJ, Stitzer MC, Ross-Ibarra J. Adaptation in plant genomes: Bigger is different. Am J Bot. 2018;105(1):16–9.

83. Srivastava D, Aydin B, Mazzoni EO, Mahony S. An interpretable bimodal neural network characterizes the sequence and preexisting chromatin predictors of induced transcription factor binding. Genome Biol. 2021;22(1).

84. Rivière Q, Corso M, Ciortan M, Noël G, Verbruggen N, Defrance M. Exploiting
Genomic Features to Improve the Prediction of Transcription Factor Binding Sites
in Plants. Plant Cell Physiol. 2022 Jul 7;pcac095.

85. Savolainen O, Lascoux M, Merilä J. Ecological genomics of local adaptation. Nat
Rev Genet. 2013 Nov;14(11):807–20.

86. Ferebee TH, Buckler E. Exploring the utility of regulatory network-based machine
learning for gene expression prediction in maize [Internet]. bioRxiv; 2023 [cited
2023 May 15]. p. 2023.05.11.540406. Available from:
https://www.biorxiv.org/content/10.1101/2023.05.11.540406v1

87. Monroe JG, McGovern C, Lasky JR, Grogan K, Beck J, McKay JK. Adaptation to
warmer climates by parallel functional evolution of CBF genes in Arabidopsis
thaliana. Mol Ecol. 2016;25(15):3632–44.

88. Wittkopp PJ, Kalay G. Cis-regulatory elements: molecular mechanisms and
evolutionary processes underlying divergence. Nat Rev Genet. 2012
Jan;13(1):59–69.

89. Buel GR, Walters KJ. Can AlphaFold2 predict the impact of missense mutations
on structure? Nat Struct Mol Biol. 2022 Jan;29(1):1–2.

90. Cheng J, Novati G, Pan J, Bycroft C, Žemgulytė A, Applebaum T, et al. Accurate
proteome-wide missense variant effect prediction with AlphaMissense. Science.
2023 Sep 19;381(6664):eadg7492.

91. Koo PK, Ploenzke M. Improving representations of genomic sequence motifs in
convolutional networks with exponential activations. Nat Mach Intell. 2021
Mar;3(3):258–66.

1029

Data Accessibility

1031 All scripts are available at the github repositories:

1032 https://github.com/mtakou/cnns_DEGpromoters and https://github.com/em-bellis/CNN_GxE.

Conflict of Interest Statement

1034 ESB is currently a full time employee of Avalo, a crop improvement company.

1035

48