Title: Long-sperm precedence and other cryptic female choices in *Drosophila melanogaster*

Authors: Brooke Peckenpaugh[1], Joanne Y. Yew[2], and Leonie C. Moyle[1]

[1] Department of Biology, Indiana University, Bloomington, Indiana 47405
[2] Pacific Biosciences Research Center, University of Hawai'i at Mānoa, Honolulu, Hawai'i 96822

Corresponding author

Leonie C. Moyle, Department of Biology, Indiana University, 1001 E 3rd Street, Bloomington, IN 47405-7000, USA.
Email: lmoyle@iu.edu

Data availability statement

The data and code underlying this article are available in the Dryad Digital Repository, at https://dx.doi.org/[doi].

Author contributions

BP and LCM conceived and designed the experiment. BP performed the experiment and statistical analyses. JYY performed the chemical analysis. BP and LCM wrote the paper. All authors read and provided feedback on the manuscript.

Funding

Indiana University, Bloomington (LCM)
NIH Common Themes in Reproductive Diversity (CTRD) training grant (BP)
National Institute of General Medical Sciences of the National Institutes of Health Grant No. P20GM125508 (JYY), P20GM139753 (JYY)

Conflict of interest statement

The authors have no conflict of interest to declare.

Acknowledgments

We thank Stuart J. Macdonald for providing the DSPR founder lines used in this study, and Brandon S. Cooper for providing the *Drosophila yakuba* line. We thank Sung Nichin for assistance scoring courtship behaviors. We thank Alison Pischedda for advice on the perfuming protocol. All GCMS measurements were performed in the Univ. of Hawai'i at Mānoa Microbial Genomics and Analytical Laboratory core. We thank Matthew W. Hahn, Kimberly A. Rosvall, and Michael J. Wade for their helpful comments on the manuscript. The research was supported by Indiana University Department of Biology funding to LCM and BP, as well as the NIH Common Themes in Reproductive Diversity (CTRD) training grant funding to BP.

LONG-SPERM PRECEDENCE AND OTHER CRYPTIC FEMALE CHOICES IN

DROSOPHILA MELANOGASTER

Abstract

Females that mate multiply make postmating choices about which sperm fertilize their

eggs (cryptic female choice); however, the male characteristics they use to make such choices

remain unclear. In this study, we sought to understand female sperm use patterns by evaluating

whether *Drosophila melanogaster* females adjust sperm use (second male paternity) in response

to four main factors: male genotype, male courtship effort, male pheromone alteration, and male

postmating reproductive morphology. Our experiment was replicated across four different *D.*

melanogaster lines, in a full factorial design, including a pheromone manipulation in which

second males were perfumed to resemble heterospecific (*D. yakuba*) males. We found that

females prefer longer sperm—regardless of mating order—in almost all contexts; this observed

pattern of 'long-sperm precedence' is consistent with female postcopulatory choice of high-

fitness male traits. Nonetheless, we also found that this general preference can be plastically

altered by females in response to effects including perfuming treatment; this differential female

sperm use is between otherwise identical males, and therefore solely female-mediated.

Furthermore, our finding that females exercise choice using diverse criteria suggests a possible

mechanism for the maintenance of variation in sexually selected male traits.

Keywords: cryptic female choice, sperm, pheromones, *Drosophila*

Teaser text

How do females make choices about which sperm will fertilize their eggs? In this study, we assess four geographically diverse *Drosophila melanogaster* lines to show that females strongly prefer long sperm—a pattern consistent with female preference for high-fitness males. In addition, we perfumed males to show that females can adjust this general preference based on other factors, including pheromones. Because this adjustment occurs between genotypically identical males, it is entirely female-mediated.

1 Introduction

Reproduction in nature is costly; organisms are therefore expected to use their reproductive resources judiciously and to adjust them according to reproductive context. Examples of such behavioral reproductive plasticity span diverse taxa, from palm trees (Barot et al. 2005) to guppies (Gasparini and Evans 2018) to *Drosophila* (Nandy et al. 2016). One specific strategy for plastically allocating reproductive resources is to adjust choices about reproductive partners. Females, in particular, are known to do this in numerous ways prior to mating. For instance, in red jungle fowl, females choose to mate with males with brighter feathers and longer tails; these traits are more prominent in unparasitized males and are therefore honest indicators of male health (Zuk et al. 1990). While many such cases involve choices exercised before mating, some organisms also have the ability to selectively allocate postmating reproductive resources; for instance, female guppies fertilize more eggs when mating with more colorful focal males in comparison to stimulus males (Pilastro et al. 2004). However, because most of these choices are made during and after mating events, it remains hard to differentiate the relative contributions of

each mating partner, the criteria on which their choices are based, and therefore the degree to

which these choices could be adaptive.

Interestingly, in *Drosophila*, postmating reproductive plasticity has primarily been

studied in males. Even under optimal conditions, *Drosophila melanogaster* males are known to

be ejaculate-limited and cannot produce progeny after two to five matings in a day (Douglas et

al. 2020). Plastic allocation of reproductive resources could therefore allow them to prioritize

sperm and seminal fluid for preferred mates. Consistent with this, *D. melanogaster* males use

premating information, such as female condition and the relative abundance of potential sexual

competitors, to modulate ejaculate composition (Sirot 2019, Wigby et al. 2009, Lüpold et al.

2011, Sirot et al. 2011, Garbaczewska et al. 2013). *D. melanogaster* males also allocate more

resources to matings with large females—an investment that reduces their success at defensive

sperm competition in subsequent matings (Anastasio et al. 2023). Together, these studies broadly

indicate that costs to *Drosophila* males drive plasticity in their reproductive allocation during

mating, as well as identify specific criteria used in making these behavioral decisions.

Like males, females are also expected to selectively allocate reproductive resources (i.e.,

eggs) at postmating stages, through the action of "cryptic female choice"—the female-mediated

physical and chemical factors that determine which sperm are used to fertilize eggs after mating

(Firman et al. 2017). In *Drosophila,* several proximate mechanisms of cryptic female choice

involve modulating the storage of sperm prior to fertilization. *Drosophila* females have two types

of sperm storage organs and can bias sperm use through differential sperm storage and/or by

preferentially drawing sperm from these alternative organs (Manier et al. 2013b). Females can

also manipulate the identity of sperm available in storage by adjusting the time until sperm

ejection after specific matings (Snook and Hosken 2004, Manier et al. 2013a, Laturney et al.

2018). Finally, it has been proposed that the size of sperm storage organs themselves might

provide a mechanism by which females make these choices. For example, females

experimentally evolved to have longer seminal receptacles have been shown to select for males

with longer sperm (Miller and Pitnick 2002). Even when the specific mechanism is unknown, it

is evident that *Drosophila* females can change sperm use patterns depending upon context,

including according to male genotype identity (Chow et al. 2010). What is comparatively poorly

understood, however, is the evolutionary explanation for these apparent female choices,

including the criteria used by females to exercise them.

One reason for this is the challenge of differentiating female choice from primarily male-

mediated factors that can also influence the outcome of postmating processes (Firman et al.

2017). For instance, *Drosophila simulans* males that are more attractive to females also sire more

offspring as second males; however, it is unclear whether this is because females preferentially

use sperm from attractive males, or because attractive males are intrinsically better sperm

competitors (Hosken et al. 2008). While the former explanation would be consistent with an

adaptive function for cryptic female choice, the latter explanation implies passive female

responses to the outcome of male-male sexual selection. One strategy for determining the

specific contribution of female choices to patterns of biased sperm use is to evaluate how these

patterns change when female perception of male identity is manipulated, without influencing

male identity itself. Although female assessment of male identity is likely to be complex, in

Drosophila, female premating preference primarily depends on auditory and chemosensory cues

(Yamamoto and Nakano 1999, Greenspan and Ferveur 2000); manipulating these cues could

therefore modify this assessment. Contact pheromones are one such cue. In insects broadly, and

flies specifically, these pheromones include cuticular hydrocarbons (CHCs)—waxy compounds

on the outer surface (cuticle) of many insects that can be used as sexual pheromones in mate

choice by males (Billeter et al. 2009) and females (Scott 1994), both within and between species

(Blows and Allan 1998). CHC composition can be artificially altered to assess its effect on mate

choice by "perfuming" focal males with the CHCs of other male genotypes (Dyer et. al. 2014,

Wang et al. 2011). Perfuming males with CHCs from unfamiliar or unpreferred males,

immediately prior to mating, can therefore manipulate female perception of male identity while

keeping male postmating quality the same.

In this study, we sought to understand the nature and relative importance of factors that

females use when exercising cryptic female choice. To do so, we evaluated whether *Drosophila*

melanogaster females adjust their sperm use patterns in response to four main factors: male

genotype, male courtship effort, male pheromone alteration, and male postmating reproductive

morphology. We manipulated male pheromones via "perfuming" with heterospecific (*D. yakuba*)

males—thereby changing the perceived, but not actual, identity of these treatment males. We also

evaluated postmating reproductive trait variation in males and females (testis length, sperm

length, male reproductive tract [MRT] mass, sperm storage organ size, female reproductive tract

[FRT] mass) to assess their contribution to sperm preference patterns. We assessed how second-

male paternity (proportion of offspring sired) changed according to each factor, by mating males

of each type to females that had previously mated with a consistent first-male tester genotype.

Our experiment was replicated across four different *D. melanogaster* lines in a full factorial

design, to evaluate whether the effects of these factors on sperm use depend upon female

genotype, male genotype, or their interaction. We find clear evidence that females adjust their

use of sperm from the second male. In particular, females show a general preference for longer

sperm, whether that sperm is from the first or the second male—i.e. 'long-sperm precedence'

rather than 'last-male sperm precedence' (Laturney et al. 2018). Nonetheless, we also find that

this general preference can be overridden in specific contexts by female-mediated changes in

response to male genotype, courtship effort, and perfuming treatment. The presence and direction

of these cryptic choices vary in complex ways, including evidence that females of some

genotypes adjust sperm use based on their perception of male identity, regardless of actual

identity. These data allow us to draw broader inferences about the possible criteria for, as well as

the evolutionary causes and consequences of, such choices.

2   Methods

2.1 Fly stocks

All stocks were reared on standard media prepared by the Bloomington Drosophila Stock

Center and kept at room temperature (~22°C). For this experiment we used four lines from

among the 15 *D. melanogaster* founder lines in the Drosophila Synthetic Population Resource

(DSPR) (King et al. 2012). These fully-sequenced founder lines represent a diverse collection of

genotypes from different locations around the world, and are known to vary in reproductive traits

(i.e., male courtship song, see Pischedda et al. 2014a). Our four DSPR founder lines (A5

[Greece], A7 [Taiwan], B4 [California], and B7 [Malaysia]) were used as focal genotypes for

females and second males in each manipulation. The GFP-labeled *D. melanogaster* line (32170)

was used as the first tester male. The *D. yakuba* line yak_S09_L38 was the heterospecific donor

male for the heterospecific-perfumed treatment (collected by B. Cooper in Sao Tome in 2009).

For each conspecific-perfumed treatment, target males were perfumed (see methods below) with

males from their own genotype.

2.2 Experimental design

To assess patterns of preferential sperm use, including after remating with conspecific-

vs. heterospecific-perfumed males, we mated *D. melanogaster* females sequentially to two

males: first, to *D. melanogaster* GFP-labeled tester males, and second, to *D. melanogaster* target

males. Using a GFP-labeled first male allowed us to assess paternity share (first [tester] versus

second [target] males) of offspring from each female) by examining the phenotypes of offspring

generated after the second mating. Any offspring not expressing GFP are sired by the second

focal male. For our CHC manipulation, the second, target males were either perfumed by

genotypically identical males (conspecific-perfumed control) or perfumed by heterospecific *D.*

yakuba males (Figure 1).

We replicated this experimental procedure across all four *D. melanogaster* genotypes, in

a full factorial design. For each line, we set up matings between females and second males from

each of the four DSPR lines, including males from the females' own genotype. This created 16

cross types in total. For each cross-type, we assessed paternity share from crosses with

conspecific-perfumed males and with heterospecific-perfumed males. Each of these cross types

was repeated until successfully replicated at least five times, resulting in over 160 total crosses.

2.3 Perfuming manipulation

Males used in the second mating were perfumed in the morning before setting up crosses.

To do so, we vortexed four target males with 30 donor males, either of the same *D. melanogaster*

genotype (for conspecific-perfumed control crosses) or with 30 heterospecific (*D. yakuba*) donor

males (for heterospecific-perfumed treatment crosses). The flies were vortexed on medium

strength for one minute total, in three 20-second increments with 20-second breaks in between

(Pischedda et al. 2014b). The target males were separated immediately after vortexing and

isolated individually in vials. They were allowed to recover for one hour before setting up

crosses.

2.4 Paternity assay

D. melanogaster males were isolated and aged for seven days, while females were

isolated and aged for two days before setting up the first cross with a GFP-labeled tester male.

We allowed mating for 24 hours before removing the males. We allowed the isolated female to

lay eggs for an additional 24 hours and verified successful matings by observing larval activity.

Unmated females (those that failed to produce larvae after the first mating) were removed from

the experiment. Mated females were then moved to a new vial to be paired with the second male.

To determine whether and the degree to which females exhibit premating discrimination against

heterospecific-perfumed males, we observed a subset of crosses for two hours to observe the

number of second matings within this period and recorded the time until remating (latency) and

copulation duration for each. Regardless of whether matings were observed in the first two

hours, seven-day-old second males were housed with females for 24 hours. Males were then

removed and the females were transferred to a new vial. Females continued to be transferred to

new vials every 24 hours for three days. Progeny in these vials were allowed to mature into

adulthood, then counted and assessed for GFP to determine paternity (see details below). All

cross types were repeated until we had five successful replicates (crosses in which there is

evidence for both a first and second mating). Conspecific- and heterospecific-perfumed treatment

crosses for each genotype were completed in parallel and in a randomized order.

Upon eclosion, all progeny from the three post-remating vials were anesthetized with

CO_2 and viewed under a Leica M205FA stereo scope equipped with a UV light for visualizing

GFP. The presence or absence of GFP in the ocelli of the eye were used as a marker of paternity;

offspring with GFP ocelli were sired by the first male and were thus counted as P1. Progeny with

wild-type ocelli were sired by the second male and were counted as P2. If no wild-type progeny

were present in the three post-remating vials, we assumed that no remating event occurred, and

these were scored as "unremated".

2.5 Cuticular hydrocarbon extraction and quantification

We confirmed the efficacy of perfuming using gas chromatography–mass spectrometry

(GC/MS) analysis. We extracted cuticular hydrocarbons from males of all four genotypes, for

both conspecific-perfumed and heterospecific-perfumed treatments. Following treatment of

seven-day old virgin males, cuticular hydrocarbons were extracted from pooled samples by

placing five males in a 1.8 mL glass vial (Wheaton 224740 E-C Clear Glass Sample Vials) with

120 μL of hexane (Sigma Aldrich, St Louis, MO, USA) spiked with 10 μg/mL of hexacosane

(Sigma Aldrich). After 20 min, 100 μL of the solution was removed to a sterilized 1.8 mL glass

vial (Wheaton 224740 E-C Clear Glass Sample Vials) and allowed to evaporate under a fume

hood. Extracts were stored at −20°C until analysis. Five replicate samples, consisting of five flies

per sample (25 flies total), were prepared for each sample type.

We performed gas chromatography–mass spectrometry (GC/MS) analysis on a 7820A

GC system equipped with a 5975 Mass Selective Detector (Agilent Technologies, Inc., Santa

Clara, CA, USA) and a HP-5ms column ((5%-Phenyl)-methylpolysiloxane, 30 m length, 250 μm

ID, 0.25 μm film thickness; Agilent Technologies, Inc.). Electron ionization (EI) energy was set

at 70 eV. One microliter of the sample was injected in splitless mode and analyzed with helium

flow at 1 mL/min. We used two different temperature gradients depending on the sample type.

The column was set at 40°C and held for 3 min, increased to 200°C at a rate of 35°C/min, then

increased to 280°C at a rate of 3°C/min and held for 15 min. Chromatograms and spectra were

analyzed using MSD ChemStation (Agilent Technologies, Inc.). We identified CHCs based on

retention time and electron ionization fragmentation pattern. The abundance of each compound

was then quantified by normalizing the area under each CHC peak to the area of the hexacosane

signal using homebuilt peak selection software (personal correspondence, Dr. Scott Pletcher,

Univ. of Michigan).

2.6 Postmating reproductive trait morphology

We measured several elements of reproductive trait morphology in each of our lines, and

in the tester line, in order to assess their potential influence on patterns of female preferential

sperm use.

2.6.1 Thorax length, testis length, SR length, and ST area

We collected both female and male virgins and aged them to reproductive maturity (N=5

replicates per sex per genotype). We measured thorax length for all flies; this is a standard

measure of body size in *Drosophila* studies because it strongly correlates with the size of other

characters such as wing length (Robertson and Reeve 1952). We then dissected the reproductive

tract into Grace's Insect Medium. For females, we separated the seminal receptacle (SR) and a

randomly chosen spermatheca (ST) from the rest of the reproductive tract and transferred them to

a slide. For males, a randomly chosen testis was separated from the rest of the reproductive tract

and transferred to a slide. In both cases, we then applied a cover slip and took images under an

EVOS FL digital inverted microscope. Using ImageJ, we measured the length of the SR, the

height and width of the ST, and the length of the testis, from these images. Measurements were

replicated across five individuals per genotype.

2.6.2 Sperm length

We dissected sperm out of the seminal vesicles of reproductively mature virgin males into

Grace's insect medium on a slide (N=5 replicates per genotype). Sperm were gently teased apart

with a probe when necessary. The slides were stained with SpermBlue (Microptic), and imaged

under an EVOS FL digital inverted microscope. In ImageJ, we measure sperm length for 5-10

randomly chosen sperm per individual.

2.6.3 Body mass and reproductive tract mass

We collected both female and male virgins and aged them to reproductive maturity (N=5

replicates per sex, per genotype). We measured body and reproductive tract mass according to

Holman et al. (2008). Flies were dissected in distilled water onto a piece of preweighed foil. For

females, the reproductive tract (including the bursa, SR, ST, parovaria, and ovaries) was removed

and transferred to another piece of pre-weighed foil. For males, the reproductive tract (including

testes, seminal vesicles, accessory glands, and the ejaculatory bulb) was removed and transferred

to another piece of pre-weighed foil. For both males and females, the foils holding the body and

reproductive systems of the flies were then placed in a drying oven at 60°C for 18 ± 1 h and

weighed on a Mettler Toledo UMX2 balance to the nearest 0.1 μg.

2.7 Second-male courtship behavior assay

To identify variation in male courtship across genotypes, and to determine whether our

experimental manipulation affects male courtship behavior, we recorded male courtship for each

cross type on video. We recorded videos using a modified FlyPi setup, which combines a

Raspberry Pi (Raspberry Pi Foundation, Cambridge, UK), pi camera, and 3D printed parts (Maia

Chagas et al. 2017). Crosses were set up in a modified cell culture plate consisting of six culture

wells, each with a 3 cm diameter and a small amount of cornmeal media in the bottom, allowing

us to record six crosses simultaneously. Male and female flies were aspirated without

anesthetization into each cell culture well, and the plate was videotaped for two hours. The full

videos were scored for successful copulations, latency to copulation, and copulation duration.

The first 30 minutes of each video was also scored in more detail for the frequency of courtship

song, male tapping, number of male licking attempts, and copulation attempts. Recorded matings

were set up until we had five replicate recordings per cross type.

2.8 Statistical analyses

All statistical analysis and figure construction in this study was performed with R version

4.3.1. All linear models were run using deviation-coded contrasts. This does not change the fit of

the model, but affects the interpretation of regression coefficients such that the intercept

corresponds to the grand mean and main effect estimates represent deviations from the mean.

Model estimates are therefore robust to whichever line is used as the baseline in each

comparison, which is preferable in our experiments where none of our genotypes were *a priori*

designated as the baseline level.

2.8.1 CHC composition

We assessed variation in CHC composition using a principal components analysis (PCA),

which we computed using the prcomp function in R. To test for significant differences among

our principal components, we performed a multivariate analysis of variance (MANOVA) using

the manova function.

2.8.2 Postmating reproductive trait morphology

We assessed variation in female and male reproductive morphology by fitting linear

models using the lm function in R. To test whether male or female genotype predicts the

variation we observed in reproductive morphology, we ran linear models for each morphological

trait separately as the dependent variable. Male or female genotype was always included as the

predictor. The GFP-labelled *D. melanogaster* line was incorporated into each model as a

baseline.

2.8.3 Second-male courtship effort, female second-mating frequency, and second-male paternity

We assessed variation in second-male courtship by fitting a linear model using the lm

function in R. Second male courtship effort was the dependent variable, with male genotype,

female genotype, their interaction, and male perfuming treatment, as predictors.

We assessed variation in female second-mating frequency by fitting a binomial

generalized linear model using the glm function in R. We coded our response as a binomial

variable, with one column representing the number of successful second-matings, and the other

column representing the number of failed second-matings for each cross type (N=32 total cross

types). The generalized linear model included male genotype, female genotype, and male

perfuming treatment as predictors (this dataset did not allow us sufficient power to assess

interaction terms).

Variation in second male siring success was also assessed with a binomial generalized

linear model. We coded our response as a binomial variable, with one column representing the

number of offspring sired by the second male ("successes"), and the other column representing

the number of offspring sired by the first male ("failures") for each cross (N=196 total crosses).

Our generalized linear model included male genotype, female genotype, the interaction between

male and female genotype, male perfuming treatment, and the interaction between female

genotype and male perfuming treatment as predictors. California males, California females, and

conspecific-perfumed males were always incorporated into the models as baselines.

To assess the overall significance of our model components, we computed type III

analyses of variance (ANOVAs) using the same models, with the aov function in R. Effect sizes

(and 95% confidence intervals) of our model components were estimated using the resi function

in R, from the RESI package (Vandekar et al. 2020).

2.8.4 Correlations among datasets

Finally, we looked for correlations among our datasets. Across all pairwise genotype

combinations we assessed the strength of association between: second-male courtship effort and

second-male mating success; second male courtship effort and paternity share (P2); female

readiness to remate and second-male paternity share (P2); and, male and female reproductive

trait variation and second-male paternity share (P2). To do this, in each case, we computed

Spearman's rank correlation coefficients using the cor.test function in R. Each correlation was

assessed separately for experiments involving conspecific-perfumed males and heterospecific-

perfumed males.

3   Results

3.1 Males differ in CHC composition based on genotype and perfuming status

We quantified CHC composition across male genotypes and perfuming treatments to

characterize the variation present across lines, as well as to verify that our perfuming treatment

was effective. We found that CHC composition varied substantially across male genotypes

((Figure 2a,b, Table S1, Table S2). For each of the first three PCs—which together account for

92.3% of the variation observed—we detected a significant genotype effect (MANOVA $p <$

0.001; Table S1). CHC composition was also shifted by our perfuming treatment, as shown by

significant treatment effects for PC3 ($p = 0.004$) and PC4 ($p < 0.001$) (MANOVA; Figure 2b,

Table S1). These two PCs account for 8.6% of the variation in CHC composition.

3.2 Postmating reproductive traits differ more among males than among females

We quantified variation in male and female traits, focusing on postmating reproductive

traits, to assess whether these might influence patterns of sperm use. We found that all measured

male traits differed among genotypes, including two overall size traits and three reproductive

size traits (Figure 3, Table S3). Differences among female lines were more modest and only

observed for overall size and one reproductive size trait, RT mass (Figure 3, Table S3).

In terms of non-reproductive traits, both male and female Taiwan individuals consistently

had shorter thoraxes but also the largest dry non-RT body mass. Other differences among lines

were sex-specific (for example, in Greece males and Malaysia females) and often more modest

(Table S3).

For reproductive traits, California females exhibited the highest RT dry mass and Taiwan

females exhibited the lowest; however, female lines did not differ for either SR length or ST area

(Figure S1). In contrast, male lines differed in RT mass (Taiwan males had lower RT mass, while

Greece males had higher RT mass, than average), as well as testis length and sperm length

(Figure 3, Figure S1). California and Greece males had longer testes than average, while

Malaysia and Taiwan males had shorter testes. Interestingly, California and Taiwan males had

longer sperm than all other lines, and Malaysia and Greece males had significantly shorter

sperm. GFP sperm length was intermediate. This indicates that sperm length and testis length are

not closely associated among our *D. melanogaster* lines: Taiwan males had short testes but also

had among the longest sperm, while the inverse pattern was observed for Greece males.

3.3 Courtship effort varies consistently among male genotypes, but not among female genotypes

being courted or with perfuming

We evaluated courtship across each male-female genotype combination to better

understand if and/or how premating processes influence cryptic female choices, including via our

perfuming treatments. We measured courtship effort as the number of courtship behaviors

exhibited per minute until copulation occurred or until our observation period ended (at two

hours). Note that only 9 of 162 scored pairings resulted in copulation within 2 hours. Courtship

intensity varied significantly among male genotypes (Table 1), with Greece and Taiwan males

courting significantly less than the others (Figure 4, Table S4). Male courtship intensity was also

marginally reduced in heterospecific-perfumed males, but this effect was not statistically

significant (Figure 4, Table S4). In addition, although female line identity had only a marginal

influence on courtship (Table 1), we did find that Taiwan females were consistently courted less

than average across male genotypes (Figure 4, Table S4). We did not detect any significant

female-by-male genotype interactions. Consistent with these observations, the estimated effect

size of male genotype on male courtship effort was more than 2.5 times that of any other effect in

our model (Table 1).

3.4 Readiness to remate differs among female genotypes and depends on male genotype, but not

on male perfuming status

For each cross type, we assessed whether females chose to mate with second males

versus refused the second-male mating. Remating was evidenced by whether a female produced

offspring from the second male. We found that female genotypes varied in their readiness to

remate following a first mating with the tester male (Table 2): California females remated more

readily while Greece and Taiwan females mated less readily than average (Figure 5, Table S5).

Male genotype also affected remating (Table 2): California males were more consistently

successful in these second matings, while Greece males were less successful on average (Figure

5, Table S5). In contrast, we detected no effect of male perfuming status on second-male

premating success (Figure 5, Table 2, Table S5).

3.5 Female genotypes vary in their overall use of sperm from the second male, and in whether

they adjust sperm use according to second-male perfuming status

We assessed how paternity of second-males (P2) was affected by genotype and by our

perfuming manipulation. We found strong and significant differences in second-male paternity

depending on both female and second-male genotype, perfuming treatment, female by male

genotype interaction effects, and female by treatment interaction effects (Table 3). The largest of

these effects was male genotype, while the smallest was the female by treatment effect; the

estimated effect size of second male genotype was ~2 times that of female genotype, and ~3.5

times that of the male by female interaction (Table 3). When evaluated with a complete linear

model across all genotype pairs, we detected several general genotype effects. In particular,

Taiwan males gained higher P2 than average, while Greece and Malaysia males had reduced

paternity share (Figure 6, Table 4). In comparison, Greece females exhibited a higher P2 than

average, while Taiwan females exhibited a lower P2 than average, across all genotype pairs

(Figure 6, Table 4). Despite these general effects, second-male paternity share also varied

substantially depending on the particular male-female combination. For instance, Greece females

exhibit lower average P2 when paired with Greece second males, while Taiwan females increase

P2 when paired with Greece second males (Figure 6, Table 4).

We also found that females adjust sperm use according to male perfuming treatment,

including in female genotype-specific ways. Compared to conspecific-perfumed males,

heterospecific-perfumed males showed a modest but significantly higher second male paternity

overall (Figure 6, Table 4), but female genotypes vary in whether and how sperm use patterns

were affected by perfuming (Figure 6, Table 4). Specifically, Taiwan females show an overall

reduction in paternity associated with male heterospecific perfuming status, while Greece and

Malaysia females exhibit slight overall increases in P2 associated with male heterospecific

perfuming (Figure 6, Table 4). In comparison, California females do not appear to respond to

male CHC perfuming when making sperm use choices.

3.6 Both pre- and postmating choices can be affected by premating behavior, but sperm length

shows the most consistent association with patterns of preferential sperm use

We next assessed correlations between datasets to better understand how these variables

might influence one another. First, we evaluated the relationship between second-male courtship

effort and second-male mating success—i.e., is premating courtship intensity associated with

premating success? We did not see a consistent association for either conspecific-perfumed males

(Figure S3a; Spearman's rank correlation $\rho = 0.308$, $p = 0.246$) or heterospecific-perfumed males

(Figure S3b; Spearman's rank correlation $\rho = 0.174$, $p = 0.519$). Instead, readiness to remate

seems to be more clearly associated with individual female line differences. For instance,

California females tend to remate most readily, and Greece and Taiwan least readily, regardless

of the courtship intensity of the male line they are paired with (Figure S3a,b).

Second, we assessed the relationship between second male courtship effort and paternity

share (P2)—i.e., do females use premating signals to make postmating decisions? We found that

when males are perfumed by conspecifics, male courtship effort is significantly correlated with

siring success (Figure 7a; Spearman's rank correlation $\rho = 0.535$, $p = 0.035$). When males are

perfumed by heterospecifics, the correlation is positive but not significant (Figure 7b;

Spearman's rank correlation $\rho = 0.353$, $p = 0.180$). This weak positive trend between courtship

effort and P2 appears to be largely driven by California males, which are the most active courters

and also received the greatest share of paternity (Figure 7a,b), an association which is unclear in

the other male lines. Thus, we infer that the relationship between these variables is likely

influenced by additional factors not measured here; for instance, Greece males, which have low

siring success, have been shown to exhibit especially long interpulse intervals during courtship

song, which are less preferred by *D. melanogaster* females (Pischedda et al. 2014a, Kyriacou and

Hall 1982).

We next assessed the relationship between female readiness to remate and second-male

paternity share—i.e., is premating choosiness associated with postmating choosiness? We found

no correlation among these factors for conspecific-perfumed males (Figure 7c; Spearman's rank

correlation $\rho = -0.095$, $p = 0.726$), and a slight association for experiments with heterospecific-

perfumed males (Figure 7d; Spearman's rank correlation $\rho = 0.502$, $p = 0.048$). These weak or

absent associations are consistent with heterogenous effects we observe among female lines. For

instance, while Greece females showed relatively high premating choosiness (remating in only

59% of trials) and the highest average P2 of any genotype (consistent with strong second male

preference), Taiwan females exhibit a similar premating choosiness (60% remating frequency)

but significantly lower average P2 (indicating no consistent postmating preference for first or

second male sperm) (Figure 7c,d). This relationship therefore appears to vary substantially

depending upon which of the four genotypes is considered.

Finally, we examined the relationship between second-male paternity share and

reproductive trait variation among lines. Of RT mass (both male and female), SR length, testis

length, sperm length, and SR length minus sperm length, only sperm length and MRT mass were

significantly associated with P2. Of these two, MRT mass was significantly negatively correlated

with P2 among heterospecific-perfumed males (Figure S3d; Spearman's rank correlation $\rho = -$

0.679, $p = 0.004$), but not among conspecific-perfumed males (Figure S3c; Spearman's rank

correlation $\rho = -0.449$, $p = 0.081$). These relationships appear to be substantially influenced by

Taiwan males, which have high P2 but comparatively low RT mass. In comparison, the

relationship between sperm length and P2 was consistently and strongly positive for both

conspecific-perfumed (Figure 7e; Spearman's rank correlation $\rho = 0.631$, $p = 0.009$) and

heterospecific-perfumed groups (Figure 7f; Spearman's rank correlation $\rho = 0.728$, $p = 0.001$).

Second males with longer sperm experience greater siring success. Specifically, when the second

male's sperm is longer than the tester male (as for California or Taiwan second males), P2 is

generally higher; whereas when the second male sperm is shorter than the tester male (as for

Malaysia and Greece), P2 is generally lower. Together, these data indicate that females

preferentially use longer sperm—a generic preference that appears as last male precedence when

the second male's sperm is longer than the first (tester) male's sperm, but can override last male

precedence when the second male's sperm is relatively shorter.

4   Discussion

In this study, we sought to understand whether and how females make plastic choices

about sperm use. We found that sperm use patterns vary based on male genotype, female

genotype, their interaction, as well as based on male pheromone profile (even when male identity

is the same). Among these effects, male sperm length, specifically the relative difference in

sperm length between the first and second male, appears to play a particularly important role in

determining female sperm use patterns. While sperm length has been recognized as an important

component of male-male competition (e.g., Snook 2005, Immler et al. 2011, Lüpold and Pitnick

2018), our analysis enables us to infer that the preference for longer sperm is also female-

mediated, context-dependent, and controlled by factors other than the size of the female sperm

storage organs, as we expand upon below.

4.1 Female sperm use choices vary among lines in their direction and magnitude, and in the criteria used to evaluate males

Consistent with previous work in *D. melanogaster*, we observe clear evidence for sperm choice mediated by females (Clark and Begun 1998, Clark et al. 1999, Miller and Pitnick 2002, Chow et al. 2010). Female lines show consistent patterns of sperm use in control crosses (with their own unmanipulated second males), and significant shifts in these patterns in response to changing either second-male genotype or perfuming treatment or both (depending on the female line). This result underscores the potential for this female-mediated process to influence genotypic representation among the resulting offspring, and to generate sexual selection on males (Pitnick et al. 2001).

We found substantial and consistent variation among our four female lines in the direction and magnitude of preferential sperm use patterns. *D. melanogaster* is conventionally described as having second (or last) male sperm precedence (Lefevre and Jonsson 1962, Pischedda and Rice 2012). However, depending on the specific male-female combination, our estimated preference for the sperm of second-males ranged from 0.2 to 0.95. Values below 0.5 (e.g., Taiwan females with Malaysia males) indicate consistent first-male precedence in these specific male-female combinations. Moreover, female lines differ in how strongly they differentiate among sperm donors. For example, with the exception of California males (which they strongly prefer; average P2 = 0.79), California females do not consistently differentiate between the first and second males' sperm when paired with the other three second male lines in our experiment (i.e., mean P2 is approximately 0.5, with a broad range of values among individual CA females; Figure 6, Table S6).

We also clearly observed among-line variation in the characteristics that females use to

evaluate males as reproductive donors. In particular, Greece females (and to some extent Taiwan

and Malaysia females) modulate their use of second male sperm in response to perfuming status,

albeit in complex ways. For example, Greece females decrease their use of Greece sperm when

these males are heterospecific-perfumed, but Taiwan females increase their use of Greece sperm

after the same treatment (Figure 6, Table S7). In comparison, male perfuming appears to have no

influence on cryptic choice by California females. This diversity in sperm choice patterns

depending on intrinsic (female genotype) and extrinsic (perfumed status) factors points to a

much greater heterogeneity of sperm use decision-making among *D. melanogaster* females than

simple last male precedence (see also Laturney et al. 2018). Given the broad geographic origin of

these lines, differences in the magnitude and direction of female choices could reflect

reproductive or other differences among lines that operated prior to their collection and lab

establishment.

4.2 Cryptic female choice is strongly shaped by male traits, especially sperm length

Within the diversity of individual sperm use criteria, however, we did observe several

broad patterns. In particular, every female line differentiated among male genotypes in at least

one postmating experimental context, indicating that one or more aspects of male genotype are

important in making paternity allocation decisions. Consistent with this, male genotype had the

largest estimated effect size on second male paternity share (P2; Table 3), nearly twofold greater

than the next largest effect (female genotype). Our analysis also identifies specific male traits

that likely mediate these male genotype effects. As noted above, some female lines adjust sperm

use in response to perfuming treatment, indicating that male cuticular hydrocarbon profile affects

their postmating choices. Among conspecific-perfumed second males, paternity share is also

associated with courtship effort suggesting that females may prefer the sperm of male genotypes

who court more, although this effect is shaped largely by California males. Most strikingly,

however—and more consistently than either chemosensory or courtship traits—we find that

sperm length is positively associated with preferential sperm use across many contexts and

conditions. The majority of female lines preferentially used the second-male's sperm when that

sperm was longer than the first-male's sperm (i.e., when California or Taiwan was the second

male). Moreover, in several cases (for example, when Taiwan females mate with Malaysia

males—see Figure 6) female lines subverted the expected precedence for last males' sperm

specifically when the first (GFP tester) male had longer sperm than the second male (i.e. Greece

and Malaysia males). That is, our results indicate that 'long-sperm precedence', in addition to

'last-male sperm precedence' (Lefevre and Jonsson 1962), jointly contribute to female sperm use

patterns in *D. melanogaster*.

Finally, although our data indicates diverse criteria for female sperm choice overlaid on a

generic preference for longer sperm, it also indicates that these patterns are not explained by at

least one studied mechanism of female sperm choice—SR length (Miller and Pitnick 2002). SR

length has been proposed as a primary mechanism via which females discriminate between

sperm of different lengths (Pitnick et al. 1999); in particular, females with longer SRs are thought

to be most selective because they can better discriminate between the sperm length of the first

and second male. However, while we observe clear variation in female discrimination of sperm

based on length—e.g., Greece females exhibit generally high P2, regardless of length, whereas

California females only exhibit high P2 for their own (long sperm) males (Figure 6, Table 4)—

we do not observe significant variation in SR length among our female lines (Figure 3, Table S3,

Table S2). Thus, SR length is not the mechanism by which these diverse *D. melanogaster* female

lines discriminate against short sperm in our experiments.

4.3 Evolutionary implications of the observed variation in female sperm choice

Consistent patterns in choice criteria among female genotypes provide an opportunity to

evaluate the nature of past drivers of female choice in our system, including hypotheses where

cryptic female choice is not beneficial to females versus where it has specific fitness benefits.

For example, one proposal is that female sperm use patterns result primarily from manipulation

by males; that is, cryptic female choice is the product of sexual conflict rather than providing an

adaptive benefit to females. In this case, female sperm use patterns are expected to be random

with respect to actual measures of male quality. Our data do not convincingly support this

hypothesis, given the general postmating female preferences we observe for males who have the

longest sperm and, under some conditions, for males that court frequently.

Alternatively, Birkhead (1998) proposed that female sperm choice might function to bias

sperm use against the sperm of closely related males. However, our findings are not consistent

with inbreeding avoidance as a primary function of cryptic female choice. If that were the case,

second-male success (against the unrelated tester first male) should consistently be lower in

crosses with males from the same line as females, compared to males from different lines.

However, female lines differed in their choice of own male sperm, ranging from consistent high

preference to low preference.

Apart from inbreeding avoidance, other hypothesized functions of sperm choice include

female benefits from sire discrimination (e.g., choice of high-quality males ['good genes'

hypothesis; Eberhard 1996], or choice of attractive males ['sexy sons' hypothesis; Weatherhead

and Robertson 1979]). We find that females have a generic preference for longer sperm (Figure 7e,f); in this case, choosing longer-spermed males increases the frequency of longer-spermed (i.e. 'sexier') sons, which themselves could benefit from the same generic preference. As studies have shown that long sperm have a competitive advantage in displacing shorter sperm (Lüpold et al. 2012), this preference likely also aligns specifically with higher reproductive fitness in males. Thus, our data support the hypothesis that females could gain indirect fitness benefits from exercising cryptic choice based on specific male traits like sperm length.

The differences we observe among female lines in the criteria they used for sperm choice also have implications for the expected future drivers of postmating reproductive evolution. Some of our lines altered sperm use patterns based on contact pheromones (especially Greece, but also Taiwan and Malaysia), indicating that pheromone bouquet can be a criterion for evaluating preferred sires. Other lines (e.g., California) were insensitive to pheromone manipulation in their sperm use choices. Variation in choice criteria makes it more challenging for males to tailor specific traits to satisfy, or manipulate (Ryan 1990), a broad range of female biases. Moreover, when the specific preferences of females differ according to male trait values and male trait identity, there is no single most preferred male genotype, and variation should be maintained at male traits. This expectation is consistent with our observation, and with previous analyses (e.g., Clark et al. 1999, Miller and Pitnick 2002, Chow et al. 2010), showing that the outcomes of cryptic female choice are demonstrably influenced by male by female interactions.

That these criteria for female sperm choice can also vary behaviorally is particularly evident in the case of Greece females. When sequentially mated with males from other lines, Greece females uniformly favor whichever genotype (tester versus second male) has the longer sperm. However, when mated with their own conspecific-perfumed second males, Greece

females preferentially use this sperm (Figure 6), even though it is significantly shorter than the

first (tester) male (Figure 3). In this case, the female preference for local males appears to

override their generic preference for longer sperm. We know that this decision is both female-

mediated and based on chemosensory assessment, because it is reversed when Greece females

are paired with heterospecific-perfumed Greece males. Greece females instead favor the longer

(tester male) sperm in these pairings (Figure 6), even though these Greece males are identical in

every sense to conspecific-perfumed males—except for their CHC profile. Flexible changes in

preference, such as this case where females specifically prefer own males despite generic

preferences for longer sperm, is another potential mechanism by which sexually selected

variation could be maintained in males.

5   Conclusion

The direction and magnitude of female-mediated sperm choice, as well as the male traits

influencing these choices, varies significantly among four female lines of *D. melanogaster*. This

diversity in female sperm choices is overlaid on a generic preference for longer sperm,

suggesting that paternity decisions may ultimately benefit female fitness indirectly through the

production of higher-quality offspring or more attractive offspring. Interestingly, at least one

proposed proximate mechanism of female sperm choice—SR length—does not account for

variation in observed sperm use. Diversity in cryptic female choices—both in the specific

modalities used to assess and choose among males, and in the plasticity of these choices—may

explain how variation in male postmating sexually selected traits is maintained over time.

References

Anastasio, O. E., Sinclair, C. S., & Pischedda, A. (2023). Cryptic male mate choice for high-quality females reduces male postcopulatory success in future matings. *Evolution*, 77(6), 1396-1407. doi:10.1093/evolut/qpad064

Barot, S., Mitja, D., Miranda, I., Meija, G. D., & Grimaldi, M. (2005). Reproductive plasticity in an Amazonian palm. *Evolutionary Ecology Research*, *7*(7), 1051-1065.

Billeter, J. C., Atallah, J., Krupp, J. J., Millar, J. G., & Levine, J. D. (2009). Specialized cells tag sexual and species identity in Drosophila melanogaster. *Nature*, 461(7266), 987-991. doi:10.1038/nature08495

Birkhead, T. R. (1998). Cryptic Female Choice: Criteria for Establishing Female Sperm Choice. *Evolution*, 52(4), 1212-1218. doi:10.1111/j.1558-5646.1998.tb01848.x

Blows, M. W., & Allan, R. A. (1998). Levels of mate recognition within and between two Drosophila species and their hybrids. *Am Nat*, 152(6), 826-837. doi:10.1086/286211

Chow, C. Y., Wolfner, M. F., & Clark, A. G. (2010). The genetic basis for male x female interactions underlying variation in reproductive phenotypes of Drosophila. *Genetics,* 186(4), 1355-1365. doi:10.1534/genetics.110.123174

Clark, A. G., & Begun, D. J. (1998). Female genotypes affect sperm displacement in Drosophila. *Genetics*, 149(3), 1487-1493. doi:10.1093/genetics/149.3.1487

Clark, A. G., Begun, D. J., & Prout, T. (1999). Female x male interactions in Drosophila sperm competition. *Science,* 283(5399), 217-220. doi:10.1126/science.283.5399.217

Douglas, T., Anderson, R., & Saltz, J. B. (2020). Limits to male reproductive potential across mating bouts in Drosophila melanogaster. *Animal Behaviour*, 160, 25-33.

Dyer, K. A., White, B. E., Sztepanacz, J. L., Bewick, E. R., & Rundle, H. D. (2014). Reproductive character displacement of epicuticular compounds and their contribution to mate choice in Drosophila subquinaria and Drosophila recens. *Evolution*, 68(4), 1163-1175.

Eberhard, W. (1996). Female control: sexual selection by cryptic female choice (Vol. 69): Princeton University Press.

Firman, R. C., Gasparini, C., Manier, M. K., & Pizzari, T. (2017). Postmating female control: 20 years of cryptic female choice. *Trends in Ecology & Evolution*, *32*(5), 368-382.

Garbaczewska, M., Billeter, J. C., & Levine, J. D. (2013). Drosophila melanogaster males increase the number of sperm in their ejaculate when perceiving rival males. *J Insect Physiol,* 59(3), 306-310. doi:10.1016/j.jinsphys.2012.08.016

Gasparini, C., & Evans, J. P. (2018). Female control over multiple matings increases the
opportunity for postcopulatory sexual selection. *Proceedings of the Royal Society*
B, *285*(1888), 20181505.

Greenspan, R. J., & Ferveur, J. F. (2000). Courtship in Drosophila. *Annu Rev Genet*, 34, 205-232.
doi:10.1146/annurev.genet.34.1.205

Holman, L., Freckleton, R. P., & Snook, R. R. (2008). What use is an infertile sperm? A
comparative study of sperm-heteromorphic Drosophila. *Evolution*, *62*(2), 374-385.

Hosken, D. J., Taylor, M. L., Hoyle, K., Higgins, S., & Wedell, N. (2008). Attractive males have
greater success in sperm competition. *Curr Biol*, 18(13), R553-554.
doi:10.1016/j.cub.2008.04.028

Immler, S., Pitnick, S., Parker, G. A., Durrant, K. L., Lupold, S., Calhim, S., & Birkhead, T. R.
(2011). Resolving variation in the reproductive tradeoff between sperm size and number.
Proc Natl Acad Sci U S A, 108(13), 5325-5330. doi:10.1073/pnas.1009059108

King, E. G., Merkes, C. M., McNeil, C. L., Hoofer, S. R., Sen, S., Broman, K. W., . . .
Macdonald, S. J. (2012). Genetic dissection of a model complex trait using the
Drosophila Synthetic Population Resource. *Genome Res*, 22(8), 1558-1566.
doi:10.1101/gr.134031.111

Kyriacou, C. P., & Hall, J. C. (1982). The function of courtship song rhythms in
Drosophila. *Animal Behaviour*, *30*(3), 794-801.

Laturney, M., van Eijk, R., & Billeter, J. C. (2018). Last male sperm precedence is modulated by
female remating rate in Drosophila melanogaster. *Evolution letters*, *2*(3), 180-189.

Lefevre, G., Jr., & Jonsson, U. B. (1962). Sperm transfer, storage, displacement, and utilization
in Drosophila melanogaster. *Genetics*, 47(12), 1719-1736.
doi:10.1093/genetics/47.12.1719

Lüpold, S., Manier, M. K., Ala-Honkola, O., Belote, J. M., & Pitnick, S. (2011). Male Drosophila
melanogaster adjust ejaculate size based on female mating status, fecundity, and
age. *Behavioral Ecology*, *22*(1), 184-191.

Lupold, S., Manier, M. K., Berben, K. S., Smith, K. J., Daley, B. D., Buckley, S. H., . . . Pitnick,
S. (2012). How multivariate ejaculate traits determine competitive fertilization success in
Drosophila melanogaster. *Curr Biol*, 22(18), 1667-1672. doi:10.1016/j.cub.2012.06.059

Lupold, S., & Pitnick, S. (2018). Sperm form and function: what do we know about the role of
sexual selection? *Reproduction*, 155(5), R229-R243. doi:10.1530/REP-17-0536

Maia Chagas, A., Prieto-Godino, L. L., Arrenberg, A. B., & Baden, T. (2017). The€ 100 lab: A
3D-printable open-source platform for fluorescence microscopy, optogenetics, and
accurate temperature control during behaviour of zebrafish, Drosophila, and
Caenorhabditis elegans. *PLoS biology*, *15*(7), e2002702.

Manier, M. K., Lupold, S., Pitnick, S., & Starmer, W. T. (2013). An analytical framework for
estimating fertilization bias and the fertilization set from multiple sperm-storage organs.
Am Nat, 182(4), 552-561. doi:10.1086/671782

Manier, M. K., Lupold, S., Belote, J. M., Starmer, W. T., Berben, K. S., Ala-Honkola, O., . . .
Pitnick, S. (2013). Postcopulatory sexual selection generates speciation phenotypes in
Drosophila. *Curr Biol*, 23(19), 1853-1862. doi:10.1016/j.cub.2013.07.086

Miller, G. T., & Pitnick, S. (2002). Sperm-female coevolution in Drosophila. *Science*, 298(5596),
1230-1233. doi:10.1126/science.1076968

Nandy, B., Dasgupta, P., Halder, S., & Verma, T. (2016). Plasticity in aggression and the
correlated changes in the cost of reproduction in male Drosophila melanogaster. *Animal
Behaviour*, *114*, 3-9.

Pilastro, A., Simonato, M., Bisazza, A., & Evans, J. P. (2004). Cryptic female preference for
colorful males in guppies. *Evolution*, *58*(3), 665-669.

Pischedda, A., & Rice, W. R. (2012). Partitioning sexual selection into its mating success and
fertilization success components. *Proc Natl Acad Sci U S A*, 109(6), 2049-2053.
doi:10.1073/pnas.1110841109

Pischedda, A., Cochrane, V. A., Cochrane, W. G., & Turner, T. (2014). Genetic dissection of
courtship song variation using the Drosophila Synthetic Population Resource. *bioRxiv*,
006643.

Pischedda, A., Shahandeh, M. P., Cochrane, W. G., Cochrane, V. A., & Turner, T. L. (2014).
Natural variation in the strength and direction of male mating preferences for female
pheromones in Drosophila melanogaster. *PLoS One*, 9(1), e87509.
doi:10.1371/journal.pone.0087509

Pitnick, S., Marrow, T., & Spicer, G. S. (1999). Evolution of Multiple Kinds of Female Sperm-
Storage Organs in Drosophila. *Evolution*, 53(6), 1804-1822. doi:10.1111/j.1558-
5646.1999.tb04564.x

Pitnick, S., Brown, W. D., & Miller, G. T. (2001). Evolution of female remating behaviour
following experimental removal of sexual selection. *Proc Biol Sci*, 268(1467), 557-563.
doi:10.1098/rspb.2000.1400

Robertson, F. W., & Reeve, E. C. (1952). Heterozygosity, environmental variation and heterosis.
Nature, 170(4320), 286. doi:10.1038/170286a0

Ryan, M. J. (1990). Sexual selection, sensory systems and sensory exploitation. *Oxford surveys*
in evolutionary biology, *7*, 157-195.

Scott, D. (1994). Genetic Variation for Female Mate Discrimination in Drosophila Melanogaster.
Evolution, 48(1), 112-121. doi:10.1111/j.1558-5646.1994.tb01298.x

Sirot, L. K., Wolfner, M. F., & Wigby, S. (2011). Protein-specific manipulation of ejaculate
composition in response to female mating status in Drosophila melanogaster. *Proc Natl*
Acad Sci U S A, 108(24), 9922-9926. doi:10.1073/pnas.1100905108

Sirot, L. K. (2019). Modulation of seminal fluid molecules by males and females. *Curr Opin*
Insect Sci, 35, 109-116. doi:10.1016/j.cois.2019.07.009

Snook, R. R., & Hosken, D. J. (2004). Sperm death and dumping in Drosophila. *Nature*,
428(6986), 939-941. doi:10.1038/nature02455

Snook, R. R. (2005). Sperm in competition: not playing by the numbers. *Trends Ecol Evol*, 20(1),
46-53. doi:10.1016/j.tree.2004.10.011

Vandekar S., Tao R., Blume J. (2020). A Robust Effect Size Index. *Psychometrika*. 85(1), 232-
246. doi:10.1007/s11336-020-09698-2.

Wang, L., Han, X., Mehren, J., Hiroi, M., Billeter, J.-C., Miyamoto, T., . . . Anderson, D. J.
(2011). Hierarchical chemosensory regulation of male-male social interactions in
Drosophila. *Nature neuroscience*, 14(6), 757.

Weatherhead, P. J., & Robertson, R. J. (1979). Offspring quality and the polygyny threshold:" the
sexy son hypothesis". *The American Naturalist*, *113*(2), 201-208.

Wigby, S., Sirot, L. K., Linklater, J. R., Buehner, N., Calboli, F. C., Bretman, A., . . . Chapman,
T. (2009). Seminal fluid protein allocation and male reproductive success. *Curr Biol*,
19(9), 751-757. doi:10.1016/j.cub.2009.03.036

Yamamoto, D., & Nakano, Y. (1999). Sexual behavior mutants revisited: molecular and cellular
basis of Drosophila mating. *Cell Mol Life Sci*, 56(7-8), 634-646.
doi:10.1007/s000180050458

Zuk, M., Johnson, K., Thornhill, R., & Ligon, J. D. (1990). Mechanisms of Female Choice in
Red Jungle Fowl. *Evolution*, 44(3), 477-485. doi:10.1111/j.1558-5646.1990.tb05933.x

Table 1. Factors affecting male courtship effort. Overall significance of model components and their effect sizes in predicting courtship effort as assessed using ANOVAs.

Model component	Sum Sq	Df	F	p	Effect size	95% CI
Female line	1.397	3	2.140	0.098	0.087	0.000-0.443
Second male line	7.349	3	11.254	<0.001 ***	0.392	0.275-0.897
Treatment	0.824	1	3.786	0.054	0.120	0.000-0.301
Female line x second male line	3.663	9	1.870	0.061	0.226	0.153-0.569

Table 2. Factors affecting female remating frequency. Overall significance of model components and their effect sizes in predicting female remating as assessed using ANOVAs.

Model component	LR Chi sq	Df	p	Effect size	95% CI
Female line	19.776	3	<0.001 ***	0.707	0.199-1.329
Second male line	13.787	3	0.003 **	0.439	0.000-1.263
Treatment	0.061	1	0.805	0.000	0.000-0.323

Table 3. Factors affecting second male paternity share (P2). Overall significance of model components and their effect sizes in predicting female use of second male sperm (P2) as assessed using ANOVAs.

Model component	LR Chi sq	Df	p	Effect size	95% CI
Female line	448.71	3	<0.001 ***	0.331	0.221-0.616
Second male line	1327.93	3	<0.001 ***	0.632	0.492-0.980
Treatment	37.91	1	<0.001 ***	0.063	0.000-0.218
Female line x second male line	297.72	9	<0.001 ***	0.170	0.020-0.507
Female line x treatment	21.79	3	<0.001 ***	0.000	0.000-0.225

Table 4. Specific genotype, treatment, and interaction effects on second male paternity share (P2). Shown is the binomial generalized linear model of variation in second male paternity share (P2), depending on female line, second male line, treatment group, the interaction between female line and second male line, and the interaction between female line and treatment group. California females, California males, and conspecific-perfumed males are incorporated into the model as a baseline.

Model component	Estimate	Std. error	z	p
Intercept	0.723	0.023	30.860	<0.001 ***
Female line (Greece)	0.889	0.051	17.372	<0.001 ***
Female line (Malaysia)	0.024	0.032	0.733	0.463
Female line (Taiwan)	-0.385	0.036	-10.842	<0.001 ***
Second male line (Greece)	-0.616	0.039	-15.688	<0.001 ***

Second male line (Malaysia)	-0.921	0.035	-26.074	<0.001 ***
Second male line (Taiwan)	0.446	0.039	11.496	<0.001 ***
Treatment (het-perfumed)	0.132	0.022	6.144	<0.001 ***
Female line (Greece) x second male line (Greece)	-0.341	0.087	-3.904	<0.001 ***
Female line (Malaysia) x second male line (Greece)	-0.012	0.053	-0.230	0.818
Female line (Taiwan) x second male line (Greece)	0.265	0.057	4.665	<0.001 ***
Female line (Greece) x second male line (Malaysia)	-0.083	0.068	-1.211	0.226
Female line (Malaysia) x second male line (Malaysia)	-0.081	0.049	-1.661	0.097
Female line (Taiwan) x second male line (Malaysia)	-0.592	0.058	-10.247	<0.001 ***
Female line (Greece) x second male line (Taiwan)	0.792	0.087	9.126	<0.001 ***
Female line (Malaysia) x second male line (Taiwan)	0.033	0.053	0.628	0.530
Female line (Taiwan) x second male line (Taiwan)	-0.028	0.058	-0.481	0.630
Female line (Greece) x treatment (het-perfumed)	0.098	0.044	2.205	0.027 *
Female line (Malaysia) x treatment (het-perfumed)	0.073	0.030	2.435	0.015 *
Female line (Taiwan) x treatment (het-perfumed)	-0.123	0.034	-3.658	<0.001 ***

Figure 1. Experimental procedure for assessing paternity share of second (focal) males. Females were first mated with a GFP tester male. After 48 hours, these females were paired for second matings with focal males, which were either perfumed by conspecifics (left, control) or by heterospecifics (right, treatment). Unmated females were removed from the experiment. In remated females, we assessed paternity for first and second males based on the presence/absence of the GFP marker in resulting offspring.

Figure 2 CHC composition varied based on genotype (G, Greece; T, Taiwan; C, California; and M, Malaysia) and perfuming status (conspecific-perfumed males shown in grey, heterospecific-perfumed males shown in purple) (Table S1). Both panels show standardized principal component (PC) values generated from multivariate CHC composition data for replicate males of each genotype and treatment. Left panel: PCs 1 and 2 largely reflect differences between male genotypes. Right panel: PCs 3 and 4 reflect additional shifts in CHC composition associated with heterospecific perfuming treatment.

Figure 3. Variation in (a) thorax length in mm and (b) reproductive trait length in mm in females and males across *D. melanogaster* lines. We observed significant variation among lines for thorax length (both males and females), testis length, and sperm length, but SR length did not significantly differ between lines (Table S3). Box plots show the median (central line)

and first and third quartiles; whiskers extend to the smallest and largest values (not including outliers >1.5x the inter-quartile range).

Figure 4. Courtship effort (courtship behaviors per minute) consistently differed between second male genotypes but did not consistently vary by the female genotype being courted or by male perfuming status (Table 1, Table S4). Conspecific-perfumed males are shown in grey, and heterospecific-perfumed males are shown in purple. The dashed line indicates the average courtship effort for each male line. Box plots show the median (central line) and first and third quartiles; whiskers extend to the smallest and largest values (not including outliers >1.5x the inter-quartile range). The y-axis has been truncated to make variation in the data easier to visualize; the non-truncated data (as used in statistical analyses) is shown in Figure S2.

Figure 5. Counts of mating successes (dark shading) and mating failures (light shading) differs across male (column) and female (row) genotypes (Table 2, Table S5). Results are shown separately for conspecific-perfumed males (grey) and heterospecific-perfumed males (purple) within each male x female panel.

Figure 6. Female use of second male sperm (P2) varies across female genotypes, male genotypes, and female by male genotype combinations (Table 3, Table 4). P2 also varies depending on male perfuming status, as shown in grey (conspecific-perfumed males) and purple (heterospecific-perfumed males). The dashed line indicates the average P2 for each male line. Box plots show the median (central line) and first and third quartiles; whiskers extend to the smallest and largest values (not including outliers >1.5x the inter-quartile range).

Figure 7. Second male courtship effort is significantly correlated with siring success when males are perfumed by conspecifics (a),but is not significant when perfumed by heterospecifics (b). Female remating rate is not correlated with second male siring success when males are conspecific-perfumed (c) but is correlated when males are heterospecific-perfumed (d). Sperm length is significantly correlated with siring success for both conspecific-perfumed (e) and heterospecific-perfumed males (f). Crosses are labeled as female x male genotype (G, Greece; T, Taiwan; C, California; and M, Malaysia).

845
846