1 **Socio-ecological context modulates significance of territorial contest**
2 **competition in *Drosophila prolongata***
3

4 Alessio N. De Nardo[1]*, Broti Biswas[1,2], Jhoniel Perdigón Ferreira[1], Abhishek Meena[1],
5 Stefan Lüpold[1]
6
7 [1]Department of Evolutionary Biology and Environmental Studies, University of Zurich,
8 Winterthurerstrasse 190, 8057 Zurich, Switzerland
9 [2]Department of Systematic and Evolutionary Botany, University of Zurich,
10 Zollikerstrasse 107, 8008 Zurich, Switzerland
11
12 *Correspondence: alessio.denardo@uzh.ch
13

14 **Abstract**
15

The intensity and direction of sexual selection is intricately linked to the social and
ecological context. Theory predicts that male-biased operational sex ratios (OSR) and
high population density should skew resource and mate monopolization toward few
successful males, thereby intensifying sexual selection on males. Here, we studied
how the mating system of the promiscuous and strongly sexually dimorphic fruit fly
Drosophila prolongata responds to changes in the OSR and population density. We
recorded groups of flies over five days and quantified territory occupancy, mating
success, and competitive fertilization success. Although general indices of the
intensity of sexual selection supported theoretical predictions, realized selection on
morphological traits did not. Rather, we found selection on these traits to intensify
under high density, but also under even as opposed to male-biased OSRs. Larger and
more territorial males achieved higher mating success, but only under even OSRs.
Similarly, competitive fertilization success, mainly determined by sperm precedence,
was biased toward large males only under even OSRs. Finally, our combined results
support a shift in the mating system from territorial contest competition to scramble
competition under male-biased OSRs and potentially at low density, where there was
no clear contribution of the measured traits to reproductive success. Our study
emphasizes the limitations of traditional selection metrics and the role of the socio-
ecological context in predicting adaptation to a changing environment.

Significance

Mating systems are complex and dynamic, adapting to ongoing ecological change. Studies often assume that changes in the socio-ecological context alter the intensity of sexual selection on traits indicating individual fitness, but our work on *Drosophila prolongata* challenges this view. By manipulating operational sex ratio and population density, and jointly investigating territorial behavior and both pre- and post-mating reproductive success over several days, we reveal the plastic mating strategies in this fly. This dynamism underscores the limitations of static classifications and the importance of studying selection across diverse socio-ecological contexts. This broader perspective advances our understanding of the tight connections between environmental change, population demographics, and the evolutionary process.

Introduction

Sexual selection, driven by competition for mates and/or fertilization (1), shapes heritable traits and behaviors that are directly related to reproductive success (2). While often stronger on males (2, 3), sexual selection can also significantly impact females (4–6), with its extent and nature on either sex being intricately linked to the ecological context (7).

One important ecological parameter affecting the intensity of sexual selection and mating system dynamics is the distribution of resources, which shapes the local population composition and so the availability of potential mates [i.e., operational sex ratio, OSR (7)]. Skewed OSRs can lead to monopolization of mating opportunities by a few successful individuals of the competing sex (7). Male-biased OSRs in particular can thus drive selection on traits that are advantageous in competition, including larger body size, the growth of weaponry, or elaborate ornamentation favored by females (7–9).

A meta-analysis across a diversity of animal taxa supports theoretical predictions that OSRs can drive the general intensity of sexual selection (10). However, studies exploring selection on specific phenotypic traits often find no differences between OSRs (11–13), or their results contrast general theory (14–17). For example, in the common lizard (*Lacerta vivipara*), positive selection on male body size is stronger in female-biased than in male-biased OSRs, possibly resulting from higher female costs

under elevated male aggression during mating (14). Overall, the effect of the OSR can
vary considerably between species and mating systems (18, 19).

OSRs can also influence post-mating processes, such as competitive fertilization
(e.g., refs. 20–22; but see ref. 23), ultimately impacting aspects like ejaculate transfer
or mating duration (24). Males tend to invest more in ejaculate production when
exposed to a higher risk or intensity of sperm competition (20), often induced by
changes in the perceived OSR (25–27). Across generations, this is expected to lead
to stronger selection on traits associated with competitive fertilization, such as
ejaculate allocation (28) or testes size (29), yet studies on the latter are inconclusive
(30–36). In a literature review (30), only one of four studies found an effect of OSR on
testes size. This one study showed an increase in testes size for male *Drosophila*
melanogaster evolving under an extreme female bias [10:1], suggesting selection
against sperm limitation rather than competition (31).

In addition to the OSR, competition for mates can also intensify as populations
become denser due to growing numbers of competitors (37). However, studies on the
impact of density on metrics of sexual or trait-specific selection have yielded mixed
results (supporting: refs. 38–41, opposing: refs. 9, 42–46) or suggest a trade-off
between pre- and post-mating sexual selection (36, 47). This implies that the exclusive
focus on pre-mating selection in many studies misses this shift toward post-mating
competition at higher densities where aggressive strategies might become less
effective (36, 37, 48). For example, across anuran species, the trade-off between
forelimb muscles and testes size is biased more strongly toward the latter when
population density is high and fertilizations are more difficult to monopolize (36).

Beyond the population structure, male reproductive tactics may also be influenced
by the distribution of resources (7). While the opportunity for monopolization is minimal
when resources are evenly distributed, highly competitive males may gain control over
spatially concentrated resources (7). When resource monopolization strongly
correlates with mating success (49–51), a skew toward few successful males can be
predicted (7). These males may become more attractive to potential mates for their
ability to defend a resource or for the benefit the resource provides. For example, in
antlered flies, *Phytalmia spp.,* males fight over oviposition sites to which females then
gain access through copulation (52). Therefore, females indirectly choose mates with
large antlers, which in turn benefit males in these contests (53). However, within a
species, diverse reproductive tactics may coexist, depending on demographic,

environmental or individual conditions (37, 43, 49, 54–60). For example, whether a
resource is defendable can hinge on the number of individuals competing for it, and
switching to an alternative strategy can be advantageous (61–63). One such case is
the European bitterling, *Rhodeus sericeus*, in which males abandon their territories as
these become indefensible, and join spawning groups instead (63).

Although OSRs and population densities have both been proposed to shape sexual
selection and mating systems (7, 37), likely mediated by resource distribution (7), to
our knowledge these factors have not been studied jointly, let alone in the context of
both pre- and post-mating sexual selection. Here, we investigated these factors in a
comprehensive laboratory experiment using the fruit fly *Drosophila prolongata.* This
South-Asian species exhibits conspicuous male-biased sexual dimorphism both in
body size and in foreleg size and coloration (64), while also showing signals of
resource defense when high-quality resources are concentrated (65). Males use their
forelegs in both fights against competitors (66, 67) and elaborate courtship behaviors
(64). These behaviors also include "leg vibration", a species-specific behavior that
involves a male stimulating the female abdomen and is often displayed immediately
before copulation (67), likely to overcome remating suppression by mated females (68,
69). Leg vibration, however, can also trigger an alternative reproductive tactic by a
nearby male, namely interception of the female (70). With highly polygamous females
(68), this species is ideal for studies of context-dependent pre- and post-mating sexual
selection.

Here, we monitored over a five-day interaction period each of 30 genetically diverse
groups of *D. prolongata* that varied in their sex ratio (1:1 or 3:1 males:females) and
density (12, 16, or 20 flies). Quantifying the distribution of flies around a centrally
located resource, pre-mating behaviors, individual copulations, and molecular
parentage provided a uniquely comprehensive understanding of reproductive
dynamics in this species, highlighting the links between the evolution of sexually
selected traits, mating systems, population ecology, and behavioral plasticity.

Results and Discussion

Potential and realized sexual selection. To quantify the selection dynamics in
D. prolongata under different population compositions, we measured Bateman

gradients and the opportunity for sexual selection (I_s) as two commonly used estimates
for the intensity of sexual selection (e.g., 10). Whereas the former represents the slope
between individuals' number of mates and resulting offspring (ref. 71; i.e., the fitness
advantage gained by additional matings), the latter is based on the sex-specific
variances in reproductive success (ref. 72; i.e., the opportunity for selection to act on
variation). The Bateman gradients for both sexes were positive (Fig. 1; males: X^2_1 =
63.65, n = 193, P < 0.001, Table S1; females: X^2_1 = 24.1, n = 175, P < 0.001, Table
S2), but more pronounced in males (X^2_1 = 7.42, P = 0.006, Table S3a). The disparity
in Bateman gradients between the sexes was greater in the male-biased populations
(X^2_1 = 5.6, n = 368, P = 0.018) but unaffected by group density (X^2_2 = 0.23, P = 0.89,
Table S3b). Similarly, I_s was higher for males compared to females in all treatments,
but particularly pronounced in the male-biased populations (6.2× higher for males)
compared to the even OSR populations (2.7× higher for males). Population density
again had no significant effect (Table S4, Fig. S1). These findings combined suggest
that sexual selection is more pronounced on males than females, and its intensity for
males is amplified under a male-biased OSR, as previously shown for a diversity of
taxa (10).
When comparing selection on specific male traits, results were more complex. For
example, we found positive selection gradients on thorax, foreleg and wing length, and
composite body size (principal component 1, PC1; see *Methods*) under even OSR,
but only on wing length in male-biased populations (Fig. 2, Table S5). Also, contrary
to expectations based on I_s, positive selection on all measured traits appeared to
intensify with density (Fig. 2, Table S6). This increasing selection on secondary sexual
traits with density aligns with theory (37), yet has been reported for only few species
(e.g., ref. 40; but see refs. 43, 73). The changing pattern of selection on phenotypic
traits across OSRs, however, contradicts both theoretical predictions (7) and insights
gained from Bateman gradients and I_s in this study.
The lack of positive selection on putative sexually selected traits in male-biased
populations might simply indicate insufficient statistical power or that other,
unmeasured behavioral or morphological traits would be favored under conditions of
mate monopolization (2, 74, 75). However, our findings might also reveal the
limitations of using metrics like Bateman gradients, I_s, or the OSR to predict realized
selection on specific sexually selected traits, as previously emphasized (18). As our
results confirm, these common measures of the intensity of sexual selection do not

necessarily correlate with the observed selection on traits and merely set an upper
limit to the potential of sexual selection (18). These results underscore the need to
integrate different measures to gain a more complete understanding of the dynamics
of a mating system. Despite their limitations, Bateman gradients or I_s remain valuable
tools for numerically characterizing mating systems. They consider all types of sexual
selection (e.g. pre- and post-mating, intra- and intersexual), acting on any relevant
morphological or behavioral traits (76). Thus, they accurately capture variation in
fitness and allow for comparisons between sexes and populations (76). For example,
we showed that reproductive success is more skewed toward a few successful males
in male-biased populations than under even OSR. The trait-specific selection
gradients indicate, however, that traits selected in male-biased populations are not the
morphological traits expected to be involved in physical contest such as resource
defense (e.g., forelegs). These insights emphasize the intricate interplay between
OSR, density, and sexual selection in shaping mating system dynamics and trait
evolution in this species.

Territorial behavior & mating success. To investigate how the territorial behavior
previously reported in this species (65) manifests in different group compositions, a
resource-dense area (RDA) was established in the center of each experimental unit.
This RDA, consisting of highly nutritious yeast paste, covered approximately 16% of
the otherwise nutrient-poor arena. Regardless of population composition, males spent
more time in the RDA than females (35.6% vs. 12.3%; $X^2_1 = 671.6$, $P < 0.001$, $n = 493$
[303 males, 190 females], Table S7), with greater variance in males ($\sigma^2 = 0.016$) than
in females ($\sigma^2 = 0.005$). Hence, the RDA appeared to be highly attractive to males, but
access was unequal among males as reflected by their varying time spent there
(indicating "territorial success"). In addition, individuals of both sexes spent less time
(on average) in the RDA at higher densities ($X^2_1 = 22.0$, $P < 0.001$) and under male-
biased OSR ($X^2_1 = 21.0$, $P < 0.001$), suggesting intensifying competition for this
resource. Although females spent considerably less time in the RDA overall, 69.4% of
all matings started there. These matings were also more likely to involve aggressive
male-male behavior than matings outside this area ($z = 2.54$, $P = 0.011$, $n = 716$
matings across 212 males and 187 females). Both sexes spent more time in the RDA
on the first day of recording, when most matings occurred, compared to the
subsequent days ($X^2_4 = 313.724$, $P < 0.001$ $n = 2379$ across 5 days and 492

individuals; Fig. S2, Table S8). Interestingly, whilst body size (PC1) had no effect on
territorial occupancy on the first day, it did on all following days (interaction: X^2_1 = 5.05,
n = 1445, P = 0.02, Table S9).

Amino & Matsuo (65) report a hierarchy in pairs of males, with one male clearly
dominating territory occupancy, particularly when the area is small and easily
defendable. Although their findings do not directly demonstrate mating success by
territory occupancy, a strong correlation between territory occupancy and courtship
rate (65), as well as a mating advantage for winners of contests over territories (77),
have been reported for *D. prolongata*. In our study, the RDA was comparable to the
larger of the two areas used by Amino & Matsuo (65), but with more males and so
reduced ability to monopolize the contested area. Nevertheless, territorial control
influenced mating success, albeit with greater complexity at higher numbers of
competitors: Compared to 72.5% at an even OSR, only 64.6% of matings were
initiated in the RDA at a male-biased OSR. Interestingly, although most matings were
initiated in the RDA at both OSRs, overall territory occupancy determined mating
success only in even (t = 2.4, P = 0.019) but not male-biased populations (t = 0.24, P
= 0.8; OSR×territoriality: t = 1.7, P = 0.09, n = 290, controlling for PC1; Fig. 3, Table
S10a-b). Similarly, PC1 was the main contributor to mating success under even (t =
3.7, P < 0.001) but not male-biased OSRs (t = 1.1, P = 0.26, OSR×PC1: t = 2.15, P =
0.03*; n* = 290, Fig. 3, Table S10a-b). In addition to these results, the occurrence of
courtship behavior before mating depended on body size and the OSR (PC1×OSR: t
= 2.93, P = 0.003, n = 760, Fig. 3, Table S14a), in that large males showed more
courtship behavior in the even OSR (z = 3.20, P = 0.001) but not in the male-biased
OSR (z = -1.27, P = 0.20). These results align with the selection gradients above, but
additionally illustrate the changing dynamics of sexual selection in response to varying
levels of competition.

Our findings support the notion that when competition intensifies (e.g., strongly
male-biased OSR), monopolization of resources becomes increasingly difficult,
potentially shifting mating strategies (36, 37, 47, 48, 57). However, we neither
observed an increase in sneak matings at higher densities nor under male-biased
OSRs (both |z| < 1.2, P > 0.26, n = 716, Table S12). In fact, leg vibration, an auditory
cue used by sneaky males to intercept a mating attempt, even decreased at high
densities (z = -1.7, P = 0.09) and in male-biased OSR (z = 1.89, P = 0.059, n = 716,
Table S11). Courtship was observed less frequently when matings occurred in the

RDA (z = -3.3, P < 0.001, Table S14), possibly due to a high local density of
competitors and so higher risk of becoming the target of aggression (z = 2.54, P =
0.011, n = 716, Table S13) or interception (z = 2.31, P = 0.021, n = 716, Table S13).
Similarly, leg vibration was observed less often before matings in the RDA (z = -1.81,
P = 0.07, n = 716, Table S11). Large males were more likely to gain matings following
leg vibration (z = 2.89, P = 0.004, n = 716), whereas those acquiring matings by
interception tended to be smaller males (t = -2.03, P = 0.042, n = 716, Table S12). As
shown previously (78), males avoid leg vibration when the risk of interception is high
(e.g., at high density). Hence, notwithstanding the lesser importance of resource-
defense and body size for mating success in male-biased populations, males did not
adopt alternative mating tactics. This raises the question of whether selection might
instead favor increased investment in post-mating processes.
250
Fertilization success. Across all treatments, mating success (MS; 48.9%) explained
most of the variation in reproductive success, followed by fertilization success (FS;
14.3%) and female fecundity (Fec; 6.4%; Fig. S3, Table S16). This pattern was
comparable across social contexts in general, but while mating success explained
three times the variation of fertilization success under an even OSR (MS: 42.5%, FS:
14.4%, Fec: 9.2%), these variance components were more similar under a male-
biased OSR (MS: 37.3%, FS: 25.4%, Fec: 7.2%). Fertilization success may thus gain
in relative importance when female monopolization is hampered.
Males that were a female's last mate until the end of the experiment (t = 4.21, P <
0.001, n = 190, Table S16) and males that were able to delay remating for longer
("remating interval", see *SI Methods; t* = 4.79, P < 0.001) had higher fertilization
success (adjusted PCS, see *SI Methods*). Analogous to our findings for mating
success, we found that larger males only had higher fertilization success under an
even OSR (t = 2.63, P = 0.009), but not under a male bias (t = 0.14, P = 0.89;
OSR×PC1: t = 1.79, P = 0.075, n = 190, Table S16). This indicates that rather than
counteracting the lack of pre-mating selection on body size pre-mating under male-
biased OSRs, the pattern seems to be reinforced in the post-mating phase. Longer
mating durations did not significantly contribute to fertilization success (t = 1.62, P =
0.11), but males mating more often had decreased fertilization success per mating (t
= -2.29, P = 0.02), indicating strategic allocation (79) or sperm depletion (80).

In close relatives of *D. prolongata*, the allocation of male seminal fluid proteins

(SFPs), such as sex-peptide, can suppress female remating (81, 82). This allocation

can vary with a male's social context (82, 83) or his own condition and genotype (84).

Although the role of sex peptide in remating suppression is less clear in *D. prolongata*

(68), for any SFP delaying remating we would expect enhanced allocation when the

perceived sperm competition risk is higher [i.e., higher density or male-biased OSR;

(83, 85, 86)]. Yet, while female remating intervals were not affected by density (t =

0.69, P = 0.49), they were prolonged under even rather than male-biased OSRs (t =

2.14, P = 0.033, n = 190, Table S17). Matings were shorter for large males (t = -3.32,

P = 0.001, n = 710, Table S19) and tended to extend remating intervals (t = 1.91, P =

0.058), making an overall last-male position more likely (t = 1.97, P = 0.049, n = 211).

Male size did not affect remating intervals (t = 0.15, P = 0.88), yet large males were

more likely to be a female's last mate overall (t = 2.27, P = 0.024). To the extent that

mating duration predicts ejaculate transfer, shorter copulations by larger males can be

interpreted as strategic ejaculate allocation in anticipation of more future matings, as

shown previously in *D. melanogaster* (79) and supported here by shorter copulations

for males with more total matings on that day (t = -2.8, P = 0.005, n = 710).

The remating interval only reflects the average delay between two matings and here

was not affected by body size. However, the ability to delay remating is captured to a

larger extent in the proportion of overall last-male positions held until the end of the

experiment. Despite their short mating durations, large males ended up in the last-

male position to a higher proportion of their encounters, and remating intervals were

extended under even OSR where pre-mating selection is stronger. It thus seems as

though females might cryptically bias paternity toward preferred males by delaying

remating (87, 88). This scenario is further supported by three arguments: First, with

SFPs such as sex-peptide playing a major role in controlling the mating rate

D. melanogaster (89), a similar function could be predicted for *D. prolongata*.

However, considering a much higher mating rate than close relatives [i.e., 2.9±1.4

matings on the first day compared to a typical remating interval of 3-5 days in

D. melanogaster (90)]*, remating suppression through SFPs may be less effective in

D. prolongata (68), similar to other species with high mating rates (91). Alternatively,

even if sex-peptide does play a limited role, female *D. prolongata* might be able to

control the scope of its activity through strategic timing of ejaculate ejection (87).

Second, no effect of the OSR or density on mating duration were found (both |t| < 1.41,

P > 0.17, *n* = 710), which could indicate no strategic SFP allocation in response to the
level of sperm competition. Third, the remating interval was longer under even OSRs
although mating durations remained unaffected. Given the positive pre- and post-
mating selection on body size and territoriality under an even OSR, the observed
patterns are consistent with females potentially influencing post-mating paternity
biases to reinforce pre-mating choices.
In sum, our results indicate that intensifying competition does not augment post-
mating investment by males. Additionally, post-mating selection does not seem to
counteract the lack of pre-mating selection on ostensibly beneficial male traits.
Instead, post-mating processes, partially under female control, seem to reinforce the
patterns of pre-mating selection.
316
Mating system dynamics. To synthesize the differences in the mating-system
dynamics of *D. prolongata* across different contexts, we built structural equation
models. We provide separate visualizations for each density level in Fig. S4 (Tables
S22-S24), but Fig. 4 presents a simplified overview of the dynamics under an even
and male-biased OSR, respectively. One of the most striking observations was the
absence of many positive (green) or negative (red) effects in the male-biased scenario
(Fig. 4b, Table S21) compared to that in even populations (Fig. 4a, Table S20). This
underlines how both pre- and post-mating sexual selection vary across social contexts.
For example, body size, a primary determinant of both pre- and post-mating success
in the even population, seemed irrelevant under male-biased OSRs.
Even though we documented a larger skew in reproductive success and a higher
potential for sexual selection in male-biased populations, we were unable to pinpoint
the specific traits under selection. Altered ecological or demographic factors can result
in plastic changes in mating systems and ultimately individual strategies (7, 37, 43,
57). When the number of rivals increases (e.g., male-biased OSR) or the encounter
rate with mates declines (e.g., low density), males might transition from territorial
contests to scramble competition (47, 48, 74). In such scenarios, traits like speed,
agility or the ability to locate receptive females could be under selection (74).
While not systematically quantifying them in this study, we did frequently observe
multi-male scrambles ['love knots'; (75)] when a receptive female had been located,
often followed by mating. Consequently, morphological traits linked to speed may also
be under selection like their behavioral counterparts (74). Incidentally, of all

morphological traits assessed here, the only one exhibiting significant positive selection under male-biased OSR was wing length, a trait often thought to be selected under scramble competition due to its link to speed (74). Similarly, small body size has been proposed to be advantageous in scramble competition (2, 74, 92), with selection on it documented in several arthropods (e.g., refs. 93, 94), including multi-male scramblers (75). Here, we did not measure the behavioral traits potentially under selection and found no significant negative selection on body size under male-biased OSR or low density. It is thus possible that reproductive success became more random in those populations due to a more limited number of selection events (95). Considering the pronounced male-biased sexual size dimorphism in *D. prolongata,* which is unique among Drosophilids (96) and uncommon for species reliant on scramble competition (92, 97), scramble competition might play a minor role in its mating system, triggered under relatively extreme (and potentially unnatural) circumstances. Thus, it is possible that a shift toward scramble competition at low densities or under a male-biased OSR does occur, but our treatments were not extreme enough or the sample size too low to conclusively detect it.

Conclusion. Our findings broaden our current understanding of how mating systems respond to complex socio-ecological pressures. The plastic mating strategies observed in *D. prolongata* challenge the notion of static classifications in favor of a dynamic framework, likely beyond this species. This dynamism emphasizes the importance of studying selection across ecological contexts, such as varying OSRs, population densities or resource distributions to understand the intricate interplay between environmental change, population demographics, and the evolutionary process. Incorporating these factors expands our perspective beyond single-environment selection metrics, with important implications. For example, traits like body size may be associated with dominance in contest competition and so assumed to be under directional selection, but this does not mean that the same traits are also advantageous in other contexts [e.g., dynamic scramble competition (74)]. Our study also highlights that integrating multiple indices of sexual selection intensity and complementing these with measures of selection on phenotypic traits offers a far more comprehensive picture of how sexual selection shapes phenotypes. Finally, the observed link between remating intervals and mating success underscores the need to evaluate selection pressures across episodes of selection. Our integrated approach

373 holds the potential to advance our understanding of evolutionary adaptation and pave
374 the way for the development of more robust predictions about how populations will
375 respond to future environmental challenges.
376
377 **Materials and Methods**
378
Study organism. The *D. prolongata* flies used in this study were descendants from
45 isofemale lines, originally collected in the Sa Pa region in Vietnam (70) maintained
in the lab for approx. 30 generations at 20±1°C and 60% relative humidity, on a 14:10
light:dark cycle. To avoid potential inbreeding effects and maximize genetic diversity
within each experimental unit, virgin males and females were crossed between
independent lines, with varying heterozygous genotypes between units. To ensure
virginity of experimental individuals, flies were collected by aspiration on the day of
eclosion before being kept in sex-specific vials (groups of 5 males or 10 females of the
same parental genotypes) on standard fly food (see *SI Methods*) until sexually mature
at 5-7 days post-eclosion. For identification, the thorax of each experimental fly was
marked with a dot of acrylic paint (Kreul El Greco Acrylic colors) in a unique color
within sex and experimental unit, three days before the start of the experiment.
Thereafter, flies were housed individually in 20mL vials with approx. 2mL of fly food.
392
Experimental units and procedure. Each experimental arena (see *SI Methods*)
consisted of a Petri dish (ø 9cm) with small ventilation holes in the lid. Petri dishes
were filled with orange juice-agar medium (see *SI Methods*) around a central Petri dish
(ø 3.5cm) that was filled with standard fly food and covered with yeast paste. This
central yeast patch served a resource-dense area (RDA), whilst the peripheral agar
medium was considered resource-poor, with small grains of dry yeast sprinkled at a
low density (Fig. S1). Each experimental unit (group of flies) was housed in one of
these Petri dishes, varying in their operational sex ratio (OSR; 1:1 or 3:1 M:F) and
population density (low [12 flies], medium [16], high [20]). Each group composition was
replicated five times.
Flies were introduced into the plates by aspiration according to the density and sex
ratio of their treatment. There was neither an overlap of isoline crosses within a plate
nor repeated color within either sex of a plate. Each plate was placed under a
Raspberry Pi 3 model B attached to a camera module (V2) in a climate chamber at

20°C and 60% humidity with a 14:10 light:dark cycle, and fans near the plates
prevented condensation of the lids. Units were photographed from above every two
seconds during the light period across five days. On days two to five, groups of flies
were moved to fresh plates within the first hour of light. All eggs and larvae of the
previous plate were transferred to a fresh plate containing standard fly food and left to
develop for 3-4 days before being collected in individual Eppendorf tubes and stored
at -20°C for later molecular parentage analysis. On the morning after the fifth night, all
experimental flies were removed and anesthetized on ice before separately storing
their heads and bodies in Eppendorf tubes at -20°C for later genotyping and
morphometry, respectively.

Morphological measurements. For each male and female, thorax length was
measured, using the reticular eyepiece of a Leica M60 stereoscope. In males, the left
foreleg and wing were removed using forceps and mounted on a microscope slide in
a drop of Euparal underneath a coverslip. These body parts were then photographed
at 25x magnification using a sCMOS Microscope camera K5 on a Leica M165 FC
stereoscope to measure wing length and total foreleg length using Fiji (98) (details in
SI Methods). Male thorax, wing and foreleg length were further combined in a principal
component analysis, with the first principal component (PC1) serving as a proxy of
overall body size (see *SI Methods*).

Collection of behavioral data. The images were visually screened for matings. For
each mating, we recorded the male and female ID's, along with the start and end times
to calculate mating durations and remating intervals. We further noted whether
copulations started in the RDA, whether they were attained by an alternative
reproductive tactic ["interception", (78)], aggression or courtship, and whether they
were preceded by leg vibration (see *SI Methods* for details). In addition to quantifying
matings and accompanying behaviors, we recorded every 10 minutes which
individuals were present in the RDA (i.e. at least their head or the thorax and abdomen
were in the RDA).

Parentage analysis. The parentage analysis was based on six microsatellite loci that
we previously determined to be highly variable among isolines (details in *SI Methods*).
The larvae or heads of parents were suspended in KAWA buffer (10 mM Tris HCl, 1

mM EDTA, 0.5% Tween 20, 50 µg/ml Proteinase K, pH 8.0) and sent to Ecogenics
(Microsynth) for DNA extraction, PCR amplification and sequencing. Approximately
50% of all larvae from each plate and day were analyzed, selected at random.
Parentage assignment was done using CERVUS 3.0.7 (99), which determines the
most likely parent-offspring trio by maximum likelihood exclusion based on LOD
(logarithm of the odds) scores and their probabilities by simulations on the allele
frequencies in the population. Parent-offspring trios were accepted as valid when at
least three loci were compared (excluded when more than three were missing), and
when these were assigned with a likelihood of 95% ("strict") or 80% ("relaxed"). As all
potential parents were known and combinations could be cross-checked with
observed matings, trios with 80% likelihood were included.
452
Statistical analyses. All statistical analyses were conducted in R version 4.3. (100).
Linear and generalized linear mixed models were conducted using the package *lme4*
(101) with experimental units as a random effect. Where the analysis was done on the
level of matings instead of individuals, the male and female ID were added as random
effects. Negative binomial model [*glmmTMB* (102)] or observation level random effects
(103) were used to control for overdispersion in Poisson or binomial models,
respectively. We bootstrapped confidence intervals (n = 1000) for the opportunity for
sexual selection and for the variance partitioning using the package *sampling* (104).
We used the package *piecewiseSEM* (105) for piecewise structural equation modeling
and *diagrammeR* (106) to visualize the structural equation models. A comprehensive
description of all statistical methods can be found in the *SI Methods*.
464

465   **Acknowledgments**
466
467   We thank Florin Allmendinger, Sina Lerch, Monica Anderson Berdal, Sonja Sbilordo
468   and Jeannine Roy for their support in the lab.
469   This project was supported by the Swiss National Science Foundation (grants
470   310030_197651 and PP00P3_202662 to S.L.) and by the University of Zurich (FK-20-
471   088 to A.D.N.).

472
473
474

References

1. C. Darwin, *The descent of man and selection in relation to sex* (Murray, 1871).

2. M. Andersson, *Sexual selection* (Princeton University Press, 1994).

3. T. Janicke, I. K. Häderer, M. J. Lajeunesse, N. Anthes, Darwinian sex roles confirmed across the animal kingdom. *Sci. Adv.* **2**, e1500983 (2016).

4. T. Clutton-Brock, Sexual selection in females. *Anim. Behav.* **77**, 3–11 (2009).

5. R. M. Hare, L. W. Simmons, Sexual selection and its evolutionary consequences in female animals. *Biol. Rev.* **94**, 929–956 (2019).

6. J. Knight, Sexual stereotypes. *Nature* **415**, 254–256 (2002).

7. S. T. Emlen, L. W. Oring, Ecology, sexual selection, and the evolution of mating systems. *Science* **197**, 215–223 (1977).

8. M. Jirotkul, Operational sex ratio influences female preference and male–male competition in guppies. *Anim. Behav.* **58**, 287–294 (1999).

9. S. Wacker, *et al.*, Operational sex ratio but not density affects sexual selection in fish: Effects of OSR and density on sexual selection. *Evolution* **67**, 1937–1949 (2013).

10. T. Janicke, E. H. Morrow, Operational sex ratio predicts the opportunity and direction of sexual selection across animals. *Ecol. Lett.* **21**, 384–391 (2018).

11. M. L. Head, A. T. Kahn, J. M. Henshaw, J. S. Keogh, M. D. Jennions, Sexual selection on male body size, genital length and heterozygosity: Consistency across habitats and social settings. *J. Anim. Ecol.* **86**, 1458–1468 (2017).

12. M. L. Head, A. K. Lindholm, R. Brooks, Operational sex ratio and density do not affect directional selection on male sexual ornaments and behavior. *Evolution* **62**, 135–144 (2007).

13.   A. G. Jones, G. Rosenqvist, A. Berglund, J. C. Avise, The measurement of sexual
selection using Bateman's principles: An experimental test in the sex-role-
reversed pipefish *Syngnathus typhle*. *Integr. Comp. Biol.* **45**, 874–884 (2005).

14.   P. S. Fitze, J.-F. Le Galliard, Operational sex ratio, sexual conflict and the
intensity of sexual selection. *Ecol. Lett.* **11**, 432–439 (2008).

15.   I. Klemme, H. Ylönen, J. A. Eccard, Reproductive success of male bank voles
(*Clethrionomys glareolus*): the effect of operational sex ratio and body size.
Behav. Ecol. Sociobiol. **61**, 1911–1918 (2007).

16.   S. C. Mills, A. Grapputo, E. Koskela, T. Mappes, Quantitative measure of sexual
selection with respect to the operational sex ratio: a comparison of selection
indices. *Proc. R. Soc. B Biol. Sci.* **274**, 143–150 (2007).

17.   G. Arnqvist, The effects of operational sex ratio on the relative mating success
of extreme male phenotypes in the water strider *Gerris odontogaster* (Zett.)
(Heteroptera; Gerridae). *Anim. Behav.* **43**, 681–683 (1992).

18.   H. Klug, J. Heuschele, M. D. Jennions, H. Kokko, The mismeasurement of sexual
selection. *J. Evol. Biol.* **23**, 447–462 (2010).

19.   R. Rios Moura, P. E. C. Peixoto, The effect of operational sex ratio on the
opportunity for sexual selection: a meta-analysis. *Anim. Behav.* **86**, 675–683
(2013).

20.   G. A. Parker, M. A. Ball, P. Stockley, M. J. G. Gage, Sperm competition games:
individual assessment of sperm competition intensity by group spawners. *Proc.
R. Soc. Lond. B Biol. Sci.* **263**, 1291–1297 (1996).

21.   U. Candolin, J. D. Reynolds, Adjustments of ejaculation rates in response to risk
of sperm competition in a fish, the bitterling (*Rhodeus sericeus*). *Proc. R. Soc.
Lond. B Biol. Sci.* **269**, 1549–1553 (2002).

22.   A. Rondeau, B. Sainte-Marie, Variable mate-guarding time and sperm allocation
by male snow crabs (*Chionoecetes opilio*) in response to sexual competition,

and their impact on the mating success of females. *Biol. Bull.* **201**, 204–217
(2001).

23. L. K. Weir, J. W. A. Grant, J. A. Hutchings, The influence of operational sex ratio
on the intensity of competition for mates. *Am. Nat.* **177**, 167–176 (2011).

24. F. Garcia-Gonzalez, M. Gomendio, Adjustment of copula duration and ejaculate
size according to the risk of sperm competition in the golden egg bug
(*Phyllomorpha laciniata*). *Behav. Ecol.* **15**, 23–30 (2004).

25. J. P. Moatt, C. Dytham, M. D. F. Thom, Sperm production responds to perceived
sperm competition risk in male *Drosophila melanogaster*. *Physiol. Behav.* **131**,
111–114 (2014).

26. S. A. Ramm, *et al.*, Sperm competition risk drives plasticity in seminal fluid
composition. *BMC Biol.* **13**, 87 (2015).

27. A. Jarrige, D. Riemann, M. Goubault, T. Schmoll, Strategic sperm allocation in
response to perceived sperm competition risk in a lekking insect. *Anim. Behav.*
109, 81–87 (2015).

28. J. R. Linklater, B. Wertheim, S. Wigby, T. Chapman, Ejaculate depletion patterns
evolve in response to experimental manipulation of sex ratio in *Drosophila*
melanogaster. *Evolution* **61**, 2027–2034 (2007).

29. Y. Gao, L. Kang, Operational Sex Ratio and Alternative Reproductive Behaviours
in Chinese Bushcricket, *Gampsocleis gratiosa*. *Ethology* **112**, 325–331 (2006).

30. K. U. Wensing, M. Koppik, C. Fricke, Precopulatory but not postcopulatory male
reproductive traits diverge in response to mating system manipulation in
Drosophila melanogaster. *Ecol. Evol.* **7**, 10361–10378 (2017).

31. M. Reuter, *et al.*, Adaptation to experimental alterations of the operational sex
ratio in populations of *Drosophila melanogaster*. *Evolution* **62**, 401–412 (2008).

32. Y. L. Cai, C. L. Mai, X. Yu, W. B. Liao, Effect of population density on relationship
between pre- and postcopulatory sexual traits. *Anim. Biol.* **69**, 281–292 (2019).

33. T. S. Chechi, Z. Ali Syed, N. G. Prasad, Virility does not imply immensity: Testis
size, accessory gland size and ejaculate depletion pattern do not evolve in
response to experimental manipulation of sex ratio in *Drosophila melanogaster*.
J. Insect Physiol. **98**, 67–73 (2017).

34. H. S. Crudgington, S. Fellows, N. S. Badcock, R. R. Snook, Experimental
manipulation of sexual selection promotes greater male mating capacity but does
not alter sperm investment. *Evolution* **63**, 926–938 (2009).

35. K. B. McNamara, *et al.*, Male-biased sex ratio does not promote increased sperm
competitiveness in the seed beetle, *Callosobruchus maculatus*. *Sci. Rep.* **6**,
28153 (2016).

36. S. Lüpold, L. Jin, W. B. Liao, Population density and structure drive differential
investment in pre- and postmating sexual traits in frogs. *Evolution* **71**, 1686–1699
(2017).

37. H. Kokko, D. J. Rankin, Lonely hearts or sex in the city? Density-dependent
effects in mating systems. *Philos. Trans. R. Soc. B Biol. Sci.* **361**, 319–334
(2006).

38. L. Winkler, R. Eilhardt, T. Janicke, Population density affects sexual selection in
an insect model. *Funct. Ecol.* **37**, 2734–2747 (2023).

39. C. M. House, J. Rapkin, J. Hunt, D. J. Hosken, Operational sex ratio and density
predict the potential for sexual selection in the broad-horned beetle. *Anim.*
Behav. **152**, 63–69 (2019).

40. R. D. Cothran, D. Schmidenberg, A. R. Stiff, G. A. Wellborn, R. A. Relyea, Effects
of density on the strength of sexual selection in the laboratory and in nature. *Biol.*
J. Linn. Soc. **140**, 504–517 (2023).

41. C. C. Taff, C. R. Freeman-Gallant, P. O. Dunn, L. A. Whittingham, Spatial
distribution of nests constrains the strength of sexual selection in a warbler. *J.*
Evol. Biol. **26**, 1392–1405 (2013).

42.   M. Jirotkul, Population density influences male–male competition in guppies.
Anim. Behav. **58**, 1169–1175 (1999).

43.   D. McLain, Population density and the intensity of sexual selection on body
length in spatially or temporally restricted natural populations of a seed bug.
Behav. Ecol. Sociobiol. **30** (1992).

44.   J. I. Morales-Mata, J. Potti, C. Camacho, J. Martínez-Padilla, D. Canal,
Phenotypic selection on an ornamental trait is not modulated by breeding density
in a pied flycatcher population. *J. Evol. Biol.* **35**, 610–620 (2022).

45.   J. Conner, Density-dependent sexual selection in the fungus beetle, *Bolitotherus*
cornutus. Evolution **43**, 1378–1386 (1989).

46.   N. P. Sharp, A. F. Agrawal, Mating density and the strength of sexual selection
against deleterious alleles in *Drosophila melanogaster*. *Evolution* **62**, 857–867
(2008).

47.   E. L. McCullough, B. A. Buzatto, L. W. Simmons, Population density mediates
the interaction between pre- and postmating sexual selection. *Evolution* **72**, 893–
905 (2018).

48.   R. J. Knell, Population density and the evolution of male aggression. *J. Zool.* **278**,
83–90 (2009).

49.   C. D. Kelly, The interrelationships between resource-holding potential, resource-
value and reproductive success in territorial males: How much variation can we
explain? *Behav. Ecol. Sociobiol.* **62**, 855–871 (2008).

50.   E. A. Lacey, J. R. Wieczorek, Territoriality and male reproductive success in
arctic ground squirrels. *Behav. Ecol.* **12**, 626–632 (2001).

51.   M. E. Roithmair, Territoriality and male mating success in the dart-poison frog,
Epipedobates femoralis (Dendrobatidae, Anura). *Ethology* **92**, 331–343 (1992).

52.   G. N. Dodson, Resource defense mating system in antlered flies, *Phytalmia spp.*
(Diptera: Tephritidae). *Ann. Entomol. Soc. Am.* **90**, 496–504 (1997).

53. G. N. Dodson, "Behavior of the Phytalmiinae and the evolution of antlers in tephritid flies" in *Fruit Flies (Tephritidae): Phylogeny and Evolution of Behavior*, (CRC Press, 1999).

54. E. Batsuren, *et al.*, Density-dependent changes of mating system and family structure in Brandt's voles (*Lasiopodomys brandtii*). *Ecol. Evol.* **12**, e9199 (2022).

55. P. G. Byrne, J. D. Roberts, Intrasexual selection and group spawning in quacking frogs (*Crinia georgiana*). *Behav. Ecol.* **15**, 872–882 (2004).

56. J. Radwan, J. Unrug, J. L. Tomkins, Status-dependence and morphological trade-offs in the expression of a sexually selected character in the mite, *Sancassania berlesei*. *J. Evol. Biol.* **15**, 744–752 (2002).

57. M. Reichard, C. Smith, W. C. Jordan, Genetic evidence reveals density-dependent mediated success of alternative mating behaviours in the European bitterling (*Rhodeus sericeus*). *Mol. Ecol.* **13**, 1569–1578 (2004).

58. G. Raihani, M. A. Serrano-Meneses, A. Córdoba-Aguilar, Male mating tactics in the American rubyspot damselfly: territoriality, nonterritoriality and switching behaviour. *Anim. Behav.* **75**, 1851–1860 (2008).

59. K. Seidelmann, Territoriality is just an option: allocation of a resource fundamental to the resource defense polygyny in the European wool carder bee, *Anthidium manicatum* (Hymenoptera: Megachilidae). *Behav. Ecol. Sociobiol.* **75**, 138 (2021).

60. B. A. Buzzatto, J. L. Tomkins, L. W. Simmons, "Alternative phenotypes within mating systems" in *The Evolution of Insect Mating Systems*, (Oxford University Press, 2014).

61. J. L. Brown, The evolution of diversity in avian territorial systems. *Wilson Bull.* **76**, 160–169 (1964).

62. J. W. A. Grant, Whether or not to defend? The influence of resource distribution. *Mar. Behav. Physiol.* **23**, 137–153 (1993).

63.   S. C. Mills, J. D. Reynolds, Operational sex ratio and alternative reproductive
behaviours in the European bitterling, *Rhodeus sericeus*. *Behav. Ecol. Sociobiol.*
54, 98–104 (2003).

64.   S. Setoguchi, *et al.*, Sexual dimorphism and courtship behavior in *Drosophila*
prolongata. *J. Ethol.* **32**, 91–102 (2014).

65.   K. Amino, T. Matsuo, Reproductive advantage of the winners of male-male
competition in *Drosophila prolongata*. *Behav. Proc.* **206**, 104831 (2023).

66.   K. Amino, T. Matsuo, Intra- versus inter-sexual selection on sexually dimorphic
traits in *Drosophila prolongata*. *Zool. Sci.* **37**, 210 (2020).

67.   A. Kudo, H. Takamori, H. Watabe, Y. Ishikawa, T. Matsuo, Variation in
morphological and behavioral traits among isofemale strains of *Drosophila*
prolongata (Diptera: Drosophilidae). *Entomol. Sci.* **18**, 221–229 (2015).

68.   K. Minekawa, K. Amino, T. Matsuo, A courtship behavior that makes
monandrous females polyandrous. *Evolution* **74**, 2483–2493 (2020).

69.   K. Minekawa, T. Miyatake, Y. Ishikawa, T. Matsuo, The adaptive role of a
species-specific courtship behaviour in coping with remating suppression of
mated females. *Anim. Behav.* **140**, 29–37 (2018).

70.   J. Perdigón Ferreira, S. Lüpold, Condition- and context-dependent alternative
reproductive tactic in *Drosophila prolongata*. *Behav. Ecol.* **33**, 213–221 (2022).

71.   S. J. Arnold, Bateman's principles and the measurement of sexual selection in
plants and animals. *Am. Nat.* **144**, S126–S149 (1994).

72.   M. J. Wade, Sexual selection and variance in reproductive success. *Am. Nat.*
114, 742–747 (1979).

73.   E. L. McCullough, D. J. Emlen, Evaluating the costs of a sexually selected
weapon: big horns at a small price. *Anim. Behav.* **86**, 977–985 (2013).

74.   M. E. Herberstein, C. J. Painting, G. I. Holwell, "Scramble Competition Polygyny
in Terrestrial Arthropods" in *Advances in the Study of Behavior*, (Elsevier, 2017),
pp. 237–295.

75.   F. V. Vencl, A. D. Carlson, Proximate Mechanisms of Sexual Selection in the
Firefly *Photinus pyralis* (Coleoptera: Lampyridae). *J. Insect Behav.* **11**, 191–207
(1998).

76.   A. H. Krakauer, M. S. Webster, E. H. Duval, A. G. Jones, S. M. Shuster, The
opportunity for sexual selection: not mismeasured, just misunderstood. *J. Evol.*
Biol. **24**, 2064–2071 (2011).

77.   N. Toyoshima, T. Matsuo, Fight outcome influences male mating success in
Drosophila prolongata. *J. Ethol.* **41**, 119–127 (2023).

78.   S. Setoguchi, A. Kudo, T. Takanashi, Y. Ishikawa, T. Matsuo, Social context-
dependent modification of courtship behaviour in *Drosophila prolongata*. *Proc.*
R. Soc. B Biol. Sci. **282**, 20151377 (2015).

79.   A. N. De Nardo, J. Roy, S. H. Sbilordo, S. Lüpold, Condition-dependent
interaction between mating success and competitive fertilization success in
Drosophila melanogaster *. *Evolution* **75**, 2014–2026 (2021).

80.   E. L. Macartney, *et al.*, Sperm depletion in relation to developmental nutrition and
genotype in *Drosophila melanogaster*. *Evolution* **75**, 2830–2841 (2021).

81.   H. Liu, E. Kubli, Sex-peptide is the molecular basis of the sperm effect in
Drosophila melanogaster. *Proc. Natl. Acad. Sci.* **100**, 9929–9933 (2003).

82.   A. Bretman, C. Fricke, T. Chapman, Plastic responses of male *Drosophila*
melanogaster to the level of sperm competition increase male reproductive
fitness. *Proc. R. Soc. B Biol. Sci.* **276**, 1705–1711 (2009).

83.   S. Wigby, *et al.*, Seminal fluid protein allocation and male reproductive success.
Curr. Biol. **19**, 751–757 (2009).

84. V. Zeender, S. Pfammatter, B. Roschitzki, S. Dorus, S. Lüpold, Genotype-by-
environment interactions influence the composition of the *Drosophila* seminal
proteome. *Proc. R. Soc. B Biol. Sci.* **290**, 20231313 (2023).

85. K. M. Fedorka, W. E. Winterhalter, B. Ware, Perceived sperm competition
intensity influences seminal fluid protein production prior to courtship and mating.
Evolution **65**, 584–590 (2011).

86. B. R. Hopkins, *et al.*, Divergent allocation of sperm and the seminal proteome
along a competition gradient in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci.*
116, 17925–17933 (2019).

87. L. K. Sirot, M. F. Wolfner, "Who's zooming who? Seminal fluids and cryptic
female choice in diptera" in *Cryptic Female Choice in Arthropods*, A. V. Peretti,
A. Aisenberg, Eds. (Springer International Publishing, 2015).

88. R. C. Firman, C. Gasparini, M. K. Manier, T. Pizzari, Postmating female control:
20 years of cryptic female choice. *Trends Ecol. Evol.* **32**, 368–382 (2017).

89. J. Morimoto, *et al.*, Sex peptide receptor-regulated polyandry modulates the
balance of pre- and post-copulatory sexual selection in *Drosophila*. *Nat.*
Commun. **10**, 283 (2019).

90. S. Pitnick, Male size influences mate fecundity and remating interval in
Drosophila melanogaster. *Anim. Behav.* **41**, 735–745 (1991).

91. B. R. Hopkins, J. C. Perry, The evolution of sex peptide: sexual conflict,
cooperation, and coevolution. *Biol. Rev.* **97**, 1426–1448 (2022).

92. M. T. Ghiselin, *The Economy of nature and the evolution of sex* (University of
California press, 1974).

93. C. D. Kelly, Sexual selection on size and shape in Japanese beetles (*Popillia*
japonica). *Behav. Ecol.* **31**, 1073–1083 (2020).

94. J. Moya-Laraño, M. E. T. El-Sayyid, C. W. Fox, Smaller beetles are better
scramble competitors at cooler temperatures. *Biol. Lett.* **3**, 475–478 (2007).

95.　H. Klug, L. Stone, More than just noise: Chance, mating success, and sexual
selection. *Ecol. Evol.* **11**, 6326–6340 (2021).

96.　P. T. Rohner, *et al.*, Interrelations of global macroecological patterns in wing and
thorax size, sexual size dimorphism, and range size of the Drosophilidae.
Ecography **41**, 1707–1717 (2018).

97.　W. U. Blanckenhorn, R. F. Preziosi, D. J. Fairbairn, Time and energy constraints
and the evolution of sexual size dimorphism — to eat or to mate? *Evol. Ecol.* **9**,
369–381 (1995).

98.　J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis.
Nat. Methods **9**, 676–682 (2012).

99.　S. T. Kalinowski, M. L. Taper, T. C. Marshall, Revising how the computer
program CERVUS accommodates genotyping error increases success in paternity
assignment. *Mol. Ecol.* **16**, 1099–1106 (2007).

100.　R Core Team, R: A Language and Environment for Statistical Computing. (2023).
Deposited 2023.

101.　D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models
using *lme4*. *J. Stat. Softw.* **67** (2015).

102.　M. Brooks E., *et al.*, glmmTMB balances speed and flexibility among packages
for zero-inflated generalized linear mixed modeling. *R J.* **9**, 378 (2017).

103.　X. A. Harrison, A comparison of observation-level random effect and beta-
binomial models for modelling overdispersion in binomial data in ecology &
evolution. *PeerJ* **3**, e1114 (2015).

104.　Y. Tillé, Matei, sampling: Survey Sampling. (2023). Deposited 2023.

105.　J. S. Lefcheck, piecewiseSEM: Piecewise structural equation modelling in r for
ecology, evolution, and systematics. *Methods Ecol. Evol.* **7**, 573–579 (2016).

106.　R. Iannone, DiagrammeR: Graph/Network Visualization. (2023). Deposited
2023.

Figures

Figure 1: Bateman gradients for males (blue) and females (red) for each combination

of density (low, medium, high) and OSR (even, male-biased) and density.

Figure 2: Linear selection gradients with 95% confidence intervals on male
morphological traits under different levels of population density and OSRs.

752 Figure 3: The effect of territoriality (time on the resource-dense area; left), and PC1
753 (middle) on mating success depends on the OSR. The effect of PC1 on courtship
754 occurrence also depends on the OSR (right).
755

756

757   Figure 4: Visualized structural equation model of male reproductive dynamics under
758   (a) an even OSR. Green and red indicate positive and negative relationships,
759   respectively. Black lines indicate neutral relationships. Dashed lines (red or green)
760   indicate trends ($P < 0.1$).
761

762   a)   b)