1 **Socio-ecological context modulates significance of territorial contest**
2 **competition in *Drosophila prolongata***
3

4 Alessio N. De Nardo[1]*, Broti Biswas[1,2], Jhoniel Perdigón Ferreira[1], Abhishek Meena[1],
5 Stefan Lüpold[1]
6
7 [1]Department of Evolutionary Biology and Environmental Studies, University of Zurich,
8 Winterthurerstrasse 190, 8057 Zurich, Switzerland
9 [2]Department of Systematic and Evolutionary Botany, University of Zurich,
10 Zollikerstrasse 107, 8008 Zurich, Switzerland
11
12 *Correspondence: alessio.denardo@uzh.ch
13

14 **Abstract**
15

The intensity and direction of sexual selection is intricately linked to the social and
ecological context. Both operational sex ratios (OSRs) and population densities can
affect the ability of males to monopolize resources and mates, and thus the form and
intensity of sexual selection on them. Here, we studied how the mating system of the
promiscuous and strongly sexually dimorphic fruit fly *Drosophila prolongata* responds
to changes in the OSR and population density. We recorded groups of flies over five
days and quantified territory occupancy, mating success, and competitive fertilization
success. Although sexual selection was stronger under male-biased than even OSRs
but unrelated to density, realized selection on morphological traits was higher under
even OSRs and increased with density. Larger and more territorial males achieved
both higher mating success and competitive fertilization success, but only under even
OSRs. Our combined results also support a shift in the mating system from territorial
contest competition to scramble competition under male-biased OSRs and potentially
at low density, where there was no clear contribution of the measured traits to
reproductive success. Our study emphasizes the limitations of traditional selection
metrics and the role of the socio-ecological context in predicting adaptation to a
changing environment.
33
34

Significance

Mating systems are complex and dynamic, adapting to ongoing ecological change. Studies often assume that changes in the socio-ecological context alter the intensity of sexual selection on traits indicating individual fitness, but our work on *Drosophila prolongata* challenges this view. By manipulating operational sex ratio and population density, and jointly investigating territorial behavior and both pre- and post-mating reproductive success over several days, we reveal the plastic mating strategies in this fly. This dynamism underscores the limitations of static classifications and the importance of studying selection across diverse socio-ecological contexts. This broader perspective advances our understanding of the tight connections between environmental change, population demographics, and the evolutionary process.

Introduction

Sexual selection, driven by competition for mates and/or fertilization (1), shapes heritable traits and behaviors that are directly related to reproductive success (2). While sexual selection can operate in both sexes, it tends to be stronger on males than females (2, 3). Much of this asymmetry in sexual selection is attributed to the typically greater investments per offspring by females compared to males, which reduces their reproductive potential and intensifies the competition among males over the limited reproductive opportunities (4, 5). The direction and intensity of sexual selection between the sexes can be influenced by the ecological context. For example, food scarcity in katydid bush crickets shifts the typical sex roles to females competing for access to the nutritious spermatophores provided by males (6).

One important ecological parameter affecting the intensity of sexual selection is the distribution and defensibility of resources, which can shape the local population composition and density. For example, variation in the operational sex ratio (OSR; i.e., the ratio of sexually receptive males to sexually receptive females (7)) can affect the availability of potential mates. A skewed OSR can lead to monopolization of mating opportunities by a few successful individuals of the competing sex (7). Male-biased OSRs in particular can thus drive selection on traits that are advantageous in competition, including larger body size, the growth of weaponry, or elaborate ornamentation favored by females (7–9).

A meta-analysis across 62 empirical studies of 58 different species generally
supports theoretical predictions that OSRs can drive the intensity of sexual selection
(10). However, studies exploring selection on specific phenotypic traits often find no
difference in the strength of selection between different OSRs (e.g., on body size (11–
13)), or they show increased selection intensity in even or female-biased populations
(14–16). For example, in the common lizard (*Lacerta vivipara*), females favor small
males under male-biased OSRs as large males leave more bite scars on them during
mating (14, 17). Overall, the effect of the OSR on the intensity of sexual selection
appears to vary considerably across species and mating systems (18, 19).

Apart from mate competition, OSRs can also influence post-mating processes, such
as competitive fertilization (20–22; but see 23), ultimately impacting aspects like
ejaculate transfer or mating duration (24). Males tend to invest more in ejaculate
production when exposed to a higher risk or intensity of sperm competition (20), often
induced by changes in the perceived OSR (25–27). Thus, traits favored in the post-
mating phase, such as enhanced ejaculate allocation (28) or larger testes (29) are
expected to be under stronger selection when OSRs are male-biased, yet studies on
the latter are inconclusive (30–36). In a literature review (30), only one of four studies
found a significant effect of OSR on testes size. This one study showed an increase
in testes size for male *Drosophila melanogaster* evolving under an extreme female
bias (10:1), suggesting selection against sperm limitation rather than sperm
competition (31).

In addition to the OSR, competition for mates can also vary with population density.
Theoretical work suggests that higher population densities increase the encounter rate
both between potential mates and competitors, which can in turn intensify sexual
selection (37). By contrast, the scope for sexual selection may be more limited at low
densities, where infrequent mate encounters may lower a female's mate discrimination
due to higher sampling costs (37–39), and rare encounters among males also limit the
potential for mate competition (37). Empirical studies on these patterns, however, have
yielded mixed results (9, 40–48). Whereas some studies point toward more intense
sexual selection at higher densities (40–42), others report the opposite (44–46) or no
relationship (9, 47, 48). One possible reason for these contrasting results is that the
traits examined in these studies are under pre-mating sexual selection, which might
trade off with traits under post-mating sexual selection in a density-dependent manner
(36, 49). For example, higher population densities might increase the relative

importance of post-mating competition if increasing costs of monopolization lower the
effectiveness of aggressive strategies (36, 37, 45, 50, 51). Such a trend is suggested
across frogs and toads, where the trade-off between forelimb muscles and testes size
is biased more strongly toward the latter as population densities increase and
competitive fertilization becomes more difficult for males to prevent (36).

Sexual selection and male reproductive tactics may also be influenced by the
distribution of resources (7). While evenly distributed resources limit the opportunity
for monopolization, highly competitive males may gain control over spatially
concentrated resources (7). These males may then become more attractive to
potential mates for their ability to defend a resource or for the benefit the resource
provides, tying mating success to resource monopolization (52–54). For example, in
antlered flies, *Phytalmia spp.,* males fight over oviposition sites to which females then
gain access through copulation (55). Therefore, females indirectly choose mates with
large antlers, which in turn benefit males in these contests (56). However, reproductive
tactics may vary between, or temporally within, individuals (57), mediated by the
number of individuals competing for a resource and how well it can be defended (58,
59). Males may follow a territorial or a non-territorial strategy, depending on the relative
benefits of each for a given competitor density (45, 51, 60) or their own resource-
holding potential (52), or they may employ a flexible strategy (61, 62). Similarly, if the
population density or the physical condition of a male limit his reproductive success by
courting or competing for females, he may adopt an alternative tactic, such as
attempting to steal copulations from courting males (63–66).

Although OSRs and population densities have been shown to shape sexual
selection and mating systems, both separately and jointly (e.g., 8, 9, 40, 41, 44, 49,
67), it remains unknown how these factors combined drive both pre- and post-
mating sexual selection, particularly in the context of uneven resource distribution.
Here, we investigated these factors in a comprehensive laboratory experiment using
the fruit fly *Drosophila prolongata.* This South-Asian species exhibits conspicuous
male-biased sexual dimorphism both in body size and in foreleg size and coloration
(68), while also showing resource-defense behavior when high-quality resources are
concentrated (69). Males use their forelegs in both fights against competitors (70, 71)
and elaborate courtship behaviors (68). These behaviors also include "leg vibration",
a species-specific behavior that involves a male stimulating the female abdomen and
is often displayed immediately before copulation (71), likely to overcome female

remating resistance (72, 73). Leg vibration, however, can also trigger an alternative
reproductive tactic by a nearby male, namely interception of the female (66). With
highly polygamous females (72), this species is ideal for studies of context-dependent
pre- and post-mating sexual selection.

Here, we monitored over a five-day interaction period each of 30 groups of
D. prolongata that varied in their sex ratio (1:1 or 3:1 males:females) and density (12,
16, or 20 flies). Over five days, we quantified the distribution of flies around a centrally
located resource, their pre-mating behaviors, individual copulations, and molecular
parentage, which provided a uniquely comprehensive understanding of reproductive
dynamics in this species. We predicted the intensity of sexual selection on males to
increase (i) under a male-biased OSR and (ii) with population density, translating into
intensifying selection on traits mediating resource defense and mate competition, such
as male body size or foreleg length. We also predicted that (iii) male-biased OSRs and
high densities should increase the relative importance of post- relative to pre-mating
sexual selection.

Results and Discussion

Descriptive statistics and skew in reproductive success. Across all 180 females
and 300 males, we observed 754 matings, with only three females (1.6%) and 83
males (27%) never mating. Most copulations occurred on the first experimental day
(67.6%), with another 13%, 10.5%, 5.7%, and 3.2% of matings on days 2–5,
respectively. Across all five days, females mated 0–13 times (mean±SD = 3.95±2.52),
with up to 7 different sires (2.36±1.39; Fig. 1). Males gained 0–14 matings (2.50±2.6)
across up to 9 females (1.47±1.59; Fig. 1). Varying numbers of possible matings
between treatments hampered direct comparisons of absolute mating numbers.
However, relative male mating success was more heavily skewed under male-biased
compared to even OSRs despite a similar mean success rate (Fig. S1), suggesting
more intense sexual selection under a male-biased OSR. These differences were
more subtle between densities, but generally suggested weaker sexual selection at
low compared to medium or high densities (Fig. S2).

Potential and realized sexual selection. To quantify the selection dynamics in *D. prolongata* under different population compositions, we measured Bateman gradients and the opportunity for sexual selection (I_s) as two commonly used estimates for the intensity of sexual selection (10). Whereas the former represents the slope between individuals' number of mates and resulting offspring (i.e., the fitness advantage gained by additional matings (74)), the latter is based on the sex-specific variances in reproductive success (i.e., the opportunity for selection to act on variation in reproductive success (75)). The Bateman gradients for both sexes were positive (Fig. 1; males: $X^2_1 = 63.65$, $n = 193$, $P < 0.001$, Table S1; females: $X^2_1 = 24.1$, $n = 175$, $P < 0.001$, Table S2), but more pronounced in males ($X^2_1 = 7.42$, $P = 0.006$, Table S3a). The disparity in Bateman gradients between the sexes was greater in the male-biased populations ($X^2_1 = 5.6$, $n = 368$, $P = 0.018$) but unaffected by group density ($X^2_2 = 0.23$, $P = 0.89$, Table S3b). Similarly, I_s was higher for males compared to females in all treatments, but particularly pronounced in the male-biased populations (6.2× higher for males) compared to the even OSR populations (2.7× higher for males). Population density again had no significant effect (Table S4, Fig. S3). These findings combined suggest that sexual selection is more pronounced on males than females, and its intensity for males is amplified under a male-biased OSR, as previously shown for a diversity of taxa (10).

When comparing selection on specific male traits, results were more complex. For example, we found positive selection gradients on thorax, foreleg and wing length, and composite body size (principal component 1, PC1; see *Methods*) under even OSR, but only on wing length in male-biased populations (Fig. 2, Table S5). Also, contrary to expectations based on I_s, positive selection on all measured traits appeared to intensify with density (Fig. 2, Table S6). This increasing selection on secondary sexual traits with density aligns with theory (37), yet has been reported for only few species (42, 76, 77; but see refs. 45, 78). The changing pattern of selection on phenotypic traits across OSRs, however, contradicts both our predictions and insights gained from the above Bateman gradients and I_s.

The lack of positive selection on putative sexually selected traits in male-biased populations might simply indicate insufficient statistical power or that other, unmeasured behavioral or morphological traits, or traits under post-mating selection would be favored when mates are more difficult to monopolize (2, 79, 80). However, our findings might also reveal the limitations of using metrics like Bateman gradients,

I_s, or the OSR to predict realized selection on specific sexually selected traits, as
previously emphasized (18). As our results confirm, these common measures of the
intensity of sexual selection do not necessarily correlate with the observed selection
on traits and merely set an upper limit to the potential of sexual selection (18). These
results underscore the need to integrate different measures to gain a more complete
understanding of the dynamics of a mating system. Despite their limitations, Bateman
gradients or I_s remain valuable tools for numerically characterizing mating systems.
They consider all types of sexual selection (e.g. pre- and post-mating, intra- and
intersexual), acting on any relevant morphological or behavioral traits (81). Thus, they
accurately capture variation in fitness and allow for comparisons between sexes and
populations (81). For example, we showed that reproductive success is more skewed
toward a few successful males in male-biased populations than under even OSR. The
trait-specific selection gradients indicate, however, that traits selected in male-biased
populations are not the morphological traits expected to be involved in physical contest
such as resource defense (e.g., forelegs). These insights emphasize the complex
interplay between OSR, density, and sexual selection in shaping mating system
dynamics and trait evolution in this species.

Territorial behavior and mating success. To investigate how the territorial behavior
previously reported in this species (69) manifests in different group compositions, a
resource-dense area (RDA) was established in the center of each experimental unit.
This RDA, consisting of highly nutritious yeast paste, covered approximately 16% of
the otherwise nutrient-poor arena. Regardless of population composition, males spent
more time in the RDA than females (35.6% vs. 12.3%; $X^2_1 = 671.6$, $P < 0.001$, $n = 493$
(303 males, 190 females), Table S7), with greater variance in males ($\sigma^2 = 0.016$) than
in females ($\sigma^2 = 0.005$). Hence, the RDA appeared to be highly attractive to males, but
access was unequal among them as reflected by their varying time spent there
(indicating "territorial success"). In addition, individuals of both sexes spent less time
(on average) in the RDA at higher densities ($X^2_1 = 22.0$, $P < 0.001$) and under male-
biased OSRs ($X^2_1 = 21.0$, $P < 0.001$), suggesting intensifying competition for this
resource. Although females spent considerably less time in the RDA overall, 69.4% of
all matings started there. These matings were also more likely to involve aggressive
male-male behavior than matings outside this area ($z = 2.54$, $P = 0.011$, $n = 716$
matings across 212 males and 187 females). Both sexes spent more time in the RDA

on the first day of recording, when most matings occurred, compared to the subsequent days (X^2_4 = 313.724, P < 0.001, n = 2379 across 5 days and 492 individuals; Fig. S4, Table S8). Interestingly, body size (PC1) had no effect on territorial occupancy on the first day, but did so on all following days (interaction: X^2_1 = 5.05, n = 1445, P = 0.02, Table S9), suggesting that males might initially need time to establish dominance hierarchies and resource access, with large males ultimately being more successful (69, 82). Additionally, given the females' heightened receptivity on day one, following females around may have been advantageous to staying in the RDA, also for large males. Indeed, matings were, if anything, less likely to start in the RDA on the first (66.9%) compared to later days (73.8%; z = 1.78, P = 0.075).

Amino & Matsuo (69) report a hierarchy in pairs of males, with one male clearly dominating territory occupancy, particularly when the area is small and easily defendable. Although their findings do not directly demonstrate mating success by territory occupancy, a strong correlation between territory occupancy and courtship rate (69), as well as a mating advantage for winners of contests over territories (83), have been reported for *D. prolongata*. In our study, the RDA was comparable to the larger of the two areas used by Amino & Matsuo (69), but with more males and so reduced ability to monopolize the contested area. Nevertheless, territorial control influenced mating success, albeit with greater complexity at higher numbers of competitors: Compared to 72.5% at an even OSR, only 64.6% (z = 1.85, P = 0.065) of matings were initiated in the RDA at a male-biased OSR. Interestingly, although most matings were initiated in the RDA at both OSRs, overall territory occupancy determined mating success only in even (t = 2.4, P = 0.019) but not male-biased populations (t = 0.24, P = 0.8; OSR×territoriality: t = 1.7, P = 0.09, n = 290, controlling for PC1; Fig. 3, Table S10a-b). Similarly, PC1 was the main contributor to mating success under even (t = 3.7, P < 0.001) but not male-biased OSRs (t = 1.1, P = 0.26, OSR×PC1: t = 2.15, P = 0.03; n = 290, Fig. 3, Table S10a-b). In addition to these results, the occurrence of courtship behavior before mating depended on body size and the OSR (PC1×OSR: t = 2.93, P = 0.003, n = 760, Fig. 3, Table S14a), in that large males showed more courtship behavior in the even OSR (z = 3.20, P = 0.001) but not in the male-biased OSR (z = -1.27, P = 0.20). These results align with the selection gradients above, but additionally illustrate the changing dynamics of sexual selection in response to varying levels of competition.

Our findings support the notion that when competition intensifies (e.g., strongly
male-biased OSR), monopolization of resources becomes increasingly difficult,
potentially shifting mating strategies (36, 37, 49, 50, 64). However, we neither
observed an increase in sneak matings at higher densities nor under male-biased
OSRs (both $|z| < 1.2$, $P > 0.26$, $n = 716$, Table S12). In fact, leg vibration, an auditory
cue used by sneaky males to intercept a mating attempt, even tended to decrease at
high densities ($z = -1.7$, $P = 0.088$) and in male-biased OSRs ($z = 1.89$, $P = 0.059$, $n =$
716, Table S11). Courtship was observed less frequently when matings occurred in
the RDA ($z = -3.3$, $P < 0.001$, Table S14), possibly due to a high local density of
competitors and so higher risk of becoming the target of aggression ($z = 2.54$, $P =$
0.011, $n = 716$, Table S13) or interception ($z = 2.31$, $P = 0.021$, $n = 716$, Table S13).
Similarly, leg vibration was observed less often before matings in the RDA ($z = -1.81$,
$P = 0.07$, $n = 716$, Table S11). Large males were more likely to gain matings following
leg vibration ($z = 2.89$, $P = 0.004$, $n = 716$), whereas those acquiring matings by
interception tended to be smaller males ($t = -2.03$, $P = 0.042$, $n = 716$, Table S12). As
shown previously (84), males avoid leg vibration when the risk of interception is high
(e.g., at high density). Hence, notwithstanding the lesser importance of resource-
defense and body size for mating success in male-biased populations, males did not
adopt alternative mating tactics. This raises the question of whether selection might
instead favor increased investment in post-mating processes.
290
Fertilization success. Across all treatments, male mating success (MS; 48.9%)
explained most of the variation in reproductive success, followed by fertilization
success (FS; 14.3%) and female fecundity (Fec; 6.4%; Fig. S5, Table S16). This
pattern was comparable across social contexts in general, but while mating success
explained three times the variation of fertilization success under an even OSR (MS:
42.5%, FS: 14.4%, Fec: 9.2%), these variance components were more similar under
a male-biased OSR (MS: 37.3%, FS: 25.4%, Fec: 7.2%) despite unchanged female
mating rates ($z = 0.01$, $P = 0.99$). Fertilization success may thus gain in relative
importance when more potential mates compete over each mating opportunity,
hampering female monopolization.
Males that were a female's last mate until the end of the experiment ($t = 4.19$, $P <$
0.001, $n = 190$, Table S16), and males that were able to delay remating for longer

("remating interval", see *SI Methods; t* = 5.00, *P* < 0.001), had higher fertilization
success (adjusted PCS, see *SI Methods*). Analogous to our findings for mating
success, larger males only had higher fertilization success under an even OSR (*t* =
2.68, *P* = 0.008), but not under a male bias (*t* = 0.14, *P* = 0.88; OSR×PC1: *t* = 1.83, *P*
= 0.069, *n* = 190, Table S16). Longer mating durations did not significantly contribute
to fertilization success (*t* = 1.36, *P* = 0.18), but males mating more often had lower
fertilization success per mating (*t* = -2.22, *P* = 0.027), suggesting strategic allocation
(85) or sperm depletion (86).
Female remating intervals were not affected by density (*t* = 0.79, *P* = 0.43), but they
were longer under even than male-biased OSRs (*t* = 1.90, *P* = 0.06, *n* = 190, Table
S17). Copulation duration was positively related with remating intervals (*t* = 1.81, *P* =
0.073), but did not vary with OSR or density (both |*t*| < 1.41, *P* > 0.17, *n* = 710). These
were unexpected to the extent that male *Drosophila* have been shown to transfer more
of the seminal fluid proteins (SFPs) known to suppress female remating when they
perceive a high sperm competition risk (e.g., 87), and this allocation may covary with
copulation duration (88). Our results, however, support previous findings suggesting
a limited role for the control of remating suppression through SFPs in this (72) and
other species with high mating rates (89). If remating intervals are under female
control, females might delay remating until a favored male is found (90, 91). Some
support for this idea comes from the fact that large males were more likely to be the
last male overall (i.e., until the end of the experiment; *t* = 2.25, *P* = 0.026, *n* = 211), but
size did not affect remating intervals up to that point (*t* = 0.05, *P* = 0.95).
In sum, our results indicate that intensifying competition does not augment post-
mating investment by males. Additionally, post-mating selection does not seem to
counteract the lack of pre-mating selection on ostensibly beneficial male traits.
Instead, post-mating processes seem to reinforce the patterns of pre-mating selection.
329
Mating system dynamics. To synthesize the differences in the mating-system
dynamics of *D. prolongata* across different contexts, we built structural equation
models. We provide separate visualizations for each density level in Fig. S6 (Tables
S22-S24), but Fig. 4 presents a simplified overview of the dynamics under even and
male-biased OSRs, respectively. One of the most striking observations was the
absence of many positive (green) or negative (red) effects in the male-biased scenario
(Fig. 4b, Table S21) compared to that under even OSR (Fig. 4a, Table S20). This

underlines how both pre- and post-mating sexual selection vary across social contexts.

For example, variation in body size, a primary determinant of both pre- and post-

mating success under even OSR, seemed inconsequential under male-biased OSRs.

Although we documented a larger skew in reproductive success and a higher

potential for sexual selection in male-biased populations, we were unable to pinpoint

the specific traits under selection. Altered ecological or demographic factors can result

in plastic changes in mating systems and ultimately individual strategies (7, 37, 45,

64). When the number of rivals increases (e.g., male-biased OSR) or the encounter

rate with mates declines (e.g., low density), males might transition from territorial

contests to an alternative mating tactic, potentially a form of scramble competition (49,

50, 79). Scramble competition polygyny is a common mating system in invertebrates

and can manifest in diverse forms (79). At low density, females may be dispersed,

generating selection on the ability to locate receptive females (79). In another type of

scramble competition, individuals of both sexes cluster around resources that cannot

be economically monopolized (e.g., if resource is widely distributed or number of

competitors is high). In this form of scramble competition, it can be a race between

competitors to locate a receptive female in a dense aggregation, possibly combined

with coercion or courtship when the female is reached (79). In such scenarios, traits

like speed, agility, or the ability to locate receptive females could be under

selection (79).

While not systematically quantifying them in this study, we did frequently observe

multi-male clusters ('love knots' (80)) when a receptive female had been located, often

followed by mating. Consequently, morphological traits linked to speed may also be

under selection like their behavioral counterparts (79). Incidentally, of all morphological

traits assessed here, the only one exhibiting significant positive selection under male-

biased OSRs was wing length, a trait often thought to be selected under scramble

competition due to its link to speed (79). Similarly, small body size has been proposed

to be advantageous in scramble competition (2, 79, 92), with selection on it

documented in several arthropods (e.g., 93, 94), including multi-male scramblers (80).

Here, we did not measure the behavioral traits potentially under selection and found

no significant negative selection on body size under male-biased OSR or low density.

However, we have unpublished data after experimentally evolving replicated lines

under different OSRs for 35 generations, which indicate a reduction in male body size

under male-biased relative to even OSR ($t = -6.58$, $n = 681$, $P < 0.001$). In combination

with the results of this study, this divergence may indicate a shift to a mating system
broadly described as scramble competition, or certainly away from territorial contest
competition. It is important to note, however, that contest and scramble competition
are not necessarily separate categories but rather "extremes on a continuum"(79,
p. 247). Considering the pronounced male-biased sexual size dimorphism in *D.*
prolongata, which is unique among Drosophilids (95) and uncommon for species
reliant on scramble competition (92, 96), scramble competition might play a minor role
in this species, triggered primarily under relatively extreme (and potentially unnatural)
circumstances. Whether a shift toward scramble competition would occur at low
densities or under a male-biased OSR remains unclear, as our treatments were not
extreme enough or the sample size too low to conclusively detect it.

Conclusion. Our findings broaden our current understanding of how mating systems
respond to complex socio-ecological pressures. The plastic mating strategies
observed in *D. prolongata* challenge the notion of static classifications in favor of a
dynamic framework, likely beyond this species. This dynamism emphasizes the
importance of studying selection across ecological contexts, such as varying OSRs,
population densities or resource distributions to understand the intricate interplay
between environmental change, population demographics, and the evolutionary
process. Incorporating these factors expands our perspective beyond single-
environment selection metrics, with important implications. For example, traits like
body size may be associated with dominance in contest competition and so assumed
to be under directional selection, but this does not mean that the same traits are also
advantageous in other contexts (e.g., dynamic scramble competition (79)) or where
post-mating selection becomes more pronounced. Our study also highlights that
integrating multiple indices of sexual selection intensity and complementing these with
measures of selection on phenotypic traits offers a far more comprehensive picture of
how sexual selection shapes phenotypes. Finally, by tracking individuals over multiple
mating events we were able to quantify how post- reinforces pre-mating selection and
how the targets of selection change across sequential matings. Our integrated
approach holds the potential to advance our understanding of evolutionary adaptation
and pave the way for the development of more robust predictions about how
populations will respond to future environmental challenges.

Materials and Methods

Study organism. The *D. prolongata* flies used in this study were descendants from 45 isofemale lines, originally collected in the Sa Pa region in Vietnam (66) maintained in the lab for approx. 30 generations at 20±1°C and 60% relative humidity, on a 14:10 light:dark cycle. To avoid potential inbreeding effects and maximize genetic diversity within each experimental unit, virgin males and females were crossed between independent lines, with varying heterozygous genotypes between units. To ensure virginity of experimental individuals, flies were collected by aspiration on the day of eclosion before being kept in sex-specific vials (groups of 5 males or 10 females of the same parental genotypes) on standard fly food (see *SI Methods*) until sexually mature at 5-7 days post-eclosion. For identification, the thorax of each experimental fly was marked with a dot of acrylic paint (Kreul El Greco Acrylic colors) in a unique color within sex and experimental unit, three days before the start of the experiment. Thereafter, flies were housed individually in 20mL vials with approx. 2mL of fly food.

Experimental units and procedure. Each experimental arena (see *SI Methods*) consisted of a Petri dish (ø 9cm) with small ventilation holes in the lid. Petri dishes were filled with orange juice-agar medium (see *SI Methods*) around a central Petri dish (ø 3.5cm) that was filled with standard fly food and covered with yeast paste. This central yeast patch served a resource-dense area (RDA), whilst the peripheral agar medium was considered resource-poor, with small grains of dry yeast sprinkled at a low density (Fig. S7). Each experimental unit (group of flies) was housed in one of these Petri dishes, varying in their operational sex ratio (OSR; 1:1 or 3:1 M:F) and population density (low [12 flies], medium [16], high [20]). Each group composition was replicated five times.

Flies were introduced into the plates by aspiration according to the density and sex ratio of their treatment. There was neither an overlap of isoline crosses within a plate nor repeated color within either sex of a plate. Each plate was placed under a Raspberry Pi 3 model B attached to a camera module (V2) in a climate chamber at 20°C and 60% humidity with a 14:10 light:dark cycle, and fans near the plates prevented condensation of the lids. Units were photographed from above every two seconds during the light period across five days. On days two to five, groups of flies were moved to fresh plates within the first hour of light. All eggs and larvae of the

previous plate were transferred to a fresh plate containing standard fly food and left to

develop for 3-4 days before being collected in individual Eppendorf tubes and stored

at -20°C for later molecular parentage analysis. On the morning after the fifth night, all

experimental flies were removed and anesthetized on ice before separately storing

their heads and bodies in Eppendorf tubes at -20°C for later genotyping and

morphometry, respectively.

Morphological measurements. For each male and female, thorax length was

measured, using the reticular eyepiece of a Leica M60 stereoscope. In males, the left

foreleg and wing were removed using forceps and mounted on a microscope slide in

a drop of Euparal underneath a coverslip. These body parts were then photographed

at 25x magnification using a sCMOS Microscope camera K5 on a Leica M165 FC

stereoscope to measure wing length and total foreleg length using Fiji (97) (details in

SI Methods). Male thorax, wing and foreleg length were further combined in a principal

component analysis, with the first principal component (PC1) serving as a proxy of

overall body size (see *SI Methods*).

Collection of behavioral data. The images were visually screened for matings. For

each mating, we recorded the male and female ID's, along with the start and end times

to calculate mating durations and remating intervals. We further noted whether

copulations started in the RDA, whether they were attained by an alternative

reproductive tactic ("interception", (84)), aggression or courtship, and whether they

were preceded by leg vibration (see *SI Methods* for details). In addition to quantifying

matings and accompanying behaviors, we recorded every 10 minutes which

individuals were present in the RDA (i.e. at least their head or the thorax and abdomen

were in the RDA).

Parentage analysis. The parentage analysis was based on six microsatellite loci that

we previously determined to be highly variable among isolines (details in *SI Methods*).

The larvae or heads of parents were suspended in KAWA buffer (10 mM Tris HCl, 1

mM EDTA, 0.5% Tween 20, 50 µg/ml Proteinase K, pH 8.0) and sent to Ecogenics

(Microsynth) for DNA extraction, PCR amplification and sequencing. Approximately

50% of all 10,667 larvae were selected at random from each plate and day for

genotyping. Parentage assignment was done using CERVUS 3.0.7 (98), which

determines the most likely parent-offspring trio by maximum likelihood exclusion
based on LOD (logarithm of the odds) scores and their probabilities by simulations on
the allele frequencies in the population. Parent-offspring trios were accepted as valid
when at least three loci were compared (excluded when more than three were
missing), and when these were assigned with a likelihood of 95% ("strict") or 80%
("relaxed"). As all potential parents were known and combinations could be cross-
checked with observed matings, trios with 80% likelihood were included.

Statistical analyses. All statistical analyses were conducted in R version 4.3. (99).
Linear and generalized linear mixed models were conducted using the package *lme4*
(100) with experimental units as a random effect. Where the analysis was done on the
level of matings instead of individuals, the unique male and female ID were added as
random effects. Negative binomial model (*glmmTMB* (101)) or observation level
random effects (102) were used to control for overdispersion in Poisson or binomial
models, respectively. We bootstrapped confidence intervals (n = 1000) for the
opportunity for sexual selection and for the variance partitioning using the package
sampling (103). The morphological measurements for the selection gradients were
scaled within each experimental unit, and relative reproductive success within
experimental units was used as a response variable. We used the package
piecewiseSEM (104) for piecewise structural equation modeling and *diagrammeR*
(105) to visualize the structural equation models. A comprehensive description of all
statistical methods can be found in the *SI Methods.*

Acknowledgments

498 We thank Florin Allmendinger, Sina Lerch, Monica Anderson Berdal, Sonja Sbilordo
499 and Jeannine Roy for their support in the lab, and Scott Pitnick for valuable comments.
500 This project was supported by the Swiss National Science Foundation (grants
501 310030_197651 and PP00P3_202662 to S.L.) and by the University of Zurich (FK-20-
502 088 to A.D.N.).

References

1. M. Andersson, L. W. Simmons, Sexual selection and mate choice. *Trends Ecol. Evol.* **21**, 296–302 (2006).

2. M. Andersson, *Sexual selection* (Princeton University Press, 1994).

3. T. Janicke, I. K. Häderer, M. J. Lajeunesse, N. Anthes, Darwinian sex roles confirmed across the animal kingdom. *Sci. Adv.* **2**, e1500983 (2016).

4. Bateman, A. J., Intra-sexual selection in *Drosophila*. *Heredity* **2**, 349–368 (1948).

5. R. L. Trivers, "Parental investment and sexual selection" in *Sexual Selection and the Descent of Man, 1871-1971*, (Aldine Publishing Company, 1972), pp. 52–95.

6. D. T. Gwynne, L. W. Simmons, Experimental reversal of courtship roles in an insect. *Nature* **346**, 172–174 (1990).

7. S. T. Emlen, L. W. Oring, Ecology, sexual selection, and the evolution of mating systems. *Science* **197**, 215–223 (1977).

8. M. Jirotkul, Operational sex ratio influences female preference and male–male competition in guppies. *Anim. Behav.* **58**, 287–294 (1999).

9. S. Wacker, *et al.*, Operational sex ratio but not density affects sexual selection in fish: Effects of OSR and density on sexual selection. *Evolution* **67**, 1937–1949 (2013).

10. T. Janicke, E. H. Morrow, Operational sex ratio predicts the opportunity and direction of sexual selection across animals. *Ecol. Lett.* **21**, 384–391 (2018).

11. M. L. Head, A. T. Kahn, J. M. Henshaw, J. S. Keogh, M. D. Jennions, Sexual selection on male body size, genital length and heterozygosity: Consistency across habitats and social settings. *J. Anim. Ecol.* **86**, 1458–1468 (2017).

12. M. L. Head, A. K. Lindholm, R. Brooks, Operational sex ratio and density do not affect directional selection on male sexual ornaments and behavior. *Evolution* **62**, 135–144 (2007).

13. A. G. Jones, G. Rosenqvist, A. Berglund, J. C. Avise, The measurement of sexual selection using Bateman's principles: an experimental test in the sex-role-reversed pipefish *Syngnathus typhle*. *Integr. Comp. Biol.* **45**, 874–884 (2005).

14. P. S. Fitze, J.-F. Le Galliard, Operational sex ratio, sexual conflict and the intensity of sexual selection. *Ecol. Lett.* **11**, 432–439 (2008).

15.   I. Klemme, H. Ylönen, J. A. Eccard, Reproductive success of male bank voles
(*Clethrionomys glareolus*): the effect of operational sex ratio and body size.
Behav. Ecol. Sociobiol. **61**, 1911–1918 (2007).

16.   G. Arnqvist, The effects of operational sex ratio on the relative mating success
of extreme male phenotypes in the water strider *Gerris odontogaster* (Zett.)
(Heteroptera; Gerridae). *Anim. Behav.* **43**, 681–683 (1992).

17.   J.-F. Le Galliard, P. S. Fitze, R. Ferrière, J. Clobert, Sex ratio bias, male
aggression, and population collapse in lizards. *Proc. Natl. Acad. Sci.* **102**,
18231–18236 (2005).

18.   H. Klug, J. Heuschele, M. D. Jennions, H. Kokko, The mismeasurement of
sexual selection. *J. Evol. Biol.* **23**, 447–462 (2010).

19.   R. Rios Moura, P. E. C. Peixoto, The effect of operational sex ratio on the
opportunity for sexual selection: a meta-analysis. *Anim. Behav.* **86**, 675–683
(2013).

20.   G. A. Parker, M. A. Ball, P. Stockley, M. J. G. Gage, Sperm competition games:
individual assessment of sperm competition intensity by group spawners. *Proc.*
R. Soc. Lond. B Biol. Sci. **263**, 1291–1297 (1996).

21.   U. Candolin, J. D. Reynolds, Adjustments of ejaculation rates in response to
risk of sperm competition in a fish, the bitterling (*Rhodeus sericeus*). *Proc. R.*
Soc. Lond. B Biol. Sci. **269**, 1549–1553 (2002).

22.   A. Rondeau, B. Sainte-Marie, Variable mate-guarding time and sperm
allocation by male snow crabs (*Chionoecetes opilio*) in response to sexual
competition, and their impact on the mating success of females. *Biol. Bull.* **201**,
204–217 (2001).

23.   L. K. Weir, J. W. A. Grant, J. A. Hutchings, The influence of operational sex
ratio on the intensity of competition for mates. *Am. Nat.* **177**, 167–176 (2011).

24.   F. Garcia-Gonzalez, M. Gomendio, Adjustment of copula duration and
ejaculate size according to the risk of sperm competition in the golden egg bug
(*Phyllomorpha laciniata*). *Behav. Ecol.* **15**, 23–30 (2004).

25.   J. P. Moatt, C. Dytham, M. D. F. Thom, Sperm production responds to
perceived sperm competition risk in male *Drosophila melanogaster*. *Physiol.*
Behav. **131**, 111–114 (2014).

26.   S. A. Ramm, *et al.*, Sperm competition risk drives plasticity in seminal fluid
composition. *BMC Biol.* **13**, 87 (2015).

27.   A. Jarrige, D. Riemann, M. Goubault, T. Schmoll, Strategic sperm allocation in
response to perceived sperm competition risk in a lekking insect. *Anim. Behav.*
109, 81–87 (2015).

28. J. R. Linklater, B. Wertheim, S. Wigby, T. Chapman, Ejaculate depletion patterns evolve in response to experimental manipulation of sex ratio in *Drosophila melanogaster*. *Evolution* **61**, 2027–2034 (2007).

29. Y. Gao, L. Kang, Operational sex ratio and alternative reproductive behaviours in Chinese bushcricket, *Gampsocleis gratiosa*. *Ethology* **112**, 325–331 (2006).

30. K. U. Wensing, M. Koppik, C. Fricke, Precopulatory but not postcopulatory male reproductive traits diverge in response to mating system manipulation in *Drosophila melanogaster*. *Ecol. Evol.* **7**, 10361–10378 (2017).

31. M. Reuter, *et al.*, Adaptation to experimental alterations of the operational sex ratio in populations of *Drosophila melanogaster*. *Evolution* **62**, 401–412 (2008).

32. Y. L. Cai, C. L. Mai, X. Yu, W. B. Liao, Effect of population density on relationship between pre- and postcopulatory sexual traits. *Anim. Biol.* **69**, 281–292 (2019).

33. T. S. Chechi, Z. Ali Syed, N. G. Prasad, Virility does not imply immensity: Testis size, accessory gland size and ejaculate depletion pattern do not evolve in response to experimental manipulation of sex ratio in *Drosophila melanogaster*. *J. Insect Physiol.* **98**, 67–73 (2017).

34. H. S. Crudgington, S. Fellows, N. S. Badcock, R. R. Snook, Experimental manipulation of sexual selection promotes greater male mating capacity but does not alter sperm investment. *Evolution* **63**, 926–938 (2009).

35. K. B. McNamara, *et al.*, Male-biased sex ratio does not promote increased sperm competitiveness in the seed beetle, *Callosobruchus maculatus*. *Sci. Rep.* **6**, 28153 (2016).

36. S. Lüpold, L. Jin, W. B. Liao, Population density and structure drive differential investment in pre- and postmating sexual traits in frogs. *Evolution* **71**, 1686–1699 (2017).

37. H. Kokko, D. J. Rankin, Lonely hearts or sex in the city? Density-dependent effects in mating systems. *Philos. Trans. R. Soc. B Biol. Sci.* **361**, 319–334 (2006).

38. G. Lehmann, Density-dependent plasticity of sequential mate choice in a bushcricket (Orthoptera: Tettigoniidae). *Aust. J. Zool.* **55**, 123–130 (2007).

39. A. Atwell, W. E. Wagner, Female mate choice plasticity is affected by the interaction between male density and female age in a field cricket. *Anim. Behav.* **98**, 177–183 (2014).

40. C. M. House, J. Rapkin, J. Hunt, D. J. Hosken, Operational sex ratio and density predict the potential for sexual selection in the broad-horned beetle. *Anim. Behav.* **152**, 63–69 (2019).

41. L. Winkler, R. Eilhardt, T. Janicke, Population density affects sexual selection in an insect model. *Funct. Ecol.* **37**, 2734–2747 (2023).

42. R. D. Cothran, D. Schmidenberg, A. R. Stiff, G. A. Wellborn, R. A. Relyea,
Effects of density on the strength of sexual selection in the laboratory and in
nature. *Biol. J. Linn. Soc.* **140**, 504–517 (2023).

43. C. C. Taff, C. R. Freeman-Gallant, P. O. Dunn, L. A. Whittingham, Spatial
distribution of nests constrains the strength of sexual selection in a warbler. *J.*
Evol. Biol. **26**, 1392–1405 (2013).

44. M. Jirotkul, Population density influences male–male competition in guppies.
Anim. Behav. **58**, 1169–1175 (1999).

45. D. McLain, Population density and the intensity of sexual selection on body
length in spatially or temporally restricted natural populations of a seed bug.
Behav. Ecol. Sociobiol. **30** (1992).

46. J. Conner, Density-dependent sexual selection in the fungus beetle,
Bolitotherus cornutus. *Evolution* **43**, 1378–1386 (1989).

47. J. I. Morales-Mata, J. Potti, C. Camacho, J. Martínez-Padilla, D. Canal,
Phenotypic selection on an ornamental trait is not modulated by breeding
density in a pied flycatcher population. *J. Evol. Biol.* **35**, 610–620 (2022).

48. N. P. Sharp, A. F. Agrawal, Mating density and the strength of sexual selection
against deleterious alleles in *Drosophila melanogaster*. *Evolution* **62**, 857–867
(2008).

49. E. L. McCullough, B. A. Buzatto, L. W. Simmons, Population density mediates
the interaction between pre- and postmating sexual selection. *Evolution* **72**,
893–905 (2018).

50. R. J. Knell, Population density and the evolution of male aggression. *J. Zool.*
278, 83–90 (2009).

51. M. Reichard, P. Jurajda, C. Smith, Male-male interference competition
decreases spawning rate in the European bitterling (Rhodeus sericeus). *Behav.*
Ecol. Sociobiol. **56**, 34–41 (2004).

52. C. D. Kelly, The interrelationships between resource-holding potential,
resource-value and reproductive success in territorial males: How much
variation can we explain? *Behav. Ecol. Sociobiol.* **62**, 855–871 (2008).

53. E. A. Lacey, J. R. Wieczorek, Territoriality and male reproductive success in
arctic ground squirrels. *Behav. Ecol.* **12**, 626–632 (2001).

54. M. E. Roithmair, Territoriality and male mating success in the dart-poison frog,
Epipedobates femoralis (Dendrobatidae, Anura). *Ethology* **92**, 331–343 (1992).

55. G. N. Dodson, Resource defense mating system in antlered flies, *Phytalmia*
spp. (Diptera: Tephritidae). *Ann. Entomol. Soc. Am.* **90**, 496–504 (1997).

56. G. N. Dodson, "Behavior of the Phytalmiinae and the evolution of antlers in
tephritid flies" in *Fruit Flies (Tephritidae): Phylogeny and Evolution of Behavior*,
(CRC Press, 1999).

57. B. A. Buzzatto, J. L. Tomkins, L. W. Simmons, "Alternative phenotypes within
mating systems" in *The Evolution of Insect Mating Systems*, (Oxford University
Press, 2014).

58. J. L. Brown, The evolution of diversity in avian territorial systems. *Wilson Bull.*
76, 160–169 (1964).

59. J. W. A. Grant, Whether or not to defend? The influence of resource distribution.
Mar. Behav. Physiol. **23**, 137–153 (1993).

60. S. C. Mills, J. D. Reynolds, Operational sex ratio and alternative reproductive
behaviours in the European bitterling, *Rhodeus sericeus. Behav. Ecol.*
Sociobiol. **54**, 98–104 (2003).

61. K. Seidelmann, Territoriality is just an option: allocation of a resource
fundamental to the resource defense polygyny in the European wool carder
bee, *Anthidium manicatum* (Hymenoptera: Megachilidae). *Behav. Ecol.*
Sociobiol. **75**, 138 (2021).

62. G. Raihani, M. A. Serrano-Meneses, A. Córdoba-Aguilar, Male mating tactics
in the American rubyspot damselfly: territoriality, nonterritoriality and switching
behaviour. *Anim. Behav.* **75**, 1851–1860 (2008).

63. P. G. Byrne, J. D. Roberts, Intrasexual selection and group spawning in
quacking frogs (*Crinia georgiana*). *Behav. Ecol.* **15**, 872–882 (2004).

64. M. Reichard, C. Smith, W. C. Jordan, Genetic evidence reveals density-
dependent mediated success of alternative mating behaviours in the European
bitterling (*Rhodeus sericeus*). *Mol. Ecol.* **13**, 1569–1578 (2004).

65. Y. Sato, M. W. Sabelis, M. Egas, Alternative male mating behaviour in the two-
spotted spider mite: dependence on age and density. *Anim. Behav.* **92**, 125–
131 (2014).

66. J. Perdigón Ferreira, S. Lüpold, Condition- and context-dependent alternative
reproductive tactic in *Drosophila prolongata. Behav. Ecol.* **33**, 213–221 (2022).

67. T. Aronsen, A. Berglund, K. B. Mobley, I. I. Ratikainen, G. Rosenqvist, Sex ratio
and density affect sexual selection in a sex-role reversed fish. *Evolution* **67**,
3243–3257 (2013).

68. S. Setoguchi, *et al.*, Sexual dimorphism and courtship behavior in *Drosophila*
prolongata. J. Ethol. **32**, 91–102 (2014).

69. K. Amino, T. Matsuo, Reproductive advantage of the winners of male-male
competition in *Drosophila prolongata. Behav. Processes* **206**, 104831 (2023).

70.   K. Amino, T. Matsuo, Intra- versus inter-sexual selection on sexually dimorphic
692   traits in *Drosophila prolongata. Zoolog. Sci.* **37**, 210 (2020).

693   71.   A. Kudo, H. Takamori, H. Watabe, Y. Ishikawa, T. Matsuo, Variation in
694   morphological and behavioral traits among isofemale strains of *Drosophila*
695   *prolongata* (Diptera: Drosophilidae). *Entomol. Sci.* **18**, 221–229 (2015).

696   72.   K. Minekawa, K. Amino, T. Matsuo, A courtship behavior that makes
697   monandrous females polyandrous. *Evolution* **74**, 2483–2493 (2020).

698   73.   K. Minekawa, T. Miyatake, Y. Ishikawa, T. Matsuo, The adaptive role of a
699   species-specific courtship behaviour in coping with remating suppression of
700   mated females. *Anim. Behav.* **140**, 29–37 (2018).

701   74.   S. J. Arnold, Bateman's principles and the measurement of sexual selection in
702   plants and animals. *Am. Nat.* **144**, S126–S149 (1994).

703   75.   M. J. Wade, Sexual selection and variance in reproductive success. *Am. Nat.*
704   **114**, 742–747 (1979).

705   76.   C. C. Rittschof, Male density affects large-male advantage in the golden silk
706   spider, Nephila clavipes. *Behav. Ecol.* **21**, 979–985 (2010).

707   77.   A. Bertin, F. Cézilly, Density-dependent influence of male characters on mate-
708   locating efficiency and pairing success in the waterlouse Asellus aquaticus: an
709   experimental study. *J. Zool.* **265**, 333–338 (2005).

710   78.   E. L. McCullough, D. J. Emlen, Evaluating the costs of a sexually selected
711   weapon: big horns at a small price. *Anim. Behav.* **86**, 977–985 (2013).

712   79.   M. E. Herberstein, C. J. Painting, G. I. Holwell, "Scramble competition polygyny
713   in terrestrial arthropods" in *Advances in the Study of Behavior*, (Elsevier, 2017),
714   pp. 237–295.

715   80.   F. V. Vencl, A. D. Carlson, Proximate mechanisms of sexual selection in the
716   firefly *Photinus pyralis* (Coleoptera: Lampyridae). *J. Insect Behav.* **11**, 191–207
717   (1998).

718   81.   A. H. Krakauer, M. S. Webster, E. H. Duval, A. G. Jones, S. M. Shuster, The
719   opportunity for sexual selection: not mismeasured, just misunderstood. *J. Evol.*
720   *Biol.* **24**, 2064–2071 (2011).

721   82.   A. Yurkovic, O. Wang, A. C. Basu, E. A. Kravitz, Learning and memory
722   associated with aggression in *Drosophila melanogaster. Proc. Natl. Acad. Sci.*
723   **103**, 17519–17524 (2006).

724   83.   N. Toyoshima, T. Matsuo, Fight outcome influences male mating success in
725   *Drosophila prolongata. J. Ethol.* **41**, 119–127 (2023).

726   84.   S. Setoguchi, A. Kudo, T. Takanashi, Y. Ishikawa, T. Matsuo, Social context-
727   dependent modification of courtship behaviour in *Drosophila prolongata. Proc.*
728   *R. Soc. B Biol. Sci.* **282**, 20151377 (2015).

85. A. N. De Nardo, J. Roy, S. H. Sbilordo, S. Lüpold, Condition-dependent
interaction between mating success and competitive fertilization success in
Drosophila melanogaster. *Evolution* **75**, 2014–2026 (2021).

86. E. L. Macartney, *et al.*, Sperm depletion in relation to developmental nutrition
and genotype in *Drosophila melanogaster*. *Evolution* **75**, 2830–2841 (2021).

87. B. R. Hopkins, *et al.*, Divergent allocation of sperm and the seminal proteome
along a competition gradient in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci.*
116, 17925–17933 (2019).

88. L. K. Sirot, M. F. Wolfner, S. Wigby, Protein-specific manipulation of ejaculate
composition in response to female mating status in *Drosophila melanogaster*.
Proc. Natl. Acad. Sci. **108**, 9922–9926 (2011).

89. B. R. Hopkins, J. C. Perry, The evolution of sex peptide: sexual conflict,
cooperation, and coevolution. *Biol. Rev.* **97**, 1426–1448 (2022).

90. L. K. Sirot, M. F. Wolfner, "Who's zooming who? Seminal fluids and cryptic
female choice in diptera" in *Cryptic Female Choice in Arthropods*, A. V. Peretti,
A. Aisenberg, Eds. (Springer International Publishing, 2015).

91. R. C. Firman, C. Gasparini, M. K. Manier, T. Pizzari, Postmating female control:
20 years of cryptic female choice. *Trends Ecol. Evol.* **32**, 368–382 (2017).

92. M. T. Ghiselin, *The Economy of nature and the evolution of sex* (University of
California press, 1974).

93. C. D. Kelly, Sexual selection on size and shape in Japanese beetles (*Popillia
japonica*). *Behav. Ecol.* **31**, 1073–1083 (2020).

94. J. Moya-Laraño, M. E. T. El-Sayyid, C. W. Fox, Smaller beetles are better
scramble competitors at cooler temperatures. *Biol. Lett.* **3**, 475–478 (2007).

95. P. T. Rohner, *et al.*, Interrelations of global macroecological patterns in wing
and thorax size, sexual size dimorphism, and range size of the Drosophilidae.
Ecography **41**, 1707–1717 (2018).

96. W. U. Blanckenhorn, R. F. Preziosi, D. J. Fairbairn, Time and energy
constraints and the evolution of sexual size dimorphism — to eat or to mate?
Evol. Ecol. **9**, 369–381 (1995).

97. J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis.
Nat. Methods **9**, 676–682 (2012).

98. S. T. Kalinowski, M. L. Taper, T. C. Marshall, Revising how the computer
program CERVUS accommodates genotyping error increases success in
paternity assignment. *Mol. Ecol.* **16**, 1099–1106 (2007).

99. R Core Team, R: A language and environment for statistical computing. (2023).
Deposited 2023.

100. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models
using lme4. *J. Stat. Softw.* **67** (2015).

101. M. Brooks E., *et al.*, glmmTMB balances speed and flexibility among packages
for zero-inflated generalized linear mixed modeling. *R J.* **9**, 378 (2017).

102. X. A. Harrison, A comparison of observation-level random effect and Beta-
Binomial models for modelling overdispersion in Binomial data in ecology &
evolution. *PeerJ* **3**, e1114 (2015).

103. Y. Tillé, Matei, sampling: Survey Sampling. (2023). Deposited 2023.

104. J. S. Lefcheck, piecewiseSEM: Piecewise structural equation modelling in R for
ecology, evolution, and systematics. *Methods Ecol. Evol.* **7**, 573–579 (2016).

105. R. Iannone, DiagrammeR: Graph/Network Visualization. (2023). Deposited
2023.

Figures

Figure 1: Bateman gradients for males (blue) and females (red) for each combination

of density (low, medium, high) and OSR (even, male-biased) and density.

 Figure 2: Linear selection gradients with 95% confidence intervals on male
 morphological traits under different levels of population density and OSRs.

791   Figure 3: The effect of territoriality (time on the resource-dense area; left), and PC1
792   (middle) on mating success depends on the OSR. The effect of PC1 on courtship
793   occurrence also depends on the OSR (right).

Figure 4: Visualized structural equation model of male reproductive dynamics under (a) an even OSR. Green and red indicate positive and negative relationships, respectively. Black lines indicate neutral relationships. Dashed lines (red or green) indicate trends ($P < 0.1$).