High-resolution detection of copy number alterations in single cells with HiScanner

Yifan Zhao[1,2], Lovelace J Luquette[1], Alexander D Veit [1], Xiaochen Wang[3], Ruibin Xi[3], Vinayak V Viswanadham[1], Diane D Shao[4,5], Christopher A Walsh[4], Hong Wei Yang[6], Mark D Johnson[6]*, Peter J Park[1]*

1 Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA
2 Program in Health Sciences & Technology, Harvard Medical School & Massachusetts Institute of Technology, Boston, MA, USA
3 Department of Probability and Statistics, School of Mathematical Sciences, Peking University, Beijing, China
4 Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA
5 Department of Neurology, Boston Children's Hospital, Boston, MA, USA
6 Department of Neurological Surgery, University of Massachusetts Chan Medical School, Worcester, MA 01655

*Correspondence to: mark.johnson3@umassmemorial.org, peter_park@hms.harvard.edu

Abstract

Improvements in single-cell whole-genome sequencing (scWGS) assays have enabled detailed characterization of somatic copy number alterations (CNAs) at the single-cell level. Yet, current computational methods are mostly designed for detecting chromosome-scale changes in cancer samples with low sequencing coverage. Here, we introduce HiScanner (High-resolution Single-Cell Allelic copy Number callER), which combines read depth, B-allele frequency, and haplotype phasing to identify CNAs with high resolution. In simulated data, HiScanner consistently outperforms state-of-the-art methods across various CNA types and sizes. When applied to high-coverage scWGS data from human brain cells, HiScanner shows a superior ability to detect smaller CNAs, uncovering distinct CNA patterns between neurons and oligodendrocytes. For 179 cells we sequenced from longitudinal meningioma samples, integration of CNAs with point mutations revealed evolutionary trajectories of tumor cells. These findings show that HiScanner enables accurate characterization of frequency, clonality, and distribution of CNAs at the single-cell level in both non-neoplastic and neoplastic cells.

Introduction

Over the past decade, advances in single-cell omics technologies have provided an opportunity to interrogate cellular heterogeneity with unprecedented resolution. Single-cell RNA-sequencing has been the most popular[1,2], but significant progress has also been made for single-cell ATAC-sequencing for profiling open chromatin[3] and single-cell Hi-C for profiling three-dimensional folding[4,5]. Another technology that has advanced is single-cell whole-genome sequencing (scWGS), which can be used to uncover cell-to-cell variation in somatic mutations that may be masked in bulk sequencing.

Various scWGS approaches have been developed for detecting copy number alterations (CNAs), primarily in cancer cells. Examples include degenerate-oligonucleotide-primed PCR (DOP-PCR)[6], direct library preparation (DLP+)[7,8], and the now-defunct 10x Genomics chromium CNV assay[9]. However, the fraction of genome covered by these approaches is low (low breadth-of-coverage) and their main application was to detect clonal patterns of chromosomal arm-sized CNAs. Single-cell whole-genome amplification (scWGA) offers high breadth-of-coverage and could improve spatial resolution, but it has been hindered by the noise introduced during amplification. Earlier scWGA methods have involved exponential or quasi-linear amplification, generating highly uneven coverage across the genome, allelic bias (in which one allele is amplified at a higher rate), and frequent allelic dropout (where only one allele is amplified). Even if the cell is sequenced at high depth, these amplification artifacts can confound CNA detection by causing large fluctuations in the read depth ratio (RDR; ratio between observed and expected read count), yielding relatively few advantages over low-coverage (<1X, often <0.1X) scWGS.

Recent advances in scWGA chemistry[10,11] and declining sequencing costs have made high-coverage (>20X) scWGS more attractive for CNA detection. In particular, scWGA by primary template-directed amplification (PTA) achieves more uniform amplification across the genome[10,12]. We have already used the high-coverage PTA data to identify somatic single nucleotide variants in the brain, tracing cellular lineage across different tissues[13], providing a more precise estimate of age-associated mutational rate[12], and characterizing the regions more prone to mutagenesis in different cells[14]. PTA data also offer a potential to improve the resolution of CNA detection beyond the usual Mb-scale[15,16]. However, current single-cell CNA callers are not designed to leverage high-coverage data and are tailored to the analysis of clonal CNAs in cancer cells.

Here, we propose HiScanner (High-resolution Single-Cell Allelic copy Number callER) for accurate, high-resolution allele-specific CNA detection in scWGS. It leverages reference-based phasing and local sequence bias to efficiently estimate B-allele frequency (BAF, the relative proportions of reads covering the two alleles at heterozygous germline single nucleotide polymorphisms, gHETs), which provides essential information to complement read depth-based CNA identification. Building upon the framework of a bulk CNA caller we previously developed[17], HiScanner performs read depth normalization at nucleotide resolution, followed by joint segmentation of single-cell profiles using the Bayesian information criterion (BIC). By simultaneously modeling

86   RDR and BAF for individual cells, HiScanner is able to infer cell ploidy and integer
87   allele-specific copy number states. It also considers assay-specific allelic dropout patterns
88   to find an optimal bin size, a nuanced yet crucial parameter in CNA analysis.
89
The use of BAF is a standard feature in best bulk CNA methods such as ASCAT[18] and
Battenberg[19] for inferring allele-specific copy number state. Two recently developed
single-cell callers, CHISEL[20] and Alleloscope[21], leverage phasing and BAF for
identifying allele-specific CNAs in low-coverage scWGS. However, Alleloscope
performs segmentation on bulk samples, not single cells, limiting its ability to detect rare
CNAs which are critical to understanding the genomic architecture of non-cancerous cells
and especially non-dividing cells. CHISEL is not tuned for the discovery of small CNAs
(e.g., <5 Mb) potentially obscuring certain biological processes. SCOVAL[22] also utilizes
BAF information, but it is primarily a validation approach and not a CNA caller.
HiScanner offers robust detection of rare CNA events across a broad spectrum of sizes,
including sub-megabase events, thereby extending the accessible landscape of CNAs in
single-cell genomes. We show that HiScanner outperforms two representative single-cell
CNA methods, CHISEL (BAF-aware) and Ginkgo (BAF-agnostic)[20,23], in real and
simulated data with greater sensitivity and precision at small size scales (200kb-5Mb).
104
For biological discovery, we analyzed two datasets. Application of HiScanner to a high-
coverage scWGS dataset from eleven neurotypical human brains uncovered a large
number of previously undetected small CNAs, revealing contrasting patterns between
neuronal and glial cells. We also sequenced 179 cells at low coverage (~0.5X on average)
from biopsies of an initial (n=88) and recurrent (n=91) meningioma tumor pair.
HiScanner revealed phases and timing of tumor progression by detecting subclone-
defining CNAs, which were subsequently corroborated by an orthogonal clustering based
on somatic point mutations[24]. Lastly, we built a web-based tool, ViScanner, to facilitate
visualization of HiScanner output. Built using the HiGlass architecture[25], ViScanner
offers users the ability to navigate genomic data with continuous zooming, allowing for
user-friendly exploration of genomic regions at various resolutions.
116
117

Results

Overview of the algorithm

HiScanner introduces three key improvements over existing single-cell CNA detection
methodologies to enhance the resolution of CNA discovery (Figure 1). First, optimal
resolution is inferred by determining the smallest permissible bin size based on the size of
amplification artifacts, rather than leaving the tuning of this critical parameter to the user.
Second, joint segmentation is performed across cells to avoid overfitting amplification
noise. Third, an efficient BAF estimator is implemented to mitigate the effects of phase-
switch errors and allelic dropout. In addition to these features, HiScanner corrects for bias
in read depth due to local GC content using a nucleotide-level model of GC bias[17]. These
improvements are described in more detail below.
130

Selecting optimal bin size based on allelic dropout events Variations in RDR across the
genome is the primary signal used to detect copy number changes, but scWGA introduces
considerable levels of uneven amplification along the genome that masquerade as CNAs.
A prevalent form of uneven amplification is allelic dropout, where failed amplification of
one allele results in an RDR decrease similar to a copy number loss (Fig S1a). Previous
single-cell CNA callers partitioned the genome into large bins to dampen the effect of
segments lost to amplification, but this limits the size of CNAs that can be detected. On
the other hand, decreasing the bin size unmasks RDR fluctuations, possibly inducing
false positive CNA calls. A key insight exploited by HiScanner is that the size
distribution of allelic dropout events, which is usually bounded above and is specific to
assay-dependent amplicon length, can be used to determine a balanced bin size that
maximizes detection resolution and minimizes the effects of amplification noise (Fig
S1b). Furthermore, because dropout sizes and frequencies can differ considerably
between scWGA protocols[26] (Fig S1c), it is important to fit the dropout size distribution
on each dataset. For example, the average size of dropout events in two cells amplified by
PTA and MDA[27] differed by two orders of magnitude (10^3 bp versus 10^5 bp, respectively,
Fig S1d). HiScanner employs a two-state hidden Markov model (HMM) to detect dropout
events using BAF at gHETs (Fig S1b). Subsequently, the predicted dropout events are
used to build a sample-specific dropout size distribution and the top 5^{th} percentile of
dropout sizes is used as the smallest permissible bin size.
151
Joint segmentation across multiple cells By leveraging signals from multiple cells,
HiScanner's joint segmentation promotes breakpoint sharing, thus improving detection
sensitivity. We extended our previous strategy[17] of merging adjacent bins with similar
copy number state (as determined by the BIC) to the multi-sample case by computing a
BIC from the joint breakpoint likelihood of a breakpoint across cells (Methods). This
merging approach retains information about bin position, avoids overfitting by penalizing
excessive segmentations, and amplifies signals of clonal CNAs. Notably, the clustering
approach proposed in CHISEL to simultaneously model BAF and RDR across all cells
does not consider the proximity of bins, making the method more susceptible to calling
false-positive breakpoints, especially in regions with complex structural variations or
relatively high amplification noise.
163
Accurate estimation of the B-allele frequency (BAF) BAF can provide additional
information for CNA inference where RDR is ambiguous; a shift in the BAF from the
expected value of 0.5 in diploid regions suggests an underlying CNA (for example, a
single copy gain shifts the BAF to 1/3 or 2/3, depending on which allele the BAF is
measured against). In bulk data, obtaining allele-specific read depth at gHET loci is an
effective way to compute BAF; however, in scWGS, uneven amplification between
homologous alleles (allelic bias) and dropout can hinder this approach. Furthermore, low-
coverage sequencing–which is more typical for single-cell CNA studies–reduces the
number of reads at gHETs, thereby increasing BAF noise with its undersampling. The
effects of both allelic bias and gHET undersampling on BAF estimation can be mitigated
by summing allele-specific read depth over bins. However, this requires phasing each
gHET to a haplotype, and phase switch errors, which can occur at rates of 1-2% [28,29] for
reference-based phasing approaches, confound bin-level BAF estimation by assigning

depth to the incorrect allele. CHISEL proposed a two-step process to calculate per-bin
BAF estimates: after an initial phasing based on an external reference panel,[30,31]
expectation-maximization (EM) is used for refinement, first aggregating phased gHETs
into haplotype blocks and then performing EM refinement of phase across cells.
Alleloscope also adopted an EM approach, but it operates on unphased, individual gHETs.
Although effective, the EM step used by both CHISEL and Alleloscope is
computationally expensive and is challenging to apply to high-resolution bins. Indeed,
CHISEL operates on ~5 Mb bins (by default) and Alleloscope only computes BAF on the
final copy number segments derived from bulk or pseudobulk.
186
HiScanner computes BAF by phasing gHETs using a reference-based method followed
by a computationally efficient, two-step procedure to reduce the effect of phase-switch
errors and dropout artifacts (Fig S2; Supplementary Note S1). In the first step, LOH is
detected by exploiting GC bias, a source of non-biological fluctuations in total read depth
that is treated solely as a nuisance. In a segment without LOH, GC-induced read depth
changes—which are roughly evenly split between homologous alleles due to similar
nucleotide content—lead to a subtle but detectable difference in the distributions of BAF
before and after a haplotype-specific *shift* operation (Supplementary Note S1). In regions
affected by LOH, this shift procedure has little effect. Thus, HiScanner predicts the
presence of LOH in each segment by a Kolmogorov-Smirnov test of the BAF and shifted
BAF distributions. In the second step, peaks (local maximums) are called in the per-bin
BAF distribution of each segment. If the segment was not predicted to contain LOH, then
any BAF peak near 0 or 1 is interpreted as dropout artifacts and removed, and the
maximum of the remaining non-zero peak is the final segment BAF. Otherwise, the BAF
is set to 0. The efficiency of this procedure is critical for CNA analysis with high
resolution bins.
203
Visualization using ViScanner To enhance the accessibility of HiScanner output, we
developed a web-based visualization platform, ViScanner (Methods). Traditional single-
cell CNA analysis tools often generate static output, including mapping coordinates for
potential CNAs and genome-wide visualizations of CNAs. However, genome-scale plots
do not adequately support the exploration of small CNAs. We developed an interactive
React-based web application, ViScanner, which leverages the HiGlass toolkit[25] to provide
a user-friendly, dynamic resolution CNA browser. ViScanner allows users to scrutinize
HiScanner output with real-time zooming, akin to the user experience in web-based
services like Google Maps. Importantly, ViScanner ensures data security by executing
JavaScript locally in the user's browser, avoiding the need to upload data, even
temporarily, to any external server.
215
Comparison with other methods
To assess HiScanner's performance, we conducted benchmarking studies using both real
and simulated data. HiScanner was compared against two widely-used single-cell CNA
methods, Ginkgo[23] and CHISEL[20]. Ginkgo is a popular web-based application that
identifies CNAs based on RDR and performs segmentation via circular binary
segmentation[32]; CHISEL is a more sophisticated algorithm that combines phasing, BAF,
and RDR to identify allele-specific CNAs. Our benchmarking analysis focused on the

detection of non-clonal CNAs (private to a single cell), which are more difficult to
identify than clonal CNAs (present in a subpopulation). Consequently, methods designed
to detect only clonal events, such as Alleloscope[21], were excluded to avoid
mischaracterizing their performance (Table 1).
227
We first evaluated the three callers on detecting non-clonal CNAs using high-coverage
simulated data. Previous benchmarking studies often simulated single-cell CNAs by
manipulating read counts within pre-determined CNA locations, thereby focusing
exclusively on RDR-based CNA callers[33,34]. However, allele-specific callers cannot be
properly assessed by this strategy because simulation at the level of read counts does not
preserve the haplotype structure. Moreover, although simulating read counts using
Lorenz curve sampling[33] can recapitulate read count variability associated with GC
content, it does not faithfully capture sampling and amplification noise characteristic of
scWGS sequencing data. To overcome these limitations, we adapted the synthetic diploid
X chromosome methodology[11] to simulate somatic CNAs. In this approach, sequencing
reads sampled from two male haploid X chromosomes from different individuals are
pooled to generate a diploid chromosome (Fig 2a). By sampling reads prior to pooling, it
is possible to insert copy gain or loss events specific to a single allele without disrupting
local haplotype structure. Further, one can explicitly control phasing accuracy and phase
switch rates, while preserving scWGS amplification artifacts.
243
We simulated three types of non-clonal CNAs with size varying from 200 kb to 5 Mb
using the synthetic diploid X: single-copy losses, single-copy gains, and copy-neutral
loss-of-heterozygosity (cnLOHs). We find that all tools performed well on larger CNAs,
but their performance was variable for smaller CNAs. In the loss detection task,
HiScanner and Ginkgo generally achieved higher precision and sensitivity than CHISEL
across the range of CNA sizes we tested (Fig 2b). However, for certain CNA sizes (e.g.,
small <500 kb deletions), the difference in performance was considerable, with
HiScanner (0.77) achieving slightly greater precision than Ginkgo (0.74) but an almost
20-fold increase over CHISEL (0.04) while maintaining higher sensitivity (Fig 2c).
Figure 2d illustrates a representative ~600 kb single copy loss. In this example, although
the simulated event was called by all three methods, Ginkgo incorrectly called a deletion
nearby and CHISEL incorrectly called three false-positive gains nearby, in regions with
higher RDR noise. In the gain detection task, HiScanner and CHISEL performed better
than Ginkgo overall, with HiScanner outperforming CHISEL in sensitivity for
events >250 kb (Fig 2e, f). A common source of false positives for CHISEL was
overfitting to RDR noise (Fig 2g), similar to what was observed in the loss task. The final
CNA event type, cnLOH, cannot be detected without the use of BAF information and
thus Ginkgo was not assessed in this task. Only HiScanner detected cnLOH events <1
Mb in size and, across the size spectrum, maintained higher precision than CHISEL.
CHISEL, however, slightly outperformed HiScanner in detection sensitivity for cnLOH
events >2 Mb (Fig 2d, g). CHISEL's increased sensitivity for large cnLOH events might
be attributed to its joint clustering approach which utilizes both RDR and BAF, whereas
HiScanner's multi-sample segmentation is based only on RDR. Overall, HiScanner
consistently outperformed Ginkgo and CHISEL in precision and sensitivity across most

CNA sizes and types, especially for single-copy losses and gains smaller than 1 Mb. It
was also the only tool capable of detecting cnLOH events less than 1 Mb.
270
271   As an illustrative example using real data, we next applied the three methods to detect
272   CNAs in a single human neuron amplified by PTA and sequenced to 29X (Fig 3a-e). This
273   neuron displays distinct signals corresponding to loss events in chromosomes 2, 5, 6, 8,
274   11, 12, and 17, encompassing a spectrum of sizes from 300 kb to 170 Mb (whole-
275   chromosome) with some potentially biallelic loss events.
276
Although the CNA calls by Ginkgo and HiScanner were generally similar, we observed
two discrepancies. First, Ginkgo tended to report focal amplifications with high copy
number near centromeres which contain highly repetitive sequences, suggesting
inadequate mappability correction of read depth or bin masking during preprocessing. In
contrast, HiScanner showed smoother read depth signals across the genome, likely due to
its use of only mappable positions and the nucleotide-level normalization strategy (Fig
3h). Second, Gingko tended to call small copy number loss events which clearly lacked
orthogonal BAF support (Fig 3f-g, j-k). CHISEL's calls differed markedly: it produced a
notably high number of breakpoints (10,322 or 0.38 breakpoints per 100 kb; Fig 3i) and
an overall tetraploid state (average CN=4, Fig 3d). The segments CHISEL identified as
cnLOH (CN=2, BAF at 0 or 1) seemed implausible without postulating a series of copy
number losses preceding a whole genome duplication event. After manual parameter
tuning (Methods), CHISEL demonstrated improved performance, though it still had
difficulties in distinguishing biallelic losses from single-copy losses and in detecting
smaller CNAs (Fig 3e).
292
In summary, our analysis of both simulated data and an aneuploid single neuron revealed
distinct characteristics of each caller. Ginkgo displayed high sensitivity to changes in
RDR but, without BAF information, lacked precision in detecting small CNAs and was
not capable of cnLOH detection. CHISEL tended to overfit when used with small initial
bin sizes and, in a single-sample context, produced an improbably large number of
breakpoints. Given that CHISEL is optimized for megabase-scale clonal CNA detection
(e.g., in cancer samples), this behavior is expected. Overall, HiScanner maintained a
balance between sensitivity and precision in detecting CNAs at high resolution.
301

Cell-type-specific CNA patterns in neurotypical human brains

Previous studies using scWGS to investigate non-diseased human brains have reported
contradictory findings regarding the frequency of CNAs in neurons. A meta-analysis of
multiple low-coverage scWGS reported a linear decline of CNA neuron frequency with
age (R^2=0.92, range 0-40.0%) in brains aged 0.36 y.o. to 95 y.o.[15] Knouse et al. reported
a significantly lower frequency (2.2%; 95% CI 0.3%–7.9%) for large CNAs in human
brain cells[35]. More recently, Kalef-Ezra et al. applied PTA to human brain cells and found
33.8% to have at least one megabase-scale CNA[36]. These discrepancies highlight the
need for further research into the prevalence of CNAs in the human brain.
311
Using HiScanner, we analyzed previously published high-coverage PTA scWGS data
encompassing 65 single neurons and oligodendrocytes from eleven neurotypical human

brains spanning an age range from 0.4 to 83 years[12,14] (Fig 4a; Table S1). HiScanner was
applied separately to each brain, as clonal CNAs should be specific to each individual.
Large-scale aneuploidy, characterized by multiple chromosome-arm level losses, was
clearly observed in two cells out of 65 (3.1%). These numbers are concordant with
Knouse et al[35] and substantially less than the others.
319
Beyond these instances, HiScanner revealed an additional 614 smaller, mostly private
CNAs (314 private losses and 57 private gains) in the rest of the cells. cnLOH calls were
excluded from this analysis because of the relatively poor performance in simulation
across both callers tested and lack of RDR support. Whereas previous approaches were
limited to a lower size bound of ~3 Mb on somatic CNA detection in single cells[15,16,34],
nearly all (609/614, 99.3%) of the CNAs identified by HiScanner were smaller than 3 Mb
(Fig 4b-c), with a median size of 612 Kb (range 300Kb-24.1Mb). Using the size-specific
sensitivities and specificities from our simulated data, we estimated a false discovery rate
of 34.5% among the 614 calls (Methods; Table S2). In total, 77% (24/31) of neurons and
53% (17/32) of oligodendrocytes contained at least one CNA. We did not observe a
significant correlation with age for the number of private gains or losses in neurons
(Methods; Figure S3).
332
Our analysis also uncovered a significant association between cell type and CNA type.
Neurons contained both gains and losses, with losses being more prevalent, whereas
nearly all CNAs in oligodendrocytes were gains (Fig 4b-c). In comparison to
oligodendrocytes, the odds ratio of observing losses compared to gains in neurons is 61
(p-value: 1e-23, Fisher's Exact Test). Classification into clonal vs non-clonal categories
revealed broad distinctions: only 5% of the losses in neurons are clonal, whereas 78% of
neuronal gains are clonal. Since neurons are typically non-proliferative, this observation
may suggest differential timings of gain and loss events: gains, which were mostly
shared, must have occurred early in development prior to terminal differentiation into a
post-mitotic neuron, whereas losses, which were mostly private, must have occurred after
development in the post-mitotic state. In oligodendrocytes, the majority of CNAs are
clonal gains (81% of gains being clonal and no losses), similar to the high clonal fractions
in neuronal gains. The differences between neuronal and oligodendrocyte CNAs suggest
differential timing and/or mutagenic processes.
347
Analysis with Ginkgo yielded a fundamentally different view of neuronal and
oligodendrocyte CNAs, even though a similar range of CNAs were detected (Fig 4d-e).
Notably, Ginkgo called nearly a 10-fold more private neuronal losses than HiScanner and
a large number of private losses in oligodendrocytes whereas HiScanner detected almost
none. We assessed the quality of these calls by inspection of the BAF distribution in the
putative CNAs and found that a considerable proportion of Ginkgo's calls disagreed with
local BAF signals (Fig 4f-j): 73% (330/451) of CNAs called in oligodendrocytes and
61% (1110/1847) of CNAs called in neurons were discordant. This suggests that
Ginkgo's calls in PTA scWGS data should be filtered based on CNA size or be subjected
to BAF-based post-call filtering strategies[22] to ensure the reliability of small CNA calls.
In contrast, HiScanner's integration of BAF and RDR signals was effective in preventing
these types of errors.

Integrative CNA and SNV analysis in longitudinal meningioma scWGS

To demonstrate the applicability of HiScanner to low-coverage scWGS, we sequenced PTA-amplified single cells from meningiomas, the most common primary intracranial tumor, to an average depth of 0.5X (Table S3). High-grade (grade II-III) meningiomas are notorious for their high recurrence rate despite treatment, and the effectiveness of postoperative radiotherapy in managing these recurrences is still a subject of debate[37,38]. Characterizing the timing of oncogenic events and their evolution in meningiomas may therefore prove helpful in assessing the impact of therapeutic interventions and guiding clinical management strategies. We obtained tumor tissue samples from initial and recurrent (grade II/III) meningiomas from a single patient. 88 and 91 single tumor cells were sorted from the grade II and grade III specimens, respectively, totaling 179 cells (Fig 5a, Methods). All cells were amplified by PTA, sequenced to 0.5X, and analyzed for CNAs by HiScanner.

Hierarchical clustering on the copy number profiles from both initial (grade II) and recurrent (grade III) tumors revealed four main clusters (Fig 5b-d). Cluster 13, which is a mixture of grade II/III cells, consists of cells that lack any large CNAs. Cluster 11 is entirely composed of grade II cells and exhibits hallmarks of high-grade meningiomas such as single-copy losses in chromosomal regions 1p and 22q (Fig 5e). Cluster 15 is dominated by grade III cells which inherited all CNAs present in Cluster 11 along with additional loss events on chromosomes 6q, 9p, and 11q (Fig 5f). Interestingly, one grade II cell was assigned to Cluster 15, suggesting that it may be closely related to the origin of a major grade III clone. If true, this would imply that the additional copy number events found in grade III tumor cells likely occurred in the natural course of tumor evolution rather than as a consequence of adjuvant therapy. Lastly, Cluster 8 represents a small, distinct group of tetraploid cells from the grade III tumor (Fig 5h), exhibiting a range of copy number states from two to six, including odd copy numbers. Tetraploidy hints at a potential tumor progression pattern involving whole-genome duplication, which is rarely reported in the meningioma literature; however, further experiments (e.g., fluorescence in situ hybridization) would be essential to validate this ploidy estimate. The heterogeneity of these clusters reflects the complex nature of meningioma evolution and underscores the importance of single-cell profiling in delineating tumor progression.

Next, we investigated whether cluster assignments derived from CNAs were consistent with patterns of somatic SNVs (sSNVs). To this end, we employed a recently developed somatic SNV-based cell clustering method, SECEDO[24]. Initially, applying SECEDO with its default settings failed to differentiate between aneuploid and diploid cells, which are a hallmark of meningiomas. However, upon refining the input by limiting to sSNV locations identified in matched bulk WGS via GATK MuTect2[39] (Methods), SECEDO yielded reasonable clustering results. SECEDO predicted three primary cell clusters which closely corresponded to HiScanner's CNA-based clustering (Fig 5i, j): SNV Cluster 4 aligned largely with CNA Cluster 11; sSNV Cluster 2 comprised a mix of cells from CNA Clusters 8 and 13; and sSNV Cluster 4 was found to be a mix of CNA Clusters 8 and 15. Notably, however, SECEDO's clustering did not distinguish diploid and tetraploid grade III cells. A possible explanation for this is that the whole-genome

duplication preceded the majority of sSNV events. SECEDO also assigned three grade II
tumor cells to its grade III major cluster, suggesting that some sSNVs preceded CNAs in
the evolution of this tumor. To quantitatively compare the sSNV and CNA cluster
assignments, we computed the Adjusted Rand Index (ARI), a score ranging from 0
(random labeling) to 1 (perfect match). The two orthogonal clusterings produced an ARI
of 0.83, indicating a high level of agreement and again supporting the validity of
HiScanner's CNA calls.
413
414

Discussion

We developed a new single-cell CNA detection algorithm, benchmarked it on real and
synthetic data, and applied it to high-depth scWGS of non-diseased human brains and
low-depth scWGS of longitudinal meningioma tumor samples. In our benchmarking,
HiScanner achieved similar performance to a BAF-aware and BAF-agnostic single-cell
CNA caller for detecting large (3-5 Mb) CNAs but outperformed them in the task of
detecting small CNAs (<1 Mb) and cnLOH events. Small and cnLOH events are
characteristic of many mutagenic processes and detecting them accurately is essential.[40]
423
In the human brains, HiScanner detected many CNAs events smaller than the broadly
assumed ~3 Mb detection limit. Although the small sample size provided insufficient
power to ascertain an association between CNA burden and age, HiScanner revealed
previously undetected differences[12] in CNA patterns between cell types. Neuronal CNAs
were predominantly private losses, in line with previous studies in human neurons[41],
whereas oligodendrocytes mostly exhibited clonal gains. This finding suggests that
neurons likely acquired these losses post-mitotically via non-replicative mechanisms such
as nonhomologous end joining and microhomology-mediated end joining which can lead
to deletions or complex genomic rearrangements without the need for DNA synthesis[42].
Indeed, neurons frequently undergo double-stranded breaks and repair due to their
synaptic activity[43]. In contrast, the clonal gains found in both neurons and
oligodendrocytes may have occurred earlier in brain development through replicative
mechanisms such as microhomology-mediated break-induced replication and fork
stalling and template switching, which can induce duplications or deletions during DNA
replication[42]. While the clonal gains in post-mitotic neurons are most likely to have
occurred during development, the clonal gains in oligodendrocytes may have occurred
later in life in ancestral oligodendrocyte precursor cells, which remain proliferative.
Further investigation into the timing and mutational mechanisms of these CNAs with
more cells is warranted.
443
Our integrative analysis of sSNVs and CNAs provided a more detailed view of the clonal
origin and evolution of a meningioma tumor, highlighting the complex and dynamic
nature of cancer progression and the importance of single-cell profiling. This approach
identified an ancestor of the grade III major clone in the grade II tumor, suggesting that
the mutagenesis was not caused by radiotherapy. However, it remains possible that the
clonal expansion of the grade III clone was influenced by selective pressure induced by
radiotherapy. Timing the progression between tumor grades, and whether this occurs in
response to or because of therapy, is critical information for clinical decision making.

452
Several studies of somatic CNAs have delineated the evolutionary trajectories of
subpopulations of tumor cells, most notably in breast cancer where chromosome-scale
aneuploidy is common. A more comprehensive set of CNAs including smaller CNVs will
allow similar analysis for other tumor types in which the large CNAs are less common.
The timing of SNV driver mutations with respect to the appearance of genome-doubling
or large CNAs can be inferred based on careful analysis of variant allele fractions in bulk
data[44]. With the growing availability of high-coverage scWGS and the accompany
development of methods for detecting SNVs, CNAs, and other variants, a timing analysis
at the single cell resolution will be possible, deepening our understanding of genomic
stability and evolution.
463
464
465
466

Methods

Preprocessing of BAF

The GATK HaplotypeCaller "Best Practices" pipeline[39] with default parameters was used to call SNPs on bulk sequencing data from each individual. Germline heterozygous SNPs (gHETs) were defined as the variants passing the FILTER column flag, marked with a heterozygous '0/1' genotype, annotated non-zero population frequencies in the 1000 Genomes Project database[45] and listed in the Single Nucleotide Polymorphism Database[46]. SHAPEIT2[47] was applied to phase the gHETs using the 1000 Genomes Phase 3 integrated haplotype panel[48]. Only the gHETs that can be phased by SHAPEIT2 and have a variant allele frequency between 0.1 and 0.9 in bulk data were retained for subsequent analysis. BCFtools[49] was used to query the allelic-specific depth at each gHET site within each single cell BAM. Next, reads associated with the same haplotypes were aggregated to obtain bin-wise haplotype-specific read counts. The phased B-allele frequency (pBAF) was computed as the fraction of haplotype-B read count divided by the total read count within a given bin. Note that the pBAF here corresponds to individual bins and can be heavily influenced by local allelic imbalance. To ensure robust allelic copy number inference, we implemented additional per-segment refinement of pBAF after the segmentation step using the LOH test (Supplementary Notes).

Preprocessing of RDR

BICseq2-norm[17] with default parameters was used to perform GC correction and binning on single-cell and bulk sequencing data from each individual. Only uniquely mappable reads with a minimum mapping quality of 30 were considered and only mappable genomic loci, as determined by the 100mer mappability track from https://www.math.pku.edu.cn/teachers/xirb/downloads/software/BICseq2/BICseq2.html, were considered.

Allelic dropout analysis

Loss or under-amplification of one allele during whole-genome amplification, known as allelic dropout, is a well-documented phenomenon in single-cell sequencing assays[26]. In the sequencing data, dropout regions manifest with reduced RDRs and imbalanced allelic ratios, closely resembling regions with copy number losses. The characteristics and prevalence of dropout can be affected by several factors, including amplicon length and sequencing depths. Consequently, different single-cell assays may exhibit varying patterns of dropout, and these nuances should be accounted for when determining the optimal bin size for CNA analysis in single cells.

To determine the distribution of dropout sizes without relying on arbitrary thresholds, we designed a hidden Markov model (HMM) with two distinct states—"Allelic dropout (ADO)" and "non-ADO"—and used Gaussian distribution probabilities to describe them (Fig. S1). This model assumes that CNAs are infrequent compared to allelic imbalance by dropout in the selected single-cell genomes (in this study we used oligodendrocytes). We used a binary representation of BAF data at gHETs on the autosomes. A value of |BAF-0.5| less than 0.2 was marked as "0" to indicate no sign of a homozygous state, while a

value of 0.2 or greater was marked as "1" to indicate a homozygous state. Only gHET
sites with >5 reads were considered. To derive optimal bin sizes for PTA- and MDA-
amplified single cells in this study, we fit two HMMs using the *hmmlearn* library in
Python on two oligodendrocytes—1278-Oligo-5 for PTA and 1278_ct_p1G9 for MDA—
which exhibited minimal aneuploidy, as determined through manual inspection of their
read coverage profiles. After fitting, the HMMs were applied to predict dropout events
across the autosomes of additional oligodendrocytes from the same brain UMB1278
(1278-Oligo-{7,8} for PTA and 1278_ct_{p1E3, p1E6, p2B9, p2E4, p2F5, p2G5} for
MDA) derive dropout length distributions. The 95th percentile of each dropout length
distribution (one for PTA, one for MDA) was selected as the minimum bin size.
521
522 Joint segmentation algorithm

523 We extended the single-sample segmentation algorithm implemented by BICseq2-seg[17]
524 to the multi-sample case. Within the original BIC-seq framework, the simplified Bayesian
525 Information Criterion (BIC) is defined as:

526
$$BIC(\lambda) = -2\sum_{j=0}^{m}[a_j(\log(\frac{a_j}{a_j + b_j}) + b_j\log(\frac{b_j}{a_j + b_j}))] + (m + 1)\lambda\log(N)$$
,

where for every breakpoint j, a_j and b_j are the number of observed reads and the number
of expected reads between breakpoints β_j and β_{j+1}; $\lambda > 0$ is the tuning parameter; m is the
number of breakpoints; and $N = \Sigma_{j=0}^{m} a_j$ is the total number of reads in the given cell.
Recognizing that the joint log-likelihood of the short reads from G single-cell samples is
just the sum of the individual log-likelihoods, and assuming independence in cells'
segmentation patterns, the multi-sample form of the simplified BIC becomes:

533
$$BIC(\lambda) = -2\sum_{k=1}^{G}\sum_{j=0}^{m}[a_{kj}(\log(\frac{a_{kj}}{a_{kj} + b_{kj}}) + b_{kj}\log(\frac{b_{kj}}{b_{kj}}))] + (m + 1)G\lambda\log(\sum_{k=1}^{G} N_k)$$
.

Given an initial set of bins, HiScanner attempts to reduce the overall BIC by merging the
neighboring bins. HiScanner calculates the BIC for each possible merge of adjacent bins.
If the calculated BIC for a merged pair is lower than it would be if they remained
separate, HiScanner considers these bins for merging. Out of all possible merges,
HiScanner selects the pair of bins that, if merged, would result in the greatest reduction in
the BIC. The selected bins are then merged. HiScanner repeats this process, continuing to
merge bins pair by pair, until no more merges would result in a lower BIC.
541
542 Ploidy and copy number inference

We extended the likelihood-based method introduced in CHISEL[20] to infer ploidy and
allelic copy number states. The procedure has two main steps: identification of all
possible values of the scale factor γ, and finding the maximum likelihood estimate of
allelic copy number and γ based on BIC. The scale factor γ is a cell-specific constant
value and describes the linear relationship between the total copy number and RDR for
each bin. It follows that knowing the true copy number and RDR of one bin is sufficient
to compute the corresponding γ. We assume that within any cell, there exists a substantial
proportion of bins unaffected by CNAs, retaining balanced allelic copy numbers. The
allelic copy number of these bins takes the form {1,1} when there is no whole genome
duplication (WGD), {2,2} with one WGD, or {3,3} with two WGDs, etc. CHISEL's
global clustering approach allows for directly selecting the largest cluster with BAF≈ 0.5

as the "balanced" bin set, even when these bins are not contiguous. In contrast,
HiScanner's bottom-up segmentation restricts bin merging to spatially adjacent bins with
similar RDR and BAF values. In HiScanner, the "balanced" bins are identified by
requiring $0.45 < BAF < 0.55$. HiScanner employs a Gaussian Mixture Model (GMM) to
further segregate these "balanced" bins into clusters with similar RDR values. To
determine the best GMM, HiScanner conducts a parameter search for the number of
Gaussian components, iterating through $\Theta = \{1,...,t\}$, where t denotes the maximum
number of allowed WGD events. Only the set of bins in the largest cluster is chosen as
the set of "balanced" bins. The list of candidates γ is then computed using Equation 38 in
CHISEL's supplementary note. Next, HiScanner introduces a homozygosity testing
framework that leverages local GC bias and allelic imbalance to derive robust per-
segment BAF estimates (Supplemental Notes). The segment-wise RDR is simply the
mean RDR of all bins in a given segment. By modeling both RDR and BAF as Gaussian
mixtures for each allelic copy number state, HiScanner applies the maximum likelihood
formulation in CHISEL (Supplementary Note 5.3) to find the most probable allelic copy
number state for each segment.
570
Simulation of male synthetic diploid X chromosomes

We follow a procedure similar to that described in Luquette *et al*[11] to construct synthetic
diploid X (SDX) chromosomes from three single oligodendrocytes from two neurotypical
male donors. For simplicity, here we denote the individuals as A and B, and the cells as
OligoA, OligoB1, and OligoB2. We simulated three types of CNAs: single copy losses,
single copy gains, and cnLOHs at 10 different sizes following a geometric progression
from 200kb to 5Mb. For each CNA type and size, we randomly generated 100 start sites
within chromosome X, after excluding pseudoautosomal regions and germline CNAs
regions. For single copy losses, we removed reads within the randomly generated start
and end sites from OligoB1. For single copy gains, we added reads mapped to CNA
coordinates in OligoB2 to OligoB1. For cnLOH, we did not make changes to the read
depth (only to BAF, described below), as both copies should be independently sampled in
real data. After the read removal or addition, the modified OligoB1-reads and OligoA-
reads were merged with SAMtools[49] to create the final SDX.
585
We next simulated gHETs and BAFs. To resemble a realistic SNP density, we randomly
selected five positions in each 10 kb region to serve as simulated gHET loci. Only the
reads uniquely mapping to the simulated gHETs are used to compute BAF. For cnLOH,
only the reads from OligoA at CNA regions were used. Additional Poisson noise was
added to allele-specific read depths at gHETs to represent sequencing errors.
Furthermore, we simulated phase-switch errors that can occur during reference-based
phasing by introducing an indicator variable i to determine whether to flip the allele-
specific read depths. Specifically, i follows a binomial distribution with a probability of
0.01.
595
Copy number calling with Ginkgo and CHISEL

Ginkgo was run with default parameters with the "binning" parameter set to "variable
100000 101 bowtie" and hg19 genome. Ginkgo's command line version was downloaded

from https://github.com/robertaboukhalil/ginkgo. For the single-neuron comparison, we
applied CHISEL's complete pipeline to both single-cell PTA data and paired bulk
sequencing data from the same individual. This was done separately for bin sizes of 5 Mb
and 100 kb. Regardless of bin size, we noticed a tendency for high ploidy predictions and
overfitting, similar to what was reported previously[22]. To address this, we fine-tuned
several parameters in the *chisel_calling* module to optimize the method performance.
Specifically, we set the *sensitivity* parameter to 10, *maxploidy* to 3 and *k* to 10. In addition
there was a strong GC bias in CHISEL-preprocessed RDR for 100 kb bins, likely due to
the lack of GC-normalization in the CHISEL pipeline. To ensure fairness in the
comparison of segmentation algorithms during the synthetic diploid experiment, we used
the same RDR and BAF data preprocessed by HiScanner as input for both HiScanner's
and CHISEL's segmentation algorithms. CHISEL's bioconda package was downloaded
from https://github.com/raphael-group/chisel. We use the "hg19/GRCh37+decoy"
reference genome for all experiments in this manuscript.
613
614

HiScanner analysis of single neurons and oligodendrocytes

We analyzed PTA-amplified scWGS data and paired bulk whole-genome sequencing
from two public datasets of eleven neurotypical brains[12,14]. Each brain includes 2 to 3
neurons and oligodendrocytes whose genomes were subject to high-depth sequencing
(20-30x per cell). HiScanner was run separately for each brain, using the PTA and bulk
sequencing data as input and a common bin size of 100kb, as determined by our allelic
dropout size model. Two out of 33 neurons were found to be aneuploid, displaying
extensive copy number losses. No oligodendrocytes were aneuploid. Given our focus on
megabase-scale mosaic CNAs, we excluded the aneuploid neurons from subsequent
analyses of CNA length and prevalence. Single-cell CNA calls that overlapped with bulk
CNA calls were removed. To maintain consistency across samples from both sexes, our
analysis was restricted to autosomal chromosomes.
627
We employed a linear mixed effects (LME) model framework to assess the relationship
between CNA count and age across different cell types and CNA types. The statistical
analyses were implemented using the *statsmodels* library in Python, using the *mixedlm*
function from the *statsmodels.formula.api* module. We analyzed four combinations: gains
in neurons, gains in oligodendrocytes, losses in neurons, and losses in oligodendrocytes.
The dependent variable in our LME models was the CNA count (*cna_count*), with age as
the predictor variable. To account for the within-individual correlation of CNA counts,
we included random intercepts by specifying the individual's identifier (*indID*) as the
grouping variable for random effects. This approach allowed us to model the intra-subject
variability and adjust for the non-independence of the data. The formula for the LME
models was set as *cna_count ~ age*, fitting each of the four specified datasets.
639
640

Estimation of false-discovery rate (FDR)

We computed mean FDR values for each CNA size and type assessed in the synthetic diploid experiment. Each CNA detected in the single neuron and oligodendrocyte data was matched with the FDR value from the same event type (e.g., gain or loss) with the largest size smaller than the CNA event. The final FDR estimate is then calculated as the mean of FDRs across all CNAs. This produces a conservative estimate of FDR since FDR increases as CNA size decreases.

Development of ViScanner

We developed a web-based visualization platform (https://parklab.github.io/ViScanner/) in order to visualize HiScanner output. It is a React-based web application hosted on Github. The interactive visualization is based on Higlass[25]. Custom plugin tracks, tailored to the output files of HiScanner, were created for HiGlass. When using ViScanner, the user's input files are loaded into the local browser cache and are not transferred to a server, thus enabling safe use with confidential data.

Meningioma data generation

The samples were collected from the operation room and directly frozen in -80°C after written informed consent was obtained under the auspices of a human subjects institutional review board protocol approved by the Brigham and Women's Hospital. The histological WHO Grade for each meningioma was determined by board-certified neuropathologists. The samples were shipped to Bioskryb, Inc. and stored at -80C until nuclei isolation. Nuclei were isolated using the NUC-V2 protocol on an S2 Genomics Singulator 100 machine. ResolveDNA reactions were run for 1 plate for each sample for whole-genome amplification. All cells passing quality control metrics (PTA yield) were then subjected to library preparation and sequenced on a NextSeq 1000 instrument targeting 8M paired-end reads at 2x150bp.

HiScanner analysis of the single-cell meningioma dataset

HiScanner was run on the entire set of initial and recurrent meningioma scWGS as well as paired tumor and blood bulk sequencing, with a bin size of 500kb. Hierarchical clustering was performed using R package *hclust* with complete linkage and canberra distance.

Somatic SNV analysis of the single-cell meningioma dataset

GATK4 Mutect2[39] with default parameters was run on tumor bulk separately for initial and recurrent samples, using the blood bulk as control. Only variants with "PASS" flag were kept. SECEDO was run on the entire set of initial and recurrent meningioma single cells, with pileup positions restricted to Mutect2-called somatic SNVs.

Data availability

Newly generated scWGS data for 179 PTA-amplified single-cells and two matched tumor/blood bulk WGS of the initial and recurrent meningioma samples will be accessible upon the publication of this work. Further details regarding the data release, including access procedures, will be provided at that time. Previously generated scWGS and matched bulk WGS of PTA neurons are available for download at dbGaP, accession number phs001485.v3.p1. scWGS of MDA and PTA oligodendrocytes are available for download at NIAGADs with accession number NG00162.

Code availability

HiScanner is open-source software and is available on https://pypi.org/project/hiscanner/. HiScanner output can be visualized through ViScanner (https://parklab.github.io/ViScanner/). Scripts to reproduce the results of the primary analyses will be made available on GitHub (https://github.com/parklab/HiScanner).

Acknowledgments

We extend our deepest gratitude to the donors and their families, whose generous contributions are pivotal to the advancement of science. This work was supported by R01HG012573 and R01CA269805 to PJP, F31CA264958 to VVV, and a Sontag Foundation grant to MDJ. We thank Teng Gao, Viktor Ljungström, Antuan Tran, and Dominika Maziec, for their invaluable input to this work.

Author contributions

Study design: Y.Z., L.J.L., V.V.V., P.J.P; HiScanner algorithm and software development: Y.Z.; synthetic data generation and caller performance benchmark: Y.Z., L.J.L.; meningioma somatic SNV clustering analysis: Y.Z.; interpretation of non-diseased brain CNA call set: D.D.S. C.A.W.; generation and interpretation of meningioma data: H.W.Y., M.D.J; implementation of multi-sample segmentation module: X.W, R.X.; implementation of ViScanner: A.D.V.; manuscript writing: Y.Z, P.J.P, L.J.L. with contributions from all authors.

Competing Interests

P.J.P. is a member of the scientific advisory board for Bioskryb Genomics, Inc.

References

1. Svensson, V., Vento-Tormo, R. & Teichmann, S. A. Exponential scaling of single-cell RNA-seq in the past decade. *Nature Protocols* vol. 13 Preprint at https://doi.org/10.1038/nprot.2017.149 (2018).
2. Saliba, A. E., Westermann, A. J., Gorski, S. A. & Vogel, J. Single-cell RNA-seq: Advances and future challenges. *Nucleic Acids Research* vol. 42 Preprint at https://doi.org/10.1093/nar/gku555 (2014).

3.   Buenrostro, J. D. *et al.* Single-cell chromatin accessibility reveals principles of
regulatory variation. *Nature* **523**, (2015).
4.   Ramani, V. *et al.* Massively multiplex single-cell Hi-C. *Nat Methods* **14**, (2017).
5.   Nagano, T. *et al.* Single-cell Hi-C reveals cell-to-cell variability in chromosome
structure. *Nature* **502**, (2013).
6.   Arneson, N., Hughes, S., Houlston, R. & Done, S. Whole-genome amplification by
degenerate oligonucleotide primed PCR (DOP-PCR). *Cold Spring Harb Protoc* **3**,
(2008).
7.   Laks, E. *et al.* Clonal decomposition and DNA replication states defined by scaled
single-cell genome sequencing. *Cell* **179**, 1207–1221 (2019).
8.   Zahn, H. *et al.* Scalable whole-genome single-cell library preparation without
preamplification. *Nat Methods* **14**, 167–173 (2017).
9.   Andor, N. *et al.* Joint single cell DNA-seq and RNA-seq of gastric cancer cell lines
reveals rules of in vitro evolution. *NAR Genom Bioinform* **2**, lqaa016 (2020).
10.   Gonzalez-Pena, V. *et al.* Accurate genomic variant detection in single cells with
primary template-directed amplification. *Proceedings of the National Academy of
Sciences* **118**, e2024176118 (2021).
11.   Luquette, L. J., Bohrson, C. L., Sherman, M. A. & Park, P. J. Identification of
somatic mutations in single cell DNA-seq using a spatial model of allelic
imbalance. *Nat Commun* **10**, 3908 (2019).
12.   Luquette, L. J. *et al.* Single-cell genome sequencing of human neurons identifies
somatic point mutation and indel enrichment in regulatory elements. *Nat Genet* **54**,
1564–1571 (2022).
13.   Bizzotto, S. *et al.* Landmarks of human embryonic development inscribed in
somatic mutations. *Science (1979)* **371**, (2021).
14.   Ganz, J. *et al.* Contrasting patterns of somatic mutations in neurons and glia reveal
differential predisposition to disease in the aging human brain. *bioRxiv* 2021–2023
(2023).
15.   Chronister, W. D. *et al.* Neurons with complex karyotypes are rare in aged human
neocortex. *Cell Rep* **26**, 825–835 (2019).
16.   McConnell, M. J. *et al.* Mosaic copy number variation in human neurons. *Science
(1979)* **342**, 632–637 (2013).
17.   Xi, R., Lee, S., Xia, Y., Kim, T.-M. & Park, P. J. Copy number analysis of whole-
genome data using BIC-seq2 and its application to detection of cancer
susceptibility variants. *Nucleic Acids Res* **44**, 6274–6286 (2016).
18.   Van Loo, P. *et al.* Allele-specific copy number analysis of tumors. *Proceedings of
the National Academy of Sciences* **107**, 16910–16915 (2010).
19.   Nik-Zainal, S. *et al.* The life history of 21 breast cancers. *Cell* **149**, 994–1007
(2012).
20.   Zaccaria, S. & Raphael, B. J. Characterizing allele-and haplotype-specific copy
numbers in single cells with CHISEL. *Nat Biotechnol* **39**, 207–214 (2021).
21.   Wu, C.-Y. *et al.* Integrative single-cell analysis of allele-specific copy number
alterations and chromatin accessibility in cancer. *Nat Biotechnol* **39**, 1259–1269
(2021).
22.   Sun, C. *et al.* Mapping the Complex Genetic Landscape of Human Neurons.
bioRxiv 2023 (2023).

23. Garvin, T. *et al.* Interactive analysis and assessment of single-cell copy-number
variations. *Nat Methods* **12**, 1058–1060 (2015).
24. Rozhoňová, H. *et al.* SECEDO: SNV-based subclone detection using ultra-low
coverage single-cell DNA sequencing. *Bioinformatics* **38**, 4293–4300 (2022).
25. Kerpedjiev, P. *et al.* HiGlass: web-based visual exploration and analysis of
genome interaction maps. *Genome Biol* **19**, 125 (2018).
26. Zhang, C. Z. *et al.* Calibrating genomic and allelic coverage bias in single-cell
sequencing. *Nat Commun* **6**, (2015).
27. Dean, F. B. *et al.* Comprehensive human genome amplification using multiple
displacement amplification. *Proceedings of the National Academy of Sciences* **99**,
5261–5266 (2002).
28. Browning, B. L. & Browning, S. R. Genotype error biases trio-based estimates of
haplotype phase accuracy. *Am J Hum Genet* **109**, (2022).
29. Loh, P. R., Palamara, P. F. & Price, A. L. Fast and accurate long-range phasing in
a UK Biobank cohort. *Nat Genet* **48**, (2016).
30. Browning, B. L., Tian, X., Zhou, Y. & Browning, S. R. Fast two-stage phasing of
large-scale sequence data. *Am J Hum Genet* **108**, (2021).
31. Loh, P.-R. *et al.* Reference-based phasing using the Haplotype Reference
Consortium panel. *Nat Genet* **48**, 1443–1448 (2016).
32. Olshen, A. B., Venkatraman, E. S., Lucito, R. & Wigler, M. Circular binary
segmentation for the analysis of array-based DNA copy number data. *Biostatistics*
5, 557–572 (2004).
33. Mallory, X. F., Edrisi, M., Navin, N. & Nakhleh, L. Assessing the performance of
methods for copy number aberration detection from single-cell DNA sequencing
data. *PLoS Comput Biol* **16**, e1008012 (2020).
34. Knouse, K. A., Wu, J. & Amon, A. Assessment of megabase-scale somatic copy
number variation using single-cell sequencing. *Genome Res* **26**, 376–384 (2016).
35. Knouse, K. A., Wu, J., Whittaker, C. A. & Amon, A. Single cell sequencing
reveals low levels of aneuploidy across mammalian tissues. *Proc Natl Acad Sci U*
S A **111**, 13409–13414 (2014).
36. Kalef-Ezra, E. *et al.* Single-cell somatic copy number variants in brain using
different amplification methods and reference genomes. *bioRxiv* (2023)
doi:10.1101/2023.08.07.552289.
37. Keric, N. *et al.* Impact of postoperative radiotherapy on recurrence of primary
intracranial atypical meningiomas. *J Neurooncol* **146**, 347–355 (2020).
38. Chen, W. C. *et al.* Radiotherapy for meningiomas. *J Neurooncol* **160**, 505–515
(2022).
39. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297–1303
(2010).
40. Steele, C. D. *et al.* Signatures of copy number alterations in human cancer. *Nature*
606, (2022).
41. Cai, X. *et al.* Single-Cell, Genome-wide Sequencing Identifies Clonal Somatic
Copy-Number Variation in the Human Brain. *Cell Rep* **8**, (2014).

42.   Hastings, P. J., Lupski, J. R., Rosenberg, S. M. & Ira, G. Mechanisms of change in
813   gene copy number. *Nature Reviews Genetics* vol. 10 Preprint at
814   https://doi.org/10.1038/nrg2593 (2009).

815   43.   Suberbielle, E. *et al.* Physiologic brain activity causes DNA double-strand breaks
816   in neurons, with exacerbation by amyloid-β. *Nat Neurosci* **16**, (2013).

817   44.   Yang, L. *et al.* An enhanced genetic model of colorectal cancer progression history.
818   *Genome Biol* **20**, (2019).

819   45.   Consortium, 1000 Genomes Project & others. A global reference for human
820   genetic variation. *Nature* **526**, 68 (2015).

821   46.   Sherry, S. T., Ward, M. & Sirotkin, K. dbSNP—database for single nucleotide
822   polymorphisms and other classes of minor genetic variation. *Genome Res* **9**, 677–
823   679 (1999).

824   47.   Delaneau, O., Zagury, J.-F. & Marchini, J. Improved whole-chromosome phasing
825   for disease and population genetic studies. *Nat Methods* **10**, 5–6 (2013).

826   48.   Delaneau, O. & Marchini, J. Integrating sequence and array data to create an
827   improved 1000 Genomes Project haplotype reference panel. *Nat Commun* **5**, 3934
828   (2014).

829   49.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,
830   giab008 (2021).

Fig 1. HiScanner algorithm overview. HiScanner starts with scWGS data as input. In the preprocessing step, gHETs are called from bulk and phased using population-level reference panels. Normalization is performed at the nucleotide level to remove GC bias from read depth signals. Subsequently, a two-state hidden Markov model is employed to estimate ADO event lengths, which are then used to find the optimal bin size by setting a cutoff at the top 5th percentile of the length distribution. For each cell, HiScanner aggregates allele- and haplotype-specific read counts per bin to compute BAF and performs semi-parametric regression to compute RDR. In the segmentation step, HiScanner jointly partitions the genomes across all cells using BIC to identify breakpoints. An LOH test is then used to further refine the BAF estimates within each segment (see Supp. Note for details). In the variant calling and interpretation step, for each cell, HiScanner simultaneously infers the cell-specific scale factor and allelic copy number for each segment by modeling the RDR and BAF as Gaussian distributions. A web-based visualization tool, ViScanner, is also provided, which enables interactive visualization of HiScanner CNA calls at any resolution. Other downstream analyses of HiScanner output include tasks such as phylogenetic inference and correlation of CNAs with SNVs and indels. HiScanner's key contributions are highlighted in light red boxes.

Fig 2. Method comparisons on simulated data. (a) Construction of synthetic diploid X chromosomes. First, two single cells are obtained from different male donors. Next, reads are extracted from chromosome X, excluding reads from pseudoautosomal regions, and germline or somatic CNAs identified in the original cells are removed. This serves as a template for creating various non-clonal CNA conditions, including single copy loss (1|0), single copy gain (2|1), and cnLOH (2|0), after which the reads are merged to create a BAM file for each mutated synthetic diploid chromosome X. Evaluations of precision and sensitivity for each caller for simulated single copy losses (b-c), single copy gains (e-f), and cnLOHs (h-i). (d-g) Examples illustrating caller discrepancies in detecting a simulated loss (d) and a simulated gain (g).

Fig 3. Method comparisons in a single, aneuploid PTA neuron. a. BAF track of single neuron 5823PFC-B using 100 kb mappable bins. Black arrows indicate the example regions in f-g. b-e. Corresponding RDR tracks from 5823PFC-B overlaid with total copy number calls reported by different methods: HiScanner on 100 kb bins (b), Ginkgo on 100 kb bins (c), CHISEL on 100 kb bins (d), and CHISEL after parameter optimization on 5 Mb bins (e). f. An example of a diploid region showing uneven read coverage. The shaded region indicates a false positive call made by Ginkgo. g. An example showing a 1 Mb loss on chromosome 13. The shaded region indicates the location where the CNA boundary misaligns with the changepoint of BAF signals. h. RDR distribution for bins inside centromeric regions. CHISEL*: CHISEL on 5 Mb bins. i. Breakpoint density of each method's call set. j-k. Length distribution of BAF-concordant or BAF-discordant loss events called by HiScanner (j) and Ginkgo (k). All HiScanner calls were BAF-concordant (i.e., y value=0 for the red bars).

Fig 4. Application to somatic mosaicism in human brain cells. a. Experimental workflow. Nuclei from neurons and oligodendrocytes were isolated from 11 human brains, followed by whole-genome amplification, sequencing, and somatic CNA detection using Ginkgo and HiScanner. Created with BioRender.com. b-c. Length distribution of HiScanner CNA calls, categorized by type and clonality, in 31 neurons (b) and 32 oligodendrocytes (c). d-e. Length distribution of Ginkgo CNA calls. f. Example of a diploid region with uneven coverage where a spurious loss is called by Ginkgo. g. Example of two adjacent single copy losses, one of which is missed by Ginkgo. h. Example of a 2Mb single copy loss. Shaded regions indicate positions where Ginkgo calls are discrepant with HiScanner and BAF. i-j. Number of Ginkgo calls that are concordant or discordant with BAF, categorized by length and cell types.

Fig 5. Application to low-coverage scWGS of a recurrent meningioma tumor pair. a. 179 single cells from a paired initial and recurrent meningioma tumor sample were amplified by PTA and sequenced. b-d. Total copy number (b), RDR (c) and BAF (d) matrices output by HiScanner. Cell ordering is determined by hierarchical clustering applied to the copy number matrix. Each row represents one cell and each column corresponds to one genomic bin (500 kb mappable). e-g. Total copy number and BAF profiles of three representative cells from major clones of each tumor grade as well as a Grade III tetraploid clone. h-i. Scatter plots of the second and third eigenvectors of the Laplacian similarity matrix output by SECEDO, colored by SECEDO-inferred SNV clusters (h) and HiScanner-based CNA hierarchical clusters (i).

	Input signal type(s)	GC correction	Bin size (Mb)	Segmentation algorithm	Joint segmentation across cells	Integration of phasing and BAF
Ginkgo	RDR	Bin-level	0.5	CBS	No	N/A
SCOVAL	RDR, phased BAF	Bin-level	0.5	CBS	No	CNA filtering step
HiScanner	RDR, phased BAF	Nucleotide-level	0.1	BIC-based	Yes	CNA calling step
CHISEL	RDR, phased BAF	Not performed	5	K-means clustering	Yes	Segmentation step
Alleloscope	RDR, phased BAF, bulk segmentation output	No performed	N/A*	HMM	No	CNA calling step

Table 1. Qualitative comparison of five single cell copy number callers (using default parameters).
RDR: read-depth ratio. BAF: B-allele frequency. CBS: circular binary segmentation. BIC: Bayesian Information Criterion. HMM: hidden Markov model. *: Alleloscope models gHETs directly and does not perform binning.